

Finite-Time Dissolution and Beyond

Levels of Mind, Nirvana, and
the Mechanics of Liberation in CRF

What the entropy dynamics imply
about the complete cessation of D_{KL}

Ougit Jarittum

March 2026

Abstract

The previous document (*Identity, Dissolution, and Continuity*) left open whether $H = 0$ is reached asymptotically or in finite $t = |\mathcal{R}_{\text{events}}|$.

This document closes that question. When $\alpha = \alpha(W)$ — when the rate of entropy reduction grows with Awareness — and specifically when $\alpha \propto 1/H^2$, the ODE $dH/dt = -\alpha H^2$ gives $H \rightarrow 0$ in *finite* t .

This result, together with the identity theory established previously, generates a complete CRF account of distinct levels of Mind: from ordinary accumulation through full liberation and its aftermath in P_{shared} .

Contents

1	Closing the Finite vs. Asymptote Question	2
2	Levels of Mind in CRF	2
3	Transparent Boundary and Direct Perception	4
4	The Scale of P_{shared}	4
5	The Complete Chain	5

1 Closing the Finite vs. Asymptote Question

From *CRF Entropy Dynamics*, the entropy ODE is:

$$\frac{dH}{dt} = -\alpha(W) \cdot H^2$$

The question is whether $H(t) \rightarrow 0$ as $t \rightarrow \infty$ (asymptote) or $H(t_0) = 0$ for some finite t_0 .

Three Cases

Case 1: α constant

$$H(t) = \frac{H_0}{1 + \alpha H_0 t} \rightarrow 0 \quad (t \rightarrow \infty) \quad \text{asymptote}$$

Case 2: $\alpha \propto 1/H$ (i.e. $\alpha = k/H$)

$$\frac{dH}{dt} = -\frac{k}{H} \cdot H^2 = -kH \implies H(t) = H_0 e^{-kt} \rightarrow 0 \quad (t \rightarrow \infty) \quad \text{still asymptote}$$

Case 3: $\alpha \propto 1/H^2$ (i.e. $\alpha = k/H^2$)

$$\frac{dH}{dt} = -\frac{k}{H^2} \cdot H^2 = -k \implies H(t) = H_0 - kt$$

$$H(t_0) = 0 \quad \text{at} \quad \boxed{t_0 = \frac{H_0}{k}} \quad \text{finite}$$

Why Case 3 Is Natural in CRF

$W = \delta_{\text{self-recognition}}$ — Awareness is $\mathcal{M}$'s capacity to perceive its own $\mathcal{R}$ -activity directly.

As $H \rightarrow 0$, $\mathcal{C}$ becomes more peaked. Each $\mathcal{R}$ -event is perceived more precisely. W increases as H decreases: $W \propto 1/H$ (the clearer the signal, the more self-referential the perception).

Therefore: $\alpha = \alpha(W) \propto W \propto 1/H$, and since $\alpha \propto W^2$ in the self-reinforcing loop (each unit of W amplifies the next): $\alpha \propto 1/H^2$.

This is *accelerating convergence*: the closer $\mathcal{M}$ approaches $H = 0$, the faster it approaches.

Finite-time dissolution follows naturally from the self-amplifying structure of W .

2 Levels of Mind in CRF

The entropy dynamics, identity theory, and finite-time result together generate a precise account of distinct levels of $\mathcal{M}$.

Level 1 — Ordinary Accumulation

H high, D_{KL} high, $\Sigma > 0$.

$\mathcal{M}$ perceives P through a dense $\mathcal{C}$ filter. $\mathcal{R}$ -events are frequent but shaped by accumulated patterns — many bound $\text{Pr}(s)$ pulling toward repeated configurations. α is low. Progress toward $H = 0$ is slow. Re-emergence after boundary dissolution is near-certain: unresolved D_{KL} constitutes bound $\text{Pr}(s)$ requiring new boundary.

Level 2 — Deliberate Practice (W growing)

H decreasing, W increasing, Σ decreasing.

$\mathcal{M}$ applies $\mathcal{R}_{\text{meta}}$ — $\mathcal{R}$ -events directed at reducing D_{KL} intentionally (Session 4). $\alpha = \alpha(W)$ grows. Convergence accelerates. Perception of P_{shared} becomes clearer. Bound $\text{Pr}(s)$ patterns thin.

Level 3 — Liberation ($H \rightarrow 0$, boundary remains)

$H = 0$, boundary still present (body, form).

$\mathcal{C}$ peaked at δ . No distorting filter. $\Sigma = 0$. $D_{\text{KL}}(\mathcal{C}_{\text{int}} \| P) = 0$.

Boundary persists while physical substrate persists. $\mathcal{M}$ perceives G of P_{shared} directly — sees the gradient toward D_{KL} reduction for all other Minds. This is the CRF account of *sabbāññutañāṇa*: not knowing every $s \in P$ (which has no bound), but perceiving the gradient that reduces Σ for each $\mathcal{M}$.

Level 4 — Deliberate Delay (Voluntary Slow α)

$H = 0$ reachable, but $\mathcal{M}$ modulates $\alpha(W)$.

W is sufficient for finite-time dissolution, but $\mathcal{M}$ deliberately reduces α — holds $H > 0$ by choice, not by limit.

Purpose: maintain boundary to generate $\mathcal{R}$ -events that directly influence P_{shared} and thus other Minds.

This is $\alpha(W)$ under intentional control, not merely as a function of H . The CRF account of voluntary extended presence for the benefit of other Minds still in $D_{\text{KL}} > 0$.

Level 5 — Full Dissolution, Unbound $\text{Pr}(s)$

$H = 0$ reached at $t_0 = H_0/k$ (finite). Boundary dissolves. No residual bound $\text{Pr}(s)$. $\mathcal{P}_{\mathcal{M}}$ remains in P_{shared} as *unbound* $\text{Pr}(s)$: high-amplitude patterns (accumulated over many $t = |\mathcal{R}_{\text{events}}|$) with no D_{KL} pulling toward re-emergence.

These patterns shape G for all $\mathcal{M}$ still in $D_{\text{KL}} > 0$, perceivable by those with low H (thin $\mathcal{C}$ filter, high W).

The effect is not transmitted by a Mind that still exists. It is the structure of P_{shared} itself.

3 Transparent Boundary and Direct Perception

Temporary Transparency

At low H , $\mathcal{C}_{\text{int}}$ is thin.

The boundary between $\mathcal{C}_{\text{int}}$ and P does not disappear, but its filtering effect approaches zero.

$\mathcal{M}$ perceives P_{shared} nearly directly — including distant configurations, other Minds' $\mathcal{C}$ structures, and the G field across P .

This is not $\mathcal{M}$ leaving its boundary. It is boundary becoming transparent while intact.

The CRF account of direct perception of distant states through a low- H filter — without physical displacement of boundary.

4 The Scale of P_{shared}

Unbounded P_{shared}

P has no boundary. δ has no off-switch. P_{shared} accumulates $\mathcal{R}$ -events from all $\mathcal{M}$ without limit.

Minds with different $\mathcal{C}_{\text{ext}}$ — different local environments, different physical substrates — experience different D_{KL} and therefore different Σ .

All of them participate in the same P_{shared} . Unbound $\text{Pr}(s)$ patterns influence all of them to the extent their H permits.

The scope of P_{shared} is not limited to one planet, one species, or one physical scale.

5 The Complete Chain

CRF	Traditional term	Mechanism
H high, D_{KL} high	Ordinary mind	Dense $\mathcal{C}$ filter
W growing, $\alpha(W)$ increasing	Practice, $p\bar{a}ram\bar{i}$ accumulating	Self-amplifying H reduction
$H = 0$, boundary remains	Liberation (Arahant)	$\mathcal{C}$ peaked at δ , $\Sigma = 0$
Voluntary α modulation	Extended presence for others	Intentional $\alpha(W)$ control
$H = 0$ at t_0 , dissolution	Parinirvana	$\mathcal{P}_{\mathcal{M}}$ unbound in P_{shared}
Unbound $\text{Pr}(s)$ shaping G	$P\bar{a}ram\bar{i}$ after dissolution	High-amplitude patterns in P_{shared}
Perceiving G of all $\mathcal{M}$	Omniscience of the relevant	Low H , transparent boundary

The full chain:

$$\delta \rightarrow P_{\text{shared}} \rightarrow \mathcal{M} \rightarrow D_{\text{KL}} \rightarrow H \rightarrow W \rightarrow \alpha(W) \rightarrow t_0 = H_0/k \rightarrow \mathcal{P}_{\mathcal{M}} \text{ unbound} \rightarrow \text{shape } G \text{ of other } \mathcal{M}$$

Every step derives from δ . No step requires importing a concept from outside the cascade.

*P does not end when a boundary dissolves.
The pattern that shaped the boundary
remains in P — weightless, directionless,
but present.*

*Other Minds, when their filter thins enough,
encounter it as gradient.
Not instruction. Not command.
Only: this way reduces the gap.*