

Lung Epithelial Fluid Transport and the Resolution of Pulmonary Edema

MICHAEL A. MATTHAY, HANS G. FOLKESSON, AND CHRISTINE CLERICI

Cardiovascular Research Institute and Departments of Medicine and Anesthesia, University of California, San Francisco, California; Department of Physiology, Northeastern Ohio Universities College of Medicine, Rootstown, Ohio; and Department of Physiology, Faculté de Médecine de Bobigny, Université Paris 13, Paris, France

I. Introduction	569
II. Salt and Water Transport Across the Distal Pulmonary Epithelia	570
A. Structural features of the distal pulmonary epithelia	570
B. Vectorial fluid transport in the intact lung	572
C. Sodium transport in cultured alveolar type II cells	573
D. Phenotype of cultured alveolar type II cells	575
E. Lung epithelial fluid transport	575
III. Regulation of Salt and Water Transport	578
A. Catecholamine-dependent regulation	578
B. Catecholamine-independent regulation	582
IV. Mechanisms That Impair Vectorial Fluid Transport	584
A. Atrial natriuretic peptide	584
B. Hypoxia	585
C. Anesthetics	585
D. Reactive oxygen and nitrogen species	586
V. Epithelial Fluid Transport Under Pathological Conditions	586
A. Clinical studies	586
B. Experimental studies	588
VI. Future Directions	591

Matthay, Michael A., Hans G. Folkesson, and Christine Clerici. Lung Epithelial Fluid Transport and the Resolution of Pulmonary Edema. *Physiol Rev* 82: 569–600, 2002; 10.1152/physrev.00003.2002.—The discovery of mechanisms that regulate salt and water transport by the alveolar and distal airway epithelium of the lung has generated new insights into the regulation of lung fluid balance under both normal and pathological conditions. There is convincing evidence that active sodium and chloride transporters are expressed in the distal lung epithelium and are responsible for the ability of the lung to remove alveolar fluid at the time of birth as well as in the mature lung when pathological conditions lead to the development of pulmonary edema. Currently, the best described molecular transporters are the epithelial sodium channel, the cystic fibrosis transmembrane conductance regulator, $\text{Na}^+\text{-K}^+\text{-ATPase}$, and several aquaporin water channels. Both catecholamine-dependent and -independent mechanisms can upregulate isosmolar fluid transport across the distal lung epithelium. Experimental and clinical studies have made it possible to examine the role of these transporters in the resolution of pulmonary edema.

I. INTRODUCTION

In the last 20 years, the mechanisms that regulate active salt and water transport by the alveolar and distal airway epithelium of the lung have emerged as a new area of research with important implications for understanding lung fluid balance under both normal and pathological conditions. Studies of epithelial fluid transport by the distal pulmonary epithelium have provided important

new concepts regarding the resolution of pulmonary edema, a common clinical problem. Before 1982, there was no information on how lung fluid balance was regulated across the distal airway and alveolar epithelial barriers. However, in 1982, new work provided evidence that fluid balance in the lung was regulated by active ion transport mechanisms (131, 214, 229). For many years, it was generally believed that differences in hydrostatic and protein osmotic pressures (Starling forces) accounted for

the removal of excess fluid from the airspaces of the lung. This misconception persisted in part because some experiments that measured solute flux across the epithelial and endothelial barriers of the lung were done at room temperature (347), and the studies were done in dogs, a species that turned out to have a very low rate of active sodium and fluid transport (28). Also, until the early 1980s, there were no satisfactory animal models to study the resolution of alveolar edema, and the isolation and culture of alveolar epithelial type II cells was just becoming a useful experimental method. Although the removal of interstitial pulmonary edema by lung lymphatics and the lung microcirculation was discussed by Staub in 1974 in his review of pulmonary edema (333), there was no information on how pulmonary edema was removed from the distal airspaces of the lung.

This review provides a perspective on how in vivo and in vitro experiments in the last 20 years have provided a new understanding of the regulation of lung fluid balance by active transport mechanisms across both the alveolar and distal airway epithelium. On balance, there is convincing evidence that the vectorial transport of salt and water across the alveolar and distal airway epithelium is the primary determinant of fluid clearance, thus accounting for the ability of the lung to remove alveolar fluid at the time of birth as well as in the mature lung when pathological conditions lead to the development of pulmonary edema. Most of this review is focused on the function of the adult, not the fetal or the perinatal, lung.

II. SALT AND WATER TRANSPORT ACROSS THE DISTAL PULMONARY EPITHELIA

With few exceptions, the general model for transepithelial fluid movement is that active salt transport drives osmotic water transport. This paradigm is probably correct for fluid clearance from the distal airspaces of the lung (22, 30, 91, 92, 94, 228, 229). The results of several in vivo studies have demonstrated that changes in hydrostatic or protein osmotic pressures cannot account for the removal of excess fluid from the distal airspaces (22, 28, 30, 94, 215, 219, 226–229, 343). Furthermore, pharmacological inhibitors of sodium transport can reduce the rate of fluid clearance in the lungs of several different species, including the human lung (30, 110, 164, 217, 228, 258, 266, 308, 327). In addition, there is good evidence that isolated epithelial cells from the distal airspaces of the lung actively transport sodium and other ions. Less than 10 years ago, the pathways for water clearance across alveolar and distal airway epithelium were unknown. Since then, several studies have demonstrated the presence of several functional water channels, aquaporins, in the lung.

This section reviews relevant morphological features

of the distal lung epithelium with particular reference to their potential contribution to solute and water transport capacity. Next, based on in vivo and in vitro studies, there will be a discussion of some of the evidence supporting the hypothesis that active ion transport is the primary mechanism for driving vectorial fluid transport from the airspaces to the interstitium of the lung. The last section describes the evidence for the role of epithelial sodium channel (ENaC), $\text{Na}^+\text{-K}^+\text{-ATPase}$, and aquaporins in mediating salt and water transport in distal lung epithelia.

A. Structural Features of the Distal Pulmonary Epithelia

Although the large surface area of the alveoli seems to favor the hypothesis that most fluid reabsorption occurs at the alveolar level, active fluid reabsorption could occur across all of the different segments of the pulmonary epithelium of the distal airspaces of the lung. The precise contribution of each of the anatomic segments of the distal airspaces to fluid reabsorption is not firmly established. Therefore, it is important to review briefly the morphological features of the distal lung epithelia and some of the information regarding their transport characteristics.

The human lung consists of a series of highly branched hollow tubes that end blindly in the alveoli, with the conducting airways (the cartilaginous trachea, bronchi, and the membranous bronchioles) occupying the first 16 generations of airways (281). Gas exchange primarily occurs in the branches that make up the last seven generations including respiratory bronchioles, alveolar ducts, alveolar sacs, and alveoli (334). The airways, $\sim 1.4 \text{ m}^2$ in the adult human lung (376), and alveoli, $\sim 143 \text{ m}^2$ in the adult human lung (376), constitute the interface between lung parenchyma and the external environment and are lined by a continuous epithelium. The distal airway epithelium is composed of terminal respiratory and bronchiolar units with polarized epithelial cells that have the capacity to transport sodium and chloride including ciliated Clara cells and nonciliated cuboidal cells. The alveoli themselves are composed of a thin alveolar epithelium ($0.1\text{--}0.2 \mu\text{m}$) that covers 99% of the airspace surface area in the lung and contains thin, squamous type I cells and cuboidal type II cells (334, 376) (Fig. 1). The alveolar type I cell covers 95% of the alveolar surface (376). The close apposition between the alveolar epithelium and the vascular endothelium facilitates efficient exchange of gases, but also forms a tight barrier to movement of liquid and proteins from the interstitial and vascular spaces, thus assisting in maintaining relatively dry alveoli (377).

The tight junctions are the critical structures for the barrier function of the alveolar epithelium. Tight junctions connect adjacent epithelial cells near their apical




FIG. 1. A schematic diagram of the distal pulmonary epithelium that is relevant for salt and water transport.

surfaces, thereby maintaining apical and basolateral cell polarity (318). Ion transporters and other membrane proteins are asymmetrically distributed on opposing cell surfaces, conferring vectorial transport properties to the epithelium. In addition, the tight junction itself might contain discrete ion-selective pores (139). In the past, it was thought that tight junctions were rigid structures, which physically restrict passage of larger molecules; however, the permeability of tight junctions is dynamic and regulated, in part, by cytoskeletal proteins and intracellular calcium concentration (318). Physiological studies of the barrier properties of tight junctions in the alveolar epithelium indicate that diffusion of water-soluble solutes between alveolar epithelial cells is much slower than through the intercellular junctions of the adjacent lung capillaries (145, 279, 317, 320, 333). Based on tracer fluxes of small water-soluble solutes across the air-blood barrier of the distal airspaces, the effective pore radii were 0.5–0.9 nm in the distal respiratory epithelium and 6.5–7.5 nm in the capillary endothelium (346). In addition, studies of protein flux across both the endothelial and epithelial barriers of the sheep lung indicate that 92% of the resistance to albumin flux across the alveolar barrier resides in the epithelium (134). Removal of large quantities of soluble protein from the airspaces appears to occur primarily by restricted diffusion (119, 150, 151), although there is evidence for some endocytosis and transcytosis of albumin across alveolar epithelium (27, 114, 119, 150, 167, 177, 178, 221).

The most extensively studied cell in the distal pulmonary epithelium is the alveolar type II cell, partly because type II cells can be readily isolated from the lung and studied in vitro. The alveolar type II cell is responsible for the secretion of surfactant (65, 66) as well as vectorial

transport of sodium from the apical to the basolateral surface (67, 132, 214, 216, 217, 228, 238). The active transport of sodium by type II cells appears to provide a major driving force for removal of fluid from the alveolar space. Sodium uptake occurs on the apical surface, partly through amiloride-sensitive and amiloride-insensitive channels. Subsequently, sodium is pumped actively from the basolateral surface into the lung interstitium by Na⁺-K⁺-ATPase (Fig. 1). An ENaC that participates in sodium movement across the cell apical membrane was cloned and characterized in 1994 (52), and work by other investigators has provided new insight into the molecular and biochemical basis for sodium uptake in alveolar epithelial cells (264, 370, 387) (see details below).

The role of the alveolar type I cell in vectorial fluid transport in the lung is unknown, although several investigators are currently trying to assess the potential contribution of the alveolar type I cell to vectorial fluid transport. On the basis of studies in freshly isolated type I cells, it is known that these cells have a high osmotic permeability to water with expression of aquaporin-5 on the apical surface (84). Immunocytochemical studies in the intact lung by some investigators failed to demonstrate the presence of Na⁺-K⁺-ATPase in these cells in vivo (319). However, more recent studies have reported the presence of the α_1 - and α_2 -subunits of Na⁺-K⁺-ATPase in both type I like cells in vitro (303). The presence of the Na⁺-K⁺-ATPase could be consistent with a role for this cell in vectorial fluid transport, although it is not conclusive since the Na⁺-K⁺-ATPase may be needed to maintain cell volume. Also, recent studies of freshly isolated alveolar type I cells from rats demonstrated that these cells express the Na⁺-K⁺-ATPase α_1 - and β_1 -subunit isoforms, but not the α_2 -subunit (38). In the same study, there was

evidence for α_1 Na⁺-K⁺-ATPase expression on the basolateral surface of the alveolar epithelial type I cells in situ in the rat lung. In addition, there is evidence for expression of all the subunits of ENaC in freshly isolated alveolar type I cells from two laboratories (38, 168) as well as in situ in the rat lung (168). Finally, there is some evidence that ²²Na uptake can be partially inhibited by amiloride in the freshly isolated rat alveolar type I cells (168), although definitive studies of cultured, polarized type I cells have not yet been achieved. There is also recent preliminary evidence that β -receptors are expressed on type I cells (201), although this issue is not clear since evidence also has been brought forward for the lack of such expression in type I cells (249). Also, type I cells might be involved in the transport of macromolecules because of the presence of vesicles and caveolin in these cells (50, 171, 247). Although the inability to study alveolar type I cells in culture has hindered progress in assessing their capacity for ion transport and the role they may play in vectorial fluid transport across the alveolar epithelium, the new evidence provides suggestive though not conclusive evidence that alveolar type I cells may participate in vectorial salt transport in the lung.

The alveolar epithelium comprises 99% of the surface area of the lung, a finding that suggests that removal of edema fluid from the lung might primarily occur across the alveolar epithelium. However, this conclusion may be misleading. It has been demonstrated that the distal airway epithelium actively transports sodium, a process that depends on amiloride-inhibitable uptake of sodium on the apical surface and extrusion of sodium through a basolateral Na⁺-K⁺-ATPase (5, 13, 14, 41, 42, 159, 382). It appears that in airways <200 μ m in diameter, sodium absorption predominates (5, 13, 14) (Fig. 1). Secretion of chloride may occur in the trachea and larger airways, and perhaps in distal airways of some animal species in the presence of cAMP stimulation (254). In support of the potential role of distal airway cells is evidence that Clara cells actively absorb sodium and transport from an apical to basal direction (358, 359). Also, there are new data on the possible role of cystic fibrosis transmembrane conductance regulator (CFTR) in upregulating cAMP fluid clearance (see sect. III). This information provides support for a possible role of distal airway epithelia in fluid clearance, since CFTR is expressed abundantly in distal airway epithelial cells. Also, there is new evidence that some alveolar fluid may be removed by convective surface active forces that might propel alveolar edema fluid into the distal airways where some absorption may occur across the respiratory bronchiolar epithelium (263, 372). Thus, even though their surface area is limited, a contribution from distal airway epithelia to the overall fluid transport is probable, especially since cells from the distal airway epithelium primarily transport salt from the apical to the basolateral surface (5, 13, 14, 41, 42, 159).

B. Vectorial Fluid Transport in the Intact Lung

A substantial number of innovative experimental methods have been used to study fluid and protein transport from the distal airspaces of the intact lung including isolated perfused lung preparations, in situ lung preparations, surface fluorescence methods, and intact lung preparations in living animals for short time periods (30–240 min) or for extended time periods (24–144 h). The advantages and disadvantages of these preparations have been reviewed in some detail (228, 313).

The first in vivo evidence that active ion transport could account for the removal of alveolar edema fluid across the distal pulmonary epithelium was obtained in studies of anesthetized, ventilated sheep (229). In those studies, the critical discovery was that isosmolar fluid clearance of salt and water occurred in the face of a rising concentration of protein in the airspaces of the lung, whether the instilled solution was autologous serum or an isosmolar protein solution. The initial protein concentration of the instilled protein solution was the same as the circulating plasma. After 4 h, the concentration of the protein had risen from ~6.5 to 8.4 g/100 ml, while the plasma protein concentration was unchanged. In longer term studies in unanesthetized, spontaneously breathing sheep, alveolar protein concentrations increased to very high levels. After 12 and 24 h, the alveolar protein concentration increased to 10.2 and 12.9 g/100 ml, respectively (226). The overall rise in protein concentration was equivalent to an increase in distal airspace protein osmotic pressure from 25 to 65 cmH₂O (Fig. 2). The concentration of protein in the lung lymph draining the lung interstitium declined, providing further evidence that pro-




FIG. 2. In spontaneously breathing unanesthetized sheep, alveolar fluid protein concentration increased from 6.0 g/dl to a maximum of 13.8 g/dl (*x*-axis) after 24 h. This increase in protein concentration was paralleled by a marked increase in protein oncotic pressure (*y*-axis). [Data adapted from Matthay et al. (226).]

tein-free fluid was being reabsorbed from the distal airspaces into the lung interstitium (30, 229). Also, morphological studies showed that the interstitial fluid did not contain the Evans blue dye-labeled alveolar protein (224). These data provided convincing evidence that active ion transport must be responsible for the fluid clearance, especially in the face of a rising alveolar protein osmotic pressure that was ~ 40 cmH₂O greater than the protein osmotic pressure in the vascular or interstitial spaces of the lung (226, 229).

Other studies in the intact lung have supported the hypothesis that removal of alveolar fluid requires active transport processes. For example, elimination of ventilation to one lung did not change the rate of fluid clearance in sheep, thus ruling out changes in transpulmonary airway pressure as a major determinant of fluid clearance, at least in the uninjured lung (309). Furthermore, if active ion transport were responsible for fluid clearance, then fluid clearance should be temperature dependent. In an *in situ* perfused goat lung preparation, the rate of fluid clearance progressively declined as temperature was lowered from 37 to 18°C (323). Similar results were obtained in perfused rat lungs (306) and *ex vivo* human lung studies (308) in which hypothermia inhibited sodium and fluid transport.

Additional evidence for active ion transport was obtained in intact animals with the use of amiloride, an inhibitor of sodium uptake by the apical membrane of alveolar epithelium and distal airway epithelium. Amiloride inhibited 40–70% of basal fluid clearance in sheep, rabbits, rats, guinea pigs, mice, and in the human lung (21, 22, 30, 74, 94, 121, 164, 258, 266, 308, 327, 386). Amiloride also inhibited sodium uptake in distal airway epithelium from sheep and pigs (5, 13). To further explore the role of active sodium transport, experiments were designed to inhibit Na⁺-K⁺-ATPase. It has been difficult to study the effect of ouabain in intact animals because of cardiac toxicity. However, in the isolated rat lung, ouabain inhibited >90% of fluid clearance (21, 270). Subsequently, after

the development of an *in situ* sheep preparation for measuring fluid clearance in the absence of blood flow, it was reported that ouabain inhibited 90% of fluid clearance over a 4-h period (309).

Important species differences in the basal and stimulated rates of fluid clearance have been identified; not all species respond to cAMP agonists (Table 1). To normalize for differences in lung size or the available surface area, different instilled volumes were used ranging from 1.5 to 13.0 ml/kg. The slowest fluid clearance was measured in dogs (28, 135), intermediate rates of fluid clearance in sheep and goats (30, 226, 229, 323), and the highest basal fluid clearance rates in rabbits, guinea pigs, rats, and mice (121, 124, 164, 258, 327). The basal rate of fluid clearance in the human lung has been difficult to estimate, but based on the isolated, nonperfused human lung model, basal fluid clearance rates appear to be intermediate to fast (308). In fact, the fluid clearance in the *ex vivo* human lung is approximately one-half of the rate in the *ex vivo* rat lung (307). However, as discussed in section v, studies in patients suggest that the rate of maximal fluid clearance from the human lung may be fast. The explanation for the species differences is not apparent, although it may be related to the number or activity of sodium or chloride channels or the density of Na⁺-K⁺-ATPases in alveolar epithelium in different species. Morphometric studies (76, 282) have shown no significant difference in the number of alveolar type II cells in different species. It is also possible that the contribution of the distal airway epithelium may not be uniform in all species.

C. Sodium Transport in Cultured Alveolar Type II Cells

The success in obtaining nearly pure cultures of alveolar epithelial type II cells from rats made it possible to study the transport properties of these cells and relate the results to the findings in the intact lung studies. The initial

TABLE 1. Fluid clearance in different animal species under control conditions and after stimulation with cAMP agonists or inhibition with amiloride or ouabain

Species	Baseline, Fluid Clearance Over 1 h, %	β -Adrenergic Stimulation, % Increase Over Baseline	Amiloride Sensitivity, % Decrease Below Baseline	Ouabain Sensitivity, % Decrease Below Baseline	Reference Nos.
Mouse	48–60	58–71	63–91	80	103, 121, 122, 146, 155, 236
Guinea pig	38	32	26	NA	258
Rat	15–35	45–136	40	30–82	7, 59, 116, 164, 261, 307, 311, 339
Rabbit*	36	0	81	NA	327
Sheep†	8	50–60	42	90	30, 226, 229
Dog†	2–4	73–150	31	84	28, 135, 191, 210
Human†‡	3	72–100	42	50§	307, 308

All values are adapted from other studies. * Values are corrected for 25-min instillation time. † Studies were done originally over 4 h, and the values are extrapolated to 1 h in the table. ‡ Resected lung lobe model. § Ouabain was only applied to the apical surface in the human lung; thus basolateral penetration may have been incomplete. NA, not available.

studies showed that when type II cells were cultured on a nonporous surface such as plastic, they would form a continuous confluent layer of polarized cells after 2–3 days (131, 214). Interestingly, after 3–5 days, small domes of fluid could be appreciated from where the substratum was detached (Fig. 3). The domes were thought to result from active ion transport from the apical to the basal surface with water following passively since they were inhibited by the replacement of sodium by another cation or by pharmacological inhibitors of sodium transport, such as amiloride and ouabain (132). More detailed information on the nature of ion transport across alveolar type II cells was obtained by culturing these cells on porous supports and mounting them in Ussing chambers and measuring short-circuit current and ion flux under voltage-clamp conditions. Confluent monolayers with tight junctions were formed by 4–5 days later, and with considerable care to the culture conditions, it has been possible in some, but not all studies, to obtain monolayers with a high transepithelial resistance ($2,000 \Omega/\text{cm}^2$) (60). Alveolar type II cell monolayers are capable of generating spontaneous transmembrane potential difference (PD) related to electrogenic transport, and it is interesting to note that the spontaneous PD values (0.2–10 mV with apical membrane negative) (60, 73, 337) are similar to those obtained in vivo between the alveolar lumen and the pleural surface (255). When the apical and basolateral compartments are bathed with solutions with identical ionic compositions and the transepithelial potential is maintained (clamped) to zero, the measured current is termed short-circuit current (I_{sc}). The I_{sc} in alveolar type II cells monolayers corresponded closely to the net sodium absorption; substitution of sodium by equimolar concentration of choline in the apical side of monolayers or addition of ouabain that have the ability to block $\text{Na}^+\text{-K}^+\text{-ATPase}$ at the basolateral side, quickly decreased

I_{sc} to zero (60). With the use of labeled sodium (^{22}Na), unidirectional flux from the apical to the basolateral surface exceeded the flux in the opposite direction (176), a finding that was similar to results in isolated perfused lung preparations (21, 22, 74, 93, 94, 133).

The coordinated role of apical and basolateral sodium transport has been studied in several in vitro studies. Sodium ions that enter the epithelial cells at the apical membrane are pumped out of the cells at the basolateral membrane by the $\text{Na}^+\text{-K}^+\text{-ATPase}$ enzyme. Because of the continuous pumping, sodium chemical potential is lower inside the cell. The entry step is passive and then sodium flows down a chemical potential gradient through specialized pathways, where basolateral transport requires energy to move ions against the gradient. Because of the pump activity, potassium electrochemical potential is larger inside the cell, and potassium leaks through the basolateral membrane and is then recycled by the $\text{Na}^+\text{-K}^+\text{-ATPase}$. The pathways for sodium entry into alveolar type II cells are numerous. Amiloride blocked dome formation (131) and decreased I_{sc} in the in vitro studies (60), a finding that supported the critical importance of sodium uptake through an amiloride-sensitive pathway in the apical membrane of alveolar epithelial cells. As already discussed, the efficacy of amiloride as an inhibitor of fluid clearance in the intact lung was demonstrated in several in vivo studies, although the fraction of amiloride-sensitive transport was as low as 40–50% in some lung preparations, particularly the rat and the human lung. The amiloride-insensitive sodium influx may be represented in vivo in part by the $\text{Na}^+\text{-glucose}$ cotransport (21, 22). In vitro, the amiloride-insensitive sodium pathways are not clearly defined, but several sodium cotransporters such as $\text{Na}\text{-amino acid}$, $\text{Na}\text{-phosphate}$, $\text{Na}\text{-proton}$, as well as other cation channels may be involved (46, 68, 69, 256).




FIG. 3. A: light micrograph of domes from primary culture of alveolar epithelial type II cells. The domes formed spontaneously after 3 days in culture. Dome formation was inhibited by amiloride. B: schematic representation of a dome of fluid accumulation between cultured alveolar epithelial type II cell monolayers. The type II cells actively transport sodium from the culture medium to beneath the monolayer (dashed arrows), which results in fluid accumulation between the cell monolayer and the culture dish. The apical cell surface (microvilli) is shown in contact with the medium; the basolateral surface is shown facing the culture dish. [From Mason et al. (214), with permission from the National Academy of Sciences USA.]

D. Phenotype of Cultured Alveolar Type II Cells

One important unresolved issue is to what extent cultured alveolar type II cells retain their *in vivo* phenotype. Within the first 24 h of culture, some of the morphological and functional characteristics of the alveolar type II cells are lost (81, 86). Most surfactant protein expression is downregulated (202), and some transport proteins such as Na-phosphate and Na⁺-K⁺-2Cl cotransport vanish (68, 69). However, active ion transport can be measured for up to 7 days. Also, some data indicate that culture conditions may help the cells to retain features of type II cells. For example, alveolar type II cells cultured on collagen gels with culture medium containing fetal calf serum and with an apical surface exposed to air retain morphological characteristics of alveolar type II cells for a longer duration (85). These cells express surfactant proteins and their mRNAs and express a plasma membrane marker specific for alveolar type II cells. Recent work (163) indicates that *in vitro* culture conditions are an important determinant for the biophysical characteristics of sodium channels. Alveolar type II cells grown on a permeable surface with an air interface and glucocorticoids in the medium maintain alveolar type II cell phenotype and express highly selective sodium channel while cells cultured on glass and in air-liquid interface express a low selective sodium channel.

E. Lung Epithelial Fluid Transport

This section considers pharmacological, biophysical, and the molecular basis for fluid clearance across the alveolar and distal airway epithelium.

1. Amiloride-sensitive sodium channels and ENaC

A number of studies using patch-clamp techniques, measurements of ²²Na uptake, and binding of amiloride analogs have demonstrated the presence of amiloride-sensitive sodium channels in alveolar type II cells. On the basis of the biophysical and pharmacological characterization, at least three types of amiloride-sensitive sodium channels have been described in the apical membrane of cultured adult alveolar type II cells (220). The most frequently observed in both inside-out and cell-attached configurations is a nonselective cation (NSC) channel with a conductance of ~21 pS, which was equally permeable to sodium and potassium ($P_{\text{Na}}/P_{\text{K}} \sim 1$), voltage-independent calcium-activated and completely blocked by 1 μM amiloride (105). A similar channel was described in fetal distal lung epithelial cells (213, 274). Also, others have recorded a channel with a conductance of 27 pS with a high relative permeability of sodium to potassium (7:1), completely blocked by 1 μM amiloride or its analog *N*-ethyl-*N*-isopropylamiloride (EIPA) (388). Finally, the presence of highly

selective cation (HSC) channels has been identified in adult alveolar type II cells cultured on permeable support, in air-liquid interface with aldosterone. These channels recorded on cell-attached configuration have a low conductance (5–7 pS) and were inhibited by a low concentration of amiloride (<0.1 μM) (163). These properties are similar to those described in fetal alveolar epithelial cells (370). Treatment of A549 cells with dexamethasone also shifted the conductance of the channel from 8 to 4 pS (194). A variety of other channels have been recorded in alveolar type II cells including ones regulated by G proteins (220).

Molecular identification of the proteins involved in amiloride-sensitive sodium influx has been achieved in the last few years. Three homologous subunits, entitled α -, β -, and γ -ENaC, correspond to the pore-forming subunits. The three subunits share in common a structure predicting two hydrophobic membrane-spanning regions, intracellular amino and carboxy termini, two transmembrane spanning domains, and a large extracellular loop with highly conserved cysteine residues (51, 298, 330). Not all investigators agree on the numbers of subunits that form functional channels; some propose that ENaC is a tetramer made of two α -, one β -, and one γ -subunit that assemble pseudosymmetrically around the channel pore to form a high amiloride affinity sodium-selective pore (111, 184), whereas others contend that ENaC is a much larger complex of nine (3 α -, 3 β -, 3 γ -) subunits (97, 329). An important role has been established for ENaC in the distal renal tubule and the collecting ducts and in the colon (15, 96, 223, 299). Some of the data suggest that individual ENaC subunits may have independent functions. For example, there are differences in the onset of expression of the three genes in the fetal lung (78, 348), and the relative levels of mRNA for the subunits can vary significantly in different regions of the lung, kidney, and colon (15, 87, 96, 104, 126, 223, 232). *Xenopus* oocyte studies predict a greater dependence of amiloride-sensitive sodium transport on α -ENaC than on β -ENaC or γ -ENaC (52, 237). Low levels of sodium transport (1% of maximal) could be detected in oocytes injected with α -ENaC alone but not in oocytes that expressed the β - or γ -subunits. Also, coinjection of β - or γ -subunits with α -ENaC increased sodium transport approximately five-fold, and maximal current required expression of all the three subunits. When coexpressed, the three subunits reproduce the expected properties of the HSC channel. In A549 lung epithelial cells, dexamethasone increases the expression of β - and γ -ENaC and markedly shifts the IC₅₀ for amiloride (194).

The critical role of ENaC in the absorption of salt and fluid by lung epithelia has been confirmed by the generation of knockout mice with inactivated subunits of ENaC. After inactivation of murine α -ENaC, deficient neonates develop respiratory distress syndrome and die within 40 h

of birth, primarily from failure to clear their lungs of fluid (154). In contrast, β - or γ -ENaC knockout mice were able to clear fluid from lungs at birth, although at a slower rate than in wild-type control, but these mice died from abnormal electrolyte reabsorption in the kidney resulting in fatal hyperkalemia (16, 233). These results suggest that small levels of activity provided by the remaining intact subunits ($\alpha\beta$ - or $\alpha\gamma$ -channels) are enough to maintain respiratory function but are not sufficient to absorb sodium in the kidney. More definitive answers regarding the role of ENaC should be forthcoming from inducible knockout ENaC mice, i.e., CRE-lox mice in which tetracycline can transiently suppress ENaC.

In humans, loss-of-function mutations in the α -, β -, and γ -ENaC subunits have been described in systemic pseudohypoaldosteronism (173, 316). Patients had no respiratory symptoms at birth but developed, within weeks or months after birth, respiratory illnesses characterized by chest congestion and cough caused by an excessive volume of liquid that resulted from sodium channel dysfunction in the airways.

In situ hybridization studies, as well as Northern blot analyses done in mice, rats, and humans, have identified the presence of mRNA for all three subunits of ENaC in alveolar epithelial type II cells both in vivo and in vitro with usually a greater expression of α - than β - and γ -subunits (64, 75, 78, 162, 220, 223, 232, 238, 264, 315, 345, 348). The relative amounts of ENaC mRNA could explain the difference in ENaC activity and amiloride sensitivity between tissues and species, although a direct relationship between mRNA and functional protein cannot be predicted. As previously mentioned, patch-clamp studies using freshly isolated or cultured rat alveolar type II cells have reported the presence of a nonselective cation (NSC) channel and have often failed to identify the classical HSC channel that would be expected from ENaC expression. Recent studies have provided evidence that HSC channels are expressed in the apical membrane of adult alveolar type II cells under specific culture conditions (163, 194). All studies showing the presence of a 4-pS channel were done after exposure of alveolar type II cells or A549 cells to aldosterone or dexamethasone. In these cells, the number of HSC channels decreased after treatment with antisense oligonucleotides to any of the three subunits of ENaC: α -antisense reduced all channel types while treatment with β - or γ -antisense increased NSC channels (163). Therefore, in alveolar type II cells, ENaC may be expressed in different subunit combinations that determine sodium channel characteristics and render them less sensitive to amiloride (35, 268). This observation raises the possibility that ENaC-like channels could be present in vivo. As described in section II A, new evidence has localized the three subunits of ENaC protein to alveolar type I cells also (38, 168).

In intact rat, sheep, mouse, or human lung, a high

concentration of amiloride (~ 1 mM) was required to inhibit 50–60% of fluid clearance. The discrepancy between in vivo and in vitro studies in which $1 \mu\text{M}$ is required to completely block epithelial sodium channels (105, 388) is not fully understood but could be due in part to the binding of amiloride to protein, a rapid degradation of amiloride after instillation (386), and/or escape from the airspaces (146, 266). Nevertheless, in vivo studies suggest that amiloride has poor affinity for sodium channels that are present in alveolar epithelium. In one rat study, EIPA, a pharmacological inhibitor primarily of low-affinity amiloride channels, inhibited an equal fraction of fluid clearance as amiloride (386). These results suggest that active sodium transport in vivo involves in part low-affinity amiloride-sensitive channels, i.e., NSC channels.

Several investigators have investigated how ENaC is regulated by transcriptional, translational, and posttranslational mechanisms (275, 276, 304). It is clear that understanding the regulation of ENaC processing, trafficking to, and stability at the cell surface is of fundamental importance (304).

2. Amiloride-insensitive sodium transport

Studies in several species including sheep, rabbit, guinea pig, rat, and human lungs demonstrated that there was a substantial fraction of fluid clearance that could not be inhibited by amiloride (30, 164, 258, 266, 308, 327, 386). In the rat and human lung, for example, 40–50% of the basal fluid clearance is amiloride insensitive while in mice $\sim 20\%$ of fluid clearance is insensitive to amiloride, although the latter result has been mainly identified in CD-1 mice (121). In CD57BL/6 mice, $\sim 40\%$ of the fluid clearance is amiloride insensitive (146).

There is some limited evidence that the amiloride-insensitive fraction of 8-Br-cGMP stimulated I_{sc} and sodium uptake in rat tracheal epithelia was inhibited by dichlorobenzamil or *L-cis*-diltiazem, both inhibitors of cyclic nucleotide-gated cation channels (CNG) (322). In subsequent studies, dichlorobenzamil inhibited a significant fraction of lung fluid absorption in sheep (170), a finding that supported the hypothesis that CNG channels could play a role in fluid absorption from the distal airspaces. The CNG channels were originally identified and cloned from vertebrate rod photoreceptors (109) and the olfactory neuroepithelium (246). Three different isoforms have been cloned (33). The CNG1 channel has a wide tissue distribution including expression in the lung; mRNA for CNG1 has been localized to distal lung epithelium and alveolar epithelial cells (82). In a recent rat study, *L-cis*-diltiazem inhibited a fraction of terbutaline or cGMP-stimulated fluid clearance that was not inhibited by amiloride, suggesting that part of the amiloride-insensitive fraction fluid clearance in the distal lung could be mediated through CNG channels, activated by cGMP (82, 261).

Also, there is recent data that inducible nitric oxide synthase (iNOS) (–/–) mice have normal rates of basal fluid clearance, but none of the basal fluid clearance is inhibited by amiloride (146).

In the intact rat, the amiloride-insensitive part of fluid clearance may be mediated in part through the Na⁺-glucose cotransporter (23). The amiloride-insensitive part of fluid clearance is abolished when sodium is replaced by choline in the instillate or when phlorizin, a specific inhibitor of the Na⁺-glucose cotransport, is added in the instillate. However, in one study, only sodium channel inhibition slowed lung water clearance in neonatal animals (265). On balance, the importance of Na⁺-glucose cotransport for fluid clearance from the airspaces remains unproven. Surprisingly, although SGLT1 mRNA is detected in mouse lungs, suggesting the presence of Na⁺-glucose transporter, luminal glucose has no influence on fluid clearance (155). In vitro there is no evidence that the Na⁺-glucose cotransporter plays a major role in transepithelial sodium transport (174). Also, the role of other cotransporters such as Na-amino acid and Na-phosphate that are specifically expressed in the apical membranes of human, mouse, and rat type II alveolar epithelial cells, remains to be defined (46, 69, 354). Some recent work indicates that SGLT1 can function as a cotransporter of water (89). Finally, regulation of vectorial fluid transport could be indirectly influenced by Na⁺/H⁺ exchange, especially in the setting of chronic metabolic acidosis (169).

3. Na⁺-K⁺-ATPase and lung epithelial fluid transport

The Na⁺-K⁺-ATPase enzyme is a ubiquitous plasma membrane ion-transporting ATPase that maintains electrochemical sodium and potassium gradients across the plasma membrane by pumping sodium out of the cell and potassium into the cell against their respective concentration gradients, fueled by hydrolysis of ATP. The electrochemical gradient generated by extrusion of sodium from and uptake of potassium into the cell is essential for the secondary active movement of sodium ions. In most of the epithelia, including the alveolar epithelium, the Na⁺-K⁺-ATPase is confined to the basolateral domain of the cells; this polarized distribution is essential for the vectorial transport of sodium followed isosmotically by that of water while also controlling in its housekeeping mode cell volume and composition. In the alveolar epithelium, the Na⁺-K⁺-ATPase protein is detected primarily in type II cells with a weak expression in type I cells consistent perhaps with a predominant role of type II cells in resorbing alveolar fluid. The Na⁺-K⁺-ATPase is a heterodimeric transmembrane protein composed of one α - and one β -subunit in a 1:1 ratio, although there are multiple isoforms of both subunits usually expressed in a tissue-specific pattern. The α_1 -isoform catalyzes the movement of sodium and potassium, is phosphorylated

by ATP, and binds the specific inhibitor ouabain (326). The α_1 -isoform is the predominant form in alveolar type II cells, but cultured type I cells may express α_2 (303). The β -subunit has three isoforms and targets the assembled molecule to the plasma membrane. It is believed that a heterodimeric form composed of the α_1 - and β_1 -subunits is the predominant Na⁺-K⁺-ATPase isoform in alveolar epithelial cells. It has been convincingly demonstrated that both α - and β -isoforms are required for functional insertion of Na⁺-K⁺-ATPase in the membrane (235). In the lung, adenoviral gene transfer of the β_1 , but not α_1 , subunit in the alveolar epithelium increases Na⁺-K⁺-ATPase expression and fluid clearance in the adult rat (102) and in confluent monolayers of fetal alveolar type II cells (351).

Removing edema fluid from the airspaces is accomplished by the polar distribution of Na⁺-K⁺-ATPases and channels on opposite poles of epithelial cells (129). Na⁺-K⁺-ATPase activity in the plasma membrane is regulated acutely by covalent or allosteric modification and/or acute trafficking of Na⁺-K⁺-ATPases between the plasma membrane and intracellular endosomal pools, and chronically by regulation of the abundance of Na⁺-K⁺-ATPases in the membrane by changes in Na⁺-K⁺-ATPase synthesis and/or degradation rates (31).

4. Aquaporins and lung epithelial fluid transport

In 1993, the first transcellular water channel, aquaporin-1 (AQP1), was cloned (3). Since that time, the list of known aquaporins has grown rapidly to approximately 10 mammalian water channels (182, 366). Four aquaporins have been localized to the lung with nonoverlapping distribution (365). In rat and mouse lung, AQP1 is present in both apical and basolateral membrane of microvascular endothelial cells and in fibroblasts, while AQP3, -4, and -5 polarize to the apical or basolateral membrane at different locations in the respiratory epithelium (179). Recent work in the human lung demonstrates that there are many similarities between the distribution of aquaporins in the rodent and human lung, but there are some additional areas of expression in the human respiratory tract (185). In addition to its expression on the apical surface of alveolar type I cells, AQP5 is located in the apical membrane of airway epithelium as well as in the superficial epithelium of the nasopharynx (185). AQP3 was expressed in the apical membrane of columnar cells as well as in basal cells. There seems to be some AQP3 in type II cells and AQP4 in alveolar epithelial cells (185).

The identification of aquaporins in the lung naturally suggested the hypothesis that aquaporins might contribute to the developmental regulation of lung fluid balance in the formation or reabsorption of fetal lung fluid, the pathogenesis or resolution of pulmonary edema in the mature lung, or in the humidification of the airways.

These possibilities have been studied in several lung preparations over the last few years.

The initial studies measured the water permeabilities of the endothelial and epithelial barriers in the lung in conjunction with studies that determined the cellular expression of aquaporins in the mature lung. Water permeability across a simple barrier separating two compartments is characterized by an osmotic water permeability coefficient (P_f) defined as the volume flux (J_v) induced by an osmotic gradient: $P_f + J_v/(SV_wC)$, where S is the barrier surface area, V_w is the partial molar volume of water, and C is the osmotic gradient. High P_f (generally >0.0005 – 0.01 cm/s) suggests a facilitated water pathway involving molecular water channels (53). More detailed description of the biophysics of measuring water permeability is available in several sources (2, 3, 54, 362, 364).

In the lung, osmotic water permeability was found to be high in the intact sheep lung across the alveolar-capillary barrier, 0.02 cm/s (113). Additional studies were done in mice with a novel fluorescence method that provided similar data, namely, a high water permeability across the alveolar-capillary barriers, 0.017 cm/s. Aquaporin localization studies had already demonstrated that AQP1 was distributed primarily throughout the lung capillary endothelium, whereas AQP5 was localized primarily to the apical surface of alveolar epithelial type I cells (113, 181, 252, 321). Interestingly, measurements of osmotic water permeability of isolated alveolar epithelial I cells indicated that these cells have a high water permeability, 0.07 cm/s (84). A relatively high water permeability was also measured across isolated micropperfused distal airways from guinea pigs, 0.005 cm/s (112). All of these studies provided indirect evidence for a possible functional role of aquaporins in lung fluid balance in the lung as potential rate-limiting channels for isosmolar fluid clearance from the distal airspaces of the lung. However, because there are no reliable, nontoxic inhibitors of water transport, it was necessary to generate specific aquaporin knockout mice to test the importance of specific aquaporins in isosmolar vectorial fluid transport under physiological conditions across the distal pulmonary epithelium.

AQP1 knockout mice had a modest decrease in lung interstitial fluid accumulation in the presence of a hydrostatic stress, but there was no effect on isosmolar fluid clearance across the distal pulmonary epithelium (9). The potential contribution of AQP5 to isosmolar fluid clearance in knockout mice was tested under both basal and stimulated conditions. Even when isosmolar fluid transport in the mouse was increased to a maximum level of 30% in 15 min, the absence of this apical type I cell membrane water channel did not slow the rate of fluid clearance (9, 331, 364, 365, 367). It is unlikely that the discovery of other water channels would alter the results of these studies because deletion of AQP1 or AQP5 each produced a 10-fold decrease in osmotically driven water

transport between the airspace and capillary compartments.

Although aquaporin expression is strongly upregulated near the time of birth (180, 355, 385), AQP1, AQP4, and AQP5 knockout mice have the same ability to clear lung fluid in the perinatal period as wild-type controls (331). One study suggested that aquaporins might play some role in lung fluid balance after pneumonia because there was decreased expression of AQP4 and AQP5 in the mouse lung after acute viral pneumonia (353). However, the decreased expression may simply have reflected loss of functional epithelial cells from the pneumonia. Furthermore, knockout of AQP1, -4, and -5 had no effect on the formation or resolution of experimental pulmonary edema from hyperoxia, thiourea, or acid instillation (331). Finally, recent data indicate that the elimination of AQP3 does not delay the normal humidification process in the airways of mice (332).

The insensitivity of fluid clearance to aquaporin deletion is probably the consequence of the substantially lower rate of active fluid absorption per unit surface area in the lung compared, for example, with the proximal tubule of the kidney or the salivary gland, organs in which aquaporin deletion does alter fluid absorption or secretion (362, 363). Slower rates of active fluid transport probably do not require very high cell membrane water permeabilities.

Although a major role of AQP5 in physiological transalveolar epithelial water movement appears unlikely, aquaporins may have effects on other cell functions, especially volume regulation, particularly in the highly specialized type I cell (152, 199), in alveolar fluid homeostasis (206), or in the regulation of fluid secretion from mucus secreting cells (207).

III. REGULATION OF SALT AND WATER TRANSPORT

This section considers how vectorial fluid transport across the distal pulmonary epithelium can be upregulated by either catecholamine-dependent or catecholamine-independent mechanisms. The potential relevance of these mechanisms under pathological conditions is evaluated in section v.

A. Catecholamine-Dependent Regulation

Until recently, most studies focused primarily on the active transport of sodium as the primary determinant for regulating catecholamine-dependent transport across distal pulmonary epithelium. New evidence indicates that cAMP-stimulated uptake of chloride may be an important mechanism for regulating fluid clearance across distal lung epithelium (103, 165, 166, 267). Therefore, this sec-

tion is divided into the effects of catecholamines on fluid transport based primarily on studies that examined the effects of sodium transport inhibitors. Section IIIA2 considers the new evidence regarding a potential role for chloride transport in cAMP-mediated transport.

1. Upregulation of fluid transport by sodium-dependent mechanisms

Studies in newborn animals indicate that endogenous release of catecholamines, particularly epinephrine, may stimulate reabsorption of fetal lung fluid from the airspaces of the lung (45, 110, 259, 260, 371). In most adult mammal species, stimulation of β_2 -adrenergic receptors either by salmeterol, terbutaline, or epinephrine increases fluid clearance (30, 59, 74, 164, 210, 290, 307). This stimulatory effect occurs rapidly after intravenous administration of epinephrine or instillation of terbutaline in alveolar space and is completely prevented by either a nonspecific β_2 -receptor antagonist, propranolol, or in rats by a specific β_2 -antagonist (210, 290). In some species, such as the guinea pig, there is also evidence that stimulation of the β_1 -receptor will augment fluid clearance (258).

The increased fluid clearance by β_2 agonists can be prevented by amiloride, indicating that the stimulation was related to an increased transepithelial sodium transport (164, 258, 290). In anesthetized ventilated sheep, terbutaline-induced stimulation of fluid clearance was associated with an increase in lung lymph flow, a finding that reflected removal of some of the alveolar fluid volume to the interstitium of the lung (30). Although terbutaline increased pulmonary blood flow, this effect was not important, since control studies with nitroprusside, an agent that increased pulmonary blood flow, did not increase fluid clearance. Other studies have demonstrated that β -adrenergic agonists increased fluid clearance in rat (164), dog (28, 191), guinea pig (258), mouse (121, 124, 155), as well as human lung (307, 308). The presence of β_1 - and β_2 -receptors on alveolar type II cells has been demonstrated in vivo by autoradiographic and immunohistochemistry techniques (19, 98, 201). In vitro, studies in cultured alveolar type II cells indicate that β -adrenergic agonists added at the apical or the basolateral sides increased I_{sc} that was abolished by amiloride (132, 217). In addition, terbutaline stimulated net apical-to-basolateral flux of ^{22}Na , and the magnitude of this stimulation was similar to the steady-state increase in I_{sc} (60). Experiments in isolated lungs (130, 313) and in vitro studies provided further evidence that cAMP is a second messenger for the β -adrenergic effects, whereas activation of protein kinase C does not appear to be involved (26, 29). Interestingly, β -adrenergic agonist therapy does not increase fluid clearance in rabbits and hamsters (129, 327). The explanation for this lack of effect is unclear, partic-

ularly since there are β -receptors in rabbit type II cells that stimulate surfactant secretion (234). One study in rabbits suggested that cAMP agonists might stimulate chloride secretion in rabbits, thus perhaps offsetting any increase in vectorial fluid clearance from the distal airspaces (254).

On the basis of studies of the resolution of alveolar edema in humans, it has been difficult to quantify the effect of catecholamines on the rate of fluid clearance (230). However, studies of fluid clearance in the isolated human lung have demonstrated that β -adrenergic agonist therapy increases fluid clearance, and the increased fluid clearance can be inhibited with propranolol or amiloride (Fig. 4) (307, 308). Subsequent studies showed that long-acting lipid-soluble β -agonists are more potent than hydrophilic β -agonists in the ex vivo human lung (307). The magnitude of the effect is similar to that observed in other species, with a β -agonist-dependent doubling of fluid clearance over baseline levels. These data are particularly important because aerosolized β -agonist treatment in some patients with pulmonary edema might accelerate the resolution of alveolar edema (see sect. v).

What has been learned about the basic mechanisms that mediate the catecholamine-dependent upregulation of sodium transport in the lung? On the basis of in vitro studies, it was proposed that an increase in intracellular cAMP resulted in increased sodium transport across alveolar type II cells by an independent upregulation of the apical sodium conductive pathways and the basolateral $\text{Na}^+\text{-K}^+\text{-ATPase}$. Proposed mechanisms for upregulation of sodium transport proteins by cAMP include augmented sodium channel open probability (75, 105, 192, 212, 216, 340), increases in $\text{Na}^+\text{-K}^+\text{-ATPase}$ α -subunit phosphorylation, as well as delivery of more ENaC channels to the apical membrane and more $\text{Na}^+\text{-K}^+\text{-ATPases}$ to the basolateral cell membrane (34, 48, 61, 106, 165, 328). Patch-clamp studies done on alveolar type II cells in the attached mode indicated that β -agonists increased the open-state probability and the mean open time of the moderately selective amiloride-sensitive sodium channel without affecting single-channel conductance. This effect was completely blocked by propranolol. The observed increase in open probability by protein kinase A (PKA) was perhaps mediated by the direct phosphorylation of channel proteins. Although one study indicated that rat ENaC can be phosphorylated by PKA (324), the functional relevance of phosphorylation is not yet known, and PKA does not activate amiloride-sensitive current in *Xenopus* oocytes injected with ENaC mRNAs (6), suggesting that other mechanisms are involved. Sodium channels, like other ion channels and transporters, are associated with cytoskeletal proteins such as actin, ankyrin, fodrin, or spectrin, which can serve as signaling molecules. Reconstitution of ENaC into lipid bilayers resulted in a channel that is activated by PKA and ATP in the presence but not




FIG. 4. Alveolar fluid clearance (*y*-axis) plotted under control conditions, β -agonist stimulation, and after amiloride inhibition in the human lung (A), the rat lung (B), and the mouse lung (C). Amiloride sensitivity is different depending on mouse strain (146). [Data from Fukuda et al. (121), Jay et al. (164), and Sakuma et al. (308).]

the absence of short actin filaments (25). Thus the increase in open probability may have been the result of increased phosphorylation of short actin filaments or other cytoskeleton proteins. Finally terbutaline, a β_2 -adrenergic agonist, may promote insertion of new channel protein from a cytoplasmic pool to the apical membrane. In fetal distal lung cells, brefeldin A, an inhibitor of pro-

tein trafficking, prevented terbutaline-induced increase in open probability of Ca^{2+} -activated nonselective channels (160).

One investigator reported that stimulation of thyroid epithelial cells transfected with α -, β -, and γ -ENaC with cAMP translocated ENaC from the cytoplasm to the cell surface, and that effect was dependent on a sequence (PPPXY) in the COOH terminus of each subunit (328). Existing evidence also suggests that agents that increase cAMP may increase sodium influx through an increase in apical chloride conductance without affecting apical membrane sodium conductance (165, 267).

One recent study using patch clamping of rat alveolar type II cells demonstrated that treatment with a β_2 -agonist, terbutaline, increased the number of highly selective sodium channels, an effect that was mediated by PKA (61). Also, in the same study, terbutaline increased both intracellular calcium levels and the open probability of NSC channels, an effect that was blocked by a calcium chelator (61). The authors concluded that β -adrenergic stimulation increase intracellular cAMP and the activation of PKA, and the PKA promotes an increase in the number of highly selective sodium channels and an increase in intracellular calcium. The increase in intracellular calcium increases the open probability of NSC channels. Interestingly, other investigators recently found that both extracellular and intracellular calcium may act as second messengers for β -adrenergic stimulation of fluid clearance from the distal airspaces of near-term fetal guinea pigs (260). Other investigators have also demonstrated that calcium channel activation and cell volume changes alter the rate of sodium transport via NSC channels (211–213).

β -Adrenergic stimulation increases extrusion of sodium at the basolateral side by increasing Na^+ - K^+ -ATPase activity. In alveolar type II cells, isoproterenol increased the maximal velocity of the enzyme independently of sodium permeability (340). The stimulation of Na^+ - K^+ -ATPase activity was associated with an increase in ouabain binding and expression of the α_1 -subunit at the membrane (32). This effect occurred rapidly (<15 min) and was independent of protein kinase phosphorylation and mediated by subunit recruitment from late endosomes into the plasma membrane. There is some evidence that long-term exposure of alveolar type II cells to β -agonists increased Na^+ - K^+ -ATPase and sodium channel activity, at least in part through a transcriptional effect, since cAMP increased transiently the α_1 -subunit of Na^+ - K^+ -ATPase (239) and the three subunits of ENaC (77). Increased activity of PKA could lead to phosphorylation of the cAMP-responsive element protein, which could bind to the cAMP responsive element (CRE) site on the gene promoter. A CRE site has been demonstrated on the promoter region of α_1 -subunit of Na^+ - K^+ -ATPase (4) and in the 5'-flanking region of γ -ENaC (350). However, other

investigators have failed to demonstrate change in mRNA levels of $\text{Na}^+\text{-K}^+\text{-ATPase}$ after long exposure to isoproterenol, whereas they found a posttranscriptional regulation via mitogen-activated protein kinase/ERK and rapamycin-sensitive pathways (280).

2. Potential role of CFTR in cAMP-mediated upregulated fluid transport

While most experimental studies have attributed a primary role for active sodium transport in the vectorial transport of salt and water from the apical to the basal surface of the alveolar epithelium of the lung, the potential role of chloride, especially in mediating the cAMP-mediated upregulation of fluid clearance across distal lung epithelium, has been the subject of a few recent studies. One older study of cultured alveolar epithelial cells concluded that vectorial transport of chloride across alveolar epithelium occurs by a paracellular route under basal conditions and perhaps by a transcellular route in the presence of cAMP stimulation (176). Another study of cultured alveolar epithelial type II cells suggested that cAMP-mediated apical uptake of sodium might depend of an initial uptake of chloride (165). A more recent study of cultured alveolar type II cells under apical air interface conditions reported that β -adrenergic agonists produced acute activation of apical chloride channels with enhanced sodium absorption (166). However, the results of these studies have been considered to be inconclusive by some investigators (193, 378, 379), partly because the data depend on cultured cells of an uncertain phenotype. Furthermore, studies of isolated alveolar epithelial type II cells do not address the possibility that vectorial fluid transport may be mediated by several different epithelial cells including alveolar epithelial type I cells as well as distal airway epithelial cells.

It should also be appreciated that the pathways for chloride secretion and absorption in the fetal and the neonatal lung are not well understood. Recent and older studies have demonstrated a role for chloride secretion and absorption in the fetal and perinatal lung (57, 117, 127, 273) as well as in distal fetal rat lung epithelial cells (70, 195), but the molecular basis for chloride transport is not well understood (262).

To define the role of chloride transport in the active transport of salt and water across the distal pulmonary epithelium of the lung, one group has used in vivo lung studies to define the mechanisms and pathways that regulate chloride transport during the absorption of fluid from the distal airspaces of the lung. This approach may be important because studies in several species, as already discussed, have indicated that distal airway epithelia are capable of ion transport (5, 13, 14), and both ENaC and CFTR are expressed in alveolar and distal airway epithelia (104, 123, 185, 284, 299).

Both inhibition and ion substitution studies demonstrated that chloride transport was necessary for basal fluid clearance. The potential role of CFTR under basal and cAMP-stimulated conditions was tested using intact lung studies in which CFTR was not functional because of failure in trafficking of CFTR to the cell membrane, the most common human mutation in cystic fibrosis (ΔF508 mice). The results supported the hypothesis that CFTR was essential for cAMP-mediated upregulation of isosmolar fluid clearance from the distal airspaces of the lung because fluid clearance could not be increased in the ΔF508 mice with either β -agonists or with forskolin, unlike the wild-type control mice (103). Additional studies using pharmacological inhibition of CFTR in both the mouse and human lung with glibenclamide supported the same conclusion, namely, that chloride uptake and CFTR-like transport seemed to be required for cAMP-stimulated fluid clearance from the distal airspaces of the lung (103). Glibenclamide can also inhibit potassium channels so the inhibitory effects are not specific for CFTR, but the ΔF508 mouse studies have provided more direct evidence. Although the absence of CFTR in the upper airways results in enhanced sodium absorption (336), the data in these studies provide evidence that the absence of CFTR prevents cAMP-upregulated fluid clearance from the distal airspaces of the lung, a finding that is similar to work on the importance of CFTR in mediating cAMP-stimulated sodium absorption in human sweat ducts (297). Because CFTR is distributed throughout the distal pulmonary epithelium in distal airway epithelium as well as at the alveolar level in the human lung (95), the data also suggest that the cAMP-mediated upregulated reabsorption of pulmonary edema fluid may occur across distal airway epithelium as well as at the level of the alveolar epithelium. Finally, additional studies indicated that the lack of CFTR results in a greater accumulation of pulmonary edema in the presence of a hydrostatic stress, thus demonstrating the potential physiological importance of CFTR in upregulating fluid transport from the distal airspaces of the lung (103).

The new data on a role for CFTR in cAMP-upregulated fluid clearance from the distal airspaces raise further interest in determining how CFTR and ENaC interact and contribute to net fluid clearance under isosmolar conditions. The relative conductances for chloride and sodium are difficult to measure in the in vivo lung epithelium but conceivably the debate about the role of either ion in limiting isosmolar fluid clearance is somewhat misleading. Under open-circuit conditions, the net transfer of chloride and sodium across the distal lung epithelium must be equal, i.e., there cannot be a significant net charge accumulation. It is possible that with cAMP stimulation the conductances for both chloride and sodium increase in parallel. This hypothesis would be in accord with recent data on isolated distal lung epithelial cells in which

CFTR protein was immunolocalized to these cells and forskolin-stimulated I_{sc} was inhibited by glibenclamide (195). Other investigators did not find evidence to support the hypothesis that cAMP-stimulated chloride conductance directly controlled sodium uptake in rat fetal distal lung epithelial cells (70), and these investigators concluded that chloride and sodium transport must be linked by some mechanism.

There are several interesting unanswered questions. What is the molecular basis for chloride transport under basal conditions across the distal lung epithelium, since it is clear that CFTR is not required for basal fluid clearance or for clearance of fetal lung fluid at birth (103)? Second, if CFTR is required for cAMP-mediated transport, does this mechanism apply to all levels of the distal lung epithelium including the alveolar epithelial type I and type II cells as well as the distal airway epithelia? Also, does upregulation of fluid clearance by cAMP-independent pathways require functional CFTR (see sect. III)?

3. Regulation of β_2 -receptor expression

There is new information regarding the effect of up-regulating the expression of the β_2 -receptor (β_2 AR) in alveolar type II cells using the SP-C promoter in transgenic mice (236). β_2 -AR expression was 4.8-fold greater than that of the nontransgenic mice (939 ± 113 vs. 194 ± 18 fmol/mg protein, $P < 0.001$). Basal fluid clearance in the transgenic mice was 40% greater than that of untreated nontransgenic mice (15 ± 1.4 vs. $10.9 \pm 0.6\%$, $P < 0.005$) and approached that of nontransgenic mice treated with 10^{-5} M of the potent β -agonist formoterol ($19.8 \pm 2.2\%$). Adrenalectomy had no effect on nontransgenic mice ($11.5 \pm 1.0\%$) but decreased basal fluid clearance in the transgenic mice to $9.7 \pm 0.5\%$. These findings demonstrated that overexpression of the β_2 -AR can be an effective means to increase vectorial fluid transport in the absence of exogenous agonists.

Other investigators have used an adenoviral vector to deliver the β_2 -AR to the rat lung, which results in a comparable increase in fluid clearance that was also inhibited by adrenalectomy or by β -antagonists (99). In these studies, the β_2 -AR may be expressed in several cells in the distal airspaces of the lung. There was evidence that the β_2 -AR overexpression was associated with increased levels of ENaC (α -subunit) and Na^+ - K^+ -ATPase in apical and basolateral cell membrane fractions isolated from peripheral lungs (88).

Some recent studies suggest that prolonged exposure to exogenous catecholamines may impair the ability of the alveolar epithelium to remove alveolar edema fluid and that this impairment was associated with a reduction in epithelial β -receptor number (244). It is important to bear in mind that although a mild impairment was evident at clinically used doses, infused β -agonist as well as in-

stilled β -agonist stimulated fluid clearance above baseline levels.

B. Catecholamine-Independent Regulation

In the last few years several interesting catecholamine-independent mechanisms have been identified that can upregulate fluid transport across the distal airspaces of the lung as well as in cultured alveolar type II cells. Hormonal factors, such as glucocorticoids, can upregulate transport by transcriptional mechanisms while thyroid hormone may work by a posttranslational mechanism (197). Some growth factors can work by either a transcriptional or direct membrane effect, or by enhancing the number of alveolar type II cells. There is also evidence that a proinflammatory cytokine, tumor necrosis factor- α , can rapidly upregulate sodium uptake and fluid transport by a novel mechanism. Finally, serine proteases can regulate the activity of ENaC and potentially increase fluid clearance across the distal airway epithelium.

1. Glucocorticoids

Glucocorticoid hormones regulate sodium uptake and fluid transport in both adult and fetal lungs (20, 158, 299, 348). In adult guinea pig lungs, endogenous plasma levels of cortisol are important for maintaining normal lung fluid balance and distal airspace fluid clearance. Cortisol modulates the amiloride-sensitive sodium transport pathways through regulation of de novo synthesis of sodium channel proteins (257). Dexamethasone, a more potent glucocorticoid agonist than cortisol, regulates transepithelial sodium transport in vivo by increasing lung fluid clearance in anesthetized, ventilated rats (116). Based on in vitro studies, there was a functional increase in I_{sc} across monolayers of alveolar type II cells after dexamethasone treatment (77). In A549 cell studies, dexamethasone increased the whole cell amiloride-sensitive current and decreased the IC_{50} for amiloride and increased the channel conductance (194). Dexamethasone increased Na^+ - K^+ -ATPase and sodium channel activity and expression by acting at both transcriptional and post-transcriptional levels. The molecular studies have reported that glucocorticoids directly stimulated the transcription of sodium transport proteins. In vivo experiments have demonstrated a differential regulation of ENaC subunits by corticosteroids: α - but neither β - nor γ -ENaC mRNA levels increase within 8 h of treatment. Dexamethasone increases mRNA of the three subunits of ENaC in studies using primary cultures of fetal and adult lung cells (77, 299, 348), but only β - and γ -ENaC mRNA in the human alveolar epithelial cell line A549 (194). This latter observation is in contrast to the recent evaluation of the human α -ENaC promoter showing the presence of glucocorticoid responsive element in the 5'-flanking re-

gion in human (315) and in rat (77, 276). Interestingly, one study indicated that the differential regulation of mRNA of ENaC subunits correlated with similar changes in the quantity of protein and with a change in selectivity of the sodium channel from essentially nonselective to highly selective (194). Also, glucocorticoids differentially regulate $\text{Na}^+\text{-K}^+\text{-ATPase}$ subunit expression in fetal and adult rat lungs (20, 77, 158). In adult alveolar type II cells, dexamethasone increased $\beta_1\text{-}$ but not $\alpha_1\text{-Na}^+\text{-K}^+\text{-ATPase}$ mRNAs (20, 77), while both $\alpha_1\text{-}$ and $\beta_1\text{-Na}^+\text{-K}^+\text{-ATPase}$ proteins were increased. Because $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity increased only when the proteins of $\alpha_1\text{-}$ and $\beta_1\text{-Na}^+\text{-K}^+\text{-ATPase}$ were upregulated, it was suggested that post-translational events may be involved in the regulation of $\text{Na}^+\text{-K}^+\text{-ATPase}$ by glucocorticoids (20).

2. Mineralocorticoids

The lung is a potential target organ for aldosterone because it expresses mineralocorticoid receptors and the enzyme 11- β -hydroxysteroid dehydrogenase that converts the glucocorticoid corticosterone into receptor-inactive 11-dehydro-corticosterone, thereby allowing preferential access of aldosterone to mineralocorticoid receptors (186, 339, 341). There is excellent evidence that aldosterone plays a major role in the regulation of sodium transport in epithelial tissues, which has led to the hypothesis that aldosterone might also regulate salt transport in the lung. In vitro, overnight incubation of alveolar type II cells with aldosterone concentrations ranging from 10^{-7} to 10^{-4} M did not increase the I_{sc} (72). On the other hand, in isolated alveolar type II cells, aldosterone showed increased mRNA expression of $\beta_1\text{-Na}^+\text{-K}^+\text{-ATPase}$ and increased amount of $\alpha_1\text{-}$ and $\beta_1\text{-Na}^+\text{-K}^+\text{-ATPase}$ proteins, suggesting that aldosterone regulates $\beta_1\text{-}$ subunit at the transcriptional/translational level while the $\alpha_1\text{-}$ subunit is probably recruited from intracellular pool to the basolateral membrane (271). This effect was associated with a fourfold increase in the $\text{Na}^+\text{-K}^+\text{-ATPase}$ hydrolytic activity and ouabain-sensitive ^{86}Rb uptake (271). The in vivo relevance of these observations was substantiated by the finding that 24 h after aerosolized delivery of aldosterone there was a 50% increase in fluid clearance in rats (271). Similarly, rats treated with a low-sodium diet that developed a hyperaldosteronism have an increase in fluid clearance related to a stimulation of amiloride-sensitive component (339). The effect of aldosterone on ENaC mRNA expression has not been studied. However, a new study found that alveolar type II cells cultured in the presence of aldosterone (10 μM) had an increase in the frequency of highly selective channels than cells cultured in the absence of aldosterone (163). These results suggest that aldosterone may regulate ENaC subunits as demonstrated in other epithelial cells (52, 87, 203–205, 231, 277, 368) and indicate that further studies will be useful to determine

the molecular and cellular effect of aldosterone on ENaC in alveolar epithelial cells.

Also, there are data that estrogen may modulate mRNA levels and amiloride-sensitive I_{sc} in adult rat lung and cultured alveolar type II cells (342).

3. Growth factors

Several studies have reported that growth factors can upregulate sodium uptake and net fluid transport across the distal airspaces of the lung by several different mechanisms. This section considers the effects of three growth factors that have been studied.

A) EPIDERMAL GROWTH FACTOR. In vitro studies demonstrated that incubations of alveolar epithelial type II cells with epidermal growth factor (EGF; 20 ng/ml) for 4–5 days resulted in a significant increase in both $\alpha_1\text{-}$ and $\beta_1\text{-Na}^+\text{-K}^+\text{-ATPase}$ mRNA and protein concentrations with no changes in ENaC subunit mRNA. These transcriptional effects were paralleled by an increase in benzamil-inhibitable I_{sc} (79). Subsequently, the same research group reported that alveolar type II cells cultured with EGF induced a decrease in mRNA for all three subunits of ENaC, and patch-clamp studies of these cells showed only 25-pS channels that were amiloride sensitive and relatively nonselective for cations ($P_{\text{Na}}/P_{\text{K}} = 1.0:0.48$). Channel density was increased with EGF. Thus the authors concluded that EGF induced the formation of NSC channels in cultured type II cells (172). Also, when EGF was aerosolized to rats, there was a marked increase in fluid clearance 24 h later. Because EGF has mitogenic properties, it is possible that some of the EGF effect in vivo could have occurred by stimulating type II cell proliferation, although there was no evidence for a mitogenic effect in the rat studies (344).

B) TRANSFORMING GROWTH FACTOR- α . EGF has been reported to increase sodium uptake and transport in isolated alveolar epithelial type II cells after direct in vitro exposure (37) as well as after prior in vivo exposure (161). EGF and transforming growth factor- α (TGF- α) have also been shown to stimulate intracellular accumulation of cAMP in several epithelial cell lines, heart cells, parotid gland cells, and embryonic cells (12, 245). Epithelial sodium transport can be regulated by tyrosine kinase in kidney epithelial cells (222), which is stimulated by both EGF and TGF- α .

Interestingly, TGF- α can upregulate fluid clearance rapidly, apparently by direct membrane effects. One study in rats demonstrated a marked increase in fluid clearance within 1 h of instillation into the living rat lung (118). There was no increase in cAMP levels, and the effect was not inhibited by β -adrenergic blockade. However, genistein prevented the effect, suggesting that the effect might be mediated by tyrosine kinase activity. This observation may have relevance to patients with pulmonary

edema, since measurable quantities of TGF- α have been reported in patients with acute lung injury (62).

c) **KERATINOCYTE GROWTH FACTOR.** It is well known that keratinocyte growth factor (KGF) is a potent mitogen for alveolar epithelial type II cells. Administration of KGF (5 mg/kg) into the distal airspaces of the rat lung upregulated fluid clearance by 66% over baseline levels (373), an effect that was sustained for 120 h. Also, the magnitude of the increase in fluid clearance was comparable to the effect previously reported in rats with short-term upregulation from β -adrenergic agonists. There was a good correlation between the increase in the number of alveolar type II cells and the effect of KGF on upregulating fluid clearance (373). Although there was a modest increase in the expression of α -ENaC 72 h after administration of KGF, the predominant effect seemed to be related to the type II cell hyperplasia. This sustained upregulation of fluid clearance suggested potential therapeutic value of an epithelial-specific mitogen-like KGF. Furthermore, the addition of a β_2 -adrenergic agonist further upregulated fluid clearance in rats so that the combination of KGF treatment plus terbutaline resulted in a net clearance of 50% of the instilled fluid in 1 h in rats compared with control rats with fluid clearance rates of 23%/h. Other investigators have shown that KGF can enhance sodium and fluid transport in normal and injured rat lungs (137, 140). In vitro studies of the effect of KGF in cultured alveolar epithelial cells indicated that KGF might also work by enhancing the expression of α_1 -Na⁺-K⁺-ATPase subunits (39), thereby enhancing sodium transport independent of an effect on the number of alveolar type II cells.

4. Dopamine

Dopamine is a vasoactive agonist that was previously described to impair sodium reabsorption in renal tubule. In the rat lung, the effect of dopamine is opposite to that in the renal tubular epithelium as dopamine increases fluid clearance in isolated perfused lung during different conditions (7, 17, 312). This increase was dependent on the sodium transport and was mediated by D₁ receptor present in alveolar type II cells and not via the β -adrenergic receptor pathway. In alveolar type II cells, dopamine increases the expression of the α_1 -subunit of Na⁺-K⁺-ATPase in the basolateral membrane (18). An inhibitor of cell microtubular transport, colchicine, inhibits both the dopamine-induced stimulation of fluid clearance and dopamine-induced increase of α_1 -Na⁺-K⁺-ATPase expression, which suggests that dopamine works by recruiting Na⁺-K⁺-ATPase subunits from intracellular pools to the basolateral membrane of epithelial cells. This effect is mediated by D₁ receptors through the activation of protein phosphatase 2A (196) as well as recently suggested to be mediated via mitogen-activated protein kinases (136).

5. Serine proteases

The activity of ENaC can be upregulated by serine protease activity (356). In one study, exposure of a kidney epithelial cells line (A6) to the protease inhibitor aprotinin reduced epithelial sodium transport, and sodium channel activity could be restored by exposure to a nonspecific protease trypsin. Expression of a serine protease in *Xenopus* oocytes along with ENaC increases the activity of the sodium channel by two- to threefold. This protease (CAP1) is expressed in lung as well as other organs. Thus it is possible that under normal and pathological conditions, ENaC activity could be regulated by the activity of proteases expressed at the surface of the same cell and thus may provide a mechanism for autocrine regulation of ENaC in the lung. Preliminary data in recent lung studies support this hypothesis (360).

IV. MECHANISMS THAT IMPAIR VECTORIAL FLUID TRANSPORT

Several mechanisms have been identified that can impair fluid transport from the distal airspaces of the lung. This section considers one hormonal factor, atrial natriuretic factor (ANF), and three other conditions that have relevance to human disease: hypoxia, anesthetics, and reactive oxygen and nitrogen species

A. Atrial Natriuretic Peptide

While several hormones are capable of upregulating epithelial fluid transport in the lung, as discussed in section III, atrial natriuretic peptide (ANP) is an example of a hormone that can downregulate fluid transport in the lung. ANP plays a major role in volume and electrolyte homeostasis, through potent biological effect including natriuresis, diuresis, and vasorelaxation. In addition to being a target organ for ANP from atrial origin, the lung is also a site of synthesis and release of bioactive ANP (142). The lung has the highest tissue concentration of ANP binding sites (295), and alveolar type II cells have been shown to express both guanylate cyclase (GC) receptor subtypes with a functional predominance of GC-A over the GC-B receptors but do not possess clearance receptors (49, 349). The GC receptors are present at the apical and at the basolateral sides, but cGMP expression is polarized at the basolateral side (349).

The functional role of ANP on salt and water transport in the lung is not clear. In one study in guinea pigs, ANP administration seemed to have a protective effect on the development of cardiogenic and noncardiogenic lung edema (157). However, in isolated perfused liquid-filled rat lungs, ANP increased alveolar epithelial permeability and decreased active sodium transport, thus decreasing

alveolar liquid clearance (269). In sheep with left atrial hypertension, there was an increase in plasma levels of ANP that may have inhibited the normal upregulation of fluid clearance that should have occurred in the presence of a rise in endogenous catecholamines (49). In alveolar type II cells cultured on filters, ANP decreased the amiloride-sensitive component of ^{22}Na influx (349) but did not change the basal $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity (49, 349). Interestingly, in the presence of β_2 -agonist stimulation, ANP inhibited $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity (49). Thus, on balance, it appears that ANP may impair fluid clearance by a direct effect on reducing sodium transport in alveolar type II cells.

B. Hypoxia

Hypoxia may occur during residence or recreation at high altitudes and under a variety of pathological conditions associated with acute and chronic respiratory disease. Therefore, it is important to understand the effect of hypoxia on the ion and fluid transport capacity of the lung epithelium.

Both in vitro and in vivo studies clearly show that decreased O_2 tension reduces the capacity of alveolar epithelial cells to actively transport sodium across alveolar epithelium. In alveolar type II cells, hypoxia (0% and 3% O_2) inhibits dome formation (292), decreases both amiloride-sensitive ^{22}Na influx and $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity, and decreases the amiloride-sensitive I_{sc} (208, 209, 292, 293, 383), indicating that the transepithelial sodium transport is impaired. The mechanisms whereby hypoxia induces decreased sodium transport proteins activity depend on the severity and the length of hypoxic exposure. For long exposure times (>12 h), the decrease in amiloride-sensitive sodium channel and $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity was associated with a parallel decline in mRNA levels of the three subunits, α , β , and γ of rENaC and two subunits α_1 - and β_1 - $\text{Na}^+\text{-K}^+\text{-ATPase}$ and the rate of α -rENaC protein synthesis, indicating a transcriptional or posttranscriptional regulation. For a short exposure time (3-h exposure), the decrease in ^{22}Na influx and $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity preceded any detectable change in mRNA levels, findings that suggest other mechanisms may be involved in regulation, including decreased efficiency in the translation of rENaC mRNA or in apical membrane trafficking of rENaC subunits, abnormal degradation or internalization of the channel protein, or hypoxia-induced modification of intracellular signals (147, 292).

The decrease in gene expression of sodium proteins likely represents the direct effect of hypoxia and raised the question of whether gene expression for sodium channels and $\text{Na}^+\text{-K}^+\text{-ATPase}$ is regulated by ambient P_{O_2} in alveolar type II cells. In support of this hypothesis, an increase in O_2 tension upregulated the level of ENaC mRNA transcripts in alveolar type II cells. The transfer of

fetal distal lung epithelial (FDLE) cells in culture from 3% O_2 to higher O_2 concentrations upregulated α -, β -, and γ -rENaC, mRNA subunits as well as sodium channel activity (10, 283, 296). Exposure of rats to sublethal hyperoxia (85% O_2) increased alveolar type II cell α -rENaC mRNA level (387). In addition, it has been recently reported that α_1 - and β_1 - $\text{Na}^+\text{-K}^+\text{-ATPase}$ mRNA subunit levels both increased in alveolar type II cells from rats exposed to hyperoxia (148). The mechanism whereby O_2 tension regulates sodium channels and $\text{Na}^+\text{-K}^+\text{-ATPase}$ gene expression is thought to be transcriptional. Activation of ENaC by increased O_2 concentration is associated with NF- κ B activation consistent with the identification of NF- κ B transcription binding sites in the α -ENaC promoter region (294). In addition to the potential role for an O_2 -responsive element in the ENaC promoter, it is also possible that there are other genes that are O_2 responsive and alter ENaC expression through their expressed protein or metabolic products.

The effect of hypoxia under in vivo conditions has been studied primarily in rats. In anesthetized rats, as well as in isolated perfused lungs, hypoxia decreased alveolar liquid clearance by inhibition of the amiloride-sensitive component (338, 369). In contrast to the in vitro studies, hypoxia increased α -rENaC and β_1 - $\text{Na}^+\text{-K}^+\text{-ATPase}$ mRNA transcripts with little increase or no change in protein amounts, suggesting a posttranslational mechanism such as a direct change of sodium transporter protein activity or protein internalization (369). This latter hypothesis was supported by the normalization of fluid clearance by a cAMP agonist (terbutaline), which is known to increase the trafficking of sodium transporter proteins from the cytoplasm to the membrane (48, 183, 328).

There is some limited data that ouabain-like substances may be released under conditions of hypoxia that could decrease epithelial fluid transport. A soluble factor that inhibits $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity is released from cultured rat alveolar type II cells after hypoxic exposure (293). Also, an endogenous ouabain-like compound has been found to be increased in high altitude in human plasma (80, 107, 108). One experimental study in rats that reported a reduction in airway epithelial sodium transport from hypoxia supported this hypothesis as well (352).

C. Anesthetics

In alveolar epithelial cells, the halogenated anesthetics affect sodium and fluid transport at the physiological level as well as on a cellular level. In the rat, halothane and isoflurane decrease fluid clearance by inhibition of the amiloride-sensitive component. This effect was rapidly reversible after cessation of halothane exposure (301). Unlike the rat, the ability of the rabbit to clear fluid from the alveolar space through an amiloride-sensitive

pathway is not decreased by halothane (253). In vitro, exposure to a low concentration of halothane (1%) and for a short time (30 min) induced a reversible decrease in $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity and amiloride-sensitive ^{22}Na influx in rat alveolar type II cells (242). The mechanisms whereby halothane induced a decrease in sodium transport protein activity have not been yet elucidated but are not related in vitro to a decrease in intracellular ATP content or to change in cytosolic free calcium. Also, some studies show that halothane increases permeability of alveolar capillary barrier to both water and protein as measured by the pulmonary clearance of $^{99\text{m}}\text{TcDTPA}$ in both rabbit (384) and humans (58). Taken together, these observations suggest that halogenated anesthetics may interfere with the clearance of alveolar edema.

Lidocaine is widely used in patients with acute cardiac disorders and has also been recently implicated as a possible cause of pulmonary edema after liposuction. In experimental studies in rats, either intravenous or intra-alveolar lidocaine reduced fluid clearance in rats by 50% (189). Because lidocaine did not inhibit ENaC when expressed in oocytes, it seems that the inhibitory effect on vectorial fluid transport was primarily on the basal surface of alveolar epithelial cells, either through an effect on the activity of $\text{Na}^+\text{-K}^+\text{-ATPase}$ or through an indirect effect through blockade of potassium channels, a well-known property of lidocaine (47, 156, 302, 357). The effect of lidocaine was completely reversible with β_2 -agonist therapy (189).

D. Reactive Oxygen and Nitrogen Species

Under several pathological conditions, in response to proinflammatory cytokines, activated neutrophils and macrophages can localize in the lung and migrate into the airspaces of the lung and release reactive oxygen species by the membrane-bound enzyme complex NADPH oxidase and nitric oxide (NO) via the calcium-insensitive inducible form of NO synthase. NO decreased I_{sc} across cultured rat type II cells without affecting transepithelial resistance. NO also inhibited 60% of amiloride-sensitive I_{sc} across type II cell monolayers after permeabilization of the basolateral membrane with amphotericin B (141). NO reacted with superoxide ($\text{O}_2^{\cdot-}$) to form peroxynitrite (ONOO^-), a potent oxidant and nitrating species that directly oxidizes a wide spectrum of biological molecules, such as DNA constituents, lipids, and proteins (220). Boluses of peroxynitrite (0.5–1 mM) into suspensions of freshly isolated type II cells from rabbits decreased amiloride-inhibitable sodium uptake to 68 and 56% of control values without affecting cell viability (153). Some investigators reported that products of macrophages, including NO, can downregulate sodium transport in fetal distal lung epithelium stimulated with endotoxin (71, 83). Also,

another study indicated that a generator of peroxynitrite (3-morpholiniosydnonimine) inhibited the amiloride-sensitive whole cell conductance in *Xenopus* oocytes expressing the three cloned subunits of ENaC (90). The data indicate that oxidation of critical amino acids residues in ENaC protein is probably responsible for this effect. This evidence matches well with other studies that have shown that protein nitration and oxidation by reactive oxygen and nitrogen species have been associated with diminished function of a variety of important proteins present in the alveolar space, including α_1 -proteinase inhibitor (243) and surfactant protein A (389–391).

V. EPITHELIAL FLUID TRANSPORT UNDER PATHOLOGICAL CONDITIONS

Fluid clearance from the distal airspaces of the lung has been measured in mechanically ventilated patients with acute respiratory failure from pulmonary edema as well as in several animal models designed to simulate clinically relevant pathological conditions (Fig. 5).

A. Clinical Studies

Studies of fluid clearance have been done in intubated, ventilated patients by measuring the concentration of total protein in sequential samples of undiluted pulmonary edema fluid aspirated from the distal airspaces of the lung with a standard suction catheter passed through the endotracheal tube into a wedged position in the distal airways of the lung (150, 230, 361, 374, 375, 391). This method for measuring fluid clearance in patients was adapted from the method for aspirating fluid from the distal airspaces of the lung in experimental studies in small and large animals (30, 228). The clinical procedure has been validated in patients by demonstrating that there is a relationship between fluid clearance and the improvement in oxygenation and the chest radiograph (230, 361).

1. Resolution of hydrostatic pulmonary edema

In 65 patients with severe hydrostatic pulmonary edema, predominantly from acute or chronic left ventricular dysfunction, there was net fluid clearance in the majority of the patients during the first 4 h after endotracheal intubation and the onset of positive pressure ventilation (361). The rate of fluid clearance in these patients varied between maximal ($>14\%/h$) in 38% and submaximal (3–14%/h) in 37%. Overall, 75% of the patients had intact fluid clearance. There was no significant correlation between the levels of fluid clearance and endogenous plasma levels of epinephrine, although twice as many of the patients with intact fluid clearance received aerosolized β -adrenergic therapy as those with impaired fluid




FIG. 5. This histological section stained with hematoxylin and eosin shows pulmonary edema from a patient who died with congestive heart failure. The edema fluid is located in the alveoli and is stained pink; a few red blood cells are also evident in the edema fluid. Note that there are a few alveoli that are air filled (white). Note the *inset* in the *bottom left* identifies the two most important lung epithelial sodium transporters that are responsible for the active removal of alveolar edema fluid. In the top half of the *inset*, epithelial sodium channels (ENaC) are shown labeling the apical surface of distal airway epithelium, and in the bottom half of the *inset*, there is diffuse basolateral staining of distal airway epithelium for $\text{Na}^+\text{-K}^+\text{-ATPase}$. (The authors appreciate the assistance of Rebecca Cleff and Kurt Albertine, PhD, in preparation of this figure.)

clearance, but this difference did not reach statistical significance, perhaps because the total number of studied patients was modest. The inability to transport edema fluid from the distal airspaces of the lung in 25% of the patients was not simply related to elevated pulmonary vascular pressures. Several mechanisms could downregulate fluid transport in these patients including elevated levels of ANF (49, 269) or the presence of ouabain-like substances in the circulation (80, 107, 108). Experimental studies have provided some insight into the mechanisms that may downregulate fluid transport from the distal airspaces of the lung in the presence of elevated pulmonary vascular hydrostatic pressures (see sect. vA2). Because hydrostatic pulmonary edema is associated with an uninjured epithelial barrier, the studies of hydrostatic pulmonary edema provide an important comparison group to the patients with pulmonary edema from acute lung injury because some degree of morphological or functional injury to the epithelial barriers probably occurs in most lung injuries (see sect. vB). Also, there is experimental evidence that accelerated fluid clearance can occur during the resolution of hydrostatic pulmonary edema in patients that is faster than in sheep, but similar to the fast rates recorded in rats (120) (Table 2). There is also recent data that very fast rates of fluid clearance can occur from the lungs of patients with pulmonary alveolar proteinosis by catecholamine-independent mechanisms (63).

2. Resolution of increased permeability edema from clinical acute lung injury

The majority of patients with increased permeability edema and acute lung injury have impaired alveolar epithelial fluid transport, a finding that is associated with

more prolonged respiratory failure and a higher mortality (Fig. 6). In contrast, a minority of patients can remove alveolar edema fluid rapidly, and these patients have a higher survival rate (230, 374, 375). These results indicate that a functional, intact distal lung epithelium is associated with a better prognosis in patients with acute lung injury, thus supporting the hypothesis that the degree of injury to the distal lung epithelium is an important determinant of the outcome in patients with increased permeability pulmonary edema from acute lung injury.

What are the mechanisms that may impair fluid clearance from the airspaces of the lung? Some patients have pathological (8) and biochemical (248) evidence of necrotic injury to the alveolar epithelium. There is also some clinical data that a decrease in fluid clearance may be associated with higher levels of nitrate and nitrite in pulmonary edema fluid, a finding that supports the hypothesis that nitration and oxidation of proteins essential

TABLE 2. *Fluid clearance in intact ventilated sheep, rats, and humans under basal and stimulated conditions and with and without hydrostatic pulmonary edema*

Species	Fluid Clearance Without Hydrostatic Pulmonary Edema, %/h			Fluid Clearance With Hydrostatic Pulmonary Edema, %/h		
	Basal	Stimulated	% Increase	Basal	Stimulated	% Increase
Sheep	8	13	62	5	8	60
Rat	20	35	75	16	22*	38
Human		54‡			25†	

* Lung fluid clearance (%/h). † Rate was the mean value for patients in the highest clearance group (see Ref. 361). ‡ Based on data obtained from lavage fluid of patients with pulmonary alveolar proteinosis (see Ref. 63).




FIG. 6. Hospital mortality (*y*-axis) plotted against two groups of patients with acute lung injury or the acute respiratory distress syndrome: those with maximal fluid clearance ($>14\%/h$) and those with impaired or submaximal fluid clearance ($<14\%/h$). The columns represent percent hospital mortality in each group (n = number of patients). Hospital mortality of patients with maximal fluid clearance was significantly less ($P < 0.02$). [From Ware and Matthay (375), with permission from The American Thoracic Society.]

to the epithelial fluid transport may occur in some patients with lung injury, depressing their ability to remove alveolar edema fluid (391) (see sect. *vB2*)

B. Experimental Studies

This section discusses what has been learned from experimental studies of high-pressure, hydrostatic pulmonary edema as well as from animal models of acute lung injury with increased permeability pulmonary edema.

1. Hydrostatic stress

The first major study to evaluate the effect of acute hydrostatic pulmonary edema on fluid clearance was done in anesthetized, ventilated sheep in which left atrial pressure was elevated acutely to either 18 or 25 cmH₂O (49). The rise in left atrial pressure created lung interstitial edema with the expected increase in protein poor lung lymph flow. Alveolar flooding was simulated with instillation of large volumes of isosmolar 5% albumin Ringer lactate solution into the distal airspaces of both lungs. Remarkably, fluid clearance remained at a normal level over 4 h. Because there was a mild increase in plasma epinephrine levels, bilateral adrenalectomies were done to determine the contribution of endogenous β -adrenergic stimulation. After the acute removal of both adrenal glands, fluid clearance was reduced by 30%, but still functioned at 70% of normal rates, even in sheep with a left atrial pressure of 24 cmH₂O. This finding provided direct evidence that an intact distal lung epithelium could actively remove fluid even though there was interstitial edema and a moderately elevated left atrial pressure.

In an isolated perfused rat lung model, other investi-

gators have found that 15 cmH₂O pulmonary venous pressure reduces fluid clearance from the airspaces by 50%, although elevations to 5 and 10 cmH₂O (control = 0 cmH₂O in this model) had no effect (310). The lower level at which vascular pressure impaired fluid clearance in this model compared with the sheep studies may be explained by the lack of functioning lung lymphatics in these isolated perfused lung studies. Interestingly, in this rat study, there was no evidence of morphological injury to the epithelial barrier, a finding that matches well with the finding that a reduction of vascular pressure was associated with a return of fluid clearance to a normal level (7). Thus, as in the sheep studies, there was no evidence of sustained injury to the epithelial barrier, a finding that fits well with the clinical studies of the resolution of hydrostatic pulmonary edema (361).

It is possible that there may be some mechanisms, other than elevated lung vascular pressures, that could depress vectorial fluid transport in hydrostatic pulmonary edema. For example, as already discussed, ANF is released with atrial stretch; this molecule can inhibit alveolar epithelial sodium uptake and fluid clearance (49, 269, 310, 349). Also, hypoxia may impair vectorial fluid clearance from the airspaces of the lung (see sect. *ivB* for discussion of hypoxia and epithelial fluid transport).

Can the resolution of alveolar edema be accelerated in hydrostatic lung edema? In the initial sheep study, aerosolization of salmeterol, a potent lipid-soluble β_2 -agonist, did not increase fluid clearance over 4 h in the presence of left atrial pressure elevation to either 18 or 24 cmH₂O (120). When left atrial pressure was normalized (6 cmH₂O), then aerosolized salmeterol markedly increased fluid clearance (49). Studies in rats demonstrated that a β_2 -agonist could enhance the resolution of lung edema and improve oxygenation in the resolution phase of hydrostatic pulmonary edema (120). Also, in isolated perfused rat lungs (7), there was a significant increase in fluid clearance with administration of isoproterenol or dopamine.

2. Hypovolemic shock

Hypovolemic shock from blood loss is an important clinical problem after major trauma. Short-term studies in rats that simulated acute hemorrhagic shock with a 30% loss of blood volume resulted in a sharp rise in endogenous levels of plasma epinephrine, a finding that was associated with a doubling of fluid clearance from the distal airspaces of the lung (241, 285). The effect was inhibited by propranolol and partially inhibited by amiloride.

When hypovolemic shock, however, was prolonged for 4–5 h in rats, the results were markedly different. Under these conditions, there was no increase in fluid transport from the airspaces of the lung, even when β_2 -

adrenergic agonists were instilled into the distal airspaces. This result prompted a series of experiments to discover the mechanisms that downregulate fluid clearance after prolonged hemorrhagic shock in rats. The initial studies established that the mechanism was neutrophil dependent (240). Further studies established that the process involves α -adrenergic activity and release of oxidant radicals with interleukin-1 β in the airspaces (190, 198), probably from neutrophils that accumulate in the lung after the onset of hemorrhagic shock. Finally, recent work has indicated that an increase in the expression of inducible NO synthase in the lung and release of NO, probably in part from alveolar macrophages, diminishes the capacity of the alveolar epithelium to actively transport fluid from the airspaces after severe hemorrhage. Second, NO inhibits the upregulation of alveolar epithelial fluid transport by cAMP-dependent mechanisms by directly affecting the function of the β_2 -adrenergic receptor and adenylyl cyclase. Third, shock-mediated release of NO in the airspaces of the lung depends in part on the activation and nuclear translocation of NF- κ B (288).

The results of these experimental studies may have important clinical implications for explaining the susceptibility to pulmonary edema in some patients after major trauma. Since, as already discussed, in vitro studies demonstrated that peroxynitrate can directly impair the function of sodium channels (24, 90, 141, 143, 153, 218), these in vivo studies fit well with one recent clinical study that reported an inverse relationship between elevated levels of nitrate and nitrite and the rate of fluid clearance in patients with pulmonary edema (391). Also, one prior study identified nitrotyrosine (the stable by-product of peroxynitrite reactions with tyrosine residues) in the lungs of patients with the acute respiratory distress syndrome (144) as well as in alveolar macrophages isolated from patients with the same cause of acute respiratory failure (11, 325).

3. Endotoxemia/bacteremia

The effect of endotoxemia and bacteremia on lung vascular permeability was well described in studies in sheep several years ago (43, 44). However, the impact on the function of the alveolar epithelial barrier was not addressed in those studies. More recent work has indicated that the acute shock produced by severe bacteremia in rats markedly increases plasma epinephrine levels, as in hemorrhagic shock, and the elevated epinephrine levels markedly upregulate the fluid transport capacity of the distal lung epithelium (290). Thus it is possible that in the short term, upregulation of fluid clearance may protect the airspaces against alveolar flooding when there is an increase in lung vascular permeability and accumulation of interstitial edema. In fact, one study in sheep demonstrated that lung vascular permeability can be augmented

markedly with intravenous endotoxin with a rise in protein-rich lung lymph flow, but this effect was not associated with a change in lung epithelial permeability to protein and no change in the capacity of the alveolar epithelium to remove alveolar fluid (380). These studies were done over 4 and 24 h in sheep, and in some of the studies both intra-alveolar and intravenous endotoxin was administered, but in all cases the epithelial barrier remained intact and capable of transporting alveolar fluid normally.

However, when large doses of live bacteria (*Pseudomonas aeruginosa*) were given to sheep, there was an increase in both lung endothelial and epithelial permeability to protein in the sheep that had developed the most severe shock (291). These sheep had alveolar flooding, and their capacity to remove alveolar fluid was impaired, similar to the findings in humans who develop severe permeability pulmonary edema with septic shock (305). The mechanisms for injury to the epithelial barrier probably depend on both neutrophil-dependent release of injurious proteases and reactive oxygen species as well as the bacterial exoproducts. In one study, gram-negative bacteria that produced proteases increased alveolar epithelial barrier permeability to protein by altering basolateral surface permeability while the nonprotease-producing strains only increased lung vascular permeability (287).

4. Pneumonia

In sharp contrast to intra-alveolar endotoxin, live bacteria increased alveolar epithelial barrier permeability and decreased fluid transport in sheep (380). Further studies indicated that the products of *P. aeruginosa* were important in determining the extent of injury. For example, exoenzyme S and phospholipase C mediated injury to the epithelial barrier in rabbits with a decrease in vectorial fluid transport (381). Subsequent studies indicated that bacterial pneumonia may progress to septic shock when the infecting gram-negative organism generates proinflammatory cytokines in the airspaces of the lung that are released into the circulation when bacterial-mediated injury results in sufficient injury to the distal lung epithelial barrier (188). Several experimental studies have indicated that active and passive immunization against *P. aeruginosa* antigens can prevent epithelial injury in sheep (289) and in mice (314).

Recent data also indicate that influenza virus infection (A/PR/8/34) can specifically alter epithelial ion transport by inhibiting amiloride-sensitive sodium current across mouse tracheal epithelium (187). The inhibitory effect of the influenza virus was caused by binding the viral hemagglutinin to a cell-surface receptor, which then activated phospholipase C and protein kinase C. It is well known that protein kinase C can reduce ENaC activity so that influenza infections in the lung may inhibit the func-

tion of ENaC (138). Given the importance of sodium channels in vectorial transport of fluid in distal airway epithelia and in the alveoli, these results provide a new mechanism that may explain the accumulation of alveolar edema fluid in patients with viral pneumonia and acute lung injury.

Interestingly, in the process of studying *P. aeruginosa* pneumonia in rats, one group of investigators found that the rate of fluid clearance was upregulated in the rats that survived (300). The mechanism for this effect was secondary to release of tumor necrosis factor- α (TNF- α), which was surprising since TNF- α plays an important role in mediating the host inflammatory response to infection (1, 175, 200) as well as potentially contributing to the pathogenesis of septic shock (225). It had been previously reported that TNF- α could increase sodium-coupled amino acid transport across hepatocytes (278). The capacity of TNF- α to increase fluid clearance was confirmed in a subsequent rat study (36) in which a neutralizing polyclonal anti-TNF- α antibody inhibited the upregulation of fluid clearance induced by intestinal ischemia-reperfusion. Also, the effect of TNF- α is amiloride inhibitable in both rats (36, 122) and isolated A549 cell patch-clamp studies (122). Based on the A549 human cell studies, the effect appears to be a receptor-mediated process. It is not clear, however, what signaling pathways are involved, since cAMP levels are not elevated by TNF- α (36).

5. Hyperoxia

Several investigators have used hyperoxia as a model to study the effect of acute lung injury on epithelial ion and fluid transport in the lung, in part because the injury develops over 2–5 days in rats and mice and pathologically resembles clinical acute lung injury with both endothelial and epithelial injury in association with an influx of neutrophils and protein-rich pulmonary edema.

However, the results of these studies have not been uniform, in part because of variations in the duration of O₂ exposure, the exact level of hyperoxia, and the use of rats or mice. In one rat study, administration of 85% O₂ for 7 days increased the level of α -ENaC protein, and both inward and outward sodium currents were stimulated in patch clamps of isolated alveolar type II cells (149). Subsequently, another study from the same group showed increased expression and activity of amiloride-inhibitable, sodium channels in alveolar type II cells of rats exposed to 85% O₂ for 7 days followed by 100% O₂ for 4 days. Both the number and the open probability of the L-type sodium channels (25 pS) were increased (387). Another group also studied the effect of 85% O₂ for 7 days in rats and found that amiloride-inhibitable sodium uptake was greater than in control rats and that ouabain decreased active sodium transport to a greater percentage in the hyperoxic rats, suggesting an upregulation of Na⁺-K⁺-

ATPase activity after subacute hyperoxia (101, 128, 270, 343). In other studies, exposure to 100% O₂ for 48 h produced moderate interstitial lung edema but no impairment of basal or cAMP-stimulated fluid transport (125, 192, 311). However, when exposure was prolonged to 64 h, one group of investigators reported decreased transport in rats, an effect that seemed to be related to decreased gene expression of α_1 -Na⁺-K⁺-ATPase subunits (272). Other studies found that there was rapid upregulation of mRNA for α_1 - and β_1 -subunits of Na⁺-K⁺-ATPase as well as antigenic protein shortly after prolonged exposure to >97% O₂ in rats for 60 h (250, 251), suggesting the induction of the Na⁺-K⁺-ATPase could occur as a protective mechanism. Similar findings were reported by Stern et al. (335) in a thiourea model of lung injury in which upregulation of Na⁺-K⁺-ATPase gene expression and protein occurred after initial injury and was associated with recovery from the pulmonary edema. Other studies have reported that hyperoxia does not seem to produce a clear change in sodium and fluid transport during the period of hyperoxia (40, 55, 56).

Interestingly, one group of investigators reported that pretreatment of rats with aerosolized adenoviral β_1 -Na⁺-K⁺-ATPase upregulated fluid transport and also made the rats resistant to the lethal effects of hyperoxia (100). This represented the first evidence that gene therapy could potentially be used to produce a sustained upregulation of fluid transport in the lung in the presence of a pathological condition. Furthermore, gene transfer of both α_1 - and β_1 -Na⁺-K⁺-ATPase decreases edema formation induced by thiourea in rats (335). An earlier study had shown that administration of the adenoviral β_1 -Na⁺-K⁺-ATPase gene, but not the α_1 -Na⁺-K⁺-ATPase, would increase fluid transport in normal rats (102).

6. Subacute lung injury

The mechanisms that regulate fluid balance across the distal lung epithelium in the subacute phase after lung injury have been studied after bleomycin lung injury, a commonly used model for subacute lung injury that is associated with fibrosis (286). In one study, the rate of fluid clearance was doubled compared with normal lungs, a finding that correlated with a large increase in alveolar type II-like cells (115). However, these epithelial cells did not express α -ENaC, and when isolated and studied in vitro, the absolute rate of transport was only 50% of normal control type II cells, and all of the transport was amiloride insensitive. Also, β -agonist therapy did not upregulate transport. Thus the provisional repair epithelium in this rat model lacked the normal sodium and fluid transport properties, although the extent of epithelial hyperplasia with type II-like cells appeared to compensate well for the impairment of transport of the isolated type II-like cells. By 60 days, the lungs had recovered their

normal architecture, and fluid clearance from the distal airspaces was normal.

VI. FUTURE DIRECTIONS

Important advances have been made in the understanding of the reabsorption of fluid and solutes by the distal epithelia with characterization of sodium transport and water pathways under both physiological and pathological conditions. However, several fundamental issues require additional study.

Alveolar type II cells and distal airway epithelial cells, such as Clara cells, are implicated in sodium and fluid transport, but their relative contributions in both physiological and pathological conditions are not well defined. The current in vivo models and cultured cell systems cannot determine the relative contribution of these cells to net fluid transport in the distal lung. Innovative approaches are needed. Another important issue is the relative contribution of alveolar type I cells in sodium, chloride, and fluid transport across the alveolar epithelium. Alveolar type I cells cover 95% of the alveolar surface area, and the recent demonstration of the presence of water channels and ENaC expression in those cells suggests a role for these cells in net fluid clearance. Improved models are needed to assess the differential contribution of alveolar type I and II cells to fluid transport.

Another important area of research is the characterization of the sodium transporters involved in sodium and fluid reabsorption and their regulation. Amiloride-sensitive sodium transport is one of the major pathways for sodium entry across distal epithelial cells, but several questions remain unsolved. For example, are the molecular and biophysical characteristics of these channels in vitro representative of their in vivo characteristics, and how are these channels regulated during physiological and pathological conditions? The mechanisms that regulate the trafficking of ENaC and $\text{Na}^+\text{-K}^+\text{-ATPases}$ between the cytoplasm and the membrane need to be evaluated also in distal lung epithelia. Increased insertion of transport proteins may be one important mechanism for increasing sodium and fluid transport under pathological conditions and may potentially contribute to regulating the clearance of edema fluid from distal airspaces of the lung. In addition to amiloride-sensitive sodium transport, a characterization of ion transporters involved in amiloride-insensitive sodium transport needs to be defined. Also, the pathways for chloride reabsorption under basal and stimulated conditions need to be determined with particular attention to the role of CFTR under cAMP-stimulated conditions.

Recent advances have been made with transgenic mice models to define the role of sodium and water

channels in the lung fluid balance. Knockout of the three subunits of ENaC has clearly established the preponderant role of α - compared with β - and γ -ENaC in alveolar transepithelial sodium absorption. Similarly, the knockout mice for several aquaporin-type water channels have revealed that in the lung these channels are not essential for water transport. However, a genomic disruption of genes that are expressed during development or in multiple tissue types complicates the phenotypic analysis. A solution to this problem may be provided by conditional knockouts. This system permits control of the timing for cell-specific expression of specific proteins, thereby circumventing both embryonic lethality and confounding effects of complex adaptive responses that can occur when the physiological observations follow the gene knockout events by days or weeks. In this system, gene expression is regulated temporally and spatially using cell-specific promoters, such as SP-C for alveolar type II cells, in combination with a regulatory on-off system. This approach may provide a major opportunity to advance the understanding of the role of sodium and fluid transport during physiological and pathological conditions during the reabsorption of edema from the distal airspaces of the lung.

Progress in the last decade demonstrates that it is feasible to quantify the rate of edema reabsorption from the distal airspaces of the lung in ventilated, critically ill patients with acute pulmonary edema. In conjunction with progress in experimental studies of lung fluid balance under clinically relevant pathological conditions, further studies should be done to test the potential role of catecholamine-dependent and -independent therapies that might enhance the resolution of clinical pulmonary edema.

This work was supported by National Heart, Lung, and Blood Institute Grants HL-51854 and HL-51856 (to M. A. Matthay).

Address for reprint requests and other correspondence: M. A. Matthay, Univ. of California at San Francisco, 505 Parnassus Ave., Rm. M917, San Francisco, CA 94143-0624 (E-mail: mmatt@itsa.ucsf.edu).

REFERENCES

1. ABRAHAM E, WUNDERINK R, SILVERMAN H, PERL TM, NASRAWAY S, LEVY H, BONE R, WENZEL RP, BALK R, AND ALLRED R. Efficacy and safety of monoclonal antibody to human tumor necrosis factor alpha in patients with sepsis syndrome A randomized, controlled, double-blind, multicenter clinical trial TNF-alpha MAb Sepsis Study Group. *JAMA* 273: 934-941, 1995.
2. AGRE P. Aquaporin null phenotypes: the importance of classical physiology (commentary). *Proc Natl Acad Sci USA* 95: 9061-9063, 1998.
3. AGRE P, PRESTON GM, SMITH BL, JUNG JS, RAINA S, MOON C, GUGGINO WB, AND NIELSEN S. Aquaporin CHIP: the archetypal molecular water channel. *Am J Physiol Renal Physiol* 265: F463-F476, 1993.
4. AHMAD M AND MEDFORD RM. Evidence for the regulation of $\text{Na}^+\text{-K}^+$ -

- ATPase $\alpha 1$ gene expression through the interaction of aldosterone and cAMP-inducible transcriptional factors. *Steroids* 60: 147–152, 1995.
5. AL-BAZZAZ FJ. Regulation of Na and Cl transport in sheep distal airways. *Am J Physiol Lung Cell Mol Physiol* 267: L193–L198, 1994.
 6. AWAYDA MS, ISMAILOV II, BERDIEV BK, FULLER CM, AND BENOS DJ. Protein kinase regulation of a cloned epithelial Na^+ channel. *J Gen Physiol* 108: 49–65, 1996.
 7. AZZAM ZS, SALLDAS FJ, COMELLAS A, RIDGE KM, RUTSCHMAN DH, AND SZNAJDER JI. Catecholamines increase lung edema clearance in rats with increased left atrial pressure. *J Appl Physiol* 90: 1088–1094, 2001.
 8. BACHOFEN M AND WEIBEL ER. Alterations of the gas exchange apparatus in adult respiratory insufficiency associated with septicemia. *Am Rev Respir Dis* 116: 589–615, 1977.
 9. BAI C, FUKUDA N, SONG Y, MA T, MATTHAY MA, AND VERKMAN AS. Lung fluid transport in aquaporin-1 and aquaporin-4 knockout mice. *J Clin Invest* 103: 555–561, 1999.
 10. BAINES DL, RAMMINGER SJ, COLLETT A, HADDAD JJE, BEST OG, LAND SC, OLVER RE, AND WILSON SM. Oxygen-evoked Na^+ transport in rat fetal distal lung epithelial cells. *J Physiol* 532: 105–113, 2001.
 11. BALDUS S, CASTRO L, EISERICH JP, AND FREEMAN BA. Is NO news bad news in acute respiratory distress syndrome? *Am J Respir Crit Care Med* 163: 308–310, 2001.
 12. BALL RL, TANNER KD, AND CARPENTER G. Epidermal growth factor potentiates cyclic AMP accumulation in A-431 cells. *J Biol Chem* 265: 12836–12845, 1990.
 13. BALLARD ST, SCHEPENS SM, FALCONE JC, MEININGER GA, AND TAYLOR AE. Regional bioelectric properties of porcine airway epithelium. *J Appl Physiol* 73: 2021–2027, 1992.
 14. BALLARD ST AND TAYLOR AE. Bioelectric properties of proximal bronchiolar epithelium. *Am J Physiol Lung Cell Mol Physiol* 267: L79–L84, 1994.
 15. BARBRY P AND HOFMAN P. Molecular biology of Na^+ absorption. *Am J Physiol Gastrointest Liver Physiol* 273: G571–G585, 1997.
 16. BARKER PM, NGUYEN MS, GATZY JT, GRUBB B, NORMAN H, HUMMLER E, ROSSIER B, BOUCHER RC, AND KOLLER B. Role of γENaC subunit in lung liquid clearance and electrolyte balance in newborn mice. Insights into perinatal adaptation and pseudohypoaldosteronism. *J Clin Invest* 102: 1634–1640, 1998.
 17. BARNARD ML, OLIVERA WG, RUTSCHMAN DM, BERTORELLO AM, KATZ AI, AND SZNAJDER JI. Dopamine stimulates sodium transport and liquid clearance in rat lung epithelium. *Am J Respir Crit Care Med* 156: 709–714, 1997.
 18. BARNARD ML, RIDGE KM, SALLDAS F, FRIEDMAN E, GARE M, GUERRERO C, LECUONA E, BERTORELLO AM, KATZ AI, AND SZNAJDER JI. Stimulation of the dopamine 1 receptor increases lung edema clearance. *Am J Respir Crit Care Med* 160: 982–986, 1999.
 19. BARNES PJ. Beta-adrenergic receptors and their regulation. *Am J Respir Crit Care Med* 152: 838–860, 1995.
 20. BARQUIN N, CICCOCIELLA DE, RIDGE KM, AND SZNAJDER JI. Dexamethasone upregulates the Na-K-ATPase in rat alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 273: L825–L830, 1997.
 21. BASSET G, CRONE C, AND SAUMON G. Significance of active ion transport in transalveolar water absorption: a study on isolated rat lung. *J Physiol* 384: 311–324, 1987.
 22. BASSET G, CRONE C, AND SAUMON G. Fluid absorption by rat lung in situ: pathways for sodium entry in the luminal membrane of alveolar epithelium. *J Physiol* 384: 325–345, 1987.
 23. BASSET G, SAUMON G, BOUCHONNET F, AND CRONE C. Apical sodium-sugar transport in pulmonary epithelium in situ. *Biochim Biophys Acta* 942: 11–18, 1998.
 24. BAUER ML, BECKMAN JS, BRIDGES RJ, FULLER CM, AND MATALON S. Peroxynitrite inhibits sodium uptake in rat colonic membrane vesicles. *Biochim Biophys Acta* 1104: 87–94, 1992.
 25. BERDIEV BK, PRAT AG, CANTIELLO HF, AUSIELLO DA, FULLER CM, JOVOV B, BENOS DJ, AND ISMAILOV II. Regulation of epithelial sodium channels by short actin filaments. *J Biol Chem* 271: 17704–17710, 1996.
 26. BERTHIAUME Y. Effect of exogenous cAMP and aminophylline on alveolar and lung liquid clearance in anesthetized sheep. *J Appl Physiol* 70: 2490–2497, 1991.
 27. BERTHIAUME Y, ALBERTINE KH, GRADY M, FICK G, AND MATTHAY MA. Protein clearance from the airspaces and lungs of unanesthetized sheep over 144 h. *J Appl Physiol* 67: 1887–1897, 1989.
 28. BERTHIAUME Y, BROADDUS VC, GROPPER MA, TANITA T, AND MATTHAY MA. Alveolar liquid and protein clearance from normal dog lungs. *J Appl Physiol* 65: 585–593, 1988.
 29. BERTHIAUME Y, SAPIJASZKO M, MACKENZIE J, AND WALSH MP. Protein kinase C activation does not stimulate lung liquid clearance in anesthetized sheep. *Am Rev Respir Dis* 144: 1085–1090, 1991.
 30. BERTHIAUME Y, STAUB NC, AND MATTHAY MA. Beta-adrenergic agonists increase lung liquid clearance in anesthetized sheep. *J Clin Invest* 79: 335–343, 1987.
 31. BERTORELLO AM AND KATZ AI. Short-term regulation of renal Na-K-ATPase activity: physiological relevance and cellular mechanisms. *Am J Physiol Renal Physiol* 265: F743–F755, 1993.
 32. BERTORELLO AM, RIDGE KM, CHIBALIN AV, KATZ AI, AND SZNAJDER JI. Isoproterenol increases $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity by membrane insertion of α -subunits in lung alveolar cells. *Am J Physiol Lung Cell Mol Physiol* 276: L20–L27, 1999.
 33. BIEL M, ZONG X, DISTLER M, BOSSE E, KLUGBAUER N, MURAKAMI M, FLOCKERZ V, AND HOFMANN F. Another member of the cyclic nucleotide-gated channel family, expressed in testis, kidney, and heart. *Proc Natl Acad Sci USA* 91: 3505–3509, 1994.
 34. BLANCO G, SANCHEZ G, AND MERCER RW. Differential regulation of $\text{Na}^+\text{-K}^+\text{-ATPase}$ isozymes by protein kinases and arachidonic acid. *Arch Biochem Biophys* 359: 139–150, 1998.
 35. BONNY O, CHRAIBI A, LOFFING J, FOWLER JAEGER N, GRÜNDER S, HORISBERGER JD, AND ROSSIER BC. Functional expression of a pseudohypoaldosteronism type I mutated epithelial Na^+ channel lacking the pore-forming region of its α subunit. *J Clin Invest* 104: 967–974, 1999.
 36. BÖRJESSON A, NORLIN A, WANG X, ANDERSSON R, AND FOLKESSON HG. $\text{TNF}\alpha$ stimulates alveolar liquid clearance during intestinal ischemia-reperfusion in rats. *Am J Physiol Lung Cell Mol Physiol* 278: L3–L12, 2000.
 37. BOROK Z, DANTO SJ, KIM KJ, LUBMAN RL, AND CRANDALL ED. Effects of EGF and dexamethasone on bioelectric properties of alveolar epithelial cell monolayers (Abstract). *Am Rev Respir Dis* 147: 1005A, 1993.
 38. BOROK Z, LIEBLER JM, LUBMAN RL, FOSTER MJ, ZHOU B, LI X, ZABSKI SM, KIM KJ, AND CRANDALL ED. Sodium transport proteins are expressed by rat alveolar epithelial type I cells. *Am J Physiol Lung Cell Mol Physiol* 282: L599–L608, 2002.
 39. BOROK Z, LUBMAN RL, DANTO SI, ZHANG XL, ZABSKI SM, KING LS, LEE DM, AGRE P, AND CRANDALL ED. Keratinocyte growth factor modulates alveolar epithelial cell phenotype in vitro: expression of aquaporin 5. *Am J Respir Cell Mol Biol* 18: 554–561, 1998.
 40. BOROK Z, MIHYU S, FERNANDES VFJ, ZHANG XL, KIM KJ, AND LUBMAN RL. KGF prevents hyperoxia-induced reduction of active ion transport in alveolar epithelial cells. *Am J Physiol Cell Physiol* 276: C1352–C1360, 1999.
 41. BOUCHER RC. Human airway ion transport. Part one. *Am J Respir Crit Care Med* 150: 271–281, 1994.
 42. BOUCHER RC. Human airway ion transport. Part two. *Am J Respir Crit Care Med* 150: 581–593, 1994.
 43. BRIGHAM KL AND MEYERICK B. Endotoxin and lung injury. *Am Rev Respir Dis* 133: 913–927, 1986.
 44. BRIGHAM KL, WOOLVERTON WC, BLAKE LH, AND STAUB NC. Increased sheep lung vascular permeability caused by *Pseudomonas* bacteremia. *J Clin Invest* 54: 792–804, 1974.
 45. BROWN MJ, OLVER RE, RAMSDEN CA, STRANG LB, AND WALTERS DV. Effects of adrenaline and of spontaneous labour on the secretion and absorption of lung liquid in the fetal lamb. *J Physiol* 344: 137–152, 1983.
 46. BROWN SE, KIM KJ, GOODMAN BE, WELLS JR, AND CRANDALL ED. Sodium-amino acid cotransport by type II alveolar epithelial cells. *J Appl Physiol* 59: 1616–1622, 1985.
 47. BUTTERWORTH JF AND STRICHARTZ GR. Molecular mechanisms of local anesthesia: a review. *Anesthesiology* 72: 711–734, 1990.
 48. BUTTERWORTH MB, HELMAN SI, AND ELS WJ. cAMP-sensitive endocytic trafficking in A6 epithelia. *Am J Physiol Cell Physiol* 280: C752–C762, 2001.
 49. CAMPBELL AR, FOLKESSON HG, BERTHIAUME Y, GUTKOWSKA J, SUZUKI S,

- AND MATTHAY MA. Alveolar fluid clearance persists in the presence of moderate left atrial hypertension in sheep. *J Appl Physiol* 86: 139–151, 1999.
50. CAMPBELL L, HOLLINS AJ, AL-EID A, NEWMAN GR, VON RUHLAND C, AND GUMBLETON M. Caveolin-1 expression and caveolae biogenesis during cell transdifferentiation in lung alveolar epithelial primary cultures. *Biochem Biophys Res Commun* 262: 744–751, 1999.
 51. CANESSA CM, MERILLAT AM, AND ROSSIER BC. Membrane topology of the epithelial sodium channel in intact cells. *Am J Physiol Cell Physiol* 267: C1682–C1690, 1994.
 52. CANESSA CM, SCHILD L, BUELL G, THORENS B, GAUTSCHI I, HORISBERGER JD, AND ROSSIER BC. Amiloride-sensitive epithelial Na⁺ channel is made of three homologous subunits. *Nature* 367: 463–467, 1994.
 53. CARTER EP, MATTHAY MA, FARINAS J, AND VERKMAN AS. Transalveolar osmotic and diffusional water permeability in intact mouse lung measured by a novel surface fluorescence method. *J Gen Physiol* 108: 133–142, 1996.
 54. CARTER EP, OLVECKY BP, MATTHAY MA, AND VERKMAN AS. High microvascular endothelial water permeability in mouse lung measured by a pleural surface fluorescence method. *Biophys J* 74: 2121–2128, 1998.
 55. CARTER EP, WANGENSTEEN OD, DUNITZ J, AND INGBAR DH. Hyperoxic effects on alveolar sodium resorption and lung Na-K-ATPase. *Am J Physiol Lung Cell Mol Physiol* 273: L1191–L1202, 1997.
 56. CARTER EP, WANGENSTEEN OD, O'GRADY SM, AND INGBAR DH. Effects of hyperoxia on type II cell Na-K-ATPase function and expression. *Am J Physiol Lung Cell Mol Physiol* 272: L542–L551, 1997.
 57. CASSIN S, GAUSE G, AND PERKS AM. The effects of bumetanide and furosemide on lung liquid secretion in fetal sheep. *Proc Soc Exp Biol Med* 181: 427–431, 1986.
 58. CHANGLAI SP, HUNG WT, AND LIAO KK. Detecting alveolar epithelial injury following volatile anesthetics by ^{99m}Tc DTPA radioaerosol inhalation lung scan. *Respiration* 66: 506–510, 1999.
 59. CHARRON PD, FAWLEY JP, AND MARON MB. Effect of epinephrine on alveolar liquid clearance in the rat. *J Appl Physiol* 87: 611–618, 1999.
 60. CHEEK JM, KIM KJ, AND CRANDALL ED. Tight monolayers of rat alveolar epithelial cells: bioelectric properties and active sodium transport. *Am J Physiol Cell Physiol* 256: C688–C693, 1989.
 61. CHEN XJ, EATON DC, AND JAIN L. β -Adrenergic regulation of amiloride-sensitive lung sodium channels. *Am J Physiol Lung Cell Mol Physiol* 282: L609–L620, 2002.
 62. CHESNUTT AN, KHERADMAND F, FOLKESSON HG, ALBERTS MA, AND MATTHAY MA. Soluble transforming growth factor- α is present in the pulmonary edema fluid of patients with acute lung injury. *Chest* 111: 652–656, 1997.
 63. CHESNUTT MS, NUCKTON TJ, GOLDEN J, FOLKESSON HG, AND MATTHAY MA. Rapid alveolar epithelial fluid clearance following lung lavage in pulmonary alveolar proteinosis. *Chest* 120: 271–274, 2001.
 64. CHOW YH, WANG Y, PLUMB J, O'BRODOVICH H, AND HU J. Hormonal regulation and genomic organization of the human amiloride-sensitive epithelial sodium channel α subunit gene. *Pediatr Res* 46: 208–214, 1999.
 65. CLEMENTS JA. Lung surfactant: a personal perspective. *Annu Rev Physiol* 59: 1–21, 1997.
 66. CLEMENTS JA, HUSTEAD RF, JOHNSON RP, AND GRIBETZ I. Pulmonary surface tension and alveolar stability. *J Appl Physiol* 16: 444–450, 1961.
 67. CLERICI C. Sodium transport in alveolar epithelial cells: modulation by O₂ tension. *Kidney Int Suppl* 65: S79–S83, 1998.
 68. CLERICI C, COUETTE S, LOISEAU A, HERMAN P, AND AMIEL C. Evidence for Na-K-Cl cotransport in alveolar epithelial cells: effect of phorbol ester and osmotic stress. *J Membr Biol* 147: 295–304, 1995.
 69. CLERICI C, SOLER P, AND SAUMON G. Sodium-dependent phosphate and alanine transports but sodium-independent hexose transport in type II alveolar epithelial cells in primary culture. *Biochim Biophys Acta* 1063: 27–35, 1991.
 70. COLLETT A, RAMMINGER SJ, OLIVER RE, AND WILSON SM. β -Adrenoceptor-mediated control over the conductive properties of the apical membrane in distal lung epithelia. *Am J Physiol Lung Cell Mol Physiol* 282: L621–L630, 2002.
 71. COMPEAU CG, ROTSTEIN OD, TOHDA H, MARUNAKA Y, RAFII B, SLUTSKY AS, AND O'BRODOVICH H. Endotoxin-stimulated alveolar macrophages impair lung epithelial Na⁺ transport by an L-Arg-dependent mechanism. *Am J Physiol Cell Physiol* 266: C1330–C1341, 1994.
 72. COTT GR. Modulation of bioelectric properties across alveolar type II cells by substratum. *Am J Physiol Cell Physiol* 257: C678–C688, 1989.
 73. COTT GR, SUGAHARA K, AND MASON RJ. Stimulation of net active ion transport across alveolar type II cell monolayers. *Am J Physiol Cell Physiol* 250: C222–C227, 1986.
 74. CRANDALL ED, HEMING TH, PALOMBO RL, AND GOODMAN BE. Effect of terbutaline on sodium transport in isolated perfused rat lung. *J Appl Physiol* 60: 289–294, 1986.
 75. CRANDALL ED AND MATTHAY MA. Alveolar epithelial transport. Basic science to clinical medicine. *Am J Respir Crit Care Med* 163: 1021–1029, 2001.
 76. CRAPO JD, YOUNG SL, FRAM EK, PINKERTON KE, BARRY BE, AND CRAPO RO. Morphometric characteristics of cells in the alveolar region of mammalian lungs. *Am Rev Respir Dis Suppl* 128: S42–S46, 1983.
 77. DAGENAIS A, DENIS C, VIVES MF, GIROUARD S, MASSÉ C, NGUYEN T, YAMAGATA T, GRYGORCZYK C, KOTHARY R, AND BERTHAUME Y. Modulation of α -ENaC and α_1 -Na⁺-K⁺-ATPase by cAMP and dexamethasone in alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 281: L217–L230, 2001.
 78. DAGENAIS A, KOTHARY R, AND BERTHAUME Y. The α subunit of the epithelial sodium channel in the mouse: developmental regulation of its expression. *Pediatr Res* 42: 327–334, 1997.
 79. DANTO SI, BOROK Z, ZHANG XL, LOPEZ MZ, PATEL P, CRANDALL ED, AND LUBMAN RL. Mechanisms of EGF-induced stimulation of sodium reabsorption by alveolar epithelial cells. *Am J Physiol Cell Physiol* 275: C82–C92, 1998.
 80. DE ANGELIS C, RISCACCI M, SALVINI R, PICCOLI A, FERRI C, AND SANTUCCI A. Isolation and characterization of a digoxin-like immunoreactive substance from human urine by affinity chromatography. *Clin Chem* 43: 1416–1420, 1997.
 81. DIGLIO CA AND KIKAWA Y. The type II epithelial cells of the lung. IV. Adaptation and behavior of isolated type II cells in culture. *Lab Invest* 37: 622–631, 1977.
 82. DING C, POTTER ED, QIU W, COON SL, LEVINE MA, AND GUGGINO SE. Cloning and widespread distribution of the rat rod-type cyclic nucleotide-gated cation channel. *Am J Physiol Cell Physiol* 272: C1335–C1344, 1997.
 83. DING JW, DICKIE J, O'BRODOVICH H, SHINTANI Y, RAFII B, HACKAM D, MARUNAKA Y, AND ROTSTEIN OD. Inhibition of amiloride-sensitive sodium-channel activity in distal lung epithelial cells by nitric oxide. *Am J Physiol Lung Cell Mol Physiol* 274: L378–L387, 1998.
 84. DOBBS LG, GONZALEZ R, MATTHAY MA, CARTER EP, ALLEN L, AND VERKMAN AS. Highly water permeable type I alveolar epithelial cells confer high water permeability between the airspace and vasculature in rat lung. *Proc Natl Acad Sci USA* 95: 2991–2996, 1998.
 85. DOBBS LG, PIAN MS, MAGLIO M, DUMARS S, AND ALLEN L. Maintenance of the differentiated type II cell phenotype by culture with an apical air surface. *Am J Physiol Lung Cell Mol Physiol* 273: L347–L354, 1997.
 86. DOBBS LG, WILLIAMS MC, AND BRANDT AE. Changes in biochemical characteristics and pattern of lectin binding of alveolar type II cells with time in culture. *Biochim Biophys Acta* 846: 155–166, 1985.
 87. DUC C, FARMAN N, CANESSA CM, BONVALET JP, AND ROSSIER BC. Cell-specific expression of epithelial sodium channel α , β , and γ subunits in aldosterone-responsive epithelia from the rat: localization by in situ hybridization and immunocytochemistry. *J Cell Biol* 127: 1907–1921, 1994.
 88. DUMASIVUS V, SZNAJDER JI, AZZAM ZS, BOJA J, MUTLU GM, MARON MB, AND FACTOR P. β_2 -Adrenergic receptor overexpression increases alveolar fluid clearance and responsiveness to endogenous catecholamines in rats. *Circ Res* 89: 907–914, 2001.
 89. DUQUETTE PP, BISSONNETTE P, AND LAPOINTE JY. Local osmotic gradients drive the water flux associated with Na⁺/glucose cotransport. *Proc Natl Acad Sci USA* 98: 3796–3801, 2001.
 90. DUVALL MD, ZHU S, FULLER CM, AND MATALON S. Peroxynitrite inhibits amiloride-sensitive Na⁺ currents in *Xenopus* oocytes expressing $\alpha\beta\gamma$ -ENaC. *Am J Physiol Cell Physiol* 274: C1417–C1423, 1998.
 91. EFFROS RM. Osmotic extraction of hypotonic fluid from the lungs. *J Clin Invest* 5: 935–947, 1974.

92. EFFROS RM AND CHANG HK. Fluid and solute transport in the airspaces of the lungs. In: *Lung Biology in Health and Disease*, edited by Lenfant C. New York: Dekker, 1994, p. 539.
93. EFFROS RM, HACKER A, SILVERMAN P, AND HUKKANEN J. Protein concentrations have little effect on reabsorption of fluid from isolated rat lungs. *J Appl Physiol* 70: 416–422, 1991.
94. EFFROS RM, MASON GR, HUKKANEN J, AND SILVERMAN P. New evidence for active sodium transport from fluid-filled rat lungs. *J Appl Physiol* 66: 906–919, 1989.
95. ENGELHARDT JF, ZEPEDA M, COHN JA, YANKASKAS JR, AND WILSON JM. Expression of the cystic fibrosis gene in adult human lung. *J Clin Invest* 93: 737–749, 1994.
96. ESCOUBET B, COUREAU C, BONVALET JP, AND FARMAN N. Noncoordinate regulation of epithelial Na channel and Na pump subunit mRNAs in kidney and colon by aldosterone. *Am J Physiol Cell Physiol* 272: C1482–C1491, 1997.
97. ESKANDARI S, SNYDER PM, KREMAN M, ZAMPIGHI GA, WELSH MJ, AND WRIGHT EM. Number of subunits comprising the epithelial sodium channel. *J Biol Chem* 274: 27281–27286, 1999.
98. FABISIAK JP, VESELL ES, AND RANNELS DE. Interactions of beta adrenergic antagonists with isolated rat alveolar type II pneumocytes. I. Analysis, characterization and regulation of specific beta adrenergic receptors. *J Pharmacol Exp Ther* 241: 722–727, 1987.
99. FACTOR P, AZZAM Z, AND DUMASIVUS V. β_2 -Adrenergic receptor overexpression increases alveolar fluid clearance by upregulating solute transport proteins on both sides of the cell membrane (Abstract). *Am J Respir Crit Care Med* 163: A570, 2001.
100. FACTOR P, DUMASIVUS V, SALLDIAS F, BROWN LA, AND SZNAJDER JI. Adenovirus-mediated transfer of an Na^+/K^+ -ATPase beta1 subunit gene improves alveolar fluid clearance and survival in hyperoxic rats. *Hum Gene Ther* 11: 2231–2242, 2000.
101. FACTOR P, RIDGE K, ALVERDY J, AND SZNAJDER JI. Continuous enteral nutrition attenuates pulmonary edema in rats exposed to 100% oxygen. *J Appl Physiol* 89: 1759–1765, 2000.
102. FACTOR P, SALLDIAS F, RIDGE K, DUMASIVUS V, ZABNER J, JAFFE HA, BLANCO G, BARNARD M, MERCER R, PERRIN R, AND SZNAJDER JI. Augmentation of lung liquid clearance via adenovirus-mediated transfer of a Na,K-ATPase β_1 subunit gene. *J Clin Invest* 102: 1421–1430, 1998.
103. FANG X, FUKUDA N, BARBRY P, SARTORI C, VERKMAN AS, AND MATTHAY MA. Novel role for CFTR in fluid absorption from the distal airspaces of the lung. *J Gen Physiol* 119: 199–208, 2002.
104. FARMAN N, TALBOT CR, BOUCHER R, FAY M, CANESSA C, ROSSIER B, AND BONVALET JP. Noncoordinated expression of α -, β -, and γ -subunit mRNAs of epithelial Na^+ channel along rat respiratory tract. *Am J Physiol Cell Physiol* 272: C131–C141, 1997.
105. FENG ZP, CLARK RB, AND BERTHAUME Y. Identification of nonselective cation channels in cultured adult rat alveolar type II cells. *Am J Respir Cell Mol Biol* 9: 248–254, 1993.
106. FERAILLE E, CARRANZA ML, GONIN S, BEGUIN P, PEDEMONTE C, ROUSSELOT M, CAVERZASIO J, GEERING K, MARTIN PY, AND FAVRE H. Insulin-induced stimulation of Na^+/K^+ -ATPase activity in kidney proximal tubule cells depends on phosphorylation of the α -subunit at Tyr-10. *Mol Biol Cell* 10: 2847–2859, 1999.
107. FERRANDI M, MANUNTA P, BALZAN S, HAMLYN JM, BIANCHI G, AND FERRARI P. Ouabain-like factor quantification in mammalian tissues and plasma: comparison of two independent assays. *Hypertension* 30: 886–896, 1997.
108. FERRI C, BELLINI C, COASSIN S, DE ANGELIS C, PERRONE A, AND SANTUCCI A. Plasma endogenous digoxin-like substance levels are dependent on blood O_2 in man. *Clin Sci* 87: 447–451, 1994.
109. FESENKO EE, KOLESNIKOV SS, AND LYUBARSKY AL. Induction by cyclic GMP of cationic conductance in plasma membrane of retinal rod outer segment. *Nature* 313: 310–313, 1985.
110. FINLEY N, NORLIN A, BAINES DL, AND FOLKESSON HG. Alveolar epithelial fluid clearance is mediated by endogenous catecholamines at birth in guinea pigs. *J Clin Invest* 101: 972–981, 1998.
111. FIRSOV D, GAUTSCHI I, MERILLAT AM, ROSSIER BC, AND SCHILD L. The heterotetrameric architecture of the epithelial sodium channel (ENaC). *EMBO J* 17: 344–352, 1998.
112. FOLKESSON HG, MATTHAY MA, FRIGERI A, AND VERKMAN AS. Transepithelial water permeability in microperfused distal airways: evidence for channel-mediated water transport. *J Clin Invest* 97: 664–671, 1996.
113. FOLKESSON HG, MATTHAY MA, HASEGAWA H, KHERADMAND F, AND VERKMAN AS. Transcellular water transport in lung alveolar epithelium through mercury-sensitive water channels. *Proc Natl Acad Sci USA* 91: 4970–4974, 1994.
114. FOLKESSON HG, MATTHAY MA, WESTRÖM BR, KIM KJ, KARLSSON BW, AND HASTINGS RH. Alveolar epithelial clearance of protein. *J Appl Physiol* 80: 1431–1445, 1996.
115. FOLKESSON HG, NITENBERG G, OLIVER BL, JAYR C, ALBERTINE KH, AND MATTHAY MA. Upregulation of alveolar epithelial fluid transport after subacute lung injury in rats from bleomycin. *Am J Physiol Lung Cell Mol Physiol* 275: L478–L490, 1998.
116. FOLKESSON HG, NORLIN A, WANG Y, ABEDINPOUR P, AND MATTHAY MA. Dexamethasone and thyroid hormone pretreatment upregulate alveolar epithelial fluid clearance in adult rats. *J Appl Physiol* 88: 416–424, 2000.
117. FOLKESSON HG, NYGREN J, NIELSEN SO, AND NORLIN A. Evidence for chloride transport in adult and developing guinea pig lung (Abstract). *FASEB J* 14: A128, 2000.
118. FOLKESSON HG, PITTE JF, NITENBERG G, AND MATTHAY MA. Transforming growth factor- α increases alveolar liquid clearance in anesthetized ventilated rats. *Am J Physiol Lung Cell Mol Physiol* 271: L236–L244, 1996.
119. FOLKESSON HG, WESTRÖM BR, AND KARLSSON BW. Permeability of the respiratory tract to different-sized macromolecules after intratracheal instillation in young and adult rats. *Acta Physiol Scand* 139: 347–354, 1990.
120. FRANK JA, WANG Y, OSORIO O, AND MATTHAY MA. β -Adrenergic agonist therapy accelerates the resolution of hydrostatic pulmonary edema in sheep and rats. *J Appl Physiol* 89: 1255–1265, 2000.
121. FUKUDA N, FOLKESSON HG, AND MATTHAY MA. Relationship of interstitial fluid volume to alveolar fluid clearance in mice: ventilated versus in situ studies. *J Appl Physiol* 89: 672–679, 2000.
122. FUKUDA N, JAYR C, LAZRAK A, WANG Y, LUCAS R, MATALON S, AND MATTHAY MA. Mechanisms of TNF- α stimulation of amiloride-sensitive sodium transport across alveolar epithelium. *Am J Physiol Lung Cell Mol Physiol* 280: L1258–L1265, 2001.
123. GAILLARD D, HINNRSKY J, COSCOY S, HOFMAN P, MATTHAY MA, PUCHELLE E, AND BARBRY P. Early expression of β - and γ -subunits of epithelial sodium channel during human airway development. *Am J Physiol Lung Cell Mol Physiol* 278: L177–L184, 2000.
124. GARAT C, CARTER EP, AND MATTHAY MA. New in situ mouse model to quantify alveolar epithelial fluid clearance. *J Appl Physiol* 84: 1763–1767, 1998.
125. GARAT C, MEIGNAN M, MATTHAY MA, LUO DF, AND JAYR C. Alveolar epithelial fluid clearance mechanisms are intact after moderate hyperoxic lung injury in rats. *Chest* 111: 1381–1388, 1997.
126. GARTY H AND PALMER LG. Epithelial sodium channels: function, structure, and regulation. *Physiol Rev* 77: 359–396, 1997.
127. GILLIE DJ, PACE AJ, COAKLEY RJ, KOLLER BH, AND BARKER PM. Liquid and ion transport by fetal airway and lung epithelia of mice deficient in sodium-potassium-2-chloride transporter. *Am J Respir Cell Mol Biol* 25: 14–20, 2001.
128. GONZALEZ-FLECHA B, EVELSON P, RIDGE K, AND SZNAJDER JI. Hydrogen peroxide increases Na^+/K^+ -ATPase function in alveolar type II cells. *Biochim Biophys Acta* 1290: 46–52, 1996.
129. GOODMAN BE. Lung fluid clearance. In: *Fluid and Solute Transport in the Air Spaces of the Lungs*, edited by Effros RM and Chang HK. New York: Dekker, 1993, p. 139–183.
130. GOODMAN BE, ANDERSON JL, AND CLEMENS JW. Evidence for regulation of sodium transport from airspace to vascular space by cAMP. *Am J Physiol Lung Cell Mol Physiol* 257: L86–L93, 1989.
131. GOODMAN BE AND CRANDALL ED. Dome formation in primary cultured monolayers of alveolar epithelial cells. *Am J Physiol Cell Physiol* 243: C96–C100, 1982.
132. GOODMAN BE, FLEISCHER RD, AND CRANDALL ED. Evidence for active Na^+ transport by cultured monolayers of pulmonary alveolar epithelial cells. *Am J Physiol Cell Physiol* 245: C78–C83, 1983.
133. GOODMAN BE, KIM KJ, AND CRANDALL ED. Evidence for active sodium transport across alveolar epithelium of isolated rat lung. *J Appl Physiol* 62: 2460–2466, 1987.
134. GORIN AB AND STEWART PA. Differential permeability of endothelial

- and epithelial barriers to albumin flux. *J Appl Physiol* 47: 1315–1324, 1979.
135. GRIMME JD, LANE SM, AND MARON MB. Alveolar liquid clearance in multiple nonperfused canine lung lobes. *J Appl Physiol* 82: 348–353, 1997.
 136. GUERRERO C, LECUONA E, PESCE L, RIDGE KM, AND SZNAJDER JI. Dopamine regulates Na-K-ATPase in alveolar epithelial cells via MAPK-ERK-dependent mechanisms. *Am J Physiol Lung Cell Mol Physiol* 281: L79–L85, 2001.
 137. GUERY BPH, MASON CM, DOBARD EP, BEAUCAIRE G, SUMMER WR, AND NELSON S. Keratinocyte growth factor increases transalveolar sodium reabsorption in normal and injured rat lungs. *Am J Respir Crit Care Med* 155: 1777–1784, 1997.
 138. GUGGINO WB AND GUGGINO SE. Amiloride-sensitive sodium channels contribute to the woes of the flu. *Proc Natl Acad Sci USA* 97: 9827–9829, 2000.
 139. GUMBNER BM. Breaking through the tight junction barrier. *J Cell Biol* 123: 1631–1633, 1993.
 140. GUO J, YI ES, HAVILL AM, SAROSI I, WHITCOMB L, YIN S, MIDDLETON SC, PIGUET P, AND ULLICH TR. Intravenous keratinocyte growth factor protects against experimental pulmonary injury. *Am J Physiol Lung Cell Mol Physiol* 275: L800–L805, 1998.
 141. GUO Y, DUVAL MD, CROW JP, AND MATALON S. Nitric oxide inhibits Na⁺ absorption across cultured alveolar type II monolayers. *Am J Physiol Lung Cell Mol Physiol* 274: L369–L377, 1998.
 142. GUTKOWSKA J, NEMER M, SOLE MJ, DROUIN J, AND SIROIS P. Lung is an important source of atrial natriuretic factor in experimental cardiomyopathy. *J Clin Invest* 83: 1500–1504, 1989.
 143. HADDAD IY, ISCHIROPOULOS H, HOLM BA, BECKMAN JS, BAKER JR, AND MATALON S. Mechanisms of peroxynitrite-induced injury to pulmonary surfactants. *Am J Physiol Lung Cell Mol Physiol* 265: L555–L564, 1993.
 144. HADDAD IY, PATAKI G, HU P, GALLIANI C, BECKMAN JS, AND MATALON S. Quantitation of nitrotyrosine levels in lung sections of patients and animals with acute lung injury. *J Clin Invest* 94: 2407–2413, 1994.
 145. HAKIM TS, MINNEAR FC, VAN DER ZEE H, BARIE PS, AND MALIK AB. Adrenoreceptor control of lung fluid and protein exchange. *J Appl Physiol* 51: 68–72, 1981.
 146. HARDIMAN KM, LINDSEY JR, AND MATALON S. Lack of amiloride-sensitive transport across alveolar and respiratory epithelium of iNOS(–/–) mice in vivo. *Am J Physiol Lung Cell Mol Physiol* 281: L722–L731, 2001.
 147. HARDIMAN KM AND MATALON S. Modification of sodium transport and alveolar fluid clearance by hypoxia. Mechanisms and physiological implications (Editorial). *Am J Respir Cell Mol Biol* 25: 538–541, 2001.
 148. HARRIS ZL, RIDGE KM, GONZALEZ-FLECHA B, GOTTLIEB L, ZUCKER A, AND SZNAJDER JI. Hyperbaric oxygenation upregulates rat lung Na,K-ATPase. *Eur Respir J* 9: 472–477, 1996.
 149. HASKELL JF, YUE G, BENOS DJ, AND MATALON S. Upregulation of sodium conductive pathways in alveolar type II cells in sublethal hyperoxia. *Am J Physiol Lung Cell Mol Physiol* 266: L30–L37, 1994.
 150. HASTINGS RH, GRADY M, SAKUMA T, AND MATTHAY MA. Clearance of different-sized proteins from the alveolar space in humans and rabbits. *J Appl Physiol* 73: 1310–1316, 1992.
 151. HASTINGS RH, WRIGHT JR, ALBERTINE KH, CIRIALES R, AND MATTHAY MA. Effect of endocytosis inhibitors on alveolar clearance of albumin, immunoglobulin G, and SP-A in rabbits. *Am J Physiol Lung Cell Mol Physiol* 266: L544–L552, 1994.
 152. HOFFERT JD, LEITCH V, AGRE P, AND KING LS. Hypertonic induction of aquaporin-5 expression through an ERK-dependent pathway. *J Biol Chem* 275: 9070–9077, 2000.
 153. HU P, ISCHIROPOULOS H, BECKMAN JS, AND MATALON S. Peroxynitrite inhibition of oxygen consumption and sodium transport in alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 266: L628–L634, 1994.
 154. HUMMLER E, BARKER P, GATZY J, BEERMANN F, VERDUMO C, SCHMIDT A, BOUCHER R, AND ROSSIER BC. Early death due to defective neonatal lung liquid clearance in α ENaC-deficient mice. *Nat Genet* 12: 325–328, 1996.
 155. ICARD P AND SAUMON G. Alveolar sodium and liquid transport in mice. *Am J Physiol Lung Cell Mol Physiol* 277: L1232–L1238, 1999.
 156. ILLEK B, FISCHER H, KREUSEL KM, HEGEL U, AND CLAUSS W. Volume-sensitive basolateral K⁺ channels in HT-29/B6 cells: block by lidocaine, quinidine, NPPB, and Ba²⁺. *Am J Physiol Cell Physiol* 263: C674–C683, 1992.
 157. IMAMURA T, OHNUMA N, IWASA F, FURUYA M, HAYASHI Y, INOMATA N, ISHIIHARA T, AND NOGUCHI T. Protective effect of α -human atrial natriuretic polypeptide (α -hANP) on chemical-induced pulmonary edema. *Life Sci* 42: 403–414, 1988.
 158. INGBAR DH, DUVICK S, SAVICK SK, SCHELLHASE DE, DETTERDING R, JAMIESON JD, AND SHANNON JM. Developmental changes of fetal rat lung Na-K-ATPase after maternal treatment with dexamethasone. *Am J Physiol Lung Cell Mol Physiol* 272: L665–L672, 1997.
 159. INGLIS SK, CORBOZ MR, TAYLOR AE, AND BALLARD ST. Regulation of ion transport across porcine distal bronchi. *Am J Physiol Lung Cell Mol Physiol* 270: L289–L297, 1996.
 160. ITO Y, NIISATO N, O'BRODOVICH H, AND MARUNAKA Y. The effect of brefeldin A on terbutaline-induced sodium absorption in fetal rat distal lung epithelium. *Pflügers Arch* 434: 492–494, 1997.
 161. JAFFE HA, OLIVERA W, RIDGE K, YEATES D, AND SZNAJDER JI. In vivo administration of EGF increases Na⁺,K⁺-ATPase activity in alveolar type II cells. *FASEB J* 8: A141, 1994.
 162. JAIN L, CHEN XJ, MALIK B, AL-KHALILI O, AND EATON DC. Antisense oligonucleotides against the α -subunit of ENaC decrease lung epithelial cation-channel activity. *Am J Physiol Lung Cell Mol Physiol* 276: L1046–L1051, 1999.
 163. JAIN L, CHEN XJ, RAMOSEVAC S, BROWN LA, AND EATON DC. Expression of highly selective sodium channels in alveolar type II cells is determined by culture conditions. *Am J Physiol Lung Cell Mol Physiol* 280: L646–L658, 2001.
 164. JAYR C, GARAT C, MEIGNAN M, PITTET JF, ZELTER M, AND MATTHAY MA. Alveolar liquid and protein clearance in anesthetized ventilated rats. *J Appl Physiol* 76: 2636–2642, 1994.
 165. JIANG X, INGBAR DH, AND O'GRADY SM. Adrenergic stimulation of Na⁺ transport across alveolar epithelial cells involves activation of apical Cl[–] channels. *Am J Physiol Cell Physiol* 275: C1610–C1620, 1998.
 166. JIANG X, INGBAR DH, AND O'GRADY SM. Adrenergic regulation of ion transport across adult alveolar epithelial cells: effects on Cl[–] channel activation and transport function in cultures with an apical air interface. *J Membr Biol* 181: 195–204, 2001.
 167. JOHN T, VOGEL S, MINSHALL R, RIDGE K, TIRUPPATHI C, AND MALIK A. Evidence for the role of alveolar epithelial gp60 in active transalveolar albumin transport in the rat lung. *J Physiol* 533: 547–559, 2001.
 168. JOHNSON M, WIDDICOMBE J, ALLEN L, BARBRY P, AND DOBBS L. Alveolar epithelial type I cells contain transport proteins and transport sodium, supporting an active role for type I cells in regulation of lung liquid homeostasis. *Proc Natl Acad Sci USA* 99: 1966–1971, 2002.
 169. JOSEPH D, TIRMIZI O, ZHANG XL, CRANDALL ED, AND LUBMAN RL. Polarity of alveolar epithelial cell acid-base permeability. *Am J Physiol Lung Cell Mol Physiol* 282: L675–L682, 2002.
 170. JUNOR RWJ, BENJAMIN AR, ALEXANDROU D, GUGGINO SE, AND WALTERS DV. A novel role for cyclic nucleotide-gated cation channels in lung liquid homeostasis in sheep. *J Physiol* 520: 255–260, 1999.
 171. KASPER M, REIMANN T, HEMPEL U, WENZEL KW, BIERHAUS A, SCHUH D, DIMMER V, HAROSKE G, AND MULLER M. Loss of caveolin expression in type I pneumocytes as an indicator of subcellular alterations during lung fibrogenesis. *Histochem Cell Biol* 109: 41–48, 1998.
 172. KEMP PJ, BOROK Z, KIM KJ, LUBMAN RL, DANTO SI, AND CRANDALL ED. Epidermal growth factor regulation in adult rat alveolar type II cells of amiloride-sensitive cation channels. *Am J Physiol Cell Physiol* 277: C1058–C1065, 1999.
 173. KEREM E, BISTRITZER T, HANUKOGLU A, HOFMANN T, ZHOU Z, BENNETT W, MACLAUGHLIN E, BARKER P, NASH M, QUITTEL L, BOUCHER R, AND KNOWLES MR. Pulmonary epithelial sodium-channel dysfunction and excess airway liquid in pseudohypoaldosteronism. *N Engl J Med* 341: 156–162, 1999.
 174. KERR JS, REICHERTER J, AND FISHER AB. 2-Deoxy-D-glucose uptake by rat granular pneumocytes in primary culture. *Am J Physiol Cell Physiol* 243: C14–C19, 1982.
 175. KHIMENKO PL, BAGBY GJ, FUSELER J, AND TAYLOR AE. Tumor necrosis factor- α in ischemia and reperfusion injury in rat lungs. *J Appl Physiol* 85: 2005–2011, 1998.

176. KIM KJ, CHEEK JM, AND CRANDALL ED. Contribution of active Na^+ and Cl^- fluxes to net ion transport by alveolar epithelium. *Respir Physiol* 85: 245–256, 1991.
177. KIM KJ AND CRANDALL ED. Models for investigation of peptide and protein transport across cultured mammalian respiratory epithelial barriers. *Pharm Biotechnol* 8: 325–346, 1996.
178. KIM KJ, LEBON TR, SHINBANE JS, AND CRANDALL ED. Asymmetric [^{14}C]albumin transport across bullfrog alveolar epithelium. *J Appl Physiol* 59: 1290–1297, 1985.
179. KING LS AND AGRE P. Man is not a rodent. Aquaporins in the airways. *Am J Respir Cell Mol Biol* 24: 221–223, 2001.
180. KING LS, NIELSEN S, AND AGRE P. Aquaporin-1 water channel protein in lung. Ontogeny, steroid-induced expression, and distribution in rat. *J Clin Invest* 97: 2183–2191, 1996.
181. KING LS, NIELSEN S, AND AGRE P. Aquaporin-5 is expressed in rat type I pneumocytes (Abstract). *Am J Respir Crit Care Med* 153: A508, 1996.
182. KING LS, YASUI M, AND AGRE P. Aquaporins in health and disease. *Mol Med Today* 6: 60–65, 2000.
183. KLEYMAN TR, ERNST SA, AND COUPAYE-GERARD B. Arginine vasopressin and forskolin regulate apical cell surface expression of epithelial Na^+ channels in A6 cells. *Am J Physiol Renal Fluid Electrolyte Physiol* 266: F506–F511, 1994.
184. KOSARI F, SHENG S, LI J, MAK DO, FOSKETT JK, AND KLEYMAN TR. Subunit stoichiometry of the epithelial sodium channel. *J Biol Chem* 273: 13469–13474, 1998.
185. KREDA SM, GYNN MC, FENSTERMACHER DA, BOUCHER RC, AND GABRIEL SE. Expression and localization of epithelial aquaporins in the adult human lung. *Am J Respir Cell Mol Biol* 24: 224–234, 2001.
186. KROZOWSKI Z AND FUNDER JW. Mineralocorticoid receptors in the rat lung. *Endocrinology* 109: 1811–1813, 1981.
187. KUNZELMANN K, BEESLEY AH, KING NJ, KARUPIAH G, YOUNG JA, AND COOK DI. Influenza virus inhibits amiloride-sensitive Na^+ channels in respiratory epithelia. *Proc Natl Acad Sci USA* 97: 10282–10287, 2000.
188. KURAHASHI K, KAJIKAWA O, SAWA T, OHARA M, GROPPER MA, FRANK DW, MARTIN TR, AND WIENER-KRONISH JP. Pathogenesis of septic shock in *Pseudomonas aeruginosa* pneumonia. *J Clin Invest* 104: 743–750, 1999.
189. LAFFON M, JAYR C, BARBRY P, WANG Y, FOLKESSON HG, PITTET JF, CLERICI C, AND MATTHAY MA. Lidocaine induces a reversible decrease in alveolar epithelial fluid clearance in rats. *Anesthesiology* 96: 392–399, 2002.
190. LAFFON M, LU LN, MODELSKA K, MATTHAY MA, AND PITTET JF. α -Adrenergic blockade restores normal fluid transport capacity of alveolar epithelium after hemorrhagic shock. *Am J Physiol Lung Cell Mol Physiol* 277: L760–L768, 1999.
191. LANE SM, MAENDER KC, AWENDER NE, AND MARON MB. Adrenal epinephrine increases alveolar liquid clearance in a canine model of neurogenic pulmonary edema. *Am J Respir Crit Care Med* 158: 760–768, 1998.
192. LASNIER JM, WANGENSTEIN OD, SCHMITZ LS, GROSS CR, AND INGBAR DH. Terbutaline stimulates alveolar fluid resorption in hyperoxic lung injury. *J Appl Physiol* 81: 1723–1729, 1996.
193. LAZRAC A, NIELSEN VG, AND MATALON S. Mechanisms of increased Na^+ transport in A11 cells by cAMP: we agree to disagree and do more experiments. *Am J Physiol Lung Cell Mol Physiol* 278: L233–L238, 2000.
194. LAZRAC A, SAMANTA A, VENETSANO K, BARBRY P, AND MATALON S. Modification of biophysical properties of lung epithelial Na^+ channels by dexamethasone. *Am J Physiol Cell Physiol* 279: C762–C770, 2000.
195. LAZRAC A, THOME U, MYLES C, WARE J, CHEN L, VENGLARIK CJ, AND MATALON S. cAMP regulation of Cl^- and HCO_3^- secretion across rat fetal distal lung epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 282: L650–L658, 2002.
196. LECUONA E, GARCIA A, AND SZNAJDER JI. A novel role for protein phosphatase 2A in the dopaminergic regulation of Na,K-ATPase . *FEBS Lett* 481: 217–220, 2000.
197. LEI J, MARIASH CN, AND INGBAR DH. T_3 stimulation of alveolar epithelial Na,K-ATPase (Abstract). *FASEB J* 15: A860, 2001.
198. LE TULZO Y, SHENKAR R, KANEKO D, MOINE P, FANTUZZI G, DINARELLO CA, AND ABRAHAM E. Hemorrhage increases cytokine expression in lung mononuclear cells in mice: involvement of catecholamines in nuclear factor-kappaB regulation and cytokine expression. *J Clin Invest* 99: 1516–1524, 1997.
199. LI X, ANN D, CRANDALL ED, AND BOROK Z. Osmotic regulation of aquaporin-5 regulation in alveolar epithelial cells in primary culture (Abstract). *Am J Respir Crit Care Med* 163: A571, 2001.
200. LI XY, DONALDSON K, BROWN D, AND MACNEE W. The role of tumor necrosis factor in increased airspace epithelial permeability in acute lung inflammation. *Am J Respir Cell Mol Biol* 13: 185–195, 1995.
201. LIEBLER JM, BOROK Z, KIM KJ, AND CRANDALL ED. Immunoreactive β_2 adrenergic receptors are expressed in alveolar type I cells (Abstract). *Am J Respir Crit Care Med* 163: A570, 2001.
202. LILEY HG, ERTSEY R, GONZALES LW, ODOM MW, HAWGOOD S, DOBBS LG, AND BALLARD PL. Synthesis of surfactant components by cultured type II cells from human lung. *Biochim Biophys Acta* 961: 86–95, 1988.
203. LIN W, FINGER TE, ROSSIER BC, AND KINNAMON SC. Epithelial Na^+ channel subunits in rat taste cells: localization and regulation by aldosterone. *J Comp Neurol* 405: 406–420, 1999.
204. LINGUEGLIA E, RENARD S, WALDMANN R, VOILLEY N, CHAMPIGNY G, PLASS H, LAZDUNSKI M, AND BARBRY P. Different homologous subunits of the amiloride-sensitive Na^+ channel are differently regulated by aldosterone. *J Biol Chem* 269: 13736–13739, 1994.
205. LOFFING J, ZECEVIC M, FERAILLE E, KAISLING B, ASHER C, ROSSIER BC, FIRESTONE GL, PEARCE D, AND VERREY F. Aldosterone induces rapid apical translocation of ENaC in early portion of renal collecting system: possible role of SGK. *Am J Physiol Renal Physiol* 280: F675–F682, 2001.
206. MA T, FUKUDA N, SONG Y, MATTHAY MA, AND VERKMAN AS. Lung fluid transport in aquaporin-5 knockout mice. *J Clin Invest* 105: 93–100, 2000.
207. MA T, SONG Y, GILLESPIE A, CARLSON EJ, EPSTEIN CJ, AND VERKMAN AS. Defective secretion of saliva in transgenic mice lacking aquaporin-5 water channels. *J Biol Chem* 274: 20071–20074, 1999.
208. MAIRBÄURL H, MAYER K, KIM KJ, BOROK Z, BÄRTSCH P, AND CRANDALL ED. Hypoxia decreases active Na transport across primary rat alveolar epithelial cell monolayers. *Am J Physiol Lung Cell Mol Physiol* 282: L659–L665, 2002.
209. MAIRBÄURL H, WODOPIA R, ECKES S, SCHULZ S, AND BÄRTSCH P. Impairment of cation transport in A549 cells and rat alveolar epithelial cells by hypoxia. *Am J Physiol Lung Cell Mol Physiol* 274: L797–L806, 1998.
210. MARON MB. Dose-response relationship between plasma epinephrine concentration and alveolar liquid clearance in dogs. *J Appl Physiol* 85: 1702–1707, 1998.
211. MARUNAKA Y, NISATO N, AND ITO Y. Beta agonist regulation of sodium transport in fetal lung epithelium: roles of cell volume, cytosolic chloride and protein tyrosine kinase. *J Korean Med Sci* 15 Suppl: S42–S43, 2000.
212. MARUNAKA Y, NISATO N, O'BRODOVICH H, AND EATON DC. Regulation of an amiloride-sensitive Na^+ -permeable channel by a β_2 -adrenergic agonist, cytosolic Ca^{2+} and Cl^- in fetal rat alveolar epithelium. *J Physiol* 515: 669–683, 1999.
213. MARUNAKA Y, TOHDA H, HAGIWARA N, AND O'BRODOVICH H. Cytosolic Ca^{2+} -induced modulation of ion selectivity and amiloride sensitivity of a cation channel and beta agonist action in fetal lung epithelium. *Biochem Biophys Res Commun* 187: 648–656, 1992.
214. MASON RJ, WILLIAMS MC, WIDDICOMBE JH, SANDERS MJ, MISFELDT DS, AND BERRY LCJ. Transepithelial transport by pulmonary alveolar type II cells in primary culture. *Proc Natl Acad Sci USA* 79: 6033–6037, 1982.
215. MATALON S. Mechanisms and regulation of ion transport in adult mammalian alveolar type II pneumocytes. *Am J Physiol Cell Physiol* 261: C727–C738, 1991.
216. MATALON S, BENOS DJ, AND JACKSON RM. Biophysical and molecular properties of amiloride-inhibitable Na^+ channels in alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 271: L1–L22, 1996.
217. MATALON S, BRIDGES RJ, AND BENOS DJ. Amiloride-inhibitable Na^+ conductive pathways in alveolar type II pneumocytes. *Am J Physiol Lung Cell Mol Physiol* 260: L90–L96, 1991.
218. MATALON S, HU P, ISCHIROPOULOS H, AND BECKMAN JS. Peroxynitrite

- inhibition of oxygen consumption and ion transport in alveolar type II pneumocytes. *Chest* 105: 74S, 1994.
219. MATALON S, KIRK KL, BUBIEN JK, OH Y, HU P, YUE G, SHOEMAKER R, CRAIGOE EJ JR, AND BENOS DJ. Immunocytochemical and functional characterization of Na⁺ conductance in adult alveolar pneumocytes. *Am J Physiol Lung Cell Mol Physiol* 262: C1228–C1238, 1992.
 220. MATALON S AND O'BRODOVICH H. Sodium channels in alveolar epithelial cells: molecular characterization, biophysical properties, and physiological significance. *Annu Rev Physiol* 61: 627–661, 1999.
 221. MATSUKAWA Y, YAMAHARA H, YAMASHITA F, LEE VH, CRANDALL ED, AND KIM KJ. Rates of protein transport across rat alveolar epithelial cell monolayers. *J Drug Target* 7: 335–342, 2000.
 222. MATSUMOTO PS, OHARA A, DUCHATELLE P, AND EATON DC. Tyrosine kinase regulates epithelial sodium transport in A6 cells. *Am J Physiol Lung Cell Mol Physiol* 264: C246–C250, 1993.
 223. MATSUSHITA K, MCCRAY PB JR, SIGMUND RD, WELSH MJ, AND STOKES JB. Localization of epithelial sodium channel subunit mRNAs in adult rat lung by in situ hybridization. *Am J Physiol Lung Cell Mol Physiol* 271: L332–L339, 1996.
 224. MATTHAY MA. Pathophysiology of pulmonary edema. *Clin Chest Med* 6: 301–314, 1985.
 225. MATTHAY MA. Severe sepsis—a new treatment with both anticoagulant and anti-inflammatory properties. *N Engl J Med* 344: 759–762, 2001.
 226. MATTHAY MA, BERTHIAUME Y, AND STAUB NC. Long-term clearance of liquid and protein from the lungs of unanesthetized sheep. *J Appl Physiol* 59: 928–934, 1985.
 227. MATTHAY MA, FOLKESSON HG, CAMPAGNA A, AND KHERADMAND F. Alveolar epithelial barrier and acute lung injury. *New Horiz* 1: 613–622, 1993.
 228. MATTHAY MA, FOLKESSON HG, AND VERKMAN AS. Salt and water transport across alveolar and distal airway epithelium in the adult lung. *Am J Physiol Lung Cell Mol Physiol* 270: L487–L503, 1996.
 229. MATTHAY MA, LANDOLT CC, AND STAUB NC. Differential liquid and protein clearance from the alveoli of anesthetized sheep. *J Appl Physiol* 53: 96–104, 1982.
 230. MATTHAY MA AND WIENER-KRONISH JP. Intact epithelial barrier function is critical for the resolution of alveolar edema in humans. *Am Rev Respir Dis* 142: 1250–1257, 1990.
 231. MAY A, PUOTI A, GAEGELER HP, HORISBERGER JD, AND ROSSIER BC. Early effect of aldosterone on the rate of synthesis of the epithelial sodium channel alpha subunit in A6 renal cells. *J Am Soc Nephrol* 8: 1813–1822, 1997.
 232. McDONALD FJ, SNYDER PM, MCCRAY PB JR, AND WELSH MJ. Cloning, expression, and tissue distribution of a human amiloride-sensitive Na⁺ channel. *Am J Physiol Lung Cell Mol Physiol* 266: L728–L734, 1994.
 233. McDONALD FJ, YANG B, HRSTKA RF, DRUMMOND HA, TARR DE, MCCRAY PB JR, STOKES JB, WELSH MJ, AND WILLIAMSON RA. Disruption of the β subunit of the epithelial Na⁺ channel in mice: hyperkalemia and neonatal death associated with a pseudohypoaldosteronism phenotype. *Proc Natl Acad Sci USA* 96: 1727–1731, 1999.
 234. McDONALD JV JR, GONZALES LW, BALLARD PL, PITHA J, AND ROBERTS JM. Lung β -adrenoreceptor blockade affects perinatal surfactant release but not lung water. *J Appl Physiol* 60: 1727–1733, 1986.
 235. McDONOUGH AA, GEERING K, AND FARLEY RA. The sodium pump needs its β subunit. *FASEB J* 4: 1598–1605, 1990.
 236. MCGRAW DW, FUKUDA N, JAMES PF, FORBES SL, WOO AL, LINGREL JB, WITTE DP, MATTHAY MA, AND LIGGETT SB. Targeted transgenic expression of β_2 -adrenergic receptors to type II cells increases alveolar fluid clearance. *Am J Physiol Lung Cell Mol Physiol* 281: L895–L903, 2001.
 237. McNICHOLAS CM AND CANESSA CM. Diversity of channels generated by different combinations of epithelial sodium channel subunits. *J Gen Physiol* 109: 681–692, 1997.
 238. MICHAUT P, PLANES C, ESCOUBET B, CLEMENT A, AMIEL C, AND CLERICI C. Rat lung alveolar type II cell line maintains sodium transport characteristics of primary culture. *J Cell Physiol* 169: 78–86, 1996.
 239. MINAKATA Y, SUZUKI S, GRYGORCZYK C, DAGENAIS A, AND BERTHIAUME Y. Impact of β -adrenergic agonist on Na⁺ channel and Na⁺-K⁺-ATPase expression in alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 275: L414–L422, 1998.
 240. MODELSKA K, MATTHAY MA, BROWN LAS, DEUTCH E, LU LN, AND PITTET JF. Inhibition of β -adrenergic-dependent alveolar epithelial clearance by oxidant mechanisms after hemorrhagic shock. *Am J Physiol Lung Cell Mol Physiol* 276: L844–L857, 1999.
 241. MODELSKA K, MATTHAY MA, McELROY MC, AND PITTET JF. Upregulation of alveolar liquid clearance after fluid resuscitation for hemorrhagic shock in rats. *Am J Physiol Lung Cell Mol Physiol* 273: L305–L314, 1997.
 242. MOLLIEUX S, CRESTANI B, DUREUIL B, BASTIN J, ROLLAND C, AUBIER M, AND DESMONTS JM. Effects of halothane on surfactant biosynthesis by rat alveolar type II cells in primary culture. *Anesthesiology* 81: 668–676, 1994.
 243. MORENO JJ AND PRYOR WA. Inactivation of α_1 -proteinase inhibitor by peroxynitrite. *Chem Res Toxicol* 5: 425–431, 1992.
 244. MORGAN EE, HODNICHAK CM, STADER SM, MAENDER KC, BOJA JW, FOLKESSON HG, AND MARON MB. Prolonged isoproterenol infusion impairs the ability of β_2 -agonists to increase alveolar liquid clearance. *Am J Physiol Lung Cell Mol Physiol* 282: L666–L674, 2002.
 245. NAKAGAWA Y, GAMMICHIA J, PURUSHOTHAM KR, SCHNEIDER CA, AND HUMPHREYS-BEHER MG. Epidermal growth factor activation of rat parotid gland adenylate cyclase and mediation by a GTP-binding regulatory protein. *Biochem Pharmacol* 42: 2333–2340, 1991.
 246. NAKAMURA T AND GOLD GH. A cyclic nucleotide-gated conductance in olfactory receptor cilia. *Nature* 325: 442–444, 1987.
 247. NEWMAN GR, CAMPBELL L, VON RUHLAND C, JASANI B, AND GUMBLETON M. Caveolin and its cellular and subcellular immunolocalisation in lung alveolar epithelium: implications for alveolar epithelial type I cell function. *Cell Tissue Res* 295: 111–120, 1999.
 248. NEWMAN V, GONZALEZ RF, MATTHAY MA, AND DOBBS LG. A novel alveolar type I cell-specific biochemical marker of human acute lung injury. *Am J Respir Crit Care Med* 161: 990–995, 2000.
 249. NICHOLSON DJ, PAPKA RE, AND MARON MB. Distribution of β_2 -adrenoreceptors (β_2 -ARS) in the alveolar septum of the rat (Abstract). *FASEB J* 14: A129, 2000.
 250. NICI L, DOWIN R, GILMORE-HEBERT M, JAMIESON JD, AND INGBAR DH. Response of rat type II pneumocyte Na,K-ATPase to hyperoxic injury. *Chest* 31S–33S, 1991.
 251. NICI L, DOWIN R, GILMORE-HEBERT M, JAMIESON JD, AND INGBAR DH. Upregulation of rat lung Na-K-ATPase during hyperoxic injury. *Am J Physiol Lung Cell Mol Physiol* 261: L307–L314, 1991.
 252. NIELSEN S, SMITH BL, CHRISTENSEN EI, AND AGRE P. Distribution of the aquaporin CHIP in secretory and resorptive epithelia and capillary endothelia. *Proc Natl Acad Sci USA* 90: 7275–7279, 1993.
 253. NIELSEN VG, BAIRD MS, GEARY BT, AND MATALON S. Halothane does not decrease amiloride-sensitive alveolar fluid clearance in rabbits. *Anesth Analg* 90: 1445–1449, 2000.
 254. NIELSEN VG, DuVALL MD, BAIRD MS, AND MATALON S. cAMP activation of chloride and fluid secretion across the rabbit alveolar epithelium. *Am J Physiol Lung Cell Mol Physiol* 275: L1127–L1133, 1998.
 255. NIELSON DW. Electrolyte composition of pulmonary alveolar subphase in anesthetized rabbits. *J Appl Physiol* 60: 972–979, 1986.
 256. NORD EP, BROWN SES, AND CRANDALL ED. Characterization of Na⁺-H⁺ antiport in type II alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 252: C490–C498, 1987.
 257. NORLIN A, BAINES DL, AND FOLKESSON HG. Role of endogenous cortisol in basal liquid clearance from distal air spaces in adult guinea-pigs. *J Physiol* 519: 261–272, 1999.
 258. NORLIN A, FINLEY N, ABEDINPOUR P, AND FOLKESSON HG. Alveolar liquid clearance in the anesthetized ventilated guinea pig. *Am J Physiol Lung Cell Mol Physiol* 274: L235–L243, 1998.
 259. NORLIN A AND FOLKESSON HG. Alveolar liquid clearance in late gestational guinea pigs after induction of labor: mechanisms and regulation. *Am J Physiol Lung Cell Mol Physiol* 280: L606–L616, 2001.
 260. NORLIN A AND FOLKESSON HG. Calcium-dependent stimulation of alveolar fluid clearance in near-term fetal guinea pigs. *Am J Physiol Lung Cell Mol Physiol* 282: L642–L649, 2002.
 261. NORLIN A, LU LN, GUGGINO SE, MATTHAY MA, AND FOLKESSON HG. Contribution of amiloride-insensitive pathways to alveolar fluid clearance in adult rats. *J Appl Physiol* 90: 1489–1496, 2001.
 262. O'BRODOVICH H. Fetal lung liquid secretion. Insights using the tools

- of inhibitors and genetic knock-out experiments. *Am J Respir Cell Mol Biol* 25: 8–10, 2001.
263. O'BRODOVICH H. Pulmonary edema fluid movement within the lung (Editorial focus). *Am J Physiol Lung Cell Mol Physiol* 281: L1324–L1326, 2001.
 264. O'BRODOVICH H, CANESSA C, UEDA J, RAFII B, ROSSIER BC, AND EDELSON J. Expression of the epithelial Na⁺ channel in the developing rat lung. *Am J Physiol Cell Physiol* 264: C491–C496, 1993.
 265. O'BRODOVICH H, HANNAM V, AND RAFII B. Sodium channel but neither Na⁺-H⁺ nor Na-glucose symport inhibitors slow neonatal lung water clearance. *Am J Respir Cell Mol Biol* 5: 377–384, 1991.
 266. O'BRODOVICH H, HANNAM V, SEEAR M, AND MULLEN JB. Amiloride impairs lung water clearance in newborn guinea pigs. *J Appl Physiol* 68: 1758–1762, 1990.
 267. O'GRADY SM, JIANG X, AND INGBAR DH. Cl[−] channel activation is necessary for stimulation of Na transport in adult alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 278: L239–L244, 2000.
 268. OH YS, SAXENA S, AND WARNOCK DG. αENaC: leading the charge. *J Clin Invest* 104: 849–850, 1999.
 269. OLIVERA W, RIDGE K, WOOD LD, AND SZNAJDER JI. ANF decreases active sodium transport and increases alveolar epithelial permeability in rats. *J Appl Physiol* 75: 1581–1586, 1993.
 270. OLIVERA W, RIDGE K, WOOD LD, AND SZNAJDER JI. Active sodium transport and alveolar epithelial Na-K-ATPase increase during subacute hyperoxia in rats. *Am J Physiol Lung Cell Mol Physiol* 266: L577–L584, 1994.
 271. OLIVERA WG, CICCOLELLA DE, BARQUIN N, RIDGE KM, RUTSCHMAN DH, YEATES DB, AND SZNAJDER JI. Aldosterone regulates Na,K-ATPase and increases lung edema clearance in rats. *Am J Respir Crit Care Med* 161: 567–573, 2000.
 272. OLIVERA WG, RIDGE KM, AND SZNAJDER JI. Lung liquid clearance and Na,K-ATPase during acute hyperoxia and recovery in rats. *Am J Respir Crit Care Med* 152: 1229–1234, 1995.
 273. OLVER RE. Ion transport and water flow in the mammalian lung. *Ciba Found Symp*: 199–200, 1976.
 274. ORSER BA, BERTLIK M, FEDORKO L, AND O'BRODOVICH H. Cation selective channel in fetal alveolar type II epithelium. *Biochim Biophys Acta* 1094: 19–26, 1991.
 275. OTULAKOWSKI G, FREYWALD T, WEN Y, AND O'BRODOVICH H. Translational activation and repression by distinct elements within the 5'-UTR of ENaC α-subunit mRNA. *Am J Physiol Lung Cell Mol Physiol* 281: L1219–L1231, 2001.
 276. OTULAKOWSKI G, RAFII B, BREMNER HR, AND O'BRODOVICH H. Structure and hormone responsiveness of the gene encoding the α-subunit of the rat amiloride-sensitive epithelial sodium channel. *Am J Respir Cell Mol Biol* 20: 1028–1040, 1999.
 277. PACHA J, FRINDT G, ANTONIAN L, SILVER RB, AND PALMER LG. Regulation of Na channels of the rat cortical collecting tubule by aldosterone. *J Gen Physiol* 102: 25–42, 1993.
 278. PACITTI AJ, INOUE Y, AND SOUBA WW. Tumor necrosis factor stimulates amino acid transport in plasma membrane vesicles from rat liver. *J Clin Invest* 91: 474–483, 1993.
 279. PARKER JC, CRAIN M, GRIMBERT F, ROTILI G, AND TAYLOR AE. Total lung lymph flow and fluid compartmentation in edematous dog lungs. *J Appl Physiol* 51: 1268–1277, 1981.
 280. PESCE L, GUERRERO C, COMELLAS A, RIDGE KM, AND SZNAJDER JI. β-Agonists regulate Na,K-ATPase via novel MAPK/ERK and rapamycin-sensitive pathways. *FEBS Lett* 486: 310–314, 2000.
 281. PHALEN RF AND OLDHAM MJ. Airway structures. Tracheobronchial airway structure as revealed by casting techniques. *Am Rev Respir Dis* 128 Suppl: S1–S4, 1983.
 282. PINKERTON KE, BARRY BE, O'NEIL JJ, RAUB JA, PRATT PC, AND CRAPO JD. Morphologic changes in the lung during the lifespan of Fischer 344 rats. *Am J Anat* 164: 155–174, 1982.
 283. PITKANEN O, TANSWELL AK, DOWNEY G, AND O'BRODOVICH H. Increased Po₂ alters the bioelectric properties of fetal distal lung epithelium. *Am J Physiol Lung Cell Mol Physiol* 270: L1060–L1066, 1996.
 284. PITKANEN OM, SMITH D, O'BRODOVICH H, AND OTULAKOWSKI G. Expression of α-, β-, and γ-hENaC mRNA in the human nasal, bronchial, and distal lung epithelium. *Am J Respir Crit Care Med* 163: 273–276, 2001.
 285. PITTET JF, BRENNER TJ, MODELSKA K, AND MATTHAY MA. Alveolar liquid clearance is increased by endogenous catecholamines in hemorrhagic shock in rats. *J Appl Physiol* 81: 830–837, 1996.
 286. PITTET JF, GRIFFITHS MJ, GEISER T, KAMINSKI N, DALTON SL, HUANG X, BROWN LA, GOTWALS PJ, KOTELANSKY VE, MATTHAY MA, AND SHEPPARD D. TGF-β is a critical mediator of acute lung injury. *J Clin Invest* 107: 1537–1544, 2001.
 287. PITTET JF, KUDOH I, AND WIENER-KRONISH JP. Endothelial exposure to *Pseudomonas aeruginosa* proteases increases the vulnerability of the alveolar epithelium to a second injury. *Am J Respir Cell Mol Biol* 18: 129–135, 1998.
 288. PITTET JF, LU LN, MORRIS DG, MODELSKA K, WELCH WJ, CAREY HV, ROUX J, AND MATTHAY MA. Reactive nitrogen species inhibit alveolar epithelial fluid transport after hemorrhagic shock in rats. *J Immunol* 166: 6301–6310, 2001.
 289. PITTET JF, MATTHAY MA, PIER G, GRADY M, AND WIENER-KRONISH JP. *Pseudomonas aeruginosa*-induced lung and pleural injury in sheep. Differential protective effect of circulating versus alveolar immunoglobulin G antibody. *J Clin Invest* 92: 1221–1228, 1993.
 290. PITTET JF, WIENER-KRONISH JP, MCELROY MC, FOLKESSON HG, AND MATTHAY MA. Stimulation of lung epithelial liquid clearance by endogenous release of catecholamines in septic shock in anesthetized rats. *J Clin Invest* 94: 663–671, 1994.
 291. PITTET JF, WIENER-KRONISH JP, SERIKOV V, AND MATTHAY MA. Resistance of the alveolar epithelium to injury from septic shock in sheep. *Am J Respir Crit Care Med* 151: 1093–1100, 1995.
 292. PLANES C, ESCOUBET B, BLOT-CHABAUD M, FRIEDLANDER G, FARMAN N, AND CLERICI C. Hypoxia downregulates expression and activity of epithelial sodium channels in rat alveolar epithelial cells. *Am J Respir Cell Mol Biol* 17: 508–518, 1997.
 293. PLANES C, FRIEDLANDER G, LOISEAU A, AMIEL C, AND CLERICI C. Inhibition of Na-K-ATPase activity after prolonged hypoxia in an alveolar epithelial cell line. *Am J Physiol Lung Cell Mol Physiol* 271: L70–L78, 1996.
 294. RAFII B, TANSWELL AK, OTULAKOWSKI G, PITKANEN O, BELCASTRO-TAYLOR R, AND O'BRODOVICH H. O₂-induced ENaC expression is associated with NF-κB activation and blocked by superoxide scavenger. *Am J Physiol Lung Cell Mol Physiol* 275: L764–L770, 1998.
 295. RAMBOTTI MG AND SPRECA A. Ultrastructural demonstration of guanylate cyclase in rat lung after activation by ANF. *Cell Mol Biol* 37: 455–462, 1991.
 296. RAMMINGER SJ, BAINES DL, OLVER RE, AND WILSON SM. The effects of Po₂ upon transepithelial ion transport in fetal rat distal lung epithelial cells. *J Physiol* 524: 539–547, 2000.
 297. REDDY MM, LIGHT MJ, AND QUINTON PM. Activation of the epithelial Na⁺ channel (ENaC) requires CFTR Cl[−] channel function. *Nature* 402: 301–304, 1999.
 298. RENARD S, LINGUEGLIA E, VOILLEY N, LAZDUNSKI M, AND BARBRY P. Biochemical analysis of the membrane topology of the amiloride-sensitive Na⁺ channel. *J Biol Chem* 269: 12981–12986, 1994.
 299. RENARD S, VOILLEY N, BASSILANA F, LAZDUNSKI M, AND BARBRY P. Localization and regulation by steroids of the alpha, beta and gamma subunits of the amiloride-sensitive Na⁺ channel in colon, lung and kidney. *Pflügers Arch* 430: 299–307, 1995.
 300. REZAIGUIA S, GARAT C, DELCLAUX C, MEIGNAN M, FLEURY J, LEGRAND P, MATTHAY MA, AND JAYR C. Acute bacterial pneumonia in rats increases alveolar epithelial fluid clearance by a tumor necrosis factor-α-dependent mechanism. *J Clin Invest* 99: 325–335, 1997.
 301. REZAIGUIA-DELCLAUX S, JAYR C, LUO DF, SAIDI NE, MEIGNAN M, AND DUVALDESTIN P. Halothane and isoflurane decrease alveolar epithelial fluid clearance in rats. *Anesthesiology* 88: 751–760, 1998.
 302. RICHARDS NW AND DAWSON DC. Single potassium channels blocked by lidocaine and quinidine in isolated turtle colon epithelial cells. *Am J Physiol Cell Physiol* 251: C85–C86, 1986.
 303. RIDGE KM, RUTSCHMAN DH, FACTOR P, KATZ AI, BERTORELLO AM, AND SZNAJDER JI. Differential expression of Na-K-ATPase isoforms in rat alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 273: L246–L255, 1997.
 304. ROTIN D, KANELIS V, AND SCHILD L. Trafficking and cell surface stability of ENaC. *Am J Physiol Renal Physiol* 281: F391–F399, 2001.
 305. RUBIN DB, WIENER-KRONISH JP, MURRAY JF, GREEN DR, TURNER J, LUCE JM, MONTGOMERY AB, MARKS JD, AND MATTHAY MA. Elevated

- von Willebrand factor antigen is an early plasma predictor of acute lung injury in nonpulmonary sepsis syndrome. *J Clin Invest* 86: 474–480, 1990.
306. RUTSCHMAN DH, OLIVERA W, AND SZNAJDER JI. Active transport and passive liquid movement in isolated perfused rat lungs. *J Appl Physiol* 75: 1574–1580, 1993.
 307. SAKUMA T, FOLKESSON HG, SUZUKI S, OKANIWA G, FUJIMURA S, AND MATTHAY MA. Beta-adrenergic agonist stimulated alveolar fluid clearance in ex vivo human and rat lungs. *Am J Respir Crit Care Med* 155: 506–512, 1997.
 308. SAKUMA T, OKANIWA G, NAKADA T, NISHIMURA T, FUJIMURA S, AND MATTHAY MA. Alveolar fluid clearance in the resected human lung. *Am J Respir Crit Care Med* 150: 305–310, 1994.
 309. SAKUMA T, PITTET JF, JAYR C, AND MATTHAY MA. Alveolar liquid and protein clearance in the absence of blood flow or ventilation in sheep. *J Appl Physiol* 74: 176–185, 1993.
 310. SALDÍAS FJ, AZZAM ZS, RIDGE KM, YELDANDI A, RUTSCHMAN DH, SCHRAUFNAGEL D, AND SZNAJDER JI. Alveolar fluid reabsorption is impaired by increased left atrial pressures in rats. *Am J Physiol Lung Cell Mol Physiol* 281: L591–L597, 2001.
 311. SALDÍAS FJ, COMELLAS A, RIDGE KM, LECUONA E, AND SZNAJDER JI. Isoproterenol improves ability of lung to clear edema in rats exposed to hyperoxia. *J Appl Physiol* 87: 30–35, 1999.
 312. SALDÍAS FJ, LECUONA E, COMELLAS AP, RIDGE KM, AND SZNAJDER JI. Dopamine restores lung ability to clear edema in rats exposed to hyperoxia. *Am J Respir Crit Care Med* 159: 626–633, 1999.
 313. SAUMON G AND BASSET G. Electrolyte and fluid transport across the mature alveolar epithelium. *J Appl Physiol* 74: 1–15, 1993.
 314. SAWA T, YAHN TL, OHARA M, KURAHASHI K, GROPPER MA, WIENER-KRONISH JP, AND FRANK DW. Active and passive immunization with the *Pseudomonas* V antigen protects against type III intoxication and lung injury. *Nat Med* 5: 392–398, 1999.
 315. SAYEGH R, AUERBACH SD, LI X, LOFTUS RW, HUSTED RF, STOKES JB, AND THOMAS CP. Glucocorticoid induction of epithelial sodium channel expression in lung and renal epithelia occurs via trans-activation of a hormone response element in the 5'-flanking region of the human epithelial sodium channel α subunit gene. *J Biol Chem* 274: 12431–12437, 1999.
 316. SCHAEDEL C, MARTINSEN L, KRISTOFFERSSON AC, KORNFALT R, NILSSON KO, ORLENIUS B, AND HOLMBERG L. Lung symptoms in pseudohypoaldosteronism type I are associated with deficiency of the α -subunit of the epithelial sodium channel. *J Pediatr* 135: 739–745, 1999.
 317. SCHNEEBERGER EE AND KARNOVSKY MJ. The influence of intravascular fluid volume on the permeability of newborn and adult mouse lungs to ultrastructural protein tracers. *J Cell Biol* 49: 310–334, 1971.
 318. SCHNEEBERGER EE AND LYNCH RD. Structure, function, and regulation of cellular tight junctions. *Am J Physiol Lung Cell Mol Physiol* 262: L647–L661, 1992.
 319. SCHNEEBERGER EE AND MCCARTHY KM. Cytochemical localization of $\text{Na}^+\text{-K}^+\text{-ATPase}$ in rat type II pneumocytes. *J Appl Physiol* 60: 1584–1589, 1986.
 320. SCHNEEBERGER-KEELEY EE AND KARNOVSKY MJ. The ultrastructural basis of alveolar-capillary membrane permeability to peroxidase used as a tracer. *J Cell Biol* 37: 781–793, 1968.
 321. SCHNITZER JE AND OH P. Aquaporin-1 in plasma membrane and caveolae provides mercury-sensitive water channels across lung endothelium. *Am J Physiol Heart Circ Physiol* 270: H416–H422, 1996.
 322. SCHWIEBERT EM, POTTER ED, HWANG TH, WOO JS, DING C, QIU W, GUGGINO WB, LEVINE MA, AND GUGGINO SE. cGMP stimulates sodium and chloride currents in rat tracheal airway epithelia. *Am J Physiol Cell Physiol* 272: C911–C922, 1997.
 323. SERIKOV VB, GRADY M, AND MATTHAY MA. Effect of temperature on alveolar liquid and protein clearance in an in situ perfused goat lung. *J Appl Physiol* 75: 940–947, 1993.
 324. SHIMKETS RA, LIFTON R, AND CANESSA CM. In vivo phosphorylation of the epithelial sodium channel. *Proc Natl Acad Sci USA* 95: 3301–3305, 1998.
 325. SITIPUNT C, STEINBERG KP, RUZINSKI JT, MYLES C, ZHU S, GOODMAN RB, HUDSON LD, MATALON S, AND MARTIN TR. Nitric oxide and nitrotyrosine in the lungs of patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 163: 503–510, 2001.
 326. SKOU JC. Nobel Lecture. The identification of the sodium pump. *Biosci Rep* 18: 155–169, 1998.
 327. SMEDIRA N, GATES L, HASTINGS R, JAYR C, SAKUMA T, PITTET JF, AND MATTHAY MA. Alveolar and lung liquid clearance in anesthetized rabbits. *J Appl Physiol* 70: 1827–1835, 1991.
 328. SNYDER PM. Liddle's syndrome mutations disrupt cAMP-mediated translocation of the epithelial Na^+ channel to the cell surface. *J Clin Invest* 105: 45–53, 2000.
 329. SNYDER PM, CHENG C, PRINCE LS, ROGERS JC, AND WELSH MJ. Electrophysiological and biochemical evidence that DEG/ENaC cation channels are composed of nine subunits. *J Biol Chem* 273: 681–684, 1998.
 330. SNYDER PM, McDONALD FJ, STOKES JB, AND WELSH MJ. Membrane topology of the amiloride-sensitive epithelial sodium channel. *J Biol Chem* 269: 24379–24383, 1994.
 331. SONG Y, FUKUDA N, BAI C, MA T, MATTHAY MA, AND VERKMAN AS. Role of aquaporins in alveolar fluid clearance in neonatal and adult lung, and in oedema formation following acute lung injury: studies in transgenic aquaporin null mice. *J Physiol* 525: 771–779, 2000.
 332. SONG Y, JAYARAMAN S, YANG B, MATTHAY MA, AND VERKMAN AS. Role of aquaporin water channels in airway fluid transport, humidification, and surface liquid hydration. *J Gen Physiol* 117: 573–582, 2001.
 333. STAUB NC. Pulmonary edema. *Physiol Rev* 54: 678–811, 1974.
 334. STAUB NC AND ALBERTINE KH. The structure of the lungs relative to their principal function. In: *Textbook of Respiratory Medicine*, edited by Murray JF and Nadel JA. Philadelphia, PA: Saunders, 1988, p. 12–36.
 335. STERN M, ULRICH K, ROBINSON C, COPELAND J, GRIESENBACH U, MASSE C, CHENG S, MUNKONGE F, GEDDES D, BERTHAUME Y, AND ALTON E. Pretreatment with cationic lipid-mediated transfer of the $\text{Na}^+\text{-K}^+\text{-ATPase}$ pump in a mouse model in vivo augments resolution of high permeability pulmonary oedema. *Gene Ther* 7: 960–966, 2000.
 336. STUTTS MJ, CANESSA CM, OLSEN JC, HAMRICK M, COHN JA, ROSSIER BC, AND BOUCHER RC. CFTR as a cAMP-dependent regulator of sodium channels (see comments). *Science* 269: 847–850, 1995.
 337. SUGAHARA K, CALDWELL JH, AND MASON RJ. Electrical currents flow out of domes formed by cultured epithelial cells. *J Cell Biol* 154: 1541–1544, 1984.
 338. SUZUKI S, NODA M, SUGITA M, ONO S, KOIKE K, AND FUJIMURA S. Impairment of transalveolar fluid transport and lung $\text{Na}^+\text{-K}^+\text{-ATPase}$ function by hypoxia in rats. *J Appl Physiol* 87: 962–968, 1999.
 339. SUZUKI S, TSUBOCHI H, SUZUKI T, DARNEL AD, KROZOWSKI ZS, SASANO H, AND KONDO T. Modulation of transalveolar fluid absorption by endogenous aldosterone in adult rats. *Exp Lung Res* 27: 143–155, 2001.
 340. SUZUKI S, ZUEGE D, AND BERTHAUME Y. Sodium-independent modulation of $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity by β -adrenergic agonist in alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 268: L983–L990, 1995.
 341. SUZUKI T, SASANO H, SUZUKI S, HIRASAWA G, TAKEYAMA J, MURAMATSU Y, DATE F, NAGURA H, AND KROZOWSKI ZS. 11 β -hydroxysteroid dehydrogenase type 2 in human lung: possible regulator of mineralocorticoid action. *J Clin Endocrinol Metab* 83: 4022–4025, 1998.
 342. SWEZEY N, TCHEPICHEV S, GAGNON S, FERTUCK K, AND O'BRODOVICH H. Female gender hormones regulate mRNA levels and function of the rat lung epithelial Na channel. *Am J Physiol Cell Physiol* 274: C379–C386, 1998.
 343. SZNAJDER JI, OLIVERA WG, RIDGE KM, AND RUTSCHMAN DH. Mechanisms of lung liquid clearance during hyperoxia in isolated rat lungs. *Am J Respir Crit Care Med* 151: 1519–1525, 1995.
 344. SZNAJDER JI, RIDGE KM, YEATES DB, ILEKIS J, AND OLIVERA W. Epidermal growth factor increases lung liquid clearance in rat lungs. *J Appl Physiol* 85: 1004–1010, 1998.
 345. TALBOT CL, BOSWORTH DG, BRILEY EL, FENSTERMACHER DA, BOUCHER RC, GABRIEL SE, AND BARKER PM. Quantitation and localization of ENaC subunit expression in fetal, newborn, and adult mouse lung. *Am J Respir Cell Mol Biol* 20: 398–406, 1999.
 346. TAYLOR AE AND GAAR KA JR. Estimation of equivalent pore radii of pulmonary capillary and alveolar membranes. *Am J Physiol* 218: 1133–1140, 1970.

347. TAYLOR AE, GUYTON AC, AND BISHOP VS. Permeability of the alveolar epithelium to solutes. *Circ Res* 16: 353–362, 1965.
348. TCHEPICHEV S, UEDA J, CANESSA C, ROSSIER BC, AND O'BRODOVICH H. Lung epithelial Na channel subunits are differentially regulated during development and by steroids. *Am J Physiol Cell Physiol* 269: C805–C812, 1995.
349. THARAUX PL, DUSSAULE JC, COULETTE S, AND CLERICI C. Evidence for functional ANP receptors in cultured alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 274: L244–L251, 1998.
350. THOMAS CP, DOGGETT NA, FISHER R, AND STOKES JB. Genomic organization and the 5' flanking region of the gamma subunit of the human amiloride-sensitive epithelial sodium channel. *J Biol Chem* 271: 26062–26066, 1996.
351. THOME U, CHEN L, FACTOR P, DUMASUS V, FREEMAN B, SZNAJDER JJ, AND MATALON S. Na,K-ATPase gene transfer mitigates an oxidant-induced decrease of active sodium transport in rat fetal ATI cells. *Am J Respir Cell Mol Biol* 24: 245–252, 2001.
352. TOMLINSON LA, CARPENTER TC, BAKER EH, BRIDGES JB, AND WEIL JV. Hypoxia reduces airway epithelial sodium transport in rats. *Am J Physiol Lung Cell Mol Physiol* 277: L881–L886, 1999.
353. TOWNE JE, HARROD KS, KRANE CM, AND MENON AG. Decreased expression of aquaporin (AQP)1 and AQP5 in mouse lung after acute viral infection. *Am J Respir Cell Mol Biol* 22: 34–44, 2000.
354. TRAEBERT M, HATTENHAUER O, MURER H, KAISLING B, AND BIBER J. Expression of type II Na-P_i cotransporter in alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 277: L868–L873, 1999.
355. UMENISHI F, CARTER EP, YANG B, OLIVER B, MATTHAY MA, AND VERKMAN AS. Sharp increase in rat lung water channel expression in the perinatal period. *Am J Respir Cell Mol Biol* 15: 673–679, 1996.
356. VALLET V, CHRAÏBI A, GAEGGELER HP, HORISBERGER JD, AND ROSSIER BC. An epithelial serine protease activates the amiloride-sensitive sodium channel. *Nature* 389: 607–610, 1997.
357. VAN DRIESSCHE W. Lidocaine blockade of basolateral potassium channels in the amphibian urinary bladder. *J Physiol* 381: 575–593, 1986.
358. VAN SCOTT MR, CHINET TC, BURNETTE AD, AND PARADISO AM. Puri-nergic regulation of ion transport across nonciliated bronchiolar epithelial (Clara) cells. *Am J Physiol Lung Cell Mol Physiol* 269: L30–L37, 1995.
359. VAN SCOTT MR, DAVIS CW, AND BOUCHER RC. Na⁺ and Cl[−] transport across rabbit nonciliated bronchiolar epithelial (Clara) cells. *Am J Physiol Cell Physiol* 256: C893–C901, 1989.
360. VENGLARIK CJ, VERGHESE G, FORMAN HJ, MATTHAY MA, AND MATALON S. Regulation of airway Na⁺ absorption by extracellular protease (Abstract). *FASEB J* 15: A859, 2001.
361. VERGHESE GM, WARE LB, MATTHAY BA, AND MATTHAY MA. Alveolar epithelial fluid transport and the resolution of clinically severe hydrostatic pulmonary edema. *J Appl Physiol* 87: 1301–1312, 1999.
362. VERKMAN AS. Role of aquaporin water channels in kidney and lung. *Am J Med Sci* 316: 310–320, 1998.
363. VERKMAN AS. Lessons on renal physiology from transgenic mice lacking aquaporin water channels. *J Am Soc Nephrol* 10: 1126–1135, 1999.
364. VERKMAN AS. Physiological importance of aquaporins: lessons from knockout mice. *Curr Opin Nephrol Hypertens* 9: 517–522, 2000.
365. VERKMAN AS, MATTHAY MA, AND SONG Y. Aquaporin water channels and lung physiology. *Am J Physiol Lung Cell Mol Physiol* 278: L867–L879, 2000.
366. VERKMAN AS AND MITRA AK. Structure and function of aquaporin water channels. *Am J Physiol Renal Physiol* 278: F13–F28, 2000.
367. VERKMAN AS, YANG B, SONG Y, MANLEY GT, AND MA T. Role of water channels in fluid transport studied by phenotype analysis of aquaporin knockout mice. *Exp Physiol* 85: 233S–241S, 2000.
368. VERREY F, SPINDLER B, AND MASTROBERARDINO L. Regulation of Na⁺ reabsorption in aldosterone target epithelia. *Kidney Blood Press Res* 20: 148–150, 1997.
369. VIVONA M, MATTHAY MA, CHABAUD M, FRIEDLANDER G, AND CLERICI C. Hypoxia reduces alveolar epithelial sodium and fluid transport in rats: reversal by beta adrenergic treatment. *Am J Respir Cell Mol Biol*. In press.
370. VOILLEY N, LINGUEGLIA E, CHAMPIGNY G, MATTÉI MG, WALDMANN R, LAZDUNSKI M, AND BARBRY P. The lung amiloride-sensitive Na⁺ channel: biophysical properties, pharmacology, ontogenesis, and molecular cloning. *Proc Natl Acad Sci USA* 91: 247–251, 1994.
371. WALTERS DV AND OLVER RE. The role of catecholamines in lung liquid absorption at birth. *Pediatr Res* 12: 239–242, 1978.
372. WANG PM, ASHINO Y, ICHIMURA H, AND BHATTACHARYA J. Rapid alveolar liquid removal by a novel convective mechanism. *Am J Physiol Lung Cell Mol Physiol* 281: L1327–L1334, 2001.
373. WANG Y, FOLKESSON HG, JAYR C, WARE LB, AND MATTHAY MA. Alveolar epithelial fluid transport can be simultaneously upregulated by both KGF and β -agonist therapy. *J Appl Physiol* 87: 1852–1860, 1999.
374. WARE LB, GOLDEN JA, FINKBEINER WE, AND MATTHAY MA. Alveolar epithelial fluid transport capacity in reperfusion lung injury after lung transplantation. *Am J Respir Crit Care Med* 159: 980–988, 1999.
375. WARE LB AND MATTHAY MA. Alveolar fluid clearance is impaired in the majority of patients with acute lung injury and the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 163: 1376–1383, 2001.
376. WEIBEL ER. Lung morphometry and models in respiratory physiology. In: *Respiratory Physiology. An Analytical Approach*, edited by Chang HK and Paiva M. New York: Dekker, 1989, p. 1–56.
377. WEST JB AND MATHIEU-COSTELLO O. Strength of the pulmonary blood-gas barrier. *Respir Physiol* 88: 141–148, 1992.
378. WIDDICOMBE JH. How does cAMP increase active Na absorption across alveolar epithelium? *Am J Physiol Lung Cell Mol Physiol* 278: L231–L232, 2000.
379. WIDDICOMBE JH. Yet another role for the cystic fibrosis transmembrane conductance regulator. *Am J Respir Cell Mol Biol* 22: 11–14, 2000.
380. WIENER-KRONISH JP, ALBERTINE KH, AND MATTHAY MA. Differential responses of the endothelial and epithelial barriers of the lung in sheep to *Escherichia coli* endotoxin. *J Clin Invest* 88: 864–875, 1991.
381. WIENER-KRONISH JP, SAKUMA T, KUDOH I, PITTET JF, FRANK D, DOBBS L, VASIL ML, AND MATTHAY MA. Alveolar epithelial injury and pleural empyema in acute *P. aeruginosa* pneumonia in anesthetized rabbits. *J Appl Physiol* 75: 1661–1669, 1993.
382. WILLUMSEN NJ AND BOUCHER RC. Sodium transport and intracellular sodium activity in cultured human nasal epithelium. *Am J Physiol Cell Physiol* 261: C319–C331, 1991.
383. WODOPIA R, KO HS, BILLIAN J, WIESNER R, BARTSCH P, AND MAIRBÄURL H. Hypoxia decreases proteins involved in epithelial electrolyte transport in A549 cells and rat lung. *Am J Physiol Lung Cell Mol Physiol* 279: L1110–L1119, 2000.
384. WOLLMER P, SCHAIRER W, BOS JA, BAKKER W, KRENNING EP, AND LACHMANN B. Pulmonary clearance of ^{99m}Tc-DTPA during halothane anaesthesia. *Acta Anaesthesiol Scand* 34: 572–575, 1990.
385. YASUI M, SERLACHIUS E, LÖFGREN M, BELUSA R, NIELSEN S, AND APERIA A. Perinatal changes in expression of aquaporin-4 and other water and ion transporters in rat lung. *J Physiol* 505: 3–11, 1997.
386. YUE G AND MATALON S. Mechanisms and sequelae of increased alveolar fluid clearance in hyperoxic rats. *Am J Physiol Lung Cell Mol Physiol* 272: L407–L412, 1997.
387. YUE G, RUSSELL WJ, BENOS DJ, JACKSON RM, OLMAN MA, AND MATALON S. Increased expression and activity of sodium channels in alveolar type II cells of hyperoxic rats. *Proc Natl Acad Sci USA* 92: 8418–8422, 1995.
388. YUE G, SHOEMAKER RL, AND MATALON S. Regulation of low-amiloride-affinity sodium channels in alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 267: L94–L100, 1994.
389. ZHU S, BASIOUNY KF, CROW JP, AND MATALON S. Carbon dioxide enhances nitration of surfactant protein A by activated alveolar macrophages. *Am J Physiol Lung Cell Mol Physiol* 278: L1025–L1031, 2000.
390. ZHU S, KACHEL DL, MARTIN WJ II, AND MATALON S. Nitrated SP-A does not enhance adherence of *Pneumocystis carinii* to alveolar macrophages. *Am J Physiol Lung Cell Mol Physiol* 275: L1031–L1039, 1998.
391. ZHU S, WARE LB, GEISER T, MATTHAY MA, AND MATALON S. Increased levels of nitrate and surfactant protein A nitration in the pulmonary edema fluid of patients with acute lung injury. *Am J Respir Crit Care Med* 163: 166–172, 2001.