

Review Article

ROLE OF ALVEOLAR MACROPHAGES IN THE INFLAMMATORY RESPONSE AFTER TRAUMA

Ulrike Niesler,* Annette Palmer,* Peter Radermacher,[†] and Markus S. Huber-Lang*

**Departments of Trauma Surgery, Hand, Plastic and Reconstructive Surgery, and [†]Anesthesiology and Intensive Care, Center for Biomedical Research, University of Ulm, Ulm, Germany*

Received 27 Jan 2014; first review completed 12 Feb 2014; accepted in final form 4 Mar 2014

ABSTRACT—Acute lung injury (ALI) and its more severe form, acute respiratory distress syndrome (ARDS), can result from both direct and indirect pulmonary damage caused by trauma and shock. In the course of ALI/ARDS, mediators released from resident cells, such as alveolar macrophages, may act as chemoattractants for invading cells and stimulate local cells to build up a proinflammatory microenvironment. Depending on the trauma setting, the role of alveolar macrophages is differentially defined. This review focuses on alveolar macrophage function after blunt chest trauma, ischemia/reperfusion, hemorrhagic shock, and thermal burns.

KEYWORDS—Alveolar macrophages, trauma, inflammation, blunt chest trauma, ischemia/reperfusion, hemorrhagic shock, burn

ABBREVIATIONS—ALI — acute lung injury; AP-1 — activator protein 1; ARDS — acute respiratory distress syndrome; AT2 cells — alveolar epithelial type 2 cells; CCR2 — CC chemokine receptor type 2; CINC-1 — cytokine-induced neutrophil chemoattractant 1; COX — cyclooxygenase; cPLA2 — cytosolic phospholipase A2; Hck — tyrosine-protein kinase Hck; HMGB1 — high-mobility group protein B1; HS/R — hemorrhagic shock/resuscitation; IL — interleukin; iNKT — invariant natural killer T cells; I/R — ischemia/reperfusion; IRAK4 — interleukin-1 receptor-associated kinase 4; JNK — c-Jun N-terminal kinase; KC — keratinocyte-derived chemokine; LPS — lipopolysaccharide; MCP-1 — monocyte chemoattractant protein-1; MIP-1 α — macrophage inflammatory protein-1 α ; MIP-2 — macrophage inflammatory protein 2; MyD88 — myeloid differentiation primary response gene (88); NF- κ B — nuclear factor kappa B; Nlrp3 — NOD-like receptor family, pyrin domain containing protein 3; NO — nitric oxide; PCA — procoagulant activity; ROS — reactive oxygen species; Src — proto-oncogene tyrosine-protein kinase Src; TLR — toll-like receptor; TNF — tumor necrosis factor

INTRODUCTION

Acute lung injury (ALI) and its more severe form acute respiratory distress syndrome (ARDS) are common sequelae after trauma and shock. Aside from lung infection (1) and sepsis (2), ALI/ARDS can result from both direct pulmonary damage (e.g., blunt chest trauma) (3) and indirect damage (e.g., after burn injury) (4). A common complication of ARDS is pneumonia based on nosocomial infections, especially in mechanically ventilated patients (5). Acute lung injury and ARDS are characterized by increased vascular permeability caused by dysfunction of the alveolar-capillary membrane. The resulting lung edema-induced impairment of blood gas exchange leads to decreased arterial oxygenation. Cytokines released from resident cells, such as alveolar macrophages or alveolar epithelial type 2 (AT2) cells, into the alveoli induce the migration of large numbers of activated inflammatory cells into the air space (6, 7).

In turn, mediator release from the accumulated inflammatory cells further contributes to the tissue damage in ALI and ARDS.

Alveolar macrophages are phagocytes (greek: *phagein*, “to eat”; *kýtos*, “hollow vessel”), which are located in the alveolar compartment of the lungs. Under physiologic conditions, alveolar macrophages play a pivotal role in the protection of the lungs against inhaled particles like dust, bacteria, and viruses. However, under inflammatory conditions, alveolar macrophages generate and release a multitude of mediators, such as cytokines, chemokines, complement factors, alarmins, and arachidonic acid metabolites (8–12) (Table 1). These mediators may act as chemoattractants for invading cells (6) or stimulate local cells (epithelial cells or interstitial macrophages) to build up a proinflammatory microenvironment (13).

Experimentally, alveolar macrophages have been analyzed in different trauma models in mice, rats, or swine (10, 14–17). Technically, alveolar macrophages are relatively easy to obtain by bronchoalveolar lavage and can be used for various functional *ex vivo/in vitro* analyses. In some cases, alveolar macrophages were depleted before the experimental insult, and changes in the inflammatory response after trauma were analyzed. However, although general mediator release in bronchoalveolar lavage fluids has been extensively analyzed in the clinical setting of trauma (18–20), there are only a few data on the specific role of alveolar macrophages after trauma. An exception is the alveolar macrophage function after burn injury, which has been studied to some extent in humans (21–23).

Address reprint requests to Ulrike Niesler, PhD, Trauma Lab, Department of Trauma Surgery, Hand, Plastic and Reconstructive Surgery, Center for Biomedical Research, Ulm University, Helmholtzstr. 8/2, 89081 Ulm, Germany. E-mail: ulrike.niesler@uniklinik-ulm.de.

No reprints will be ordered.

This work was supported by grants from the Deutsche Forschungsgemeinschaft (DFG KFO 200, KN 475/5-2; DFG KFO 200, HU 823/3-2).

None of the authors have any financial interests or affiliations with commercial organizations whose products or services are related to the subject matter of this article (no existing conflicts of interest).

DOI: 10.1097/SHK.0000000000000167

Copyright © 2014 by the Shock Society

TABLE 1. Proinflammatory and anti-inflammatory mediators released by alveolar macrophages after trauma

Proinflammatory mediators	Anti-inflammatory mediators
IL-1β	IL-10
TNF-α	TGF-β
IL-6	NO
MCP-1	
MIP-2	
MIP-1α	
CINC-1/KC/IL-8	
sFasL	
HMGB1	
Complement factors	
Thromboxanes	
Leukotrienes	
ROS (O ₂ ⁻ , H ₂ O ₂ , OH ⁻)	
NO	

HMGB1 indicates high-mobility group protein B1; ROS, reactive oxygen species; sFasL, soluble Fas ligand.

The role of alveolar macrophage function in the global inflammatory response is still not fully understood. Because of their location in the immediate vicinity of the lung capillaries, mediators released from alveolar macrophages (or from other stimulated lung cells) may be transported via the bloodstream to distant organs. At these sites, the lung-derived mediators may stimulate cells for cytokine release, thereby modifying the in-

flammatory balance. For example, the systemic inflammatory response after blunt chest trauma leads to impaired fracture healing by altering the recruitment of inflammatory cells to the fracture site (24). The cellular source of the causative mediators, however, has not been investigated yet.

Depending on the trauma setting, the role of alveolar macrophages is differentially defined. Therefore, this review focuses on alveolar macrophage function after blunt chest trauma, ischemia/reperfusion, hemorrhagic shock, and thermal burns.

ALVEOLAR MACROPHAGES DECISIVELY MODULATE THE PROINFLAMMATORY AND ANTI-INFLAMMATORY RESPONSE AFTER BLUNT CHEST TRAUMA

Lung contusion induced by blunt chest trauma is a common injury that can occur in an isolated manner but more often is observed in patients with multiple injuries (25). A frequent complication of lung contusion is ALI/ARDS (25).

Alveolar macrophages activated by blunt chest trauma release chemokines (monocyte chemotactic protein 1 [MCP-1], macrophage inflammatory protein 2 [MIP-2]) and proinflammatory (interleukin 1β [IL-1β], IL-6, tumor necrosis factor-α [TNF-α]) and anti-inflammatory (IL-10) cytokines. Monocyte chemotactic protein 1 and MIP-2 generated by alveolar macrophages stimulate the migration of monocytes (6) and neutrophils (7, 14, 26, 27) into the contused lung (Fig. 1). However, neutrophils rapidly undergo apoptosis and accumulated apoptotic neutrophils are phagocytosed by alveolar macrophages (14). Fas ligand, which is expressed by alveolar macrophages




FIG. 1. Role of alveolar macrophages after blunt chest trauma. AMΦ indicates alveolar macrophage; AT1 cell, alveolar type 1 epithelial cell; AT2 cell, alveolar type 2 epithelial cell; CCR2, CC chemokine receptor 2; COX, cyclooxygenase; PG, prostaglandins; PMN, neutrophil; sFasL, soluble Fas ligand; TLR, toll-like receptor.

and released into the alveoli, induces apoptosis of alveolar epithelial type 2 cells (28–30), which are then also phagocytosed by alveolar macrophages (14). The binding of Fas ligand to Fas receptor on alveolar macrophages does not induce apoptosis but instead modifies the cytokine release (31). In the setting of additive *ex vivo* stimulation with soluble Fas ligand, IL-6 levels are enhanced whereas IL-10 levels are decreased, and MCP-1 expression remains unchanged (29).

Interestingly, 1 h after blunt chest trauma, alveolar macrophages from the contused lungs of swine show an increased expression of the constitutive isoform of the key enzyme in prostaglandin biosynthesis, cyclooxygenase 1 (COX-1), but not of the inducible isoform COX-2 (17). The significance of this phenomenon in the inflammation after blunt chest trauma is not clear, however, COX-1 inhibition during influenza A viral infection was detrimental to the host (32).

Toll-like receptor 4 (TLR4) expression on alveolar macrophages is significantly increased 48 h after blunt chest trauma in mice (33), priming them for the defense against gram-negative bacteria such as *Escherichia coli* or *Pseudomonas aeruginosa*. At this late time point after trauma, TNF- α and IL-6 concentrations in bronchoalveolar lavage fluids are not increased compared with those in control mice (33), possibly because cytokine concentrations have already returned to sham levels. Infection with *P. aeruginosa* causes a significant increase of TNF- α and IL-6 concentrations in bronchoalveolar lavage fluids of traumatized animals (33). This reaction shows that even at late time points, the alveolar macrophages are still

in a “primed” state, although the primary inflammation after blunt chest trauma has already been resolved.

In the contused lung, alveolar macrophages exhibit not only a proinflammatory role (6) but also regulate cytokine release locally within the lungs as well as systemically (34). The binding of MCP-1 to its receptor CCR2 on alveolar macrophages may play a regulatory role as cytokine release in the lungs of CCR2^{-/-} mice is significantly increased (35). The regulatory role of alveolar macrophages after blunt chest trauma has also been studied by our group (34). Using a nonlethal blast injury model of blunt chest trauma (36) and alveolar macrophage depletion indicated that alveolar macrophages ameliorate the local inflammatory response after lung contusion (34). This effect may be mediated by suppressing the cytokine release of AT2 cells and interstitial macrophages (34). Systemically, alveolar macrophages seem to suppress Kupffer cell chemokine release but are not involved in the splenic immunosuppression after blunt chest trauma (34, 37).

ALVEOLAR MACROPHAGES PLAY A CENTRAL ROLE IN ISCHEMIA/REPERFUSION INJURY IN LUNGS

Diminished blood flow during trauma or transplantation can lead to ischemia, which, together with the following restoration of blood flow (reperfusion), results in the clinical picture of ischemia/reperfusion injury (I/R). According to the temperature of the ischemic organ, I/R can be divided into warm




FIG. 2. **Role of alveolar macrophages after ischemia/reperfusion.** AMΦ indicates alveolar macrophage; AP1, activator protein 1; AT1 cell, alveolar type 1 epithelial cell; AT2 cell, alveolar type 2 epithelial cell; CCR2, CC chemokine receptor 2; COX, cyclooxygenase; cPLA₂, cytosolic phospholipase A2; HMGB1, high-mobility group protein B1; iNKT, invariant natural killer T cell; JNK, c-Jun N-terminal kinase; NF-κB, nuclear factor kappa B; PMN, neutrophil; RAGE, receptor for advanced glycation end products; sFasL, soluble Fas ligand; TLR, toll-like receptor.

TABLE 2. Extrapulmonary I/R

Authors	Model	Rat strain	AMΦ
LaNoue et al. (1998)	Gut I/R	Sprague Dawley	PGE ₂ , TxB ₂ , O ₂ ⁻ , PCA upregulated
Liu et al. (2001)	Liver I/R	Fischer	TNF-α, IL-1β upregulated
McGuire et al. (1996)	Liver I/R + endotoxemia	Fischer	AMΦ not activated
Moraes et al. (2008)	Gut I/R	Wistar	proinflammatory, not further specified
Silva et al. (2013)	Kidney I/R + OVA stimulation	Wistar	Phagocytosis and bacterial killing downregulated
Souza et al. (2000)	Gut I/R	Wistar	TNF-α, H ₂ O ₂ upregulated

PCA indicates procoagulant activity.

and cold I/R. Examples of warm I/R lung injury are pulmonary sleeve resection during surgical treatment of lung cancer or cardiopulmonary bypass during heart surgery (38, 39). Cold I/R injury is the most common cause of respiratory failure after lung transplantation (40–42).

Intrapulmonary I/R

In terms of inflammation, however, temperature does not seem to be a major critical factor. Both warm and cold I/R lead to activation of alveolar macrophages with upregulation of gene expression via NF-κB and JNK/AP-1 (43) in the early phase of reperfusion (Fig. 2). Alveolar macrophages release a multitude of mediators, such as proinflammatory cytokines (IL-1β, IL-6, TNF-α), chemokines (keratinocyte-derived chemokine [KC], MIP-2, MCP-1, MIP-1α), alarmins (high-mobility group protein B1 [HMGB1]), and arachidonic acid metabolites (8–12, 44). Through the release of TNF-α, AT2 cells are stimulated to release neutrophil chemoattractants such

as KC and MIP-2 (13). Along the chemokine gradient, neutrophils and monocytes migrate into the lungs.

Cytosolic phospholipase A₂ (cPLA₂) activity in alveolar macrophages is increased by I/R. The resulting release of thromboxane contributes to increased pulmonary artery pressure, which, together with leukotriene-induced vascular leakage and neutrophil infiltration, exacerbates lung edema (9).

Opposite results regarding alveolar macrophage function in two models of warm *ex vivo* I/R with preceding alveolar macrophage depletion (10, 45) may be well attributed to different fluid reperfusion protocols. Although alveolar macrophages seem to exhibit a proinflammatory profile when Krebs-Henseleit buffer was used (10), they present a rather suppressive functional profile in the model using diluted whole blood (45).

A new role of alveolar macrophages in the initiation of I/R injury has been described in a recent publication: the binding of alveolar macrophage-derived HMGB1 to the receptor for advanced glycation end products (RAGE) on invariant natural killer T cells (iNKT) results in the release of the




FIG. 3. Role of alveolar macrophages after hemorrhagic shock. AMΦ indicates alveolar macrophage; AT1 cell, alveolar type 1 epithelial cell; AT2 cell, alveolar type 2 epithelial cell; CCR2, CC chemokine receptor 2; COX, cyclooxygenase; Hck, tyrosine-protein kinase Hck; LPS, lipopolysaccharide; NF-κB, nuclear factor kappa B; NO, nitric oxide; PMN, neutrophil; ROS, reactive oxygen species; sFasL, soluble Fas ligand; TF, tissue factor; TLR, toll-like receptor; ⊕, exposure to.

neutrophil-chemoattractant IL-17 and subsequent neutrophil infiltration (46). Furthermore, toll-like receptor 4 (TLR4) on alveolar macrophages plays an important role in I/R-induced inflammation, as a TLR4 knockout leads to reduced cytokine expression and an inability to recruit neutrophils into the lung (12). Considering the fact that HMGB1 binds specifically to TLR4 (47), it may activate alveolar macrophages (48) in an autocrine or paracrine manner.

Extrapulmonary I/R

The majority of the studies focused on the alveolar macrophage function after pulmonary I/R, however, alveolar macrophages are also influenced by extrapulmonary I/R, for example, in the gut, kidneys, or liver (Table 2). Three studies on I/R in the gut and liver (49–51) showed activation of alveolar macrophages, with the release of TNF- α , IL-1 β , and arachidonic acid metabolites (prostaglandin E₂, thromboxane A₂), which seems to be comparable to the alveolar macrophage reaction to pulmonary I/R. In addition, production of reactive oxygen species (O₂⁻, H₂O₂) by alveolar macrophages has been described (50, 51). The proinflammatory activity of alveolar macrophages after gut I/R mediates the development of ALI, as depletion of alveolar macrophages leads to decreased lung permeability and blocks the occurrence of ALI (52).

However, a combination of hepatic I/R and endotoxemia did not activate alveolar macrophages, as determined by unchanged superoxide formation of alveolar macrophages (53). Interestingly, kidney I/R led to reduced phagocytic and microbicidal capacity of alveolar macrophages in ovalbumin-immunized rats (54).

HEMORRHAGIC SHOCK/RESUSCITATION INDUCES MULTIPLE CHANGES IN ALVEOLAR MACROPHAGES

Hemorrhage with a significant loss of intravascular blood volume and consequential hypovolemia, tachycardia, and reduced blood pressure is known as hemorrhagic shock (HS). During severe hypovolemia, decreases in blood flow in all organs leads to reduced delivery of oxygen and nutrients to the tissues and subsequent cellular hypoxia. Hypoxia in turn stimulates the production and release of reactive oxygen species (ROS). Fluid resuscitation can cause further injury because of the release of increasing amounts of ROS (e.g., by activated neutrophils) (55).

Surprisingly, despite hypoxic conditions in the lung tissue (comparable to I/R injury), alveolar macrophages are not fully activated early (up to 3 h) after HS/resuscitation (HS/R), as reflected by unaltered cytokine concentrations and neutrophil cell counts in the bronchoalveolar lavage fluids (56, 57). However, oxidative stress during HS/R primes alveolar macrophages, resulting in an enhanced inflammatory response to lipopolysaccharide (LPS) (Fig. 3). As a consequence, minor lung infections may result in the development of posthemorrhage global lung inflammation, a common cause of morbidity after HS/R. The sensitization of alveolar macrophages to LPS is caused by redistribution of TLR4 from the Golgi apparatus or from endosomes to lipid rafts in the plasma membrane and its colocalization with the adaptor protein MyD88 in a common signaling complex (58). In addition, the Src family kinase Hck, which plays an important role in LPS-induced cytokine




FIG. 4. **Role of alveolar macrophages after thermal burns.** AM Φ indicates alveolar macrophage; AT1 cell, alveolar type 1 epithelial cell; AT2 cell, alveolar type 2 epithelial cell; CCR2, CC chemokine receptor 2; COX, cyclooxygenase; LPS, lipopolysaccharide; PMN, neutrophil; sFasL, soluble Fas ligand; TLR, toll-like receptor; \oplus , exposure to.

production, is phosphorylated and activated in an oxidant-dependent manner (59).

Hemorrhagic shock/resuscitation in general does not seem to result in mediator release from unstimulated alveolar macrophages. However, alveolar macrophages are capable of releasing NO (60, 61) and TNF- α (60) in a 24-h culture when isolated 4.5 to 6 h after hemorrhage and short resuscitation time of 30 min or less.

On stimulation with LPS, alveolar macrophages show a MyD88-dependent increase of IRAK4 activity (55) as well as the translocation of the transcription factor NF- κ B from the cytoplasm to the nucleus (62, 63). Nuclear factor- κ B activation leads to the production and release of proinflammatory mediators such as TNF- α (56, 57, 64) and CINC-1 (15, 62, 65). Contrasting results regarding the release of the anti-inflammatory cytokine IL-10 may be caused by the duration of the resuscitation time. When resuscitation lasts 20 min, LPS-induced IL-10 release is noticeably reduced (15, 66), whereas after a prolonged resuscitation time of 2 h, LPS-induced IL-10 release is enhanced (56). In turn, IL-10 suppresses generation of proinflammatory IL-1 β via negative-feedback regulation of the Nlrp3 inflammasome (66).

The increased expression of tissue factor in alveolar macrophages after HS/R leads to increased procoagulant activity (PCA), that is, the activation of the extrinsic pathway of coagulation (57), which is associated with pulmonary fibrin deposition, a pathologic feature of acute lung injury.

The increased NO release from LPS-stimulated alveolar macrophages into the alveoli (61) results in the inhibition of a Na⁺ channel, with a subsequent decrease in the fluid transport capacity of the alveolar epithelium, and consequently the development of pulmonary edema (67).

Regarding TLR4 expression, which is normally suppressed by LPS, HS/R stabilizes TLR4 gene transcription and protein levels (68). Through TLR4 signaling, the expression of the peptidoglycan receptor TLR2 is upregulated (63, 69), "priming" the alveolar macrophages for infections with gram-positive bacteria.

THERMAL BURN INJURY PRIMES ALVEOLAR MACROPHAGES FOR AN EXCESSIVE INFLAMMATORY RESPONSE

Thermal burns can be caused by a variety of factors including fire, hot liquids, steam, or hot objects. Respiratory complications (pneumonia, ARDS, pulmonary embolism) are major causes of death after burn injury (70, 71). The role of alveolar macrophages in the development of ALI/ARDS in burn-injured patients has not been fully understood.

In the initial hyperinflammatory phase after burn, macrophages, monocytes, and neutrophils are activated by toxic metabolites. Clinically and experimentally, the inflammatory response of alveolar macrophages after burn injury is somehow delayed until 3 to 5 days after injury (Fig. 4). Although IL-6 concentrations in lung tissue (72) and bronchoalveolar lavage fluids (22) are increased on days 3 and 4 after burn, respectively, there is no corresponding IL-6 expression in alveolar macrophages (72). Nonstimulated alveolar macrophages release barely any inflammatory mediators in the first

7 days after thermal injury (16). In contrast, on stimulation with the TLR4 agonist LPS, alveolar macrophages release CINC-1/IL-8 early (day 1) after burn injury, whereas TNF- α concentrations and bioactivity as well as IL-6 concentrations are increased on days 3 and 4 (22, 73). Thus, alveolar macrophages may contribute to significantly elevated levels of IL-6 and TNF- α in lung tissue on days 3 and 4 of an *in vivo* model of burn and *P. aeruginosa* infection (74). Interestingly, 7 days after burn injury, LPS-stimulated alveolar macrophages release increased concentrations of IL-6, MIP-1 β , and RANTES but not TNF- α , KC (CINC-1), MCP-1, or MIP-1 α (16). However, stimulation with the TLR2 agonist zymosan leads not only to the release of IL-6 but also to generation of TNF- α , IL-17, MIP-1 β , and MCP-1 (16). Reaction of alveolar macrophages to both TLR4 and TLR2 agonists points to a priming for an excessive inflammatory response to gram-positive and gram-negative bacteria, which makes burn patients more susceptible to pulmonary complications in the later phases after burn injury. Especially if burn injury occurs in combination with inhalation injury, alveolar macrophages are primed for an increased inflammatory response to LPS and cytokine release is increased already early after injury (22).

The functional response of alveolar macrophages (chemotaxis, phagocytotic activity, respiratory burst, and microbicidal activity) is also significantly influenced by burn injury. Although overall microbicidal activity is enhanced in alveolar macrophages after burn injury, the phagocytotic activity is significantly depressed (75). The respiratory burst after phagocytosis of *P. aeruginosa* or latex beads is suppressed, even after stimulation with phorbol myristate acetate (PMA) (76). Under these conditions, alveolar macrophages are incapable of locally clearing a secondary infection with *P. aeruginosa*, resulting in the spread of the infection to the periphery and development of systemic inflammatory response and sepsis (77). Chemotaxis to zymosan-activated serum as well as random migration of alveolar macrophages is significantly impaired after burn injury (78). An exception from this is the unaltered migration of alveolar macrophages from burn-injured smokers toward casein- and zymosan-activated serum (23). The defective chemotaxis is accompanied by increases in RNA, protein, the lysosomal enzymes beta-glucuronidase and acid phosphatase, as well as the ectoenzyme membrane marker 5'-nucleotidase (78).

Smoke inhalation in addition to burn injury induces a significant hypersensitivity of alveolar macrophages to LPS with an increased release of IL-1RA, IL-6, IL-8, IL-10, G-CSF, IFN- γ , RANTES, TNF- α , and VEGF on day 1 after injury (21, 22). There is also increased random migration as well as increased target-specific migration of alveolar macrophages toward casein, zymosan-activated serum, and *N*-formylmethionyl-leucyl-phenylalanine (fMLP) (23).

CONCLUSIONS

Depending on the initial damage, the role of alveolar macrophages may be proinflammatory and anti-inflammatory and their inflammatory response may be immediate (blunt chest trauma, ischemia/reperfusion) or delayed (hemorrhagic shock, burn). "Priming" of alveolar macrophages for defense against

pathogens in the delayed inflammatory response is an important step in the development of ALI or ARDS. Besides own generation and release of inflammatory mediators, alveolar macrophages also significantly regulate the inflammatory response after trauma and shock.

At present, there is a lack of data on the systemic extrapulmonary consequence of alveolar macrophage activation after trauma, except for a suppressive effect on Kupffer cells after blunt chest trauma. Therefore, studies addressing systemic effects of alveolar macrophages in lung injury are needed. It is noteworthy that there are also no direct comparative studies on signal transduction pathways in alveolar macrophages after various traumas. Nevertheless, it is likely that generation of reactive oxygen species in the lung and TLR4 expression, as well as the release of inflammatory cytokines by alveolar macrophages, are similar between the different trauma settings. However, despite some similarities, such as increased TLR4 expression, a "one-fits-all" model of alveolar macrophage function after trauma and shock is neither feasible nor realistic. Moreover, the complex response after a combination of blunt chest trauma with hemorrhagic shock, ischemia/reperfusion, or burn injury has so far not been investigated. Further experimental and clinical studies on alveolar macrophage function after trauma are needed, especially with regard to preexisting conditions, such as nicotine-induced chronic obstructive pulmonary disease or asthma.

REFERENCES

- Traylor ZP, Aeffner F, Davis IC: Influenza A H1N1 induces declines in alveolar gas exchange in mice consistent with rapid post-infection progression from acute lung injury to ARDS. *Influenza Other Respir Viruses* 7(3):472–479, 2013.
- Mikkelsen ME, Shah CV, Meyer NJ, Gaieski DF, Lyon S, Miltiades AN, Goyal M, Fuchs BD, Bellamy SL, Christie JD: The epidemiology of acute respiratory distress syndrome in patients presenting to the emergency department with severe sepsis. *Shock* 40(5):375–381, 2013.
- Miller PR, Croce MA, Bee TK, Qaisi WG, Smith CP, Collins GL, Fabian TC: ARDS after pulmonary contusion: accurate measurement of contusion volume identifies high-risk patients. *J Trauma* 51(2):223–228, 2001.
- Dancey DR, Hayes J, Gomez M, Schouten D, Fish J, Peters W, Slutsky AS, Stewart TE: ARDS in patients with thermal injury. *Intensive Care Med* 25(11):1231–1236, 1999.
- Chastre J, Trouillet JL, Vaugnat A, Joly-Guillou ML, Clavier H, Dombret MC, Gibert C: Nosocomial pneumonia in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 157(4 Pt 1):1165–1172, 1998.
- Seitz DH, Niesler U, Palmer A, Sulger M, Braumüller ST, Perl M, Gebhard F, Knöferl MW: Blunt chest trauma induces mediator-dependent monocyte migration to the lung. *Crit Care Med* 38(9):1852–1859, 2010.
- Hoth JJ, Wells JD, Hiltbold EM, McCall CE, Yoza BK: Mechanism of neutrophil recruitment to the lung after pulmonary contusion. *Shock* 35(6):604–609, 2011.
- Gazoni LM, Tribble CG, Zhao MQ, Unger EB, Farrar RA, Ellman PI, Fernandez LG, Laubach VE, Kron IL: Pulmonary macrophage inhibition and inhaled nitric oxide attenuate lung ischemia-reperfusion injury. *Ann Thorac Surg* 84(1):247–253, 2007.
- Bellido-Reyes YA, Akamatsu H, Kojima K, Arai H, Tanaka H, Sunamori M: Cytosolic phospholipase A2 inhibition attenuates ischemia-reperfusion injury in an isolated rat lung model. *Transplantation* 81(12):1700–1707, 2006.
- Zhao M, Fernandez LG, Doctor A, Sharma AK, Zarbock A, Tribble CG, Kron IL, Laubach VE: Alveolar macrophage activation is a key initiation signal for acute lung ischemia-reperfusion injury. *Am J Physiol Lung Cell Mol Physiol* 291(5):L1018–L1026, 2006.
- Naidu BV, Woolley SM, Farivar AS, Thomas R, Fraga CH, Goss CH, Mulligan MS: Early tumor necrosis factor- α release from the pulmonary macrophage in lung ischemia-reperfusion injury. *J Thorac Cardiovasc Surg* 127(5):1502–1508, 2004.
- Prakash A, Mesa KR, Wilhelmsen K, Xu F, Dodd-o JM, Hellman J: Alveolar macrophages and Toll-like receptor 4 mediate ventilated lung ischemia reperfusion injury in mice. *Anesthesiology* 117(4):822–835, 2012.
- Sharma AK, Fernandez LG, Awad AS, Kron IL, Laubach VE: Proinflammatory response of alveolar epithelial cells is enhanced by alveolar macrophage-produced TNF- α during pulmonary ischemia-reperfusion injury. *Am J Physiol Lung Cell Mol Physiol* 293(1):L105–L113, 2007.
- Seitz DH, Palmer A, Niesler U, Fröba JS, Heidemann V, Rittlinger A, Braumüller ST, Zhou S, Gebhard F, Knöferl MW: Alveolar macrophage phagocytosis is enhanced after blunt chest trauma and alters the posttraumatic mediator release. *Shock* 36(6):621–627, 2011.
- Khadaroo RG, Fan J, Powers KA, Fann B, Kapus A, Rotstein OD: Impaired induction of IL-10 expression in the lung following hemorrhagic shock. *Shock* 22(4):333–339, 2004.
- Oppeltz RF, Rani M, Zhang Q, Schwacha MG: Burn-induced alterations in toll-like receptor-mediated responses by bronchoalveolar lavage cells. *Cytokine* 55(3):396–401, 2011.
- Desselle WJ, Greenhaw JJ, Trentham LL, Fabian TC, Proctor KG: Macrophage cyclooxygenase expression, immunosuppression, and cardiopulmonary dysfunction after blunt chest trauma. *J Trauma* 51(2):239–251, 2001.
- Raymondos K, Martin MU, Schmudlach T, Baus S, Weibach C, Welte T, Krettek C, Frink M, Hildebrand F: Early alveolar and systemic mediator release in patients at different risks for ARDS after multiple trauma. *Injury* 43(2):189–195, 2012.
- Hirani N, Antonicelli F, Strieter RM, Wiesener MS, Ratcliffe PJ, Haslett C, Donnelly SC: The regulation of interleukin-8 by hypoxia in human macrophages—a potential role in the pathogenesis of the acute respiratory distress syndrome (ARDS). *Mol Med* 7(10):685–697, 2001.
- Bitto A, Barone M, David A, Polito F, Familiari D, Monaco F, Giardina M, David T, Messina R, Noto A, et al.: High mobility group box-1 expression correlates with poor outcome in lung injury patients. *Pharmacol Res* 61(2):116–120, 2010.
- Davis CS, Albright JM, Carter SR, Ramirez L, Kim H, Gamelli RL, Kovacs EJ: Early pulmonary immune hyporesponsiveness is associated with mortality after burn and smoke inhalation injury. *J Burn Care Res* 33(1):26–35, 2012.
- Wright MJ, Murphy JT: Smoke inhalation enhances early alveolar leukocyte responsiveness to endotoxin. *J Trauma* 59(1):64–70, 2005.
- Riyami BM, Kinsella J, Pollok AJ, Clark C, Stevenson RD, Reid WH, Campbell D, Gemmell CG: Alveolar macrophage chemotaxis in fire victims with smoke inhalation and burns injury. *Eur J Clin Invest* 21(5):485–489, 1991.
- Recknagel S, Bindl R, Brochhausen C, Gockelmann M, Wehner T, Schoengraf P, Huber-Lang M, Claes L, Ignatius A: Systemic inflammation induced by a thoracic trauma alters the cellular composition of the early fracture callus. *J Trauma Acute Care Surg* 74(2):531–537, 2013.
- Cohn SM: Pulmonary contusion: review of the clinical entity. *J Trauma* 42(5):973–979, 1997.
- Chavko M, Adeb S, Ahlers ST, McCarron RM: Attenuation of pulmonary inflammation after exposure to blast overpressure by *N*-acetylcysteine amide. *Shock* 32(3):325–331, 2009.
- Raghavendran K, Davidson BA, Woytash JA, Helinski JD, Marschke CJ, Manderscheid PA, Notter RH, Knight PR: The evolution of isolated bilateral lung contusion from blunt chest trauma in rats: cellular and cytokine responses. *Shock* 24(2):132–138, 2005.
- Liener UC, Knöferl MW, Sträter J, Barth TF, Pauser EM, Nüssler AK, Kinzl L, Brückner UB, Gebhard F: Induction of apoptosis following blunt chest trauma. *Shock* 20(6):511–516, 2003.
- Seitz DH, Palmer A, Niesler U, Braumüller ST, Bauknecht S, Gebhard F, Knöferl MW: Altered expression of Fas receptor on alveolar macrophages and inflammatory effects of soluble Fas ligand following blunt chest trauma. *Shock* 35(6):610–617, 2011.
- Seitz DH, Perl M, Mangold S, Neddermann A, Braumüller ST, Zhou S, Bachem MG, Huber-Lang MS, Knöferl MW: Pulmonary contusion induces alveolar type 2 epithelial cell apoptosis: role of alveolar macrophages and neutrophils. *Shock* 30(5):537–544, 2008.
- Park DR, Thomsen AR, Frevert CW, Pham U, Skerrett SJ, Kiener PA, Liles WC: Fas (CD95) induces proinflammatory cytokine responses by human monocytes and monocyte-derived macrophages. *J Immunol* 170(12):6209–6216, 2003.
- Carey MA, Bradbury JA, Rebolloso YD, Graves JP, Zeldin DC, Germolec DR: Pharmacologic inhibition of COX-1 and COX-2 in influenza A viral infection in mice. *PLoS One* 5(7):e11610, 2010.
- Southard R, Ghosh S, Hilliard J, Davis C, Mazuski C, Walton A, Hotchkiss R: Pulmonary contusion is associated with toll-like receptor 4 upregulation and decreased susceptibility to pseudomonas pneumonia in a mouse model. *Shock* 37(6):629–633, 2012.
- Niesler U, Palmer A, Fröba JS, Braumüller ST, Zhou S, Gebhard F, Knöferl MW, Seitz DH: Role of alveolar macrophages in the regulation of local and systemic inflammation after lung contusion. *Journal of Trauma* 76(2):386–393, 2014.
- Suresh MV, Yu B, Machado-Aranda D, Bender MD, Ochoa-Frongia L, Helinski JD, Davidson BA, Knight PR, Hogaboam CM, Moore BB, et al.: Role of macrophage chemoattractant protein-1 in acute inflammation after lung contusion. *Am J Respir Cell Mol Biol* 46(6):797–806, 2012.

36. Knöferl MW, Liener UC, Seitz DH, Perl M, Brückner UB, Kinzl L, Gebhard F: Cardiopulmonary, histological, and inflammatory alterations after lung contusion in a novel mouse model of blunt chest trauma. *Shock* 19(6):519–525, 2003.
37. Knöferl MW, Liener UC, Perl M, Brückner UB, Kinzl L, Gebhard F: Blunt chest trauma induces delayed splenic immunosuppression. *Shock* 22(1):51–56, 2004.
38. Jordan S, Mitchell JA, Quinlan GJ, Goldstraw P, Evans TW: The pathogenesis of lung injury following pulmonary resection. *Eur Respir J* 15(4):790–799, 2000.
39. Ng CS, Wan S, Yim AP, Arifi AA: Pulmonary dysfunction after cardiac surgery. *Chest* 121(4):1269–1277, 2002.
40. Laubach VE, Kron IL: Pulmonary inflammation after lung transplantation. *Surgery* 146(1):1–4, 2009.
41. Granton J: Update of early respiratory failure in the lung transplant recipient. *Curr Opin Crit Care* 12(1):19–24, 2006.
42. Chatila WM, Furukawa S, Gaughan JP, Criner GJ: Respiratory failure after lung transplantation. *Chest* 123(1):165–173, 2003.
43. Ishii M, Suzuki Y, Takeshita K, Miyao N, Kudo H, Hiraoka R, Nishio K, Sato N, Naoki K, Aoki T, et al.: Inhibition of c-Jun NH2-terminal kinase activity improves ischemia/reperfusion injury in rat lungs. *J Immunol* 172(4):2569–2577, 2004.
44. Naidu BV, Krishnadason B, Farivar AS, Woolley SM, Thomas R, van RN, Verrier ED, Mulligan MS: Early activation of the alveolar macrophage is critical to the development of lung ischemia-reperfusion injury. *J Thorac Cardiovasc Surg* 126(1):200–207, 2003.
45. Nakamura T, Abu-Dahab R, Menger MD, Schafer U, Vollmar B, Wada H, Lehr CM, Schafers HJ: Depletion of alveolar macrophages by clodronate-liposomes aggravates ischemia-reperfusion injury of the lung. *J Heart Lung Transplant* 24(1):38–45, 2005.
46. Sharma AK, LaPar DJ, Stone ML, Zhao Y, Kron IL, Laubach VE: Receptor for advanced glycation end products (RAGE) on iNKT cells mediates lung ischemia-reperfusion injury. *Am J Transplant* 13(9):2255–2267, 2013.
47. Yang H, Hreggvidsdottir HS, Palmblad K, Wang H, Ochani M, Li J, Lu B, Chavan S, Rosas-Ballina M, Al-Abed Y, et al.: A critical cysteine is required for HMGB1 binding to Toll-like receptor 4 and activation of macrophage cytokine release. *Proc Natl Acad Sci U S A* 107(26):11942–11947, 2010.
48. Deng Y, Yang Z, Gao Y, Xu H, Zheng B, Jiang M, Xu J, He Z, Wang X: Toll-like receptor 4 mediates acute lung injury induced by high mobility group box-1. *PLoS One* 8(5):e64375, 2013.
49. Liu P, Xu B, Hock CE: Inhibition of nitric oxide synthesis by L-name exacerbates acute lung injury induced by hepatic ischemia-reperfusion. *Shock* 16(3):211–217, 2001.
50. Souza AL Jr, Poggetti RS, Fontes B, Birolini D: Gut ischemia/reperfusion activates lung macrophages for tumor necrosis factor and hydrogen peroxide production. *J Trauma* 49(2):232–236, 2000.
51. LaNoue JL, Iglesias JL, Rogers TE, Kim LT, Meng Y, Myers SI, Turnage RH: Alveolar macrophage response to remote organ injury. *Shock* 9(4):261–265, 1998.
52. Moraes LB, Murakami AH, Fontes B, Poggetti RS, van RN, Younes RN, Heimbecker AM, Birolini D: Gut ischemia/reperfusion induced acute lung injury is an alveolar macrophage dependent event. *J Trauma* 64(5):1196–1200, 2008.
53. McGuire GM, Liu P, Jaeschke H: Neutrophil-induced lung damage after hepatic ischemia and endotoxemia. *Free Radic Biol Med* 20(2):189–197, 1996.
54. Silva RC, Landgraf MA, Corrêa-Costa M, Semedo P, Cenedeze MA, Pacheco-Silva A, Landgraf RG, Câmara NOS: Acute kidney injury reduces phagocytic and microbicidal capacities of alveolar macrophages. *Cell Physiol Biochem* 31(2–3):179–188, 2013.
55. Fan J, Li Y, Levy RM, Fan JJ, Hackam DJ, Vodovotz Y, Yang H, Tracey KJ, Billiar TR, Wilson MA: Hemorrhagic shock induces NAD(P)H oxidase activation in neutrophils: role of HMGB1-TLR4 signaling. *J Immunol* 178(10):6573–6580, 2007.
56. Powers KA, Woo J, Khadaroo RG, Papia G, Kapus A, Rotstein OD: Hypertonic resuscitation of hemorrhagic shock upregulates the anti-inflammatory response by alveolar macrophages. *Surgery* 134(2):312–318, 2003.
57. Fan J, Kapus A, Li YH, Rizoli S, Marshall JC, Rotstein OD: Priming for enhanced alveolar fibrin deposition after hemorrhagic shock: role of tumor necrosis factor. *Am J Respir Cell Mol Biol* 22(4):412–421, 2000.
58. Powers KA, Szaszi K, Khadaroo RG, Tawadros PS, Marshall JC, Kapus A, Rotstein OD: Oxidative stress generated by hemorrhagic shock recruits Toll-like receptor 4 to the plasma membrane in macrophages. *J Exp Med* 203(8):1951–1961, 2006.
59. Khadaroo RG, He R, Parodo J, Powers KA, Marshall JC, Kapus A, Rotstein OD: The role of the Src family of tyrosine kinases after oxidant-induced lung injury in vivo. *Surgery* 136(2):483–488, 2004.
60. Naziri W, Pietsch JD, Appel SH, Cheadle WG, Bergamini TM, Polk HC Jr: Hemorrhagic shock-induced alterations in circulating and bronchoalveolar macrophage nitric oxide production. *J Surg Res* 59(1):146–152, 1995.
61. Lee H, Pespeni M, Roux J, Dennery PA, Matthay MA, Pittet JF: HO-1 induction restores c-AMP-dependent lung epithelial fluid transport following severe hemorrhage in rats. *FASEB J* 19(2):287–289, 2005.
62. Papia G, Fan J, Kapus A, Szaszi K, Marshall JC, Tawadros P, Ailenberg M, Rotstein OD: Altered inhibitory kappaB expression in LPS-stimulated alveolar macrophages following resuscitated hemorrhagic shock. *Shock* 35(2):171–177, 2011.
63. Fan J, Li Y, Vodovotz Y, Billiar TR, Wilson MA: Hemorrhagic shock-activated neutrophils augment TLR4 signaling-induced TLR2 upregulation in alveolar macrophages: role in hemorrhage-primed lung inflammation. *Am J Physiol Lung Cell Mol Physiol* 290(4):L738–L746, 2006.
64. Molina PE, Bagby GJ, Stahls P: Hemorrhage alters neuroendocrine, hemodynamic, and compartment-specific TNF responses to LPS. *Shock* 16(6):459–465, 2001.
65. Fan J, Marshall JC, Jimenez M, Shek PN, Zagorski J, Rotstein OD: Hemorrhagic shock primes for increased expression of cytokine-induced neutrophil chemoattractant in the lung: role in pulmonary inflammation following lipopolysaccharide. *J Immunol* 161(1):440–447, 1998.
66. Xu P, Wen Z, Shi X, Li Y, Fan L, Xiang M, Li A, Scott MJ, Xiao G, Li S, et al.: Hemorrhagic shock augments Nlrp3 inflammasome activation in the lung through impaired pyrin induction. *J Immunol* 190(10):5247–5255, 2013.
67. Althaus M, Clauss WG, Fronius M: Amiloride-sensitive sodium channels and pulmonary edema. *Pulm Med* 2011:830320, 2011.
68. Fan J, Kapus A, Marsden PA, Li YH, Oreopoulos G, Marshall JC, Frantz S, Kelly RA, Medzhitov R, Rotstein OD: Regulation of Toll-like receptor 4 expression in the lung following hemorrhagic shock and lipopolysaccharide. *J Immunol* 168(10):5252–5259, 2002.
69. Fan J, Li Y, Vodovotz Y, Billiar TR, Wilson MA: Neutrophil NAD(P)H oxidase is required for hemorrhagic shock-enhanced TLR2 up-regulation in alveolar macrophages in response to LPS. *Shock* 28(2):213–218, 2007.
70. Williams FN, Herndon DN, Hawkins HK, Lee JO, Cox RA, Kulp GA, Finnerty CC, Chinkes DL, Jeschke MG: The leading causes of death after burn injury in a single pediatric burn center. *Crit Care* 13(6):R183, 2009.
71. Brusselaers N, Monstrey S, Vogelaers D, Hoste E, Blot S: Severe burn injury in Europe: a systematic review of the incidence, etiology, morbidity, and mortality. *Crit Care* 14(5):R188, 2010.
72. Bankey PE, Williams JG, Guice KS, Taylor SN: Interleukin-6 production after thermal injury: evidence for nonmacrophage sources in the lung and liver. *Surgery* 118(2):431–438, 1995.
73. Williams JG, Bankey P, Minei JP, McIntyre K, Turbeville T: Burn injury enhances alveolar macrophage endotoxin sensitivity. *J Burn Care Rehabil* 15(6):493–498, 1994.
74. Li N, Hu X, Liu Y, Wang Y, Wang Y, Liu J, Cai W, Bai X, Zhu X, Han J, et al.: Systemic inflammatory responses and multiple organ dysfunction syndrome following skin burn wound and *Pseudomonas aeruginosa* infection in mice. *Shock* 40(2):152–159, 2013.
75. Loose LD, Turinsky J: Macrophage dysfunction after burn injury. *Infect Immun* 26(1):157–162, 1979.
76. Loose LD, Turinsky J: Depression of the respiratory burst in alveolar and peritoneal macrophages after thermal injury. *Infect Immun* 30(3):718–722, 1980.
77. Davis KA, Santaniello JM, He LK, Muthu K, Sen S, Jones SB, Gamelli RL, Shankar R: Burn injury and pulmonary sepsis: development of a clinically relevant model. *J Trauma* 56(2):272–278, 2004.
78. Loose LD, Megirian R, Turinsky J: Biochemical and functional alterations in macrophages after thermal injury. *Infect Immun* 44(3):554–558, 1984.