

COMMONWEALTH OF AUSTRALIA
Copyright Regulations 1969
WARNING

**This material has been copied and communicated to
you by or on behalf of the University of Sydney
pursuant to Part VB of the Copyright Act 1968. (The Act).**

**The material in this communication may be subject
to copyright under the Act. Any further copying or
communication of this material by you may be the
subject of copyright protection under the Act.**

Do not remove this notice.

Pharmacogenomics

Michael Murray

Pharmacogenomics and Drug Development
Discipline of Pharmacology


Focus of this lecture

- Drug toxicities and adverse drug reactions (ADRs)
- Type A (predictable) ADRs
 - *Genetic factors and mechanisms*
- Type B (drug hypersensitivity) ADRs
 - *Reactive metabolites and MHC molecules*
- Development of pharmacogenomic tests to prevent ADRs

All Chemicals are Potentially Toxic

- “All substances are poisonous; there are none that are not poisons. The right DOSE differentiates a poison and a remedy.”
 - Paracelsus (1493-1541)
- This is true for most ADRs but some are unrelated to dose
- ADRs are responses to drugs that are unintended and that occur in man at near-normal therapeutic doses

Therapeutic window



- Therapeutic index is the ratio of the toxic dose to the therapeutic dose
- Narrow for drugs like digoxin, warfarin, clozapine and lithium
- Therapeutic drug monitoring important for toxic drugs

Drugs and iatrogenic injury

- Drug treatment is itself a major cause of adverse events
- Recent estimate: up to 2 million patients are hospitalised annually in the US because of severe ADRs
- ADRs are considered to be the fourth to sixth leading cause of death in the US
- ~15% of courses of drug treatment may lead to an ADR

Types of ADRs



- **Type A**
 - Predictable
 - Usually related to the principal pharmacological actions of the drug
 - However, higher doses of some drugs cause predictable ADRs that are unrelated to their pharmacological actions
 - eg paracetamol hepatotoxicity and doxorubicin cardiotoxicity
- Treatment is to reduce dose or withhold drug

Types of ADRs

- **Type B**
 - Unpredictable or idiosyncratic
 - Less common than type A reactions
 - Unrelated to the principal pharmacological actions of the drug
 - High mortality
- Treatment is to withhold drug and avoid rechallenge, even with very low doses

ADRs and the therapeutic window


Type A



Type B


Pharmacogenetic variation (ADME factors)

- Transporter proteins and biotransformation enzymes determine drug influx into and efflux from cells and the duration of drug action
- Genes encoding these proteins are often highly polymorphic
- Defective proteins may promote drug accumulation and toxicity



Type A ADRs – pharmacogenetic background

- The human genome contains 3.1 billion nucleotide bases
- Estimated total number of genes between 20,000 and 25,000
- 99.9% of nucleotide bases are identical between individuals
- Alleles are alternate forms of the same gene or genetic locus



Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

Common polymorphisms - definitions

- Polymorphisms occur in at least 1% of population and exhibit a *stable inheritance pattern*
- Even though genomes are 99.9% identical this still allows 3 to 10 million sequence differences between any 2 genomes

Single Nucleotide Polymorphism (SNP):

GAATT**T**AAG
GAATT**C**AAG


Variable number of tandem repeats (VNTR): (here repeat of 2bp-CA)

(4) CACACACA
(7) CACACACACACACA
(6) CACACACACACA

Insertions/Deletions:

GAAA -TCCAAG
GAAAT**T**CCAAG


SNPs and their impact on gene function



- Codons are base triplets that encode an amino acid
- Coding region variants usually alter amino acid sequence of the encoded protein
- Regulatory region variants usually alter the amount of the protein produced


Pharmacogenomics and Type A ADRs: Predictable toxicity related to drug action – anticancer agents

- Most established anticancer agents have low therapeutic indexes
- 25-50% of cancer patients may not be treated effectively by chemotherapy
- Unpredictable toxicity in many patients, but also
- Under-treatment that leads to failure of therapy is quite common
- To date there has been a strong focus on variants of transporters and drug metabolising enzymes that affect drug disposition




Dosing of anticancer agents: current approaches

- Body surface area used to calculate initial dose
- Adjust dosage upwards until toxicity, then decrease
- If toxicity occurs at initial dose level adjust downwards
- Severe toxicity: dose interruption/cessation (risk of under-treatment and failure)
- NOTE: No patient- or tumour-specific factors considered in dose determination




Thiopurine methyltransferase (TPMT)

- 6-mercaptopurine (6MP) used for treatment of ALL (acute lymphoblastic leukaemia)
- Azathioprine is a prodrug for 6MP used in immunosuppression
- Cytotoxic effect due to formation of 6-thioguanine nucleotides (6TGN) that inhibit gene replication
- TPMT deactivates 6MP so when it is deficient 6TGN are over-produced leading to ADRs




TPMT pharmacogenomics

- Can measure 6TGN or TPMT activity in erythrocytes as a surrogate for overall TPMT status
- Wide inter-patient variation in activity
- 1 in 300 patients homozygous for low activity alleles
- Now genotyping for TPMT alleles is also possible




TPMT pharmacogenomics

- Patients who have low activity: frequent dose adjustments with AZA
- Those with normal activity: tolerate usual doses of AZA
- Those with high activity: low incidence of neutropenia, ?possible under-dosing




TPMT pharmacogenomics and therapeutic window

- Low TPMT activity
 - Defective alleles
 - Over-production of toxic 6TGN with standard doses of AZA
 - Lower production of non-toxic 6MMPR and 6MMP




Irinotecan

- Topoisomerase inhibitor
- Topoisomerase repairs DNA that gets tangled in rapidly dividing cells
- Used in colorectal cancer but neutropaenia and delayed diarrhoea occurs in up to 40% of patients
- May require hospitalisation
- Irinotecan biotransformation is complex. The metabolite SN-38 is the active cytotoxic agent



UGT1A1 variants and SN-38 conjugation

- UGT1A1 mediates SN-38 conjugation
- Transcription is initiated at the TATA box in the UGT1A1 promoter
- *28 variant has 7 TA repeats instead of 6 (VNTR polymorphism)
- *28 variant – less transcription – less enzyme
- *28 homozygotes exhibit decreased SN-38 conjugation capacity and higher grade neutropaenia
- Another variant – the *6 allele – carries a coding region SNP – results in the amino acid substitution Gly71Arg – the encoded enzyme has decreased function



(Desai *et al.*, Oncogene, 2002;
Tukey *et al.*, Mol Pharmacol, 2002)



An exon is a DNA region that usually codes for protein (numbered above)

Ethnicity-related incidence of UGT1A1 variants

- Frequency of *28 allele >40% in sub-Saharan Africans and 30-40% in Caucasians, but lower in Asians (Haga *et al.*, Pharmacogenet Genom 2006)
- However, the low activity *6 allele is as frequent as *28 in Japanese, but is absent in Caucasians and Africans
- Whereas Caucasian *28 homozygotes are at risk from irinotecan toxicity susceptible Asians may carry either *28/*28, *6/*6 or *6/*28
- The US FDA and Japanese/Singaporean regulatory bodies have different warnings for the use of irinotecan in relation to UGT1A1 variants and ADRs


Type A ADRs unrelated to drug action: paracetamol hepatotoxicity

- Major metabolites are stable, non-toxic and readily eliminated (left side of diagram)
- Normally only ~5% of paracetamol is metabolized by CYP2E1 to the quinoid intermediate (right side)
- This amount of the intermediate is detoxified by glutathione.
- Alcohol increases formation of the toxic intermediate.
- High doses of paracetamol forms more of the toxic intermediate, depleting glutathione and causing hepatic injury.



glutathione

Nomogram to predict liver damage from plasma paracetamol concentrations




- Rumack-Matthew nomogram - doses greater than 150 mg/kg/24 hr (>10 g).
- Potential for hepatotoxicity after acute overdose is best predicted from serum paracetamol concentrations plotted against the time elapsed since ingestion.
- This is only a useful predictor if the person gets to hospital within 10 hrs of overdose.
- Treatment: N-acetylcysteine


<http://members.ozemail.com.au/~ouad/doc/images/Paracetamol%20nomogram.gif>

Type A ADRs unrelated to drug action: doxorubicin cardiotoxicity

Some of the clinically used anthracycline anti-tumor drugs.



www.chem2.bham.ac.uk/labs/hannon/3wayjunction.htm




- Anthracycline type anticancer agents
- Bind to DNA and prevent replication
- Dose-related cardiotoxicity
- Heart failure


<http://stripe.colorado.edu/~kocht/Anthracycline.drugs.GIF>

Doxorubicin toxicity

- Anthracyclines accumulate in the heart
- Accumulation produces oxidative stress and lipid peroxidation
- Heart tissue is very susceptible to oxidative stress
- Low levels of glutathione and antioxidants



Overview of lipid Peroxidation



<http://www.biochemsoctrans.org/bst/029/0358/bst0290358a01.gif>

- Radicals initiate lipid peroxidation
- After oxygen uptake reactive peroxy radicals form
- Radicals are propagated
- Liberate reactive breakdown products
- Also directly attack membranes and proteins

Predictability of Type A ADRs


- Paracetamol: ingestion of high dose (range)
 - *Predictable hepatotoxicity*
- Doxorubicin: ingestion of cumulative dose (range)
 - *Predictable cardiotoxicity*
- Irinotecan: presence of UGT1A variants
 - *Not yet fully predictive of neutropaenia or other toxicities*

Types of ADRs

- **Type B**
 - Unpredictable or idiosyncratic
 - Less common than type A reactions
 - Unrelated to the principal pharmacological actions of the drug
 - High mortality
- Treatment is to withhold drug and avoid rechallenge, even with very low doses


Type B ADRs

- Up to one-third of ADRs are due to unpredictable hypersensitivities
- Concept: drugs are not themselves immunogenic
- Instead a drug or metabolite interacts as a hapten with a carrier protein to form a stable conjugate
- Produces a modified protein no longer recognised as “self”
- Production of drug-specific cytotoxic T-cells



The major histocompatibility complex (MHC)


- The MHC system is a family of highly polymorphic genes that encode the human lymphocyte antigens (HLA)
- One of these genes (HLA-B) alone has >1000 alleles
- HLA molecules bind short peptides formed by intracellular proteolysis and present them on the surface of specialised cells to be detected by cytotoxic T-cells
- Peptide binding specificity is dictated by the amino acid sequence of the HLA binding groove
- So some peptides will only bind if the individual carries the appropriate HLA allele



Nature Reviews | Immunology



Abacavir

- A reverse transcriptase inhibitor used to treat HIV-AIDs
- Has a number of common toxicities that are not severe
- However, it also produces a hypersensitivity reaction in 5-10% of patients in the first few weeks of treatment
- Fever, dermatological, GI and respiratory symptoms that eventually resolve on removal but return if drug is resumed
- Occasionally fatal




Abacavir hypersensitivity syndrome

- associated with an HLA-B allele - several studies have demonstrated that testing for this allele is *predictive* (Mallal *et al.*, Lancet, 2002)
- Abacavir binds and distorts the binding cleft of the HLA-B molecule – which alters the peptides that can bind
- This exposes the self-tolerant T-cell compartment to modified peptides (Illing *et al.*, Nature, 2012)



Stern, MIT 2013

Carbamazepine and severe cutaneous reactions



- Carbamazepine is an anticonvulsant used to treat seizures
- Stevens-Johnson-Syndrome (SJS) and toxic epidermal necrolysis (TEN) are severe cutaneous reactions induced by carbamazepine that have significant mortalities
- Strong association with an HLA-B allele in Chinese subjects (Chung *et al.*, Nature, 2004) - prior genotyping recommended
- In Europe the incidence of SJS/TEN from carbamazepine is ~5-6% (~30% in Chinese) but incidence of the HLA-B allele is low
- Instead an HLA-A allele has been implicated in Japanese and Europeans
- Note: the HLA-B alleles implicated in abacavir and carbamazepine hypersensitivities are different

Whether to develop a pharmacogenetic test

(Based on Maitland *et al.*, TIPS, 2006)

- If there is unexplained variability in drug toxicity or efficacy
- Toxicity seems to be associated with a polymorphism that is common in the population
- A pharmacogenomic mechanism is implicated in toxicity
- Finally, in order to change practice, evidence from prospective studies is required
- The test should predict ADRs when the polymorphism is present and should predict no ADR if it is absent

Key points

- Simple ADRs (predictable) are often related to drug concentrations in serum
- Usually, but not always, an extension of pharmacological actions
- Defective ADME genes may account for many such ADRs (possible genotyping to detect susceptible individuals)
- Drug hypersensitivities usually have an immune basis
- Genetic variation in specific HLA genes now associated with type B ADRs with certain drugs
- Genetic tests now being developed to identify at risk individuals before therapy with certain drugs that have major toxicity profiles
- In future the unpredictable nature of Type B reactions may be modified

References

- Casarett and Doull's Toxicology, Ch 12 and Ch 13
- TPMT p277
- Drug hypersensitivity: *Curr Opin Allerg Clin Immunol* 7: 317, 2007 and *Curr Opin Immunol* 25: 81, 2013 (more detailed)
- UGT1A1 variants: *Ann Oncol* 19: 2089, 2008 and *Mol Pharmacol* 62: 446, 2002