ECON 124: Midterm #2

Due on Jul 9, 2025

Dr. Deniz Baglan

Alejandro Ouslan

Problem 1

Use the data in **FERTIL2.XLSX** to answer this question.

(a) Estimate the model

$$children = \beta_0 + \beta_1 aqe + \beta_2 aqe^2 \beta_3 educ + \beta_4 electricity + \beta_5 urban + \epsilon$$

And report the usual and heteroskedasticity-robust standard errors. Are the robust standard errors always bigger than the non robust ones?

- (b) Add the three religious dummy variables and test whether they are jointly significant. What are the p-values for the nonrobust and robust tests?
- (c) From the regression in part (b), obtain the fitted values \hat{y} and the residuals, $\hat{\epsilon}$. Regress $\hat{\epsilon}^2 \sim \hat{y}$, and $\hat{\epsilon}^2 \sim \hat{y}^2$ and test the joint significance of the two regressors.

Problem 2

Use the data set Movies

Does viewing a violent movie lead to violent behavior? If so, the incidence of violent crimes, such as assaults, should rise following the release of a violent movie that attracts many viewers. Alternatively, movie viewing may substitute for other activities (such as alcohol consumption) that lead to violent behavior, so that assaults should fall when more viewers are attracted to the cinema. The dataset includes weekend U.S. attendance for strongly violent movies (such as Hannibal), mildly violent movies (such as Spider-Man), and nonviolent movies (such as Finding Nemo). The dataset also includes a count of the number of assaults for the same weekend in a subset of counties in the United States. Finally, the dataset includes indicators for year, month, whether the weekend is a holiday, and various measures of the weather.

(a) (i) Regress the logarithm of the number of assaults [ln_(assaults) = ln(assaults)] on the year and month indicators. Is there evidence of seasonality in assaults? That is, do there tend to be more assaults in some months than others? Explain.

Answer: It seems that the robust standard errors are generally larger than the non robust ones, but not necessarily always the case.

Table 1: OLS Regression Results: Children ~Age, Age², Education, Electricity, Urban

Variable	Coef.	Std. Err.	\mathbf{t}	$\mathbf{P} > t $	[0.025]	0.975]
const	-4.2225	0.240	-17.580	0.000	-4.693	-3.752
age	0.3409	0.017	20.652	0.000	0.309	0.373
age^2	-0.0027	0.000	-10.086	0.000	-0.003	-0.002
education	-0.0752	0.006	-11.948	0.000	-0.088	-0.063
electricity	-0.3100	0.069	-4.493	0.000	-0.445	-0.175
urban	-0.2000	0.047	-4.301	0.000	-0.291	-0.109
Model state	istics:					
R-squared		0.573				
Adj. R-squ	ared	0.573				
F-statistic		1170	(Prob F-statistic = 0.000)			
No. Observations		4358				
Df Residua	ls	4352				
Df Model		5				
Log-Likelih	lood	-7806.3				
AIC		1.562e + 04				
BIC		1.566e + 04				
Durbin-Wa	tson: 1.88	3				
Omnibus:	203 155 P	rob(Omnibus	a) • 0 000			

Omnibus: 203.155, Prob(Omnibus): 0.000

Jarque-Bera (JB): 715.135, Prob(JB): 5.13×10^{-156}

Skew: 0.014, Kurtosis: 4.984

Cond. No.: 1.07e+04

Table 2: F-test Results				
Statistic	Value	Notes		
F-statistic	2.196	-		
p-value	0.0864	-		
Degrees of Freedom (denominator)	4350	-		
Degrees of Freedom (numerator)	3	-		

Table 3: OLS Regression Results: Children ~Age, Age², Education, Electricity, Urban

Variable	Coef.	Std. Err.	z	P > z	[0.025]	0.975]
const	-4.2225	0.244	-17.316	0.000	-4.700	-3.745
age	0.3409	0.019	17.780	0.000	0.303	0.379
$ m age^2$	-0.0027	0.000	-7.821	0.000	-0.003	-0.002
education	-0.0752	0.006	-11.927	0.000	-0.088	-0.063
electricity	-0.3100	0.064	-4.848	0.000	-0.435	-0.185
urban	-0.2000	0.045	-4.399	0.000	-0.289	-0.111
Model statistics:						
R-squared		0.573				
Adj. R-squared		0.573				
F-statistic		1161	(Prob F-statistic = 0.000)			
No. Observations		4358				
Df Residua	ls	4352				
Df Model		5				
Log-Likelih	ood	-7806.3				
AIC		1.562e + 04				
BIC		1.566e + 04				
Durbin-Watson: 1.883						

Omnibus: 203.155, Prob(Omnibus): 0.000

Jarque-Bera (JB): 715.135, Prob(JB): 5.13×10^{-156}

Skew: 0.014, Kurtosis: 4.984

Cond. No.: 1.07e+04

Table 4: F-test Results				
Statistic	Value	Notes		
F-statistic	2.156	-		
p-value	0.0911	-		
Degrees of Freedom (denominator)	4350	-		
Degrees of Freedom (numerator)	3	-		

Python Code

```
import polars as pl
import statsmodels.formula.api as smf
# Question
# 1a
df = pl.read_excel("data/fertil2.xlsx")
df = df.select(
    pl.col(
        "children",
            "age",
            "educ",
            "electric",
            "urban",
            "spirit",
            "protest",
            "catholic",
        ]
    )
df = df.with_columns(age2=pl.col("age") ** 2)
df = df.to_pandas()
model = smf.ols("children ~ age + age2 + educ + electric + urban", data=df).fit()
print(model.summary())
model = smf.ols("children ~ age + age2 + educ + electric + urban", data=df).fit
    cov_type="HC1"
print(model.summary())
```

- (ii) Regress total movie attendance (attend = attend_v + attend_m + attend_n) on the year and month indicators. Is there evidence of seasonality in movie attendance? Explain.
- (b) Regress ln_assaults on attend_v, attend_m, attend_n, the year and month indicators, and the weather and holiday control variables available in the data set
 - (i) Based on the regression, does viewing a strongly violent movie increase or decrease assaults? By how much? Is the estimated effect statistically significant?
 - (ii) Does attendance at strongly violent movies affect assaults differently than attendance at moderately violent movies? Differently than attendance at nonviolent movies?
 - (iii) A strongly violent blockbuster movie is released, and the weekend's attendance at strongly violent movies increases by 6 million; meanwhile, attendance falls by 2 million for moderately violent movies and by 1 million for nonviolent movies. What is the predicted effect on assaults? Construct a 95% confidence interval for the change in assaults.
 - (iv) It is difficult to control for all the variables that affect assaults and that might be correlated with movie attendance. For example, the effect of the weather on assaults and movie attendance is only crudely approximated by the weather variables in the data set. However, the data set does include a set of instruments, pr_attend_v, pr_attend_m, and pr_attend_n, that are correlated with

attendance but are (arguably) uncorrelated with weekend-specific factors (such as the weather) that affect both assaults and movie attendance. These instruments use historical attendance patterns, not information on a particular weekend, to predict a film's attendance in a given weekend. For example, if a film's attendance is high in the second week of its release, then this can be used to predict that its attendance was also high in the first week of its release. (The details of the construction of these instruments are available in the Dahl and DellaVigna's paper on Canvas) Run the regression from part (b) (including year, month, holiday, and weather controls) but now using pr_attend_v, pr_attend_m, and pr_attend_n as instruments for attend_v, attend_m, and attend_n. Use this regression to answer (b)(i)-(b)(iii).

- (v) It is difficult to control for all the variables that affect assaults and that might be correlated with movie attendance. For example, the effect of the weather on assaults and movie attendance is only crudely approximated by the weather variables in the data set. However, the data set does include a set of instruments, pr_attend_v, pr_attend_m, and pr_attend_n, that are correlated with attendance but are (arguably) uncorrelated with weekend-specific factors (such as the weather) that affect both assaults and movie attendance. These instruments use historical attendance patterns, not information on a particular weekend, to predict a film's attendance in a given weekend. For example, if a film's attendance is high in the second week of its release, then this can be used to predict that its attendance was also high in the first week of its release. (The details of the construction of these instruments are available in the Dahl and DellaVigna's paper on Canvas) Run the regression from part (b) (including year, month, holiday, and weather controls) but now using pr_attend_v, pr_attend_m, and pr_attend_n as instruments for attend_v, attend_m, and attend_n. Use this regression to answer (b)(i)-(b)(iii).
- (vi) Based on your analysis, what do you conclude about the effect of violent movies on (short-run) violent behavior?

Problem 3

We examined Koop and Tobias's data on wages, education, ability, and so on. We considered the model.

```
\begin{split} \ln wage &= \beta_0 + \beta_1 educ + \beta_2 ability + \beta_3 experience \\ &+ \beta_4 mother Educ + \beta_5 Father Educ + \beta_6 broken \\ &+ \beta_7 sinlings + \epsilon \end{split}
```

(a) We are interested in possible non-linearities in the effect of education on ln Wage. (Koop and Tobias focused on experience. As before, we are not attempting to replicate their results.) A histogram of the education variable shows values from 9 to 20, a spike at 12 years (high school graduation), and a second at 15. Consider aggregating the education variable into a set of dummy variables:

```
\mathrm{HS} = 1 if \mathrm{Educ} \leq 12, 0 otherwise \mathrm{Col} = 1 if 12 < \mathrm{Educ} \leq 16, 0 otherwise \mathrm{Grad} = 1 if \mathrm{Educ} > 16, 0 otherwise
```

replace Educ in the model with (Col, Grad), making high school (HS) the base category, and recompute the model. Report all results. How do the results change? Based on your results, what is the marginal value of a college degree? What is the marginal impact on ln Wage of a graduate degree?

(b) The aggregation in part (a) actually loses quite a bit of information. Another way to introduce non-linearity in education is through the function itself. Add $educ^2$ to the equation in part (a) and recompute the model. Again, report all results. What changes are suggested? Test the hypothesis that the quadratic term in the equation is not needed—that is, that its coefficient is zero. Based on your results, sketch a profile of log wages as a function of education.

(c) One might suspect that the value of education is enhanced by greater ability. We could examine this effect by introducing an interaction of the two variables in the equation. Add the variable

$$EducAb = Educ \times ability$$

- to the base model in part a. Now, what is the marginal value of an additional year of education? The sample mean value of ability is 0.052374. Compute a confidence interval for the marginal impact on ln Wage of an additional year of education for a person of average ability.
- (d) Combine the models in (b) and (c). Add both $educ^2$ and EducAb to the base model in the beginning of the question and re-estimate. As before, report all results and describe your findings. If we define low ability as less than the mean and high ability as greater than the mean, the sample averages are -0.798563 for the 7,864 low-ability individuals in the sample and +0.717891 for the 10,055 high-ability individuals in the sample. Using the formulation in part (b), with this new functional form, sketch, describe, and compare the log wage profiles for low- and high-ability individuals.
- (e) Suppose that you are now given the following regression model:

$$\begin{split} \ln(\text{wage}) &= \beta_0 + \beta_1 \text{educ} \times \mathbf{1}(\text{educ} < \tau) + \beta_2 \text{educ} \times \mathbf{1}(\text{educ} \ge \tau) + \beta_3 \text{exp} \\ &+ \beta_4 \text{MotherEduc} + \beta_5 \text{FatherEduc} + \beta_6 \text{broken} \\ &+ \beta_7 \text{siblings} + \epsilon \end{split}$$

where τ is the threshold parameter, and

$$\mathbf{1}(\mathrm{Educ} < \tau) = \begin{cases} 1 & \text{if Educ} < \tau, \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad \mathbf{1}(\mathrm{Educ} \geq \tau) = \begin{cases} 1 & \text{if Educ} \geq \tau, \\ 0 & \text{otherwise.} \end{cases}$$

Problem 4

Using the California test score date, estimate the regression below using Nonlinear Least Squares. Report you coefficient estimates and standard errors.

$$TestScore = \beta_0(1 - e^{\beta_1(income - \beta_2)}) + \epsilon$$

Problem 5

Use the **Consumption.xlsx**. We have previously estimated the nonlinear consuption function below using nonlinear least squares in class:

$$C = \alpha + \beta Y^{\gamma} + \epsilon$$

Where C is the real consumption and Y is the real disposable income. Alternatively, we can assume that the error term has a normal distribution and estimate the nonlinear regression above using the maximum likeligood estimation (MLE) approach, In particular, the MLE approach maximizes the log-likelihood function given by:

$$L(\alpha, \beta, \gamma, \sigma^2) = -\frac{n}{2}\log(\sigma^2) - \log(2\pi) - \frac{1}{2\sigma^2} \sum_{i=1}^n (C_i - \alpha - \beta Y^{\gamma})^2$$

Where σ^2 is the variance of the error term. Using a statistical programming language of you choice, estimate the regression model using the maximum likelihood estimation approach. Your estimate are expected to be similar to those in Table 7.1 of Green's textbook. Please submit the following:

- (a) Your code used to preform the estimation.
- (b) The output of your estimation, including the estimated parameters and the error variance.