

# **MATE 5150: Asignacion #3**

Due on Septiembre 26, 2024

*Dr. Pedro Vasquez*

**Alejandro Ouslan**

## Problem 1

For each of the following homogeneous systems of linear equations, find the dimensions of and a basis for the solution space.

$$\begin{aligned}
 x_1 + 2x_2 + x_3 + x_4 &= 0 \\
 x_2 - x_3 + x_4 &= 0 \\
 \begin{bmatrix} 1 & 2 & 1 & 1 & 0 \\ 0 & 1 & -1 & 1 & 0 \end{bmatrix} \\
 x_2 &= x_3 - x_4 \\
 x_1 &= -x_3 - x_4 \\
 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} &= \begin{bmatrix} -1 \\ 1 \\ 1 \\ 0 \end{bmatrix} x_3 + \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \end{bmatrix} x_4
 \end{aligned}$$

The rank of the matrix is 2, and dimension of the solution space is 2. A basis for the solution space is  $\{(1, 1, 1, 0), (1, -1, 0, 1)\}$ .

## Problem 2

Using the results of Exercise 2, find all solutions to the following system of linear equations.

$$\begin{aligned}
 x_1 + 2x_2 + x_3 + x_4 &= 0 \\
 x_2 - x_3 + x_4 &= 0 \\
 K &= \left\{ t_1 \begin{bmatrix} -1 \\ 1 \\ 1 \\ 0 \end{bmatrix} + t_2 \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \end{bmatrix} : t_1, t_2 \in \mathbb{R} \right\}
 \end{aligned}$$

## Problem 3

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  be defined by  $T(a, b, c) = (a + b, 2a - c)$ . Determine  $T^{-1}(1, 11)$ .

$$\begin{aligned}
 T(a, b, c) &= (a + b, 2a - c) \\
 1 &= a + b \quad 11 = 2a - c \\
 b &= 1 - a \quad c = 2a - 11 \\
 (a, b, c) &= (a, 1 - a, 2a - 11) \\
 T^{-1}(1, 11) &= \{(a, 1 - a, 2a - 11) : a \in \mathbb{R}\}
 \end{aligned}$$

## Problem 4

Determine which of the following systems of linear equations has a solution.

$$\begin{aligned}
 x_1 + x_2 + 3x_3 - x_4 &= 0 \\
 x_1 + x_2 + x_3 + x_4 &= 1 \\
 x_1 - 2x_2 + x_3 - x_4 &= 1 \\
 4x_1 + x_2 + 8x_3 - x_4 &= 0
 \end{aligned}$$

$$A = \begin{bmatrix} 1 & 1 & 3 & -1 & 0 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & -2 & 1 & -1 & 1 \\ 4 & 1 & 8 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 & \frac{25}{6} \\ 0 & 1 & 0 & 0 & \frac{4}{3} \\ 0 & 0 & 1 & 0 & -\frac{5}{2} \\ 0 & 0 & 0 & 1 & -2 \end{bmatrix}$$

$$s = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \frac{25}{6} \\ \frac{4}{3} \\ -\frac{5}{2} \\ -2 \end{bmatrix}$$

## Problem 5

u A certain economy consists of two sectors: goods and services. Suppose that 60% of all goods and 30% of all services are used in the production of goods. What proportion of the total economic output is used in the production of goods?

$$P = \frac{.6G + .3S}{G + S} = \frac{.6G + .3S}{T_2}$$

$$P = \frac{.9G}{T_2} = .45, \text{ where } G = 1, S = 1, T_2 = 2$$

## Problem 6

Use Gaussian elimination to solve the following systems of linear equations.

$$\begin{aligned} 2x_1 - 2x_2 - x_3 + 6x_4 - 2x_5 &= 1 \\ x_1 - x_2 + x_3 + 2x_4 - x_5 &= 2 \\ 4x_1 - 4x_2 + 5x_3 + 7x_4 - x_5 &= 6 \end{aligned}$$

$$A = \begin{bmatrix} 2 & -2 & -1 & 6 & -2 & 1 \\ 1 & -1 & 1 & 2 & -1 & 2 \\ 4 & -4 & 5 & 7 & -1 & 6 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 0 & 0 & 23 & -23 \\ 0 & 0 & 1 & 0 & -6 & 7 \\ 0 & 0 & 0 & 1 & -9 & 9 \end{bmatrix}$$

$$S = \begin{bmatrix} x_2 - 23x_5 - 23 \\ x_2 \\ 6x_5 + 7 \\ 9x_5 + 9 \\ x_5 \end{bmatrix}$$

## Problem 7

Let the reduce row echolon form of  $A$  be

$$\begin{bmatrix} 1 & 0 & 2 & 0 & -2 \\ 0 & 1 & -5 & 0 & -3 \\ 0 & 0 & 0 & 1 & 6 \end{bmatrix}$$

Determine  $A$  if the first, second, and forth columns of  $A$  are

$$\begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}, \text{ and } \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix}$$

## Problem 8

Let  $W$  be the subspace of  $M_{2 \times 2}(\mathbb{R})$  consisting of the symmetric  $2 \times 2$  matrices. Then set

$$S = \left\{ \begin{bmatrix} 0 & -1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ 1 & 9 \end{bmatrix}, \begin{bmatrix} 1 & -2 \\ -2 & 4 \end{bmatrix}, \begin{bmatrix} -1 & -2 \\ 2 & -1 \end{bmatrix} \right\}$$

generates  $W$ . Find a subset of  $S$  that is a basis for  $W$ .

$$A_1 = \begin{bmatrix} 0 & -1 \\ -1 & 1 \end{bmatrix},$$

$$A_2 = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix},$$

$$A_3 = \begin{bmatrix} 2 & 1 \\ 1 & 9 \end{bmatrix},$$

$$A_4 = \begin{bmatrix} 1 & -2 \\ -2 & 4 \end{bmatrix},$$

$$A_5 = \begin{bmatrix} -1 & -2 \\ 2 & -1 \end{bmatrix}.$$

$$M = \begin{bmatrix} 0 & 1 & 2 & 1 & -1 \\ -1 & 2 & 1 & -2 & -2 \\ 1 & 3 & 9 & 4 & -1 \end{bmatrix}.$$

$$1. R_2 \leftarrow R_2 - 2R_1, \quad R_3 \leftarrow R_3 - 1R_1$$

$$\Rightarrow M \sim \begin{bmatrix} 0 & 1 & 2 & 1 & -1 \\ -1 & 0 & -3 & -4 & 1 \\ 1 & 3 & 9 & 4 & -1 \end{bmatrix}.$$

$$\Rightarrow \begin{bmatrix} -1 & 0 & -3 & -4 & 1 \\ 0 & 1 & 2 & 1 & -1 \\ 1 & 3 & 9 & 4 & -1 \end{bmatrix}.$$

$$3. R_1 \leftarrow -R_1 \text{ and then } R_3 \leftarrow R_3 + R_1$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 3 & 4 & -1 \\ 0 & 1 & 2 & 1 & -1 \\ 0 & 3 & 12 & 8 & 0 \end{bmatrix}.$$

$$4. R_3 \leftarrow R_3 - 3R_2$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 3 & 4 & -1 \\ 0 & 1 & 2 & 1 & -1 \\ 0 & 0 & 6 & 5 & 3 \end{bmatrix}.$$

This shows there are 3 non-zero rows, hence dimension of span = 3.

Thus, a basis for  $W$  is  $\{A_1, A_2, A_3\}$ .

**Problem 9**

Let  $V$  be a in Exercise 12

$$\begin{aligned}x_1 - x_2 + 2x_4 - 3x_5 + x_6 &= 0 \\2x_1 - x_2 - x_3 + 3x_4 - 4x_5 + 4x_6 &= 0\end{aligned}$$

1. Show that  $S = \{(0, 1, 0, 1, 1, 0), (0, 2, 1, 1, 0, 0)\}$  is a linearly independent subset of  $V$ .
2. Extend  $S$  to a basis for  $V$ .