

# MATE 5150: Elementary Matrix Operations and Systems of Linear Equations

Alejandro Ouslan

Academic Year 2024-2025

## Contents

<b>1</b>	<b>Elementary Matrix Operations and Elementary Matrices</b>	<b>1</b>
1.1	Elementary Operations . . . . .	1
1.1.1	Example Elementary Matrices . . . . .	2
1.2	Properties of Elementary Matrices . . . . .	2
1.2.1	Example . . . . .	2
1.3	Matrix Multiplication and Elementary Matrices . . . . .	2
<b>2</b>	<b>The Rank of a Matrix and Matrix Inverses</b>	<b>3</b>
2.0.1	Example . . . . .	3
2.0.2	Example . . . . .	3
2.0.3	Example . . . . .	4
2.0.4	Example . . . . .	4
2.1	The inverse of a Matrix . . . . .	4
2.1.1	Example . . . . .	5

## 1 Elementary Matrix Operations and Elementary Matrices

In this section, we define the elementary operations that are used throughout the chapter. In subsequent sections, we use these operations to obtain simple computational methods for determining the rank of a linear transformation and the solution of a system of linear equations. There are two types of elementary operations -row operations and column operations. As we will see, the row operations are more useful. They arise from the three operations that can be used to eliminate variables in a system of linear equations.

### 1.1 Elementary Operations

**Definition 1.** Let  $A$  be an  $m \times n$  matrix. Any of the following three operations on the rows [columns] of  $A$  is called an **elementary row [column] operation**:

1. interchanging any two rows [columns] of  $A$ ;
2. multiplying any row [column] of  $A$  by a nonzero scalar;
3. adding any scalar multiple of a row [column] of  $A$  to another row [column] of  $A$ .

Any of these three operations is called an **elementary operation**. Elementary operations are of **type 1**, **type 2**, or **type 3** depending on whether they are obtained by (1), (2), or (3) above.

**Definition 2.** An  $n \times n$  **elementary matrix** is a matrix obtained by performing an elementary operation on  $I_n$ . The elementary matrix is said to be of **type 1**, **type 2**, or **type 3** according to whether the elementary operation performed on  $I_n$  is of type 1, type 2, or type 3.

### 1.1.1 Example Elementary Matrices

Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 & 3 \\ 1 & -2 & 1 \\ 1 & -3 & 1 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix}$$

Find an elementary operation that transforms  $A$  into  $B$  and an elementary operation that transforms  $B$  into  $C$ . By means of several additional operations, transform  $C$  into  $I_3$ .

$$AE = B \quad \text{where} \quad E = \begin{bmatrix} 1 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$EB = C \quad \text{where} \quad E = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$E_1 E_2 E_3 E_4 = I_3 \quad \text{where} \quad E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix}, \quad E_4 = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

## 1.2 Properties of Elementary Matrices

**Theorem 1.** *Elementary matrices are invertible, and the inverse of an elementary matrix is an elementary matrix of the same type.*

*Proof.*

Let  $E$  be an elementary matrix  $n \times n$ . The  $E$  is defined by an elementary operation on  $I_n$ .

□

### 1.2.1 Example

Use the proof in Theorem 1 to obtain the inverse of each of the following elementary matrices:

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

Finding the inverse of each of the elementary matrices in the example above, we have:

$$E = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}, \quad \text{Therefore} \quad A^{-1} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad \text{Therefore} \quad B^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}, \quad \text{Therefore} \quad C^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

## 1.3 Matrix Multiplication and Elementary Matrices

Let  $A$  be an  $m \times n$  matrix. Prove that if  $E$  can be obtained from  $A$  by an elementary row [column] operation, then  $B^T$  can be obtained from  $A^T$  by the corresponding elementary column [row] operation.

*Proof.*

$$\begin{aligned}(E_R B)^T &= (A)^T \\ B^T E_R^T &= A^T\end{aligned}$$

Therefore,  $B^T$  can be obtained from  $A^T$  by the corresponding elementary column operation.  $\square$

## 2 The Rank of a Matrix and Matrix Inverses

In this section, we define the *rank* of a matrix. We then use elementary operations to compute the rank of a matrix and a linear transformation. The section concludes with a procedure for computing the inverse of an invertible matrix.

**Definition 3.** If  $A \in M_{m \times n}(F)$ , We define the **rank** of  $A$ , denoted  $\text{rank}(A)$ , to be the rank of the linear transformation  $L_A : F^n \rightarrow F^m$ .

Every matrix  $A$  is the matrix representation of the linear transformation  $L_A$  with respect to the appropriate standard ordered bases. Thus the rank of the linear transformation  $L_A$  is the same as the rank of one of its matrix representations, namely,  $A$ . The next theorem extends this fact to any matrix representation of any linear transformation defined on finite-dimensional vector spaces.

**Theorem 2.** Let  $T : V \rightarrow W$  be a linear transformation between finite dimensional vector spaces, and let  $\beta$  and  $\gamma$  be ordered bases for  $V$  and  $W$ , respectively. Then  $\text{rank}(T) = \text{rank}([T]_{\beta}^{\gamma})$ .

**Theorem 3.** Let  $A$  be an  $m \times n$  matrix. If  $P$  and  $Q$  are invertible matrices of sizes  $m \times m$  and  $n \times n$ , respectively, then

1.  $\text{rank}(AQ) = \text{rank}(A)$
2.  $\text{rank}(PA) = \text{rank}(A)$
3.  $\text{rank}(PAQ) = \text{rank}(A)$

**Corollary 1.** Elementary row and column operations on a matrix are rank-preserving.

Now that we have a class of matrix operations that preserve rank, we need a way of examining a transformed matrix to ascertain its rank. The next theorem is the first of several in this direction.

**Theorem 4.** The rank of any matrix equals the maximum number of its linearly independent columns; that is the rank of a matrix is the dimension of the subspace generated by its columns.

### 2.0.1 Example

Find the rank of the following matrix:

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 1 & 0 \\ 2 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

### 2.0.2 Example

Prove that for any  $m \times n$  matrix  $A$ ,  $\text{rank}(A) = 0$  if and only if  $A$  is the zero matrix.

The next theorem uses this process to transform a matrix into a particularly simple form. The power of this theorem can be seen in its corollaries.

**Theorem 5.** Let  $A$  be an  $m \times n$  matrix of rank  $r$ . Then  $r \leq m$ ,  $r \leq n$ , and by means of a finite number of elementary row and column operations,  $A$  can be transformed into the matrix

$$D = \begin{bmatrix} I_r & O_1 \\ O_2 & O_3 \end{bmatrix}$$

Where  $O_1$ ,  $O_2$ , and  $O_3$  are zero matrices. Thus  $D_{ii} = 1$  for  $1 \leq i \leq r$  and  $D_{ij} = 0$  otherwise.

**Corollary 2.** Let  $A$  be an  $m \times n$  matrix of rank  $r$ . Then there exist invertible matrices  $B$  and  $C$  of sizes  $m \times m$  and  $n \times n$ , respectively, such that  $D = BAC$ , where

$$D = \begin{bmatrix} I_r & O_1 \\ O_2 & O_3 \end{bmatrix}$$

is the  $m \times n$  matrix in which  $O_1$ ,  $O_2$ , and  $O_3$  are zero matrices.

**Corollary 3.** Let  $A$  be an  $m \times n$  matrix. Then

1.  $\text{rank}(A^T) = \text{rank}(A)$ .
2. The rank of any matrix equals the maximum number of its linearly independent rows; that is, the rank of a matrix is the dimension of the subspace generated by its rows.
3. The rows and columns of any matrix generate subspace of the same dimensions, numerically equal to the rank of the matrix.

**Corollary 4.** Every invertible matrix is a product of elementary matrices.

*Proof.*

$d$

□

**Theorem 6.** Let  $T : V \rightarrow W$  and  $U : W \rightarrow Z$  be linear transformations on finite-dimensional vector spaces  $V$ ,  $W$ , and  $Z$  and let  $A$  and  $B$  be matrices such that the product  $AB$  is defined. Then

1.  $\text{rank}(UT) \leq \text{rank}(U)$
2.  $\text{rank}(UT) \leq \text{rank}(T)$
3.  $\text{rank}(AB) \leq \text{rank}(A)$
4.  $\text{rank}(AB) \leq \text{rank}(B)$

### 2.0.3 Example

Use elementary row and column operations to transform each of the following matrices into a matrix  $D$  satisfying the conditions of Theorem 5, and then determine the rank of each matrix:

$$A = \begin{bmatrix} 1 & 1 & 1 & 2 \\ 2 & 0 & -1 & 2 \\ 1 & 1 & 1 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 1 \\ -1 & 2 \\ 2 & 1 \end{bmatrix}$$

### 2.0.4 Example

For each of the following linear transformation  $T$ , determine whether  $T$  is invertible, and compute  $T^{-1}$  if it exists:

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be defined by

$$T(a_1, a_2, a_3) = (a_1 + 2a_2 + a_3, -a_1 + a_2 + 2a_3, a_1 + a_3)$$

## 2.1 The inverse of a Matrix

We have remarked that an  $n \times n$  matrix is invertible if and only if its rank is  $n$ . Since we know how to compute the rank of any matrix, we can always test a matrix to determine whether it is invertible. We now provide a simple technique for computing the inverse of a matrix that utilizes elementary row operations.

**Theorem 7.** Let  $A$  and  $B$  be  $m \times n$  and  $m \times p$  matrices, respectively. By the **augmented matrix**  $[A|B]$ , we mean the  $m \times (n + p)$  matrix, that is, the matrix whose first  $n$  columns are the columns of  $A$ , and whose last  $p$  columns are the columns of  $B$ .

Conversely, suppose that  $A$  is invertible and that, for some  $m \times p$  matrix  $B$ , the matrix  $[A|I_n]$  can be transformed into the matrix  $[I_n|B]$  by a finite number of elementary row operations. Let  $E_1, E_2, \dots, E_p$  be the elementary matrices associated with these elementary row operations as in Theorem 1. Then

### 2.1.1 Example

Express the invertible matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

As a product of elementary matrices.