

MATE 5150: Asignacion #3

Due on Septiembre 26, 2024

Dr. Pedro Vasquez

Alejandro Ouslan

Problem 1

Prove that T is a linear transformation, and find bases for both $N(T)$ and $R(T)$. Then compute the nullity and rank of T , and verify the dimension theorem. Finally use the appropriate theorems in this section to determine wheather T is one-to-one or onto.

$$T : R^2 \rightarrow R^3 \text{ defined by } T(a_1, a_2) = (a_1 + 2a_2, 0, 2a_1 - a_2)$$

Problem 2

Prove that T is a linear transformation, and find bases for both $N(T)$ and $R(T)$. Then compute the nullity and rank of T , and verify the dimension theorem. Finally use the appropriate theorems in this section to determine wheather T is one-to-one or onto.

$$T : M_{2 \times 3}(F) \rightarrow M_{2 \times 2}(F) \text{ defined by } T \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{pmatrix} = \begin{pmatrix} 2a_{11} - a_{12} & a_{13} + 2a_{12} \\ 0 & 0 \end{pmatrix}$$

Problem 3

In this exercise, $T : R^2 \rightarrow R^2$ is a function. State why T is not linear.

$$T(a_1, a_2) = (a_1 + 1, a_2)$$

Problem 4

Is there a linear transformation $T : R^3 \rightarrow R^2$ such that $T(1, 0, 3) = (1, 1)$ and $T(-2, 0, -6) = (2, 1)$?

Problem 5

Let V and W be vector spaces, let $T : V \rightarrow W$ be a linear, and let $\{w_1, w_2, \dots, w_k\}$ be a linearly independent set of k vectors from $R(T)$. Prove that if $S = \{v_1, v_2, \dots, v_k\}$ is chosen so that $T(v_i) = w_i$ for $i = 1, 2, \dots, k$, then S is linearly independent. Visit goo.gl/kmaQS2 for a solution.

Problem 6

Let β and γ be the standard ordered bases for R^n and R^m , respectively. For each linear transformation $T : R^n \rightarrow R^m$, compute $[T]_{\beta}^{\gamma}$.

1. $T : R^3 \rightarrow R$ defined by $T(a_1, a_2, a_3) = 2a_1 + a_2 - 3a_3$
2. $T : R^n \rightarrow R^n$ defined by $T(a_1, a_2, \dots, a_n) = (a_n, a_{n-1}, \dots, a_1)$

Problem 7