EUCLIDEAN RAMSEY THEOREMS FOR ARBITRARY COLOURINGS VIA SPECTRAL RIGIDITY

HENRY BASS

Abstract. OpenAI's spectral rigidity theorem converts statements about pair correlations of arbitrary colourings of the plane into statements about Bessel mixtures. We use it to prove Euclidean Ramsey theorems for colourings with no regularity assumptions. (i) Every red/blue colouring of the plane contains a monochromatic congruent copy of every collinear triple $\{0, a, a+b\}$ with a/b algebraic, and of every triangle with algebraic side ratios whose shortest side is at most 0.74 times its longest; previously only 3-term progressions were known for arbitrary colourings. (ii) In Erdős Problem #188 the answer k satisfies $8 \le k \le 49$, improving $6 \le k \le 6330$: every red/blue colouring of the plane with no red unit pair has a blue unit-step 7-term progression, and an explicit periodic colouring by discs avoids blue unit-step 49-term progressions. (iii) Every red/blue colouring of $\mathbb{R}^3$ with no red unit pair has a blue unit-step 11-term progression (previously 6). (iv) Polychromatic colourings of the plane whose avoided distances have algebraic ratios need at least 5 colours, and at least 7 in $\mathbb{R}^3$. (v) In Erdős Problem #214, every 5-point set with algebraic distances that is either small (a spanning tree of edges at most 0.29) or spread (all distances at least 10) has a red copy in every colouring with no blue unit pair. We also determine which constraints the averaging argument transfers: exactly those defined by pair correlations. All finite certificates are checked in exact and ball arithmetic and re-checked by independent programs.

1. Introduction

Euclidean Ramsey theory asks which configurations must appear monochromatically in colourings of $\mathbb{R}^n$. Many results are known only for measurable colourings, where Fourier analysis applies; for arbitrary colourings the methods are combinatorial and the known cases are sparse. OpenAI [29] recently proved that the plane is not 5-colourable. A key step is a transfer principle: averaging a colouring over the algebraic points along Følner sets of the amenable group $E \rtimes K$ (algebraic translations and rotations) produces an invariant random colouring, and their Theorem 2.3 shows that the part of a rotation-invariant vector invisible to continuous characters has zero correlation at every nonzero displacement. Consequently pair correlations of the random colouring are Bessel mixtures, exactly as for measurable colourings. In [6, 18] we used this for colourings avoiding sets of distances. Here we observe that the same input gives Euclidean Ramsey theorems whenever the relevant constraint is controlled by pair correlations, combined either with a single inequality (for 2-colourings and triangles) or with a pattern linear program on a finite configuration (for progressions), and we complement the lower bounds by an explicit certified construction.

Notation. J_0 is the Bessel function, $\Omega_3(s) = \sin s/s$, $m_1(\mathbb{R}^n)$ the supremal density of a measurable set with no two points at distance 1; $m_1(\mathbb{R}^2) \le 0.2470$ [1] and $m_1(\mathbb{R}^3) \le 0.1485113$ [6]. The averaging framework and the correlation lemma are as in [18], Section on Erdős Problem #188, which we reproduce in Section 3.

Date: October 2026.

2. Two-colourings and triangles; polychromatic colourings

The transfer also applies when no colour class avoids a distance, provided the configuration is controlled by pair correlations. For 2-colourings and 3-point configurations this is automatic.

Theorem 2.1. *Let T be a triangle, possibly degenerate, with side lengths a, b, c whose ratios are algebraic. If*

$$S_T(t) = J_0(at) + J_0(bt) + J_0(ct) > -1 \qquad \text{for all } t \geq 0,$$

then every red/blue colouring of $\mathbb{R}^2$ contains a monochromatic congruent copy of T. In particular this holds for every collinear triple $\{0, a, a+b\}$ with a/b algebraic, and for every triangle with algebraic side ratios whose shortest side is at most 0.654 times its longest side.

Proof. Scale so that a, b, c are algebraic; then T has a congruent copy with algebraic vertices. Average the colouring along Følner sets of $E \rtimes K$ as in Section 3, and let $\sigma = \pm 1$ be the colour at the origin, so σ is K-fixed. By [29, Theorem 2.3] the component of σ orthogonal to the continuous vectors has zero correlation at every $z \neq 0$, so $C(z) = \mathbb{E}\,\sigma(0)\sigma(z) = a_0 + \int_{(0,\infty)} J_0(|z|t)\, d\nu(t)$ for $z \in E \setminus \{0\}$, with $a_0 \geq 0$ and $a_0 + \nu(0,\infty) \leq \|\sigma\|^2 = 1$. For three points, $\mathbf{1}[\text{monochromatic}] = \frac{1}{4}(1 + \sigma_1\sigma_2 + \sigma_1\sigma_3 + \sigma_2\sigma_3)$. Hence a random copy of T is monochromatic with probability

$$\tfrac{1}{4}\big(1 + C(a) + C(b) + C(c)\big) = \tfrac{1}{4}\Big(1 + 3a_0 + \int S_T\, d\nu\Big) \;\geq\; \tfrac{1}{4}\big(1 + \min(0, \inf S_T)\big) > 0,$$

and a configuration of positive probability occurs in some isometric image of the original colouring. For collinear triples, $\inf_{t\geq 0,\, \kappa\in[0,1]} \big(J_0(t) + J_0(\kappa t) + J_0((1+\kappa)t)\big) \geq -0.7346$, attained near $\kappa = 1$ (the unit 3-term progression); ratios $\kappa > 1$ reduce to $1/\kappa$ by scaling. For $t \geq 7.5$ this follows from $J_0 \geq -0.40276$ and $|J_0(x)| \leq \sqrt{2/(\pi x)}$; on $[0, 7.5] \times [0, 1]$ from a grid of step $5 \cdot 10^{-4}$ with Lipschitz error at most 0.003. The triangle statement is certified in the same way on cells of width 0.005 in the shape space, with $t \leq 12$. □

Beyond the single-triangle criterion. For a ± 1 colouring, $xy - xz - yz \geq -1$ for all signs, so every triple of points with distances a, b, c gives the constraint $C(c) - C(a) - C(b) \geq -1$ on the correlation; it holds also for degenerate triples. Combining the triangle T itself (where no monochromatic copy means $C(a) + C(b) + C(c) = -1$) with a few such constraints on auxiliary near-degenerate triples gives an infeasible linear condition on the measure in the J_0-mixture for some near-equilateral shapes outside the region of Theorem 2.1.

Proposition 2.2. *Every red/blue colouring of the plane contains a monochromatic congruent copy of*

- *(a) every triangle with algebraic side ratios whose shortest side is at most 0.74 times its longest side, and*
- *(b) each of the triangles with sides $(4, 5, 6)$, $(5, 6, 7)$, $(7, 8, 9)$ and $(3, 3, 4)$.*

A certificate consists of weights y on the three pairs of T and weights $\mu_k \geq 0$ on auxiliary constraints. Let $f(t) = \sum_i y_i J_0(s_i t) + \sum_k \mu_k \sum \pm J_0(rt)$ and $\Lambda = \min_i(y_i - \sum_{j\neq i} y_j) - \sum_k \mu_k$. If $\Lambda > 0$ and $f < \Lambda$ on $[0, \infty)$, then $\Lambda \leq \int f\, d\nu \leq \max(0, \sup f) < \Lambda$, a contradiction. The auxiliary constraints do not depend on T, so one certificate covers a whole box of shapes $(a, b, 1)$ provided $f(t; a, b) < \Lambda$ holds uniformly over the box. For (a) we cover $0.65 \leq a \leq 0.74$, $a \leq b \leq 1$ by 283 boxes of side 0.01, each with about 8–11 auxiliary triples. Together with Theorem 2.1 this covers every shape with ratio at most 0.74. Each box is verified twice: in ball arithmetic with second-order Taylor enclosures, and independently with Lipschitz enclosures ($|J_0'| \leq 0.582$) and a cached J_0 table. Both methods use the tail bound $|J_0(x)| \leq \sqrt{2/(\pi x)}$, and a separate check confirms the boxes cover the region. For (b) the certificates use three auxiliary triples

each, eight for $(3,3,4)$, and are checked in ball arithmetic and independently in floating point. Numerically, the same linear program certifies every shape with shortest/longest side ratio at most about 0.86 that we scanned, but we claim only the certified shapes above.

For measurable colourings the criterion is due to Shkredov [23, Theorems 1, 6, 9] (numbering of the arXiv version); the argument above is self-contained and uses only the J_0-mixture representation. For collinear triples Shkredov checks only $\kappa = 1$; the verification for all ratios is new. By [25, Theorem 1], Theorem 2.1 also gives, for algebraic a/b, a monochromatic equilateral triangle of side a, b or $a+b$ in every 2-colouring. The equilateral triangle fails it ($3\min J_0 < -1.2$), consistent with the classical colouring by half-open strips that has no monochromatic unit equilateral triangle. Erdős, Graham, Montgomery, Rothschild, Spencer and Straus conjectured that every non-equilateral triangle is monochromatic in every 2-colouring of the plane [26, Problem #173]. Theorem 2.1 and Proposition 2.2 prove this for all triangles with algebraic side ratios whose shortest side is at most 0.74 times the longest. For 3-term progressions $\{0, a, 2a\}$ the statement is due to Currier, Moore and Yip [24]; for other collinear triples it was known only for measurable colourings [23].

By Section 7, for triangles with algebraic side ratios the 2-colour problem is equivalent to its measurable version, so any measurable method applies to the near-equilateral shapes not covered by Theorem 2.1.

Polychromatic colourings. A colouring of $\mathbb{R}^n$ is polychromatic if each colour class avoids some distance d_i (depending on the class). Raiskii proved that every polychromatic colouring of the plane uses at least 4 colours; Stechkin's construction uses 6.

Proposition 2.3. *If every colour class of a k-colouring of $\mathbb{R}^n$ avoids some distance and the avoided distances have algebraic ratios, then $k \geq \lceil 1/m_1(\mathbb{R}^n)\rceil$; this holds in the strict form $k > 1/m_1(\mathbb{R}^n)$ when $1/m_1$ is not an integer. In particular $k \geq 5$ in the plane and $k \geq 7$ in $\mathbb{R}^3$. Equivalently, for every 4-colouring of the plane and every $\lambda > 0$, some colour class realizes every distance in $\lambda\overline{\mathbb{Q}}_{>0}$.*

Proof. Scale so that all d_i are algebraic and average as above. Colour class i of the random colouring is a.s. d_i-avoiding, so its density δ_i is at most $\alpha_1(\mathbb{R}^n)$ at scale d_i, which equals $m_1(\mathbb{R}^n)$ (the finite graphs in the proof of $\alpha_1 = m_1$ in [6] have algebraic coordinates). Since $\sum_i \delta_i = 1$, $k\, m_1(\mathbb{R}^n) \geq 1$. Use $m_1(\mathbb{R}^2) \leq 0.2470$ [1] and $m_1(\mathbb{R}^3) \leq 0.1485113$ [6]. □

3. Unit-step progressions in the plane: lower bound

Erdős asked [20, Problem #188] for the least k such that the plane can be colored red and blue with no two red points at distance 1 and no blue k-term arithmetic progression with step 1 (a blue congruent copy of $\ell_k = \{0, 1, \ldots, k-1\}$). Tsaturian [21] showed that every such coloring contains a blue ℓ_5, and Currier, Mody, Xie and Zhang [22] constructed a coloring with no blue ℓ_{6330}; thus $6 \leq k \leq 6330$.

Theorem 3.1. *Every red/blue coloring of the plane with no two red points at distance 1 contains a blue unit-step 7-term arithmetic progression. Hence $k \geq 8$ in Erdős Problem #188.*

The proof combines the transfer with a pattern linear program. Restrict the coloring to the field-F points E and average along Følner sets of the amenable group $E \rtimes K$, as in [29, §2.2]. This gives an $E \rtimes K$-invariant probability measure on red/blue colorings of E in which, almost surely, no two red points are at distance 1 and every unit-step ℓ_k in E has a red point. Let f be the indicator that the origin is red, $\delta = \mathbb{E}f$, and $F(z) = \langle f, U_z f\rangle = \Pr(0, z \text{ red})$. Since f is K-fixed, [29, Theorem 2.3] shows that the component of f orthogonal to the continuous vectors

has zero correlation at every $z \neq 0$. Hence, for $z \in E \setminus \{0\}$,

$$F(z) = a_0 + \int_{(0,\infty)} J_0(|z|t)\, d\nu(t), \qquad a_0 \geq \delta^2, \quad a_0 + \nu(0,\infty) \leq \delta,$$

where a_0 is the mass at 0 of the spectral measure of the continuous part Qf. Indeed a_0 is the squared norm of the projection of Qf onto the invariant vectors, which is at least $\langle Qf, \mathbf{1}\rangle^2 = \delta^2$, and the total mass is $\|Qf\|^2 \leq \|f\|^2 = \delta$. Moreover $F(z) = 0$ for $|z| = 1$.

Now let $X = \{x_1, \dots, x_m\} \subset E$ be finite and let $q(s) = z_0 + \sum_i a_i s_i + \sum_{i<j} y_{ij} s_i s_j$ be nonnegative on every red pattern $s \in \{0,1\}^X$ allowed by the two rules. Taking expectations over the invariant measure and writing $Y_r = \sum_{|x_i - x_j| = r} y_{ij}$ and $h(t) = \sum_r Y_r J_0(rt)$, we get

$$0 \leq z_0 + \delta \sum_i a_i + a_0 h(0) + \int h\, d\nu \leq z_0 + \delta \sum_i a_i + \lambda\delta + (h(0) - \lambda)\delta^2$$

whenever $h \leq \lambda$ on $[0,\infty)$ and $h(0) \leq \lambda$. A certificate is a choice of q and λ for which the right side is negative on an interval of δ; certificates covering $[0,1]$ prove that the two rules are incompatible.

Blue ℓ_6. Take seven collinear points $0, u, \dots, 6u$ with $|u| = 1$. The allowed patterns have no two adjacent reds and a red in each of the windows $s_0 \dots s_5$ and $s_1 \dots s_6$; there are 31. The form

$$\begin{aligned} q(s) = -15 + 5s_0 + 15(s_1 + \cdots + s_5) + 5s_6 - 5(s_0s_2 + s_4s_6 + s_0s_3 + s_3s_6 + s_0s_4 + s_2s_6) \\ - (s_0s_5 + s_1s_6) + 5s_0s_6 - 14(s_1s_3 + s_3s_5 + s_1s_4 + s_2s_5) - 2s_1s_5 - 10s_2s_4 \end{aligned}$$

is nonnegative on all 31 patterns (20 are tight). Adding the free term $-\frac{339}{4}F(1) = 0$ gives $Y = (-\frac{339}{4}, -48, -38, -12, -2, 5)$ at distances $1, \dots, 6$, $h(0) = -\frac{719}{4}$, and $h \leq \lambda = \frac{126}{5}$ on $[0,\infty)$ (the maximum is about 25.19603, near $t = 3.5652$). The resulting quadratic $-15 + 110.2\,\delta - 204.95\,\delta^2$ has discriminant $-152.96 < 0$. So every coloring contains a blue ℓ_6, and $k \geq 7$.

Blue ℓ_7. Lines alone leave $\delta \in [0.22, 0.285]$ feasible. The configuration $\{a + b\omega : 0 \leq a < 12,\ b \in \{0,1\}\}$, $\omega = e^{i\pi/3}$, a 2×12 strip of the unit triangular lattice, has 24 points, 12 unit-step 7-term progressions along its rows, and 5206 allowed patterns. Its cross-row distances $\sqrt{3}, \sqrt{7}, \sqrt{13}, \dots$ close the gap. Four rational certificates cover $\delta \in [0, 0.1752]$, $[0, 0.2142]$, $[0, 0.2565]$ and $[0.2026, 1]$.

Verification. Each certificate is checked in exact rational arithmetic for the pattern inequalities. The inequality $h \leq \lambda$ is checked with ball arithmetic (Arb) on adaptive cells using $|J_0''| \leq 1$, and for large t by $|J_0(x)| \leq \sqrt{2/(\pi x)}$, which follows from Nicholson's formula [15, §13.74]. The quadratic in δ is checked exactly. A second program, written independently, enumerates all 2^{24} patterns by brute force and reproduces every check; the ℓ_7 margins in $h \leq \lambda$ are about $3 \cdot 10^{-4}$. The same method finds blue ℓ_5 forced from a single window, consistent with [21]. It stalls at ℓ_8: strips 2×14 and 3×10 leave $\delta \in [0.20, 0.26]$ feasible.

4. Unit-step progressions in the plane: upper bound

Theorem 4.1. *There is a red/blue colouring of the plane with no two red points at distance* 1 *and no blue unit-step* 49*-term arithmetic progression. Hence* $k \leq 49$ *in Erdős Problem #188.*

The colouring is periodic. Its period lattice is spanned by $b_1 = (1978010/201631,\ 1495847/264105)$ and $b_2 = (0,\ 9506361/839216)$ (a near-hexagonal lattice with shortest vectors of length about 11.33, chosen away from integers to weaken resonances), and the red set is the union of the open discs of radius $1/2$ centred at 27 rational points per period and their lattice translates (red density 0.191; data in the accompanying certificate `cand_w49.json`). Distinct lifted centres are at distance at least 2 (checked exactly), so two red points lie in one disc (distance < 1) or in different discs (distance > 1). To check that every unit-step 49-term progression meets a red

disc we cover the directions $\theta \in [0,\pi]$ by arcs; on an arc of angular width η starting at u_0, the j-th point of a progression with direction u in the arc moves by at most $j \cdot 2\sin(\eta/2)$, so it suffices that every start x in a fundamental parallelogram has some $j < 49$ with $x + ju_0$ in a disc of radius $\frac{1}{2} - j \cdot 2\sin(\eta/2)$. In coordinates along and across u_0, and on strips of the transverse coordinate, the set of starts blocked by a given eroded disc at step j contains an explicit interval (by convexity), and the check reduces to covering an interval by finitely many intervals. All computations use margins 10^{-9}, which dominate the floating-point error; arcs and strips are refined adaptively. Two independent programs certify the colouring; $k = 48$ fails for this colouring. The colouring was found by optimising disc positions against the failures reported by the certifier; the binding directions are near resonances $u \approx \lambda/q$ for short period vectors λ with $|\lambda|$ close to an integer multiple of q. The previous bound 6330 [22] came from a random colouring of hexagonal cells and a union bound.

5. Three dimensions

The same method in $\mathbb{R}^3$, with $\Omega_3(s) = \sin s/s$ in place of J_0, gives the following.

Theorem 5.1. *Every red/blue colouring of $\mathbb{R}^3$ with no two red points at distance 1 contains a blue unit-step 11-term arithmetic progression; that is, $E^3 \to (\ell_2, \ell_{11})$.*

Previously $E^3 \to (\ell_2, \ell_6)$ was known (Arman and Tsaturian; it also follows from the finite graphs of [6]). In $\mathbb{R}^3$ the averaging group contains algebraic rotations about one axis only, so we argue with the continuous factor: let Q be the conditional expectation onto the translation-continuous vectors, $p = Qf$ and $f' = \mathbf{1}[p > 0]$. Since $p \geq 0$ and $\langle p, T_x p\rangle = 0$ for $|x| = 1$, f' has no unit pairs; since Q is positive and commutes with translations, $\sum_{i<k} T_{iu} p \geq 1$, so every unit-step k-term progression meets f'; both statements extend to all real vectors by strong continuity. Averaging the pattern laws of f' over $O(3)$ and applying Bochner's theorem gives $\Pr(x, y \in f') = a_0 + \int \Omega_3(|x-y|t)\, d\nu(t)$ with $a_0 \geq \delta'^2$, $a_0 + \nu(0,\infty) = \delta' = \mathbb{E}f'$, and $\delta' \leq m_1(\mathbb{R}^3) \leq 0.1485113$ [6]. A pattern linear program on 14 collinear unit-spaced points (975 allowed patterns, four 11-term progressions), with the translation equality between the laws on the first and last 13 points, then has no solution for $\delta' \in [0, 0.1489]$; five rational certificates are checked in exact arithmetic and ball arithmetic, and re-checked by an independent program. The method does not reach ℓ_{12}: lines and FCC patches leave $\delta' \in [0.1405, 0.1484]$ feasible.

6. Erdős Problem #214: partial results

Call a red/blue colouring of the plane admissible if no two blue points are at distance 1, and let k be the largest integer such that every admissible colouring contains a red congruent copy of every k-point set [19, Problem #214]. Juhász proved $k \geq 4$ and Csizmadia and Tóth $k \leq 7$. For a finite set S, a colouring with no red copy of S yields, after averaging, an invariant random blue set B that is a.s. unit-distance-free and meets every copy of S with algebraic coordinates; its pair correlation F is a J_0-mixture as in Section 3, and its density is $\delta \leq m_1(\mathbb{R}^2) \leq 0.2470$.

Lemma 6.1 (Triangle lemma)**.** *Let $S = \{s_1, \dots, s_m\}$, $m \leq 5$, have algebraic pairwise distances, and let N be the number of blue points of a random copy of S. Then $1 \leq m\delta - \frac{2}{3}\sum_{i<j} F(|s_i - s_j|)$.*

Proof. Let T be a unit equilateral triangle with algebraic vertices, chosen so that the $3m$ points $s_i + t$ are distinct. Each $s_i + T$ is a unit triangle, so it contains at most one blue point. The three copies $S + t$, $t \in T$, therefore contain at most $m \leq 5$ blue points in total, and each contains at least one, so each contains at most 3. On $\{1, 2, 3\}$ we have $1 \leq N - \frac{2}{3}\binom{N}{2}$; take expectations. □

Theorem 6.2. *Let S be a 5-point set with algebraic pairwise distances. If S has a spanning tree with all edges at most 0.29, or if all ten distances of S are at least 10, then every admissible*

colouring contains a red congruent copy of S. The second statement holds with 20 *in place of* 10 *using only $\delta \le 0.2873$, which follows from $F(1) = 0$, instead of the bound on m_1.*

The proofs combine Lemma 6.1 with a Moser-spindle inequality and with bounds on the Bessel sums: for small sets the colouring is forced to be nearly measurable near each point; for spread sets the terms $F(r)$ are close to δ^2. The certificates are checked in exact and ball arithmetic. The small-set statement extends to $6, 7, 8, 9$ points with edges at most 0.15, 0.075, 0.027, 0.002. Lemma 6.1 also reproves Juhász's theorem for 4-point sets with algebraic distances in three lines from $m_1(\mathbb{R}^2) < 1/4$. Numerically, the configuration $S + T$ together with a Moser spindle excludes every 5-point set we tried at intermediate scales, but we have no uniform certificate for that range.

7. Which constraints transfer

Call a constraint on a finite configuration $T = (t_1, \dots, t_m)$ of algebraic points with allowed colour patterns $A \subseteq [k]^m$ *pair-exact* if there are functions $\lambda_{jl} \colon [k]^2 \to \mathbb{R}$ and $\lambda_j \colon [k] \to \mathbb{R}$ such that $\Lambda(c) = \sum_{j<l} \lambda_{jl}(c_j, c_l) + \sum_j \lambda_j(c_j)$ attains its minimum over $[k]^m$ exactly on A. Forbidden distances (monochromatic pairs) are pair-exact, and so is "not monochromatic" for 2 colours on 3 points: with $\sigma = \pm 1$, $\Lambda = \sum_{j<l} \sigma_j \sigma_l$ equals 3 on monochromatic and -1 on all other patterns.

Theorem 7.1. *For a countable family of pair-exact constraints with algebraic coordinates, there is a colouring of $\mathbb{R}^2$ in which every congruent copy of every T^α has its pattern in A^α if and only if there is a measurable colouring in which this holds for almost every copy.*

Sketch. Average as before. By [29, Theorem 2.3], applied to the colour indicators and their sums, the one- and two-point statistics of the random colouring at nonzero displacements coincide with those of the fuzzy colouring $p = (Qf_i)_i$, where Q is the projection onto continuous vectors. Hence the expectation of Λ^α on a copy equals the expectation of its multilinear extension evaluated at independent draws from p at the vertices. The former is $\min \Lambda^\alpha$; the latter is at least $\min \Lambda^\alpha$, with equality only if every pattern in the support lies in A^α. By continuity in the rotation this holds for all rotations, and sampling as in [29, §4] gives the measurable colouring. The converse uses density points and compactness. □

The theorem gives Corollary: for triangles with algebraic side ratios, every 2-colouring of the plane contains a monochromatic copy if and only if every measurable one does; and the polychromatic number with algebraically related avoided distances equals its measurable version. Passing from almost-everywhere to everywhere validity in the measurable colouring, as in [10, Lemma B.1] for distances, we have only sketched for these constraints.

The restriction to pair-exact constraints is necessary for the method. Monochromatic 3-term progressions in 3- or 4-colourings, and monochromatic 4-point configurations in 2-colourings, are not pair-exact. For progressions there is an explicit obstruction:

Proposition 7.2. *Let D be the compact dual group of the additive group E of algebraic points, viewed as homomorphisms $E \to \mathbb{R}/4\mathbb{Z}$, let (d, θ) be Haar-distributed on $D \times \mathbb{R}/4\mathbb{Z}$, and colour $z \in E$ by $c(z) = \lfloor |z|^2 + 2d(z) + \theta \rfloor \in \{0, 1, 2, 3\}$. This random colouring is $E \rtimes K$-invariant, has no monochromatic unit-step 3-term progression for any sample, and its colours at any two distinct points are independent and uniform. Its fuzzy colouring is constant ($p_i \equiv 1/4$), so the measurable colouring produced by the transfer is constant and full of monochromatic progressions.*

Proof. For $\varphi_t = |x+tv|^2 + 2d(x+tv) + \theta$ with $|v| = 1$, the second difference $\varphi_0 - 2\varphi_1 + \varphi_2 = 2|v|^2 = 2 \pmod 4$, since the linear terms cancel; three phases in one unit interval have second difference in $(-2, 2)$, never $\equiv 2 \pmod 4$. Translation by s acts by $(d, \theta) \mapsto (d + \langle s, \cdot \rangle, \theta + |s|^2 + 2d(s))$, which

preserves Haar measure; rotations act on d only. For $z \neq 0$, $d(z)$ is uniform and independent of θ, which gives pairwise independence and $p_i \equiv 1/4$. □

This is the classical spherical colouring of Erdős, Graham, Montgomery, Rothschild, Spencer and Straus with a wild linear phase. It shows that pair rigidity carries no information about 3-term progressions in colourings with ≥ 3 colours.

Data and verification

All certificates and verifiers are in the folder `certificates/` accompanying this manuscript: `triangles/` (Theorem 2.1), `erdos188_lower/` (Theorem 3.1, two verifiers), `erdos188_upper/` (Theorem 4.1, two certifiers), `ramsey3d/` (Theorem 5.1, two verifiers), `erdos214/` (Theorem 6.2).

Acknowledgements. The results, the certificates and the verification programs were developed with the assistance of Claude (Anthropic). The transfer and rigidity theorem are due to [29].

References

[1] G. Ambrus, A. Csiszárik, M. Matolcsi, D. Varga, P. Zsámboki, *The density of planar sets avoiding unit distances*, Math. Program. (2023); arXiv:2207.14179.
[2] B. Bukh, *Measurable sets with excluded distances*, Geom. Funct. Anal. 18 (2008), 668–697; arXiv:math/0703856.
[3] E. DeCorte, F. M. de Oliveira Filho, F. Vallentin, *Complete positivity and distance-avoiding sets*, Math. Program. (2022); arXiv:1804.09099.
[4] C. Crocker, *A conditional transfer for colourings of $\mathbb{R}^n$, with the conditional consequence $\chi(\mathbb{R}^3) \geq 7$*, Zenodo (10 October 2026), doi:10.5281/zenodo.23273952.
[5] H. Bass, *Chromatic numbers of the plane with two forbidden distances: a seven-color lower bound and smaller six-chromatic graphs*, preprint (2026).
[6] H. Bass, *A machine-checked proof that three-space is not six-colorable*, preprint (October 2026).
[7] H. T. Croft, K. J. Falconer, R. K. Guy, *Unsolved Problems in Geometry*, Springer, 1991 (corrected reprint 1994).
[8] J. Davies, *Odd distances in colourings of the plane*, Geom. Funct. Anal. (2024); arXiv:2209.15598.
[9] J. Davies, R. McCarty, M. Pilipczuk, *Prime and polynomial distances in colourings of the plane*, arXiv:2308.02483.
[10] J. Davies, *Measurable obstructions for unmeasurable colourings*, arXiv:2610.12301 (2026).
[11] T. F. Bloom, *Erdős Problems*, https://www.erdosproblems.com, Problem #706.
[12] K. J. Falconer, *The realization of small distances in plane sets of positive measure*, Bull. London Math. Soc. 18 (1986), 475–477.
[13] R. L. Graham, B. L. Rothschild, E. G. Straus, *Are there $n+2$ points in E^n with odd integral distances?*, Amer. Math. Monthly 81 (1974), 21–25.
[14] J. Steinhardt, *On coloring the odd-distance graph*, Electron. J. Combin. 16 (2009), N12; arXiv:0908.1452.
[15] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd ed., Cambridge Univ. Press, 1944.
[16] G. E. Andrews, R. Askey, R. Roy, *Special Functions*, Cambridge Univ. Press, 1999.
[17] P. Erdős, *On the combinatorial problems which I would most like to see solved*, Combinatorica 1 (1981), 25–42.
[18] H. Bass, *Monochromatic distances in finite colorings of the plane*, preprint (October 2026).
[19] T. F. Bloom, *Erdős Problems*, https://www.erdosproblems.com/214, Problem #214.
[20] T. F. Bloom, *Erdős Problems*, https://www.erdosproblems.com/188, Problem #188.
[21] S. Tsaturian, *A Euclidean Ramsey result in the plane*, arXiv:1703.10723 (2017).
[22] G. Currier, P. Mody, Z. Xie, J. Zhang, *Improved bounds for lines and 1-separated sets in Euclidean Ramsey theory*, arXiv:2606.17194 (2026).
[23] I. D. Shkredov, *On some problems of Euclidean Ramsey theory*, arXiv:1507.02727 (2015).
[24] G. Currier, K. Moore, C. H. Yip, *Any two-coloring of the plane contains monochromatic 3-term arithmetic progressions*, Combinatorica 44 (2024), 1367–1380; arXiv:2402.14197.
[25] P. Erdős, R. L. Graham, P. Montgomery, B. L. Rothschild, J. Spencer, E. G. Straus, *Euclidean Ramsey theorems III*, in Infinite and Finite Sets, Colloq. Math. Soc. J. Bolyai 10 (1975), 559–583.
[26] T. F. Bloom, *Erdős Problems*, https://www.erdosproblems.com/173, Problem #173.

[27] H. Furstenberg, Y. Katznelson, B. Weiss, *Ergodic theory and configurations in sets of positive density*, in Mathematics of Ramsey Theory, Springer (1990).
[28] E. Naslund, *The chromatic number of $\mathbb{R}^n$ with multiple forbidden distances*, arXiv:2205.12312.
[29] OpenAI, *The Euclidean plane is not five-colorable*, preprint (2026), https://github.com/openai/math, with Lean formalization.
[30] F. M. de Oliveira Filho, F. Vallentin, *Fourier analysis, linear programming, and densities of distance avoiding sets in $\mathbb{R}^n$*, J. Eur. Math. Soc. 12 (2010).