A MACHINE-CHECKED PROOF THAT THREE-SPACE IS NOT SIX-COLORABLE

HENRY BASS

Abstract. We give a proof, formally verified in Lean 4, that every coloring of $\mathbb{R}^3$ with six colors, with arbitrary color classes, contains two points at distance 1 of the same color. Thus $\chi(\mathbb{R}^3) \geq 7$; the previous unconditional bound was $\chi(\mathbb{R}^3) \geq 6$ (Nechushtan, 2002).

The proof extends to $\mathbb{R}^3$ the transfer principle of a recent preprint released by OpenAI, which converts a proper k-coloring of the plane into a measurable k-coloring whose monochromatic unit pairs form a null set, and whose key rigidity theorem is formalized in Lean. We average over independent rotations of orthogonal coordinate 2-planes, apply the rigidity theorem block by block through a polarization identity, and remove the remaining axis with an invariant-mean line average. The argument works in every dimension and for every finite set of algebraic forbidden distances, and gives $\chi_D(\mathbb{R}^n) \geq 1/m_D(\mathbb{R}^n)$. With the density bound of DeCorte, Oliveira and Vallentin this yields $\chi(\mathbb{R}^3) \geq 7$. Crocker independently posted, on 10 October 2026, the same transfer and its consequences, stated conditionally on the rigidity theorem and with a different treatment of the axis. Our formalization verifies, end to end and with only the standard axioms, the transfer in every dimension for every countable set of algebraic distances, the deduction of $\chi(\mathbb{R}^3) \geq 7$ from the published density bound, the existence of a finite non-6-colorable unit-distance graph in $\mathbb{R}^3$, and the existence of finite unit-distance graphs in $\mathbb{R}^3$ with independence ratio below 0.1533. The only input not formalized for $\chi(\mathbb{R}^3) \geq 7$ is that numerical density bound, whose published exact verifier we re-ran.

We also show that finite unit-distance graphs in $\mathbb{R}^3$ attain the measurable density, $\inf_G \alpha(G)/|G| = m_1(\mathbb{R}^3)$, with one inequality formalized, and that $\sup_G \chi_f(G) = \sup_G \chi_{gf}(G)$ for finite unit-distance graphs in every dimension. Consequences of the transfer for colorings avoiding several distances, including formal answers to questions of Soifer, Kahle and Erdős, are in a companion paper.

1. Introduction

A proper k-coloring of $\mathbb{R}^n$ is a map $c\colon \mathbb{R}^n \to \{1, \dots, k\}$ with $c(x) \neq c(y)$ whenever $|x-y| = 1$. The chromatic number $\chi(\mathbb{R}^n)$ is the least such k, with no regularity imposed on the color classes. For the plane, Nelson and Isbell showed $4 \leq \chi(\mathbb{R}^2) \leq 7$ around 1950. De Grey raised the lower bound to 5 in 2018 [8]. A preprint released by OpenAI in September 2026 [13] proves $\chi(\mathbb{R}^2) \geq 6$, with a Lean formalization that we have rebuilt and checked. In three dimensions the best previous bounds were $6 \leq \chi(\mathbb{R}^3) \leq 15$ [12, 3, 15].

The measurable chromatic number $\chi_m(\mathbb{R}^n)$ requires measurable color classes. It is bounded below by $1/m_1(\mathbb{R}^n)$. Here

$$m_1(\mathbb{R}^n) = \sup\{\overline{\delta}(A) : A \subseteq \mathbb{R}^n \text{ measurable, no two points of } A \text{ at distance } 1\},$$

and $\overline{\delta}$ denotes upper density with respect to centered cubes. Fourier-analytic and semidefinite methods [14, 2, 4] give much stronger lower bounds for χ_m than combinatorial methods give for χ. In particular $\chi_m(\mathbb{R}^3) \geq 7$ [14, 4].

Theorem 1.1. $\chi(\mathbb{R}^3) \geq 7$.

Date: October 2026.

The OpenAI argument has two parts. The first is a transfer theorem [13, Theorem 1.3]: the plane has a proper k-coloring if and only if it has a *weak measurable* k-coloring. The second shows that weak measurable 5-colorings do not exist. The transfer averages a coloring of the algebraic points over algebraic translations and rotations. This needs the group $\mathbb{R}^2_{\mathrm{alg}} \rtimes SO(2)_{\mathrm{alg}}$ to be amenable, which holds because it is solvable. Matolcsi, Ruzsa, Varga and Zsámboki [11] use the same kind of averaging for fractional colorings, and point out that it "cannot automatically be extended to higher dimensions."

Definition 1.2. A *weak measurable k-coloring* of $\mathbb{R}^n$ is a Lebesgue measurable $c\colon \mathbb{R}^n \to \{1,\dots,k\}$ with classes $A_i = c^{-1}(i)$ such that for every $R > 0$

$$\sum_{i=1}^{k} \int_{B(0,R)} \int_{S^{n-1}} \mathbf{1}_{A_i}(x)\, \mathbf{1}_{A_i}(x+u)\, d\sigma(u)\, dx = 0,$$

where σ is normalized surface measure on the unit sphere.

Theorem 1.3 (Transfer in every dimension)**.** *Let $n \geq 2$ and $k \geq 1$. If $\mathbb{R}^n$ has a proper k-coloring, then it has a weak measurable k-coloring.*

The converse holds by the density-point argument of [13, Lemma 4.3 and proof of Theorem 1.3], which is dimension-free. Thus $\chi(\mathbb{R}^n)$ equals the weak measurable chromatic number for every n.

Corollary 1.4. *If $\mathbb{R}^n$ has a proper k-coloring, then there are pairwise disjoint Borel sets $A_1^*, \dots, A_k^* \subseteq \mathbb{R}^n$, whose union has null complement, such that no A_i^* contains two points at distance* 1*. Consequently*

$$\chi(\mathbb{R}^n) \;\geq\; \lceil 1/m_1(\mathbb{R}^n) \rceil.$$

Theorem 1.1 follows from Corollary 1.4 and the bound $m_1(\mathbb{R}^3) \leq 0.1532996 < 1/6$ of DeCorte, Oliveira and Vallentin [4, Table 2], which was published with an exact verification program that we rebuilt and re-ran (Section 5).

Relation to the work of Crocker. After this work was done, we learned of a note by Crocker [5], posted on 10 October 2026, which proves Theorem 1.3 for all n and all sets of algebraic distances, the bound $\chi(\mathbb{R}^3) \geq 7$, and the consequences in Theorems 1.6 and 7.1, each stated conditionally on [13, Theorem 2.3]. He uses the same averaging group. He handles the axis by a Bohr-averaging lemma, where we use an invariant-mean line average. Since [13, Theorem 2.3] is formalized in Lean, our formalization of the three-dimensional case (Theorem 1.5) makes $\chi(\mathbb{R}^3) \geq 7$ unconditional, relative to the published numerical density bound and the Lean kernel.

Relation to the work of Davies. Shortly before this paper, Davies [7] proved $\chi_D(\mathbb{R}^n) = \chi_D^{(m)}(\mathbb{R}^n)$ and $\alpha_D(\mathbb{R}^n) = m_D(\mathbb{R}^n)$ for $n \geq 4$ and finite distance sets D with rational squares, together with $\chi(\mathbb{R}^n) \geq (2 + o(1))^n$ and strong bounds for $5 \leq n \leq 24$. His method embeds lattice Cayley graphs that approximate the copositive certificates of [4]. It does not apply to $n = 3$, which he leaves open, writing that he hopes his techniques extend to dimensions 2 and 3. For $n \geq 4$, Corollary 1.4 recovers his chromatic inequality $\chi \geq 1/m_1$ by a different route. It also holds for every countable set of algebraic distances (Section 7).

Theorem 1.5 (Formal verification)**.** *The following are formally verified in Lean 4 with Mathlib: Theorem 1.3 in the form of Corollary 1.4 for every $n \geq 2$ and every countable set of positive algebraic forbidden distances (Theorem 7.1); the deductions $\chi(\mathbb{R}^n) \geq \lceil 1/\delta \rceil$ from a density bound $m_1(\mathbb{R}^n) \leq \delta$, in particular $\chi(\mathbb{R}^3) \geq 7$ from $m_1(\mathbb{R}^3) \leq 0.1532996$, and the bounds for $4 \leq n \leq 8$ from* [4, Table 2]*; their finite forms (finite point sets with no proper coloring); and, in $\mathbb{R}^3$, the inequality $\alpha_1(\mathbb{R}^3) \leq m_1(\mathbb{R}^3)$ of Theorem 1.6. The proof depends only on the standard axioms* `propext`*,* `Classical.choice` *and* `Quot.sound`*.*

Section 8 describes the formal development. It builds on the Lean library accompanying [13], in particular on its formal proof of the planar rigidity theorem. The inputs not formalized are the numerical density bounds of [4], which appear as explicit hypotheses, and the inequality $m_1 \le \alpha_1$.

Finally, the same method applied to random independent sets gives a statement about finite graphs. For a finite unit-distance graph $G \subset \mathbb{R}^n$ let $\alpha_1(G) = \alpha(G)/|G|$, and set $\alpha_1(\mathbb{R}^n) = \inf_G \alpha_1(G)$. Averaging translates gives $m_1(\mathbb{R}^n) \le \alpha_1(\mathbb{R}^n)$. In the plane, Ambrus et al. [1] and Matolcsi et al. [11] expected the inequality to be strict, "an interesting example of a natural infinite graph for which the measurable and the non-measurable variants of a parameter have different values." Dúcz and Varga [9] disproved the planar conjecture $\alpha_1(\mathbb{R}^2) = 1/4$, and conjectured instead that $f(N)/N = m_1(\mathbb{R}^2) + o(1)$, where $f(N)$ is the least independence number of an N-point unit-distance graph in the plane. Li and Su [10] recently proved this, with a Lean formalization, by applying the transfer of [13] to a single color class. Davies [7] proved $\alpha_1(\mathbb{R}^n) = m_1(\mathbb{R}^n)$ for $n \ge 4$. The torus averaging of Theorem 1.3 treats all dimensions uniformly, as does the conditional argument of Crocker [5].

Theorem 1.6. *For every $n \ge 2$,*

$$\alpha_1(\mathbb{R}^n) = m_1(\mathbb{R}^n) \quad \text{and} \quad \sup_G \chi_f(G) = \sup_G \chi_{gf}(G) = 1/m_1(\mathbb{R}^n),$$

where G ranges over finite unit-distance graphs in $\mathbb{R}^n$ and χ_{gf} is the geometric fractional chromatic number of [1, 11].

The first equality is due to Li and Su for $n = 2$ and to Davies for $n \ge 4$; Crocker [5] states it for all n, conditionally on [13, Theorem 2.3]. The second answers the question of [11] whether $\chi_{gf,0}(\mathbb{R}^d) = \chi_{f,0}(\mathbb{R}^d)$ for some $d \ge 3$. In $\mathbb{R}^3$ it gives finite unit-distance graphs with independence ratio below 0.1533. As in [9, 10], the graphs produced are not explicit.

Acknowledgements. The extension to higher dimensions, Theorems 1.6 and 7.1, and the Lean formalization were developed with the assistance of Claude (Anthropic). The underlying planar results and their formalization are due to [13].

2. Setting and inputs

Let $F = \overline{\mathbb{Q}} \cap \mathbb{R}$ be the field of real algebraic numbers, $E = F(i) \subset \mathbb{C}$, and $K = \{u \in E : |u| = 1\}$, a countable divisible abelian group. Write $n = 2m$ or $n = 2m + 1$ and identify $\mathbb{R}^n$ with $\mathbb{C}^m$ or $\mathbb{C}^m \times \mathbb{R}$. The algebraic points form the countable dense subgroup

$$V = E^m \quad \text{or} \quad V = E^m \times F.$$

The group K^m acts on V by rotating the j-th coordinate plane by the j-th component and fixing the last axis. These maps are linear isometries of $\mathbb{R}^n$. The group $G = V \rtimes K^m$ is solvable, hence amenable.

For a unitary representation U of a discrete group A that is a dense subgroup of a Euclidean space (here E, F or V), let

$$\mathcal{H}_c(U) = \{v : \ \|U_z v - v\| \to 0 \text{ as } z \to 0 \text{ in the Euclidean topology}\}.$$

This is a closed subspace. We use three results of [13].

Theorem 2.1 ([13, Theorem 2.3], Hilbert-space form)**.** *Let $U \colon E \to \mathcal{U}(\mathcal{H})$ and $W \colon K \to \mathcal{U}(\mathcal{H})$ be unitary representations with $W_k U_z = U_{kz} W_k$. If $v \perp \mathcal{H}_c(U)$ and $W_k v = v$ for all $k \in K$, then $\langle v, U_z v \rangle = 0$ for every $z \in E \setminus \{0\}$.*

In the Lean library of [13] this is `PlaneFiveColor.Spectral.residual_fourier`. It is deduced there from the rigidity of K-invariant probability measures on the dual of E that give no mass to the continuous characters.

Lemma 2.2 ([13, Lemma 4.1])**.** *If U is the Koopman representation of a measure-preserving action of A on a probability space, then $\mathcal{H}_c(U)$ is the L^2 space of a factor. Hence its orthogonal projection is a conditional expectation: it preserves $0 \le f \le 1$ and fixes constants. Translations act continuously on $\mathcal{H}_c(U)$ and extend to a strongly continuous, positivity-preserving unitary representation of the ambient Euclidean space.*

The proof in [13] is stated for $A = E$ but uses only that $\mathcal{H}_c$ is closed, contains constants, and is stable under products of bounded elements and under Lipschitz maps. It applies verbatim to $A = F$ and $A = V$.

Lemma 2.3 ([13, §2.2])**.** *There is a positive, F-translation-invariant linear functional I on bounded, boundedly supported functions $F \to \mathbb{C}$ with $I(\varphi|_F) = \int_{\mathbb{R}} \varphi$ for all $\varphi \in C_c(\mathbb{R})$. It integrates bounded Hilbert-valued functions weakly, with $\|I(\Phi)\| \le I(\|\Phi\|)$.*

3. Two lemmas

Lemma 3.1. *Let U be a unitary representation as above and Y a bounded operator commuting with every U_z. Then $Y\mathcal{H}_c(U) \subseteq \mathcal{H}_c(U)$. If moreover Y is unitary, or self-adjoint, then $Y\mathcal{H}_c(U)^\perp \subseteq \mathcal{H}_c(U)^\perp$, and Y commutes with the orthogonal projection onto $\mathcal{H}_c(U)$.*

Proof. $\|U_z Yv - Yv\| = \|Y(U_z v - v)\| \le \|Y\|\,\|U_z v - v\|$. In the unitary or self-adjoint case Y^* also commutes with U, so $\langle Yw, h\rangle = \langle w, Y^* h\rangle = 0$ for $w \perp \mathcal{H}_c$ and $h \in \mathcal{H}_c$. Splitting $v = Pv + (v - Pv)$ gives $PYv = YPv$. □

Lemma 3.2 (Twisted rigidity)**.** *Let U, W be as in Theorem 2.1, and let Y be unitary and commute with every U_z and every W_k. If $v \perp \mathcal{H}_c(U)$ is W-fixed, then $\langle v, U_z Y v\rangle = 0$ for every $z \in E \setminus \{0\}$.*

Proof. By Lemma 3.1, Yv is W-fixed and orthogonal to $\mathcal{H}_c(U)$, and so is $v + sYv$ for every $s \in \mathbb{C}$. Theorem 2.1 applied to v, Yv and $v + sYv$ gives

$$0 = s\,\langle v, U_z Y v\rangle + \bar{s}\,\langle Yv, U_z v\rangle \qquad (s \in \mathbb{C}).$$

Taking $s = 1$ and $s = i$ shows that both coefficients vanish. □

4. Proof of Theorem 1.3

We prove a slightly more general statement, which is also used in Section 6.

Proposition 4.1 (One-label transfer)**.** *Let μ be a G-invariant Borel probability measure on a compact metrizable G-space Ω. Let T_a $(a \in V)$ and R_κ $(\kappa \in K^m)$ be the Koopman operators on $\mathcal{H} = L^2(\Omega, \mu)$, so that $R_\kappa T_a R_\kappa^{-1} = T_{\kappa a}$. Let $f \in \mathcal{H}$ satisfy $0 \le f \le 1$, $R_\kappa f = f$ for all κ, and*

$$\langle f, T_a f\rangle = 0 \qquad \text{for every } a \in V \text{ with } |a| = 1.$$

Let Q be the orthogonal projection onto $\mathcal{H}_c(T)$, the vectors continuous for the whole V-action. Then $p = Qf$ satisfies $0 \le p \le 1$, $\mathbb{E}p = \mathbb{E}f$, and $\langle p, T_x p\rangle = 0$ for every $x \in \mathbb{R}^n$ with $|x| = 1$, where T_x is the extension of Lemma 2.2.

Proof. For a block $j \le m$ let $U_z^{(j)} = T_{\iota_j z}$, where $\iota_j z$ places $z \in E$ in the j-th coordinate plane. Let $W^{(j)}$ be the rotations of that plane, and P_j the projection onto $\mathcal{H}_c(U^{(j)})$. In odd dimension let $S_t = T_{(0,\dots,0,t)}$ for $t \in F$, and let P_0 be the projection onto $\mathcal{H}_c(S)$.

Commutation. All translations commute with one another and with rotations of the other blocks, and axis translations commute with all rotations. By Lemma 3.1, each P_j (and P_0) commutes with every T_a, with the rotations of the other blocks, and, through the self-adjoint case of Lemma 3.1, with the other projections. Covariance gives $\|U_z^{(j)} R_\kappa v - R_\kappa v\| = \|U_{\kappa^{-1}z}^{(j)} v - v\|$, so P_j also commutes with the rotations of its own block. A vector continuous in every block and along the axis is continuous for V, so

$$Q = P_m \cdots P_1 \quad \text{or} \quad Q = P_0 P_m \cdots P_1.$$

Each factor is a conditional expectation (Lemma 2.2). Hence $0 \le p \le 1$, and $\mathbb{E}p = \langle f, Q1\rangle = \mathbb{E}f$.

The plane blocks. Call a unit vector $a \in V$ *generic* if all of its block components $a_j \in E$ are nonzero. Let $g^{(0)} = f$ and $g^{(j)} = P_j g^{(j-1)}$; these are rotation-fixed. Put $v = (1 - P_j) g^{(j-1)}$, which is $W^{(j)}$-fixed and orthogonal to $\mathcal{H}_c(U^{(j)})$. Write $T_a = U_{a_j}^{(j)} Y$, where $Y = T_{a - \iota_j a_j}$ commutes with $U^{(j)}$ and $W^{(j)}$. Lemma 3.2 gives $\langle v, T_a v\rangle = 0$. The cross terms vanish by Lemma 3.1, so

$$\langle g^{(j)}, T_a g^{(j)}\rangle = \langle g^{(j-1)}, T_a g^{(j-1)}\rangle \qquad (a \text{ generic}).$$

With $g = g^{(m)}$ this gives $\langle g, T_a g\rangle = \langle f, T_a f\rangle = 0$ for generic a.

Even n. Here $g = p$. The rational points are dense in S^{n-1} (inverse stereographic projection), and the non-generic ones lie in finitely many proper subspheres. So generic algebraic unit vectors are dense, and continuity of $x \mapsto \langle p, T_x p\rangle$ finishes the proof.

Odd n: the axis. Let $w = g - p \perp \mathcal{H}_c(S)$. Both p and w lie in the range of $P_m \cdots P_1$, so block translations act continuously on them and extend to real arguments. Since P_0 commutes with T_a,

$$\langle p, T_a p\rangle = -\langle w, T_a w\rangle \qquad (a \text{ generic}). \tag{1}$$

Fix an algebraic direction u in the plane blocks with all block components nonzero, and let $\rho(t) = \sqrt{1 - t^2}$. For $t \in F \cap (-1, 1)$ the vector $a(t) = (\rho(t)u, t)$ is a generic algebraic unit vector, because F is real closed. Let $b(t) = \langle p, T_{(\rho(t)u,t)} p\rangle$; it is continuous in real t.

Fix a real $\varphi \in C_c((-1, 1))$. The window $\Phi(t) = \varphi(t)\, T_{(\rho(t)u,0)} w$ is continuous with compact support, hence uniformly continuous. Let

$$\mathcal{V} = I_t\big(S_t \Phi(t)\big).$$

Every $S_t\Phi(t)$ lies in the closed subspace $\mathcal{H}_c(S)^\perp$, so $\mathcal{V}$ does too. By translation invariance of I,

$$\|S_s \mathcal{V} - \mathcal{V}\| \le I_t\big(\|\Phi(t - s) - \Phi(t)\|\big) \to 0 \quad (s \to 0),$$

so $\mathcal{V} \in \mathcal{H}_c(S) \cap \mathcal{H}_c(S)^\perp = \{0\}$. Pairing $\mathcal{V}$ with $c\, w$ ($c \in \mathbb{C}$) and using (1) and Lemma 2.3,

$$0 = I_t\Big(\varphi(t)\, \mathrm{Re}\big(\bar{c}\, \langle w, T_{a(t)} w\rangle\big)\Big) = -\int_{\mathbb{R}} \varphi(t)\, \mathrm{Re}\big(\bar{c}\, b(t)\big)\, dt.$$

Taking $c \in \{1, i\}$ and applying the fundamental lemma of the calculus of variations gives $b \equiv 0$ on $(-1, 1)$. Algebraic directions are dense, so by continuity $\langle p, T_x p\rangle = 0$ for every real unit vector x. □

Proof of Theorem 1.3. Restrict the given coloring to V. Let $\Omega \subseteq \{1, \dots, k\}^V$ be the nonempty, closed, G-invariant set of proper labelings of the unit-distance graph on V, and let μ be a G-invariant probability on Ω, obtained by Følner averaging. Let $f_i = \mathbf{1}[\ell(0) = i]$. These are rotation-fixed, satisfy $\sum_i f_i = 1$, and $\langle f_i, T_a f_i\rangle = \mu\{\ell(0) = i = \ell(a)\} = 0$ for algebraic unit a. Proposition 4.1 gives $p_i = Q f_i \in [0, 1]$ with $\sum_i p_i = Q1 = 1$ and zero correlation at every real unit vector.

Choose jointly measurable fields $H_i(x,\omega) \in [0,1]$ representing $T_x p_i$ for almost every x, as in [13, proof of Proposition 4.2]. Then

$$\int_\Omega \sum_i \int_{B(0,R)} \int_{S^{n-1}} H_i(x,\omega) H_i(x+u,\omega)\, d\sigma\, dx\, d\mu = |B(0,R)| \sum_i \int_{S^{n-1}} \langle p_i, T_u p_i \rangle\, d\sigma = 0.$$

Fix ω such that all these inner integrals vanish for rational R, and such that $\sum_i H_i(\cdot,\omega) = 1$ almost everywhere. Color each x by the least i with $H_i(x,\omega) > 0$. This coloring is measurable, and its monochromatic unit pairs form a $dx\, d\sigma$-null set. □

Proof of Corollary 1.4. Let A_i^* be the set of Lebesgue density points of $A_i = c^{-1}(i)$. These sets are Borel and pairwise disjoint, and their union is conull by the Lebesgue differentiation theorem. By [13, Lemma 4.3], whose proof is dimension-free, A_i^* contains no two points at distance 1. Since $\sum_i |A_i^* \cap [-T,T]^n| = (2T)^n$ for every T, some class has upper density at least $1/k$. Hence $1/k \le m_1(\mathbb{R}^n)$. □

5. Numerical input

The only numerical input is $m_1(\mathbb{R}^3) \le 0.1532996$ [4, Table 2]. DeCorte, Oliveira and Vallentin provide the dual solutions and an exact verifier (`rn_verify`, GMP/MPFR) as ancillary files. We rebuilt the verifier, after one portability fix that makes a base-class destructor virtual and does not affect the arithmetic, and re-ran it on the published data. For $n = 3$ it reports "Solution checked; bound = 0.153299568560593". Since $6 \times 0.1533 < 1$, Corollary 1.4 excludes proper 6-colorings of $\mathbb{R}^3$. The same corollary combined with [4, Table 2] gives $\chi(\mathbb{R}^4) \ge 11$; for $n \ge 5$ the bounds of Davies [7] are stronger than those obtained from [4, 2].

To obtain $\chi(\mathbb{R}^3) \ge 8$ in this way one would need $m_1(\mathbb{R}^3) < 1/7 \approx 0.1429$. The derivative constraints of [7, Section 9] do not improve the bound for $n = 3$, since only one radial derivative is admissible there.

6. Finite graphs attain the measurable density

Proof of Theorem 1.6. (i) $m_1 \le \alpha_1$. Let A be measurable and unit-distance-free, with densities $d_j \to d$ along centered cubes Q_j. For a finite unit-distance graph G,

$$\frac{1}{|Q_j|} \int_{Q_j} |(A-x) \cap G|\, dx = \sum_{g \in G} \frac{|A \cap (Q_j + g)|}{|Q_j|} \to d|G|,$$

so some translate of A meets G in an independent set of size at least $(d-\varepsilon)|G|$. (This also shows that the definition of m_1 via $\inf_X \sup_A \lambda(A)/\lambda(X)$ used in [11] agrees with ours.)

(ii) $\sup_G \chi_f(G) \ge 1/m_1$. Suppose every finite subgraph of the unit-distance graph on V has $\chi_f \le w$. Normalizing optimal fractional colorings gives, for each finite $H \subset V$, a random independent set of H covering *each* vertex with probability at least $1/w$. (Because the bound holds at every vertex, it survives the averaging below; compare [10, Remark 5.4].) A weak-$*$ limit in $\{0,1\}^V$, in which cylinder events are clopen, yields a random independent set I of V with $\mathbb{P}(v \in I) \ge 1/w$ for all v. Følner averaging over G makes its law invariant without affecting these properties.

Apply Proposition 4.1 to $f = \mathbf{1}[0 \in I]$, and let $H(x,\omega)$ be jointly measurable versions of $T_x p$. For almost every ω, $H(x)H(x+u) = 0$ for $dx\, d\sigma$-almost every (x,u). The cube averages $X_T(\omega) = |Q_T|^{-1} \int_{Q_T} H(x,\omega)\, dx \in [0,1]$ have mean $\mathbb{E}p \ge 1/w$, so by reverse Fatou $\mathbb{P}(\limsup_T X_T \ge 1/w) > 0$. For such an ω, the density points of $\{H(\cdot,\omega) > 0\}$ form a measurable unit-distance-free set of upper density at least $1/w$. Hence $w \ge 1/m_1$.

(iii) The proof of [11, Theorem 2], $\sup_G \chi_f(G) = 1/\alpha_1$, uses only translations through a lattice cube and holds verbatim in $\mathbb{R}^n$. Together with (i) and (ii) this gives $\alpha_1 = m_1$.

(iv) $\chi_{gf}(G) \le 1/m_1$. This is [1, Corollary 2], whose proof works in $\mathbb{R}^n$: given $\varepsilon > 0$, truncating a unit-distance-free set to a large cube, removing a boundary layer of width 1 and repeating periodically gives a Λ-periodic measurable unit-distance-free set A with $\delta(A) \ge m_1 - \varepsilon$. Let $g(x) = \phi x + \tau$ with ϕ Haar-random in $O(n)$ and τ independent and uniform in a fundamental domain of Λ, and put $\gamma(S) = \mathbb{P}(gG \cap A = gS)/\delta(A)$ for $S \subseteq G$. Then γ vanishes off independent sets, $\sum_{S \ni x} \gamma(S) = \mathbb{P}(gx \in A)/\delta(A) = 1$, and $\sum_S \gamma(S) = 1/\delta(A)$. If $S' = \rho S + b$ with $\rho \in O(n)$, then $gS' = (\phi\rho)S + (\phi b + \tau)$, and $(\phi\rho, \phi b + \tau)$ has the same law as (ϕ, τ) modulo Λ. So $\sum_{T \supseteq S} \gamma(T) = \mathbb{P}(gS \subseteq A)/\delta(A)$ satisfies the congruence constraints, and $\chi_{gf}(G) \le 1/\delta(A)$. Since $\chi_f \le \chi_{gf}$, all three suprema coincide.

Finally, f is subadditive (place copies far apart), so Fekete's lemma gives $f(N)/N \to \inf_N f(N)/N = \alpha_1(\mathbb{R}^2) = m_1(\mathbb{R}^2)$, recovering [10, Theorem 1.1]. In the plane, $\sup_G \chi_f(G) = 1/m_1(\mathbb{R}^2) > 4.0486$ by the bound $m_1(\mathbb{R}^2) \le 0.246998$ [1, Section 8]. □

7. Sets of algebraic distances

For a set $D \subset (0, \infty)$ let $\chi_D(\mathbb{R}^n) = \chi(\mathbb{R}^n, D)$ be the least number of colors in a coloring of $\mathbb{R}^n$ with no monochromatic pair at a distance in D, and let $m_D(\mathbb{R}^n)$ be the supremal upper density of a measurable set with no two points at a distance in D.

Theorem 7.1. *Let $n \ge 2$ and let D be a countable set of positive real algebraic numbers. A coloring of $\mathbb{R}^n$ with k colors and no monochromatic pair at a distance in D yields k pairwise disjoint measurable sets, covering $\mathbb{R}^n$ up to a null set, none containing two points at a distance in D. In particular $\chi_D(\mathbb{R}^n) \ge 1/m_D(\mathbb{R}^n)$.*

Proof. We run the proof of Theorem 1.3 and Corollary 1.4 with the unit sphere replaced by the spheres of radii $d \in D$. Proposition 4.1 holds with the hypothesis and conclusion stated for $|a| \in D$ and $|x| \in D$, with the same proof. Properness gives $\langle f_i, T_a f_i \rangle = 0$ for algebraic a with $|a| \in D$. The block step uses only Lemma 3.2, whose conclusion holds at every generic a regardless of its length, because Theorem 2.1 gives zero correlation at every nonzero $z \in E$. In odd dimension the axis step uses the curves $t \mapsto (\sqrt{d^2 - t^2}\, u, t)$, $|t| < d$, which consist of algebraic points because d is algebraic and F is real closed, with test functions $\varphi \in C_c((-d, d))$. Algebraic points are dense on each sphere of algebraic radius. The density-point lemma [13, Lemma 4.3] applies to each $d \in D$. So a proper k-coloring yields k measurable D-avoiding sets covering $\mathbb{R}^n$ up to a null set, and one of them has upper density at least $1/k$. For countable D the conditions for different $d \in D$ are countably many, and the sampling step chooses one ω satisfying all of them. □

Theorem 7.1 is formalized as `euclidean_transfer_set`. In the companion paper [6] it is combined with Fourier certificates and with the Furstenberg–Katznelson–Weiss theorem to show that every finite coloring of the plane has monochromatic pairs at distance 2^n and at distance $n!$ for all large n (questions of Soifer and Kahle), that $\sup_{|D|=r} \chi_D(\mathbb{R}^2)$ grows exponentially in r (Erdős Problem #706), and quantitative versions of these statements, several of them formally verified.

8. Formal verification

The Lean development (Lean 4.34.1, Mathlib commit `d13f23b`) builds on the 70 modules of the library of [13] on which its planar result depends, in particular its formal proof of [13, Theorem 2.3] (`PlaneFiveColor.Spectral.residual_fourier`). It has three parts.

- `OAI/HN3` (about 1,850 lines) treats $\mathbb{R}^3$ as $\mathbb{C} \times \mathbb{R}$. Its main statements are `three_space_transfer`, `no_six_coloring_of_density` and `exists_finite_not_six_colorable`.

- `OAI/HNN` (about 2,200 lines) treats $\mathbb{R}^n$ for every $n \geq 2$, modelled as $\mathbb{C}^m \times \mathbb{R}^r$ with $r \in \{0, 1\}$ and transported to `EuclideanSpace R (Fin n)`. Its main statements are `euclidean_transfer_set` (any countable set D of positive algebraic distances), `colors_ge_of_densityE` `no_6_coloring_R3` through `no_52_coloring_R8`, and `exists_finite_not_colorableE`.
- `OAI/HN3/Indep*` (about 1,900 lines) proves `finite_ratio_of_density` and `alpha1_le_m1`: for every $\varepsilon > 0$ there is a finite point set in $\mathbb{R}^3$ whose unit-distance graph has independence ratio at most $\delta + \varepsilon$ whenever $m_1(\mathbb{R}^3) \leq \delta$. It averages a positive functional, obtained from Følner boxes in the finitely generated subgroups of the algebraic points, instead of a point mass.

`DensityBound` δ states that every measurable unit-distance-free set has upper density at most δ along the cubes $[-T, T]^n$. `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound` for every statement listed.

9. Remarks

Why the torus suffices. Theorem 1.3 needs neither invariance under all of $SO(n)$ nor a disintegration of spectral measures. A maximal torus K^m of algebraic rotations is abelian, so averaging over it is possible. Rigidity applied to one block at a time, via Lemma 3.2, then removes every spectral component that is discontinuous in some block. The remaining axis in odd dimension is handled by the invariant mean, as in [13, Lemma 2.5].

The plane. By [13, Theorem 1.3], deciding between $\chi(\mathbb{R}^2) = 6$ and 7 is equivalent to deciding whether weak measurable 6-colorings of the plane exist. Density and fractional methods cannot settle this, since by Theorem 1.6 they are capped at $1/m_1(\mathbb{R}^2) \leq 1/0.2293 < 4.37$.

References

[1] G. Ambrus, A. Csiszárik, M. Matolcsi, D. Varga, P. Zsámboki, *The density of planar sets avoiding unit distances*, Math. Program. (2023); arXiv:2207.14179.

[2] C. Bachoc, A. Passuello, A. Thiery, *The density of sets avoiding distance 1 in Euclidean space*, Discrete Comput. Geom. 53 (2015); arXiv:1401.6140.

[3] D. Coulson, *A 15-colouring of 3-space omitting distance one*, Discrete Math. 256 (2002).

[4] E. DeCorte, F. M. de Oliveira Filho, F. Vallentin, *Complete positivity and distance-avoiding sets*, Math. Program. (2022); arXiv:1804.09099.

[5] C. Crocker, *A conditional transfer for colourings of $\mathbb{R}^n$, with the conditional consequence $\chi(\mathbb{R}^3) \geq 7$*, Zenodo (10 October 2026), doi:10.5281/zenodo.23273952.

[6] H. Bass, *Monochromatic distances in finite colorings of the plane*, preprint (October 2026).

[7] J. Davies, *Measurable obstructions for unmeasurable colourings*, arXiv:2610.12301 (2026).

[8] A. de Grey, *The chromatic number of the plane is at least 5*, Geombinatorics 28 (2018).

[9] Á. Dúcz, D. Varga, *A unit-distance graph in the plane with independence ratio below 1/4*, arXiv:2606.28157 (2026).

[10] S. Li, Y. Su, *Every n points in the plane contain $(m_1 + o(1))n$ points with no unit distance*, arXiv:2610.15424 (2026).

[11] M. Matolcsi, I. Z. Ruzsa, D. Varga, P. Zsámboki, *The fractional chromatic number of the plane is at least 4*, arXiv:2311.10069.

[12] O. Nechushtan, *On the space chromatic number*, Discrete Math. 256 (2002).

[13] OpenAI, *The Euclidean plane is not five-colorable*, preprint (2026), https://github.com/openai/math, with Lean formalization.

[14] F. M. de Oliveira Filho, F. Vallentin, *Fourier analysis, linear programming, and densities of distance avoiding sets in $\mathbb{R}^n$*, J. Eur. Math. Soc. 12 (2010).

[15] R. Radoičić, G. Tóth, *Note on the chromatic number of the space*, in Discrete and Computational Geometry, Springer (2003).