

# Affine Jump-Diffusions: Stochastic Stability and Limit Theorems

Xiaowei Zhang

Department of Industrial Engineering and Logistics Management  
The Hong Kong University of Science and Technology

May 11, 2017


1. Introduction
2. Stochastic Stability
3. Limit Theorems
4. Application to Affine Point Processes
5. Conclusions

- ▶ Widely used in finance and econometrics
  - Vasicek (1977)
  - Cox et al. (1985)
  - Heston (1993)
  - Bates (2000), Duffie et al. (2000), Barndorff-Nielsen and Shephard (2001), Cheridito et al. (2007), Errais et al. (2010), etc.

- ▶ Widely used in finance and econometrics
  - Vasicek (1977)
  - Cox et al. (1985)
  - Heston (1993)
  - Bates (2000), Duffie et al. (2000), Barndorff-Nielsen and Shephard (2001), Cheridito et al. (2007), Errais et al. (2010), etc.
- ▶ Computationally tractable
  - Fourier transform is easy to compute (by solving **ODEs**)
- ▶ Defining SDE has “affine” structure
  - drift
  - variance
  - jump intensityare all affine in the state variable

# AJD Examples


- ▶ Ornstein-Uhlenbeck (OU) process in Vasicek model
  - $dX(t) = (b - \beta X(t)) dt + \sigma dW(t)$



# AJD Examples


- ▶ Ornstein-Uhlenbeck (OU) process in Vasicek model

- $dX(t) = (b - \beta X(t)) dt + \sigma dW(t)$




- ▶ square-root diffusion process in CIR model

- $dX(t) = (b - \beta X(t)) dt + \sigma \sqrt{X(t)} dW(t)$




- ▶ Ornstein-Uhlenbeck (OU) process in Vasicek model

- $dX(t) = (b - \beta X(t)) dt + \sigma dW(t)$



- ▶ square-root diffusion process in CIR model

- $dX(t) = (b - \beta X(t)) dt + \sigma \sqrt{X(t)} dW(t)$



- ▶ Mean-reversion:  $\beta > 0$

- ▶ Long-term behavior
  - Does there exist an equilibrium?
  - How fast does the process converge to the equilibrium?
  - Can LLN or CLT be established?



- ▶ Long-term behavior
  - Does there exist an equilibrium?
  - How fast does the process converge to the equilibrium?
  - Can LLN or CLT be established?
- ▶ Parameter estimation based on large-time asymptotics
  - $X$  is observed at times  $0, \Delta, \dots, n\Delta$
  - Estimating equation for the unknown parameter  $\Theta$

$$\frac{1}{n} \sum_{k=1}^n h(X(\Delta_{k-1}), X(\Delta_k); \hat{\Theta}_n) = 0$$

- Maximum likelihood, (generalized) method of moments, least squares, etc.
- LLN  $\Rightarrow$  consistency of  $\hat{\Theta}$
- CLT  $\Rightarrow$  asymptotic normality of  $\hat{\Theta}$

- ▶ Sato and Yamazato (1984), Masuda (2004): Lévy-driven OU process

$$dX(t) = -\beta X(t) dt + dJ(t)$$

- ▶ Glasserman and Kim (2010), Jena et al. (2012): affine diffusions (without jumps)
- ▶ Keller-Ressel (2011): a one-dimensional AJD
- ▶ Barczy et al. (2014): a two-dimensional AJD with Lévy-type jumps

- ▶ Consider the following 1-dimensional process

$$dX(t) = (b - \beta X(t)) dt + \sigma \sqrt{X(t)} dW(t) + dJ(t)$$

- ▶  $J(t) = \sum_{i=1}^{N(t)} Z_i$ ,  $Z_i$ 's are iid with  $\mathbb{E}|Z_1|^p < \infty$  for some  $p > 0$
- ▶  $N(t)$  is a counting process with intensity  $\kappa + \lambda X(t)$

- ▶ Consider the following 1-dimensional process

$$dX(t) = (b - \beta X(t)) dt + \sigma \sqrt{X(t)} dW(t) + dJ(t)$$

- ▶  $J(t) = \sum_{i=1}^{N(t)} Z_i$ ,  $Z_i$ 's are iid with  $\mathbb{E}|Z_1|^p < \infty$  for some  $p > 0$
- ▶  $N(t)$  is a counting process with intensity  $\kappa + \lambda X(t)$
- ▶ Two factors that determine stability
  - mean-reversion: good
  - jumps: detrimental

- ▶ Consider the following 1-dimensional process




$$dX(t) = (b - \beta X(t)) dt + \sigma \sqrt{X(t)} dW(t) + dJ(t)$$

- ▶  $J(t) = \sum_{i=1}^{N(t)} Z_i$ ,  $Z_i$ 's are iid with  $\mathbb{E}|Z_1|^p < \infty$  for some  $p > 0$
- ▶  $N(t)$  is a counting process with intensity  $\kappa + \lambda X(t)$
- ▶ Two factors that determine stability
  - mean-reversion: good
  - jumps: detrimental

**Takeaway:** If  $\beta > \lambda \mathbb{E}(Z_1)$ , then  $X$  is “stable”.

## Special Case: Compound Poisson Jumps

- ▶ Suppose  $N(t)$  is independent of  $X(t)$ , i.e.  $\lambda = 0$ 
  - $J(t) = \sum_{i=1}^{N(t)} Z_i$  is a compound Poisson process
- ▶  $X(t)$  is stable if  $\beta > 0$



- ▶ Follow the formulation in Duffie et al. (2003)
- ▶  $X(t) \in \mathcal{X} = \mathbb{R}_+^m \times \mathbb{R}^{d-m}$  satisfies

$$\begin{cases} dX(t) = \mu(X(t)) dt + \sigma(X(t)) dW(t) + dJ(t) \\ J(t) = \sum_{i=1}^{N(t)} Z_i \end{cases}$$

$N(t)$  is a counting process with intensity  $\Lambda(X(t))$ , where

$$\mu(x) = b - \beta x, \quad b \in \mathbb{R}^d, \beta \in \mathbb{R}^{d \times d}$$

$$\sigma(x)\sigma(x)^\top = a + \sum_{i=1}^d x_j \alpha_i, \quad a \in \mathbb{R}^{d \times d}, \alpha_i \in \mathbb{R}^{d \times d}, i = 1, \dots, d$$

$$\Lambda(x) = \kappa + \lambda^\top x, \quad \kappa \in \mathbb{R}, \lambda \in \mathbb{R}^d$$

- ▶ Assume  $\mathbb{E}\|Z_1\|^p < \infty$  for some  $p > 0$

# Insights from 1-D Case

1. Mean-reversion, i.e.  $\beta$  is “positive”
2. Mean-reversion “dominates” jumps



1. Mean-reversion, i.e.  $\beta$  is “positive”
2. Mean-reversion “dominates” jumps

How to define “positivity” for matrices?

How to represent “dominance” for matrices?

1. Mean-reversion, i.e.  $\beta$  is “positive”
2. Mean-reversion “dominates” jumps

How to define “positivity” for matrices?

- ▶ ODE analog:  $X'(t) = b - \beta X(t)$
- ▶  $X$  is asymptotically stable if and only if  $\beta$  is **positive stable**
  - all the eigenvalues of  $\beta$  have positive real parts

How to represent “dominance” for matrices?

1. Mean-reversion, i.e.  $\beta$  is “positive”
2. Mean-reversion “dominates” jumps

How to define “positivity” for matrices?

- ▶ ODE analog:  $X'(t) = b - \beta X(t)$
- ▶  $X$  is asymptotically stable if and only if  $\beta$  is **positive stable**
  - all the eigenvalues of  $\beta$  have positive real parts

How to represent “dominance” for matrices?

- ▶  $\beta - (\mathbb{E}Z_1)\lambda^\top$  is positive stable
  - **strong mean-reversion** condition

## Theorem

If  $\beta - \mathbb{E}(Z_1)\lambda^\top$  is positive stable, then  $X$  is **exponentially ergodic**, i.e.,

$$\|\mathbb{P}_x(X(t) \in \cdot) - \pi(\cdot)\| \leq c(x)e^{-\rho t}, \quad t \geq 0,$$

for some positive function  $c(\cdot)$  and some positive constant  $\rho$ , where  $\pi$  is the unique stationary distribution of  $X$ .

# Main Result

## Theorem

If  $\beta - \mathbb{E}(Z_1)\lambda^\top$  is positive stable, then  $X$  is **exponentially ergodic**, i.e.,

$$\|\mathbb{P}_x(X(t) \in \cdot) - \pi(\cdot)\| \leq c(x)e^{-\rho t}, \quad t \geq 0,$$

for some positive function  $c(\cdot)$  and some positive constant  $\rho$ , where  $\pi$  is the unique stationary distribution of  $X$ .

- Proof relies on the “Lyapunov approach” (Meyn and Tweedie, 1993)

# Lyapunov Approach

- ▶ Originated in the study of stability of ODE's
- ▶ Extended to stochastic stability of Markov processes in 1970's (Meyn and Tweedie, 2009)
- ▶ Establish a Lyapunov inequality: find constants  $c > 0$ ,  $d < \infty$  and a norm-like function  $V \geq 0$ , for which

$$\mathcal{A}V(x) \leq -cV(x) + d, \quad \text{for all } \|x\| \text{ large enough,}$$

where  $\mathcal{A}$  is the infinitesimal generator of  $X$

$$\mathcal{A}g(x) \triangleq \mathcal{G}g(x) + \mathcal{L}g(x)$$

$$\mathcal{G}g(x) \triangleq \nabla g(x) \cdot (b + \beta x) + \frac{1}{2} \sum_{i,j=1}^d \frac{\partial^2 g(x)}{\partial x_i \partial x_j} \left( a_{i,j} + \sum_{k=1}^d \alpha_{k,ij} x_k \right)$$

$$\mathcal{L}g(x) \triangleq (\lambda + \kappa^\top x) \int_x (g(x+z) - g(x)) \nu(dz)$$

- ▶ If  $B = \beta - (\mathbb{E}Z_1)\lambda^\top$  is positive stable, then there exists a positive definite matrix  $H$ , denoted as  $H \succ 0$ , such that

$$HB + B^\top H \succ 0$$

# Lyapunov Function

- ▶ If  $B = \beta - (\mathbb{E}Z_1)\lambda^\top$  is positive stable, then there exists a positive definite matrix  $H$ , denoted as  $H \succ 0$ , such that

$$HB + B^\top H \succ 0$$

- ▶ Set  $V(x) = \|x\|_H^p$ , where  $\|x\|_H = (x^\top Hx)^{\frac{1}{2}}$ , and show

$$\mathcal{A}V(x) = pV(x) \left( -\frac{x^\top (HB + B^\top H)x}{2\|x\|_H^2} + o(1) \right)$$

as  $\|x\| \rightarrow \infty$



## Remark on Strong Mean-Reversion

- ▶ Positive stability of  $\beta - \mathbb{E}(Z_1)\lambda^\top$  cannot be relaxed in general

## Remark on Strong Mean-Reversion

- ▶ Positive stability of  $\beta - \mathbb{E}(Z_1)\lambda^\top$  cannot be relaxed in general
- ▶ Revisit the following 1-d process

$$dX(t) = (b - \beta X(t)) dt + \sigma \sqrt{X(t)} dW(t) + dJ(t)$$

- If  $\beta - \lambda \mathbb{E}(Z_1) < 0$ , then this process is *transient*
- Proof also relies on the Lyapunov approach

## Theorem (SLLN)

If  $\beta - \mathbb{E}(Z_1)\lambda^\top$  is positive stable and  $|h(x)| \leq C\|x\|^p$ , then

$$\mathbb{P}_x \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n h(X(i\Delta)) = \pi(h) \right) = 1, \quad x \in \mathcal{X}.$$

## Theorem (FCLT)


If  $\beta - \mathbb{E}(Z_1)\lambda^\top$  is positive stable and  $|h(x)|^{2+\epsilon} \leq C\|x\|^p$ , then

$$n^{1/2} \left( \frac{1}{n} \sum_{i=1}^{\lfloor n \cdot \rfloor} h(X(i\Delta)) - \pi(h) \right) \Rightarrow \gamma_h W(\cdot)$$


as  $n \rightarrow \infty$   $\mathbb{P}_x$ -weakly in  $\mathcal{D}[0, 1]$  for each  $x \in \mathcal{X}$ .

# Affine Point Processes

- ▶ The jump process in an AJD
- ▶ Hawkes process is a popular example (limit order book, credit default)
- ▶ Jump intensity is state-dependent, capturing the clustering of jumps




# Annual Defaults of Moodys rated U.S. Firms



# CLT for Affine Point Processes

- ▶  $J(t) \overset{\mathcal{D}}{\approx} rt + \mathcal{N}(0, \eta^2 t)$  for large  $t$



- ▶ Proved exponential ergodicity of AJDs under a simple condition
  - Strong mean-reversion cannot be relaxed in general
- ▶ Proved SLLN and FCLT
  - provide theoretical support for many estimation methods for AJDs

**Thanks!**



- M. Barczy, L. Döring, Z. Li, and G. Pap. Stationarity and ergodicity for an affine two-factor models. *Adv. Appl. Probab.*, 46(3):878–898, 2014.
- O. E. Barndorff-Nielsen and N. Shephard. Non-Gaussian Ornstein-Uhlenbeck-based models and some of their uses in financial economics. *J. R. Statist. Soc. B*, 63(2):167–241, 2001.
- D. S. Bates. Post-'87 crash fears in the S&P 500 futures option market. *J. Econometrics*, 94(1-2):181–238, 2000.
- P. Cheridito, D. Filipović, and R. L. Kimmel. Market price of risk specification for affine models: Theory and evidence. *J. Financ. Econ.*, 83:123–170, 2007.
- J. C. Cox, J. E. Ingersoll, and S. A. Ross. A theory of the term structure of interest rates. *Econometrica*, 53(2):385–407, 1985.
- D. Duffie, J. Pan, and K. J. Singleton. Transform analysis and asset pricing for affine jump-diffusions. *Econometrica*, 68(6):1343–1376, 2000.
- D. Duffie, D. Filipović, and W. Schachermayer. Affine processes and applications in finance. *Ann. Appl. Probab.*, 13(3):984–1053, 2003.
- E. Errais, K. Giesecke, and L. R. Goldberg. Affine point processes and portfolio credit risk. *SIAM J. Finan. Math.*, 1:642–665, 2010.

- P. Glasserman and K.-K. Kim. Moment explosions and stationary distributions in affine diffusion models. *Math. Finance*, 20(1):1–33, 2010.
- S. L. Heston. A closed-form solution for options with stochastic volatility with applications to bond and currency options. *Rev. Financ. Stud.*, 6: 327–343, 1993.
- R. P. Jena, K.-K. Kim, and H. Xing. Long-term and blow-up behaviors of exponential moments in multi-dimensional affine diffusions. *Stoch. Proc. Appl.*, 122:2961–2993, 2012.
- M. Keller-Ressel. Moment explosions and long-term behavior of affine stochastic volatility model. *Math. Finance*, 21(1):73–98, 2011.
- H. Masuda. On multidimensional Ornstein-Uhlenbeck processes driven by a general Lévy process. *Bernoulli*, 10:97–120, 2004.
- S. P. Meyn and R. L. Tweedie. Stability of Markovian processes III: Foster-Lyapunov criteria for continuous-time processes. *Adv. Appl. Probab.*, 25:518–548, 1993.
- S. P. Meyn and R. L. Tweedie. *Markov Chains and Stochastic Stability*. Cambridge University Press, 2 edition, 2009.

## References III

- K.-i. Sato and M. Yamazato. Operator-self-decomposable distributions as limit distributions of processes of Ornstein–Uhlenbeck type. *Stoch. Proc. Appl.*, 17:73–100, 1984.
- O. Vasicek. An equilibrium characterization of the term structure. *J. Financ. Econ.*, 5:177–188, 1977.