

# 18.102 Assignment 6

Octavio Vega

April 4, 2023

## Problem 1

(a)

*Proof.* Let  $\epsilon > 0$ , and define  $c_1 := a$  and  $c_{n+1} := b$ .

Since  $\psi$  is a step function on  $[a, b]$ ,  $\exists c_1 \leq c_2 \leq \dots \leq c_n \leq c_{n+1} \in [a, b]$  such that  $\forall i = 1, \dots, n$ ,

$$\psi^{-1}(\{a_i\}) = (c_i, c_{i+1}], \quad (1)$$

where each  $a_i$  is one of the finitely many values that  $\psi$  takes on.

Choose  $\delta > 0$  such that  $\delta < \frac{\epsilon}{2n}$ . Define  $g : [a, b] \rightarrow \mathbb{R}$  via

$$g(x) := \begin{cases} \frac{a_i + a_{i+1}}{2} + \left( \frac{a_i - a_{i+1}}{2\delta} \right) (x - c_i), & x \in (c_i - \delta, c_i + \delta) \\ a_i, & x \in [c_i + \delta, c_{i+1} - \delta] \\ -\frac{a_n}{2\delta} (x - b), & x \in (c_{n-1} - \delta, b], \end{cases} \quad (2)$$

where  $a_0 := -a_1$ . Then

$$g(a) = \frac{a_1 + a_0}{2} + \left( \frac{a_1 - a_0}{2\delta} \right) (c_1 - c_1) = \frac{a_1 - a_1}{2} = 0, \quad (3)$$

and

$$g(b) = -\frac{a_n}{2\delta} (b - b) = 0, \quad (4)$$

as desired. We also see that since  $g$  is piecewise linear, it is continuous.

Now consider the difference  $|\psi(x) - g(x)|$ .

Case 1:  $x \in [c_i + \delta, c_{i+1} - \delta]$ . Then by (1), we have

$$|\psi(x) - g(x)| = |\psi((c_i, c_{i+1} - \delta)) - a_i| = |a_i - a_i| = 0. \quad (5)$$

Case 2:  $x \in (c_i - \delta, c_i + \delta)$ . Then

$$|\psi(x) - g(x)| = \left| \frac{a_i + a_{i-1}}{2} + \left( \frac{a_i - a_{i-1}}{2\delta} \right) (x - c_i) - \psi(x) \right| \quad (6)$$

$$< \left| \frac{a_i + a_{i-1}}{2} + \left( \frac{a_i - a_{i-1}}{2\delta} \right) \delta - \psi(x) \right| \quad (7)$$

$$= \left| \frac{a_i + a_{i-1}}{2} + \frac{a_i + a_{i-1}}{2} - \psi(x) \right| \quad (8)$$

$$= |a_i - \psi(x)| \quad (9)$$

$$= 0 \text{ or } |a_{i+1} - a_i|. \quad (10)$$

Case 3:  $x \in (c_n - \delta, b)$ . Then

$$|\psi(x) - g(x)| = \left| -\frac{a_n}{2\delta}(x - b) - \psi(x) \right| \quad (11)$$

$$< \left| -\frac{a_n}{2\delta}\delta - \psi(x) \right| \quad (12)$$

$$= \left| -\frac{a_n}{2} - \psi(x) \right| \quad (13)$$

$$= \frac{3a_n}{2}. \quad (14)$$

So in all three cases, we have that either  $|g(x) - \psi(x)| = 0$ , or  $|g(x) - \psi(x)| < \frac{3a}{2}$ , or  $|g(x) - \psi(x)| < |a_{i+1} - a_i|$ .

Define the set  $E$  to be the collection of points in  $[a, b]$  for which  $|\psi(x) - g(x)| \neq 0$ . Then

$$E := \bigcup_{k=1}^n (c_k - \delta, c_k + \delta). \quad (15)$$

By definition,  $\forall x \in E^c$ ,

$$|\psi(x) - g(x)| = 0 < \epsilon. \quad (16)$$

Since  $E$  is a countable union of intervals, we have

$$m(E) = m \left[ \bigcup_{k=1}^n (c_k - \delta, c_k + \delta) \right] \quad (17)$$

$$\leq \sum_{k=1}^n m[(c_k - \delta, c_k + \delta)] \quad (18)$$

$$= \sum_{k=1}^n \ell(c_k - \delta, c_k + \delta) \quad (19)$$

$$= \sum_{k=1}^n 2\delta \quad (20)$$

$$= 2n\delta \quad (21)$$

$$< 2n \frac{\epsilon}{2n} \quad (22)$$

$$= \epsilon, \quad (23)$$

as desired.  $\square$