

18.100A Assignment 10

Octavio Vega

June 6, 2023

Problem 1

(a)

Proof. Suppose $\exists C \geq 0$ such that $\forall x, y \in I$,

$$|f(x) - f(y)| \leq C|x - y|^\alpha. \quad (1)$$

Let $\epsilon > 0$. Choose $\delta = \left(\frac{\epsilon}{C}\right)^{\frac{1}{\alpha}}$. Then if $|x - y| < \delta$, we get

$$|f(x) - f(y)| \leq C|x - y|^\alpha \quad (2)$$

$$< C\delta^\alpha \quad (3)$$

$$= C\frac{\epsilon}{C} \quad (4)$$

$$= \epsilon. \quad (5)$$

Therefore f is uniformly continuous on I . \square

(b)

Proof. Suppose $\exists C \geq 0$ such that $\forall x, y \in I$, $|f(x) - f(y)| \leq C|x - y|^\alpha$.

Since $\alpha > 1$, then $\alpha = 1 + r$ for some $0 < r$, we have

$$\implies 0 \leq |f(x) - f(y)| \leq C|x - y|^{1+r} \quad (6)$$

$$\implies 0 \leq \frac{|f(x) - f(y)|}{|x - y|} \leq C|x - y|^r \quad (7)$$

$$\implies \lim_{x \rightarrow y} 0 \leq \lim_{x \rightarrow y} \frac{|f(x) - f(y)|}{|x - y|} \leq C \lim_{x \rightarrow y} |x - y|^r \quad (8)$$

$$\implies 0 \leq \lim_{x \rightarrow y} \frac{|f(x) - f(y)|}{|x - y|} \leq 0. \quad (9)$$

Then by the squeeze theorem, $\lim_{x \rightarrow y} \frac{|f(x) - f(y)|}{|x - y|} = 0$. Thus $\forall y \in I$, $f'(y) = 0$.

Therefore f is constant. \square