

# 18.100A Assignment 3

Octavio Vega

February 16, 2023

## Problem 1

*Proof.* Let  $x, y \in \mathbb{R}$  with  $x < y$ . By the density of  $\mathbb{Q}$ , we have that  $\exists r \in \mathbb{Q}$  such that  $x < r < y$ .

Then  $x + \sqrt{2} < y + \sqrt{2}$ . Then  $\exists r \in \mathbb{Q}$  such that

$$x + \sqrt{2} < r < y + \sqrt{2} \quad (1)$$

$$\implies x < r - \sqrt{2} < y. \quad (2)$$

But since  $r \in \mathbb{Q}$  and  $\sqrt{2} \notin \mathbb{Q}$ , then the number  $i := r - \sqrt{2} \notin \mathbb{Q}$ .

So  $x < i < y$  with  $i \in \mathbb{R} \setminus \mathbb{Q}$ , as desired.  $\square$

## Problem 2

*Proof.* Define the function  $f : E \rightarrow \wp(\mathbb{N})$  such that if  $x = 0.d_{-1}d_{-2}\dots$ , then

$$f(x) = \{j \in \mathbb{N} \mid d_{-j} = 2\}. \quad (3)$$

We want to show that  $f$  is a bijection. First, we show that  $f$  is injective.

Let  $x_1 = 0.d_{-1}^{(1)}d_{-2}^{(1)}\dots$  and  $x_2 = 0.d_{-1}^{(2)}d_{-2}^{(2)}\dots$  for  $x_1, x_2 \in E$ . Suppose  $f(x_1) = f(x_2)$ . Then

$$\{j \in \mathbb{N} \mid d_{-j}^{(1)} = 2\} = \{k \in \mathbb{N} \mid d_{-k}^{(2)} = 2\}. \quad (4)$$

Since each digit  $d_{-j} \in \{1, 2\}$ , then the sets of digits must be the same:

$$\{d_{-j}^{(1)} \mid j \in \mathbb{N}\} = \{d_{-k}^{(2)} \mid k \in \mathbb{N}\}. \quad (5)$$

But by the theorem from class, we know that for every set of digits  $\exists! x \in [0, 1]$  such that  $x = 0.d_{-1}d_{-2}\dots$ . So if all of the digits are the same, then the numbers must be the same, i.e.

$$f(x_1) = f(x_2) \implies x_1 = x_2. \quad (6)$$

Thus  $f$  is injective.

Next, we show that  $f$  is surjective.

Let  $S \in \wp(\mathbb{N})$  with

$$S := \{j \in \mathbb{N} \mid d_{-j} = 2\}. \quad (7)$$

Since this corresponds to the indices for a set of digits, then by the theorem from class  $\exists x \in [0, 1]$  such that  $x = 0.d_{-1}d_{-2}\dots$ ; i.e. for any  $S \in \wp(\mathbb{N})$ ,  $\exists x \in E$  such that  $f(x) = S$ .

Hence,  $f$  is also surjective, which means that it is bijective.

Therefore we conclude that  $|E| = |\wp(\mathbb{N})|$ .  $\square$

### Problem 3

(a)

*Proof.* We want to show that there exists a bijection  $h : A \cup B \rightarrow \mathbb{N}$ .

Recall that we can construct the sets of even and odd natural numbers as follows:

$$\mathcal{O} := \{2n + 1 \mid n \in \mathbb{N}\}, \text{ and } \mathcal{E} := \{2n \mid n \in \mathbb{N}\}. \quad (8)$$

We also know that  $|\mathcal{O}| = |\mathcal{E}| = |\mathbb{N}|$  because the functions defined via  $f_e(n) = 2n$  and  $f_o(n) = 2n + 1$ , mapping  $\mathcal{E}$  to  $\mathbb{N}$  and  $\mathcal{O}$  to  $\mathbb{N}$ , respectively, are both bijective. Finally, we note that  $\mathcal{O} \cup \mathcal{E} = \mathbb{N}$ .

Since  $A$  and  $B$  are both countably infinite, i.e.  $|A| = |B| = |\mathbb{N}|$ , then  $\exists$  bijections  $f, g$  such that

$$f : A \rightarrow \mathbb{N}, \text{ and } g : B \rightarrow \mathbb{N}. \quad (9)$$

Then  $\forall a \in A$ ,  $2f(a)$  is even, and  $\forall b \in B$ ,  $2g(b) + 1$  is odd. So we define the even and odd sets

$$\mathcal{E} := \{2f(a) \mid a \in A\}, \text{ and } \mathcal{O} := \{2g(b) + 1 \mid b \in B\}. \quad (10)$$

Then we construct the function  $h$  defined via

$$h(x) = \begin{cases} 2f(x) & \text{if } x \in A \\ 2g(x) + 1 & \text{if } x \in B. \end{cases} \quad (11)$$

First, we show that  $h$  is injective.

Case 1:  $x, y \in A$ . Then

$$h(x) = h(y) \quad (12)$$

$$\implies 2f(x) = 2f(y) \quad (13)$$

$$\implies f(x) = f(y) \quad (14)$$

$$\implies x = y, \quad (15)$$

Since  $f$  is injective.

Case 2:  $x, y \in B$ . Then

$$h(x) = h(y) \tag{16}$$

$$\implies 2g(x) + 1 = 2g(y) + 1 \tag{17}$$

$$\implies g(x) = g(y) \tag{18}$$

$$\implies x = y, \tag{19}$$

since  $g$  is injective.

Case 3:  $x \in A, y \in B$  WOLOG, and  $h(x) = h(y)$ . This case is vacuous because  $\mathcal{E}$  and  $\mathcal{O}$  are disjoint, whereas  $2f(x) \in \mathcal{E}$  while  $2g(y) + 1 \in \mathcal{O}$ .

Thus, according to cases 1 and 2,  $h$  is injective.

Now we show that  $h$  is surjective. Let  $n \in \mathbb{N}$ .

Case 1:  $n$  is even. Then by surjectivity of  $f$ ,  $\exists a \in A$  such that  $f(a) = \frac{n}{2}$ , i.e.  $h(a) = n$ .

Case 2:  $n$  is odd. Then by surjectivity of  $g$ ,  $\exists b \in B$  such that  $g(b) = \frac{n-1}{2}$ , i.e.  $h(b) = n$ .

Thus,  $h$  is surjective as well.

Therefore we conclude that  $h$  is a bijection, hence  $|A \cup B| = |\mathbb{N}|$ .  $\square$