CJC-1295 + IPAMORELIN

Reconstitution, Storage & Protocol Reference

RESEARCH REFERENCE This document organizes the supplied protocol information for educational and research-reference purposes. The protocols below are not derived from published dose-finding studies and are not medical advice.

Reconstitution

1. Remove the caps from both the peptide vial and the bacteriostatic water vial.
2. Wipe both rubber stoppers with an alcohol swab.
3. Draw 2 mL of bacteriostatic water into a syringe.
4. Inject the water slowly against the inside wall of the peptide vial; do not spray it directly onto the powder.
5. Allow the powder to dissolve naturally. Do not shake the vial.
6. After a few minutes, gently swirl if needed.

Storage

Condition	Storage guidance
Before reconstitution	Room temperature or refrigerated
After reconstitution	Refrigerated at 36–46°F (2–8°C)
Important	Do not freeze after reconstitution
Suggested use window	Within 4–6 weeks of reconstitution

Practical Protocols

The following protocols are not derived from published dose-finding studies. They represent clinical-practice patterns and community consensus.

Standard Protocol

Item	Reference range / timing
CJC-1295 (no DAC)	100–300 mcg per day
Ipamorelin	100–300 mcg per day
Timing	Once daily, fasted (at least 2 hours after eating)
Preferred timing	Before bed on an empty stomach
Schedule	Daily
Cycle length	12–16 weeks, followed by 4 weeks off

Reference by Body Weight

Body weight	Amount of each peptide
Under 150 lb	100–150 mcg
150–200 lb	200 mcg
200–250 lb	250–300 mcg
Over 250 lb	300 mcg

Split Dosing (Advanced)

Timing	Amount of each peptide
Morning (fasted)	100–150 mcg
Before bed	100–150 mcg

RATIONALE This approach is intended to create two separate growth-hormone pulses per day.

Standard Blend Vial: Reconstitution & Units

Specification	Value
Vial size	10 mg total: 5 mg CJC-1295 (no DAC) + 5 mg Ipamorelin
Bacteriostatic water	2 mL
Concentration	250 mcg of each peptide per 10 units on a U-100 insulin syringe
Approximate vial duration	20 daily administrations at 250 mcg of each peptide

CALCULATION CHECK Each peptide: 5 mg ÷ 2 mL = 2.5 mg/mL. On a U-100 syringe, 10 units = 0.10 mL, which provides 250 mcg of CJC-1295 and 250 mcg of Ipamorelin.

Important Notes

- Confirm that the CJC-1295 component is the no-DAC form; DAC and no-DAC products have different pharmacologic profiles.
- Use sterile handling practices and label the vial with the reconstitution date.
- Discard the solution if it appears contaminated, unusually cloudy, or contains unexpected particles.
- Products sold for laboratory research should be handled only within their stated intended-use restrictions.