
GLP3 “Retatrutide”

10mg Vial

2ml BAC Water	1mg Dose	2mg Dose	2.5mg Dose
	20 units	40 units	50 units

Recommendation for 10mg vial: Add 2ml of BAC water.

20mg Vial

2ml BAC Water	1mg Dose	2mg Dose	4mg Dose
	10 units	20 units	40 units

Recommendation for 20mg vial: Add 2ml of BAC water.

Step-by-Step Reconstitution Process

Gather supplies: BAC water, alcohol swabs, insulin syringe, vial

Removing plastic cap. Carefully pop off the plastic cap, do not remove metal flange.

Clean both vials: Wipe the rubber stoppers with alcohol swabs

Draw BAC water: Pull your chosen amount into the syringe

Inject slowly: Add the water to the peptide vial, aiming it at the glass wall—not directly onto the powder

Swirl gently: Roll the vial between your palms. Never shake.

Wait: Let it sit 5-10 minutes. Solution should be crystal clear.

Store: Refrigerate immediately. Good for 28-30 days once reconstituted.

If it's cloudy or won't dissolve: Let it sit longer. Try gentle warming (under your arm or in lukewarm water). If still cloudy after 30 minutes, contact your source.

Starting Doses: How to Titrate Safely

Retatrutide is potent. Starting too high leads to severe GI side effects. Here's the standard titration protocol:

Conservative Titration (Recommended)

Week	Weekly Dose	Notes
1	0.5mg	Assess tolerance
2	1.0mg	Most notice appetite changes here
3	1.5mg	Solid fat loss begins
4	2.0mg	Sweet spot for many
5+	2.0-2.5mg	Adjust based on response

What to Expect at Each Stage

0.5-1mg: Mild appetite suppression, minimal side effects. You might wonder if it's working—this is the adaptation phase.

1.5-2mg: Noticeable appetite reduction, early fat loss visible on the scale. Some GI adjustment (mild nausea, fullness) is normal and passes within 2-4 weeks.

2.5mg+: Strong effects. Only increase if lower doses plateau and you tolerate well. Higher isn't always better.

Injection Basics

Where: Subcutaneous—belly fat (2 inches from navel), thigh, or upper arm

When: Same day each week, any time of day. Morning works well.

Rotate sites: Don't inject the same spot twice in a row

Needle: 29-31 gauge insulin syringe, 1/2 inch needle

Common Beginner Mistakes to Avoid

Starting too high: 2mg+ on week 1 = miserable GI sides. Trust the titration.

Shaking the vial: Denatures the peptide. Always swirl.

Not refrigerating: Reconstituted peptide degrades at room temperature.

Mixing vials: Keep BAC water separate. Don't combine different peptides.

Impatience: Retatrutide's glucagon mechanism takes 4-6 weeks to fully activate. Give it time.
