

Continuing the provious question, we got Now how long the epinde bill last will also play a role in comparing. The longer the episode last, the never wand we will get, as (1-r) decrease universes with wirear in Tas Over < 1 7 T=2 2 G + C.(1-r) = G+ ((1+r)

 $\frac{Q_{S} O My'}{V_{*}(S)} = \max_{Q \in ACS} Q_{\Pi_{*}} (S_{C}a)^{2} - q_{C}ACS)$

In lince Y(S, 95') is an expectation be need to figure out the order sented for each probability of finding a can or how finding a can.

P = find a can q =

Nearer = P = P × 1 + (1-P).x0 mobability y finding a cen is B. Trearch. Given p(s', x | S(x) = d x renauch. PCS, 7 S > find-ean, a) = drumanel p(5(5(4) = 2 & So., p(s, v > 0 | s, a) =+ p(s, x > 1 | s, a) > p(s, s, a) > p(5,700 | S(0) = 2 - 2 march.

| Dorng | this for o | Thier. | 1. | Account to |
|----------|--|--|-----|-----------------------|
| | / | | | |
| 2 | ٩ | . 5' | 8 | P(5,815,9) |
| H | 2 | 1 h | | 2 ruard |
| h | 2 | h. | , 0 | d-d means |
| h | 5 | 人 | | (1-2) I search |
| | 3 | h | -3 | 1 – β |
| | 2 | 1 | | Bxrrance |
| <u> </u> | 5 | | ٥ | B-Bxruarch |
| • h | 5 | 1 | | (1-d) - (1-d) rreach |
| e h | LU | h | 1 | to Ewait |
| | W | sl | | Arsen I want |
| | ω | 1 | 0 | B = B * France 1- You |
| | γ | h | 0 | \$2000 ar 1 |
| o h | W | 人 | 0 | 1- ruait |
| | | | | 1 |
| | | | | |
| | | | | |
| | | | | |
| • | | | , | |
| 9 | | | | |
| | , | | | |
| • | | | | |
| | depression of the second secon | And the high representative about the production of the land and the second and t | | |
| | | | | |