Manual of KRASS_VS_assistant

open-source software

Appearance of the main window of the KRASS_VS_assistant web application:

Step 1. *Draw molecule or select input molecular files.*

If you want to draw the structure of a chemical compound, you can use the capabilities of the built-in chemical editor Ketcher (https://github.com/epam/ketcher). After creating the structure of a chemical compound, it can be controlled using two buttons: 1)"reset" - deleting the structure to create a new one 2) "Apply" - transferring the structure of the compound for further analysis, forecasting to step 2.



After clicking on the 'Apply' button, make sure that the structure has been created. If the structure is successfully created, its smiles will be displayed under the chemical editor window



If you choose smiles, please, directly paste the SMILES representation of the desired chemical structure and press Ctrl+Enter. If the entered chemical structure is correct, the application will generate a 2D image of the studied compound. DON'T FORGET TO CLICK THE "APPLY" BUTTON



If the entered structure is incorrect, the application reports an error.



If you choose a file *sdf or *csv, that may contain a different number of chemical structures, please specify the path to this file on your computer's hard drive. In this case, you need to click the "Browse files" button. It is important to note that if you choose a file with the *csv extension, the file should contain a column with the name "SMILES"



If incorrect structures are detected in the file *sdf or *csv, the corresponding information will appear in the section "CHEMICAL STRUCTURE VALIDATION AND STANDARDIZATION"



No.	No. failed molecule in original set	SMILES of wrong structure:
1	80	COC1=C/C2=C(OC3=CC=C(NC(=O)C4=C(=O)C=CN(C5=CC=CC=C5)=N4)C=C3
2	88	O=C(/C=C/C1=CC=C(CCNC(=O)C2=CC(C/C3=N/N=C(=O)C4=CC=CC=C43)=CC
3	90	COC1=C\C(OC)=C2\C(=O)=NC(C3=CC(C)=C(OCCCCC(=O)NO)C(C)=C3)=N\C2
4	92	COC1=C/C2=C(OC3=CC=C(NC(=O)C4=C(=O)C(C)=CN(C5=CC=CC=C5)=N4)C=
5	93	CCC1=CN(C2=CC=CC=C2)=NC(C(=O)NC2=CC=C(O/C3=C/C=N\C4=CC(OCCCC
6	153	O=C(NO)C1=CC=C(CN2CCN(C(=O)C3=CC(C/C4=N/N=C(=O)C5=CC=CC=C54)
7	158	O=C(CC1=CC=C(CN2CCN(C(=O)C3=CC(C/C4=N/N=C(=O)C5=CC=CC=C54)=C
8	193	O=C(/C=C/C1=CC=C(CN2CCN(C(=O)C3=CC(C/C4=N/N=C(=O)C5=CC=CC=C54
9	203	COC1=C\C(OC)=C2\C(=O)=NC(C3=CC(C)=C(OCC4=CC=C(/C=C/C(=O)NO)C=C

Step 2. *Select prediction of KRASS inhibitor activity, ADMET or substructural search for preferred or undesirable fragments.*

Step 2. Select prediction of KRASS inhibitor activity, ADMET or substructural search for preferred or undesirable fragments

KRASS

Run predictions!

© Ligand Pro, 2025

Click on the “Run predictions!” button for prediction.

Step 3. *Prediction results*.

The form of presentation of the results depends on the type of descriptors selected, as well as the format of the input chemical data. For example, when selecting SMILES, the results will be displayed for a single molecule. When displaying the results on the screen, it is taken into account whether there are experimental values of activity and toxicity for the studied compound.

Step 2. Select prediction of KRAS inhibitor activity or substructural search for preferred or undesirable fragments

KRAS

Run predictions!

The final table ‘Prediction results’ contains the following columns:

1) ***SMILES*** - the chemical structure is displayed in the SMILES notation

2) ***Predicted value pIC50*** - the predicted value of the activity to inhibit the KRASS enzyme, expressed in pIC_{50}, where pIC_{50} is the negative logarithm of IC_{50} in molar concentration. If experimental data is available in the ChEMBL database, the label "see experimental value" is displayed in this cell.

3) ***Applicability domain_KRASS*** - compliance of the chemical compound with Applicability domain. If experimental data is available in the ChEMBL database, the label "-" is displayed in this cell.

4) ***Experimental value, pIC50***- experimental data presented in the ChEMBL database. Where pIC_{50} is the negative logarithm of IC_{50} in molar concentration. If there is more than one value in the database, then the average value is given in the cell.

5) ***STD*** - ehe cell indicates the standard deviation of the experimental activity values presented in the ChEMBL database.

6) ***chemble_ID*** - identifier from the ChEMBL chemical database of molecule

If you choose a file *sdf or *csv, the prediction results for correct chemical structures are displayed in a table that can be downloaded.

Original data: 11 molecules

Failed data: 0 molecules

Kept data: 11 molecules

Run predictions!

	SMILES	Predicted value, pIC50	Ap
2	CCOc1cc(NC(=O)CCCCCCC(=O)Nc2cc(-c3ccccc3)ccc2N)cc2c1OCC([N+](=O)[O-])=C2	see experimental value	-
3	[N-]=[N+]=NCCC(=O)Nc1ccc(C(=O)Nc2cc(N=[N+]=[N-])ccc2N)cc1	see experimental value	-
4	[N-]=[N+]=Nc1cccc(COC(=O)NCc2ccc(C(=O)Nc3cc(-c4ccccc4)ccc3N)cc2)c1	see experimental value	-
5	[N-]=[N+]=NCc1cc(N=[N+]=[N-])cc(C(=O)Nc2cc(-c3cccs3)ccc2N)c1	see experimental value	-
6	C#CCOc1ccc(C=NNC(=O)c2cccc(C(=O)NO)c2)cc1	see experimental value	-
7	O=C(NO)c1cnc(NC2(c3ccccc3)CCC(F)(F)CC2)nc1	see experimental value	-
8	CN1CCc2c(c3ccccc3n2Cc2ccc(C(=O)NO)cc2)C1	see experimental value	-
9	CCCNNC(=O)c1ccc(CNC(=O)C(Cc2c[nH]c3ccccc23)NC(=O)c2ccc(OC)cc2)cc1	see experimental value	-
10	COc1ccc(C(=O)c2ccc3c(ccn3Cc3ccc(C=CC(=O)NO)cc3)c2)cc1	see experimental value	-
11	CC(C=CC(=O)NO)=CC(C)C(=O)c1ccc(N(C)C)cc1	see experimental value	-

Download results of prediction as CSV

The KRASS_VS_assistan application can be used to evaluate the compliance of the compound to be studied with Muegge's rules, and the result is displayed both in tabular numerical form and graphically, in the form of the so-called Bioavailability Radar, which can be used to visually evaluate the compliance of the calculated values of lipophilicity (log P), molecular weight (MW), and the number of hydrogen bond donors (HBD) and hydrogen bond acceptors (HBA) and other parameters with the limits determined by Muegge's rules.

The Drug-likeness Radar: compliance with Muegge's rules



	Values for the test substance	Reference values of Muegge rules
Molecular weight(MW), Da	499.484	200-600
Octanol-water coefficient(LogP)	3.8765	≤5
Number of hydrogen bond donors (HBD)	2	≤5
Number of hydrogen bond acceptors(HBAs)	7	≤10
Number of rotatable bonds	4	≤15
Topological polar surface area (TPSA), Å²	99.52	≤150
Number of rings	5	≤7

In addition, the KRASS_VS_assistant web application integrates detection of such widely used in medicinal chemistry structural alerts as Brenk filters and PAINS when analyzing single structures. Brenk filters which consists in a list of 105 fragments identified by Brenk et al. [Brenk, R. et al. Lessons learnt from assembling screening libraries for drug discovery for neglected diseases. ChemMedChem 3, 435–444 (2008).] to be putatively toxic, chemically reactive, metabolically unstable or to bear properties responsible for poor pharmacokinetics. PAINS (for pan assay interference compounds, a.k.a. frequent hitters or promiscuous compounds) are molecules containing substructures showing potent response in assays irrespective of the protein target. Such fragments, yielding false positive biological output, have been identified by Baell et al. [Baell, J. B. & Holloway, G. A. New substructure filters for removal of pan assay interference compounds (PAINS) from screening libraries and for their exclusion in bioassays. J. Med. Chem. 53, 2719–2740 (2010). KRASS_VS_assistant returns warnings if such moieties are found in the molecule under evaluation.

The found Brenk filter: Michael-acceptor



The PAINS are not found in the molecule.