

Lecture Notes on **General Relativity**

Owen Lin
owen2006lin@gmail.com

These notes cover general relativity at the undergraduate level. The sources used are:

- Schutz, *A First Course in General Relativity*
- Carroll, *Spacetime and Geometry*

Potential sources to consult in the future:

- Hartle, *Gravity: An Introduction to Einstein's General Relativity*
- Misner, Thorne, Wheeler, *Gravitation*
- Weinberg, *Gravitation and Cosmology*
- Wald, *General Relativity*

Contents

1	Preliminaries	3
1.1	Coordinate Transformations	3
1.2	Tensor Analysis	4
2	Gravitation	6
2.1	The Newtonian Limit	6
2.2	Stress-Energy Tensor	6
2.3	Gravitational Waves	7
3	Black Holes	8
3.1	Schwarzschild Metric	8
3.2	Coordinates Systems	8
3.3	Orbits	9
3.4	Massless Particle	10
4	Cosmology	11
4.1	The Present Universe	11
4.2	History of the Universe	12
4.3	Singularity	13
4.4	The FLRW Universe	13
4.5	The CMB	13
4.6	The Early Universe	13

1 Preliminaries

1.1 Coordinate Transformations

First, we'll briefly recap special relativity using index notation.

- Let's define a four vector x^μ where x^0 is the temporal component and x^i ($i \neq 0$) are the spacial coordinates. In these notes, we'll write the representation of a vector in coordinate transformation as $x^{\mu'}$ (each source has their own convention).
- The matrix to transform the vector x^ν into the new primed coordinates is

$$x^{\mu'} = \Lambda^{\mu'}_{\nu} x^\nu$$

An object that transforms like vector components is said to transform **contravariantly**. Note we are using the Einstein convention where summations over repeated indices drop the Σ .

- A Lorentz Transformation is one that keeps the spacetime interval constant and fix the origin so this requires $\eta = \Lambda^T \eta \Lambda$. In index notation, this is:

$$\eta_{\alpha\beta} = \Lambda^{\mu'}_{\alpha} \Lambda^{\nu'}_{\beta} \eta_{\mu'\nu'}$$

Note that when writing the equation using matrices, the order is important because of the limitations of matrix multiplication. However, the order is irrelevant in index notation. These matrices are **Lorentz Transformations** and are elements of the **Lorentz Group**.

- A brief aside, these matrices may seem very similar to the rotation group SO(3). The Orthogonal Group is all 3x3 matrices that satisfy $\mathbf{R}^T \mathbf{R} = \mathbf{1}$ (SO(3) then are all special matrices in O(3), ie determinant 1). The difference between O(3) and the Lorentz group is that instead of the identity matrix, it is equal to the metric η . The identity matrix is simply the metric in **Euclidean Space**, and η is similarly the metric for a **Lorentzian Space**.
- Lorentz transformations come in two forms: rotations and boosts. Below is an example of a boost in the x direction:

$$\Lambda^{\mu'}_{\nu'} = \begin{pmatrix} \cosh \phi & -\sinh \phi & 0 & 0 \\ -\sinh \phi & \cosh \phi & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

The transformed quantity for t' and x' are then $t' = t \cosh \phi - x \sinh \phi$ and $x' = -t \sinh \phi + x \cosh \phi$. The velocity follows as $v = \tanh \phi$, and we can then rewrite in more familiar form

$$t' = \gamma(t - vx)$$

$$x' = \gamma(x - vt)$$

where $\gamma = 1/\sqrt{1 - v^2}$. From here, we can proceed and obtain time dilation, length contraction, etc.

- Another transformation that preserves the spacetime interval is a pure translation:
 $x^\mu \rightarrow x^{\mu'} = (x^\mu + a^\mu)$ The set of all translations form an Abelian group but generally Lorentz transforms do not commute so the Lorentz Group is nonabelian. The set of all Lorentz transforms and translations form a larger nonabelian group: the **Poincaré Group**.

Now we'll take a look at how vectors and their components transform under a change of coordinates.

- Suppose in an arbitrary vector space we have a basis of vectors $\hat{e}_\mu$. Then any arbitrary vector can be written as $A = A^\mu \hat{e}_\mu$ where A^μ are the components of the vector. Often we will just call A^μ the vector for shorthand.
- An example is the tangent vector $V^\mu = \frac{dx^\mu}{d\lambda}$. Under a lorentz transformation the components change by $\frac{dx^{\mu'}}{dx^\mu}$ by the chain rule so the components transform as $V^\mu \rightarrow V^{\mu'} = \Lambda^{\mu'}_\nu V^\nu$ but the vector itself is invariant under transformation. So we must have $V^\mu \hat{e}_\mu = V^{\nu'} \hat{e}_{\nu'} = \Lambda^{\nu'}_\mu V^\mu \hat{e}_{\nu'}$. But this must hold for any vector no matter the components so we must have $\hat{e}_\mu = \Lambda^{\nu'}_\mu \hat{e}_{\nu'}$. This means basis vectors transform opposite to their components.
- Note that applying a transformation and then its inverse should result in the identity. In index notation, this means $\Lambda^\mu_{\nu'} \Lambda^{\nu'}_\rho = \delta^\mu_\rho$
- We can define a vector space dual to our previous vector space, which is the set of all linear maps from the original vector space to $\mathbb{R}$. We can write a set of basis dual vectors as $\hat{\theta}^\nu(\hat{e}_\mu) = \delta^\nu_\mu$. Then every dual vector can be written in component form $\omega = \omega_\mu \hat{\theta}^\mu$. (we also refer to dual vectors as "1-forms").
- The components of dual vectors transform like basis vectors (that is, inverse to vector components)

$$\omega_{\mu'} = \Lambda^\nu_{\mu'} \omega_\nu$$

An object that transforms like this is said to transform **covariantly**. Meanwhile, dual basis vectors transform like vector components $\hat{\theta}^{\rho'} = \Lambda^{\rho'}_\sigma \hat{\theta}^\sigma$. The key takeaway is that the location of the index tells us how the object transforms. Upper index transform like vector components, while lower transforms inverse to that. Accordingly, scalars are invariant under transformation.

- An example is the gradient, with components $\frac{\partial}{\partial x^\mu}$. Under transformation, this becomes $\frac{\partial}{\partial x^{\mu'}} = \frac{\partial x^\mu}{\partial x^{\mu'}} \frac{\partial}{\partial x^\mu}$ by the chain rule, but this is equivalent to just $\Lambda^\mu_{\mu'} \frac{\partial}{\partial x^\mu}$. Since the gradient is a dual vector, we tend to write it with lower index as ∂_μ .

1.2 Tensor Analysis

A tensor is a multilinear map of dual vectors and vectors to $\mathbb{R}$. For example, $T(a\omega, cV) = acT(\omega, V)$. A tensor that takes in k vectors and l dual vectors is a type (k, l) tensor. It's easy to see that scalars are $(0, 0)$ tensors and vectors and dual vectors are $(1, 0)$ and $(0, 1)$ tensors accordingly. Some important facts about tensors:

- Suppose T is a (k, l) tensor and S is a (m, n) tensor, Then we define the tensor product

$$T \otimes S(\omega^1, \dots, \omega^{k+m}, V^1, \dots, V^{l+n}) = \\ T(\omega^1, \dots, \omega^k, V^1, \dots, V^l) \times S(\omega^{k+1}, \dots, \omega^{k+m}, V^{l+1}, \dots, V^{l+n})$$

In general, the tensor product does not commute.

- In index notation, we can write T as $T^{\mu_1 \dots \mu_k}_{\nu_1 \dots \nu_l}$. Then in index notation, we can write it as $T^{\mu_1 \dots \mu_k}_{\nu_1 \dots \nu_l}$. The basis for the vector space T resides in can be expressed with basis $\hat{e}_{\mu_1} \otimes \dots \otimes \hat{e}_{\mu_k} \otimes \hat{\theta}^{\nu_1} \otimes \dots \otimes \hat{\theta}^{\nu_l}$. The components can be found by acting the tensor T on the basis vectors and dual vectors.
- T transforms k times contravariantly and l times covariantly:

$$T^{\mu'_1 \dots \mu'_k}_{\nu'_1 \dots \nu'_l} = \Lambda^{\mu'_1}_{\mu_1} \dots \Lambda^{\mu'_k}_{\mu_k} \Lambda^{\nu_1}_{\nu'_1} \dots \Lambda^{\nu_l}_{\nu'_l} T^{\mu_1 \dots \mu_k}_{\nu_1 \dots \nu_l}.$$

- We can contract a vector by summing over a repeated upper and lower index. With this, we can use the metric to raise and lower indices: $\eta_{\alpha\beta} T^\alpha = T_\beta$ and $\eta^{\alpha\beta} T_\alpha = T^\beta$. This is how you would convert a vector to its dual form and vice versa. In euclidean space, the metric is its own inverse so the gradient is actually both a vector and 1-form at the same time.
- A **symmetric** tensor is one where any change of indices leaves the tensor unchanged. For example, if $S_{\mu\nu\rho} = S_{\nu\mu\rho}$ then we say S is symmetric in the first two indices. A tensor is **antisymmetric** if it changes sign when indices are changed. $A_{\mu\nu\rho} = -A_{\nu\mu\rho}$ is antisymmetric in its first two indices.
- Symmetric and antisymmetric tensors are very useful in that the contraction of a symmetric and antisymmetric tensor along the corresponding indices is always 0. The symmetric part is $T_{(\mu\nu)} = \frac{1}{2}(T_{\mu\nu} + T_{\nu\mu})$ and the antisymmetric part is $T_{[\mu\nu]} = \frac{1}{2}(T_{\mu\nu} - T_{\nu\mu})$. We sometimes want to exclude indices from being symmetrized or antisymmetrized so we use the vertical bars. For example, to write the antisymmetrized $R_{\alpha\beta\gamma\delta}$ in only the first and third indices we use $R_{[\alpha|\beta|\gamma]\delta} = \frac{1}{2}(R_{\alpha\beta\gamma\delta} - R_{\gamma\beta\alpha\delta})$
- For a $(1,1)$ tensor, we write the trace by leaving off the indices (just the contraction along the repeated index)

$$X = X^\mu{}_\mu$$

We can similarly define the trace of a $(0,2)$ or $(2,0)$ tensor by raising/lowering into a $(1,1)$ tensor and then contracting. Note that in this case, the trace is not just the naive sum over the diagonal; the components change when raising/lowering the index. The reason behind this is because the sum over the diagonal is not a true scalar; ie it's not invariant under coordinate transform.

Example. The metric tensor $\eta_{\mu\nu}$ is a $(0,2)$ tensor that takes in two vectors and outputs a scalar. The metric acting on two vectors returns the inner product

$$\eta(V, W) = \eta_{\mu\nu} V^\mu W^\nu = V \cdot W$$

Properties of the inner product should be familiar. Two vectors are orthogonal if their inner product is 0. Note that because the inner product is a scalar, it's invariant under Lorentz transformations. This means that the cartesian basis is always orthogonal under Lorentz transformation.

Additionally, the norm, or "length" of a vector is defined by its inner product with itself $\eta_{\mu\nu} V^\mu V^\nu$. Unlike in Euclidean space, there is no restriction for the norm to be strictly positive. A positive norm is **spacelike**, a negative norm is **timelike** and 0 is **lightlike** or null.

Example. The Kronecker delta $\delta^\mu{}_\nu$ is a $(1,1)$ tensor. It inputs a vector and a dual vector and outputs a scalar, but can also be thought of as a map from dual vectors to dual vectors and vectors

to vectors. In this framing, the Kronecker delta is just the identity map. We can define the inverse metric $\eta^{\mu\nu}$ as satisfying the property

$$\eta^{\mu\nu}\eta_{\nu\rho} = \eta^{\nu\mu}_{\rho\nu} = \delta^{\mu}_{\rho}$$

2 Gravitation

2.1 The Newtonian Limit

2.2 Stress-Energy Tensor

We'll be working with fluids in this section. The definition of fluids is not precise so we'll just say a fluid has friction force negligible in comparison to the pressure.

- Fluid particles travelling with four-velocity $\mathbf{u}^{\alpha}$. We can choose to move to inertial frame where $\mathbf{u}^{\mu} = (1, 0, 0, 0)$. This is called the **comoving frame**.
- The **number current** is $\mathbf{j}^{\mu} = n\mathbf{u}^{\mu} = n(u^0, \mathbf{u}^i)$ where n is the number density (particles per unit volume). The components of the number current is the number of particles travelling through unit area per unit time which is the particle flux.
- The rest **mass density** is $\mu_0 = m_0 n$ where m_0 is the rest mass of a single particle. If there is internal energy, then we have $\mu = \mu_0(1 + \delta)$ where δ is the internal energy fraction.
- The **momentum density** is $\rho^i = m_0 n \mathbf{u}^i$. A nice exercise is to show this is equivalent to the energy flux in special relativity.

We'll now introduce the **stress-energy tensor** $\mathbf{T}^{ij}$.

- In newtonian mechanics, we have the stress tensor $\mathbf{t}^{ij}$. This represents shear: the i th component of force per unit area (pressure) exerted across a surface normal to the j th direction.
- For a fluid at rest, the force exerted across a surface is always along the normal and the same in all directions, equivalent to pressure $\mathbf{t}^{ij} = p\delta^{ij}$
- For a perfect fluid, $\mathbf{T}^{\alpha\beta} = (\rho + p)u^{\alpha}u^{\beta} + p\mathbf{g}^{\alpha\beta}$. Careful, ρ is the energy density! Meanwhile, $\mathbf{T}^{\alpha}_{\beta} = (\rho + p)u^{\alpha}u_{\beta} + p\delta^{\alpha}_{\beta}$

Here, we'll read off some of the physical interpretations of the components of the stress-energy tensor:

- $\mathbf{T}^{\alpha}_{\beta}u^{\beta} = -\rho u^{\alpha}$. Physically, this quantity's spacial components are the energy flux seen comoving with the fluid. The temporal component $\mathbf{T}^0_{\beta}u^{\beta} = -\rho$ is just the rest mass energy density.
- $\mathbf{T}^0_i$ is the flow of the i th component of momentum in the temporal ($i = 0$) direction. This is the energy flux. Consequently, $\mathbf{T}^i_0$ is the energy crossing through the i th surface, so the energy flux. Thus, these two quantities are equivalent. In the comoving frame, $\mathbf{T}^0_i = \mathbf{T}^i_0 = 0$ because the fluid is at rest.

Now, we'll see some results that arise from the Newtonian Limit. Firstly, we assume that the fluid is travelling with low velocity $|u^{\alpha}| \ll 1$. Additionally, we have low pressure $\frac{p}{\rho} \ll 1$ and the speed is non-relativistic $u^i \approx v^i$.

- Firstly, conservation of energy-momentum means that the stress-energy tensor is divergence-free. This gives us $\nabla \cdot \mathbf{T} = 0$ or equivalently $\mathbf{T}_{;\beta}^{\alpha\beta} = 0$. (why???)
- Next, reading off the stress-energy tensor components, we have:
 - $\mathbf{T}^{00} \approx \rho$
 - $\mathbf{T}^{j0} \approx \rho v^j$
 - $\mathbf{T}^{jk} \approx \rho v^j v^k + p \delta^{jk}$
- The continuity equation $\frac{\partial \rho}{\partial t} + (\rho v^k)_{;k} = 0$ effectively states conservation of mass. The rate of change of density plus the net mass flow out is 0. Combining with the divergence free condition, we obtain $\frac{\partial v^j}{\partial t} + (v \cdot \nabla)v = -\frac{1}{\rho} \nabla p$. This is the **Euler Equation**.
- The equations of state relate the pressure and the energy density for a fluid. There are several possible cases:
 - Vacuum : $p = \rho = 0$ (Inflationary universe)
 - Dust : $p = 0$ (FLRW universe)
 - Fluid : $p = \gamma \rho$
 - Polytopic : $p = n \rho^{(1+\frac{1}{n})}$ where n is the polytopic index.

2.3 Gravitational Waves

Recall the form of the metric in the linearized theory $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$.

- The einstein equations give us $2(R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R) = 16\pi GT_{\alpha\beta}$
- Introducing the trace-reversed metric perturbation, $\bar{h}_{\alpha\beta} \equiv h_{\alpha\beta} - \frac{1}{2}h$ and expanding the einstein equation, we can simplify this to $\square \bar{h}_{\alpha\beta} = \bar{h}_{\alpha\beta,\mu}^{\mu} = -16\pi GT_{\alpha\beta}$. The box is the d'Alembert operator and is the equivalent of the laplacian in spacetime. This nicely gives us a wave equation for $\bar{h}$.
- The solution for this wave equation is $\bar{h}_{\mu\nu} = \text{Re}(A_{\mu\nu} e^{ik_{\alpha}x^{\alpha}})$ where $A_{\mu\nu}$ gives the amplitude and k_{α} is the wavevector ($\omega, \mathbf{k}$). From the d'Alembert operator, we have $k_{\alpha}k^{\alpha} = 0$, or k is a null vector. This means gravitational waves propagate at the speed of light.
- We can choose a Lorentz Gauge to simplify this further (what does this mean?) with $\bar{h}_{\mu\nu}^{\nu} = 0$. This gives us $A_{\mu\alpha}k^{\alpha} = 0$ meaning these are transverse waves (why?)
- Starting with the Symmetric 4x4 tensor, we have 10 independent coordinates. Being tranverse and traceless takes away 4 degrees of freedom (why?) and the gauge condition takes away 4 more so we are left with 2 polarizations. Imposing all conditions gives us the matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & b & 0 \\ 0 & b & -a & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

- Both linear and circular polarization like light (This section needs work)

Now we'll derive the metric from a weakly gravitating source. Suppose we have a static point mass M located at point $r = 0$ at all times in the weak field limit. In the body's frame we have $|h_{\mu\nu}| \ll 1$.

- Because the body is stationary, the stress energy tensor has components $T_{00} = \delta(x)M$, and all other components are 0.
- From the einstein equation and poisson equations, we have $\bar{h}_{00} = \frac{4M}{r}$ and all other components are 0.
- Converting from $\bar{h}$ back to h and substituting into the equation for the metric $ds^2 = (\eta_{\mu\nu} + h_{\mu\nu})dx^\mu dx^\nu$ we obtain the metric:

$$ds^2 = 1 \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) (dx^2 + dy^2 + dz^2)$$

3 Black Holes

Rev. John Mitchell found in 1783 that an object with $500\odot$ but of same density will have its escape velocity to be greater than the speed of light. Then "all light emitted from such a body would be made to return towards it."

- Every mass has a radius such that if the mass is small enough than this radius, it will collapse into a black hole. This is the **Schwarzschild Radius**. $R_s = \frac{2GM}{c^2}$

3.1 Schwarzschild Metric

The Schwarzschild Metric has two forms:

- The timelike form $d\tau^2 = (1 - \frac{2M}{r})dt^2 - \frac{dr^2}{(1 - \frac{2M}{r})} - r^2(d\theta^2 + \sin^2\theta d\phi^2)$
- The spacelike form $d\tau^2 = (1 - \frac{2M}{r})dt^2 - \frac{dr^2}{(1 - \frac{2M}{r})} - r^2d\Omega^2$
- We can see the metric reduces to flat spacetime in the case where $r \rightarrow \infty$ and $m \rightarrow 0$
- Upon examining the temporal component of the metric, we that its less than that of the temporal component in flat spacetime, so this properly predicts gravitational redshift. Similarly, the spacial component is larger than that in flat spacetime, so there's spacetime curvature.

3.2 Coordinates Systems

These are three commonly used frames:

- Free-Float Frame: Frame free-floating in space, think unpowered spacecraft. Locally its flat minkowski space. For observers on shells, they see spacecraft infalling towards the black hole.
- Spherical-Shell Frame: Frame standing still on a shell of fixed radius (eg: Earth's frame). For small time lapses and regions, special relativity applies. Notice that inside a black hole, dr becomes time like while dt becomes spacelike. This means staying on a shell at constant radius is impossible, so this frame cannot be used to describe events inside a black hole.

- Schwarzschild Frame: Global book-keeping coordinates. Idea is that neither spherical shell frame nor free float frame can know all information so both send information to a Global frame that puts it together.

A common result from black holes is that from the perspective of someone outside of the black hole, we see an object travel into the black hole past the event horizon. But from the perspective of the falling object, it takes infinite time to reach the event horizon!

- We can calculate the Riemman tensor in the shell frame: $R_{trtr} = -\frac{2M}{r^3} = -R_{\theta\phi\theta\phi}$ and $R_{t\theta t\theta} = R_{t\phi t\phi} = \frac{M}{r^3} = -R_{r\theta r\theta} = -R_{r\phi r\phi}$
- Meanwhile in the free falling frame we have $R_{trtr} = -\frac{2M}{r^3}(1 - \frac{2M}{r})^{-1}$. This isn't singular at $r = 2M$ but it is at $r = 0$. The curvature invariant (also known as the Kretschmann scalar) $R = R_{trtr}R^{trtr} = \frac{48M^2}{r^6}$. This gives a measure of the strength of the gravitational field using curvature and this also blows up to infinity at $r = 0$.

3.3 Orbits

Because the Schwarzschild metric is time independent and spherically symmetric, we must have killing vectors $\xi_t = (1, 0, 0, 0)$ and $\xi_\phi = (0, 0, 0, 1)$. For any four-velocity u , $\xi \cdot u$ is constant. combining this with $p^\mu = mu^\mu$, we obtain $\frac{E}{m_0c^2}$ is constant along a path.

- Radial Infall: Let $d\tau$ be the proper time for an observer at distance r from the center of a black hole (shell frame). At $r \rightarrow \infty$, we have $\frac{E_\infty}{m_0c^2} = \frac{dt}{d\tau}$. But E_∞ is a constant. So any trajectory starting at rest from $r = \infty$ will have same energy per unit mass throughout. This gives us $d\tau^2 = (1 - \frac{2M}{r})^2 dt^2$.
- With this expression we can obtain the velocity recorded by a Schwarzschild observer: $\frac{dr}{dt} = -(1 - \frac{2M}{r})(\frac{2M}{r})^{\frac{1}{2}}$. At $r = 2M$, this goes to 0 so we see that the bookkeeper observes the particle slowing down as it approaches the event horizon, taking infinite time to reach there.
- If we instead moved to the frame of the shell observer, we can obtain from the Schwarzschild metric: $\frac{dr_{shell}}{dt_{shell}} = -(\frac{2M}{r})^{\frac{1}{2}}$. For an observer at the event horizon, they would see the object falling at the speed of light! The energy observed also depends on the distance from the center (while E_∞ is constant, E_{shell} is not).
- Meanwhile, for the free falling frame, we can actually compute the time it takes for the observer to reach $r = 0$. Using the chain rule, $\frac{dr}{d\tau} = \frac{dr}{dt} \frac{dt}{d\tau}$ we can obtain $\tau = \frac{4}{3}M$. In seconds this is $\tau_{second} = \frac{4}{3} \frac{M}{M_{odot}} (5\mu sec)$. Interestingly, the time scales linearly with mass, so for larger black holes it actually takes more time to fall! For reference, Sagittarius A, the supermassive black hole in the center of the galaxy, it takes roughly 1 minute to fall. Meanwhile, TON618, one of the largest black holes takes 11 days to fall to the center.

Now we'll go over orbits. In the case with $r > 20R_s$, the newtonian approximation is still accurate. For $r \leq 6M$, no free orbits are possible. The mass spirals in but still may exert enough energy to escape. Beyond $r \leq 2M$, no escape is possible. Inside the horizon, as mentioned before, space and time components are interchanged. Travelling to smaller r is as "inevitable" as outside a black hole we travel to greater t .

- Recall the conserved quantity from earlier $\xi_t \cdot u^t$. We call this the "energy per unit mass" and we'll just set this equal to $\frac{E}{m}$. The free fall case assumes this is one but this is not true generally. Write the angular momentum per unit mass as $\frac{L}{m} = r^2 \frac{d\phi}{d\tau}$. Using these, we have the equation of motion

$$\left(\frac{dr}{d\tau}\right)^2 = \left(\frac{E}{m}\right)^2 - \left(1 - \frac{2M}{r}\right) \left[1 + \frac{(L/m)^2}{r^2}\right] \equiv \left(\frac{E}{m}\right)^2 - \left(\frac{V}{m}\right)^2$$

. This gives us an effective potential of

$$\left(\frac{V(r)}{m}\right)^2 \equiv \left(1 - \frac{2M}{r}\right) \left[1 + \frac{(L/m)^2}{r^2}\right]$$

In the Newtonian case, we have an elliptical orbit. But for black holes, orbits tend to spend more time at smaller radii, the additional time near the inner edge of the orbit is the "**dwelt time**."

- Stable circular orbits at V_{min} and unstable orbits occur at V_{max} . Beyond V_{max} the orbit precesses because of the dwelt time. (this section needs work)

3.4 Massless Particle

We'll now cover orbits of massless particles. This is especially important for describing the behavior of light near a black hole. The procedure for massive particles won't work here for several reasons.

- The spacetime interval for massless particles is 0 (why?). Solving for the coordinate velocity of light in these coordinates gives us $r \frac{d\theta}{dt} = \pm(1 - \frac{2M}{r})^{\frac{1}{2}}$ tangentially and $\frac{dr}{dt} = \pm(1 - \frac{2M}{r})$ radially. This looks like it means the speed of light can even approach 0 but this actually isn't a problem because this interpretation is unphysical. Even if the Schwarzschild observer "sees" the speed of light as this, any shell observer still sees it at constant c.
- At $r = \infty$, the special relativity case we have $p = p_\infty$. The angular momentum L can be written using this momentum as $L = bp_\infty$ where b is known as the **impact parameter**. (Why?) Using the expression for P in special relativity, we can write $b = \frac{L/m}{E/m}$

- The impact parameter conveniently allows us to evaluate the speed even with 0 mass:

The radial velocity: $\left(\frac{dr}{dt}\right)^2 = (1 - \frac{2M}{r})^2 - (1 - \frac{2M}{r})^3 \frac{b_{photon}^2}{r^2}$

The tangential velocity: $r \frac{d\phi}{dt} = \frac{b_{photon}}{r} (1 - \frac{2M}{r})$

Thus the total speed is $v_{photon} = (1 - \frac{2M}{r}) [1 + \frac{2Mb^2}{r^3}]^{\frac{1}{2}}$.

With a similar technique, we can verify that in the shell coordinates, the speed of the particle is constant 1.

- The equivalent equation of motion in the shell frame is $\frac{1}{b^2} \left(\frac{dr_{shell}}{dt_{shell}}\right)^2 = \frac{1}{b^2} - \frac{(1 - \frac{2M}{r})}{r^2}$. $\frac{1}{b^2}$ plays the role of the energy squared while the second term is the equivalent of the effective potential.
- This potential has no stable orbits. A photon may enter a circular "knife edge" orbit before either escaping or plunging.

4 Cosmology

4.1 The Present Universe

In the last 100 years, there were many important observations about our current universe.

- Hubble discovered the universe is expanding in 1929. The speed at which galaxies recessed from us is proportional to their distance $v = Hr$ where H is the hubble constant.
- In 1965, Penzias and Wilson discovered the cosmic microwave background has temperature $T = 273^\circ K$. It is also highly isotropic and homogeneous.
- Matter density upper bounds at $10^{-29} g/cm^3$ and lower bound at $5 \times 10^{-31} g/cm^3$. The Baryon number Photon number ratio is $10^{-8} - 10^{-10}$
- From stars, galaxies and globular clusters, the Helium-4 mass fraction is 0.2-0.3, while Deutereum is 3×10^{-5}
- There is a high degree of local structure, with galaxies, galaxy clusters and galaxy filaments. Supermassive objects like galaxy filaments, voids and walls put the cosmological principle into question.
- Dark energy (or vacuum energy) is at 0.69, corresponding to 69% of total energy density in the universe.

Now we'll derive the Einstein Equations for a Robertson-Walker Universe, and show that this satisfies the isotropy and homogeneity conditions at all points.

- Consider an infinite, homogeneous gas cloud with mass density ρ . Choose an arbitrary point A as the origin and consider a thin spherical shell centered around A with radius r . By conservation of energy, we have $(\frac{\dot{r}}{r})^2 + \kappa(r) = \frac{8G\rho}{3}$. κ is $\frac{k}{r^2}$ where k is the total energy.
- By using the First Law of Thermodynamics and assuming constant entropy, we can obtain $\frac{d\rho}{dt} + \frac{3}{r} \frac{dr}{dt} (\rho + p) = 0$. Taking the first derivative of the previous line and combining, we have $\frac{\dot{r}}{r} + \frac{4\pi G}{3} (\rho + 3p) = 0$.
- Choosing a coordinate comoving with the gas cloud, we can define $\mathbf{r}(t) = a(t)\mathbf{r}_0$. $a(t)$ is the scale factor that depends only on time (no dependence on space because isotropy). From this, we can rewrite our earlier equations to obtain $\frac{\ddot{a}}{a} + \frac{4\pi G}{3} (\rho + 3p) = 0$
 $(\frac{\dot{a}}{a})^2 + \frac{\kappa(r)}{a^2} = \frac{8G\rho}{3}$. These are the the einstein equations in component form.
- The continuity equation of the stress energy tensor can similarly be rewritten $\dot{\rho} + 3(\frac{\dot{a}}{a})(\rho + p) = 0$.
- These equations can be further simplified using equations of state.
 Dust: $p = 0$
 Radiation: $p = \frac{1}{3}\rho$
 Vacuum Energy: $p = -\rho$

4.2 History of the Universe

Hubble's discovery of the expansion of the universe and Penzias & Wilson's discovery of the cosmic microwave background suggest the existence of the universe in a "hot, dense" state in the very beginning, before expansion. This is the popularly named "**Big Bang**" model (Gamow 1948, Dicke, Peebles 1965).

Furthermore the existence of General Relativity combined with the energy dominance condition on matter (energy is positive, and energy-momentum doesn't travel faster than light) implies the existence of a cosmological singularity (Penrose, Hawking, Geroch 1966). (Why?)

- Starting with the Friedmann universe, we can do a convenient change of variables by replacing the cumbersome $(1 - kr^2)^{-1}$ with χ . This is known as the comoving distance. As the universe expands, the proper distance between locations change with time but χ remains constant $D(t) = a(t)\chi$.

$$ds^2 = -dt^2 + a^2(t)[d\chi^2 + f^2(\chi)(d\theta^2 + \sin^2\theta d\phi^2)]$$

where $f(\chi) = r$. This gives us nice forms for χ :

- i $k = 1$ Closed universe: $f(\chi) = \sin(\chi)$
- ii $k = 0$ Flat universe: $f(\chi) = \sinh \chi$
- iii $k = -1$ Open universe: $f(\chi) = \chi$

In the case where $f(\chi) = \sin(\chi)$ and $0 \leq \chi \leq 2\pi$ we have a closed universe. For $f(\chi) = \chi$ and $0 \leq \chi \leq \infty$ this is a flat universe. Finally, for $f(\chi) = \sinh(\chi)$ then this is an open universe. (What do these mean?)

- Substituting the equations of state into the continuity equation we can obtain scaling factors of the energy density with a In the flat universe,

- i dust : $\rho_m \sim a^{-3}$
- ii radiation : $\rho_\gamma \sim a^{-4}$
- iii vacuum : $\rho \sim \text{const}$

- Solving the Einstein equation with the Friedmann metric we obtain the following:

$$G_{00} \left(\frac{\dot{a}}{a} \right)^2 + \frac{k}{a^2} - \frac{\Lambda}{3} = \frac{8\pi G\rho}{3}$$

This is the energy conservation equation, which relates the rate to energy density and

$$G_{ii} \rightarrow 3 \left(\frac{\ddot{a}}{a} \right) = -4\pi G(\rho + 3p)$$

This is the acceleration equation.

- If we solve these equations using the equations of state in the flat case ($k = 0$) we obtain the following:

- i In a matter dominated universe: $a \sim t^{\frac{2}{3}}$
- ii Radiation dominated: $a \sim t^{\frac{1}{2}}$
- iii Vacuum dominated: $a \sim e^{Ht}$

4.3 Singularity

A big bang corresponds to a scale factor of $a = 0$. We can ask the question if our universe was ever at this point ie: does it take finite time to reach $a = 0$ in the past?

- First, assume the universe is radiation-dominated (which it would've been at that time) and with $\Lambda = 0$. Then from the G_0^0 equation, we obtain $a^2 = \left(\frac{32}{3}\pi GB\right)^{\frac{1}{2}}(t - t_0)$ where $\rho = Ba^{-4}$. This means it takes finite time to reach $a = 0$.
- A similar analysis for the G_i^i equation gives us $\frac{\dot{a}}{a} \rightarrow \infty$ at $t = t_0$, which implies this time has infinite expansion rate.
- Causality laws: the speed of sound is always less than the speed of light imply that $\rho \rightarrow \infty$ at this time.

Since the universe began at $R = 0$ a finite time in the past, this makes the Big Bang inevitable. This point in time is called the cosmological singularity. Penrose and Hawking's singularity theorems (1973) show that the universe must have been a singularity even if the conditions of isotropy and homogeneity were relaxed.

4.4 The FLRW Universe

- We define the proper distance between 2 galaxies as the integral

$$d(t) = a(t) \int_0^r \frac{dr}{\sqrt{1 - kr^2}} = a(t) \begin{cases} \sin^{-1} r & \text{for } k = 1 \\ r & k = 0 \\ \sinh^{-1} r & k = -1 \end{cases}$$

Meanwhile the proper velocity is

$$v(t) \equiv \dot{d}(t) = \frac{\dot{a}(t)}{a(t)} d(t) = H(t) d(t)$$

this is Hubble's law, where $H_0(t_0)$ is the present value of the hubble constant.

- Let r_{ph} be the coordinate of the most distant object observable, with the earliest light coming from time t_{min} . For light rays, we can set $ds = 0$ giving us

$$\chi_p = c \int_{t_{min}}^{t_0} \frac{dt}{a(t)} = \int_0^{r_{ph}} \frac{dr}{\sqrt{1 - kr^2}}$$

The closed universe model has $r_{ph} = \sin(\chi_p)$ which is a finite value, so there exists a horizon beyond which you cannot see past (why?). This is called the particle horizon.

4.5 The CMB

4.6 The Early Universe