

MATH 323 Class Notes

Owen Lewis

Summer 2018

Contents

1	May 1, 2018	3
1.1	Definitions	3
1.2	How do we Apply this?	4
2	May 2, 2018	5
2.1	Properties of \mathbb{P}	5
2.2	Equiprobability	5
2.3	Counting Tools	5
2.3.1	The Cartesian Product	6
2.3.2	Permutations	6
2.3.3	Combinations	7
2.4	Properties of Combinations	8
2.4.1	Pascal's Triangle	8
3	May 3, 2018	9
3.1	Binomial Theorem	9
3.2	Conditional Probability	9
3.3	Independent Events	10
3.4	Baye's Rule	10
4	May 7, 2018	12
5	May 8, 2018	13
6	May 9, 2018	14
7	May 10, 2018	15
8	May 14, 2018	16
9	May 16, 2018	17
10	May 17, 2018	18
11	May 21, 2018	19
12	May 22, 2018	20
13	May 23, 2018	21
14	May 24, 2018	22
15	May 28, 2018	23
16	May 29, 2018	24

17 May 30, 2018	25
18 May 31, 2018	26

1 May 1, 2018

1.1 Definitions

Let Ω be the set of all possible outcomes. We call Ω the *Sample Space*.

Ex 1: Flipping a coin. The possible outcomes are H and T
 $\Rightarrow \Omega = \{H, T\}$.

Ex 2: Tossing a die. We list all the outcomes as ω_i where i is the face of the die that we land on. We'll assume a normal 6-sided die
 $\Rightarrow \Omega = \{\omega_1, \omega_2, \dots, \omega_6\}$.

Ex 3: Flipping a coin until an H appears. The possible outcomes are $H, TH, TTH, \dots, TT \dots TH$ (with $n - 1$ T s), \dots ad infinitum.

$\Rightarrow \Omega = \{H, TH, TTH, TTTH, TTTTH, \dots\}$.

Note that in this case, Ω is a (countably) infinite set!

Let Ω be a sample space. Any subset A of Ω is called an *Event*.

- If $A = \emptyset$, then we call A the *Null Event*
- If $A = \Omega$, then we call A the *Certain Event*
- If $|A| = 1$, then we call A an *Elementary Event*

Ex: Tossing a die. Let $\Omega := \{\omega_1, \omega_2, \dots, \omega_6\}$, $A := \{\omega_1, \omega_2\}$. Then A is an event but is not an elementary event.

If A is an event, then A^c is also an event called the *complement event* of A .

If A, B are two disjoint events then we call A , and B *mutually exclusive*, or *disjoint*.

Let Ω be a sample space, \mathcal{P} be the power set of Ω . A *Probability* \mathbb{P} on Ω is a function $\mathbb{P} : \mathcal{P}(\Omega) \rightarrow [0, 1]$, such that:

1. $\forall A \subseteq \Omega, 0 \leq \mathbb{P}(A) \leq 1$
2. $\mathbb{P}(\Omega) = 1$
3. If $A_1, A_2, \dots, A_n, \dots$ is a sequence of pairwise disjoint events then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$$

1.2 How do we Apply this?

Let Ω be a discrete set, $E_i = \{\omega_i\}$ be an elemental event, with $E_i \subseteq \Omega$. A probability on Ω is given by a sequence $\mathbb{P}_1, \mathbb{P}_2, \dots, \mathbb{P}_n, \dots$ of positive numbers such that

$$\mathbb{P}(E_i) = p_i, \text{ and } \sum_i \mathbb{P}(p_i) = 1$$

If $A \subseteq \Omega$, then

$$\mathbb{P}(A) = \sum_{\omega_i \in A} \mathbb{P}(p_i)$$

Ex 1: Toss a die.

a) Given that $\mathbb{P}(\omega_2) = \mathbb{P}(\omega_4) = \mathbb{P}(\omega_5) = \mathbb{P}(\omega_6) = \frac{1}{6}$, $\mathbb{P}(\omega_1) = \frac{1}{4}$, find $\mathbb{P}(\omega_3)$.

b) Find the probability that the die will land on an odd face.

Solution:

a) $\Omega = \{\omega_1, \omega_2, \dots, \omega_6\}$. $\mathbb{P}(\omega_3)$ is a singleton, and as we know $\sum_i \mathbb{P}(p_i) = 1$ then $\sum_{i=1}^6 \mathbb{P}(\omega_i) = 1 \implies \mathbb{P}(\omega_2) + \mathbb{P}(\omega_4) + \mathbb{P}(\omega_5) + \mathbb{P}(\omega_6) + \mathbb{P}(\omega_1) + \mathbb{P}(\omega_3) = 1 \implies \frac{4}{6} + \frac{1}{4} + \mathbb{P}(\omega_3) = 1 \implies \mathbb{P}(\omega_3) = 1 - \frac{11}{12} \implies \mathbb{P}(\omega_3) = \frac{1}{12}$.

b) Let $A \subseteq \Omega$ be the subset of Ω containing all the odd faces. The total probability of A is therefore the sum of all the probabilities of the elements of $A \Rightarrow \mathbb{P}(A) = \frac{1}{4} + \frac{1}{12} + \frac{1}{6} = \frac{1}{2}$.

Ex 2: Given a countably infinite sample space, find a constant c such that $\mathbb{P}(\{\omega_n\}) = c(\frac{1}{5})^n$ for some n .

Solution: $\sum_{n=1}^{\infty} \mathbb{P}(\{\omega_n\}) = 1 \Rightarrow \sum_{n=1}^{\infty} c(\frac{1}{5})^n = 1 \Rightarrow \sum_{n=1}^{\infty} (\frac{1}{5})^n = \frac{1}{c}$. Notice how $\sum_{n=1}^{\infty} (\frac{1}{5})^n$ is a geometric series that converges to $\frac{1}{4} \Rightarrow$ therefore $\frac{1}{c} = \frac{1}{4} \Rightarrow c = 4$.

2 May 2, 2018

2.1 Properties of \mathbb{P}

Let Ω be a sample space and let \mathbb{P} be a probability on Ω . Then:

1. $\mathbb{P}(\emptyset) = 0$
2. $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
3. $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$

Proof:

1. Set $A_1 = \Omega$, $A_2 = \emptyset$. Then $A_1 \cup A_2 = \Omega$, and $A_1 \cap A_2 = \emptyset$. Therefore $\mathbb{P}(A_1 \cup A_2) = \mathbb{P}(A_1) + \mathbb{P}(A_2) \Rightarrow \mathbb{P}(\Omega) = \mathbb{P}(\emptyset) + \mathbb{P}(\Omega) \Rightarrow 1 = \mathbb{P}(\emptyset) + 1 \Rightarrow 0 = \mathbb{P}(\emptyset)$. \square
2. $A^c \cup A = \Omega$, and $A^c \cap A = \emptyset$. Then $\mathbb{P}(\Omega) \Rightarrow 1 = \mathbb{P}(A^c \cup A) \Rightarrow 1 = \mathbb{P}(A) + \mathbb{P}(A^c) \Rightarrow \mathbb{P}(A^c) = 1 - \mathbb{P}(A)$. \square
3. It's easy to show that $A = (A \setminus B) \cup (A \cap B)$, and $\emptyset = (A \setminus B) \cap (A \cap B)$. Similarly, $\mathbb{P}(A) = \mathbb{P}(A \setminus B) + \mathbb{P}(A \cap B)$. From these, it follows that $A \cup B = A \cup (B \setminus A) \Rightarrow \mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B \setminus A) \Rightarrow \mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$. \square

2.2 Equiprobability

Let Ω be a finite sample space. Set $N := |\Omega|$. Equiprobability means that all outcomes have the same probability $\mathbb{P} = \frac{1}{N}$. Let $A \subseteq \Omega$ be an event. Then we have $\mathbb{P}(A) = |A| \cdot \frac{1}{N}$, or

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|}$$

This is great because it means that in **equiprobability** problems we just need to count the cardinality of A , count the cardinality of Ω , and divide them, and we're done. Too bad counting isn't really all that easy.

2.3 Counting Tools

We have here three tools to help us calculate the cardinalities of huge (finite) subsets of huge sample spaces, each with their own specific situations that require its use

1. The Cartesian Product
2. Permutations
3. Combinations

2.3.1 The Cartesian Product

We all know what the cartesian product *is*. In probability we use it in our experiment when we have more than one "input" each with its own possible outcome, for example, we roll three dice, or flip two coins.

Let A, B , be sets such that $|A| = a$, and $|B| = b$. Then the cardinality of the cartesian product $|(A \times B)|$ is $|(A \times B)| = a \cdot b$.

Ex 1: Suppose we roll a die twice. What is the cardinality of the sample space Ω ?

Solution: If we roll a die once we have $\Omega = \{\omega_1, \dots, \omega_6\}$. Therefore, the sample space for rolling a die twice is $\Omega \times \Omega$. The cardinality of our sample space $\Omega \times \Omega$ is $|\Omega \times \Omega| = 6 \cdot 6 = 36$.

Ex 2: Suppose we roll a fair die twice. What is the probability that the outcome is even?

Solution: If the sum of the two outcomes is even then both must either be even or both must be odd. Let A be the event where the sum of the rolls is even, and let A_1 be event where both individual rolls are even, A_2 be the event where both individual rolls are odd. For example,

$$A_1 = \{(2, 2), (2, 4), (2, 6), (4, 2), (4, 4), (4, 6), (6, 2), (6, 4), (6, 6)\}$$

Where the 1st element in each ordered pair is the outcome of the 1st roll and the 2nd element in each ordered pair is the outcome of the 2nd roll. Therefore, we have that $|A_1| = |A_2| = 9$.

Since naturally $A = A_1 \cup A_2$ then $\mathbb{P}(A) = \mathbb{P}(A_1) + \mathbb{P}(A_2)$. We also know that the die was fair, so we can use our equiprobability formula here.

$$\mathbb{P}(A) = \mathbb{P}(A_1) + \mathbb{P}(A_2) = \frac{|A_1|}{|\Omega|} + \frac{|A_2|}{|\Omega|} = \frac{9}{36} + \frac{9}{36} = \frac{18}{36} = \frac{1}{2}.$$

2.3.2 Permutations

A permutation of r integer elements chosen from n (possible elements) is equivalent to a successive draw, without replacement, of r elements from a list of n elements. We denote the number of possibilities by P_r^n . The general formula for P_r^n is

$$P_r^n = n \cdot (n - 1) \cdot \dots \cdot (n - r + 1) = \frac{n!}{(n - r)!}$$

Ex:

a) A thick black bag contains 4 balls: 1 green, 1 blue, 1 red, 1 yellow. The bag is made of lead, or something, and also light cannot exist in this bag. You couldn't see into this bag if your life depended on it. Draw successively two balls from the bag without putting them back in. What is the probability that

the second ball drawn is green?

b) What is the probability that one of the two balls drawn will be green?

Solution:

a) Let Ω be the set of permutations of 2 balls chosen from the bag containing 4 balls. Then, $|P_2^4| = \frac{4!}{2!} = \frac{24}{2} = 12$. Then let A be the event where the second ball is green. The cardinality of A is 3, as if we take for granted that the second ball is green, then there are 3 other different-coloured balls in the bag to accompany it. Since the bag is so dark that it's physically impossible to see inside it, all the balls in the bag have an equal probability of being drawn. Therefore,

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|} = \frac{3}{12} = \frac{1}{4}$$

b) Let B be the event where one of the balls drawn is green. Suppose B_1 is the event where the first ball drawn is green, and B_2 is the event where the second ball drawn is green. In part a) we found that $|B_2| = 3$, and a parallel argument shows that $|B_1| = 3$ as well. Thus,

$$\mathbb{P}(B) = \frac{|B_1|}{|\Omega|} + \frac{|B_2|}{|\Omega|} = \frac{3}{12} + \frac{3}{12} = \frac{6}{12} = \frac{1}{2}$$

2.3.3 Combinations

Consider a set Ω with n elements. Let r be an integer such that $0 \leq r \leq n$. A combination C_r^n , also denoted $\binom{n}{r}$ (pronounced n choose r), is the number of subsets of Ω containing r elements. The general formula is

$$C_r^n = \frac{P_n^r}{r!} = \frac{n!}{r! \cdot (n-r)!}$$

Ex:

a) Out of a deck of 52 cards how many distinct 5-card hands are possible?

b) What is the probability that a given hand contains at least one ace?

Solution:

a) $\binom{52}{5} = \frac{52!}{5! \cdot (47!)} = 2,598,960$

b) In this case it is easier to calculate the probability where the hand contains no aces and then subtract that from 1 to find the probability that we have an ace. If A is the event where the hand contains an ace, then A^c is the event where a hand contains no ace. Then

$$\mathbb{P}(A^c) = \frac{|A^c|}{|\Omega|} = \frac{C_5^{48}}{C_5^{52}} = \frac{1,712,304}{2,598,960} = \frac{35,673}{54,145}$$

Then we need to subtract this from 1 and we're done

$$\mathbb{P}(A) = 1 - \frac{35,673}{54,145} \approx 0.3412$$

2.4 Properties of Combinations

Here are some properties of combinations


1. $C_0^n = C_n^n = 1$
2. $C_r^n = C_{n-r}^n$ $0 \leq r \leq n$
3. $C_r^n = C_r^{n-1} + C_{r-1}^{n-1}$ $0 \leq r \leq n$

The proofs are really easy and I don't want to bother typing them out but here is the idea behind each of them:

1. Trivial
2. Induction on n
3. Also induction on n

2.4.1 Pascal's Triangle

Pascal's triangle is a table of combinations C_r^n . The rows of the triangle represent n , starting at 0 at the tip and working down, and the r^{th} element from the left (starting at 0) represents r . Each number in the triangle is determined by summing the two numbers directly above it.



Pascal's Triangle

3 May 3, 2018

3.1 Binomial Theorem

$$(a+b)^n = \sum_{k=0}^n C_k^n \cdot a^k \cdot b^{n-k}$$

$$\text{Ex 1: } (a+b)^3 = C_0^3 \cdot a^6 b^3 + C_1^3 \cdot a b^2 + C_2^3 \cdot a^2 b + C_3^3 \cdot a^3 b = b^3 + 3ab^2 + 3ba^3 + a^3$$

Ex 2: Find the coefficient of x^6 in the expansion of $(x^2 + 2)^7$.

Solution: If we let $x^2 := a$ and $2 := b$, then from the binomial theorem:

$$(a+b)^n = \sum_{k=0}^n C_k^n \cdot a^k \cdot b^{n-k} = \sum_{k=0}^n C_k^7 \cdot a^k \cdot b^{7-k} = C_3^7 \cdot 2^4 = 560$$

3.2 Conditional Probability

Let Ω be a sample space and let \mathbb{P} be a probability on Ω . Let $A \subseteq \Omega$ be an event such that $\mathbb{P}(A) > 0$, and let $B \subseteq \Omega$ be another event. The *conditional probability of B given A*, is defined as

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B \cap A)}{\mathbb{P}(A)}$$

Ex 1: We have 2 urns. Urn 1 contains 7 red balls and 4 blue balls, and urn 2 contains 5 red balls and 6 blue balls. First, we choose a ball from urn 1 and place it into urn 2, then we remove a ball from urn 2. What is the probability that the ball that we remove from urn 2 will be blue?

Solution: Let B be the event where the ball drawn from urn 2 is blue, and let A_r, A_b be the events where we draw a red or a blue ball from urn 1 respectively. Then $\mathbb{P}(A_r) = \frac{7}{11}$ and $\mathbb{P}(A_b) = \frac{4}{11}$. Also,

$$B = B \cap \Omega \iff B = B \cap (A_r \cup A_b) \iff B = (B \cap A_r) \cup (B \cap A_b)$$

Which then implies

$$\mathbb{P}(B) = \mathbb{P}(B \cap A_r) + \mathbb{P}(B \cap A_b) \iff \mathbb{P}(B) = \mathbb{P}(B|A_r) \cdot \mathbb{P}(A_r) + \mathbb{P}(B|A_b) \cdot \mathbb{P}(A_b)$$

So all that is left is to compute $\mathbb{P}(B|A_r)$, and $\mathbb{P}(B|A_b)$.

From the definition, we can see that $\mathbb{P}(B|A_r) = \frac{1}{2}$, and $\mathbb{P}(B|A_b) = \frac{7}{12}$, so finally

$$\mathbb{P}(B) = \frac{6}{12} \cdot \frac{7}{11} + \frac{7}{12} \cdot \frac{4}{11} = \frac{70}{132}$$

Ex: 2 Roll a fair die. Let $A \subseteq \Omega$ be the event where the outcome is even, $B \subseteq \Omega$ be the event where the outcome is odd, $C \subseteq \Omega$ be the event where the outcome is either a 1 or a 2. Compute all the conditional probabilities.

Solution First $\mathbb{P}(A) = \frac{1}{2}$, $\mathbb{P}(B) = \frac{1}{2}$, and $\mathbb{P}(C) = \frac{1}{3}$ Then:

$$\begin{aligned}\mathbb{P}(A|B) &= \frac{\frac{1}{6}}{\frac{1}{2}} = \frac{1}{3}, \text{ and } \mathbb{P}(B|A) = \frac{\frac{1}{6}}{\frac{1}{2}} = \frac{1}{3} \\ \mathbb{P}(A|C) &= \frac{1}{2}, \text{ and } \mathbb{P}(C|A) = \frac{1}{3} \\ \mathbb{P}(B|C) &= 1, \text{ and } \mathbb{P}(C|B) = \frac{2}{3}\end{aligned}$$

3.3 Independent Events

Let A , and B be two events over some sample space Ω . A , and B are said to be *independent* if and only if:

1. $\mathbb{P}(A|B) = \mathbb{P}(A)$
2. $\mathbb{P}(B|A) = \mathbb{P}(B)$
3. $\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$

Ex:

- a) From the previous example, are A and B independent?
- b) What about A and C ?
- c) B and C ?

Answers:

- a) No.
- b) Yes.
- c) No.

3.4 Baye's Rule

Let $A, B \subseteq \Omega$ be events on a sample space Ω . Then the equation

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(B|A) \cdot \mathbb{P}(A)}{\mathbb{P}(B)}$$

is called Baye's Rule, or Baye's Theorem.

Theorem: Total Probability Rule

Let Ω be a sample space, and let $A \subseteq \Omega$ be an event. Suppose we partition Ω like $\Omega = \{B_1, B_2, \dots, B_n\}$, where the B_i s are pairwise disjoint events such that

$$\bigcup_{i=1}^n B_i = \Omega$$

Then:

1. $\mathbb{P}(A) = \sum_{i=1}^n \mathbb{P}(A|B_i) \cdot \mathbb{P}(B_i)$, and
2. If $k \in \mathbb{N}$ is fixed, then $\mathbb{P}(B_k|A) = \frac{\mathbb{P}(A|B_k) \cdot \mathbb{P}(B_k)}{\sum_{i=1}^n \mathbb{P}(A|B_i) \cdot \mathbb{P}(B_i)}$

Proof:

1. We know that $A = A \cap \Omega \iff A = A \cap (\bigcup_{i=1}^n B_i) \iff A = \bigcup_{i=1}^n A \cap B_i$.
Therefore, $\mathbb{P}(A) = \sum_{i=1}^n \mathbb{P}(A \cap B_i) \iff \mathbb{P}(A) = \sum_{i=1}^n \mathbb{P}(A|B_i) \cdot \mathbb{P}(B_i)$ \square
2. $\mathbb{P}(B_k|A) = \frac{\mathbb{P}(B_k \cap A)}{\mathbb{P}(A)} = \frac{\mathbb{P}(A|B_k) \cdot \mathbb{P}(B_k)}{\mathbb{P}(A)} = \frac{\mathbb{P}(A|B_k) \cdot \mathbb{P}(B_k)}{\sum_{i=1}^n \mathbb{P}(A|B_i) \cdot \mathbb{P}(B_i)}$ \square

Ex: A (very simple) forecast model

Suppose that on a given day the weather is one of two states: sunny or rainy. Let $R \subseteq \Omega$ be the event where it's rainy and $S \subseteq \Omega$ be the event where it's sunny. If today is rainy then the probability that tomorrow will also be rainy is 60%. On the other hand, if today is sunny then the probability that tomorrow will be sunny is 70%.

a) If Monday is sunny then what is the probability that Wednesday will also be sunny?

b) If Wednesday is sunny then what is the probability that Tuesday was rainy?

Solutions:

a) Let A be the event where it is sunny on Wednesday. Then

$$\mathbb{P}(A) = \mathbb{P}(A|A_1) \cdot \mathbb{P}(A_1) + \mathbb{P}(A|A_2) \cdot \mathbb{P}(A_2) = 0.7 \cdot 0.7 + 0.4 \cdot 0.3 = 0.61$$

where A_1 is the event where it's sunny on Tuesday, and A_2 is the event where it's rainy on Tuesday.

$$\text{b) } \mathbb{P}(A_2|A) = \frac{\mathbb{P}(A|A_2) \cdot \mathbb{P}(A_2)}{\mathbb{P}(A)} = \frac{0.12}{0.61} = \frac{12}{61}$$

4 May 7, 2018

5 May 8, 2018

6 May 9, 2018

7 May 10, 2018

8 May 14, 2018

9 May 16, 2018

10 May 17, 2018

11 May 21, 2018

12 May 22, 2018

13 May 23, 2018

14 May 24, 2018

15 May 28, 2018

16 May 29, 2018

17 May 30, 2018

18 May 31, 2018