of discriminant 5 to the canonical form X^2 +¿ XY −¿ Y^2, where s_5 is the odd root of $x^2 = 5 \bmod p$.

This reduction algorithm, outlined by Gauss and cleverly reported in Buell [3], was described in a masterly fashion in Mathews' book [13], and thus it will be referred to as Mathews' reduction. It can be applied to any quadratic form of a non-perfect square discriminant. The square root s5 may be computed with deterministic polynomial complexity by distinguishing whether p is congruent 3 or 1 modulo 4. In the first case, s_5 is one of the two square roots $\pm 5^{\frac{p+1}{4}} \bmod p$, and the computation clearly has deterministic polynomial complexity. In the second case Schoof's algorithm [18, 21] is applied to compute both $\sqrt{-1}$ and $\sqrt{-5}$ modulo p, thus s_5 is one of the two values $\pm\sqrt{-1}\sqrt{-5} \bmod p$. Schoof's algorithm evaluates a square root modulo p in a number of arithmetical operations of order $O(\log^t p)$ with $t \leq 10$, that is with deterministic polynomial complexity [18]. A description of Schoof's algorithm can be found in the original paper [18] or in Washington's book [21], and is based on properties of elliptic curves that are briefly summarized.

.....

1. Representation of primes in Z(ω)

.....

2. Computational remarks

The representations of primes $\pi \in Q(\omega)$ as the sums of two squares, established by Theorems 2 and 3, are computed with deterministic poly-nomial complexity using procedures that will be described separately for the different cases.

4.1 *Primes* $p \equiv 13, 17 \bmod 20$

The representation p = a2 + b2 does not need any further comment, since p = 1 mod 4 is the sum of two integer squares by Fermat's theorem, and its computation was shown in Section 2.

4.2 Primes p

+¿ $y_0 F_{2k+1}$)

.....

5. Conclusions

The representation of primes as sums of two squares in $Z(\omega)$ allows us to formulate a theorem concerning the representation of any integer in $Z(\omega)$:

Theorem 4. *In the field $Q(\omega)$, any totally positive integer*

.....

The computation of the two-square representation of any element in $Z(\omega)$ is equivalent to factoring, as in Z, therefore at the present time only the two-square representations of primes can be computed with deterministic polynomial complexity.

References

[1] B. Berndt and R. Evans, Sums of Gauss, Jacobi and Jacobsthal, *Journal of Number Theory*, Vol. 11 (1979), pp. 349–398.

[2] J. Brillhart, Note on representing a prime as a sum of two squares, *Mathematics of Computation*, Vol. 26 (1972), pp. 1011–1013.

[3] D. A. *Buell, Binary Quadratic Forms*, Springer-Verlag, New York, 1989.

[4] D. A. Cox, *Primes of the Form*, Wiley, New York, 1989.

[5] F. W. Dodd, *Number Theory in the Quadratic Field with Golden Section Unit*, Polygonal Pub. House, Paissac, 1983.

[6] C.F. Gauss, *Disquisitiones Arithmeticae*, Springer-Verlag, New York, 1986.

[7] G. H. Hardy and E. M. Wright, An *Introduction to the Theory of Numbers*, Oxford University Press, Oxford, 1971.

[8] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, New York, 1998.

[9] D. E. Knuth, *The Art of Computer Programming: Seminumerical Algorithms*, Addison-Wesley, Reading, 1970.

[10] S. Lang, *Elliptic Functions*, Springer-Verlag, New York, 1987.

[11] F. Lemmermeyer, *Reciprocity Laws*, Springer-Verlag, New York, 2000.

[12] R. Lidl and H Niederreiter, *Finite Fields*, Addison-Wesley, Reading