

Charles M. Lizza
William C. Baton
Sarah A. Sullivan
SAUL EWING ARNSTEIN & LEHR LLP
One Riverfront Plaza, Suite 1520
Newark, NJ 07102-5426
(973) 286-6700
clizza@saul.com

*Attorneys for Plaintiff
Celgene Corporation*

**UNITED STATES DISTRICT COURT
DISTRICT OF NEW JERSEY**

CELGENE CORPORATION,

Plaintiff,

v.

QILU PHARMACEUTICAL CO. LTD.,

Defendant.

Civil Action No. _____

**COMPLAINT FOR
PATENT INFRINGEMENT**

(Filed Electronically)

Plaintiff Celgene Corporation (“Celgene”), by its undersigned attorneys, for its Complaint against Defendant Qilu Pharmaceutical Co. Ltd. (“Qilu”), alleges as follows:

Nature of the Action

1. This is an action for patent infringement under the patent laws of the United States, 35 U.S.C. § 100, *et seq.*, arising from Qilu’s submission of Abbreviated New Drug Application (“ANDA”) No. 217265 (“Qilu’s ANDA”) to the United States Food and Drug Administration (“FDA”) seeking approval to manufacture, use, import, distribute, offer to sell, and/or sell generic versions of Celgene’s Revlimid® drug products prior to the expiration of United States Patent Nos. 7,465,800 (the “‘800 patent”) and 7,855,217 (the “‘217 patent”) (together, “the patents-in-suit”), both owned by Celgene.

The Parties

2. Plaintiff Celgene is a biopharmaceutical company committed to improving the lives of patients worldwide. Celgene focuses on, and invests heavily in, the discovery and development of products for the treatment of severe and life-threatening conditions. Celgene is a world leader in the treatment of many such diseases, including cancer. Celgene is a corporation organized and existing under the laws of the State of Delaware, having a principal place of business at 86 Morris Avenue, Summit, New Jersey 07901.

3. On information and belief, Defendant Qilu Pharmaceutical Co. Ltd. is a corporation organized and existing under the laws of China, having a principal place of business at 8888 Lvyou Road, High-tech Zone, Jinan, 250104, China.

The Patents-in-Suit

4. On December 16, 2008, the United States Patent and Trademark Office (“USPTO”) duly and lawfully issued the ’800 patent, entitled, “Polymorphic Forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.” A copy of the ’800 patent is attached hereto as Exhibit A.

5. On December 21, 2010, the USPTO duly and lawfully issued the ’217 patent, entitled, “Polymorphic Forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.” A copy of the ’217 patent is attached hereto as Exhibit B.

The Revlimid® Drug Product

6. Celgene holds an approved New Drug Application (“NDA”) under Section 505(a) of the Federal Food Drug and Cosmetic Act (“FFDCA”), 21 U.S.C. § 355(a), for lenalidomide capsules (NDA No. 021880), which it sells under the trade name Revlimid®. The claims of the

patents-in-suit cover, *inter alia*, solid forms of lenalidomide and pharmaceutical compositions containing lenalidomide.

7. Pursuant to 21 U.S.C. § 355(b)(1) and attendant FDA regulations, the patents-in-suit are listed in the FDA publication, “Approved Drug Products with Therapeutic Equivalence Evaluations” (the “Orange Book”), with respect to Revlimid®.

Jurisdiction and Venue

8. This Court has jurisdiction over the subject matter of this action pursuant to 28 U.S.C. §§ 1331, 1338(a), 2201, and 2202.

9. This Court has personal jurisdiction over Qilu by virtue of, *inter alia*, its systematic and continuous contacts with the State of New Jersey. On information and belief, Qilu’s subsidiary, Qilu Healthcare Inc., maintains a regular and established, physical place of business in Princeton, New Jersey.

10. On information and belief, Qilu Healthcare Inc. is registered with the State of New Jersey’s Division of Revenue and Enterprise Services as a business operating in New Jersey under Business ID No. 101042575. On information and belief, Qilu Healthcare Inc. is registered with the State of New Jersey’s Department of Health as a drug manufacturer and wholesaler under Registration No. 5005245.

11. On information and belief, Qilu purposefully has conducted and continues to conduct business in this Judicial District.

12. On information and belief, Qilu is in the business of, among other things, manufacturing, marketing, importing, offering for sale, and selling pharmaceutical products, including generic drug products, throughout the United States, including in this Judicial District.

On information and belief, Qilu also prepares and/or aids in the preparation and submission of ANDAs to the FDA, including Qilu's ANDA.

13. On information and belief, this Judicial District is a likely destination for the generic drug products described in Qilu's ANDA.

14. This Court has personal jurisdiction over Qilu because, *inter alia*, it: (1) has purposefully availed itself of the privilege of doing business in New Jersey, including directly or indirectly through its subsidiary, agent, and/or alter egos, including Qilu Healthcare, Inc., a company with its principal place of business in New Jersey; and (2) maintains extensive and systematic contacts with the State of New Jersey, including through the marketing, distribution, and/or sale of generic pharmaceutical drugs in New Jersey, including through, directly or indirectly, Qilu Healthcare, Inc.

15. On information and belief, Qilu derives substantial revenue from directly or indirectly selling generic pharmaceutical products and/or active pharmaceutical ingredient(s) used in generic pharmaceutical products sold throughout the United States, including in this Judicial District.

16. In the alternative, this Court has personal jurisdiction over Qilu because the requirements of Federal Rule of Civil Procedure 4(k)(2) are met as (a) Celgene's claims arise under federal law; (b) Qilu is a foreign defendant not subject to general personal jurisdiction in the courts of any state; and (c) Qilu has sufficient contacts with the United States as a whole, including, but not limited to, preparing and submitting ANDAs to the FDA and/or manufacturing, importing, offering to sell, and/or selling pharmaceutical products that are distributed throughout the United States, such that this Court's exercise of jurisdiction over Qilu satisfies due process.

17. On information and belief, Qilu works in privity and/or concert either directly or indirectly through one or more of its wholly owned subsidiaries with respect to the regulatory approval, manufacturing, use, importation, marketing, offer for sale, sale, and distribution of generic pharmaceutical products throughout the United States, including in this Judicial District.

18. On information and belief, Qilu submitted and/or actively participated in the submission of its ANDA. On information and belief, Qilu will work in privity and/or concert with Qilu Healthcare, Inc., and/or other related entities towards the regulatory approval, manufacturing, use, importation, marketing, offer for sale, sale, and distribution of generic pharmaceutical products, including the 2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg, and 25 mg lenalidomide capsules that are the subject of Qilu's ANDA ("Qilu's Proposed Products"), throughout the United States, including in New Jersey and in this Judicial District, prior to the expiration of the patents-in-suit.

19. On information and belief, Qilu intends to benefit directly if its ANDA is approved by participating in the manufacture, importation, distribution, and/or sale of the generic drug products that are the subject of Qilu's ANDA.

20. Venue is proper in this Judicial District for Qilu pursuant to 28 U.S.C. §§ 1391 and/or 1400(b), including, for example, because Qilu is a company organized and existing under the laws of China and may be sued in any judicial district.

Acts Giving Rise To This Suit

21. Pursuant to Section 505 of the FFDCA, Qilu submitted its ANDA seeking approval to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Qilu's Proposed Products before the patents-in-suit expire.

22. On information and belief, following FDA approval of Qilu's ANDA, Qilu will make, use, sell, or offer to sell Qilu's Proposed Products throughout the United States, or import such generic products into the United States.

23. On information and belief, in connection with the submission of its ANDA as described above, Qilu provided a written certification to the FDA pursuant to Section 505 of the FFDCA, 21 U.S.C. § 355(j)(2)(A)(vii)(IV) ("Qilu's Paragraph IV Certification"), alleging that the claims of the patents-in-suit are invalid and/or will not be infringed by the activities described in Qilu's ANDA.

24. No earlier than April 29, 2022, Qilu sent a written notice of its Paragraph IV Certification to Celgene ("Qilu's Notice Letter"). Qilu's Notice Letter alleged that the claims of the '800 and '217 patents are invalid and/or will not be infringed by the activities described in Qilu's ANDA. Qilu's Notice Letter also informed Celgene that Qilu seeks approval to market Qilu's Proposed Products before the patents-in-suit expire. Qilu specifically directed Qilu's Notice Letter to Celgene's headquarters in Summit, New Jersey, in this Judicial District.

Count I: Infringement of the '800 Patent

25. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

26. Qilu's submission of its ANDA, with the accompanying Paragraph IV Certification and notice to Celgene of same, to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Qilu's Proposed Products, prior to the expiration of the '800 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

27. There is a justiciable controversy between the parties hereto as to the infringement of the '800 patent.

28. Unless enjoined by this Court, upon FDA approval of Qilu's ANDA, Qilu will infringe one or more claims of the '800 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Qilu's Proposed Products in the United States.

29. Unless enjoined by this Court, upon FDA approval of Qilu's ANDA, Qilu will induce infringement of one or more claims of the '800 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Qilu's Proposed Products in the United States. On information and belief, upon FDA approval of Qilu's ANDA, Qilu will intentionally encourage acts of direct infringement with knowledge of the '800 patent and knowledge that its acts are encouraging infringement.

30. Unless enjoined by this Court, upon FDA approval of Qilu's ANDA, Qilu will contributorily infringe one or more claims of the '800 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Qilu's Proposed Products in the United States. On information and belief, Qilu has had and continues to have knowledge that Qilu's Proposed Products are especially adapted for a use that infringes one or more claims of the '800 patent and that there is no substantial non-infringing use for Qilu's Proposed Products.

31. Celgene will be substantially and irreparably damaged and harmed if Qilu's infringement of the '800 patent is not enjoined.

32. Celgene does not have an adequate remedy at law.

33. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

Count II: Infringement of the '217 Patent

34. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

35. Qilu's submission of its ANDA, with the accompanying Paragraph IV Certification and notice to Celgene of same, to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Qilu's Proposed Products, prior to the expiration of the '217 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

36. There is a justiciable controversy between the parties hereto as to the infringement of the '217 patent.

37. Unless enjoined by this Court, upon FDA approval of Qilu's ANDA, Qilu will infringe one or more claims of the '217 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Qilu's Proposed Products in the United States.

38. Unless enjoined by this Court, upon FDA approval of Qilu's ANDA, Qilu will induce infringement of one or more claims of the '217 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Qilu's Proposed Products in the United States. On information and belief, upon FDA approval of Qilu's ANDA, Qilu will intentionally encourage acts of direct infringement with knowledge of the '217 patent and knowledge that its acts are encouraging infringement.

39. Unless enjoined by this Court, upon FDA approval of Qilu's ANDA, Qilu will contributorily infringe one or more claims of the '217 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Qilu's Proposed Products in the United States. On information and belief, Qilu has had and continues to have knowledge that Qilu's

Proposed Products are especially adapted for a use that infringes one or more claims of the '217 patent and that there is no substantial non-infringing use for Qilu's Proposed Products.

40. Celgene will be substantially and irreparably damaged and harmed if Qilu's infringement of the '217 patent is not enjoined.

41. Celgene does not have an adequate remedy at law.

42. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

PRAAYER FOR RELIEF

WHEREFORE, Plaintiff Celgene respectfully requests the following relief:

(A) A Judgment that Qilu has infringed the patents-in-suit by submitting ANDA No. 217265 with the accompanying Paragraph IV Certification and notice to Celgene of same;

(B) A Judgment that Qilu has infringed, and that Qilu's making, using, selling, offering to sell, or importing Qilu's Proposed Products will infringe one or more claims of the patents-in-suit;

(C) An Order that the effective date of FDA approval of ANDA No. 217265 be a date which is not earlier than the later of the expiration of the patents-in-suit, or any later expiration of exclusivity to which Celgene is or becomes entitled;

(D) Preliminary and permanent injunctions enjoining Qilu and its officers, agents, attorneys and employees, and those acting in privity and/or concert with them, from making, using, offering to sell, selling, or importing Qilu's Proposed Products until after the expiration of the patents-in-suit, or any later expiration of exclusivity to which Celgene is or becomes entitled;

(E) A permanent injunction, pursuant to 35 U.S.C. § 271(e)(4)(B), restraining and enjoining Qilu, its officers, agents, attorneys and employees, and those acting in privity and/or

concert with them, from practicing any solid forms of lenalidomide or compositions claimed in the patents-in-suit, or from actively inducing or contributing to the infringement of any claim of the patents-in-suit, until after the expiration of the patents-in-suit, or any later expiration of exclusivity to which Celgene is or becomes entitled;

- (F) A Judgment that the commercial manufacture, use, offer for sale, sale, and/or importation into the United States of Qilu's Proposed Products will directly infringe, induce infringement of, and/or contribute to infringement of the patents-in-suit;
- (G) To the extent that Qilu, its officers, agents, attorneys and/or employees, or those acting in privity and/or concert with them, has committed any acts with respect to the solid forms of lenalidomide or compositions, other than those acts expressly exempted by 35 U.S.C. § 271(e)(1), a Judgment awarding Celgene damages for such acts;
- (H) If Qilu, its officers, agents, attorneys, and/or employees, or those acting in privity and/or concert with them, engages in the commercial manufacture, use, offer for sale, sale, and/or importation into the United States of Qilu's Proposed Products prior to the expiration of the patents-in-suit, a Judgment awarding damages to Celgene resulting from such infringement, together with interest;
- (I) A Judgment declaring that the patents-in-suit remain valid and enforceable;
- (J) A Judgment that this is an exceptional case pursuant to 35 U.S.C. § 285 and awarding Celgene its attorneys' fees incurred in this action;
- (K) A Judgment awarding Celgene its costs and expenses incurred in this action; and
- (L) Such further and other relief as this Court may deem just and proper.

Dated: June 13, 2022

Of Counsel:

F. Dominic Cerrito
Eric C. Stops
Andrew S. Chalson
QUINN EMANUEL URQUHART & SULLIVAN,
LLP
51 Madison Avenue, 22nd Floor
New York, New York 10010
(212) 849-7000

By: s/ Charles M. Lizza
Charles M. Lizza
William C. Baton
Sarah A. Sullivan
SAUL EWING ARNSTEIN & LEHR LLP
One Riverfront Plaza, Suite 1520
Newark, New Jersey 07102-5426
(973) 286-6700
clizza@saul.com

*Attorneys for Plaintiff
Celgene Corporation*

Anthony M. Insogna
Cary Miller, Ph.D.
JONES DAY
4655 Executive Drive
San Diego, CA 92121
(858) 314-1200

Matthew J. Hertko
JONES DAY
77 W. Wacker Drive
Chicago, IL 60601
(312) 782-3939

CERTIFICATION PURSUANT TO L. CIV. R. 11.2 & 40.1

Pursuant to Local Civil Rules 11.2 and 40.1, I hereby certify that the matter captioned *Celgene Corporation v. Alembic Pharmaceuticals Ltd.*, Civil Action 21-20099 (SDW)(LDW) (D.N.J.) is related to the matter in controversy because the matter in controversy involves the same plaintiff and some of the same patents, and because Qilu is seeking FDA approval to market generic versions of the same pharmaceutical product.

I further certify that, to the best of my knowledge, the matter in controversy is not the subject of any other action pending in any court, or of any pending arbitration or administrative proceeding.

Dated: June 13, 2022

By: s/ Charles M. Lizza

Charles M. Lizza

William C. Baton

Sarah A. Sullivan

SAUL EWING ARNSTEIN & LEHR LLP

One Riverfront Plaza, Suite 1520

Newark, New Jersey 07102-5426

(973) 286-6700

clizza@saul.com

*Attorneys for Plaintiff
Celgene Corporation*

Of Counsel:
F. Dominic Cerrito
Eric C. Stops
Andrew S. Chalson
QUINN EMANUEL URQUHART &
SULLIVAN, LLP
51 Madison Avenue, 22nd Floor
New York, New York 10010
(212) 849-7000

Anthony M. Insogna
Cary Miller, Ph.D.
JONES DAY
4655 Executive Drive
San Diego, CA 92121
(858) 314-1200

Matthew J. Hertko
JONES DAY
77 W. Wacker Drive
Chicago, IL 60601
(312) 782-3939

EXHIBIT A



US007465800B2

(12) **United States Patent**
Jaworsky et al.

(10) **Patent No.:** US 7,465,800 B2
(45) **Date of Patent:** Dec. 16, 2008

(54) **POLYMORPHIC FORMS OF**
3-(4-AMINO-1-OXO-1,3
DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-
2,6-DIONE

(75) Inventors: **Markian S. Jaworsky**, Hopewell, NJ (US); **Roger Shen-Chu Chen**, Edison, NJ (US); **George W. Muller**, Bridgewater, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 596 days.

(21) Appl. No.: **10/934,863**

(22) Filed: **Sep. 3, 2004**

(65) **Prior Publication Data**

US 2005/0096351 A1 May 5, 2005

5,059,595 A 10/1991 Le Grazie
5,073,543 A 12/1991 Marshall et al.
5,120,548 A 6/1992 McClelland et al.
5,229,496 A 7/1993 Deeley et al.
5,354,556 A 10/1994 Sparks et al.
5,385,901 A 1/1995 Kaplan et al.
5,391,485 A 2/1995 Deeley et al.
5,393,870 A 2/1995 Deeley et al.
5,528,823 A 6/1996 Rudy, Jr. et al.
5,580,755 A 12/1996 Souza
5,591,767 A 1/1997 Mohr et al.
5,593,990 A 1/1997 D'Amato
5,629,327 A 5/1997 D'Amato
5,635,517 A 6/1997 Muller et al.

(Continued)

FOREIGN PATENT DOCUMENTS

WO WO 98/03502 1/1998

Related U.S. Application Data

(60) Provisional application No. 60/499,723, filed on Sep. 4, 2003.

(51) **Int. Cl.**
C07D 401/04 (2006.01)

(52) **U.S. Cl.** **546/200; 514/323**
(58) **Field of Classification Search** **546/200;**
514/323

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,536,809 A	10/1970	Applezweig
3,598,123 A	8/1971	Zaffaroni et al.
3,845,770 A	11/1974	Theeuwes et al.
3,916,899 A	11/1975	Theeuwes et al.
4,008,719 A	2/1977	Theeuwes et al.
4,810,643 A	3/1989	Souza
4,999,291 A	3/1991	Souza

(Continued)

OTHER PUBLICATIONS

Seddon "Pseudopolymorph: a polemic" Crystal Growth & design v.4(6) 1087, Web release Oct. 19, 2004.*


(Continued)

Primary Examiner—Celia Chang
(74) Attorney, Agent, or Firm—Jones Day

(57) **ABSTRACT**

Polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoin-dol-2-yl)-piperidine-2,6-dione are disclosed. Compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use are also disclosed.

14 Claims, 48 Drawing Sheets



US 7,465,800 B2

Page 2

U.S. PATENT DOCUMENTS

5,639,476 A	6/1997	Oshlack et al.	Halebian, J.; McCrone, W.; <i>Pharmaceutical Applications of Polymorphism</i> , Journal of Pharmaceutical Sciences, vol. 58., No. 8, Aug. 1969.
5,674,533 A	10/1997	Santus et al.	U.S. Appl. No. 60/372,348, Hariri et al.
5,698,579 A	12/1997	Muller	U.S. Appl. No. 10/732,867, D'Amato et al.
5,712,291 A	1/1998	D'Amato	U.S. Appl. No. 09/545,654, D'Amato.
5,731,325 A	3/1998	Andrusis, Jr. et al.	U.S. Appl. No. 09/287,377, D'Amato.
5,733,566 A	3/1998	Lewis	Anderson, "Moving disease biology from the laboratory to the clinic," <i>Seminars in Oncology</i> , 2002 29:17-20.
5,798,368 A	8/1998	Muller et al.	Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," <i>Seminars in Oncology</i> , 2002, 29 (6):26-33.
5,874,448 A	2/1999	Muller et al.	Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," <i>Seminars in Hematology</i> , 2003, 40 (4):1-2.
5,877,200 A	3/1999	Muller	Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor trial total therapy I (TTI) (N=231)," <i>Blood</i> , Abstract # 2857, Dec. 7-11, 2001, <i>American Society of Hematology</i> .
5,929,117 A	7/1999	Muller et al.	Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," <i>Seminars in Hematology</i> , 2003, 40 (4):33-38.
5,955,476 A	9/1999	Muller et al.	Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," <i>British Journal of Cancer</i> , 2004, 90:955-961.
6,020,358 A	2/2000	Muller et al.	Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," <i>Nature Reviews Cancer</i> , 2004, 4 (4):1-9.
6,071,948 A	6/2000	D'Amato	Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," <i>J. Mol. Med.</i> , 1995, 73:333-346.
6,114,355 A	9/2000	D'Amato	Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVd-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," <i>Blood</i> , Abstract # 2559, <i>American Society of Hematology</i> , Dec. 10-13, 2005.
6,140,346 A	10/2000	Andrusis, Jr. et al.	Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," <i>Clinical Prostate Cancer</i> , 2004, 3 (1):54-61.
6,235,756 B1	5/2001	D'Amato	Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45 th annual meeting," Press Release, Dec. 2003.
6,281,230 B1	8/2001	Muller et al.	Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.
6,316,471 B1	11/2001	Muller et al.	Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.
6,326,388 B1	12/2001	Man et al.	Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.
6,335,349 B1	1/2002	Muller et al.	Celgene Corporation, "Celgene corporation advances ACTIMID™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.
6,380,239 B1	4/2002	Muller et al.	Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.
6,395,754 B1	5/2002	Muller et al.	Celgene Corporation, "Celgene Corporation announces third quarter results. THALOMID® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for THALOMID and REVIMID™ finalized. Peer-reviewed publications of THALOMID and REVIMID data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.
6,403,613 B1	6/2002	Man et al.	Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.
6,420,414 B1	7/2002	D'Amato	Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.
6,458,810 B1	10/2002	Muller et al.	Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.
6,469,045 B1	10/2002	D'Amato	Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.
6,476,052 B1	11/2002	Muller et al.	
6,518,298 B2	2/2003	Green et al.	
6,555,554 B2	4/2003	Muller et al.	
2001/0018445 A1	8/2001	Huang et al.	
2001/0056114 A1	12/2001	D'Amato	
2002/0035090 A1	3/2002	Zeldis et al.	
2002/0045643 A1	4/2002	Muller et al.	
2002/0052398 A1	5/2002	D'Amato	
2002/0054899 A1	5/2002	Zeldis	
2002/0061923 A1	5/2002	D'Amato	
2002/0128228 A1	9/2002	Hwu	
2002/0161023 A1	10/2002	D'Amato	
2002/0173658 A1	11/2002	Muller et al.	
2002/0183360 A1	12/2002	Muller et al.	
2003/0013739 A1	1/2003	Masferrer et al.	
2003/0028028 A1	2/2003	Man et al.	
2003/0045552 A1	3/2003	Robarge et al.	
2003/0069428 A1	4/2003	Muller et al.	
2003/0096841 A1	5/2003	Robarge et al.	
2003/0139451 A1	7/2003	Shah et al.	
2003/0144325 A1	7/2003	Muller et al.	
2003/0181428 A1	9/2003	Green et al.	
2003/0187024 A1	10/2003	D'Amato	
2003/0191098 A1	10/2003	D'Amato	
2003/0235909 A1	12/2003	Hariri et al.	
2004/0029832 A1	2/2004	Zeldis	
2004/0077685 A1	4/2004	Figg et al.	
2004/0077686 A1	4/2004	Dannenberg et al.	
2004/0087546 A1	5/2004	Zeldis	
2004/0091455 A1	5/2004	Zeldis	
2004/0122052 A1	6/2004	Muller et al.	
2004/0220144 A1	11/2004	Zeldis	
2004/0266809 A1	12/2004	Emanuel et al.	

FOREIGN PATENT DOCUMENTS

WO	WO 98/54170	12/1998
WO	WO 01/70275	9/2001
WO	WO 01/87307	11/2001
WO	WO 02/059106	8/2002
WO	WO 02/064083	8/2002
WO	PCT/US03/11578	1/2003

OTHER PUBLICATIONS

Chen et al. "Polymorphic forms of . . ." CA 142:303619 (2005).*

US 7,465,800 B2

Page 3

- Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.
- Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.
- Celgene Corporation, "Celgene expands clinical development program for Revimid". Five additional trials of Revimid initiated in hematological and solid tumors cancers," Press Release, Jun. 2002.
- Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for ACTIMID™, CC-8490. SelCID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myloma," Press Release, Feb. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.
- Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead IMiD™, Revimid™," Press Release, Jun. 2001.
- Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for multiple myloma," Press Release, Feb. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.
- Chaudhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1)1884-86.
- Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.
- Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10: 1071-1073.
- D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myloma," *Semin. Oncol.* 28:597-601.
- Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl I), S25-S34.
- Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," Abstract # P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.
- Davies et al., "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract # 1284, *Circulation*, 1998.
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract # P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)", Abstract # 6, *American Society of Hematology*, Dec. 10-13, 2005.
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23)4444-4454.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma \geq 75 years of age," *Am. Soc. Hematol. 46th Ann. Meeting* Dec. 4-7, 2004, San Diego, CA Abstract #1482.
- Dredge et al., "A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer, Abstract # 491, *American Association for Cancer Research*, Apr. 6-10, 2002.
- Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.
- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.
- Glaspy et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8+ T cells in vitro," *J. Infect. Dis.* 187(6):946-955.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood*, Abstract #3219, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers, 206th American Chemical Society, Chicago, IL: Med. Chem., paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung/Drug Research*, 1981, 31 (I)941-949.
- Hernandez-Ilizaliturri et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," Abstract # 235, *American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," Abstract 1313, *American Society of Hematology*, Dec. 1-5, 2000.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidomide, in relapsed myeloma," *Blood*, Abstract # 3216, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood*, Abstract #208, *American Society of Hematology*, Dec. 4-7, 2004.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.

US 7,465,800 B2

Page 4

- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," Abstract # 3073, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," Abstract # 3456, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop*, Abstract #P225, May 4-8, 2001.
- Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology*, Abstract #927, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α - Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins¹," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Immunotherapeutic and antitumour potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues: anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4 $^+$ and CD8 $^+$ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood*, Abstract # 1548, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptic signaling induced by immunomodulatory thalidomide analogs (Imids) in human multiple myeloma cells: therapeutic implications," Abstract # 3224, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," Abstract # 3740, *American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed/refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," Abstract # 1640, *American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology*, Abstract # 992, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood*, Abstract #2487, Dec. 1-5, 2000, *American Society of Hematology*.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Payvandi et al., "Thalidomide and IMiDS inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood*, Abstract # 5046, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood*, Abstract # 2689, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," Abstract # 1793, *American Association for Cancer Research*, Mar. 24-28, 2001.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2005, 104 (13):4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13):4050-4053.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood*, Abstract # 825, *American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma: preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood*, Abstract #3225, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "A phase I trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood*, Abstract # 365, *American Society of Hematology*, Dec. 10-13, 2005.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16):3212-3214.
- Richardson et al., "Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002, 100:3063-3067, *American Society of Hematology*.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3):1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology*, Abstract #248, 2002.
- Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.

US 7,465,800 B2

Page 5

- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," Abstract # 2485, *American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streetly et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," Abstract #829, *American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," Abstract # 2543, *American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," Abstract # 367, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory- oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2):93.
- Teo et al., "Chiral inversion of the second generation IMiD™ CC-4047 (ACTIMID™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT Pace autotransplant regimen for Multiple Myeloma (MM)- safety and efficacy data in pilot study of 15 patients," *Blood*, Abstract # 2869, *American Society of Hematology*, Dec. 7-11, 2001.
- Tohnya et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-Pace results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood*, Abstract # 3531, *American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6):1887-1895.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," Abstract # PO.738, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber, "Lenalidomide (CC-5013), Revlimid™ and other ImiDs," Abstract # PL5.02, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood*, Abstract #4099, *American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology*, Abstract #3226, Dec. 1-5, 2001.
- Zangari et al., "Revimid 25 mg (REV 25) \times 20 versus 50 mg (REV 50) \times 10 q 28 days with bridging of 5 mg \times 10 versus 10 mg \times 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood*, Abstract # 1642, *American Society of Hematology*, Dec. 6-9, 2003.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience," *Blood*, Abstract # 681, *American Society of Hematology*, Dec. 7-11, 2001.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zeldis et al., "Update on the evolution of the IMiD™," *International Society for Biological Therapy of Cancer*, Oral Abstract, 2003.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," Abstract # B012, *International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- U.S. Appl. No. 60/499,723, filed Sep. 4, 2003, Jaworsky.
- U.S. Appl. No. 10/693,794, filed Oct. 23, 2003, Zeldis.
- DiMartino et al., 1997, "Preparation and physical characterization of forms II and III of paracetamol," *J. Thermal Analysis* 48:447-458.
- Jonsson, 1972, "Chemical structure and teratogenic properties. 3. A review of available data on structure-activity relationships and mechanism of action of thalidomide analogues," *Acta. Pharm. Suec.* 9(6):521-542.
- Wilén et al., 1977, "Strategies in optical resolutions," *Tetrahedron* 33:2725-2736.
- Wilén, 1972, *Tables of Resolving Agents and Optical Resolutions*, Eliel, ed., U. Notre Dame Press, Notre Dame, IN pp. 268.
- Wolff, ed., 1995, 1 Burger's Medicinal Chemistry and Drug Discovery, 5th ed., pp. 172-178 and 949-982.
- Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogs," *Ann. Rheum. Dis.* 58(Supp. 1):1107-1113.
- Grant, 1999, "Theory and Origin of Polymorphism," in *Polymorphism in Pharmaceutical Solids*, Minneapolis, MN, Ch. 1, pp. 1-10.
- He, W., et al., 1993, 206th American Chemical Society, Chicago, IL; Med. Chem., paper 216.
- Lentzsch et al., 2002, "S-3-Amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice," *Cancer Research* 62:2300-2305.
- Muller et al., 1999, "Amino-substituted thalidomide analogs: Potent inhibitors of TNF- α production," *Bioorg. Med. Chem. Lett.* 9:1625-1630.
- Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8:2669-2674.
- Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39:3238-3240.

* cited by examiner

U.S. Patent

Dec. 16, 2008

Sheet 1 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM A




Fig. 1

U.S. Patent

Dec. 16, 2008

Sheet 2 of 48

US 7,465,800 B2

IR SPECTRUM OF FORM A




Fig. 2

U.S. Patent

Dec. 16, 2008

Sheet 3 of 48

US 7,465,800 B2

RAMAN SPECTRUM OF FORM A




Fig. 3

U.S. Patent

Dec. 16, 2008

Sheet 4 of 48

US 7,465,800 B2




Fig. 4

U.S. Patent

Dec. 16, 2008

Sheet 5 of 48

US 7,465,800 B2

MOISTURE SORPTION/DESORPTION ISOTHERM FOR FORM A




Fig. 5

U.S. Patent

Dec. 16, 2008

Sheet 6 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM B




Fig. 6

U.S. Patent

Dec. 16, 2008

Sheet 7 of 48

US 7,465,800 B2

IR SPECTRUM OF FORM B




Fig. 7

U.S. Patent

Dec. 16, 2008

Sheet 8 of 48

US 7,465,800 B2

RAMAN SPECTRUM OF FORM B




Fig. 8

U.S. Patent

Dec. 16, 2008

Sheet 9 of 48

US 7,465,800 B2




Fig. 9

U.S. Patent

Dec. 16, 2008

Sheet 10 of 48

US 7,465,800 B2

TG-IR RESULTS FOR FORM B




Fig. 10

U.S. Patent

Dec. 16, 2008

Sheet 11 of 48

US 7,465,800 B2




Fig. 11

U.S. Patent

Dec. 16, 2008

Sheet 12 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM C




Fig. 12

U.S. Patent

Dec. 16, 2008

Sheet 13 of 48

US 7,465,800 B2

IR SPECTRUM OF FORM C




Fig. 13

U.S. Patent

Dec. 16, 2008

Sheet 14 of 48

US 7,465,800 B2

RAMAN SPECTRUM OF FORM C




Fig. 14

U.S. Patent

Dec. 16, 2008

Sheet 15 of 48

US 7,465,800 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM C




Fig. 15

U.S. Patent

Dec. 16, 2008

Sheet 16 of 48

US 7,465,800 B2

TG-IR RESULTS FOR FORM C




Fig. 16

U.S. Patent

Dec. 16, 2008

Sheet 17 of 48

US 7,465,800 B2




Fig. 17

U.S. Patent

Dec. 16, 2008

Sheet 18 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM D




Fig. 18

U.S. Patent

Dec. 16, 2008

Sheet 19 of 48

US 7,465,800 B2

IR SPECTRUM OF FORM D




Fig. 19

U.S. Patent

Dec. 16, 2008

Sheet 20 of 48

US 7,465,800 B2

RAMAN SPECTRUM OF FORM D




Fig. 20

U.S. Patent

Dec. 16, 2008

Sheet 21 of 48

US 7,465,800 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM D




Fig. 21

U.S. Patent

Dec. 16, 2008

Sheet 22 of 48

US 7,465,800 B2

MOISTURE SORPTION/DESORPTION ISOTHERM FOR FORM D




Fig. 22

U.S. Patent

Dec. 16, 2008

Sheet 23 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM E




Fig. 23

U.S. Patent

Dec. 16, 2008

Sheet 24 of 48

US 7,465,800 B2




Fig. 24

U.S. Patent

Dec. 16, 2008

Sheet 25 of 48

US 7,465,800 B2




Fig. 25

U.S. Patent

Dec. 16, 2008

Sheet 26 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM F




Fig. 26

U.S. Patent

Dec. 16, 2008

Sheet 27 of 48

US 7,465,800 B2

DSC THERMOGRAM FOR FORM F




Fig. 27

U.S. Patent

Dec. 16, 2008

Sheet 28 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM G




Fig. 28

U.S. Patent

Dec. 16, 2008

Sheet 29 of 48

US 7,465,800 B2

DSC THERMOGRAM FOR FORM G




Fig. 29

U.S. Patent

Dec. 16, 2008

Sheet 30 of 48

US 7,465,800 B2

XRPD PATTERN OF FORM H




Fig. 30

U.S. Patent

Dec. 16, 2008

Sheet 31 of 48

US 7,465,800 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM H




Fig. 31

U.S. Patent

Dec. 16, 2008

Sheet 32 of 48

US 7,465,800 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5274-104-B

Date: 06-04-04 16:10 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 32

File: Process 5274-100-C XRPD PATTERN OF POLYMORPH B

Date: 06-04-04 16:11 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 33

U.S. Patent

Dec. 16, 2008

Sheet 33 of 48

US 7,465,800 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5274-104-B

Date: 06-04-04 16:10 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 34

XRPD PATTERN OF POLYMORPH E

File: Process 5274-100-C

Date: 06-04-04 16:11 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 35

U.S. Patent

Dec. 16, 2008

Sheet 34 of 48

US 7,465,800 B2

XRPD PATTERN OF POLYMORPH MIXTURE




Fig. 36

U.S. Patent

Dec. 16, 2008

Sheet 35 of 48

US 7,465,800 B2

TGA CURVE OF POLYMORPH B

Sample: 5274-104-B

Size: 7.8320 mg

Method: Ramp

TGA

Run Date: 509-Jun-04 17:04

Instrument: TGA 0.500 V5.3 Build 171




Fig. 37

TGA CURVE OF POLYMORPH B

Sample: 5274-100-C

Size: 10.9430 mg

Method: Ramp

Comment: Evotec Batch 1675C H2O

TGA

Run Date: 509-Jun-04 11:20

Instrument: TGA 0.500 V5.3 Build 171




Fig. 38

U.S. Patent

Dec. 16, 2008

Sheet 36 of 48

US 7,465,800 B2




Fig. 39




Fig. 40

U.S. Patent

Dec. 16, 2008

Sheet 37 of 48

US 7,465,800 B2

TGA CURVE OF POLYMORPH MIXTURE






Fig. 41

U.S. Patent

Dec. 16, 2008

Sheet 38 of 48

US 7,465,800 B2



*Fig. 42**Fig. 43*

U.S. Patent

Dec. 16, 2008

Sheet 39 of 48

US 7,465,800 B2

*Fig. 44**Fig. 45*

U.S. Patent

Dec. 16, 2008

Sheet 40 of 48

US 7,465,800 B2




Fig. 46

U.S. Patent

Dec. 16, 2008

Sheet 41 of 48

US 7,465,800 B2




Fig. 47

U.S. Patent

Dec. 16, 2008

Sheet 42 of 48

US 7,465,800 B2

UV-VIS SCAN OF 0.04MG/ML SOLUTION

0.04mg/mL in Dissolution Medium




Fig. 48

U.S. Patent

Dec. 16, 2008

Sheet 43 of 48

US 7,465,800 B2

UV-VIS SCAN OF 0.008MG/ML SOLUTION

0.008mg/mL in Dissolution Medium




Fig. 49

U.S. Patent

Dec. 16, 2008

Sheet 44 of 48

US 7,465,800 B2

CALIBRATION CURVE FOR
3-(4-AMINO-1-OXO-1,3 DIHYDROISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE




Fig. 50

U.S. Patent

Dec. 16, 2008

Sheet 45 of 48

US 7,465,800 B2

SOLUBILITY OF FORM A




Fig. 51

U.S. Patent

Dec. 16, 2008

Sheet 46 of 48

US 7,465,800 B2




Fig. 52

U.S. Patent

Dec. 16, 2008

Sheet 47 of 48

US 7,465,800 B2




FIG. 53

U.S. Patent

Dec. 16, 2008

Sheet 48 of 48

US 7,465,800 B2

INTRINSIC DISSOLUTION OF FORMS A, B, AND E (EXPERIMENT NO.2)




FIG. 54

US 7,465,800 B2

1

**POLYMORPHIC FORMS OF
3-(4-AMINO-1-OXO-1,3
DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-2,6-
IONE**

This application claims the benefit of U.S. provisional application 60/499,723, filed Sep. 4, 2003, the contents of which are incorporated by reference herein their entirety.

1. FIELD OF THE INVENTION

This invention relates to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use for the treatment of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer.

2. BACKGROUND OF THE INVENTION

Many compounds can exist in different crystal forms, or polymorphs, which exhibit different physical, chemical, and spectroscopic properties. For example, certain polymorphs of a compound may be more readily soluble in particular solvents, may flow more readily, or may compress more easily than others. See, e.g., P. DiMartino, et al., *J. Thermal Anal.*, 48:447-458 (1997). In the case of drugs, certain solid forms may be more bioavailable than others, while others may be more stable under certain manufacturing, storage, and biological conditions. This is particularly important from a regulatory standpoint, since drugs are approved by agencies such as the U.S. Food and Drug Administration only if they meet exacting purity and characterization standards. Indeed, the regulatory approval of one polymorph of a compound, which exhibits certain solubility and physico-chemical (including spectroscopic) properties, typically does not imply the ready approval of other polymorphs of that same compound.

Polymorphic forms of a compound are known in the pharmaceutical arts to affect, for example, the solubility, stability, flowability, fractability, and compressibility of the compound, as well as the safety and efficacy of drug products comprising it. See, e.g., Knapman, K. *Modern Drug Discoveries*, 2000, 53. Therefore, the discovery of new polymorphs of a drug can provide a variety of advantages.

U.S. Pat. Nos. 5,635,517 and 6,281,230, both to Muller et al., disclose 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which is useful in treating and preventing a wide range of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer. New polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can further the development of formulations for the treatment of these chronic illnesses, and may yield numerous formulation, manufacturing and therapeutic benefits.

3. SUMMARY OF THE INVENTION

This invention encompasses polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. In certain aspects, the invention provides polymorphs of the compound identified herein as forms A, B, C, D, E, F, G, and H. The invention also encompasses mixtures of these forms. In further embodiments, this invention provides methods of making, isolating and characterizing the polymorphs.

This invention also provides pharmaceutical compositions and single unit dosage forms comprising a polymorph of

2

3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The invention further provides methods for the treatment or prevention of a variety of diseases and disorders, which comprise administering to a patient in need of such treatment or prevention a therapeutically effective amount of a polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

4. BRIEF DESCRIPTION OF THE DRAWINGS

- 10 Specific aspects of the invention can be understood with reference to the attached figures:
- FIG. 1 provides a representative X-ray powder diffraction (XRPD) pattern of Form A;
- 15 FIG. 2 provides a representative IR spectrum of Form A;
- FIG. 3 provides a representative Raman spectrum of Form A;
- 20 FIG. 4 provides a representative thermogravimetric analysis (TGA) curve and a representative differential scanning calorimeter (DSC) thermogram of Form A;
- FIG. 5 provides a representative moisture sorption/desorption isotherm of Form A;
- 25 FIG. 6 provides a representative XRPD pattern of Form B;
- FIG. 7 provides a representative IR spectrum of Form B;
- FIG. 8 provides a representative Raman spectrum of Form B;
- FIG. 9 provides a representative TGA curve and a representative DSC thermogram of Form B;
- 30 FIG. 10 provides representative TG-IR results of Form B;
- FIG. 11 provides a representative moisture sorption/desorption isotherm of Form B;
- FIG. 12 provides a representative XRPD pattern of Form C;
- FIG. 13 provides a representative IR spectrum of Form C;
- 35 FIG. 14 provides a representative Raman spectrum of Form C;
- FIG. 15 provides a representative TGA curve and a representative DSC thermogram of Form C;
- 40 FIG. 16 provides representative TG-IR results of Form C;
- FIG. 17 provides a representative moisture sorption/desorption isotherm of Form C;
- FIG. 18 provides a representative XRPD pattern of Form D;
- 45 FIG. 19 provides a representative IR spectrum of Form D;
- FIG. 20 provides a representative Raman spectrum of Form D;
- FIG. 21 provides a representative TGA curve and a representative DSC thermogram of Form D;
- 50 FIG. 22 provides a representative moisture sorption/desorption isotherm of Form D;
- FIG. 23 provides a representative XRPD pattern of Form E;
- FIG. 24 provides a representative TGA curve and a representative DSC thermogram of Form E;
- 55 FIG. 25 provides a representative moisture sorption/desorption isotherm of Form E;
- FIG. 26 provides a representative XRPD pattern for a sample of Form F;
- FIG. 27 provides a representative thermogram of Form F;
- 60 FIG. 28 provides a representative XRPD pattern of Form G;
- FIG. 29 provides a representative DSC thermogram for a sample of Form G;
- FIG. 30 provides a representative XRPD pattern of Form H;
- 65 FIG. 31 provides a representative TGA curve and a representative DSC thermogram of Form H;

US 7,465,800 B2

3

FIG. 32 provides a representative XRPD pattern of Form B;

FIG. 33 provides a representative XRPD pattern of Form B;

FIG. 34 provides a representative XRPD pattern of Form B;

FIG. 35 provides a representative XRPD pattern of Form E;

FIG. 36 provides a representative XRPD pattern of polymorph mixture;

FIG. 37 provides a representative TGA curve of Form B;

FIG. 38 provides a representative TGA curve of Form B;

FIG. 39 provides a representative TGA curve of Form B;

FIG. 40 provides a representative TGA curve of Form E;

FIG. 41 provides a representative TGA curve of polymorph mixture;

FIG. 42 provides a representative DSC thermogram of Form B;

FIG. 43 provides a representative DSC thermogram of Form B;

FIG. 44 provides a representative DSC thermogram of Form B;

FIG. 45 provides a representative DSC thermogram of Form E;

FIG. 46 provides a representative DSC thermogram of polymorph mixture;

FIG. 47 provides a UV-Vis scan of dissolution medium;

FIG. 48 provides a UV-Vis scan of 0.04 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 49 provides a UV-Vis scan of 0.008 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 50 provides a calibration curve for 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione;

FIG. 51 provides a solubility curve of Form A;

FIG. 52 provides a solubility curve of Form B;

FIG. 53 provides an intrinsic dissolution of Forms A, B and E; and

FIG. 54 provides an intrinsic dissolution of Forms A, B and E.

5. DETAILED DESCRIPTION OF THE INVENTION

5.1 Definitions

As used herein and unless otherwise indicated, the terms “treat,” “treating” and “treatment” refer to the alleviation of a disease or disorder and/or at least one of its attendant symptoms.

As used herein and unless otherwise indicated, the terms “prevent,” “preventing” and “prevention” refer to the inhibition of a symptom of a disease or disorder or the disease itself.

As used herein and unless otherwise indicated, the terms “polymorph” and “polymorphic form” refer to solid crystalline forms of a compound or complex. Different polymorphs of the same compound can exhibit different physical, chemical and/or spectroscopic properties. Different physical properties include, but are not limited to stability (e.g., to heat or light), compressibility and density (important in formulation and product manufacturing), and dissolution rates (which can affect bioavailability). Differences in stability can result from changes in chemical reactivity (e.g., differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical characteristics (e.g., tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both

4

(e.g., tablets of one polymorph are more susceptible to breakdown at high humidity). Different physical properties of polymorphs can affect their processing. For example, one polymorph might be more likely to form solvates or might be more difficult to filter or wash free of impurities than another due to, for example, the shape or size distribution of particles of it.

Polymorphs of a molecule can be obtained by a number of methods known in the art. Such methods include, but are not

10 limited to, melt recrystallization, melt cooling, solvent recrystallization, desolvation, rapid evaporation, rapid cooling, slow cooling, vapor diffusion and sublimation. Polymorphs can be detected, identified, classified and characterized using well-known techniques such as, but not limited to,


15 differential scanning calorimetry (DSC), thermogravimetry (TGA), X-ray powder diffractometry (XRPD), single crystal X-ray diffractometry, vibrational spectroscopy, solution calorimetry, solid state nuclear magnetic resonance (NMR), infrared (IR) spectroscopy, Raman spectroscopy, hot stage optical microscopy, scanning electron microscopy (SEM), electron crystallography and quantitative analysis, particle size analysis (PSA), surface area analysis, solubility, and rate of dissolution.

As used herein to refer to the spectra or data presented in 25 graphical form (e.g., XRPD, IR, Raman and NMR spectra), and unless otherwise indicated, the term “peak” refers to a peak or other special feature that one skilled in the art would recognize as not attributable to background noise. The term “significant peaks” refers to peaks at least the median size (e.g., height) of other peaks in the spectrum or data, or at least 1.5, 2, or 2.5 times the median size of other peaks in the spectrum or data.

As used herein and unless otherwise indicated, the term “substantially pure” when used to describe a polymorph of a 35 compound means a solid form of the compound that comprises that polymorph and is substantially free of other polymorphs of the compound. A representative substantially pure polymorph comprises greater than about 80% by weight of one polymorphic form of the compound and less than about 20% by weight of other polymorphic forms of the compound, more preferably greater than about 90% by weight of one polymorphic form of the compound and less than about 10% by weight of the other polymorphic forms of the compound, even more preferably greater than about 95% by weight of one polymorphic form of the compound and less than about 5% by weight of the other polymorphic forms of the compound, and most preferably greater than about 97% by weight of one polymorphic form of the compound and less than about 3% by weight of the other polymorphic forms of the compound.

5.2 Polymorphic Forms

This invention is directed to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which has the structure shown below:



60
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
2510
2511
2512
2513
2514
2515
2516
2517
2518
2519
2520
2521
2522
2523
2524
2525
2526
2527
2528
2529
2530
2531
2532
2533
2534
2535
2536
2537
2538
2539
2540
2541
2542
2543
2544
2545
2546
2547
2548
2549
2550
2551
2552
2553
2554
2555
2556
2557
2558
2559
2560
2561
2562
2563
2564
2565
2566
2567
2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582
2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595
2596
2597
2598
2599
25100
25101
25102
25103
25104
25105
25106
25107
25108
25109
25110
25111
25112
25113
25114
25115
25116
25117
25118
25119
25120
25121
25122
25123
25124
25125
25126
25127
25128
25129
25130
25131
25132
25133
25134
25135
25136
25137
25138
25139
25140
25141
25142
25143
25144
25145
25146
25147
25148
25149
25150
25151
25152
25153
25154
25155
25156
25157
25158
25159
25160
25161
25162
25163
25164
25165
25166
25167
25168
25169
25170
25171
25172
25173
25174
25175
25176
25177
25178
25179
25180
25181
25182
25183
25184
25185
25186
25187
25188
25189
25190
25191
25192
25193
25194
25195
25196
25197
25198
25199
25200
25201
25202
25203
25204
25205
25206
25207
25208
25209
25210
25211
25212
25213
25214
25215
25216
25217
25218
25219
25220
25221
25222
25223
25224
25225
25226
25227
25228
25229
25230
25231
25232
25233
25234
25235
25236
25237
25238
25239
25240
25241
25242
25243
25244
25245
25246
25247
25248
25249
25250
25251
25252
25253
25254
25255
25256
25257
25258
25259
25260
25261
25262
25263
25264
25265
25266
25267
25268
25269
25270
25271
25272
25273
25274
25275
25276
25277
25278
25279
25280
25281
25282
25283
25284
25285
25286
25287
25288
25289
25290
25291
25292
25293
25294
25295
25296
25297
25298
25299
252100
252101
252102
252103
252104
252105
252106
252107
252108
252109
252110
252111
252112
252113
252114
252115
252116
252117
252118
252119
252120
252121
252122
252123
252124
252125
252126
252127
252128
252129
252130
252131
252132
252133
252134
252135
252136
252137
252138
252139
252140
252141
252142
252143
252144
252145
252146
252147
252148
252149
252150
252151
252152
252153
252154
252155
252156
252157
252158
252159
252160
252161
252162
252163
252164
252165
252166
252167
252168
252169
252170
252171
252172
252173
252174
252175
252176
252177
252178
252179
252180
252181
252182
252183
252184
252185
252186
252187
252188
252189
252190
252191
252192
252193
252194
252195
252196
252197
252198
252199
252200
252201
252202
252203
252204
252205
252206
252207
252208
252209
252210
252211
252212
252213
252214
252215
252216
252217
252218
252219
252220
252221
252222
252223
252224
252225
252226
252227
252228
252229
252230
252231
252232
252233
252234
252235
252236
252237
252238
252239
252240
252241
252242
252243
252244
252245
252246
252247
252248
252249
252250
252251
252252
252253
252254
252255
252256
252257
252258
252259
252260
252261
252262
252263
252264
252265
252266
252267
252268
252269
252270
252271
252272
252273
252274
252275
252276
252277
252278
252279
252280
252281
252282
252283
252284
252285
252286
252287
252288
252289
252290
252291
252292
252293
252294
252295
252296
252297
252298
252299
252300
252301
252302
252303
252304
252305
252306
252307
252308
252309
252310
252311
252312
252313
252314
252315
252316
252317
252318
252319
252320
252321
252322
252323
252324
252325
252326
252327
252328
252329
252330
252331
252332
252333
252334
252335
252336
252337
252338
252339
252340
252341
252342
252343
252344
252345
252346
252347
252348
252349
252350
252351
252352
252353
252354
252355
252356
252357
252358
252359
252360
252361
252362
252363
252364
252365
252366
252367
252368
252369
252370
252371
252372
252373
252374
252375
252376
252377
252378
252379
252380
252381
252382
252383
252384
252385
252386
252387
252388
252389
252390
252391
252392
252393
252394
252395
252396
252397
252398
252399
252400
252401
252402
252403
252404
252405
252406
252407
252408
252409
252410
252411
252412
252413
252414
252415
252416
252417
252418
252419
252420
252421
252422
252423
252424
252425
252426
252427
252428
252429
252430
252431
252432
252433
252434
252435
252436
252437
252438
252439
252440
252441
252442
252443
252444
252445
252446
252447
252448
252449
252450
252451
252452
252453
252454
252455
252456
252457
252458
252459
252460
252461
252462
252463
252464
252465
252466
252467
252468
252469
252470
252471
252472
252473
252474
252475
252476
252477
252478
252479
252480
252481
252482
252483
252484
252485
252486
252487
252488
252489
252490
252491
252492
252493
252494
252495
252496
252497
252498
252499
252500
252501
252502
252503
252504
252505
252506
252507
252508
252509
252510
252511
252512
252513
252514
252515
252516
252517
252518
252519
252520
252521
252522
252523
252524
252525
252526
252527
252528
252529
252530
252531
252532
252533
252534
252535
252536
252537
252538
252539
252540
252541
252542
252543
252544
252545
252546
252547
252548
252549
252550
252551
252552
252553
252554
252555
252556
252557
252558
252559
252560
252561
252562
252563
252564
252565
252566
252567
252568
252569
252570
252571
252572
252573
252574
252575
252576
252577
252578
252579
252580
252581
252582
252583
252584
252585
252586
252587
252588
252589
252590
252591
252592
252593
252594
252595
252596
252597
252598
252599
252600
252601
252602
252603
252604
252605
252606
252607
252608
252609
252610
252611
252612
252613
252614
252615
252616
252617
252618
252619
252620
252621
252622
252623
252624
252625
252626
252627
252628
252629
252630
252631
252632
252633
252634
252635
252636
252637
252638
252639
252640
252641
252642
252643
252644
252645
252646
252647
252648
252649
252650
252651
252652
252653
252654
252655
252656
252657
252658
252659
252660
252661
252662
252663
252664
252665
252666
252667
252668
252669
252670
252671
252672
252673
252674
252675
252676
252677
252678
252679
252680
252681
252682
252683
252684
252685
252686
252687
252688
252689
252690
252691
252692
252693
252694
252695
252696
252697
252698
252699
252700
252701
252702
252703
252704
252705
252706
252707
252708
252709
252710
252711
252712
252713
252714
252715
252716
252717
252718
252719
252720
252721
252722
252723
252724
252725
252726
252727
252728
252729
252730
252731
252732
252733
252734
252735
252736
252737
252738
252739
252740
252741
252742
252743
252744
252745
252746
252747
252748
252749
252750
252751
252752
252753
252754
252755
252756
252757
252758
252759
252760
252761
252762
252763
252764
252765
252766
252767
252768
252769
252770
252771
252772
252773
252774
252775
252776
252777
252778
252779
252780
252781
252782
252783
252784
252785<br

US 7,465,800 B2

5

entireties of which are incorporated herein by reference. For example, the compound can be prepared through catalytic hydrogenation of 3-(4-nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. 3-(4-Nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by allowing 2,6-dioxopiperidin-3-ammonium chloride to react with methyl 2-bromomethyl-4-nitrobenzoate in dimethylformamide in the presence of triethylamine. The methyl 2-bromomethyl-4-nitrobenzoate in turn is obtained from the corresponding methyl ester of nitro-ortho-toluic acid by conventional bromination with N-bromosuccinimide under the influence of light.

Polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by techniques known in the art, including solvent recrystallization, desolvation, vapor diffusion, rapid evaporation, slow evaporation, rapid cooling and slow cooling. Polymorphs can be made by dissolving a weighed quantity of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in various solvents at elevated temperatures. The solutions of the compound can then be filtered and allowed to evaporate either in an open vial (for fast hot evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation). Polymorphs can also be obtained from slurries. Polymorphs can be crystallized from solutions or slurries using several methods. For example, a solution created at an elevated temperature (e.g., 60° C.) can be filtered quickly then allowed to cool to room temperature. Once at room temperature, the sample that did not crystallize can be moved to a refrigerator then filtered. Alternatively, the solutions can be crash cooled by dissolving the solid in a solvent at an increased temperature (e.g., 45-65° C.) followed by cooling in a dry ice/solvent bath.

One embodiment of the invention encompasses Form A of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form A is an unsolvated, crystalline material that can be obtained from non-aqueous solvent systems. Another embodiment of the invention encompasses Form B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form B is a hemihydrated, crystalline material that can be obtained from various solvent systems. Another embodiment of the invention encompasses Form C of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form C is a hemisolvated crystalline material that can be obtained from solvents such as, but not limited to, acetone. Another embodiment of the invention encompasses Form D of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form D is a crystalline, solvated polymorph prepared from a mixture of acetonitrile and water. Another embodiment of the invention encompasses Form E of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form E is a dihydrated, crystalline material. Another embodiment of the invention encompasses Form F of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form F is an unsolvated, crystalline material that can be obtained from the dehydration of Form E. Another embodiment of the invention encompasses Form G of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form G is an unsolvated, crystalline material that can be obtained from slurring forms B and E in a solvent such as, but not limited to, tetrahydrofuran (THF). Another embodiment of the invention encompasses Form H of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form H is a partially hydrated crystalline material that can be obtained by exposing Form E to 0% relative humidity. Each of these forms is discussed in detail below.

Another embodiment of the invention encompasses a composition comprising amorphous 3-(4-amino-1-oxo-1,3 dihy-

6

dro-isoindol-2-yl)-piperidine-2,6-dione and crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione of form A, B, C, D, E, F, G or H. Specific compositions can comprise greater than about 50, 75, 90 or 95 weight percent crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Another embodiment of the invention encompasses a composition comprising at least two crystalline forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (e.g., a mixture of polymorph forms B and E).

5.2.1 Form A

The data described herein for Form A, as well as for Forms B-H, were obtained using the experimental methods described in Examples 6.3-6.7, provided below.

Form A can be obtained from various solvents, including, but not limited to 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and THF. FIG. 1 shows a representative XRPD pattern of Form A. The pattern is characterized by peaks, preferably significant peaks, at approximately 8, 14.5, 16, 17.5, 20.5, 24, and 26 degrees 2θ. Representative IR and Raman spectra data are provided in FIGS. 2 and 3.

Representative thermal characteristics of Form A are shown in FIG. 4. TGA data show a small weight increase up to about 150° C., indicating an unsolvated material. Weight loss above 150° C. is attributed to decomposition. The DSC curve of Form A exhibits an endotherm at about 270° C.

Representative moisture sorption and desorption data are plotted in FIG. 5. Form A does not exhibit a significant weight gain from 5 to 95% relative humidity. Equilibrium can be obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it has typically lost only about 0.003% by weight from start to finish. Form A is capable of remaining a crystalline solid for about 11 days when stored at about 22, 45, 58, and 84% relative humidity.

Interconversion studies show that Form A can convert to Form B in aqueous solvent systems and can convert to Form C in acetone solvent systems. Form A tends to be stable in anhydrous solvent systems. In water systems and in the presence of Form E, Form A tends to convert to Form E.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% relative humidity (RH) and 40° C./93% RH), Form A typically does not convert to a different form.

In sum, Form A is a crystalline, unsolvated solid that melts at approximately 270° C. Form A is weakly or not hygroscopic and appears to be the most thermodynamically stable anhydrous polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione discovered thus far.

5.2.2 Form B

Form B can be obtained from many solvents, including, but not limited to, hexane, toluene, and water. FIG. 6 shows a representative XRPD pattern of Form B, characterized by peaks at approximately 16, 18, 22 and 27 degrees 2θ.

Solution proton NMR confirm that Form B is a form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Representative IR and Raman spectra are shown in FIGS. 7 and 8, respectively. Compared to Form A, the IR spectrum for Form B has peaks at approximately 3513 and 1960 cm⁻¹.

Representative DSC and TGA data for Form B are shown in FIG. 9. The DSC curve exhibits endotherms at about 146 and 268° C. These events are identified as dehydration and melting by hot stage microscopy experiments. Form B typi-

US 7,465,800 B2

7

cally loses about 3.1% volatiles up to about 175° C. (per approximately 0.46 moles of water). Comparison of the IR spectrum of the volatiles with that of water indicates that they are water (See FIG. 10). Calculations from TGA data indicate that Form B is a hemihydrate. Karl Fischer water analysis also supports this conclusion.

Representative moisture sorption and desorption data are shown in FIG. 11. Form B typically does not exhibit a significant weight gain from 5% to 95% relative humidity, when equilibrium is obtained at each relative humidity step. As Form B dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it typically has gained only about 0.022% by weight (about 0.003 mg) from start to finish. Form B does not convert to a different form upon exposure to about 84% relative humidity for about ten days.

Interconversion studies show that Form B typically converts to Form A in a THF solvent system, and typically converts to Form C in an acetone solvent system. In aqueous solvent systems such as pure water and 10% water solutions, Form B is the most stable of the polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. However, it can convert to Form E in the presence of water. Desolvation experiments show that upon heating at about 175° C. for about five minutes, Form B typically converts to Form A.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH), Form B does not convert to a different form.

In sum, Form B is a hemihydrated, crystalline solid which has a DSC thermogram exhibiting endotherms at about 146 and about 268° C. Interconversion studies show that Form B converts to Form E in aqueous solvent systems, and converts to other forms in acetone and other anhydrous systems.

5.2.3 Form C

Form C can be obtained from evaporation, slurries and slow cools in acetone solvent systems. A representative XRPD pattern of this form is shown in FIG. 12. The data are characterized by peaks at approximately 15.5 and 25 degrees 20.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 13 and 14, respectively. The IR spectrum of Form C is characterized by peaks at approximately 3466, 3373, and 3318 cm⁻¹. The Raman spectrum of Form C is characterized by peaks at about 3366, 3321, 1101, and 595 cm⁻¹.

Representative thermal characteristics for Form C are plotted in FIG. 15. Form C loses about 10.02% volatiles up to about 175° C., indicating it is a solvated material. Weight loss above about 175° C. is attributed to decomposition. Identification of volatiles in Form C can be accomplished with TG-IR experiments. The representative IR spectrum captured after several minutes of heating, as depicted in FIG. 13, when compared with a spectral library, shows acetone to be the best match. Calculations from TGA data show that Form C is a hemisolvate (approximately 0.497 moles of acetone). The DSC curve for Form C, shown in FIG. 15, exhibits endotherms at about 150 and about 269° C. The endotherm at about 150° C. is attributed to solvent loss based on observations made during hot stage microscopy experiments. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption balance data are shown in FIG. 17. Form C does not exhibit a significant weight gain from 5 to 85% relative humidity, when

8

equilibrium is obtained at each relative humidity step up to 85% relative humidity. At 95% relative humidity, Form C experiences a significant weight loss of about 6.03%. As the sample dries from 95% back down to 5% relative humidity, the sample maintains the weight achieved at the end of the adsorption phase at each step down to 5% relative humidity. Form C is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form C typically converts to Form A in a THF solvent system and typically converts to Form E in an aqueous solvent system. In an acetone solvent system, Form C is the most stable form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Desolvation experiments performed on Form C show that upon heating at about 150° C. for about five minutes, Form C will typically convert to Form A.

In sum, Form C is a crystalline, hemisolvated solid, which has a DSC thermogram exhibiting endotherms at about 150 and about 269° C. Form C is not hygroscopic below about 85% RH, but can convert to Form B at higher relative humidities.

5.2.4 Form D

Form D can be obtained from evaporation in acetonitrile solvent systems. A representative XRPD pattern of the form is shown in FIG. 18. The pattern is characterized by peaks at approximately 27 and 28 degrees 20.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 19 and 20, respectively. The IR spectrum of Form D is characterized by peaks at approximately 3509, 2299, and 2256 cm⁻¹. The Raman spectrum of Form D is characterized by peaks at approximately 2943, 2889, 2297, 2260, 1646, and 1150 cm⁻¹.

Representative thermal characteristics for Form D are plotted in FIG. 21. Form D loses about 6.75% volatiles up to about 175° C., indicating a solvated material. Weight loss above about 175° C. is attributed to decomposition. TG-IR experiments indicate that the volatiles are water and acetonitrile. Calculations from TG data show that about one mole of water is present in the sample. A representative DSC curve for Form D exhibits endotherms at about 122 and about 270° C. The endotherm at about 122° C. is attributed to loss of volatiles based on observations made during hot stage microscopy experiments. The endotherm at about 270° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 22. Form D does not exhibit a significant weight gain from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it maintains its weight such that at 5% relative humidity the form has typically gained only about 0.39% by weight (about 0.012 mg) from start to finish. Form A is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form D is capable of converting to Form A in a THF solvent system, to Form E in an aqueous solvent system, and to Form C in an acetone solvent system. Desolvation experiments performed on Form D show that upon heating at about 150° C. for about five minutes Form D will typically convert to Form A.

In sum, Form D is a crystalline solid, solvated with both water and acetonitrile, which has a DSC thermogram exhibiting endotherms at about 122 and about 270° C. Form D is either weakly or not hygroscopic, but will typically convert to Form B when stressed at higher relative humidities.

US 7,465,800 B2

9

10

5.2.5 Form E

Form E can be obtained by slurring 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in water and by a slow evaporation of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in a solvent system with a ratio of about 9:1 acetone:water. A representative XRPD pattern is shown in FIG. 23. The data are characterized by peaks at approximately 20, 24.5 and 29 degrees 2θ.

Representative thermal characteristics of Form E are plotted in FIG. 24. Form E typically loses about 10.58% volatiles up to about 125° C., indicating that it is a solvated material. A second weight loss of an additional about 1.38% was observed between about 125° C. and about 175° C. Weight loss above about 175° C. is attributed to decomposition. Karl Fischer and TG-IR experiments support the conclusion that the volatile weight loss in Form E is due to water. The representative DSC curve for Form E exhibits endotherms at about 99, 161 and 269° C. Based on observations made during hot stage microscopy experiments, the endotherms at about 99 and about 161° C. are attributed to loss of volatiles. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 25. Form E typically does not exhibit a significant weight change from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the sample dried from 95% back down to 5% relative humidity, the sample continues to maintain weight such that at 5% relative humidity the sample has lost only about 0.0528% by weight from start to finish.

Interconversion studies show that Form E can convert to Form C in an acetone solvent system and to Form G in a THF solvent system. In aqueous solvent systems, Form E appears to be the most stable form. Desolvation experiments performed on Form E show that upon heating at about 125° C. for about five minutes, Form E can convert to Form B. Upon heating at 175° C. for about five minutes, Form B can convert to Form F.

When stored for a period of 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH) Form E typically does not convert to a different form. When stored for seven days at room temperature/0% RH, Form E can convert to a new form, Form H.

5.2.6 Form F

Form F can be obtained by complete dehydration of Form E. A representative XRPD pattern of Form F, shown in FIG. 26, is characterized by peaks at approximately 19, 19.5 and 25 degrees 2θ.

Representative thermal characteristics of Form F are shown in FIG. 27. The representative DSC curve for Form F exhibits an endotherm at about 269° C. preceded directly by two smaller endotherms indicative of a crystallized form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The DSC thermogram does not show any thermal events prior to the melt, suggesting that it is an unsolvated material.

5.2.7 Form G

Form G can be obtained by slurring forms B and E in THF. A representative XRPD pattern of this form, shown in FIG. 28, is characterized by a peak at approximately 23 degrees 2θ. Two other peaks unique to Form G appear at approximately 21 and 24.5 degrees 2θ.

Representative thermal characteristics of Form G are plotted in FIG. 29. A representative DSC curve for Form G exhibits an endotherm at about 248° C. followed by a small, broad exotherm at about 267° C. No thermal events are seen

in the DSC thermogram at lower temperatures, suggesting that it is an unsolvated material.

5.2.8 Form H

Form H can be obtained by storing Form E at room temperature and 0% RH for about 7 days. A representative XRPD pattern is shown in FIG. 30. The pattern is characterized by a peak at 15 degrees 2θ, and two other peaks at 26 and 31 degrees 2θ.

Representative thermal characteristics are shown in FIG. 31. Form H loses about 1.67% volatiles up to about 150° C. Weight loss above about 150° C. is attributed to decomposition. Karl Fischer data shows that Form H typically contains about 1.77% water (about 0.26 moles), suggesting that the weight loss seen in the TG is due to dehydration. The DSC thermogram shows a broad endotherm between about 50° C. and about 125° C., corresponding to the dehydration of Form H and a sharp endotherm at about 269° C., which is likely due to a melt.

When slurried in water with either Forms A or B, after about 14 days Form H can convert to Form E. When slurried in THF, Form H can convert to Form A. When slurried in acetone, Form H can convert to Form C.

In sum, Form H is a crystalline solid, hydrated with about 0.25 moles of water, which has a DSC thermogram exhibiting an endotherm between about 50 and 125° C. and an endotherm at about 269° C.

5.3 Methods of Use and Pharmaceutical Compositions

Polymorphs of the invention exhibit physical characteristics that are beneficial for drug manufacture, storage or use. All polymorphs of the invention have utility as pharmaceutically active ingredients or intermediates thereof.

This invention encompasses methods of treating and preventing a wide variety of diseases and conditions using polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. In each of the methods, a therapeutically or prophylactically effective amount of the compound is administered to a patient in need of such treatment or prevention. Examples of such disease and conditions include, but are not limited to, diseases associated with undesired angiogenesis, cancer (e.g., solid and blood borne tumors), inflammatory diseases, autoimmune diseases, and immune diseases. Examples of cancers and pre-cancerous conditions include those described in U.S. Pat. Nos. 6,281,230 and 5,635,517 to Muller et al. and in various U.S. patent applications to Zeldis, including application Ser. No. 10/411,649, filed Apr. 11, 2003 (Treatment of Myelodysplastic Syndrome); Ser. No. 10/438,213 filed May 15, 2003 (Treatment of Various Types of Cancer); Ser. No. 10/411,656, filed Apr. 11, 2003 (Treatment of Myeloproliferative Diseases). Examples of other diseases and disorders that can be treated or prevented using compositions of the invention are described in U.S. Pat. Nos. 6,235,756 and 6,114,335 to D'Amato and in other U.S. patent applications to Zeldis, including Ser. No. 10/693,794, filed Oct. 23, 2003 (Treatment of Pain Syndrome) and Ser. No. 10/699,154, filed Oct. 30, 2003 (Treatment of Macular Degeneration). The entirety of each of the patents and patent applications cited herein is incorporated herein by reference.

Depending on the disease to be treated and the subject's condition, polymorphs of the invention can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implantation), inhalation spray, nasal, vaginal, rectal, sublingual, or topical routes of administration and may be formulated, alone or together, in suitable dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of administration. Because individual poly-

US 7,465,800 B2

11

morphs have different dissolution, stability, and other properties, the optimal polymorph used in methods of treatment may depend on the route of administration. For example, forms that are readily soluble in aqueous solutions are preferably used to provide liquid dosage forms, whereas forms that exhibit great thermal stability may be preferred in the manufacture of solid dosage forms (e.g., tablets and capsules).

Although the physical characteristics of polymorphs can, in some cases, affect their bioavailability, amounts of the polymorphs that are therapeutically or prophylactically effective in the treatment of various disease and conditions can be readily determined by those of ordinary skill in the pharmacy or medical arts. In certain embodiments of the invention, a polymorph is administered orally and in a single or divided daily doses in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day. In other embodiments, a polymorph is administered every other day in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day.

The invention encompasses pharmaceutical compositions and single unit dosage forms that can be used in methods of treatment and prevention, which comprise one or more polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and optionally one or more excipients or diluents. Specific compositions and dosage forms are disclosed in the various patents and patent applications incorporated herein by reference. In one embodiment, a single dosage form comprises a polymorph (e.g., Form B) in an amount of about 5, 10, 25 or 50 mg.

6. EXAMPLES**6.1 Polymorph Screen**

A polymorph screen to generate the different solid forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was carried out as follows.

A weighed sample of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (usually about 10 mg) was treated with aliquots of the test solvent. Solvents were either reagent or HPLC grade. The aliquots were usually about 200 µL. Between additions, the mixture was usually shaken or sonicated. When the solids dissolved, as judged by visual inspection, estimated solubilities were calculated. Solubilities were estimated from these experiments based on the total solvent used to provide a solution. Actual solubilities may have been greater than those calculated due to the use of too-large solvent aliquots or to a slow rate of dissolution.

Samples were created by generating solutions (usually about 30 mg in 20 mL) at elevated temperatures, filtering, and allowing the solution to evaporate whether in an open vial (hot fast evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation).

Slurry experiments were also performed. Usually about 25 mg of solid was placed in either 3 or 5 mL of solvent. The samples were then placed on orbital shakers at either ambient temperature or 40° C. for 4-10 days.

Crystallizations were performed using various cooling methods. Solid was dissolved in a solvent at an elevated temperature (e.g., about 60° C.), filtered quickly and allowed to cool to room temperature. Once at room temperature, samples that did not crystallize were moved to a refrigerator. Solids were removed by filtration or decantation and allowed to dry in the air. Crash cools were performed by dissolving solid in a solvent at an increased temperature (e.g., about 45-65° C.) followed by cooling in a dry ice/acetone bath.

12

Hygroscopicity studies were performed by placing portions of each polymorph in an 84% relative humidity chamber for approximately one week.

Desolvation studies were carried out by heating each polymorph in a 70° C. oven for approximately one week.

Interconversion experiments were carried out by making slurries containing two forms in a saturated solvent. The slurries were agitated for approximately 7-20 days at ambient temperature. The insoluble solids were recovered by filtration and analyzed using XRPD.

6.2 Preparation of Polymorphic Forms

Eight solid forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione were prepared as described below.

Form A was obtained by crystallization from various non-aqueous solvents including 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and tetrahydrofuran. Form B was also obtained by crystallization from the solvents hexane, toluene and water. Form C was obtained from evaporation, slurries, and slow cools in acetone solvent systems. Form D was obtained from evaporation in acetonitrile solvent systems. Form E was obtained most readily by slurring 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in water. Form F was obtained by complete desolvation of Form E. It is found to be an unsolvated, crystalline material that melts at about 269° C. Form G was obtained by slurring forms B and E in THF. Form H was obtained by stressing Form E at room temperature and 0% RH for 7 days.

6.2.1 Synthesis of Polymorphs B and E

Form B is the desired polymorph for the active pharmaceutical ingredient (API) of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. This form has been used in the formulation of API into drug product for clinical studies. Three batches were produced as apparent mixtures of polymorphs in the non-micronized API of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Development work was carried out to define a process that would generate polymorph B from this mixture of polymorphs and could be implemented for strict polymorphic controls in the validation batches and future manufacturing of API of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Characterization of polymorphic forms produced during the work was performed by XRPD, DSC, TGA and KF.

A process was also developed for the large-scale preparation of Form E. Polymorph E material was prepared in order to carry out a comparison with polymorph B drug product in capsule dissolution testing of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. 150 g of a mixture of polymorphs in 3 L of water was stirred at room temperature for 48 hours. The product was collected by filtration and dried at 25° C. for 24 hours under vacuum. XRPD, DSC, TGA, KF and HPLC analyses confirmed that the material isolated was polymorph E.

In a preliminary work, it was demonstrated that stirring a suspension of a mixture of polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione with water at high temperature (75° C.) for an extended period of time converted this mixture of polymorphs exclusively to form B. Several specific parameters were identified including temperature, solvent volume and drying parameters (temperature and vacuum). XRPD, DSC, TGA, KF and HPLC analyses were used to characterize all of the batches. After completing the optimization work, the optimized process was scaled-up to 100-200 g on three lots of API. Drying studies were carried

US 7,465,800 B2

13

out at 20° C., 30° C. and 40° C., and 65° C. with a vacuum of 150 mm of Hg. The results are shown in Tables 1-5.

The cooling and holding periods of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione slurry were studied. The experimental laboratory data suggests that polymorph B seems to be forming first, and overtime equilibration to polymorph E at RT conditions occurs, therefore generating a mixture of polymorphs B and E. This result supports the fact that polymorph B seems to be a kinetic product, and that prolonged processing time converts the material to polymorph E resulting in a mixture of polymorphs B and E.

A laboratory procedure was developed to exclusively produce polymorph B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The procedure includes a stirred 10 volume water slurry at ~75° C. for 6-24 hours. The following preferred process parameters have been identified:

1. Hot slurry temperature of 70-75° C.
2. Product filtration of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at 65-75° C.
3. Drying under vacuum at 60-70° C. is preferred for an efficient removal of unbound water in 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione wet cake.
4. The filtration step of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione may be a time sensitive operation. The use of efficient solid-liquid separation equipment is preferred.
5. Holding periods of water-wet cake of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at KF higher than 5% may cause the kinetic equilibrations of polymorph B to mixed polymorphs of E and B.

Drying to KF <4.0% water was achieved in ~3 hours (30-70° C., 152 mm Hg). Polymorphs B and E were distinguished by the water levels as measured by KF and TGA. The reference sample of polymorph B is micronized API. In order to make accurate comparison by XRPD samples were gently grinded before submission for analysis. This increases the clarity of the identification of the polymorphic form. All samples were analyzed for XRPD, DSC, TGA, KF and HPLC.

TABLE 1

<u>Preliminary Studies</u>			
Amount	Reaction conditions	Analysis	Results/conclusion
2 g	Water, rt, 48 h	XRPD, DSC, TGA, KF	Polymorph E
25 g	Water, rt, 48 h	XRPD, DSC, TGA, KF	Polymorph E
5 g	Water, 70-75° C., 24 h then rt 24 h	XRPD, DSC, TGA, KF	Polymorph B
1 g	9:1 Acetone - water, Slow evpo.	XRPD, DSC, TGA, KF	Mixture
1 g	175° C. 1 h in an oven	XRPD, DSC, TGA, KF	Polymorph A
0.5 g (polymorph A)	Water, rt, 24 h	XRPD, DSC, TGA, KF	Polymorph E
1 g polymorph B	Water, rt, 48 h	XRPD, DSC, TGA, KF	Polymorph E
1 g polymorph E	Water, 70-75° C., 24 h	XRPD, DSC, TGA, KF	Polymorph B
1 g	Slurry in heptane	XRPD, DSC, TGA, KF	No change

14

TABLE 2

<u>Optimization of Temperature, Time and Solvent Volume</u>					
	Amount	Amount Water (mL)	Temp (° C.)	Time (h)	Results/conclusion
10	10 g	50	75	6	Mix
	10 g	50	75	24	Polymorph B
	10 g	100	70	6	Polymorph B
	10 g	100	70	14	Polymorph B
	10 g	100	70	21	Polymorph B
	10 g	100	75	6	Polymorph B
	10 g	100	75	24	Polymorph B
	10 g	100	75	6	Polymorph B
	10 g	100	75	19	Polymorph B
	10 g	100	75	14	Polymorph B
	10 g	100	75	24	Polymorph B
	5 g	100	75	18	Polymorph B
	10 g	100	80	6	Polymorph B
	10 g	100	80	20	Polymorph B
20	10 g	200	45	6	Polymorph B + E
	10 g	200	45	24	Polymorph E
	10 g	200	60	48	Polymorph B
	10 g	200	75	6	Mix
	10 g	200	75	24	Polymorph B
	10 g	200	75	13	Polymorph B
	10 g	200	75	24	Polymorph B
	10 g	200	75	24	Polymorph B

25

Optimum conditions were determined to be 10 volumes of solvent (H_2O), 70-80° C. for 6-24 hours.

30

TABLE 3

<u>Holding Time</u>					
	Amount	Reaction Conditions	Holding Time (h)	Holding Temp (° C.)	Results/Conclusion
35	5 g	Water, 70-75° C., 24 h	24	23-25	Polymorph B
	1 g	Water, 70-75° C., 24 h	48	23-25	Polymorph E
	Polymorph B				
	2 g	Water, 40 mL	16	23-25	Polymorph E
	150 g	Water, 3.0 L	24	23-25	Polymorph E
	150 g	Water, 3.0 L	48	23-25	Polymorph E
	10 g	Water, 100 mL, 24 h, 75° C.	18	23-25	Polymorph B
	10 g	Water, 100 mL, 24 h, 75° C.	18	40	Polymorph B
	10 g	Water, 200 mL, 24 h, 75° C.	14	-5	Mix
	10 g	Water, 200 mL, 24 h, 75° C.	14	23-25	Polymorph E
40	10 g	Water, 200 mL, 24 h, 75° C.	14	40	Mix
	10 g	Water, 200 mL, 24 h, 75° C.	14	23-25	Polymorph E
	10 g	Water, 200 mL, 24 h, 75° C.	21	23-25	Polymorph E
	10 g	Water, 100 mL, 24 h, 75° C.	21	40	Mix
	10 g	Water, 100 mL, 24 h, 75° C.	21	40	Mix
	10 g	Water, 100 mL, 14 h, 75° C.	2	23-25	Mix
	10 g	Water, 100 mL, 14 h, 75° C.	2	23-25	Mix
	10 g	Water, 100 mL, 14 h, 75° C.	2	23-25	Mix
	10 g	Water, 100 mL, 14 h, 75° C.	2	23-25	Mix

55

Holding time gave mixed results and it was determined that the material should be filtered at 60-65° C. and the material washed with 0.5 volume of warm (50-60° C.) water.

60

TABLE 4

<u>Scale-up Experiments</u>					
	Amount	Amount Water (L)	Temp (° C.)	Time (h)	Results/Conclusion
65	100 g	1.0	75	6	Polymorph B
	100 g	1.0	75	22	Polymorph B

US 7,465,800 B2

15

TABLE 4-continued

<u>Scale-up Experiments</u>				
Amount	Amount Water (L)	Temp (° C.)	Time (h)	Results/ Conclusion
100 g	1.0	75	6	Polymorph B
100 g	1.0	75	24	Polymorph B
100 g	1.0	75	6	Polymorph B
100 g	1.0	75	22	Polymorph B

TABLE 5

<u>Drying Studies</u>					
Amount	Drying Time (h)	Drying Temp (° C.)	Vacuum (mm Hg)	KF\$ (%)	Results/ Conclusion
100 g	0	—	—	3.690	Polymorph B
100 g	3	30	152	3.452	Polymorph B
100 g	8	30	152	3.599	Polymorph B
100 g	0	—	—	3.917	Polymorph B
100 g	5	40	152	3.482	Polymorph B
100 g	22	40	152	3.516	Polymorph B
100 g	3	40	152	3.67	Polymorph B
100 g	22	40	152	3.55	Polymorph B

* Reaction Conditions: Water 1 L, 75° C., 22-24 h;

§ Average of 2 runs.

Drying studies determined that the material should be dried at 35-40° C., 125-152 mm Hg for 3 to 22 h or until the water content reaches $\leq 4\%$ w/w.

For a large scale preparation of polymorph E (5222-152-B), a 5-L round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (150 g, 0.579 mol) and water (3000 mL, 20 vol). The mixture was mechanically stirred at room temperature (23-25° C.) for 48 h under nitrogen atmosphere.

Samples were taken after 24 h and 48 h before the mixture was filtered and air-dried on the filter for 1 h. The material was transferred to a drying tray and dried at room temperature (23-25° C.) for 24 h. KF analysis on the dried material showed water content of 11.9%. The material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph E.

For a large scale preparation of polymorph B (5274-104), a 2 L-3-necked round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (polymorph mixture, 100 g, 0.386 mol) and water (1000 mL, 10.0 vol). The mixture was heated to 75° C. over approximately 30 minutes with mechanical stirring under nitrogen atmosphere.

Samples were taken after 6 h and 24 h before the mixture was allowed to cool to 60-65° C., filtered and the material washed with warm (50-60° C.) water (50 mL, 0.5 vol). The material was transferred to a drying tray and dried at 30° C., 152 mm Hg for 8 h. KF analysis on the dried material showed water content of 3.6%. After grinding the material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph B. The results of the analyses are shown in FIGS. 32-46.

6.3 X-Ray Powder Diffraction Measurements

X-ray powder diffraction analyses were carried out on a Shimadzu XRD-6000 X-ray powder diffractometer using Cu K α radiation. The instrument is equipped with a fine-focus X-ray tube. The tube voltage and amperage were set at 40 kV and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm.

16

Diffracted radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 2.5 degrees 2θ to 40 degrees 2θ was used. A silicon standard was analyzed each day to check the instrument alignment.

X-ray powder diffraction analyses were also carried out using Cu K α radiation on an Inel XRG-3000 diffractometer equipped with a curved position-sensitive detector. Data were collected in real time over a theta-two theta range of 120° at a resolution of 0.030°. The tube voltage and current were 40 kV and 30 mA, respectively. A silicon standard was analyzed each day to check for instrument alignment. Only the region between 2.5 and 40 degrees 2θ is shown in the figures.

6.4 Thermal Analysis

TG analyses were carried out on a TA Instrument TGA 2050 or 2950. The calibration standards were nickel and alumel. Approximately 5 mg of sample was placed on a pan, accurately weighed, and inserted into the TG furnace. The samples were heated in nitrogen at a rate of 10° C./min, up to a final temperature of 300 or 350° C.

DSC data were obtained on a TA 2920 instrument. The calibration standard was indium. Approximately 2-5 mg samples were placed into a DSC pan and the weight accurately recorded. Crimped pans with one pinhole were used for analysis and the samples were heated under nitrogen at a rate of 10° C./min, up to a final temperature of 350° C.

Hot-stage microscopy was carried out using a Kofler hot stage mounted on a Leica Microscope. The instrument was calibrated using USP standards.

A TA Instruments TGA 2050 interfaced with a Nicolet model 560 Fourier transform IR spectrophotometer, equipped with a globar source, XT/KBr beamsplitter, and deuterated triglycine sulfate (DTGS) detector, was utilized for TG-IR experiments. The IR spectrometer was wavelength calibrated with polystyrene on the day of use while the TG was temperature and weight calibrated biweekly, using indium for the temperature calibration. A sample of approximately 10 mg of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was weighed into an aluminum pan and heated from 25 to 30° C. to 200° C. at a rate of 20° C./min with a helium purge. IR spectra were obtained in series, with each spectrum representing 32 co-added scans at a resolution of 4 cm⁻¹. Spectra were collected with a 17-second repeat time. TG/IR analysis data are presented as Gram-Schmidt plots and IR spectra linked to the time. Gram-Schmidt plots show total IR intensity vs. time; hence, the volatiles can be identified at each time point. They also show when the volatiles are detected. From the Gram-Schmidt plots, time points were selected and the IR spectra of these time points are presented in the stacked linked spectra. Each spectrum identifies volatiles evolving at that time point. Volatiles were identified from a search of the HR Nicolet TGA vapor phase spectral library. The library match results are also presented to show the identified vapor.

6.5 Spectroscopy Measurements

Raman spectra were acquired on a Nicolet model 750 Fourier transform Raman spectrometer utilizing an excitation wavelength of 1064 nm and approximately 0.5 W of Nd:YAG laser power. The spectra represent 128 to 256 co-added scans acquired at 4 cm⁻¹ resolution. The samples were prepared for analysis by placing the material in a sample holder and positioning this in the spectrometer. The spectrometer was wavelength calibrated using sulfur and cyclohexane at the time of use.

The mid-IR spectra were acquired on a Nicolet model 860 Fourier transform IR spectrophotometer equipped with a glo-

US 7,465,800 B2

17

bar source XT/KBr beamsplitter and a deuterated triglycine sulfate (DTGS) detector. A Spectra-Tech, Inc. diffuse reflectance accessory was utilized for sampling. Each spectrum represents 128 co-added scans at a spectral resolution of 4 cm⁻¹. A background data set was acquired with an alignment mirror in place. A single beam sample data set was then acquired. Subsequently, a log 1/R (where R=reflectance) spectrum was acquired by rationing the two data sets against each other. The spectrophotometer was calibrated (wavelength) with polystyrene at the time of use.

6.6 Moisture Sorption/Desorption Measurements

Moisture sorption/desorption data were collected on a VTI SGA-100 moisture balance system. For sorption isotherms, a sorption range of 5 to 95% relative humidity (RH) and a desorption range of 95 to 5% RH in 10% RH increments was used for analysis. The sample was not dried prior to analysis. Equilibrium criteria used for analysis were less than 0.0100 weight percent change in 5 minutes with a maximum equilibration time of 3 hours if the weight criterion was not met. Data were not corrected for the initial moisture content of the samples.

6.7 Solution Proton NMR Measurements

NMR spectra not previously reported were collected at SSCI, Inc, 3065 Kent Avenue, West Lafayette, Ind. Solution phase ¹H NMR spectra were acquired at ambient temperature on a Bruker model AM spectrometer. The ¹H NMR spectrum represents 128 co-added transients collected with a 4 μ sec pulse and a relaxation delay time of 5 seconds. The free induction decay (FID) was exponentially multiplied with a 0.1 Hz Lorentzian line broadening factor to improve the signal-to-noise ratio. The NMR spectrum was processed utilizing GRAMS software, version 5.24. Samples were dissolved in dimethyl sulfoxide-d₆.

The scope of this invention can be understood with reference to the appended claims.

6.8 Intrinsic Dissolution and Solubility Studies

Intrinsic dissolution experiments were conducted on Form A (anhydrous), Form B (hemihydrate), and Form E (dihydrate) of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Equilibrium solubility experiments were conducted on Forms A and B. Aliquots were analyzed by ultraviolet-visible spectrophotometry, and the solids remaining from each experiment were analyzed by X-ray powder diffraction (XRPD).

6.8.1 Experimental

6.8.1.1 Dissolution

Dissolution experiments were carried out in a VanKel VK6010-8 dissolution apparatus equipped with a VK650A heater/circulator. An intrinsic dissolution apparatus (Woods apparatus) was used. Samples were compressed at 1.5 metric tons (1000 psi) for 1 min using the Woods apparatus in a hydraulic press, giving a sample surface of 0.50 cm². A dissolution medium consisting of 900 mL HCl buffer, pH 1.8, with 1% sodium lauryl sulfate, was used for each experiment. The medium was degassed by vacuum filtration through a 0.22- μ m nylon filter disk and maintained at 37° C. The apparatus was rotated at 50 rpm for each experiment. Aliquots were filtered immediately using 0.2- μ m nylon syringe filters. In some cases, the undissolved solids were recovered and analyzed by X-ray powder diffraction (XRPD).

6.8.1.2 Solubility

Equilibrium solubility experiments were conducted in a 100-mL, three-neck, round-bottom flask immersed in a constant temperature oil bath maintained at 25° C. A solid sample of 400-450 mg was stirred in 50 mL of dissolution medium (HCl buffer, pH 1.8, with 1% sodium lauryl sulfate) using a mechanical stir rod. Aliquots were filtered using 0.2- μ m

18

nylon syringe filters and immediately diluted 1 mL → 50 mL, then 5 mL → 25 mL with dissolution medium in Class A glassware, a final dilution factor of 250.

6.8.1.3 UV-Vis Spectrophotometry

Dissolution and solubility samples solutions were analyzed by a Beckman DU 640 single-beam spectrophotometer. A 1.000-cm quartz cuvette and an analysis wavelength of 228.40 nm were utilized. The detector was zeroed with a cuvette filled with dissolution medium.

10 6.8.1.4 X-Ray Powder Diffraction

XRPD analyses were carried out on a Shimadzu XRD-6000 X-ray powder diffractometer using Cu K α radiation. The instrument is equipped with a fine focus X-ray tube. The tube power and amperage were set at 40 kV and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm. Diffracted radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 2.5 to 40° 20 was used. A silicon standard was analyzed each day to check the instrument alignment. Samples were packed in an aluminum holder with silicon insert.

6.8.2 Results

The results of these solubility and intrinsic studies are summarized in Table 6. Both the solubility and dissolution experiments were conducted in a medium of HCl buffer, pH 1.8, containing 1% sodium lauryl sulfate. Form A was found to be unstable in the medium, converting to Form E. The solubilities of Forms A, B, and E were estimated to be 6.2, 5.8, and 4.7 mg/mL, respectively. The dissolution rates of Forms A, B, and E were estimated to be 0.35, 0.34, and 0.23 mg/mL, respectively.

6.8.2.1 UV-Vis Spectrophotometry Method Development

A UV-Vis scan of the dissolution medium (blanked with an empty cuvette) was done to identify any interfering peaks. A small peak at 225 nm was present as shown in FIG. 47.

Solutions of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at varying concentrations were analyzed by UV-Vis spectrophotometry. A preliminary scan of a 1.0 mg/mL solution was done, with the instrument blanked with dissolution medium. The solution was highly absorbing and noisy from 200-280 nm, making dilution necessary.

A 0.04 mg/mL solution of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was then scanned from 200-300 nm. The plot was still noisy between 200 and 230 nm as shown in FIG. 48. The sample was further diluted to 0.008 mg/mL. A wavelength scan of 200-350 nm for this sample showed a peak a 228.4 nm with no interference, as shown in FIG. 49. Therefore, a wavelength of 228.4 was chosen for analysis of the solubility and dissolution samples.

A six-point calibration curve was generated with standards of the following concentrations: 0.001 mg/mL, 0.002 mg/mL, 0.005 mg/mL, 0.010 mg/mL, 0.015 mg/mL, and 0.020 mg/mL (Notebook 569-90). A linearity coefficient of R²=0.9999 was obtained as shown in FIG. 50.

55 6.8.2.2 Solubility

A sample consisting of 449.4 mg Form A was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, and 150 min. The concentration reached 6.0 mg/mL by the first time point. The highest concentration reached was 6.2 mg/mL, at 30 min. From that point the concentration decreased, reaching 4.7 mg/mL at 150 min as in FIG. 51. The solids remaining at the final time point were analyzed by XRPD and found to be Form E as shown in Table 7. No peaks attributed to Form A can be seen in the pattern. Since the concentration did not plateau at 4.7 mg/mL, the solubility of Form E may be lower than that.

US 7,465,800 B2

19

A sample consisting of 401.4 mg Form B was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, 180, 420, and 650 min. Form B dissolved much more slowly than Form A, reaching 3.3 mg/mL in 90 min. The concentration stabilized at 5.6–5.7 mg/mL at the final three time points as in FIG. 52. The remaining solids were shown to be Form B as in Table 7, suggesting Form B has good stability in water.

A summary of the solubilities is given in Table 6. The amounts dissolved at each time point are shown in Tables 8 and 9.

TABLE 6

Form	Solubility	Summary of Results		
		Intrinsic Dissolution #1	Intrinsic Dissolution #2	Average Intrinsic Dissolution Rate
Form A	6.2 mg/mL	0.35	0.22 ^a	0.29 ^a
Form B	5.8 mg/mL	0.35	0.32	0.34
Form E	4.7 mg/mL	0.21	0.25	0.23

^aThe Form A dissolution experiment #2 may have converted to Form E on the surface of the disk, skewing the average rate lower.

TABLE 7

Experimental Details	
Experiment	Final Form
Pressed Form A	A
Pressed Form B	B
Form A Solubility	E
Form B Solubility	B
Form A Dissolution	—
Form A Dissolution	A
Form B Dissolution	—
Form B Dissolution	B
Form E Dissolution	E
Form E Dissolution	—

TABLE 8

Form A Solubility	
Time Point (min)	Concentration (mg/mL)
7	6.00
15	6.11
30	6.16
60	6.10
90	5.46
150	4.73

TABLE 9

Form B Solubility	
Time Point (min)	Concentration (mg/mL)
7	1.63
15	2.14
30	2.33
60	2.94
90	3.34
180	5.67
420	5.76
650	5.61

20

6.8.2.3 Intrinsic Dissolution

Approximately 200 mg each of Forms A and B were compressed into disks in the Woods apparatus using 2 metric tons of pressure. The samples were subsequently scraped out, ground gently, and analyzed by XRPD. The study showed that compression and grinding does not cause a form change in either case. (See Table 7).

Two preliminary dissolution runs were performed. The 10 disks fractured to some extent in both experiments, compromising the requirement of constant surface area.

The first experiment of intrinsic dissolution that strictly followed the USP chapter on intrinsic dissolution utilized 15 approximately 150 mg each of Forms A and B. Seven aliquots, beginning at 5 min and ending at 90 min, were taken to maintain sink conditions. The experiment resulted in linear 20 dissolution profiles, with a rate of 0.35 mg per cm² per minute for both forms. The Form E experiment was done later under 25 the same conditions and added to the graph for comparison. (See FIG. 53). The Form E dissolution rate was 0.21 mg per cm² per minute, significantly lower than the dissolution rate of Forms A and B. This is in line with expectations based on the solubility data. The crystal form of the remaining solids did not change in any case.

The second experiment utilized approximately 250 mg 30 each of Forms A and B. The Form E experiment (135 mg) was done later and added to the graph for comparison. (See FIG. 35 54). Nine aliquots were taken, beginning at 5 min and ending at 150 min. The dissolution rates were 0.22, 0.32, and 0.25 mg per cm² per minute, respectively, for Forms A, B, and E. The dissolution rate for Form A in this experiment was low, while 35 the rates for Forms B and E were similar to those found in the first experiment. It is believed that in this case, a thin layer of the Form A sample disk may have converted to Form E upon exposure to water. This is supported by the evidence of rapid conversion of Form A to Form E in the solubility experiment. 40 The diffraction pattern of the undissolved solids does not indicate a form change. However, the bulk of the sample disk is not exposed to water. Therefore, the true intrinsic dissolution rate of Form A is believed to be close to 0.35 mg per cm² 45 per minute. An insufficient quantity of Form A was available to repeat the experiment.

A summary of the intrinsic dissolution rates is given in 50 Table 6. The amounts dissolved at each time point are summarized in Tables 10 and 11.

TABLE 10

Intrinsic Dissolution Experiment #1 Results			
Time Point	Form A ^a	Form B ^a	Form E ^a
55	5 min	5.76	10.80 ^b
	10 min	7.73	6.85
	20 min	11.31	10.25
	30 min	15.59	14.35
	45 min	21.98	20.57
	60 min	27.11	25.70
60	90 min	34.17	34.34
			20.82

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)

^bThis data point not included in graph since the value is higher than the next two data points.

US 7,465,800 B2

21

TABLE 11

Intrinsic Dissolution Experiment #2 Results			
Time Point	Form A ^a	Form B ^a	Form E ^a
5 min	4.50	5.04	3.06
10 min	5.22	6.12	4.31
20 min	7.54	7.73	11.40
30 min	11.46	12.72	11.93
45 min	15.01	17.33	14.72
60 min	18.38	21.93	18.52
90 min	24.38	31.64	26.24
120 min	30.35	41.31	33.56
150 min	35.26	49.54	40.82

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)

6.9 Analyses of Mixtures of Polymorphs

This invention encompasses mixtures of different polymorphs. For example, an X-ray diffraction analysis of one production sample yielded a pattern that contained two small peaks seen at approximately 12.6° and 25.8° 2θ in addition to those representative of Form B. In order to determine the composition of that sample, the following steps were performed:

- 1) Matching of the new production pattern to known forms along with common pharmaceutical excipients and contaminants;
- 2) Cluster analysis of the additional peaks to identify if any unknown phase is mixed with the original Form B;
- 3) Harmonic analysis of the additional peaks to identify if any preferred orientation may be present or if any changes in the crystal habit may have occurred; and
- 4) Indexing of the unit cells for both Form B and the new production sample to identify any possible crystallographic relationships.

Based on these tests, which can be adapted for the analysis of any mixture of polymorphs, it was determined that the sample contained a mixture of polymorph forms B and E.

6.10 Dosage Form

Table 12 illustrates a batch formulation and single dosage formulation for a 25 mg single dose unit of a polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

TABLE 12

Formulation for a 25 mg capsule			
Material	Percent By Weight	Quantity (mg/tablet)	Quantity (kg/batch)
Polymorphic Form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione	40.0%	25 mg	16.80 kg
Pregelatinized Corn Starch, NF	59.5%	37.2 mg	24.99 kg
Magnesium Stearate	0.5%	0.31 mg	0.21 kg
Total	100.0%	62.5 mg	42.00 kg

22

The pregelatinized corn starch (SPRESS B-820) and polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione components are passed through a screen (i.e., a 710 μm screen) and then loaded into a Diffusion Mixer with a baffle insert and blended for about 15 minutes. The magnesium stearate is passed through a screen (i.e., a 210 μm screen) and added to the Diffusion Mixer. The blend is then encapsulated in capsules using a Dosator type capsule filling machine.

10 The entire scope of this invention is not limited by the specific examples described herein, but is more readily understood with reference to the appended claims.

What is claimed is:

1. Crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate.
2. The hemihydrate of claim 1 having an X-ray powder diffraction pattern comprising peaks at approximately 16, 22 and 27 degrees 2θ.
3. The hemihydrate of claim 2 wherein the pattern further comprises a peak at approximately 18 degrees 2θ.
4. The hemihydrate of claim 1 having an X-ray powder diffraction pattern comprising peaks at 15.8, 22.2 and 26.7 degrees 2θ.
5. The hemihydrate of claim 4 wherein the pattern further comprises a peak at 18.2 degrees 2θ.
6. The hemihydrate of claim 1 having an X-ray powder diffraction pattern corresponding to the representative X-ray powder diffraction patterns depicted in FIG. 6, FIG. 32, FIG. 33 and FIG. 34.
7. The hemihydrate of claim 1 having a differential scanning calorimetry thermogram comprising an endotherm with a maximum at about 268° C.
8. The hemihydrate of claim 7 wherein the thermogram further comprises an endotherm corresponding to dehydration.
9. The hemihydrate of claim 1 having a differential scanning calorimetry thermogram corresponding to the representative differential scanning calorimetry thermograms depicted in FIG. 9, FIG. 42, FIG. 43 and FIG. 44.
10. The hemihydrate of claim 1 having between approximately 0.46 and approximately 0.59 moles of water per mole of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.
11. The hemihydrate of claim 1 having a thermogravimetric analysis thermogram comprising a weight loss of between about 3.1% and about 4.0% when heated from about 30° C. to about 175° C.
12. The hemihydrate of claim 1 having an infrared spectrum comprising peaks at approximately 3513 and 1960 cm⁻¹.
13. The hemihydrate of claim 1 having an infrared spectrum corresponding to the representative infrared spectrum depicted in FIG. 7.
14. The hemihydrate of claim 1 having a Raman spectrum corresponding to the representative Raman spectrum depicted in FIG. 8.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,465,800 B2
APPLICATION NO. : 10/934863
DATED : December 16, 2008
INVENTOR(S) : Jaworsky et al.

Page 1 of 11

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Please replace Drawing Sheets 15, 26, 31, 32, 33, 35, 36, 38, 39, and 40 (of 48) with the Replacement Drawing Sheets on the following pages of this Certificate of Correction:

Page 2 of 11: Replacement Drawing Sheet 15 of 48 (Fig. 15);

Page 3 of 11: Replacement Drawing Sheet 26 of 48 (Fig. 26);

Page 4 of 11: Replacement Drawing Sheet 31 of 48 (Fig. 31);

Page 5 of 11: Replacement Drawing Sheet 32 of 48 (Figs. 32 and 33);

Page 6 of 11: Replacement Drawing Sheet 33 of 48 (Figs. 34 and 35);

Page 7 of 11: Replacement Drawing Sheet 35 of 48 (Figs. 37 and 38);

Page 8 of 11: Replacement Drawing Sheet 36 of 48 (Figs. 39 and 40);

Page 9 of 11: Replacement Drawing Sheet 38 of 48 (Figs. 42 and 43);

Page 10 of 11: Replacement Drawing Sheet 39 of 48 (Figs. 44 and 45);

Page 11 of 11: Replacement Drawing Sheet 40 of 48 (Fig. 46).

Signed and Sealed this
Nineteenth Day of April, 2011



David J. Kappos
Director of the United States Patent and Trademark Office


CERTIFICATE OF CORRECTION (continued)

Page 2 of 11

U.S. Patent

Dec. 16, 2008

Sheet 15 of 48

7,465,800 B2**FIG. 15**

CERTIFICATE OF CORRECTION (continued)

Page 3 of 11

U.S. Patent

Dec. 16, 2008

Sheet 26 of 48

7,465,800 B2




FIG.26

CERTIFICATE OF CORRECTION (continued)

Page 4 of 11

U.S. Patent

Dec. 16, 2008

Sheet 31 of 48

7,465,800 B2




FIG.31

CERTIFICATE OF CORRECTION (continued)

Page 5 of 11

U.S. Patent

Dec. 16, 2008

Sheet 32 of 48

7,465,800 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5274-104-B

Date: 06-04-04 16:10 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.32

XRPD PATTERN OF POLYMORPH B

File: Process 5274-100-C

Date: 06-02-04 16:11 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.33

CERTIFICATE OF CORRECTION (continued)

Page 6 of 11

U.S. Patent

Dec. 16, 2008

Sheet 33 of 48

7,465,800 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5222-157-C

Date: 06/04/04 15:07 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.34

XRPD PATTERN OF POLYMORPH E

File: Process 5222-152-B Form E

Date: 05/21/04 10:46 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.35

CERTIFICATE OF CORRECTION (continued)



Page 7 of 11

U.S. Patent

Dec. 16, 2008

Sheet 35 of 48

7,465,800 B2

**FIG.37****FIG.38**

CERTIFICATE OF CORRECTION (continued)

Page 8 of 11

U.S. Patent

Dec. 16, 2008

Sheet 36 of 48

7,465,800 B2

TGA CURVE OF POLYMORPH B
 Sample: 5222-157-C TGA
 Size: 24.4610mg
 Method: Ramp Run Date: 07-Jun-04 09:46
 Comment: Evotec Batch 17 Instrument: TGA 0.500 V5.3 Build 171




FIG.39

TGA CURVE OF POLYMORPH E
 Sample: 5222-152-B TGA
 Size: 11.3850mg
 Method: Ramp Run Date: 21-May-04 09:34
 Comment: Form E Instrument: TGA 0.500 V5.3 Build 171




FIG.40

CERTIFICATE OF CORRECTION (continued)

Page 9 of 11

U.S. Patent

Dec. 16, 2008


Sheet 38 of 48

7,465,800 B2

DSC OF POLYMORPH B
DSC

Sample: 5274-104-B
Size: 0.0000mg
Method: Cell constant calibration
Comment: OAI batch 15 22b


Run Date: 07-Jun-04 11:30
Instrument: DSC Q 1000 V7.3 Build 249

**FIG.42**

DSC OF POLYMORPH B
DSC

Sample: 5274-100-C
Size: 0.0000mg
Method: Cell constant calibration
Comment: Evotec Batch 1675C 10vol H₂O 24h 3h dry

Run Date: 02-Jun-04 17:01
Instrument: DSC Q 1000 V7.3 Build 249

**FIG.43**

CERTIFICATE OF CORRECTION (continued)

Page 10 of 11

U.S. Patent

Dec. 16, 2008

Sheet 39 of 48

7,465,800 B2

Sample: 5222-157-C DSC OF POLYMORPH B
 Size: 0.0000mg DSC
 Method: Cell constant calibration Run Date: 07-Jun-04 09:45
 Comment: OAI batch 17 Instrument: DSC Q 1000 V7.3 Build 249




FIG.44

Sample: 5222-152-B DSC OF POLYMORPH E
 Size: 3.4500mg DSC
 Method: Cell constant calibration Run Date: 21-May-04 09:32
 Comment: Evotec batch 15 Form E Instrument: DSC Q 1000 V7.3 Build 249




FIG.45

CERTIFICATE OF CORRECTION (continued)

Page 11 of 11

U.S. Patent

Dec. 16, 2008

Sheet 40 of 48

7,465,800 B2

Sample: 5222-161-A
Size: 0.0000mg

Method: Cell constant calibration
Comment: OAI batch 15

DSC OF POLYMORPH MIXTURE
DSC

Run Date: 11-Jun-04 12:44
Instrument: DSC Q 1000 V7.3 Build 249




FIG.46

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,465,800 B2
APPLICATION NO. : 10/934863
DATED : December 16, 2008
INVENTOR(S) : Jaworsky et al.

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page:

The first or sole Notice should read --

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b)
by 966 days.

Signed and Sealed this
Fifteenth Day of May, 2012



David J. Kappos
Director of the United States Patent and Trademark Office

EXHIBIT B



US007855217B2

(12) **United States Patent**
Jaworsky et al.

(10) **Patent No.:** US 7,855,217 B2
(45) **Date of Patent:** *Dec. 21, 2010

(54) **POLYMORPHIC FORMS OF
3-(4-AMINO-1-OXO-1,3
DIHYDRO-ISOINDOL-2-YL)-
PIPERIDINE-2,6-DIONE**

(75) Inventors: **Markian S. Jaworsky**, Hopewell, NJ (US); **Roger Shen-Chu Chen**, Edison, NJ (US); **George W. Muller**, Bridgewater, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 82 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/335,395**

(22) Filed: **Dec. 15, 2008**

(65) **Prior Publication Data**

US 2009/0149500 A1 Jun. 11, 2009

Related U.S. Application Data

(62) Division of application No. 10/934,863, filed on Sep. 3, 2004, now Pat. No. 7,465,800.

(60) Provisional application No. 60/499,723, filed on Sep. 4, 2003.

(51) **Int. Cl.**

A61K 31/454 (2006.01)
C07D 401/04 (2006.01)

(52) **U.S. Cl.** **514/323; 546/200**

(58) **Field of Classification Search** **514/323;
546/200**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,536,809 A	10/1970	Applezwig
3,598,123 A	8/1971	Zaffaroni
3,845,770 A	11/1974	Theeuwes et al.
3,916,899 A	11/1975	Theeuwes et al.
4,008,719 A	2/1977	Theeuwes et al.
4,810,643 A	3/1989	Souza
4,999,291 A	3/1991	Souza
5,059,595 A	10/1991	Le Grazie
5,073,543 A	12/1991	Marshall et al.
5,120,548 A	6/1992	McClelland et al.
5,229,496 A	7/1993	Deeley et al.
5,354,556 A	10/1994	Sparks et al.
5,385,901 A	1/1995	Kaplan et al.
5,391,485 A	2/1995	Deeley et al.
5,393,870 A	2/1995	Deeley et al.
5,528,823 A	6/1996	Rudy, Jr. et al.
5,580,755 A	12/1996	Souza
5,591,767 A	1/1997	Mohr et al.
5,593,990 A	1/1997	D'Amato
5,629,327 A	5/1997	D'Amato
5,635,517 A	6/1997	Muller et al.

5,639,476 A	6/1997	Oshlack et al.
5,674,533 A	10/1997	Santus et al.
5,698,579 A	12/1997	Muller
5,712,291 A	1/1998	D'Amato
5,731,325 A	3/1998	Andrusis, Jr. et al.
5,733,566 A	3/1998	Lewis
5,798,368 A	8/1998	Muller et al.
5,874,448 A	2/1999	Muller et al.
5,877,200 A	3/1999	Muller
5,929,117 A	7/1999	Muller et al.
5,955,476 A	9/1999	Muller et al.
6,020,358 A	2/2000	Muller et al.
6,071,948 A	6/2000	D'Amato
6,114,335 A	9/2000	D'Amato
6,114,355 A	9/2000	D'Amato
6,140,346 A	10/2000	Andrusis, Jr. et al.
6,235,756 B1	5/2001	D'Amato
6,281,230 B1	8/2001	Muller et al.
6,316,471 B1	11/2001	Muller et al.
6,326,388 B1	12/2001	Man et al.
6,335,349 B1	1/2002	Muller et al.
6,380,239 B1	4/2002	Muller et al.
6,395,754 B1	5/2002	Muller et al.
6,403,613 B1	6/2002	Man et al.
6,420,414 B1	7/2002	D'Amato
6,458,810 B1	10/2002	Muller et al.
6,469,045 B1	10/2002	D'Amato
6,476,052 B1	11/2002	Muller et al.
6,518,298 B2	2/2003	Green et al.
6,555,554 B2	4/2003	Muller et al.

(Continued)

FOREIGN PATENT DOCUMENTS

JP A H10-53576 2/1998

(Continued)

OTHER PUBLICATIONS

Dimartino et al., 1997, "Preparation and physical characterization of forms II and III of paracetamol," J. Thermal Analysis 48:447-458.

(Continued)

Primary Examiner—Celia Chang
(74) Attorney, Agent, or Firm—Jones Day

(57) **ABSTRACT**

Polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isooindol-2-yl)-piperidine-2,6-dione are disclosed. Compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use are also disclosed.

US 7,855,217 B2

Page 2

U.S. PATENT DOCUMENTS

7,112,602	B2	9/2006	D'Amato et al.
7,465,800	B2	12/2008	Jaworsky et al.
2001/0018445	A1	8/2001	Huang et al.
2001/0056114	A1	12/2001	D'Amato
2002/0035090	A1	3/2002	Zeldis et al.
2002/0045643	A1	4/2002	Muller et al.
2002/0052398	A1	5/2002	D'Amato
2002/0054899	A1	5/2002	Zeldis
2002/0061923	A1	5/2002	D'Amato
2002/0128228	A1	9/2002	Hwu
2002/0161023	A1	10/2002	D'Amato
2002/0173658	A1	11/2002	Muller et al.
2002/0183360	A1	12/2002	Muller et al.
2003/0013739	A1	1/2003	Masferrer et al.
2003/0028028	A1	2/2003	Man et al.
2003/0045552	A1	3/2003	Robarge et al.
2003/0069428	A1	4/2003	Muller et al.
2003/0096841	A1	5/2003	Robarge et al.
2003/0139451	A1	7/2003	Shah et al.
2003/0144325	A1	7/2003	Muller et al.
2003/0181428	A1	9/2003	Green et al.
2003/0187024	A1	10/2003	D'Amato
2003/0191098	A1	10/2003	D'Amato
2003/0235909	A1	12/2003	Hariri et al.
2004/0029832	A1	2/2004	Zeldis
2004/0077685	A1	4/2004	Figg et al.
2004/0077686	A1	4/2004	Dannenberg et al.
2004/0087546	A1	5/2004	Zeldis
2004/0091455	A1	5/2004	Zeldis
2004/0122052	A1	6/2004	Muller et al.
2004/0220144	A1	11/2004	Zeldis
2004/0266809	A1	12/2004	Emanuel et al.
2005/0203142	A1	9/2005	Zeldis
2006/0052609	A1	3/2006	Muller et al.
2008/0064876	A1	3/2008	Muller et al.
2008/0132541	A1	6/2008	Zeldis et al.
2009/0062343	A1	3/2009	Jaworsky et al.
2009/0149449	A1	6/2009	Liu et al.
2009/0149499	A1	6/2009	Jaworsky et al.
2009/0176832	A1	7/2009	Jaworsky et al.
2009/0187023	A1	7/2009	Jaworsky et al.

FOREIGN PATENT DOCUMENTS

JP	A 2001-503384	3/2001
WO	WO97/46526	A1 12/1997
WO	WO 98/03502	1/1998
WO	WO98/03502	A1 1/1998
WO	WO 98/54170	12/1998
WO	WO 01/70275	9/2001
WO	WO 01/87307	11/2001
WO	WO 02/059106	8/2002
WO	WO 02/064083	8/2002
WO	PCT/US03/11578	1/2003
WO	WO 03097052	11/2003
WO	WO02/26737	A1 4/2004
WO	WO 2004103274	12/2004
WO	WO 2005023192	3/2005
WO	WO 2006/028964	A1 3/2006
WO	WO 2007/136640	A2 11/2007
WO	WO 2009/111948	A1 9/2009
WO	WO 2009/114601	A2 9/2009
WO	WO 2010/054833	5/2010
WO	WO 2010/056384	5/2010
WO	WO 2010/061209	6/2010

WO WO 2010/100476 9/2010

OTHER PUBLICATIONS

- Jonsson, 1972, "Chemical structure and teratogenic properties. 3. A review of available data on structure-activity relationships and mechanism of action of thalidomide analogues," *Acta. Pharm. Suec.* 9(6):521-542.
- Wilens et al., 1977, "Strategies in optical resolutions," *Tetrahedron* 33:2725-2736.
- Wilens, 1972, *Tables of Resolving Agents and Optical Resolutions*, Eliel, ed., U. Notre Dame Press, Notre Dame, IN pp. 268.
- Wolff, ed., 1995, *1 Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178 and 949-982.
- Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogs," *Ann. Rheum. Dis.* 58(Supp. 1):1107-1113.
- Grant, 1999, "Theory and Origin of Polymorphism," in *Polymorphism in Pharmaceutical Solids*, Minneapolis, MN, Ch. 1, pp. 1-10.
- He, W., et al., 1993, 206th American Chemical Society, Chicago, IL; Med. Chem., paper 216.
- Lentzsch et al., 2002, "S-3-Amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice," *Cancer Research* 62:2300-2305.
- Muller et al., 1999, "Amino-substituted thalidomide analogs: Potent inhibitors of TNF- γ production," *Bioorg. Med. Chem. Lett.* 9:1625-1630.
- Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8:2669-2674.
- Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39:3238-3240.
- Anderson, "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology*, 2002 29:17-20.
- Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.
- Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.
- Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients: comparison with predecessor trial total therapy I (TTI) (N=231)," *Blood, Abstract # 2857*, Dec. 7-11, 2001, *American Society of Hematology*.
- Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.
- Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.
- Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4):1-9.
- Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.
- Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVd-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood, Abstract # 2559*, *American Society of Hematology*, Dec. 10-13, 2005.
- Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.
- Celgene Corporation, "Additional clinical data presented on Revlimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.
- Celgene Corporation, "Blood reports Revlimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.
- Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.
- Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.

US 7,855,217 B2

Page 3

- Celgene Corporation, "Celgene corporation advances ACTIMID™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.
- Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene Corporation announces third quarter results. THALOMID® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for THALOMID and REVIMID™ finalized. Peer-reviewed publications of THALOMID and REVIMID data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.
- Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.
- Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.
- Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.
- Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.
- Celgene Corporation, "Celgene expands clinical development program for Revimid™. Five additional trials of Revimid initiated in hematological and solid tumor cancers," Press Release, Jun. 2002.
- Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for ACTIMID™, CC-8490. SeICID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.
- Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead IMiD™, Revimid™," Press Release, Jun. 2001.
- Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes." Press Release, Mar. 2004.
- Chaudhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1)1884-86.
- Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.
- Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents* III 10: 1071-1073.
- D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin. Oncol.* 28:597-601.
- Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl I). S25-S54.
- Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma (~MM)," Abstract # P222, *VIIth International Myeloma Workshop*, May 4-8, 2001.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.
- Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," *Abstract # 1284, Circulation*, 1998.
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," *Abstract # P12.1.4, International Multiple Myeloma Workshop*, May 23-27, 2003.
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," *Abstract # 6, American Society of Hematology*, Dec. 10-13, 2005.
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23)4444-4454.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma \leq 75 years of age," *Am. Soc. Hematol. 46th Ann. Meeting* Dec. 4-7, 2004, San Diego, CA Abstract # 1482.
- Dredge et al., A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer, *Abstract # 491, American Association for Cancer Research*, Apr. 6-10, 2002.
- Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol Immunother.*, 2002, 51:521-531.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.
- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.
- Glaspy et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8 + T cells in vitro," *J. Infect. Dis.* 187(6):946-955.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood*, Abstract # 3219, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers. 206th American Chemical Society, Chicago, IL; Med. Chem., paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally arid pharmacologically related to thalidomide," *Arzneimittel Forschung Drug Research*, 1981, 31 (1)941-949.

US 7,855,217 B2

Page 4

- Hernandez-Illizalitur et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," *Abstract # 235, American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," *Abstract 1313, American Society of Hematology*, Dec. 1-5, 2000.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidomide, in relapsed myeloma," *Blood, Abstract # 3216*, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood, Abstract #208, American Society of Hematology*, Dec. 4-7, 2004.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3):175-180.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," *Abstract # 3073, American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," *Abstract # 3456, American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop, Abstract #P225*, May 4-8, 2001.
- Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Oncology, Abstract #927*, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α - Fluoro-substituted thalidomide analogues," *Biorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Immunotherapeutic and antitumour potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4 $^+$ and CD8 $^+$ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood, Abstract # 1548*, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptic signaling induced by immunomodulatory thalidomide analogs (IMiDs) in human multiple myeloma cells: therapeutic implications," *Abstract # 3224, Dec. 7-11, 2001, American Society of Hematology*.
- Mitsiades et al., "Apoptic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," *Abstract # 3740, American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," *Abstract # 1640, American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology, Abstract # 992*, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood, Abstract #2487*. Dec. 1-5, 2000, *American Society of Hematology*.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL- β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Payvandi et al., "Thalidomide and IMiDS inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood, Abstract # 5046*, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood, Abstract # 2689*, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi Yvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," *Abstract # 1793, American Association for Cancer Research*, Mar. 24-28, 2001.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13):4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13):4050-4053.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood, Abstract # 825, American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma; preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood, Abstract #3225*, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "A phase 1 trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood, Abstract # 365, American Society of Hematology*, Dec. 10-13, 2005.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16):3212-3214.

US 7,855,217 B2

Page 5

- Richardson et al., "Immunomodulatory drug CC-55013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, *American Society of Hematology*.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3):1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology, Abstract #248*, 2002.
- Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," *Abstract # 2485, The American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streetly et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," *Abstract #829, American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," *Abstract # 2543, American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," *Abstract # 367, International Multiple Myeloma Workshop*, May 23-27, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2)93.
- Teo et al., "Chiral inversion of the second generation IMiD™ CC-4047 (ACTIMID™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)- safety and efficacy data in pilot study of 15 patients," *Blood, Abstract # 2869, American Society of Hematology*, Dec. 7-11, 2001.
- Tohnya et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood, Abstract # 3531, American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6):1887-1895.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," *Abstract # PO.738, International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber, "Lenalidomide (CC-5013, Revlimid™) and other IMiDs," *Abstract # PL5.02, International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood, Abstract #4099, American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology, Abstract # 3226*, 2001.
- Zangari et al., "Revimid 25 mg (Rev 25) × 20 versus 50 mg (REV 50) × 10 q 28 days with bridging of 5 mg × 10 versus 10 mg × 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood, Abstract # 1642, American Society of Hematology*, Dec. 6-9, 2003.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (PTS) treated with thalidomide (Thal): The Arkansas Experience," *Blood, Abstract # 681, American Society of Hematology*, Dec. 7-11, 2001.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zeldis et al, "Update on the evolution of the IMiD™," *International Society for Biological Therapy of Cancer: Oral Abstract*, 2003.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," *Abstract # B012, International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- Halebian, J.; McCrone, W; "Pharmaceutical Applications of Polymorphism", *Journal of Pharmaceutical Sciences*, Aug. 1969, 58(8):911-929.
- Davies, F.E.; et al., "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood*, Jul. 2001, 98(1):210-216.
- Kyle, R.A.; Rajkumar, S.V., "Therapeutic Application of Thalidomide in Multiple Myeloma," *Seminars in Oncology*, Dec. 2001, 28(6):583-587.
- USPTO Office Action dated Dec. 31, 2007 in currently pending U.S. Appl. No. 10/438,213.
- Office Action dated Apr. 16, 2010 in corresponding Japanese Patent Application No. 2006-525471 (with summary).
- USPTO Office Action dated Oct. 10, 2007 in U.S. Appl. No. 7,465,800.
- Bernstein, 2002, Polymorphism in Molecular Crystals, Clarendon Press, 115-18.
- Bernstein, 2004, "Crystal Structure Prediction and Polymorphism," *ACA Transactions*, 39, 14-23.
- Bond et al., 2002, "Controlling Crystal Architecture in Molecular Solids: the Supramolecular Approach," in *Supramolecular Organization and Materials Design*, Jones & Rao eds., Cambridge University Press, Chapter 12, pp. 391-443.
- Brittain, 1999, "Methods for the Characterization of Polymorphs and Solvates," in *Polymorphism in Pharmaceutical Solids*, Brittain ed., Chapter 6, pp. 227-278.
- Brittain, 2002, "Polymorphism: Pharmaceutical Aspects," in *Encyclopedia of Pharmaceutical Technology*, 2nd Edition, vol. 3, Swarbrick & Boylan eds., pp. 2239-2249.
- U.S. Appl. No. 60/499,723, filed Sep. 4, 2003, Jaworsky.
- U.S. Appl. No. 60/372,348, filed Apr. 12, 2002, Hariri et al.
- U.S. Appl. No. 09/545,654, filed Apr. 10, 2000, D'Amato.
- U.S. Appl. No. 09/287,377, filed Apr. 7, 1999, D'Amato.
- Brittain, 2007, "Polymorphism and Solvatomorphism 2005," *Journal of Pharmaceutical Sciences*, 96(4), 705-28.
- Bruns et al., 1984, "Thermochemical Investigation of Theophylline, Theophylline Hydrate and Their Aqueous Solutions," *Thermochimica Acta*, 72, 31-40.
- Byrn et al., 1999, "Drugs as Molecular Solids," in *Solid-State Chemistry of Drugs*, 2nd Edition, Chapter 1, pp. 3-43.
- Cruz Cabeza et al., 2006, "Prediction and Observation of Isostructural Induced by Solvent Incorporation in Multicomponent Crystals," *J. Am. Chem. Soc.*, 128, 14466-67.
- Cruz Cabeza et al., 2007, "Importance of Molecular Shape for the Overall Stability of Hydrogen Bond Motifs in the Crystal Structures of Various Carbamazepine-Type Drug Molecules," *Crystal Growth & Design*, 7(1), 100-07.

US 7,855,217 B2Page 6

- Day et al., 2004, "An Assessment of Lattice Energy Minimization for the Prediction of Molecular Organic Crystal Structures," *Crystal Growth & Design*, 4(6), 1327-40.
- Day et al., 2006, "Investigating the Latent Polymorphism of Maleic Acid," *Chemical Communications*, 54-56.
- Dean, 1993, *Analytical Chemistry Handbook*, 10.24-10.26.
- Gadamasetti et al., 2007, "Process Chemistry in the Pharmaceutical Industry," in *Encyclopedia of Pharmaceutical Technology*, 3rd Edition, vol. 5, Swarbrick ed., pp. 2993-3007.
- Jones et al., 2006, "Pharmaceutical Cocrystals: An Emerging Approach to Physical Property Enhancement," *MRS Bulletin*, 31, 875-79.
- Kirk-Othmer, 2002, "Crystallization," in *Encyclopedia of Chemical Technology*, vol. 8, pp. 95-147.
- Knapman, 2000, "Polymorphic Predictions," *Modern Drug Discovery*, 3(2), 53-57.
- Office Action dated Jun. 18, 2010 in U.S. Appl. No. 12/220,336.
- Richardson et al., "A Multi-Center, Randomized, Phase II Study To Evaluate the Efficacy and Safety of Two CDC-5013 Dose Regimens When Used Alone or in Combination with Dexamethasone (Dex) for the Treatment of Relapsed or Refractory Multiple Myeloma (MM)," *Blood*, Journal of the American Society of Hematology, Dec. 6-10, 2002, 100(11):Abstract #386.
- Seddon, 2004, "Pseudopolymorph: A Polemic," *Crystal Growth & Design*, 4(6), 1087.
- Sharma et al., 2006, "Toxicity Profile of the Immunomodulatory Thalidomide Analogue, Lenalidomide: Phase I Clinical Trial of Three Dosing Schedules in Patients with Solid Malignancies," *European Journal of Cancer*, 42, 2318-25.
- Vippagunta et al., 2001, "Crystalline Solids," *Advanced Drug Delivery Reviews*, 48, 3-26.
- Caira, 1998, "Crystalline Polymorphism of Organic Compounds," *Topics in Current Chemistry*, 198, 163-208.
- Fauci et al, 1998, "Harrison's Principles of Internal Medicine, Chronic Visual Loss," in *Harrison's Principles of Internal Medicine*, 168.
- Notice of Allowability mailed Nov. 3, 2008 in U.S. Appl. No. 10/934,863.
- Office Action mailed Jun. 17, 2010 in U.S. Appl. No. 10/557,302.

U.S. Patent

Dec. 21, 2010

Sheet 1 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM A




Fig. 1

U.S. Patent

Dec. 21, 2010

Sheet 2 of 48

US 7,855,217 B2

IR SPECTRUM OF FORM A




Fig. 2

U.S. Patent

Dec. 21, 2010

Sheet 3 of 48

US 7,855,217 B2

RAMAN SPECTRUM OF FORM A




Fig. 3

U.S. Patent

Dec. 21, 2010

Sheet 4 of 48

US 7,855,217 B2




Fig. 4

U.S. Patent

Dec. 21, 2010

Sheet 5 of 48

US 7,855,217 B2

MOISTURE SORPTION/DESORPTION ISOTHERM FOR FORM A




Fig. 5

U.S. Patent

Dec. 21, 2010

Sheet 6 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM B




Fig. 6

U.S. Patent

Dec. 21, 2010

Sheet 7 of 48

US 7,855,217 B2

IR SPECTRUM OF FORM B




Fig. 7

U.S. Patent

Dec. 21, 2010

Sheet 8 of 48

US 7,855,217 B2

RAMAN SPECTRUM OF FORM B




Fig. 8

U.S. Patent

Dec. 21, 2010

Sheet 9 of 48

US 7,855,217 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM B




Fig. 9

U.S. Patent

Dec. 21, 2010

Sheet 10 of 48

US 7,855,217 B2

TG-IR RESULTS FOR FORM B





Fig. 10

U.S. Patent

Dec. 21, 2010

Sheet 11 of 48

US 7,855,217 B2



U.S. Patent

Dec. 21, 2010

Sheet 12 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM C




Fig. 12

U.S. Patent

Dec. 21, 2010

Sheet 13 of 48

US 7,855,217 B2

IR SPECTRUM OF FORM C




Fig. 13

U.S. Patent

Dec. 21, 2010

Sheet 14 of 48

US 7,855,217 B2

RAMAN SPECTRUM OF FORM C




Fig. 14

U.S. Patent

Dec. 21, 2010

Sheet 15 of 48

US 7,855,217 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM C




Fig. 15

U.S. Patent

Dec. 21, 2010

Sheet 16 of 48

US 7,855,217 B2

TG-IR RESULTS FOR FORM C




Fig. 16

U.S. Patent

Dec. 21, 2010

Sheet 17 of 48

US 7,855,217 B2




Fig. 17

U.S. Patent

Dec. 21, 2010

Sheet 18 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM D




Fig. 18

U.S. Patent

Dec. 21, 2010

Sheet 19 of 48

US 7,855,217 B2

IR SPECTRUM OF FORM D




Fig. 19

U.S. Patent

Dec. 21, 2010

Sheet 20 of 48

US 7,855,217 B2

RAMAN SPECTRUM OF FORM D




Fig. 20

U.S. Patent

Dec. 21, 2010

Sheet 21 of 48

US 7,855,217 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM D




Fig. 21

U.S. Patent

Dec. 21, 2010

Sheet 22 of 48

US 7,855,217 B2

MOISTURE SORPTION/DESORPTION ISOTHERM FOR FORM D




Fig. 22

U.S. Patent

Dec. 21, 2010

Sheet 23 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM E





Fig. 23

U.S. Patent

Dec. 21, 2010

Sheet 24 of 48

US 7,855,217 B2

*Fig. 24*

U.S. Patent

Dec. 21, 2010

Sheet 25 of 48

US 7,855,217 B2




Fig. 25

U.S. Patent

Dec. 21, 2010

Sheet 26 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM F




Fig. 26

U.S. Patent

Dec. 21, 2010

Sheet 27 of 48

US 7,855,217 B2

DSC THERMOGRAM FOR FORM F




Fig. 27

U.S. Patent

Dec. 21, 2010

Sheet 28 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM G




Fig. 28

U.S. Patent

Dec. 21, 2010

Sheet 29 of 48

US 7,855,217 B2

DSC THERMOGRAM FOR FORM G




Fig. 29

U.S. Patent

Dec. 21, 2010

Sheet 30 of 48

US 7,855,217 B2

XRPD PATTERN OF FORM H




Fig. 30

U.S. Patent

Dec. 21, 2010

Sheet 31 of 48

US 7,855,217 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM H





Fig. 31

U.S. Patent

Dec. 21, 2010

Sheet 32 of 48

US 7,855,217 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5274-104-B

Date: 06-04-04 16:10 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 32

File: Process 5274-100-C XRPD PATTERN OF POLYMORPH B

Date: 06-04-04 16:11 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 33

U.S. Patent

Dec. 21, 2010

Sheet 33 of 48

US 7,855,217 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5274-104-B

Date: 06-04-04 16:10 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 34

XRPD PATTERN OF POLYMORPH E

File: Process 5274-100-C

Date: 06-04-04 16:11 Step: 0.020° Cnt Time: 0.500 Sec.

CPS Range: 1.50 - 40.00 (Deg) Step Scan Rate: 2.40 Deg/min




Fig. 35

U.S. Patent

Dec. 21, 2010

Sheet 34 of 48

US 7,855,217 B2

XRPD PATTERN OF POLYMORPH MIXTURE




Fig. 36

U.S. Patent

Dec. 21, 2010

Sheet 35 of 48

US 7,855,217 B2

TGA CURVE OF POLYMORPH B




Fig. 37




Fig. 38

U.S. Patent

Dec. 21, 2010

Sheet 36 of 48

US 7,855,217 B2




Fig. 39




Fig. 40

U.S. Patent

Dec. 21, 2010

Sheet 37 of 48

US 7,855,217 B2

TGA CURVE OF POLYMORPH MIXTURE





Fig. 41

U.S. Patent


Dec. 21, 2010

Sheet 38 of 48

US 7,855,217 B2

*Fig. 42***DSC OF POLYMORPH B**

Sample: 5274-100-C DSC
 Size: 0.0000 mg Run Date: 02-Jun-04 17:01
 Method: Cell constant calibration Instrument: DSC Q 1000 V7.3 Build 249
 Comment: OAI batch 1675C 10vol H₂O 24b 3b dry



*Fig. 43*

U.S. Patent

Dec. 21, 2010

Sheet 39 of 48

US 7,855,217 B2

*Fig. 44**Fig. 45*

U.S. Patent

Dec. 21, 2010

Sheet 40 of 48

US 7,855,217 B2




Fig. 46

U.S. Patent

Dec. 21, 2010

Sheet 41 of 48

US 7,855,217 B2




Fig. 47

U.S. Patent

Dec. 21, 2010

Sheet 42 of 48

US 7,855,217 B2

UV-VIS SCAN OF 0.04MG/ML SOLUTION

0.04mg/mL in Dissolution Medium




Fig. 48

U.S. Patent

Dec. 21, 2010

Sheet 43 of 48

US 7,855,217 B2

UV-VIS SCAN OF 0.008MG/ML SOLUTION

0.008mg/mL in Dissolution Medium




Fig. 49

U.S. Patent

Dec. 21, 2010

Sheet 44 of 48

US 7,855,217 B2

CALIBRATION CURVE FOR
3-(4-AMINO-1-OXO-1,3-DIHYDROISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE




Fig. 50

U.S. Patent

Dec. 21, 2010

Sheet 45 of 48

US 7,855,217 B2




Fig. 51

U.S. Patent

Dec. 21, 2010

Sheet 46 of 48

US 7,855,217 B2

SOLUBILITY OF FORM B




Fig. 52

U.S. Patent

Dec. 21, 2010

Sheet 47 of 48

US 7,855,217 B2




FIG. 53

U.S. Patent

Dec. 21, 2010

Sheet 48 of 48

US 7,855,217 B2

INTRINSIC DISSOLUTION OF FORMS A, B, AND E (EXPERIMENT NO.2)




FIG. 54

US 7,855,217 B2

1

**POLYMORPHIC FORMS OF
3-(4-AMINO-1-OXO-1,3
DIHYDRO-ISOINDOL-2-YL)-
PIPERIDINE-2,6-DIONE**

This application is a divisional application of U.S. patent application Ser. No. 10/934,863, filed Sep. 3, 2004, now U.S. Pat. No. 7,465,800 presently pending, which claims the benefit of U.S. provisional application 60/499,723, filed Sep. 4, 2003, the contents of each of which are incorporated by reference herein in their entireties.

1. FIELD OF THE INVENTION

This invention relates to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use for the treatment of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer.

2. BACKGROUND OF THE INVENTION

Many compounds can exist in different crystal forms, or polymorphs, which exhibit different physical, chemical, and spectroscopic properties. For example, certain polymorphs of a compound may be more readily soluble in particular solvents, may flow more readily, or may compress more easily than others. See, e.g., P. DiMartino, et al., *J. Thermal Anal.*, 48:447-458 (1997). In the case of drugs, certain solid forms may be more bioavailable than others, while others may be more stable under certain manufacturing, storage, and biological conditions. This is particularly important from a regulatory standpoint, since drugs are approved by agencies such as the U.S. Food and Drug Administration only if they meet exacting purity and characterization standards. Indeed, the regulatory approval of one polymorph of a compound, which exhibits certain solubility and physico-chemical (including spectroscopic) properties, typically does not imply the ready approval of other polymorphs of that same compound.

Polymorphic forms of a compound are known in the pharmaceutical arts to affect, for example, the solubility, stability, flowability, fractability, and compressibility of the compound, as well as the safety and efficacy of drug products comprising it. See, e.g., Knapman, K. *Modern Drug Discoveries*, 2000, 53. Therefore, the discovery of new polymorphs of a drug can provide a variety of advantages.

U.S. Pat. Nos. 5,635,517 and 6,281,230, both to Muller et al., disclose 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which is useful in treating and preventing a wide range of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer. New polymorphic forms of 3-(4 amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can further the development of formulations for the treatment of these chronic illnesses, and may yield numerous formulation, manufacturing and therapeutic benefits.

3. SUMMARY OF THE INVENTION

This invention encompasses polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. In certain aspects, the invention provides polymorphs of the compound identified herein as forms A, B, C, D, E, F, G, and H. The invention also encompasses mixtures of these forms. In further embodiments, this invention provides methods of making, isolating and characterizing the polymorphs.

2

This invention also provides pharmaceutical compositions and single unit dosage forms comprising a polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The invention further provides methods for the treatment or prevention of a variety of diseases and disorders, which comprise administering to a patient in need of such treatment or prevention a therapeutically effective amount of a polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

4. BRIEF DESCRIPTION OF THE DRAWINGS

Specific aspects of the invention can be understood with reference to the attached figures:

FIG. 1 provides a representative X-ray powder diffraction (XRPD) pattern of Form A;

FIG. 2 provides a representative IR spectrum of Form A;

FIG. 3 provides a representative Raman spectrum of Form A;

FIG. 4 provides a representative thermogravimetric analysis (TGA) curve and a representative differential scanning calorimeter (DSC) thermogram of Form A;

FIG. 5 provides a representative moisture sorption/desorption isotherm of Form A;

FIG. 6 provides a representative XRPD pattern of Form B;

FIG. 7 provides a representative IR spectrum of Form B;

FIG. 8 provides a representative Raman spectrum of Form B;

FIG. 9 provides a representative TGA curve and a representative DSC thermogram of Form B;

FIG. 10 provides representative TG-IR results of Form B;

FIG. 11 provides a representative moisture sorption/desorption isotherm of Form B;

FIG. 12 provides a representative XRPD pattern of Form C;

FIG. 13 provides a representative IR spectrum of Form C;

FIG. 14 provides a representative Raman spectrum of Form C;

FIG. 15 provides a representative TGA curve and a representative DSC thermogram of Form C;

FIG. 16 provides representative TG-IR results of Form C;

FIG. 17 provides a representative moisture sorption/desorption isotherm of Form C;

FIG. 18 provides a representative XRPD pattern of Form D;

FIG. 19 provides a representative IR spectrum of Form D;

FIG. 20 provides a representative Raman spectrum of Form D;

FIG. 21 provides a representative TGA curve and a representative DSC thermogram of Form D;

FIG. 22 provides a representative moisture sorption/desorption isotherm of Form D;

FIG. 23 provides a representative XRPD pattern of Form E;

FIG. 24 provides a representative TGA curve and a representative DSC thermogram of Form E;

FIG. 25 provides a representative moisture sorption/desorption isotherm of Form E;

FIG. 26 provides a representative XRPD pattern for a sample of Form F;

FIG. 27 provides a representative thermogram of Form F;

FIG. 28 provides a representative XRPD pattern of Form G;

FIG. 29 provides a representative DSC thermogram for a sample of Form G;

FIG. 30 provides a representative XRPD pattern of Form H;

US 7,855,217 B2

3

FIG. 31 provides a representative TGA curve and a representative DSC thermogram of Form H;

FIG. 32 provides a representative XRPD pattern of Form B;

FIG. 33 provides a representative XRPD pattern of Form B;

FIG. 34 provides a representative XRPD pattern of Form B;

FIG. 35 provides a representative XRPD pattern of Form E;

FIG. 36 provides a representative XRPD pattern of polymorph mixture;

FIG. 37 provides a representative TGA curve of Form B;

FIG. 38 provides a representative TGA curve of Form B;

FIG. 39 provides a representative TGA curve of Form B;

FIG. 40 provides a representative TGA curve of Form E;

FIG. 41 provides a representative TGA curve of polymorph mixture;

FIG. 42 provides a representative DSC thermogram of Form B;

FIG. 43 provides a representative DSC thermogram of Form B;

FIG. 44 provides a representative DSC thermogram of Form B;

FIG. 45 provides a representative DSC thermogram of Form E;

FIG. 46 provides a representative DSC thermogram of polymorph mixture;

FIG. 47 provides a UV-V is scan of dissolution medium;

FIG. 48 provides a UV-Vis scan of 0.04 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 49 provides a UV-Vis scan of 0.008 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 50 provides a calibration curve for 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione;

FIG. 51 provides a solubility curve of Form A;

FIG. 52 provides a solubility curve of Form B;

FIG. 53 provides an intrinsic dissolution of Forms A, B and E; and

FIG. 54 provides an intrinsic dissolution of Forms A, B and E.

5. DETAILED DESCRIPTION OF THE INVENTION

5.1 Definitions

As used herein and unless otherwise indicated, the terms “treat,” “treating” and “treatment” refer to the alleviation of a disease or disorder and/or at least one of its attendant symptoms.

As used herein and unless otherwise indicated, the terms “prevent,” “preventing” and “prevention” refer to the inhibition of a symptom of a disease or disorder or the disease itself.

As used herein and unless otherwise indicated, the terms “polymorph” and “polymorphic form” refer to solid crystalline forms of a compound or complex. Different polymorphs of the same compound can exhibit different physical, chemical and/or spectroscopic properties. Different physical properties include, but are not limited to stability (e.g., to heat or light), compressibility and density (important in formulation and product manufacturing), and dissolution rates (which can affect bioavailability). Differences in stability can result from changes in chemical reactivity (e.g., differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical characteristics (e.g., tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both

4

(e.g., tablets of one polymorph are more susceptible to breakdown at high humidity). Different physical properties of polymorphs can affect their processing. For example, one polymorph might be more likely to form solvates or might be more difficult to filter or wash free of impurities than another due to, for example, the shape or size distribution of particles of it.


Polymorphs of a molecule can be obtained by a number of methods known in the art. Such methods include, but are not limited to, melt recrystallization, melt cooling, solvent recrystallization, desolvation, rapid evaporation, rapid cooling, slow cooling, vapor diffusion and sublimation. Polymorphs can be detected, identified, classified and characterized using well-known techniques such as, but not limited to, differential scanning calorimetry (DSC), thermogravimetry (TGA), X-ray powder diffractometry (XRPD), single crystal X-ray diffractometry, vibrational spectroscopy, solution calorimetry, solid state nuclear magnetic resonance (NMR), infrared (IR) spectroscopy, Raman spectroscopy, hot stage optical microscopy, scanning electron microscopy (SEM), electron crystallography and quantitative analysis, particle size analysis (PSA), surface area analysis, solubility, and rate of dissolution.

As used herein to refer to the spectra or data presented in graphical form (e.g., XRPD, IR, Raman and NMR spectra), and unless otherwise indicated, the term “peak” refers to a peak or other special feature that one skilled in the art would recognize as not attributable to background noise. The term “significant peaks” refers to peaks at least the median size (e.g., height) of other peaks in the spectrum or data, or at least 1.5, 2, or 2.5 times the median size of other peaks in the spectrum or data.

As used herein and unless otherwise indicated, the term “substantially pure” when used to describe a polymorph of a compound means a solid form of the compound that comprises that polymorph and is substantially free of other polymorphs of the compound. A representative substantially pure polymorph comprises greater than about 80% by weight of one polymorphic form of the compound and less than about 20% by weight of other polymorphic forms of the compound, more preferably greater than about 90% by weight of one polymorphic form of the compound and less than about 10% by weight of the other polymorphic forms of the compound, even more preferably greater than about 95% by weight of one polymorphic form of the compound and less than about 5% by weight of the other polymorphic forms of the compound, and most preferably greater than about 97% by weight of one polymorphic form of the compound and less than about 3% by weight of the other polymorphic forms of the compound.

5.2 Polymorphic Forms

This invention is directed to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which has the structure shown below:



This compound can be prepared according to the methods described in U.S. Pat. Nos. 6,281,230 and 5,635,517, the entireties of which are incorporated herein by reference. For

US 7,855,217 B2

5

example, the compound can be prepared through catalytic hydrogenation of 3-(4-nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. 3-(4-Nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by allowing 2,6-dioxopiperidin-3-ammonium chloride to react with methyl 2-bromomethyl-4-nitrobenzoate in dimethylformamide in the presence of triethylamine. The methyl 2-bromomethyl-4-nitrobenzoate in turn is obtained from the corresponding methyl ester of nitro-ortho-toluic acid by conventional bromination with N-bromosuccinimide under the influence of light.

Polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by techniques known in the art, including solvent recrystallization, desolvation, vapor diffusion, rapid evaporation, slow evaporation, rapid cooling and slow cooling. Polymorphs can be made by dissolving a weighed quantity of 3-(4-amino-1-oxo-1,3 dihydro isoindol-2-yl)-piperidine-2,6-dione in various solvents at elevated temperatures. The solutions of the compound can then be filtered and allowed to evaporate either in an open vial (for fast hot evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation). Polymorphs can also be obtained from slurries. Polymorphs can be crystallized from solutions or slurries using several methods. For example, a solution created at an elevated temperature (e.g., 60° C.) can be filtered quickly then allowed to cool to room temperature. Once at room temperature, the sample that did not crystallize can be moved to a refrigerator then filtered. Alternatively, the solutions can be crash cooled by dissolving the solid in a solvent at an increased temperature (e.g., 45-65° C.) followed by cooling in a dry ice/solvent bath.

One embodiment of the invention encompasses Form A of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form A is an unsolvated, crystalline material that can be obtained from non-aqueous solvent systems. Another embodiment of the invention encompasses Form B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form B is a hemihydrated, crystalline material that can be obtained from various solvent systems. Another embodiment of the invention encompasses Form C of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form C is a hemisolvated crystalline material that can be obtained from solvents such as, but not limited to, acetone. Another embodiment of the invention encompasses Form D of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form D is a crystalline, solvated polymorph prepared from a mixture of acetonitrile and water. Another embodiment of the invention encompasses Form E of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form E is a dihydrated, crystalline material. Another embodiment of the invention encompasses Form F of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form F is an unsolvated, crystalline material that can be obtained from the dehydration of Form E. Another embodiment of the invention encompasses Form G of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form G is an unsolvated, crystalline material that can be obtained from slurring forms B and E in a solvent such as, but not limited to, tetrahydrofuran (THF). Another embodiment of the invention encompasses Form H of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form H is a partially hydrated crystalline material that can be obtained by exposing Form E to 0% relative humidity. Each of these forms is discussed in detail below.

Another embodiment of the invention encompasses a composition comprising amorphous 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione and crystalline 3-(4-

6

amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione of form A, B, C, D, E, F, G or H. Specific compositions can comprise greater than about 50, 75, 90 or 95 weight percent crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Another embodiment of the invention encompasses a composition comprising at least two crystalline forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (e.g., a mixture of polymorph forms B and E).

5.2.1 Form A

The data described herein for Form A, as well as for Forms B-H, were obtained using the experimental methods described in Examples 6.3-6.7, provided below.

Form A can be obtained from various solvents, including, but not limited to 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and THF. FIG. 1 shows a representative XRPD pattern of Form A. The pattern is characterized by peaks, preferably significant peaks, at approximately 8, 14.5, 16, 17.5, 20.5, 24, and 26 degrees 2θ.

Representative IR and Raman spectra data are provided in FIGS. 2 and 3.

Representative thermal characteristics of Form A are shown in FIG. 4. TGA data show a small weight increase up to about 150° C., indicating an unsolvated material. Weight loss above 150° C. is attributed to decomposition. The DSC curve of Form A exhibits an endotherm at about 270° C.

Representative moisture sorption and desorption data are plotted in FIG. 5. Form A does not exhibit a significant weight gain from 5 to 95% relative humidity. Equilibrium can be obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it has typically lost only about 0.003% by weight from start to finish. Form A is capable of remaining a crystalline solid for about 11 days when stored at about 22, 45, 58, and 84% relative humidity.

Interconversion studies show that Form A can convert to Form B in aqueous solvent systems and can convert to Form C in acetone solvent systems. Form A tends to be stable in anhydrous solvent systems. In water systems and in the presence of Form E, Form A tends to convert to Form E.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% relative humidity (RH) and 40° C./93% RH), Form A typically does not convert to a different form.

In sum, Form A is a crystalline, unsolvated solid that melts at approximately 270° C. Form A is weakly or not hygroscopic and appears to be the most thermodynamically stable anhydrous polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione discovered thus far.

5.2.2 Form B

Form B can be obtained from many solvents, including, but not limited to, hexane, toluene, and water. FIG. 6 shows a representative XRPD pattern of Form B, characterized by peaks at approximately 16, 18, 22 and 27 degrees 2θ.

Solution proton NMR confirm that Form B is a form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Representative IR and Raman spectra are shown in FIGS. 7 and 8, respectively. Compared to Form A, the IR spectrum for Form B has peaks at approximately 3513 and 1960 cm⁻¹.

Representative DSC and TGA data for Form B are shown in FIG. 9. The DSC curve exhibits endotherms at about 146 and 268° C. These events are identified as dehydration and melting by hot stage microscopy experiments. Form B typically loses about 3.1% volatiles up to about 175° C. (per

US 7,855,217 B2

7

approximately 0.46 moles of water). Comparison of the IR spectrum of the volatiles with that of water indicates that they are water (See FIG. 10). Calculations from TGA data indicate that Form B is a hemihydrate. Karl Fischer water analysis also supports this conclusion.

Representative moisture sorption and desorption data are shown in FIG. 11. Form B typically does not exhibit a significant weight gain from 5% to 95% relative humidity, when equilibrium is obtained at each relative humidity step. As Form B dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it typically has gained only about 0.022% by weight (about 0.003 mg) from start to finish Form B does not convert to a different form upon exposure to about 84% relative humidity for about ten days.

Interconversion studies show that Form B typically converts to Form A in a THF solvent system, and typically converts to Form C in an acetone solvent system. In aqueous solvent systems such as pure water and 10% water solutions, Form B is the most stable of the polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. However, it can convert to Form E in the presence of water. Desolvation experiments show that upon heating at about 175° C. for about five minutes, Form B typically converts to Form A.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH), Form B does not convert to a different form.

In sum, Form B is a hemihydrated, crystalline solid which has a DSC thermogram exhibiting endotherms at about 146 and about 268° C. Interconversion studies show that Form B converts to Form E in aqueous solvent systems, and converts to other forms in acetone and other anhydrous systems.

5.2.3 Form C

Form C can be obtained from evaporation, slurries and slow cools in acetone solvent systems. A representative XRPD pattern of this form is shown in FIG. 12. The data are characterized by peaks at approximately 15.5 and 25 degrees 20.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 13 and 14, respectively. The IR spectrum of Form C is characterized by peaks at approximately 3466, 3373, and 3318 cm⁻¹. The Raman spectrum of Form C is characterized by peaks at about 3366, 3321, 1101, and 595 cm⁻¹.

Representative thermal characteristics for Form C are plotted in FIG. 15. Form C loses about 10.02% volatiles up to about 175° C., indicating it is a solvated material. Weight loss above about 175° C. is attributed to decomposition. Identification of volatiles in Form C can be accomplished with TG-IR experiments. The representative IR spectrum captured after several minutes of heating, as depicted in FIG. 13, when compared with a spectral library, shows acetone to be the best match. Calculations from TGA data show that Form C is a hemisolvate (approximately 0.497 moles of acetone). The DSC curve for Form C, shown in FIG. 15, exhibits endotherms at about 150 and about 269° C. The endotherm at about 150° C. is attributed to solvent loss based on observations made during hot stage microscopy experiments. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption balance data are shown in FIG. 17. Form C does not exhibit a significant weight gain from 5 to 85% relative humidity, when equilibrium is obtained at each relative humidity step up to

8

85% relative humidity. At 95% relative humidity, Form C experiences a significant weight loss of about 6.03%. As the sample dries from 95% back down to 5% relative humidity, the sample maintains the weight achieved at the end of the adsorption phase at each step down to 5% relative humidity. Form C is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form C typically converts to Form A in a THF solvent system and typically converts to Form E in an aqueous solvent system. In an acetone solvent system, Form C is the most stable form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Desolvation experiments performed on Form C show that upon heating at about 150° C. for about five minutes, Form C will typically convert to Form A.

In sum, Form C is a crystalline, hemisolvated solid, which has a DSC thermogram exhibiting endotherms at about 150 and about 269° C. Form C is not hygroscopic below about 85% RH, but can convert to Form B at higher relative humidities.

5.2.4 Form D

Form D can be obtained from evaporation in acetonitrile solvent systems. A representative XRPD pattern of the form is shown in FIG. 18. The pattern is characterized by peaks at approximately 27 and 28 degrees 20.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 19 and 20, respectively. The IR spectrum of Form D is characterized by peaks at approximately 3509, 2299, and 2256 cm⁻¹. The Raman spectrum of Form D is characterized by peaks at approximately 2943, 2889, 2297, 2260, 1646, and 1150 cm⁻¹.

Representative thermal characteristics for Form D are plotted in FIG. 21. Form D loses about 6.75% volatiles up to about 175° C., indicating a solvated material. Weight loss above about 175° C. is attributed to decomposition. TG-IR experiments indicate that the volatiles are water and acetonitrile. Calculations from TG data show that about one mole of water 40 is present in the sample. A representative DSC curve for Form D exhibits endotherms at about 122 and about 270° C. The endotherm at about 122° C. is attributed to loss of volatiles based on observations made during hot stage microscopy experiments. The endotherm at about 270° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 22. Form D does not exhibit a significant weight gain from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it maintains its weight such that at 5% relative humidity the form has typically gained only about 0.39% by weight (about 0.012 mg) from start to finish. Form A is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form D is capable of converting to Form A in a THF solvent system, to Form E in an aqueous solvent system, and to Form C in an acetone solvent system. Desolvation experiments performed on Form D show that upon heating at about 150° C. for about five minutes Form D will typically convert to Form A.

In sum, Form D is a crystalline solid, solvated with both water and acetonitrile, which has a DSC thermogram exhibiting endotherms at about 122 and about 270° C. Form D is either weakly or not hygroscopic, but will typically convert to Form B when stressed at higher relative humidities.

US 7,855,217 B2

9

5.2.5 Form E

Form E can be obtained by slurring 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in water and by a slow evaporation of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in a solvent system with a ratio of about 9:1 acetone:water. A representative XRPD pattern is shown in FIG. 23. The data are characterized by peaks at approximately 20, 24.5 and 29 degrees 2θ.

Representative thermal characteristics of Form E are plotted in FIG. 24. Form E typically loses about 10.58% volatiles up to about 125° C., indicating that it is a solvated material. A second weight loss of an additional about 1.38% was observed between about 125° C. and about 175° C. Weight loss above about 175° C. is attributed to decomposition. Karl Fischer and TG-IR experiments support the conclusion that the volatile weight loss in Form E is due to water. The representative DSC curve for Form E exhibits endotherms at about 99, 161 and 269° C. Based on observations made during hot stage microscopy experiments, the endotherms at about 99 and about 161° C. are attributed to loss of volatiles. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 25. Form E typically does not exhibit a significant weight change from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the sample dried from 95% back down to 5% relative humidity, the sample continues to maintain weight such that at 5% relative humidity the sample has lost only about 0.0528% by weight from start to finish.

Interconversion studies show that Form E can convert to Form C in an acetone solvent system and to Form G in a THF solvent system. In aqueous solvent systems, Form E appears to be the most stable form. Desolvation experiments performed on Form E show that upon heating at about 125° C. for about five minutes, Form E can convert to Form B. Upon heating at 175° C. for about five minutes, Form B can convert to Form F.

When stored for a period of 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH) Form E typically does not convert to a different form. When stored for seven days at room temperature/0% RH, Form E can convert to a new form, Form H.

5.2.6 Form F

Form F can be obtained by complete dehydration of Form E. A representative XRPD pattern of Form F, shown in FIG. 26, is characterized by peaks at approximately 19, 19.5 and 25 degrees 2θ.

Representative thermal characteristics of Form F are shown in FIG. 27. The representative DSC curve for Form F exhibits an endotherm at about 269° C. preceded directly by two smaller endotherms indicative of a crystallized form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The DSC thermogram does not show any thermal events prior to the melt, suggesting that it is an unsolvated material.

5.2.7 Form G

Form G can be obtained by slurring forms B and E in THF. A representative XRPD pattern of this form, shown in FIG. 28, is characterized by a peak at approximately 23 degrees 2θ. Two other peaks unique to Form G appear at approximately 21 and 24.5 degrees 2θ.

Representative thermal characteristics of Form G are plotted in FIG. 29. A representative DSC curve for Form G exhibits an endotherm at about 248° C. followed by a small, broad exotherm at about 267° C. No thermal events are seen

10

in the DSC thermogram at lower temperatures, suggesting that it is an unsolvated material.

5.2.8 Form H

Form H can be obtained by storing Form E at room temperature and 0% RH for about 7 days. A representative XRPD pattern is shown in FIG. 30. The pattern is characterized by a peak at 15 degrees 2θ, and two other peaks at 26 and 31 degrees 2θ.

Representative thermal characteristics are shown in FIG. 31. Form H loses about 1.67% volatiles up to about 150° C. Weight loss above about 150° C. is attributed to decomposition. Karl Fischer data shows that Form H typically contains about 1.77% water (about 0.26 moles), suggesting that the weight loss seen in the TG is due to dehydration. The DSC thermogram shows a broad endotherm between about 50° C. and about 125° C., corresponding to the dehydration of Form H and a sharp endotherm at about 269° C., which is likely due to a melt.

When slurried in water with either Forms A or B, after about 14 days Form H can convert to Form E. When slurried in THF, Form H can convert to Form A. When slurried in acetone, Form H can convert to Form C.

In sum, Form H is a crystalline solid, hydrated with about 0.25 moles of water, which has a DSC thermogram exhibiting an endotherm between about 50 and 125° C. and an endotherm at about 269° C.

5.3 Methods of Use and Pharmaceutical Compositions

Polymorphs of the invention exhibit physical characteristics that are beneficial for drug manufacture, storage or use. All polymorphs of the invention have utility as pharmaceutically active ingredients or intermediates thereof.

This invention encompasses methods of treating and preventing a wide variety of diseases and conditions using polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. In each of the methods, a therapeutically or prophylactically effective amount of the compound is administered to a patient in need of such treatment or prevention. Examples of such disease and conditions include, but are not limited to, diseases associated with undesired angiogenesis, cancer (e.g., solid and blood borne tumors), inflammatory diseases, autoimmune diseases, and immune diseases.

Examples of cancers and pre-cancerous conditions include those described in U.S. Pat. Nos. 6,281,230 and 5,635,517 to Muller et al. and in various U.S. patent applications to Zeldis, including application Ser. Nos. 10/411,649, filed Apr. 11, 2003 (Treatment of Myelodysplastic Syndrome); 10/438,213 filed May 15, 2003 (Treatment of Various Types of Cancer); 10/411,656, filed Apr. 11, 2003 (Treatment of Myeloproliferative Diseases). Examples of other diseases and disorders that can be treated or prevented using compositions of the invention are described in U.S. Pat. Nos. 6,235,756 and 6,114,335 to D'Amato and in other U.S. patent applications to Zeldis, including Ser. Nos. 10/693,794, filed Oct. 23, 2003 (Treatment of Pain Syndrome) and 10/699,154, filed Oct. 30, 2003 (Treatment of Macular Degeneration). The entirety of each of the patents and patent applications cited herein is incorporated herein by reference.

Depending on the disease to be treated and the subject's condition, polymorphs of the invention can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implantation), inhalation spray, nasal, vaginal, rectal, sublingual, or topical routes of administration and may be formulated, alone or together, in suitable dosage unit

US 7,855,217 B2

11

formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of administration. Because individual polymorphs have different dissolution, stability, and other properties, the optimal polymorph used in methods of treatment may depend on the route of administration. For example, forms that are readily soluble in aqueous solutions are preferably used to provide liquid dosage forms, whereas forms that exhibit great thermal stability may be preferred in the manufacture of solid dosage forms (e.g., tablets and capsules).

Although the physical characteristics of polymorphs can, in some cases, affect their bioavailability, amounts of the polymorphs that are therapeutically or prophylactically effective in the treatment of various disease and conditions can be readily determined by those of ordinary skill in the pharmacy or medical arts. In certain embodiments of the invention, a polymorph is administered orally and in a single or divided daily doses in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day. In other embodiments, a polymorph is administered every other day in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day.

The invention encompasses pharmaceutical compositions and single unit dosage forms that can be used in methods of treatment and prevention, which comprise one or more polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and optionally one or more excipients or diluents. Specific compositions and dosage forms are disclosed in the various patents and patent applications incorporated herein by reference. In one embodiment, a single dosage form comprises a polymorph (e.g., Form B) in an amount of about 5, 10, 25 or 50 mg.

6. EXAMPLES**6.1 Polymorph Screen**

A polymorph screen to generate the different solid forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was carried out as follows.

A weighed sample of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (usually about 10 mg) was treated with aliquots of the test solvent. Solvents were either reagent or HPLC grade. The aliquots were usually about 200 µL. Between additions, the mixture was usually shaken or sonicated. When the solids dissolved, as judged by visual inspection, estimated solubilities were calculated. Solubilities were estimated from these experiments based on the total solvent used to provide a solution. Actual solubilities may have been greater than those calculated due to the use of too-large solvent aliquots or to a slow rate of dissolution.

Samples were created by generating solutions (usually about 30 mg in 20 mL) at elevated temperatures, filtering, and allowing the solution to evaporate whether in an open vial (hot fast evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation).

Slurry experiments were also performed. Usually about 25 mg of solid was placed in either 3 or 5 mL of solvent. The samples were then placed on orbital shakers at either ambient temperature or 40° C. for 410 days.

Crystallizations were performed using various cooling methods. Solid was dissolved in a solvent at an elevated temperature (e.g., about 60° C.), filtered quickly and allowed to cool to room temperature. Once at room temperature, samples that did not crystallize were moved to a refrigerator. Solids were removed by filtration or decantation and allowed

12

to dry in the air. Crash cools were performed by dissolving solid in a solvent at an increased temperature (e.g., about 45-65° C.) followed by cooling in a dry ice/acetone bath.

Hygroscopicity studies were performed by placing portions of each polymorph in an 84% relative humidity chamber for approximately one week.

Desolvation studies were carried out by heating each polymorph in a 70° C. oven for approximately one week.

Interconversion experiments were carried out by making slurries containing two forms in a saturated solvent. The slurries were agitated for approximately 7-20 days at ambient temperature. The insoluble solids were recovered by filtration and analyzed using XRPD.

15 6.2 Preparation of Polymorphic Forms

Eight solid forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione were prepared as described below.

20 Form A was obtained by crystallization from various non-aqueous solvents including 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and tetrahydrofuran. Form B was also obtained by crystallization from the solvents hexane, toluene and water. Form C was obtained from evaporation, slurries, and slow cools in acetone solvent systems. Form D was obtained from evaporation in acetonitrile solvent systems. Form E was obtained most readily by slurring 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in water. Form F was obtained by complete desolvation of Form E. It is found to be an unsolvated, crystalline material that melts at about 269° C. Form G was obtained by slurring forms B and E in THF. Form H was obtained by stressing Form E at room temperature and 0% RH for 7 days.

25 35 6.2.1 Synthesis of Polymorphs B and E

Form B is the desired polymorph for the active pharmaceutical ingredient (API) of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. This form has been used in the formulation of API into drug product for clinical studies. Three batches were produced as apparent mixtures of polymorphs in the non-micronized API of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

40 Development work was carried out to define a process that would generate polymorph B from this mixture of polymorphs and could be implemented for strict polymorphic controls in the validation batches and future manufacturing of API of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Characterization of polymorphic forms produced during the work was performed by XRPD, DSC, TGA and KF.

45 **50** A process was also developed for the large-scale preparation of Form E. Polymorph E material was prepared in order to carry out a comparison with polymorph B drug product in capsule dissolution testing of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. 150 g of a mixture of polymorphs in 3 L of water was stirred at room temperature for 48 hours. The product was collected by filtration and dried at 25° C. for 24 hours under vacuum. XRPD, DSC, TGA, KF and HPLC analyses confirmed that the material isolated was polymorph E.

55 In a preliminary work, it was demonstrated that stirring a suspension of a mixture of polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione with water at high temperature (75° C.) for an extended period of time converted this mixture of polymorphs exclusively to form B. Several specific parameters were identified including temperature, solvent volume and drying parameters (temperature

US 7,855,217 B2

13

and vacuum). XRPD, DSC, TGA, KF and HPLC analyses were used to characterize all of the batches. After completing the optimization work, the optimized process was scaled-up to 100-200 g on three lots of API. Drying studies were carried out at 20° C., 30° C. and 40° C., and 65° C. with a vacuum of 150 mm of Hg. The results are shown in Tables 1-5.

The cooling and holding periods of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione slurry were studied. The experimental laboratory data suggests that polymorph B seems to be forming first, and overtime equilibration to polymorph E at RT conditions occurs, therefore generating a mixture of polymorphs B and E. This result supports the fact that polymorph B seems to be a kinetic product, and that prolonged processing time converts the material to polymorph E resulting in a mixture of polymorphs B and E.

A laboratory procedure was developed to exclusively produce polymorph B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The procedure includes a stirred 10 volume water slurry at ~75° C. for 6-24 hours. The following preferred process parameters have been identified:

1. Hot slurry temperature of 70-75° C.
2. Product filtration of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)piperidine-2,6-dione at 65-75° C.
3. Drying under vacuum at 60-70° C. is preferred for an efficient removal of unbound water in 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione wet cake.
4. The filtration step of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione may be a time sensitive operation. The use of efficient solid-liquid separation equipment is preferred.
5. Holding periods of water-wet cake of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at KF higher than 5% may cause the kinetic equilibrations of polymorph B to mixed polymorphs of E and B.

Drying to KF<4.0% water was achieved in ~3 hours (30-70° C., 152 mm Hg). Polymorphs B and E were distinguished by the water levels as measured by KF and TGA. The reference sample of polymorph B is micronized API. In order to make accurate comparison by XRPD samples were gently grinded before submission for analysis. This increases the clarity of the identification of the polymorphic form. All samples were analyzed for XRPD, DSC, TGA, KF and HPLC.

TABLE 1

Preliminary Studies			
Amount	Reaction conditions	Analysis	Results/conclusion
2 g	Water, rt, 48 h	XRPD, DSC, TGA, KF	Polymorph E
25 g	Water, rt, 48 h	XRPD, DSC, TGA, KF	Polymorph E
5 g	Water, 70-75° C., 24 h then rt 24 h	XRPD, DSC, TGA, KF	Polymorph B
1 g	9:1 Acetone - water, Slow evapo.	XRPD, DSC, TGA, KF	Polymorph Mixture
1 g	175° C. 1 h in an oven	XRPD, DSC, TGA, KF	Polymorph A
0.5 g (polymorph A)	Water, rt, 24 h	XRPD, DSC, TGA, KF	Polymorph E
1 g polymorph B	Water, rt, 48 h	XRPD, DSC, TGA, KF	Polymorph E
1 g polymorph E	Water, 70-75° C., 24 h	XRPD, DSC, TGA, KF	Polymorph B
1 g	Slurry in heptane	XRPD, DSC, TGA, KF	No change

14

TABLE 2

Optimization of Temperature, Time and Solvent Volume				
Amount	Amount Water (mL)	Temp (° C.)	Time (h)	Results/conclusion
10 g	50	75	6	Mix
10 g	50	75	24	Polymorph B
10 g	100	70	6	Polymorph B
10 g	100	70	14	Polymorph B
10 g	100	70	21	Polymorph B
10 g	100	75	6	Polymorph B
10 g	100	75	24	Polymorph B
10 g	100	75	6	Polymorph B
10 g	100	75	19	Polymorph B
10 g	100	75	14	Polymorph B
10 g	100	75	24	Polymorph B
5 g	100	75	18	Polymorph B
10 g	100	80	6	Polymorph B
10 g	100	80	20	Polymorph B
10 g	200	45	6	Polymorph B + E
10 g	200	45	24	Polymorph E
10 g	200	60	48	Polymorph B
10 g	200	75	6	Mix
10 g	200	75	24	Polymorph B
10 g	200	75	13	Polymorph B
10 g	200	75	24	Polymorph B

Optimum conditions were determined to be 10 volumes of solvent (H_2O), 70-80° C. for 6-24 hours.

TABLE 3

Holding Time				
Amount	Reaction Conditions	Holding Time (h)	Holding Temp (° C.)	Results/Conclusion
5 g	Water, 70-75° C., 24 h	24	23-25	Polymorph B
1 g	Water, 70-75° C., 24 h	48	23-25	Polymorph E
Polymorph B	2 g Water, 40 mL	16	23-25	Polymorph E
	150 g Water, 3.0 L	24	23-25	Polymorph E
	150 g Water, 3.0 L	48	23-25	Polymorph E
	10 g Water, 100 mL, 24 h, 75° C.	18	23-25	Polymorph B
	10 g Water, 100 mL, 24 h, 75° C.	18	40	Polymorph B
	10 g Water, 200 mL, 24 h, 75° C.	14	-5	Mix
	10 g Water, 200 mL, 24 h, 75° C.	14	23-25	Polymorph E
	10 g Water, 200 mL, 24 h, 75° C.	14	40	Mix
	10 g Water, 100 mL, 24 h, 75° C.	21	23-25	Polymorph E
	10 g Water, 100 mL, 24 h, 75° C.	21	40	Mix

Holding time gave mixed results and it was determined that the material should be filtered at 60-65° C. and the material washed with 0.5 volume of warm (50-60° C.) water.

TABLE 4

Scale-up Experiments				
Amount	Amount Water (L)	Temp (° C.)	Time (h)	Results/Conclusion
100 g	1.0	75	6	Polymorph B
100 g	1.0	75	22	Polymorph B

US 7,855,217 B2

15

TABLE 4-continued

Scale-up Experiments				
Amount	Amount Water (L)	Temp (° C.)	Time (h)	Results/Conclusion
100 g	1.0	75	6	Polymorph B
100 g	1.0	75	24	Polymorph B
100 g	1.0	75	6	Polymorph B
100 g	1.0	75	22	Polymorph B

TABLE 5

Drying Studies					
Amount	Drying Time (h)	Drying Temp (° C.)	Vacuum (mm Hg)	KF§ (%)	Results/Conclusion
100 g	0	—	—	3.690	Polymorph B
100 g	3	30	152	3.452	Polymorph B
100 g	8	30	152	3.599	Polymorph B
100 g	0	—	—	3.917	Polymorph B
100 g	5	40	152	3.482	Polymorph B
100 g	22	40	152	3.516	Polymorph B
100 g	3	40	152	3.67	Polymorph B
100 g	22	40	152	3.55	Polymorph B

*Reaction Conditions: Water 1 L, 75° C., 22-24 h;

§Average of 2 runs.

Drying studies determined that the material should be dried at 35-40° C., 125-152 mm Hg for 3 to 22 h or until the water content reaches ≤4% w/w.

For a large scale preparation of polymorph E (5222-152-B), a 5-L round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (150 g, 0.579 mol) and water (3000 mL, 20 vol). The mixture was mechanically stirred at room temperature (23-25° C.) for 48 h under nitrogen atmosphere.

Samples were taken after 24 h and 48 h before the mixture was filtered and air-dried on the filter for 1 h. The material was transferred to a drying tray and dried at room temperature (23-25° C.) for 24 h. KF analysis on the dried material showed water content of 11.9%. The material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph E.

For a large scale preparation of polymorph B (5274-104), a 2 L-3-necked round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (polymorph mixture, 100 g, 0.386 mol) and water (1000 mL, 10.0 vol). The mixture was heated to 75° C. over approximately 30 minutes with mechanical stirring under nitrogen atmosphere.

Samples were taken after 6 h and 24 h before the mixture was allowed to cool to 60-65° C., filtered and the material washed with warm (50-60° C.) water (50 mL, 0.5 vol). The material was transferred to a drying tray and dried at 30° C., 152 mm Hg for 8 h. KF analysis on the dried material showed water content of 3.6%. After grinding the material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph B. The results of the analyses are shown in FIGS. 32-46.

6.3 X-Ray Powder Diffraction Measurements

X-ray powder diffraction analyses were carried out on a Shimadzu XRD 6000 X-ray powder diffractometer using Cu K α radiation. The instrument is equipped with a fine-focus X-ray tube. The tube voltage and amperage were set at 40 kV

16

and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm. Diffracted radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 2.5 degrees 2θ to 40 degrees 2θ was used. A silicon standard was analyzed each day to check the instrument alignment.

X-ray powder diffraction analyses were also carried out using Cu K α radiation on an Inel XRG-3000 diffractometer 10 equipped with a curved position-sensitive detector. Data were collected in real time over a theta-two theta range of 120° at a resolution of 0.03°. The tube voltage and current were 40 kV and 30 mA, respectively. A silicon standard was analyzed each day to check for instrument alignment. Only the region between 2.5 and 40 degrees 2θ is shown in the figures.

6.4 Thermal Analysis

TG analyses were carried out on a TA Instrument TGA 2050 or 2950. The calibration standards were nickel and alumel. Approximately 5 mg of sample was placed on a pan, accurately weighed, and inserted into the TG furnace. The samples were heated in nitrogen at a rate of 10° C./min, up to a final temperature of 300 or 350° C.

DSC data were obtained on a TA 2920 instrument. The calibration standard was indium. Approximately 2-5 mg samples were placed into a DSC pan and the weight accurately recorded. Crimped pans with one pinhole were used for analysis and the samples were heated under nitrogen at a rate of 10° C./min, up to a final temperature of 350° C.

Hot-stage microscopy was carried out using a Kofler hot stage mounted on a Leica Microscope. The instrument was calibrated using USP standards.

A TA Instruments TGA 2050 interfaced with a Nicolet 35 model 560 Fourier transform IR spectrophotometer, equipped with a globar source, XT/KBr beamsplitter, and deuterated triglycine sulfate (DTGS) detector, was utilized for TG-IR experiments. The IR spectrometer was wavelength calibrated with polystyrene on the day of use while the TG 40 was temperature and weight calibrated biweekly, using indium for the temperature calibration. A sample of approximately 10 mg of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was weighed into an aluminum pan and heated from 25 to 30° C. to 200° C. at a rate of 20° C./min 45 with a helium purge. IR spectra were obtained in series, with each spectrum representing 32 co-added scans at a resolution of 4 cm $^{-1}$. Spectra were collected with a 17-second repeat time. TG/IR analysis data are presented as Gram-Schmidt plots and IR spectra linked to the time. Gram-Schmidt plots show total IR intensity vs. time; hence, the volatiles can be identified at each time point. They also show when the volatiles are detected. From the Gram-Schmidt plots, time points 50 were selected and the IR spectra of these time points are presented in the stacked linked spectra. Each spectrum identifies volatiles evolving at that time point. Volatiles were identified from a search of the HR Nicolet TGA vapor phase spectral library. The library match results are also presented 55 to show the identified vapor.

6.5 Spectroscopy Measurements

Raman spectra were acquired on a Nicolet model 750 Fourier transform Raman spectrometer utilizing an excitation wavelength of 1064 nm and approximately 0.5 W of Nd:YAG 65 laser power. The spectra represent 128 to 256 co-added scans acquired at 4 cm $^{-1}$ resolution. The samples were prepared for analysis by placing the material in a sample holder and posi-

US 7,855,217 B2

17

tioning this in the spectrometer. The spectrometer was wavelength calibrated using sulfur and cyclohexane at the time of use.

The mid-IR spectra were acquired on a Nicolet model 860 Fourier transform IR spectrophotometer equipped with a globar source XT/KBr beamsplitter and a deuterated triglycine sulfate (DTGS) detector. A Spectra-Tech, Inc. diffuse reflectance accessory was utilized for sampling. Each spectrum represents 128 co-added scans at a spectral resolution of 4 cm⁻¹. A background data set was acquired with an alignment mirror in place. A single beam sample data set was then acquired. Subsequently, a log 1/R (where R=reflectance) spectrum was acquired by rationing the two data sets against each other. The spectrophotometer was calibrated (wavelength) with polystyrene at the time of use.

6.6 Moisture Sorption/Desorption Measurements

Moisture sorption/desorption data were collected on a VTI SGA-100 moisture balance system. For sorption isotherms, a sorption range of 5 to 95% relative humidity (RH) and a desorption range of 95 to 5% RH in 10% RH increments was used for analysis. The sample was not dried prior to analysis. Equilibrium criteria used for analysis were less than 0.0100 weight percent change in 5 minutes with a maximum equilibration time of 3 hours if the weight criterion was not met. Data were not corrected for the initial moisture content of the samples.

6.7 Solution Proton NMR Measurements

NMR spectra not previously reported were collected at SSCI, Inc, 3065 Kent Avenue, West Lafayette, Ind. Solution phase ¹H NMR spectra were acquired at ambient temperature on a Bruker model AM spectrometer. The ¹H NMR spectrum represents 128 co added transients collected with a 4 μ sec pulse and a relaxation delay time of 5 seconds. The free induction decay (FID) was exponentially multiplied with a 0.1 Hz Lorentzian line broadening factor to improve the signal-to-noise ratio. The NMR spectrum was processed utilizing GRAMS software, version 5.24. Samples were dissolved in dimethyl sulfoxide-d₆.

The scope of this invention can be understood with reference to the appended claims.

6.8 Intrinsic Dissolution and Solubility Studies

Intrinsic dissolution experiments were conducted on Form A (anhydrous), Form B (hemihydrate), and Form E (dihydrate) of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Equilibrium solubility experiments were conducted on Forms A and B. Aliquots were analyzed by ultraviolet-visible spectrophotometry, and the solids remaining from each experiment were analyzed by X-ray powder diffraction (XRPD).

6.8.1 Experimental

6.8.1.1 Dissolution

Dissolution experiments were carried out in a VanKel VK6010-8 dissolution apparatus equipped with a VK650A heater/circulator. An intrinsic dissolution apparatus (Woods apparatus) was used. Samples were compressed at 1.5 metric tons (1000 psi) for 1 min using the Woods apparatus in a hydraulic press, giving a sample surface of 0.50 cm². A dissolution medium consisting of 900 mL HCl buffer, pH 1.8, with 1% sodium lauryl sulfate, was used for each experiment. The medium was degassed by vacuum filtration through a 0.22- μ m nylon filter disk and maintained at 37° C. The appa-

18

ratus was rotated at 50 rpm for each experiment. Aliquots were filtered immediately using 0.2- μ M nylon syringe filters. In some cases, the undissolved solids were recovered and analyzed by X-ray powder diffraction (XRPD).

6.8.1.2 Solubility

Equilibrium solubility experiments were conducted in a 100-mL, three-neck, round-bottom flask immersed in a constant temperature oil bath maintained at 25° C. A solid sample of 400-450 mg was stirred in 50 mL of dissolution medium (HCl buffer, pH 1.8, with 1% sodium lauryl sulfate) using a mechanical stir rod. Aliquots were filtered using 0.2- μ m nylon syringe filters and immediately diluted 1 mL → 50 mL, then 5 mL → 25 mL with dissolution medium in Class A glassware, a final dilution factor of 250.

6.8.1.3 UV-Vis Spectrophotometry

Dissolution and solubility samples solutions were analyzed by a Beckman DU 640 single-beam spectrophotometer. A 1.000-cm quartz cuvette and an analysis wavelength of 228.40 nm were utilized. The detector was zeroed with a cuvette filled with dissolution medium.

6.8.1.4 X-Ray Powder Diffraction

XRPD analyses were carried out on a Shimadzu XRD-6000 X-ray powder diffractometer using Cu K α radiation. The instrument is equipped with a fine focus X-ray tube. The tube power and amperage were set at 40 kV and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm. Diffracted radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 2.5 to 40° was used. A silicon standard was analyzed each day to check the instrument alignment. Samples were packed in an aluminum holder with silicon insert.

6.8.2 Results

The results of these solubility and intrinsic studies are summarized in Table 6. Both the solubility and dissolution experiments were conducted in a medium of HCl buffer, pH 1.8, containing 1% sodium lauryl sulfate. Form A was found to be unstable in the medium, converting to Form E. The solubilities of Forms A, B, and E were estimated to be 6.2, 5.8, and 4.7 mg/mL, respectively. The dissolution rates of Forms A, B, and E were estimated to be 0.35, 0.34, and 0.23 mg/mL, respectively.

6.8.2.1 UV-Vis Spectrophotometry Method Development

A UV-V is scan of the dissolution medium (blanked with an empty cuvette) was done to identify any interfering peaks. A small peak at 225 nm was present as shown in FIG. 47.

Solutions of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at varying concentrations were analyzed by UV-V is spectrophotometry. A preliminary scan of a 1.0 mg/mL solution was done, with the instrument blanked with dissolution medium. The solution was highly absorbing and noisy from 200-280 nm, making dilution necessary.

A 0.04 mg/mL solution of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was then scanned from 200-300 nm. The plot was still noisy between 200 and 230 nm as shown in FIG. 48. The sample was further diluted to 0.008 mg/mL. A wavelength scan of 200-350 nm for this sample showed a peak a 228.4 nm with no interference, as shown in FIG. 49. Therefore, a wavelength of 228.4 was chosen for analysis of the solubility and dissolution samples.

A six-point calibration curve was generated with standards of the following concentrations: 0.001 mg/mL, 0.002 mg/mL, 0.005 mg/mL, 0.010 mg/mL, 0.015 mg/mL, and 0.020 mg/mL (Notebook 569-90). A linearity coefficient of R²=0.9999 was obtained as shown in FIG. 50.

US 7,855,217 B2

19

20

6.8.2.2 Solubility

A sample consisting of 449.4 mg Form A was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, and 150 min. The concentration reached 6.0 mg/mL by the first time point. The highest concentration reached was 6.2 mg/mL, at 30 min. From that point the concentration decreased, reaching 4.7 mg/mL at 150 min as in FIG. 51. The solids remaining at the final time point were analyzed by XRPD and found to be Form E as shown in Table 7. No peaks attributed to Form A can be seen in the pattern. Since the concentration did not plateau at 4.7 mg/mL, the solubility of Form E may be lower than that.

A sample consisting of 401.4 mg Form B was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, 180, 420, and 650 min. Form B dissolved much more slowly than Form A, reaching 3.3 mg/mL in 90 min. The concentration stabilized at 5.6-5.7 mg/mL at the final three time points as in FIG. 52. The remaining solids were shown to be Form B as in Table 7, suggesting Form B has good stability in water.

A summary of the solubilities is given in Table 6. The amounts dissolved at each time point are shown in Tables 8 and 9.

TABLE 6

Summary of Results				
Form	Solubility	Intrinsic Dissolution #1	Intrinsic Dissolution #2	Average Intrinsic Dissolution Rate
Form A	6.2 mg/mL	0.35	0.22 ^a	0.29 ^a
Form B	5.8 mg/mL	0.35	0.32	0.34
Form E	4.7 mg/mL	0.21	0.25	0.23

^aThe Form A dissolution experiment #2 may have converted to Form E on the surface of the disk, skewing the average rate lower.

TABLE 7

Experimental Details	
Experiment	Final Form
Pressed Form A	A
Pressed Form B	B
Form A Solubility	E
Form B Solubility	B
Form A Dissolution	—
Form A Dissolution	A
Form B Dissolution	—
Form B Dissolution	B
Form E Dissolution	E
Form E Dissolution	—

TABLE 8

Form A Solubility	
Time Point (min)	Concentration (mg/mL)
7	6.00
15	6.11
30	6.16
60	6.10
90	5.46
150	4.73

TABLE 9

Form B Solubility	
Time Point (min)	Concentration (mg/mL)
7	1.63
15	2.14
30	2.33
60	2.94
90	3.34
180	5.67
420	5.76
650	5.61

6.8.2.3 Intrinsic Dissolution

Approximately 200 mg each of Forms A and B were compressed into disks in the Woods apparatus using 2 metric tons of pressure. The samples were subsequently scraped out, ground gently, and analyzed by XRPD. The study showed that compression and grinding does not cause a form change in either case. (See Table 7).

Two preliminary dissolution runs were performed. The disks fractured to some extent in both experiments, compromising the requirement of constant surface area.

The first experiment of intrinsic dissolution that strictly followed the USP chapter on intrinsic dissolution utilized approximately 150 mg each of Forms A and B. Seven aliquots, beginning at 5 min and ending at 90 min, were taken to maintain sink conditions. The experiment resulted in linear dissolution profiles, with a rate of 0.35 mg per cm² per minute for both forms. The Form E experiment was done later under the same conditions and added to the graph for comparison. (See FIG. 53). The Form E dissolution rate was 0.21 mg per cm² per minute, significantly lower than the dissolution rate of Forms A and B. This is in line with expectations based on the solubility data. The crystal form of the remaining solids did not change in any case.

The second experiment utilized approximately 250 mg each of Forms A and B. The Form E experiment (135 mg) was done later and added to the graph for comparison. (See FIG. 54). Nine aliquots were taken, beginning at 5 min and ending at 150 min. The dissolution rates were 0.22, 0.32, and 0.25 mg per cm² per minute, respectively, for Forms A, B, and E. The dissolution rate for Form A in this experiment was low, while the rates for Forms B and E were similar to those found in the first experiment. It is believed that in this case, a thin layer of the Form A sample disk may have converted to Form E upon exposure to water. This is supported by the evidence of rapid conversion of Form A to Form E in the solubility experiment. The diffraction pattern of the undissolved solids does not indicate a form change. However, the bulk of the sample disk is not exposed to water. Therefore, the true intrinsic dissolution rate of Form A is believed to be close to 0.35 mg per cm² per minute. An insufficient quantity of Form A was available to repeat the experiment.

A summary of the intrinsic dissolution rates is given in Table 6. The amounts dissolved at each time point are summarized in Tables 10 and 11.

TABLE 10

Intrinsic Dissolution Experiment #1 Results			
Time Point	Form A ^a	Form B ^a	Form E ^a
5 min	5.76	10.80 ^b	2.70
10 min	7.73	6.85	4.13

US 7,855,217 B2

21

22

TABLE 10-continued

Intrinsic Dissolution Experiment #1 Results			
Time Point	Form A ^a	Form B ^a	Form E ^a
20 min	11.31	10.25	6.96
30 min	15.59	14.35	9.60
45 min	21.98	20.57	12.57
60 min	27.11	25.70	15.16
90 min	34.17	34.34	20.82

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)^bThis data point not included in graph since the value is higher than the next two data points.

TABLE 11

Intrinsic Dissolution Experiment #2 Results			
Time Point	Form A ^a	Form B ^a	Form E ^a
5 min	4.50	5.04	3.06
10 min	5.22	6.12	4.31
20 min	7.54	7.73	11.40
30 min	11.46	12.72	11.93
45 min	15.01	17.33	14.72
60 min	18.38	21.93	18.52
90 min	24.38	31.64	26.24
120 min	30.35	41.31	33.56
150 min	35.26	49.54	40.82

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)

6.9 Analyses of Mixtures of Polymorphs

This invention encompasses mixtures of different polymorphs. For example, an X-ray diffraction analysis of one production sample yielded a pattern that contained two small peaks seen at approximately 12.6° and 25.8° 2θ in addition to those representative of Form B. In order to determine the composition of that sample, the following steps were performed.

- 1) Matching of the new production pattern to known forms along with common pharmaceutical excipients and contaminants;
- 2) Cluster analysis of the additional peaks to identify if any unknown phase is mixed with the original Form B;
- 3) Harmonic analysis of the additional peaks to identify if any preferred orientation may be present or if any changes in the crystal habit may have occurred; and
- 4) Indexing of the unit cells for both Form B and the new production sample to identify any possible crystallographic relationships.

Based on these tests, which can be adapted for the analysis of any mixture of polymorphs, it was determined that the sample contained a mixture of polymorph forms B and E.

6.10 Dosage Form

Table 12 illustrates a batch formulation and single dosage formulation for a 25 mg single dose unit of a polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

TABLE 12

Formulation for a 25 mg capsule				
Material	Percent By Weight	Quantity (mg/tablet)	Quantity (kg/batch)	
Polymorphic Form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione	40.0%	25 mg	16.80 kg	
Pregelatinized Corn Starch, NF	59.5%	37.2 mg	24.99 kg	
Magnesium Stearate	0.5%	0.31 mg	0.21 kg	
Total	100.0%	62.5 mg	42.00 kg	

The pregelatinized corn starch (SPRESS B-820) and polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione components are passed through a screen (i.e., a 710 μm screen) and then loaded into a Diffusion Mixer with a baffle insert and blended for about 15 minutes. The magnesium stearate is passed through a screen (i.e., a 210 μm screen) and added to the Diffusion Mixer. The blend is then encapsulated in capsules using a Dosator type capsule filling machine.

The entire scope of this invention is not limited by the specific examples described herein, but is more readily understood with reference to the appended claims.

What is claimed is:

1. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 80% by weight of the solid form.

2. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 90% by weight of the solid form.

3. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 95% by weight of the solid form.

4. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 97% by weight of the solid form.

5. A pharmaceutical composition comprising a therapeutically effective amount of the solid form of claim 1, 2, 3, or 4, and a pharmaceutically acceptable excipient or carrier.

6. The pharmaceutical composition of claim 5, which is a single unit dosage form.

7. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 5 mg.

8. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 10 mg.

9. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 25 mg.

10. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 50 mg.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,855,217 B2
APPLICATION NO. : 12/335395
DATED : December 21, 2010
INVENTOR(S) : Jaworsky et al.

Page 1 of 11

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Please replace Drawing Sheets 15, 26, 31, 32, 33, 35, 36, 38, 39, and 40 (of 48) with the Replacement Drawing Sheets on the following pages of this Certificate of Correction:

Page 2 of 11: Replacement Drawing Sheet 15 of 48 (Fig. 15);

Page 3 of 11: Replacement Drawing Sheet 26 of 48 (Fig. 26);

Page 4 of 11: Replacement Drawing Sheet 31 of 48 (Fig. 31);

Page 5 of 11: Replacement Drawing Sheet 32 of 48 (Figs. 32 and 33);

Page 6 of 11: Replacement Drawing Sheet 33 of 48 (Figs. 34 and 35);

Page 7 of 11: Replacement Drawing Sheet 35 of 48 (Figs. 37 and 38);

Page 8 of 11: Replacement Drawing Sheet 36 of 48 (Figs. 39 and 40);

Page 9 of 11: Replacement Drawing Sheet 38 of 48 (Figs. 42 and 43);

Page 10 of 11: Replacement Drawing Sheet 39 of 48 (Figs. 44 and 45);

Page 11 of 11: Replacement Drawing Sheet 40 of 48 (Fig. 46).

Signed and Sealed this
Third Day of May, 2011



David J. Kappos
Director of the United States Patent and Trademark Office

CERTIFICATE OF CORRECTION (continued)


Page 2 of 11

U.S. Patent

Dec. 21, 2010

Sheet 15 of 48

7,855,217 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM C
TGA

DSC




FIG. 15

CERTIFICATE OF CORRECTION (continued)

Page 3 of 11

U.S. Patent

Dec. 21, 2010

Sheet 26 of 48

7,855,217 B2

XRPD PATTERN OF FORM F




FIG. 26

CERTIFICATE OF CORRECTION (continued)

Page 4 of 11

U.S. Patent

Dec. 21, 2010

Sheet 31 of 48

7,855,217 B2

TGA (TOP) AND DSC (BOTTOM) OF FORM H




FIG.31

CERTIFICATE OF CORRECTION (continued)

Page 5 of 11

U.S. Patent

Dec. 21, 2010

Sheet 32 of 48


7,855,217 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5274-104-B

Date: 06-04-04 16:10 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.


**FIG.32**

XRPD PATTERN OF POLYMORPH B

File: Process 5274-100-C

Date: 06-02-04 16:11 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.

**FIG.33**

CERTIFICATE OF CORRECTION (continued)

Page 6 of 11

U.S. Patent

Dec. 21, 2010

Sheet 33 of 48

7,855,217 B2

XRPD PATTERN OF POLYMORPH B

File: Process 5222-157-C
Date: 06/04/04 15:07 Step: 0.020° Cnt Time: 0.500 Sec.
Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.34

XRPD PATTERN OF POLYMORPH E

File: Process 5222-152-B Form E
Date: 05/21/04 10:46 Step: 0.020° Cnt Time: 0.500 Sec.
Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.35

CERTIFICATE OF CORRECTION (continued)

Page 7 of 11

U.S. Patent

Dec. 21, 2010

Sheet 35 of 48

7,855,217 B2

TGA CURVE OF POLYMORPH B
 Sample: 5274-104-B
 Size 7.8320mg
 Method: Ramp
 TGA
 Run Date: 09-Jun-04 17:04
 Instrument: TGA 0.500 V5.3 Build 171




FIG.37

TGA CURVE OF POLYMORPH B
 Sample: 5274-100-C
 Size 10.9430mg
 Method: Ramp
 Comment: Evotec Batch 1675C H2O
 TGA
 Run Date: 03-Jun-04 11:20
 Instrument: TGA 0.500 V5.3 Build 171




FIG.38

CERTIFICATE OF CORRECTION (continued)

Page 8 of 11


U.S. Patent

Dec. 21, 2010


Sheet 36 of 48

7,855,217 B2

TGA CURVE OF POLYMORPH B
 Sample: 5222-157-C TGA
 Size: 24.4610mg
 Method: Ramp
 Comment: Evotec Batch 17 Run Date: 07-Jun-04 09:46
 Instrument: TGA 0.500 V5.3 Build 171

**FIG.39**

TGA CURVE OF POLYMORPH E
 Sample: 5222-152-B TGA
 Size: 11.3850mg
 Method: Ramp
 Comment: Form E Run Date: 21-May-04 09:34
 Instrument: TGA 0.500 V5.3 Build 171

**FIG.40**

CERTIFICATE OF CORRECTION (continued)

Page 9 of 11

U.S. Patent

Dec. 21, 2010


Sheet 38 of 48

7,855,217 B2

DSC OF POLYMORPH B
DSC

Sample: 5274-104-B
Size: 0.0000mg
Method: Cell constant calibration
Comment: O/W batch 15 22b


Run Date: 07-Jun-04 11:30
Instrument: DSC Q 1000 V7.3 Build 249

**FIG.42**

DSC OF POLYMORPH B
DSC

Sample: 5274-100-C
Size: 0.0000mg
Method: Cell constant calibration
Comment: Evotec Batch 1675C 10vol H₂O 24h 3h dry

Run Date: 02-Jun-04 17:01
Instrument: DSC Q 1000 V7.3 Build 249

**FIG.43**

CERTIFICATE OF CORRECTION (continued)

Page 10 of 11

U.S. Patent

Dec. 21, 2010


Sheet 39 of 48

7,855,217 B2

DSC OF POLYMORTH B
DSC

Sample: 5222-157-C
Size: 0.0000mg
Method: Cell constant calibration
Comment: OAN batch 17


Run Date: 07-Jun-04 09:45
Instrument: DSC Q 1000 V7.3 Build 249

**FIG.44**

DSC OF POLYMORPH E
DSC

Sample: 5222-152-B
Size: 3.4500mg
Method: Cell constant calibration
Comment: Evotec batch 15 Form E

Run Date: 21-May-04 09:32
Instrument: DSC Q 1000 V7.3 Build 249

**FIG.45**

CERTIFICATE OF CORRECTION (continued)

Page 11 of 11

U.S. Patent

Dec. 21, 2010

Sheet 40 of 48

7,855,217 B2

Sample: 5222-161-A

DSC OF POLYMORPH MIXTURE
DSC

Size: 0.0000mg

Method: Cell constant calibration

Run Date: 11-Jun-04 12:44

Comment: OAI batch 15

Instrument: DSC Q 1000 V7.3 Build 249




FIG.46