

# Linear Algebra HW #2

Ozaner Hansha

February 10, 2020

## Problem 1

**Problem:** Recall that an  $m \times n$  matrix  $A$  is called upper triangular if all entries lying below the diagonal entries are zero, that is, if  $a_{ij} = 0$  whenever  $i > j$ . Prove that the set of upper triangular matrices  $U$  form a subspace of  $\mathbb{F}^{m \times n}$ .

**Solution:** To show that  $U$  forms a subspace, we must verify that it:

- a) contains the zero element
- b) is closed under addition
- c) is closed under scalar multiplication

a) is easy to prove since the zero vector of  $\mathbb{F}^{m \times n}$  is simply the  $m \times n$  zero matrix  $\mathbf{0}$ . All entries of this matrix are the zero scalar  $0 \in \mathbb{F}$ , meaning that  $\mathbf{0}_{ij} = 0$  for all  $i, j$  and in particular when  $i > j$ . Thus  $\mathbf{0} \in U$ .

b) Given two upper triangular matrices  $\mathbf{a}, \mathbf{b} \in U$ , we have that for all  $i, j$  s.t.  $i > j$ :

$$a_{ij} = b_{ij} = 0$$

As is the definition of upper triangular matrix. And so their sum, defined element wise, is also 0 for all  $i > j$ :

$$a_{ij} + b_{ij} = 0 + 0 = 0$$

And thus  $\mathbf{a} + \mathbf{b} \in U$ , i.e.  $U$  is closed under addition.

c) Given an upper triangular matrix  $\mathbf{a} \in U$ , and a scalar  $\lambda \in \mathbb{F}$  we have that for all  $i, j$  s.t.  $i > j$ :

$$\lambda a_{ij} = \lambda \cdot 0 = 0$$

Which is precisely the definition of an upper triangular matrix. Thus  $\lambda \mathbf{a} \in U$ , i.e.  $U$  is closed under scalar multiplication.

## Problem 2

**Problem:** Let  $u$  and  $v$  be distinct vectors in a vector space  $V$ . Show that  $\{u, v\}$  is linearly dependent if and only if  $u$  or  $v$  is a multiple of the other.

**Solution:** Now recall that two vectors  $\mathbf{u}, \mathbf{v}$  are linearly independent iff there exist scalars  $\lambda_1, \lambda_2$ , where at least one is non-zero, such that:

$$\lambda_1 \mathbf{u} + \lambda_2 \mathbf{v} = \mathbf{0}$$

(Lemma 1) First note that if  $\mathbf{u}$  is a scalar multiple of  $\mathbf{v}$ , then  $\mathbf{v}$  is a scalar multiple of  $\mathbf{u}$  since all non-zero scalars have multiplicative inverses:

$$\mathbf{u} = \lambda \mathbf{v} \implies \mathbf{v} = \lambda^{-1} \mathbf{u}$$

And so it will suffice to show that at least one vector is a multiple of another.

( $\implies$ ) for the forward direction we will assume w.l.o.g. that  $\lambda_1 \neq 0$ . This nets us the following:

$$\begin{aligned} \lambda_1 \mathbf{u} + \lambda_2 \mathbf{v} = \mathbf{0} &\implies \lambda_1 \mathbf{u} = -\lambda_2 \mathbf{v} \\ &\implies \mathbf{u} = -\frac{\lambda_2}{\lambda_1} \mathbf{v} \end{aligned}$$

And so  $\mathbf{u}$  is a scalar multiple of  $\mathbf{v}$ . And by Lemma 1 we know this argument can be reversed for  $\mathbf{v}$  as well.

( $\impliedby$ ) for the backwards direction we will assume w.l.o.g. that  $\mathbf{u}$  is a scalar multiple of  $\mathbf{v}$ . This nets us the following:

$$\begin{aligned} \mathbf{u} = \lambda \mathbf{v} &\implies \mathbf{u} - \lambda \mathbf{v} = \mathbf{0} \\ &\implies 1 \cdot \mathbf{u} - \lambda \mathbf{v} = \mathbf{0} \end{aligned}$$

And so there is a pair of coefficients, namely  $\lambda_1 = 1$  and  $\lambda_2 = -\lambda$ , such that  $\lambda_1 \mathbf{u} + \lambda_2 \mathbf{v} = \mathbf{0}$ .

( $\Leftrightarrow$ ) And so these two implications, along with Lemma 1, give us the desired equivalence:

$$\lambda_1 \mathbf{u} + \lambda_2 \mathbf{v} = \mathbf{0} \iff (\exists \lambda_1) \mathbf{u} = \lambda_1 \mathbf{v} \vee (\exists \lambda_2) \mathbf{v} = \lambda_2 \mathbf{u}$$

### Problem 3

**Problem:** Show that if  $S_1$  and  $S_2$  are subsets of a vector space  $V$  then:

$$\text{span}(S_1 \cup S_2) = \text{span}(S_1) + \text{span}(S_2)$$

Where  $+$  denotes the sum of two subspaces.

**Solution:** To prove that these two sets are equal we must show that each is a subset of the other.

( $\subseteq$ ) Let  $\mathbf{v} \in \text{span}(S_1 \cup S_2)$ . This means that, by definition,  $\mathbf{v}$  is a linear combination of the vectors in  $S_1 \cup S_2$ :

$$\mathbf{v} = \overbrace{\sum_i a_i \mathbf{x}_i}^{\mathbf{x}} + \overbrace{\sum_j b_j \mathbf{y}_j}^{\mathbf{y}}$$

Where  $\mathbf{x}_i \in S_1$  and  $\mathbf{y}_j \in S_2$ . And since  $\mathbf{x}$  is a linear combination of vectors in  $S_1$  it is by definition a member of its span, and the same goes for  $\mathbf{y}$  and  $S_2$ :

$$\mathbf{x} \in \text{span}(S_1) \quad \mathbf{y} \in \text{span}(S_2)$$

And so  $\mathbf{v}$  can be expressed as the sum of a vector from  $\text{span}(S_1)$  and  $\text{span}(S_2)$  and thus is, by definition, an element of their sum:

$$\mathbf{v} = \mathbf{x} + \mathbf{y} \in \text{span}(S_1) + \text{span}(S_2)$$

( $\supseteq$ ) Now we instead let  $\mathbf{v} \in \text{span}(S_1) + \text{span}(S_2)$ . We can now write  $\mathbf{v}$  as the sum of two vectors  $\mathbf{x} \in \text{span}(S_1)$  and  $\mathbf{y} \in \text{span}(S_2)$ :

$$\mathbf{v} = \mathbf{x} + \mathbf{y}$$

Note that a generic element  $\mathbf{x}$  of  $S_1$  is just some linear combination of its vectors and similarly for  $\mathbf{y}$ :

$$\mathbf{v} = \overbrace{\sum_i a_i \mathbf{x}_i}^{\mathbf{x}} + \overbrace{\sum_j b_j \mathbf{y}_j}^{\mathbf{y}}$$

And at this point it should be clear that this is a generic element of  $\text{span}(S_1 \cup S_2)$  as it's a linear combination of vectors from  $S_1 \cup S_2$ . Thus  $\mathbf{v} \in \text{span}(S_1 \cup S_2)$

#### Problem 4

**Problem:** Is the following set linearly independent or dependent?

$$\left\{ \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix}, \begin{bmatrix} -2 & 6 \\ 4 & -8 \end{bmatrix} \right\} \subseteq \mathbb{R}^{2 \times 2}$$

**Solution:** As we've shown in problem 2, the linear dependence of a set of two vectors can be demonstrated by showing that one is a scalar multiple of the other. In this case, that scalar is given by  $-2 \in \mathbb{R}$ :

$$-2 \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} = \begin{bmatrix} -2 & 6 \\ 4 & -8 \end{bmatrix}$$

And so the set is linearly dependent.

#### Problem 5

**Problem:** Is the following set a basis for  $\mathcal{P}_2(\mathbb{R})$ ?

$$\{-x^2 + 2x + 1, x^2 - 2x + 4, -9x^2 + 18x - 1\}$$

**Solution:** Recalling that  $\mathcal{P}_2(\mathbb{R})$  is isomorphic to  $\mathbb{R}^3$ , we can identify each polynomial as a vector of its coefficients. Putting these vectors in a matrix, we can perform Gaussian elimination to determine the matrix's rank and thus the dimension the polynomials span:

$$\begin{aligned} \left[ \begin{array}{ccc|c} -1 & 2 & 1 & 0 \\ 1 & -2 & 4 & 0 \\ -9 & 18 & -1 & 0 \end{array} \right] & \xrightarrow{r_2+r_1} \left[ \begin{array}{ccc|c} -1 & 2 & 1 & 0 \\ 0 & 0 & 5 & 0 \\ -9 & 18 & -1 & 0 \end{array} \right] \\ & \xrightarrow{r_3-9r_1} \left[ \begin{array}{ccc|c} -1 & 2 & 1 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & -10 & 0 \end{array} \right] \\ & \xrightarrow{r_3+2r_2} \left[ \begin{array}{ccc|c} -1 & 2 & 1 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \end{aligned}$$

Once reduced, we can see that the matrix has only 2 pivot rows and thus is of rank 2. Clearly then, this set does not form a basis of the 3-dimensional  $\mathcal{P}_2(\mathbb{R})$  as it only spans 2 dimensions.

## Problem 6

**Problem:** Find a basis, and give the dimension, of the following two subspaces of  $\mathbb{F}^5$ :

- a)  $W_1 = \{(a_1, a_2, a_3, a_4, a_5)^\top \in \mathbb{F}^5 \mid a_1 - a_3 - a_4 = 0\}$   
 b)  $W_2 = \{(a_1, a_2, a_3, a_4, a_5)^\top \in \mathbb{F}^5 \mid a_2 = a_3 = a_4 \wedge a_1 + a_5 = 0\}$

**Solution:** a) For  $W_1$  we have the following basis:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Clearly each vector satisfies  $a_1 - a_3 - a_4$  and no vector is a linear combination of the others, i.e. the set is linearly independent. Now we will show that every vector  $\mathbf{v} \in W_1$  can be written as a linear combination of these 4 vectors:

$$\begin{aligned} \mathbf{v} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \end{bmatrix} &= \begin{bmatrix} a_3 + a_4 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \end{bmatrix} && (a_1 - a_3 - a_4 = 0) \\ &= a_3 \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + a_2 \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + a_4 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + a_5 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

And so these vectors form a basis of 4 vectors, thus  $W_1$  is 4-dimensional.

b) For  $W_2$  we have the following basis:

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} \right\}$$

Clearly both vectors satisfy  $a_2 = a_3 = a_4$  and  $a_1 + a_5 = 0$  and neither vector is a scalar multiple of the other, i.e. the set is linearly independent. Now we will show that every vector  $\mathbf{v} \in W_2$  can be written as a linear combination of these 2 vectors:

$$\begin{aligned}
\mathbf{v} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \end{bmatrix} &= \begin{bmatrix} a_1 \\ a_2 \\ a_2 \\ a_2 \\ a_5 \end{bmatrix} && (a_2 = a_3 = a_4) \\
&= \begin{bmatrix} a_1 \\ a_2 \\ a_2 \\ a_2 \\ -a_1 \end{bmatrix} && (a_1 + a_5 = 0) \\
&= a_1 \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ -1 \end{bmatrix} + a_2 \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}
\end{aligned}$$

And so these vectors form a basis of 2 vectors, thus  $W_2$  is 2-dimensional.