

CMP 694 Graph Theory  
Hacettepe University

## Lecture 9: Edge-coloring of graphs and Hamiltonian Cycles

Lecturer:  
Lale Özkahya

Resources:  
“Introduction to Graph Theory” by Douglas B. West

A  $k$ -edge coloring of a graph  $G$  is a coloring (labeling) of the edges of  $G$  using  $k$  colors.

# Definitions

A  $k$ -edge coloring of a graph  $G$  is a coloring (labeling) of the edges of  $G$  using  $k$  colors.

A coloring is called proper if incident edges have different colors.

# Definitions

A  *$k$ -edge coloring* of a graph  $G$  is a coloring (labeling) of the edges of  $G$  using  $k$  colors.

A coloring is called *proper* if incident edges have different colors.

A graph is  *$k$ -edge-colorable* if it has a proper  $k$ -edge coloring.

# Definitions

A  $k$ -edge coloring of a graph  $G$  is a coloring (labeling) of the edges of  $G$  using  $k$  colors.

A coloring is called proper if incident edges have different colors.

A graph is  $k$ -edge-colorable if it has a proper  $k$ -edge coloring. The edge chromatic number of  $G$ ,  $\chi'(G)$ , is the least  $k$  such that  $G$  is  $k$ -edge-colorable.

# Definitions

A  **$k$ -edge coloring** of a graph  $G$  is a coloring (labeling) of the edges of  $G$  using  $k$  colors.

A coloring is called **proper** if incident edges have different colors.

A graph is  **$k$ -edge-colorable** if it has a proper  $k$ -edge coloring. The **edge chromatic number of  $G$ ,  $\chi'(G)$** , is the least  $k$  such that  $G$  is  $k$ -edge-colorable.

**Observation:**  $\chi'(G) \geq \Delta(G)$  for all graphs.

# Definitions

A  **$k$ -edge coloring** of a graph  $G$  is a coloring (labeling) of the edges of  $G$  using  $k$  colors.

A coloring is called **proper** if incident edges have different colors.

A graph is  **$k$ -edge-colorable** if it has a proper  $k$ -edge coloring. The **edge chromatic number of  $G$ ,  $\chi'(G)$** , is the least  $k$  such that  $G$  is  $k$ -edge-colorable.

**Observation:**  $\chi'(G) \geq \Delta(G)$  for all graphs.

**Example:** Edge-coloring of  $K_{2n}$  is a modeling of scheduling problem.

# Bipartite Graphs, Petersen Graph

Theorem (König, 1916)

*If  $G$  is bipartite, then  $\chi'(G) = \Delta(G)$ .*

- Note that every bipartite graph is contained in a  $\Delta(G)$ -regular bipartite graph, call this larger graph  $G'$ .



# Bipartite Graphs, Petersen Graph

## Theorem (König, 1916)

*If  $G$  is bipartite, then  $\chi'(G) = \Delta(G)$ .*

- Note that every bipartite graph is contained in a  $\Delta(G)$ -regular bipartite graph, call this larger graph  $G'$ .
- Every regular bipartite graph has a 1-factor.

# Bipartite Graphs, Petersen Graph

## Theorem (König, 1916)

*If  $G$  is bipartite, then  $\chi'(G) = \Delta(G)$ .*

- Note that every bipartite graph is contained in a  $\Delta(G)$ -regular bipartite graph, call this larger graph  $G'$ .
- Every regular bipartite graph has a 1-factor.
- Remove 1-factors of  $G'$  one by one and let every one factor be the edges of one color class.

# Bipartite Graphs, Petersen Graph

## Theorem (König, 1916)

*If  $G$  is bipartite, then  $\chi'(G) = \Delta(G)$ .*

- Note that every bipartite graph is contained in a  $\Delta(G)$ -regular bipartite graph, call this larger graph  $G'$ .
- Every regular bipartite graph has a 1-factor.
- Remove 1-factors of  $G'$  one by one and let every one factor be the edges of one color class.
- This yields a proper  $\Delta(G)$ -coloring of  $G'$  and  $G$ .

# Bipartite Graphs, Petersen Graph

## Theorem (König, 1916)

*If  $G$  is bipartite, then  $\chi'(G) = \Delta(G)$ .*

- Note that every bipartite graph is contained in a  $\Delta(G)$ -regular bipartite graph, call this larger graph  $G'$ .
- Every regular bipartite graph has a 1-factor.
- Remove 1-factors of  $G'$  one by one and let every one factor be the edges of one color class.
- This yields a proper  $\Delta(G)$ -coloring of  $G'$  and  $G$ .

### Observation:

The chromatic number of Petersen graph is 4. (Note that if 3 colors were enough, then every color class would contain exactly five edges. Remove one matching and discuss the remaining graph.)

# Bipartite Graphs, Petersen Graph

## Theorem (König, 1916)

*If  $G$  is bipartite, then  $\chi'(G) = \Delta(G)$ .*

- Note that every bipartite graph is contained in a  $\Delta(G)$ -regular bipartite graph, call this larger graph  $G'$ .
- Every regular bipartite graph has a 1-factor.
- Remove 1-factors of  $G'$  one by one and let every one factor be the edges of one color class.
- This yields a proper  $\Delta(G)$ -coloring of  $G'$  and  $G$ .

### Observation:

The chromatic number of Petersen graph is 4. (Note that if 3 colors were enough, then every color class would contain exactly five edges. Remove one matching and discuss the remaining graph.)

## Theorem (Vizing, 1964)

*If  $G$  is a simple graph, then  $\chi'(G) \leq \Delta(G) + 1$ .*

# Bipartite Graphs, Petersen Graph

## Theorem (König, 1916)

*If  $G$  is bipartite, then  $\chi'(G) = \Delta(G)$ .*

- Note that every bipartite graph is contained in a  $\Delta(G)$ -regular bipartite graph, call this larger graph  $G'$ .
- Every regular bipartite graph has a 1-factor.
- Remove 1-factors of  $G'$  one by one and let every one factor be the edges of one color class.
- This yields a proper  $\Delta(G)$ -coloring of  $G'$  and  $G$ .

## Observation:

The chromatic number of Petersen graph is 4. (Note that if 3 colors were enough, then every color class would contain exactly five edges. Remove one matching and discuss the remaining graph.)

## Theorem (Vizing, 1964)

*If  $G$  is a simple graph, then  $\chi'(G) \leq \Delta(G) + 1$ .*

**Thus, there are two types of graphs: the ones that have edge-chromatic number  $\Delta(G)$  or  $\Delta(G) + 1$ .**

# Hamiltonian Cycles

A **Hamiltonian graph** is a graph with a spanning cycle, also called a **Hamiltonian cycle**.

# Hamiltonian Cycles

A **Hamiltonian graph** is a graph with a spanning cycle, also called a **Hamiltonian cycle**.

The problem on deciding whether a graph is hamiltonian or not is an **NP-complete** problem (no algorithm exists that runs in polynomial time).

So, there are known necessary conditions needed for a graph to be hamiltonian. Also, we know some sufficient conditions.

But, no “necessary and sufficient (if and only if)” is known.



# Hamiltonian Cycles

A **Hamiltonian graph** is a graph with a spanning cycle, also called a **Hamiltonian cycle**.

The problem on deciding whether a graph is hamiltonian or not is an **NP-complete** problem (no algorithm exists that runs in polynomial time).

So, there are known necessary conditions needed for a graph to be hamiltonian. Also, we know some sufficient conditions.

But, no “necessary and sufficient (if and only if)” is known.

**Proposition (A necessary condition)**

*If  $G$  has a Hamilton cycle, then for each nonempty set  $S \subset V$ , the graph  $G - S$  has at most  $|S|$  components.*

See Example 7.2.5 in West.

# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

Theorem (Dirac, 1952)

*If  $G$  is a simple graph with at least three vertices and  $\delta(G) \geq n(G)/2$ , then  $G$  is Hamiltonian.*

# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

**Theorem (Dirac, 1952)**

*If  $G$  is a simple graph with at least three vertices and  $\delta(G) \geq n(G)/2$ , then  $G$  is Hamiltonian.*

- Assume on the contrary that  $G$  is a maximal non-Hamiltonian graph that satisfies the minimum degree condition.

# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

## Theorem (Dirac, 1952)

*If  $G$  is a simple graph with at least three vertices and  $\delta(G) \geq n(G)/2$ , then  $G$  is Hamiltonian.*

- Assume on the contrary that  $G$  is a maximal non-Hamiltonian graph that satisfies the minimum degree condition.
- By the maximality of  $G$ , adding any other edge to  $G$  would create a Hamiltonian cycle. So, let  $uv \notin E(G)$ . There is a Ham. path  $v_1, v_2, \dots, v_n$  with ends  $u = v_1$  and  $v = v_n$ .

# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

## Theorem (Dirac, 1952)

*If  $G$  is a simple graph with at least three vertices and  $\delta(G) \geq n(G)/2$ , then  $G$  is Hamiltonian.*

- Assume on the contrary that  $G$  is a maximal non-Hamiltonian graph that satisfies the minimum degree condition.
- By the maximality of  $G$ , adding any other edge to  $G$  would create a Hamiltonian cycle. So, let  $uv \notin E(G)$ . There is a Ham. path  $v_1, v_2, \dots, v_n$  with ends  $u = v_1$  and  $v = v_n$ .
- **Fact:** If  $v_i \in N(v)$  and  $v_{i+1} \in N(u)$  for some  $1 < i < n-1$ , done.

# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

## Theorem (Dirac, 1952)

*If  $G$  is a simple graph with at least three vertices and  $\delta(G) \geq n(G)/2$ , then  $G$  is Hamiltonian.*

- Assume on the contrary that  $G$  is a maximal non-Hamiltonian graph that satisfies the minimum degree condition.
- By the maximality of  $G$ , adding any other edge to  $G$  would create a Hamiltonian cycle. So, let  $uv \notin E(G)$ . There is a Ham. path  $v_1, v_2, \dots, v_n$  with ends  $u = v_1$  and  $v = v_n$ .
- **Fact:** If  $v_i \in N(v)$  and  $v_{i+1} \in N(u)$  for some  $1 < i < n-1$ , done.
- We claim that there is such an  $i$ , let  $S = \{i : v_{i+1} \in N(u)\}$  and  $T = \{i : v_i \in N(v)\}$ .

# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

## Theorem (Dirac, 1952)

*If  $G$  is a simple graph with at least three vertices and  $\delta(G) \geq n(G)/2$ , then  $G$  is Hamiltonian.*

- Assume on the contrary that  $G$  is a maximal non-Hamiltonian graph that satisfies the minimum degree condition.
- By the maximality of  $G$ , adding any other edge to  $G$  would create a Hamiltonian cycle. So, let  $uv \notin E(G)$ . There is a Ham. path  $v_1, v_2, \dots, v_n$  with ends  $u = v_1$  and  $v = v_n$ .
- **Fact:** If  $v_i \in N(v)$  and  $v_{i+1} \in N(u)$  for some  $1 < i < n-1$ , done.
- We claim that there is such an  $i$ , let  $S = \{i : v_{i+1} \in N(u)\}$  and  $T = \{i : v_i \in N(v)\}$ .

$$|S \cup T| + |S \cap T| = |S| + |T| = \deg(u) + \deg(v) \geq n.$$



# Sufficient Conditions for being Hamiltonian

**Example:** Two cliques of order  $\lceil (n+1)/2 \rceil$  and  $\lfloor (n+1)/2 \rfloor$  merged at one vertex. This graph has a very high minimum degree, but it is not hamiltonian.

## Theorem (Dirac, 1952)

*If  $G$  is a simple graph with at least three vertices and  $\delta(G) \geq n(G)/2$ , then  $G$  is Hamiltonian.*

- Assume on the contrary that  $G$  is a maximal non-Hamiltonian graph that satisfies the minimum degree condition.
- By the maximality of  $G$ , adding any other edge to  $G$  would create a Hamiltonian cycle. So, let  $uv \notin E(G)$ . There is a Ham. path  $v_1, v_2, \dots, v_n$  with ends  $u = v_1$  and  $v = v_n$ .
- **Fact:** If  $v_i \in N(v)$  and  $v_{i+1} \in N(u)$  for some  $1 < i < n-1$ , done.
- We claim that there is such an  $i$ , let  $S = \{i : v_{i+1} \in N(u)\}$  and  $T = \{i : v_i \in N(v)\}$ .

$$|S \cup T| + |S \cap T| = |S| + |T| = \deg(u) + \deg(v) \geq n.$$

Since  $n \notin S \cup T$ ,  $|S \cup T| \leq n-1$ , done.

# Sufficient Conditions for being Hamiltonian

## Theorem (Ore, 1960)

*Let  $G$  be a simple graph. If  $u$  and  $v$  are distinct non-adjacent vertices such that  $\deg(u) + \deg(v) \geq n(G)$ , then  $G$  is Hamiltonian iff  $G + uv$  is Hamiltonian.*

# Sufficient Conditions for being Hamiltonian

## Theorem (Ore, 1960)

*Let  $G$  be a simple graph. If  $u$  and  $v$  are distinct non-adjacent vertices such that  $\deg(u) + \deg(v) \geq n(G)$ , then  $G$  is Hamiltonian iff  $G + uv$  is Hamiltonian.*

The **closure** of a graph  $G$ , denoted by  $C(G)$ , is the graph with the same vertex set as  $G$  that is obtained by iteratively adding the edges to  $G$  whose endvertices are a non-adjacent pair with degree sum at least  $n$ .

# Sufficient Conditions for being Hamiltonian

## Theorem (Ore, 1960)

*Let  $G$  be a simple graph. If  $u$  and  $v$  are distinct non-adjacent vertices such that  $\deg(u) + \deg(v) \geq n(G)$ , then  $G$  is Hamiltonian iff  $G + uv$  is Hamiltonian.*

The **closure** of a graph  $G$ , denoted by  $C(G)$ , is the graph with the same vertex set as  $G$  that is obtained by iteratively adding the edges to  $G$  whose endvertices are a non-adjacent pair with degree sum at least  $n$ .

## Theorem (Bondy-Chvátal, 1976)

*A simple graph on  $n$  vertices is Hamiltonian iff its closure is Hamiltonian.*

# Sufficient Conditions for being Hamiltonian

## Theorem (Ore, 1960)

*Let  $G$  be a simple graph. If  $u$  and  $v$  are distinct non-adjacent vertices such that  $\deg(u) + \deg(v) \geq n(G)$ , then  $G$  is Hamiltonian iff  $G + uv$  is Hamiltonian.*

The **closure** of a graph  $G$ , denoted by  $C(G)$ , is the graph with the same vertex set as  $G$  that is obtained by iteratively adding the edges to  $G$  whose endvertices are a non-adjacent pair with degree sum at least  $n$ .

## Theorem (Bondy-Chvátal, 1976)

*A simple graph on  $n$  vertices is Hamiltonian iff its closure is Hamiltonian.*

## Theorem (Chvátal's condition, 1972)

*Let  $G$  be a simple graph with vertex degrees  $d_1 \leq \dots \leq d_n$ , where  $n \geq 3$ . If for each  $i < n/2$ ,  $d_i > i$  or  $d_{n-i} \geq n - i$ , then  $G$  is Hamiltonian.*

# Chvátal's Condition

## Theorem (Chvátal's condition, 1972)

*Let  $G$  be a simple graph with vertex degrees  $d_1 \leq \dots \leq d_n$ , where  $n \geq 3$ . If for each  $i < n/2$ ,  $d_i > i$  or  $d_{n-i} \geq n - i$ , then  $G$  is Hamiltonian.*

- By using Bondy-Chvátal condition (**BCC**), we will show that  $C(G)$  is Hamiltonian under these assumptions and thus  $G$  is Ham.

# Chvátal's Condition

## Theorem (Chvátal's condition, 1972)

*Let  $G$  be a simple graph with vertex degrees  $d_1 \leq \dots \leq d_n$ , where  $n \geq 3$ . If for each  $i < n/2$ ,  $d_i > i$  or  $d_{n-i} \geq n - i$ , then  $G$  is Hamiltonian.*

- By using Bondy-Chvátal condition (**BCC**), we will show that  $C(G)$  is Hamiltonian under these assumptions and thus  $G$  is Ham.
- **Claim:**  $C(G) = K_n$ .  
To prove this, again assume on the contrary that  $C(G) \neq K_n$ . We will show that there is an  $i$  for which BCC does not hold, i.e.  
for some  $i$ , at least  $i$  vertices have degree at most  $i$  and at least  $n - i$  vertices have degree less than  $n - i$ .

# Chvátal's Condition

## Theorem (Chvátal's condition, 1972)

Let  $G$  be a simple graph with vertex degrees  $d_1 \leq \dots \leq d_n$ , where  $n \geq 3$ . If for each  $i < n/2$ ,  $d_i > i$  or  $d_{n-i} \geq n - i$ , then  $G$  is Hamiltonian.

- By using Bondy-Chvátal condition (**BCC**), we will show that  $C(G)$  is Hamiltonian under these assumptions and thus  $G$  is Ham.
- **Claim:**  $C(G) = K_n$ .  
To prove this, again assume on the contrary that  $C(G) \neq K_n$ . We will show that there is an  $i$  for which BCC does not hold, i.e.  
for some  $i$ , at least  $i$  vertices have degree at most  $i$  and at least  $n - i$  vertices have degree less than  $n - i$ .
- Details left for reading.



# Chvátal's Condition

## Theorem (Chvátal's condition, 1972)

Let  $G$  be a simple graph with vertex degrees  $d_1 \leq \dots \leq d_n$ , where  $n \geq 3$ . If for each  $i < n/2$ ,  $d_i > i$  or  $d_{n-i} \geq n - i$ , then  $G$  is Hamiltonian.

- By using Bondy-Chvátal condition (**BCC**), we will show that  $C(G)$  is Hamiltonian under these assumptions and thus  $G$  is Ham.
- **Claim:**  $C(G) = K_n$ .  
To prove this, again assume on the contrary that  $C(G) \neq K_n$ . We will show that there is an  $i$  for which BCC does not hold, i.e.  
for some  $i$ , at least  $i$  vertices have degree at most  $i$  and at least  $n - i$  vertices have degree less than  $n - i$ .
- Details left for reading.

**Example:** The graph  $K_i \vee (\bar{K}_i + K_{n-2i})$  is an example where Chvátal's condition is not satisfied, but still the degrees are high.