

	Node	Logic functions
EGFR pathway	Egfr	input
	Rl	Egfr
	Pnt	Rl
	pGro	Rl pGro
	Gro	!pGro
Wingless pathway	Wg	input
	Fz	Wg
	Arm	Fz
	Pan	Arm
Hedgehog pathway	Hh	input
	Ptc	!Hh
	Smo	!Ptc
	CiA	Smo & !en
	CiR	!Smo & !en
Markers	Dr	!vnd & !ind
	ind	!vnd & !Gro
	vnd	Pnt & !ind
	en	Pan
	hh	!CiR & en
	wg	CiA & !CiR
	gsb	!Ptc
	svp	ac ind vnd
	ac	Pan & !ind
	hkb	!Gro & !gsb
Neuroblast phenotypes	7-4	en & Dr & ac & svp
	7-1	en & vnd & ac & gsb & svp
	6-2	en & ind & gsb & svp
	5-6	wg & Dr & gsb
	5-3	wg & ind & gsb & svp
	5-2	wg & vnd & gsb & svp

Documentation of the logic functions

The EGFR Pathway

Egfr = input

Spi encodes ligand that activates Egfr. Aos functions as an inhibitor of the signaling triggered by Egfr by forming a clamp-like structure and sequestering Spi^{9,10}. In the neuroectoderm model, we consider Egfr as an input to eliminate the redundancy originating from Spi inhibition by Aos. This way we can eliminate the redundant attractor where Egfr activation is not possible even though Spi is active.

RI = Egfr

RI encodes the mitogen activated protein (MAP) kinase, a core component of the RAS/MAPK pathway. Egfr activation induces RAS/MAPK pathway activation. Cells with a loss of function in RI produce the same cell-death phenotype as seen in an EGF loss of function¹¹.

Pnt = RI

Activated RI phosphorylates and activates transcription factor Pnt^{11,12}.

pGro = RI | pGro

Gro is phosphorylated by MAPK. Modification of Gro downregulates its repressor activity, causing derepression of pathway target genes. MAPK is no longer active after RTK signaling has been turned off, yet Gro remains stably phosphorylated and its activity attenuated, allowing for sustained RTK target gene expression. Phosphorylated Groucho is a nuclear and stable protein. This was captured in our model via self-loop of pGro and Gro inhibition¹³.

Gro = ! pGro

Please refer to the function of pGro for the explanation of the Boolean update rule of Gro.

Wingless Pathway

Wg = input

Since the initial activations of *hh* and *wg* are independent of each other, Wg is provided as an input node⁶

Fz=Wg

The Wg signal is transduced across the membrane involving Frizzled proteins (such as Fz and DFz2) in the adjacent receiving cells¹.

Arm = Fz

Wg binding to Fz and Arrow brings them together, thereby recruiting Dsh to the membrane. When the destruction complex is inactivated by receptor/Dsh activity, Arm/beta-catenin translocates to the nucleus where it binds the N-terminus of Pan, displacing the Groucho co-repressor and recruiting activators to drive target gene expression. The recruitment of Dsh to the membrane is not included in the model. We considered the main event that Wg binding to Fz activates Arm translocation to the nucleus where it binds to the N-terminus of Pan, thereby recruiting activators to drive target gene expression^{7,8}.

Pan = Arm

Please refer to the function of Arm.

Hedgehog Pathway**Hh = input**

Hh is an input node as it is a morphogen and can activate Hh signal transduction pathway. 2.

Ptc=!Hh

Secreted Hh interacts with its receptor Ptc, thus relieving the repression of Ptc on Smo^{1,2}.

Smo=!Ptc

Please refer to the function of Ptc. In the absence of Hh ligand, repression of Smo allows Drosophila Protein Kinase A (PKA) phosphorylation of Ci on several sites, and these phosphorylation events are required for the cleavage of Ci into Ci^R³.

Ci^A=Smo&!en

Ci is a cytoplasmic protein with no known function in this form. It can be cleaved to generate Ci^R, a 75 kD transcriptional repressor (henceforth Ci^R) or full-length Ci, a 155 kD transcriptional activator (Ci^A). Smo is freed of the inhibitory effects of Ptc, Smo signals through unknown mechanisms to the Fu/Cos2/Ci complex, causing hyperphosphorylation of Fu and Cos2. This results in the complex loosening its hold on microtubules and leads to the stabilization of Ci^A. Stabilized Ci^A can then travel to the nucleus and functions as a transcriptional activator, upregulating transcription of Hh target genes^{4,5}.

Ci^R=!Smo&!en

Pka phosphorylates Ci on several sites and these phosphorylation events are required for the cleavage of Ci into Ci^R. Ci is repressed by *en*. Since the exact mechanism leading to the generation of the different forms of Ci is not fully known, these reactions were omitted from the model. Ci repression by *en* was introduced into the model via two inhibitory edges to both forms of Ci^{4,5}.

Markers**Dr = !vnd & !ind**

Please refer to reaction 21 for the explanation of the Boolean update rule of *Dr*.

ind = !vnd & !Gro

Vnd represses *ind* and *Dr* in the ventral neuroectoderm, and *ind* represses *Dr* in the intermediate neuroectoderm. Gro is a nuclear repressor and represses *ind* transcription¹⁵⁻¹⁷.

vnd = Pnt & !ind

EGF signaling and Pnt either directly or indirectly maintain the expression of several genes in the neurogenic ectoderm including *ind* and *vnd*, which encode regulatory proteins that pattern the future ventral nerve cord. EGF signaling maintains the expression of Pnt transcription factor, which, in turn, sustains the expression of *vnd* (previously activated by Dorsal and Twi), Rho, and Vn¹³⁻¹⁵

en = Pan

Please refer to reaction 9 for the explanation of the Boolean update rule of *en*.

wg= CiA & !CiR

Ci^R represses *wg*, *ptc*, and *hh* transcription whereas Ci^A induces transcription of *ptc* and *wg*^{4,5}.

hh = en & ! Ci^R

One of the functions of *en* is to maintain *hh* expression. Ci^R, a 75 kD transcriptional repressor moves to the nucleus and represses *hh*¹.

gsb =! Ptc

Ptc represses the expression of *gsb*¹⁴.

svp = ac | ind | vnd

Vnd controls activation of two other genes, *hkb* and *svp*. The row 5 NB5-2 expresses *svp-lacZ* and divides asymmetrically multiple times. In *gsb*- embryos, both row 3 and row 5 medial neuroblasts express the row 3 markers *ftz* and *odd*, do not express the row 5 marker *svp-lacZ*, and divide nearly symmetrically²⁷.

ac = Pan & lind

Ac is repressed in the neuroectoderm and neuroblasts by *ind*. *Arm* and *pan* regulate the transcriptional activation of numerous target genes such as *ac*¹⁸. Since *arm* is a transcriptional activator of *pan*, *arm* indirectly involved in the activation of *ac*. Therefore, in the neuroectoderm model, *pan* activates *ac*. In addition to activation of AS-C¹⁹, *vnd* also controls activation of two other genes, *hkb* and *svp*, in the proneural clusters²⁰.

hkb = !Gro & !gsb

Both Wg and Hh activate *hkb* expression in the neuroectoderm and NBs, and multiple repressors act to restrict its expression. *En* is expressed in neuroectodermal rows 6/7 and partially represses *hkb* expression in those rows²¹. *Vnd* controls activation of *hkb*²⁰.

Neuroblast fates**7-4 = en & Dr & ac & svp**

In *Drosophila* NB7-4 expresses *en* and *ac* as it delaminates²²⁻²⁴.

7-1 = en & vnd & gsb & ac & svp

NB7-1 expresses *gsb*, *en*, *vnd*, and *ac* as it forms during S1^{19,25-27}.

6-2 =en & ind & gsb & svp

At S2, NB 6-2 expresses *en* and *ind*, *svp*. *Gsb* expression is detected by S3^{24,25,29}

5-6 = wg & Dr & gsb

At S1, NB5-6 expresses *Dr*, *wg* and *gsb-d*^{23,25,27,28}

5-3 = wg & ind & gsb & svp

At S1 NB 5-3 expresses *wg*^{25,28}, *ind*²⁹, and *gsb-d*^{23,25,27}. At S2 it adds *svp* expression²⁵.

5-2 = wg & vnd & gsb & svp

NB 5-2 expresses *svp-lacZ*, *gsb-d*, *vnd*^{26,30} and *wg* as it forms^{25,27,31}.

Neuroectodermal (NE) fates

Neuroectoderm patterns by the expression of three genes- *vnd*, *ind*, *Dr* in the neuroectodermal cells. (We expect at least three attractors with the activity of *Dr*, *ind* and *vnd* separately.) Other transcriptional factors expressed by these cells but not spatially restricted include *hkb* and *svp*²⁰. *Egfr* and *pGro* input combinations reflect the neuroectodermal cell states as it leads to the activation of transcriptional factors: *vnd*, *ind*, *Dr*. The EGFR pathway is sufficient to generate multiple neuroectodermal cell fates by both internal (via phosphorylation of *Gro*) or external signals (via receptor activation).

Attractors associated with the NE fate include the following transcription factor activities:

Dr

ind, *svp*, *hkb*

vnd, *svp*, *hkb*

Non-physiological (NP) fates

The model revealed four attractors representing non-physiological states, characterized by combinations of marker expressions not observed in neuroblasts and neuroectoderm under normal conditions. Among these non-physiological attractors, one stands out with the predicted active genes *Dr*, *en*, *svp*, *ac*, and *gsb*. Typically, these gene expressions are not observed in neuroblasts, except for the exclusion of *gsb* expression in NB7-4.

Attractors associated with the NP fate include the following transcription factor activities:

Dr, *en*, *gsb*, *svp*, *ac*

ind, *en*, *svp*, *hkb*

vnd, *en*, *svp*, *ac*, *hkb*

In addition, any other combinations of marker gene activities that have not been defined as NE or cell type (e.g.7-1).

Reference

1. Bhat, K. M. Segment polarity genes in neuroblast formation and identity specification during *Drosophila* neurogenesis. *BioEssays* **21**, 472–485 (1999).

2. Im, S. H. *et al.* Tachykinin acts upstream of autocrine Hedgehog signaling during nociceptive sensitization in *Drosophila*. *Elife* **4**, 1–27 (2015).
3. Ingham, P. W. & McMahon, A. P. Hedgehog signaling in animal development: Paradigms and principles. *Genes Dev.* **15**, 3059–3087 (2001).
4. Nybakken, K. & Perrimon, N. Hedgehog signal transduction: Recent findings. *Curr. Opin. Genet. Dev.* **12**, 503–511 (2002).
5. Eaton, S. & Kornberg, T. B. Repression of *ci-D* in posterior compartments of *Drosophila* by engrailed. *Genes Dev.* **4**, 1068–1077 (1990).
6. DiNardo, S., Heemskerk, J., Dougan, S. & O’Farrell, P. H. The making of a maggot: patterning the *Drosophila* embryonic epidermis. *Curr. Opin. Genet. Dev.* **4**, 529–534 (1994).
7. Bejsovec, A. Wingless/Wnt signaling in *Drosophila*: The pattern and the pathway. *Mol. Reprod. Dev.* **80**, 882–894 (2013).
8. Swarup, S. & Verheyen, E. M. Wnt/wingless signaling in *drosophila*. *Cold Spring Harb. Perspect. Biol.* (2012) doi:10.1101/cshperspect.a007930.
9. Golembo, M., Schweitzer, R., Freeman, M. & Shilo, B. Z. Argos transcription is induced by the *Drosophila* EGF receptor pathway to form an inhibitory feedback loop. *Development* **122**, 223–230 (1996).
10. Hong, J. W., Hendrix, D. A., Papatsenko, D. & Levine, M. S. How the Dorsal gradient works: Insights from postgenome technologies. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 20072–20076 (2008).
11. Brunner, D. *et al.* A gain-of-function mutation in *Drosophila* MAP kinase activates multiple receptor tyrosine kinase signaling pathways. *Cell* **76**, 875–888 (1994).
12. Oellers, N. & Hafen, E. Biochemical characterization of rolled(Sem) an activated form of *Drosophila* mitogen-activated protein kinase. *J. Biol. Chem.* **271**, 24939–24944 (1996).
13. Cinnamon, E. *et al.* Multiple RTK pathways downregulate Groucho-mediated repression in *Drosophila* embryogenesis. *Development* **135**, 829–837 (2008).
14. Li, X. & Noll, M. Role of the gooseberry gene in *Drosophila* embryos: Maintenance of wingless expression by a wingless-geeseberry autoregulatory loop. *EMBO J.* **12**, 4499–4509 (1993).
15. Cowden, J. & Levine, M. Ventral dominance governs sequential patterns of gene expression across the dorsal-ventral axis of the neuroectoderm in the *Drosophila* embryo. *Dev. Biol.* **262**, 335–349 (2003).
16. Golembo, M., Yarnitzky, T., Volk, T. & Shilo, B. Z. Vein expression is induced by the EGF receptor pathway to provide a positive feedback loop in patterning the *Drosophila* embryonic ventral ectoderm. *Genes Dev.* **13**, 158–162 (1999).
17. Levine, M. & Davidson, E. H. Gene regulatory networks for development. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 4936–4942 (2005).
18. Zeng, Y. A., Rahnama, M., Wang, S., Lee, W. & Verheyen, E. M. Inhibition of *drosophila* Wg signaling involves competition between Mad and Armadillo/ β -catenin for dTcf binding. *PLoS One* **3**, (2008).

19. Skeath, J. B., Panganiban, G. F. & Carroll, S. B. The ventral nervous system defective gene controls proneural gene expression at two distinct steps during neuroblast formation in *Drosophila*. *Development* **120**, 1517–1524 (1994).
20. Chu, H., Parras, C., White, K. & Jiménez, F. Formation and specification of ventral neuroblasts is controlled by *vnd* in *Drosophila* neurogenesis. *Genes Dev.* **12**, 3613–3624 (1998).
21. McDonald, J. A. & Doe, C. Q. Establishing neuroblast-specific gene expression in the *Drosophila* CNS: *huckebein* is activated by *Wingless* and *Hedgehog* and repressed by *Engrailed* and *Gooseberry*. *Development* **124**, 1079–1087 (1997).
22. Lundell, M. J., Chu-Lagraff, Q., Doe, C. Q. & Hirsh, J. The *engrailed* and *huckebein* genes are essential for development of serotonin neurons in the *Drosophila* CNS. *Mol. Cell. Neurosci.* **7**, 46–61 (1996).
23. Doe, C. Q. Molecular markers for identified neuroblasts and ganglion mother cells in the. *Development* **116**, 855–863 (1992).
24. Skeath, J. B. & Carroll, S. B. Regulation of proneural gene expression and cell fate during neuroblast segregation in the *Drosophila* embryo. *Development* **114**, 939–946 (1992).
25. Broadus, J. *et al.* New neuroblast markers and the origin of the aCC/pCC neurons in the *Drosophila* central nervous system. *Mech. Dev.* **53**, 393–402 (1995).
26. McDonald, J. A. *et al.* Dorsoventral patterning in the *Drosophila* central nervous system: The *vnd* homeobox gene specifies ventral column identity. *Genes Dev.* (1998) doi:10.1101/gad.12.22.3603.
27. Skeath, J. B., Zhang, Y., Holmgren, R., Carroll, S. B. & Doe, C. Q. Specification of neuroblast identity in the *drosophila* embryonic central nervous system by *gooseberry-distal*. *Nature* **376**, 427–430 (1995).
28. Chu-LaGraff, Q. & Doe, C. Q. Neuroblast specification and formation regulated by *wingless* in the *Drosophila* CNS. *Science* (80-.). (1993) doi:10.1126/science.8372355.
29. Weiss, J. B. *et al.* Dorsoventral patterning in the *Drosophila* central nervous system: The intermediate neuroblasts defective homeobox gene specifies intermediate column identity. *Genes Dev.* **12**, 3591–3602 (1998).
30. White, K., DeCelles, N. L. & Enlow, T. C. Genetic and developmental analysis of the locus *vnd* in *Drosophila melanogaster*. *Genetics* **104**, 433–448 (1983).
31. Chu-LaGraff, Q. & Doe, C. Q. Neuroblast specification and formation regulated by *wingless* in the *Drosophila* CNS. *Science* (80-.). **261**, 1594–1597 (1993).