Regev Reduction in a Nutshell

Goal: Given an LWE solver for $\mathcal{L}$, solve SIS, *i.e.* find a short vector in $\mathcal{L}^*$.

Regev Reduction in a Nutshell

Goal: Given an LWE solver for $\mathcal{L}$, solve SIS, *i.e.* find a short vector in $\mathcal{L}^*$.

1. Using the LWE solver, prepare

$$\sum_{\mathbf{x}\in\mathcal{L}} \sum_{\mathbf{e}\in\mathbb{Z}_q^m} f(\mathbf{e}) \left|\mathbf{x}+\mathbf{e}\right\rangle .$$

Regev Reduction in a Nutshell

Goal: Given an LWE solver for $\mathcal{L}$, solve SIS, *i.e.* find a short vector in $\mathcal{L}^*$.

1. Using the LWE solver, prepare

$$\sum_{\mathbf{x}\in\mathcal{L}} \sum_{\mathbf{e}\in\mathbb{Z}_q^m} f(\mathbf{e}) \left|\mathbf{x}+\mathbf{e}\right\rangle .$$

2. Apply a QFT:

$$\sum_{\mathbf{y}\in\mathcal{L}^*} \widehat{f}(\mathbf{y}) \left|\mathbf{y}\right\rangle .$$

where $\widehat{f} = \mathsf{FT}(f)$.

Regev Reduction in a Nutshell

Goal: Given an LWE solver for $\mathcal{L}$, solve SIS, *i.e.* find a short vector in $\mathcal{L}^*$.

1. Using the LWE solver, prepare

$$\sum_{\mathbf{x}\in\mathcal{L}} \sum_{\mathbf{e}\in\mathbb{Z}_q^m} f(\mathbf{e}) \left|\mathbf{x}+\mathbf{e}\right\rangle .$$

2. Apply a QFT:

$$\sum_{\mathbf{y}\in\mathcal{L}^*} \widehat{f}(\mathbf{y}) \left|\mathbf{y}\right\rangle .$$

where $\widehat{f} = \mathsf{FT}(f)$.

Regev Reduction in a Nutshell

Goal: Given an LWE solver for $\mathcal{L}$, solve SIS, *i.e.* find a short vector in $\mathcal{L}^*$.

1. Using the LWE solver, prepare

$$\sum_{\mathbf{x}\in\mathcal{L}} \sum_{\mathbf{e}\in\mathbb{Z}_q^m} f(\mathbf{e}) \left|\mathbf{x}+\mathbf{e}\right\rangle .$$

2. Apply a QFT:

$$\sum_{\mathbf{y}\in\mathcal{L}^*} \widehat{f}(\mathbf{y}) \left|\mathbf{y}\right\rangle .$$

   where $\widehat{f} = \mathsf{FT}(f)$.

3. Measure and obtain $\mathbf{y}$ w.p. $|\widehat{f}(\mathbf{y})|^2$.