

Wildfire risk modeling

Sandra Oliveira¹, Jorge Rocha¹ and Ana Sá²

Abstract

Wildfires occur worldwide and affect different territories in diverse ways. New challenges in wildfire management are emerging, fueled by climatic, land cover, and sociodemographic changes, fostering wildfire science to respond with advanced and inclusive approaches. Supported by technological improvements, wildfire risk modeling is earning relevance in several wildfire phases, covering prevention, suppression, and recovery, and encouraging the development of different products and tools. Most wildfire risk approaches reflect a broad conceptual framework that includes different but interconnected components, which can also be analyzed separately, such as likelihood, hazard, exposure, vulnerability, and coping capacity. This article presents a concise and nonexhaustive review of the methods and approaches applied to model wildfire risk and its key components, in particular since 2018, and how they can contribute to wildfire management.

Addresses

¹Centre for Geographical Studies, Institute of Geography and Spatial Planning, Universidade de Lisboa, 1600-276, Lisbon, Portugal

²Forest Research Centre, School of Agriculture, Universidade de Lisboa, 1349-017, Lisbon, Portugal

Corresponding author: Oliveira, Sandra (sandra.oliveira1@campus.ul.pt)

Current Opinion in Environmental Science & Health 2021, **23**:100274

This review comes from a themed issue on **Environmental impact assessment: Fire. Prevention, management and challenges**

Edited by **Paulo Pereira** and **Edivaldo L Thomaz**

For a complete overview see the [Article Collection](#)

<https://doi.org/10.1016/j.coesh.2021.100274>

2468-5844/© 2021 Elsevier B.V. All rights reserved.

Keywords

Wildfire hazard, Exposure, Vulnerability, Coping capacity, Simulation approaches.

Introduction

Wildfires occur globally and affect different territories in distinct ways. In the last decades, the incidence of large and uncontrolled fires has risen, and extreme events are expected to become more frequent and destructive [1]. Exceptionally large and damaging fires have recently occurred, for example, in central Portugal in 2017, in the western USA in 2018, and in Brazil and eastern Australia

in 2019 [2–5]. Changes in land cover, climate, and sociodemographic conditions are pointed out as the main reasons behind these unparalleled events [6], but further understanding is needed on how these changes drive wildfire risk in different territorial contexts and how this knowledge can be integrated into suitable mitigation strategies. New challenges in wildfire management are emerging, prompting the scientific community to respond with integrative approaches and innovative modeling applications, fostered by technological improvements in computation capabilities and an increase in data availability [7].


Wildfire risk modeling is a key approach to uncover the complex relationships between fire occurrence, its driving factors and potential impacts and is relevant for different wildfire phases: prevention, suppression, and recovery. The methodologies used for risk assessment and modeling have evolved over time, and nowadays, most wildfire risk modeling approaches reflect a broad conceptual framework, which includes several distinct but interconnected components, that can be analyzed separately [8] and incorporated into decision-making processes at different levels, such as landscape management [9] or household protection and evacuation [10] (see [Figure 1](#)).

This paper presents a concise review of the methods and approaches applied to model wildfire risk in the last few years, focusing on scientific and technical developments since 2018. We describe the tools used to assess the key risk components, how they are combined and mapped, and how they contribute to develop valuable products for wildfire management.

Likelihood and hazard

Wildfire likelihood and hazard reflect the characteristics of the fire process, responding to specific questions: (1) how it starts, (2) when it happens, (3) where it occurs, and (4) how it will grow. Wildfire occurrence modeling and fire spread simulations have been implemented in different territorial contexts. The former are covered mainly in probability analysis and susceptibility assessments (1, 2, and 3), whereas the latter is used in building scenarios of fire ignition, intensity, and propagation (4). Regression models, easy to use and interpret, have been extensively applied [11,12]. The approaches used to model specifically human-caused fires have been previously reviewed by Costafreda-Aumedes et al. [7]

Figure 1



Wildfire risk modeling framework, with separate components and some of the approaches and methods used for their assessment.

who pointed out that the selection of methods depends as well on the available data and on the temporal and spatial scales of the analysis. When models cover large geographical areas and to capture nonstationary patterns of wildfire drivers, geographically weighted regression has been used [13,14], and generalized additive regression has also been tested [15].

In the last decade, wildfire modeling has evolved to use advanced machine learning approaches based on algorithms that can improve automatically through experience [16]. Their use presents some advantages because they do not require *a priori* knowledge of the phenomenon, and they can be used in nonparametric data, being able to identify nonlinear patterns from large data sets. On the other hand, they use complex and sophisticated algorithms, and the interpretation of the importance of the independent variables is not straightforward as in traditional regression methods [17,18]. The use of such methods, among which Artificial Neural Networks, Random Forests, or Support Vector Machines, is becoming increasingly frequent [19–22], occasionally focusing on comparative evaluations of different methods [16,23,24]. The application of different machine learning methods in wildfire science has been recently reviewed by Jain et al. [16]. Another approach gaining relevance is Maximum Entropy (MaxEnt), an algorithm that joins the predictive capability of machine learning methods with the

explicative ability of traditional regression models, originally used in species distribution and ecological niche modeling and lately applied to wildfires in several regions [25–27].

The response to how wildfires will grow (4) has been increasingly relying on simulation approaches to quantitatively estimate wildfire likelihood and impacts. Different fire simulation systems are available, such as FARSITE, FSim, FlamMap, or Burn-P3 [28,29], used to estimate fire ignition and spread in different scenarios of fuel and weather conditions, considering potential landscape interventions, such as fuel treatments and forest management levels [30–32], and eventually combined with remote sensing data [33,34]. These results can support operational decisions, regarding fuel treatment locations, surveillance and suppression activities, and mitigation measures [35,36]. Coupling simulation models with Geographic Information Systems (GIS) has been a key issue, and it can be done at three levels: loose, moderate, or tight/close coupling. The loose level uses GIS only as a preprocessing tool, information provider, and a visualizer of results, whereas moderate coupling implies remote procedure calls among GIS and simulation models, such as agent-based modeling approaches [37]. The close coupling simultaneously joins operations of both systems, and it has been developed mainly with the use of cellular automata models within a GIS environment [38].

Exposure

Wildfire exposure evaluates which assets, and to what extent, are located in fire-prone areas [28]. It results from the analysis of fire occurrence, likelihood, or hazard in relation to highly valued resources and assets (HVRAs), using historical fire data [39] or by coupling stochastic and probabilistic wildfire simulations with the spatial distribution of people and assets, such as forests, built-up areas, agricultural lands, or protected areas [40–42]. The combination of simulation outputs with GIS has led to models of wildfire transmission to human communities based on the concept of fireshed, the areas with potential to expose communities to fires, according to ignition probabilities and propagation patterns. Using this concept, wildfire exposure was estimated in Spain by combining simulation outputs with spatial layers representing residential areas, built-up structures, and land cover types [43]; land tenure types were mapped, and their relationship with wildfire transmission and exposure levels in Greece was assessed [44]; and wildfire exposure and fireshed extent were obtained in the United States, integrating as well other geospatial attributes, such as travel cost and distance between urban centers and wildland–urban interface areas [45].

Vulnerability

Vulnerability represents the potential for loss as a result of wildfires [46]. As a measure of potential wildfire impacts, it depends on the conditions of the wildfire and on the characteristics of the affected assets. When measuring the vulnerability of forests and natural areas, models focus on ecological impacts, simulating the effects of burnt area, heat release, and other fire behavior parameters in the biophysical attributes, mortality rate, or survival response of species in particular ecosystems, such as Alpine forests in France [47] and mountainous grasslands in South Africa [48], or by assessing successional trajectories in Mediterranean pine forests in Spain, via sensitivity analysis and a persistence index [49].

In parallel, many studies in the past decade emphasized the social component to assess the vulnerability of human communities to wildfires, following the research first developed by Cutter *et al.* [50] with the Social Vulnerability Index. Social attributes of people exposed to wildfires are weighted, aggregated, and mapped based on geospatial or cluster analysis [32,51]. Other recent works have coupled social data with wildfire simulations [46,52], showing which human communities have more or less capacity to absorb the effects and to recover from the impacts of wildfires.

Coping capacity and response

Lately, particular attention has been given to the implementation of self-protection and mitigation strategies by people exposed to wildfires [53]. Models of

alternative actions and behavior choices that can modify the response capacity to wildfire events have emerged, assessing evacuation, local sheltering, or stay-at-home options [54–56]. By applying the Protective Action Decision Model (PADM), by Lindell & Perry [57], a decision-making model that integrates environmental and social cues within a framework of information flows, several studies have evaluated people's behavioral choices in Australia and in the United States, for example, as reviewed by McLeenan *et al.* [58]. Other modeling procedures have been tested, with new methods or coupled with PADM, generally divided into two categories: conceptual models, which develop frameworks to explain people's behavioral steps when deciding what to do in a wildfire emergency; and engineering models, which assess the factors affecting human behavior (choice models) or simulate evacuation travel conditions (traffic models) [59]. For example, a mathematical approach to create a Wildfire Decision Model has been recently proposed, incorporating human behavior assumptions with simulations of people's actions using a binary logit model [10]. In other cases, geospatial tools are used to integrate wildfire and traffic simulations to assess evacuation triggers [60]. Other studies have been focusing on assessing how the characteristics, experience, and social context of exposed communities can influence mitigation and adaptation actions, using interactional approaches to document and compare different combinations of local characteristics. These can allow identifying tailored strategies for wildfire recovery and adaptation, according to the capabilities and needs of distinct human communities [61,62].

Wildfire risk

Wildfire risk assessment intends to provide an integrated view of fire likelihood and consequences [63]. The relative contribution of each component, previously described, is evaluated, then aggregated and summarized in quantitative indices or categories, providing a rating of wildfire occurrence conditions and potential impacts, and further combined with mapping and zoning tools to identify priority areas for intervention. In Portugal, a composite risk index in five categories, which combined burn probability, exposure, and social vulnerability, was applied to the villages of a civil parish [32]. In Australia, a Bayesian decision network model was applied to specifically assess the efficiency of prescribed burning strategies in wildfire risk levels regarding valued assets [9]. In Brazil, a fire risk map was obtained by combining historical fire data with a Kernel density estimator to determine critical areas [64]. In the United States, a mixed integer program model was used to assess fire risk variations in view of potential suppression activities at the operational level [65]. In Canada, wildfire risk has been calculated based on a specific equation that sums probability linked to a range

of fire behavior conditions, multiplied by the difference between potential benefits and losses [28]. A European initiative has also proposed a composite risk index that aggregates three dimensions: hazard and exposure, vulnerability, and lack of coping capacity, applicable at different spatial scales and for multiple hazards [66]. Because of the complexity of the phenomenon and the numerous approaches available, the comparison and interpretation of wildfire risk models is still challenging. Also, risk ratings may cover only part of the components, varying in data and methods, weighting, type of outputs obtained, or scale of application [8]. Despite this diversity, common ground among risk models is found on their main purpose of obtaining a grading system for the wildfire threat and on their spatially explicit dimension, to which geospatial tools are particularly suitable [67].

Final remarks

Wildfire science is continuously evolving to respond to environmental changes and societal needs. Currently, we are facing a paradigm shift regarding the development of sustainable planning and operational frameworks, with the aim of reducing fire-induced damages and losses, instead of focusing on burned area reduction [6]. Risk models are developed with the ultimate goal to support wildfire management and decision-making at multiple levels. The need to integrate the scientific knowledge and technical tools with the regulations and practices of wildfire agencies and operational systems is a key issue [63,65]. The involvement of citizens and stakeholders in the decision-making process is also highlighted [68], as a means to improve their preparedness and adaptation to wildfires and to ensure the applicability and acceptance of risk models. Simplification, aggregation, and the standardization of models outputs is further required [9,67] to enhance communication, knowledge transfer and co-creation. Other challenges include the integration of data that can efficiently represent extreme fire conditions and climate change effects in wildfire activity [69]. Ongoing technological developments will keep offering access to better data, from satellite imagery and drones [5], as well as advanced modeling tools based, for example, on machine learning [16], implying the frequent upgrade of modeling procedures. Comparative evaluations of advanced methods with well-established models will continue to be relevant, and the combination of different methods and their joint benefits could be particularly useful [8]. Research works linked to governmental or private initiatives associated with wildfire management, as those emerged in Portugal after the large wildfires of 2017 [32,70], can help strengthening the crucial link between science, decision-makers, and citizens.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The work developed by S.O. and J.R. was supported by the Centre for Geographical Studies, a research unit funded by Fundação para a Ciência e a Tecnologia (FCT), Portugal (UIDB/00295/2020-UIDP/00295/2020). The work developed by A.S. was supported by the Forest Research Centre, a research unit funded by FCT, Portugal (UIDB/00239/2020), and under the framework of the contract-program nr. 1382 (DL 57/2016/CP1382/CT0003).

References

Papers of particular interest, published within the period of review, have been highlighted as:

* of special interest

** of outstanding interest

1. Turco M, Rosa-Cánovas JJ, Bedia J, Jerez S, Montávez JP, Llasat MC, Provenzale A: **Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models**. *Nat Commun* 2018, **9**:1–9.
2. Nauslar N, Abatzoglou J, Marsh P: **The 2017 North Bay and southern California fires: a case study**. *Fire* 2018, **1**:18.
3. Turco M, Jerez S, Augusto S, Tarín-Carrasco P, Ratola N, Jiménez-Guerrero P, Trigo RM: **Climate drivers of the 2017 devastating fires in Portugal**. *Sci Rep* 2019, **9**:1–8.
4. Boer MM, Dios VR de, Bradstock RA: **Unprecedented burn area of Australian mega forest fires**. *Nat Clim Change* 2020, **10**: 170–172.
5. Kganyago M, Shikwambana L: **Assessment of the characteristics of recent major wildfires in the USA, Australia and Brazil in 2018-2019 using multi-source satellite products**. *Rem Sens* 2020, **12**:1803.
6. Moreira F, Ascoli D, Safford H, Adams MA, Moreno JM, Pereira JMC, Catry FX, Armesto J, Bond W, González ME, et al.: **Wildfire management in Mediterranean-type regions: paradigm change needed**. *Environ Res Lett* 2020, **15**, 011001.
7. Costafreda-Aumedes S, Comas C, Vega-Garcia C: **Human-caused fire occurrence modelling in perspective: a review**. *Int J Wildland Fire* 2017, **26**:983.
8. Xi DDZ, Taylor SW, Woolford DG, Dean CB: **Statistical models of key components of wildfire risk**. *Annu Rev Stat Its Appl* 2019, **6**:197–222.
9. Penman TD, Cirulis B, Marcot BG: **Bayesian decision network modeling for environmental risk management: a wildfire case study**. *J Environ Manag* 2020, **270**:110735.
10. Lovreglio R, Kuligowski E, Gwynne S, Strahan K: **A modelling framework for householder decision-making for wildfire emergencies**. *Int J Disaster Risk Reduct* 2019, **41**:101274.
- * This article presents a brief literature review on existing human behavior and decision-making models in face of wildfire emergency. On the basis of the PADM and other specific evacuation models, it develops a new mathematical predictive framework for households.
11. Mancini LD, Elia M, Barbati A, Salvati L, Corona P, Laforzezza R, Sanesi G: **Are wildfires knocking on the built-up areas door?** *Forests* 2018, **9**:1–13.
12. Jaafari A, Mafi-Gholami D, Thai Pham B, Tien Bui D: **Wildfire probability mapping: bivariate vs. multivariate statistics**. *Rem Sens* 2019, **11**:618.
13. Rodrigues M, Jiménez-Ruano A, Peña-Angulo D, de la Riva J: **A comprehensive spatial-temporal analysis of driving factors of human-caused wildfires in Spain using Geographically Weighted Logistic Regression**. *J Environ Manag* 2018, **225**:177–192.

14. Cao Q, Zhang L, Su Z, Wang G, Guo F: **Exploring spatially varying relationships between forest fire and environmental factors at different quantile levels.** *Int J Wildland Fire* 2020, **29**: 486–498.
 15. Sá ACL, Turkman MAA, Pereira JMC: **Exploring fire incidence in Portugal using generalized additive models for location, scale and shape (GAMLSS).** *Model Earth Syst Environ* 2018, **4**: 199–220.
 16. Jain P, Coogan SCP, Subramanian SG, Crowley M, Taylor S, Flannigan MD: **A review of machine learning applications in wildfire science and management.** *Environ Rev* 2020, **28**: 478–505.
- This paper presents a comprehensive review of machine learning methods used in wildfire science, divided into six focus areas that broadly cover the wildfire risk assessment framework and its main components.
17. Sayad YO, Mousannif H, Al Moatassime H: **Predictive modeling of wildfires: a new dataset and machine learning approach.** *Fire Saf J* 2019, **104**:130–146.
 18. Leuenberger M, Parente J, Tonini M, Pereira MG, Kanevski M: **Wildfire susceptibility mapping: deterministic vs. stochastic approaches.** *Environ Model Software* 2018, **101**:194–203.
 19. Elia M, D'Este M, Ascoli D, Giannico V, Spano G, Ganga A, Colangelo G, Laforzezza R, Sanesi G: **Estimating the probability of wildfire occurrence in Mediterranean landscapes using artificial neural networks.** *Environ Impact Assess Rev* 2020, **85**:106474.
 20. Tonini M, D'Andrea M, Biondi G, Esposti SD, Trucchia A, Fiorucci P: **A machine learning-based approach for wildfire susceptibility mapping. The case study of the Liguria region in Italy.** *Geosciences* 2020, **10**.
 21. Hong H, Tsangaratos P, Ilia I, Liu J, Zhu AX, Xu C: **Applying genetic algorithms to set the optimal combination of forest fire related variables and model forest fire susceptibility based on data mining models. The case of Dayu County, China.** *Sci Total Environ* 2018, **630**:1044–1056.
 22. Tehrany MS, Jones S, Shabani F, Martínez-Álvarez F, Tien Bui D: **A novel ensemble modeling approach for the spatial prediction of tropical forest fire susceptibility using LogitBoost machine learning classifier and multi-source geospatial data.** *Theor Appl Climatol* 2019, **137**:637–653.
 23. Gholamnia K, Gudiyangada Nachappa T, Ghorbanzadeh O, Blaschke T: **Comparisons of diverse machine learning approaches for wildfire susceptibility mapping.** *Symmetry* 2020, **12**:604.
 24. Jaafari A, Zenner EK, Pham BT: **Wildfire spatial pattern analysis in the Zagros Mountains, Iran: a comparative study of decision tree based classifiers.** *Ecol Inf* 2018, **43**:200–211.
 25. Adab H, Atabati A, Oliveira S, Moghaddam Gheshlagh A: **Assessing fire hazard potential and its main drivers in Mazandaran province, Iran: a data-driven approach.** *Environ Monit Assess* 2018, **190**:1–20.
 26. Tracy JL, Trabucco A, Lawing AM, Giermakowski JT, Tchakerian M, Drus GM, Coulson RN: **Random subset feature selection for ecological niche models of wildfire activity in Western North America.** *Ecol Model* 2018, **383**:52–68.
 27. Vacchiano G, Foderi C, Berretti R, Marchi E, Motta R: **Modeling anthropogenic and natural fire ignitions in an inner-Alpine valley.** *Nat Hazards Earth Syst Sci* 2018, **18**:935–948.
 28. Johnston LM, Wang X, Erni S, Taylor SW, McFayden CB, Oliver JA, Stockdale C, Christianson A, Boulanger Y, Gauthier S, et al.: **Wildland fire risk research in Canada.** *Environ Rev* 2020, **28**:164–186.
- This article presents a synthesis of the wildfire risk knowledge based on the Canadian experience, as a contribution for the development of a risk assessment framework at the country level. It presents the most relevant methods applied to assess each risk component, as well as to calculate the combined wildfire risk index.
29. Zigner K, Carvalho LMV, Peterson S, Fujioka F, Duine GJ, Jones C, Roberts D, Moritz M: **Evaluating the ability of FARSITE to simulate wildfires influenced by extreme, down-slope winds in Santa Barbara, California.** *Fire* 2020, **3**:1–23.
 30. Salis M, Del Giudice L, Arca B, Ager AA, Alcasena FJ, Lozano O, Bacciu V, Spano D, Duce P: **Modeling the effects of different fuel treatment mosaics on wildfire spread and behavior in a Mediterranean agro-pastoral area.** *J Environ Manag* 2018, **212**:490–505.
 31. Keane RE, Gray K, Davis B, Holsinger LM, Loehman R: **Evaluating ecological resilience across wildfire suppression levels under climate and fuel treatment scenarios using landscape simulation modelling.** *Int J Wildland Fire* 2019, **28**:533–549.
 32. Oliveira S, Gonçalves A, Benali A, Sá A, Zêzere JL, Pereira JM: **Assessing risk and prioritizing safety interventions in human settlements affected by large wildfires.** *Forests* 2020, **11**:859.
 33. Xofis P, Konstantinidis P, Papadopoulos I, Tsiourlis G: **Integrating remote sensing methods and fire simulation models to estimate fire hazard in a south-east mediterranean protected area.** *Fire* 2020, **3**:1–23.
 34. Sá ACL, Benali A, Fernandes PM, Pinto RMS, Trigo RM, Salis M, Russo A, Jerez S, Soares PMM, Schroeder W, et al.: **Evaluating fire growth simulations using satellite active fire data.** *Remote Sens Environ* 2017, **190**:302–317.
 35. Parisien MA, Dawe DA, Miller C, Stockdale CA, Armitage OB: **Applications of simulation-based burn probability modelling: a review.** *Int J Wildland Fire* 2019, **28**:913–926.
 36. Sakellariou S, Parisien MA, Flannigan M, Wang X, de Groot B, Tampekis S, Samara F, Sfougaris A, Christopoulou O: **Spatial planning of fire-agency stations as a function of wildfire likelihood in Thasos, Greece.** *Sci Total Environ* 2020, **729**: 139004.
 37. Ager AA, Barros AMG, Day MA, Preisler HK, Spies TA, Bolte J: **Analyzing fine-scale spatiotemporal drivers of wildfire in a forest landscape model.** *Ecol Model* 2018, **384**:87–102.
 38. Jiang W, Wang F, Fang L, Zheng X, Qiao X, Li Z, Meng Q: **Modelling of wildland-urban interface fire spread with the heterogeneous cellular automata model.** *Environ Model Software* 2021, **135**:104895.
 39. Chas-Amil ML, García-Martínez E, Touza J: **Iberian Peninsula October 2017 wildfires: burned area and population exposure in Galicia (NW of Spain).** *Int J Disaster Risk Reduct* 2020, **48**: 101623.
 40. Jahdi R, Salis M, Alcasena FJ, Arabi M, Arca B, Duce P: **Evaluating landscape-scale wildfire exposure in northwestern Iran.** *Nat Hazards* 2020, **101**:911–932.
 41. Alcasena FJFJ, Ager AA, Bailey JD, Pineda N, Vega-García C: **Towards a comprehensive wildfire management strategy for Mediterranean areas: framework development and implementation in Catalonia, Spain.** *J Environ Manag* 2019, **231**: 303–320.
- This paper is of particular interest because it presents an integrated approach to wildfire exposure assessment, using advanced simulation modeling and adjusted to inform fire and fuel management at the local level.
42. Salis M, Arca B, Del Giudice L, Palaiologou P, Alcasena-Urdiroz F, Ager A, Fiori M, Pellizzaro G, Scarpa C, Schirru M, et al.: **Application of simulation modeling for wildfire exposure and transmission assessment in Sardinia, Italy.** *Int J Disaster Risk Reduct* 2021, **58**:102189.
 43. Alcasena FJ, Salis M, Ager AA, Castell R, Vega-García C: **Assessing wildland fire risk transmission to communities in northern Spain.** *Forests* 2017, **8**.
 44. Palaiologou P, Ager AA, Nielsen-Pincus M, Evers CR, Kalabokidis K: **Using transboundary wildfire exposure assessments to improve fire management programs: a case study in Greece.** *Int J Wildland Fire* 2018, **27**:501–513.
 45. Ager AA, Palaiologou P, Evers CR, Day MA, Ringo C, Short K: **Wildfire exposure to the wildland urban interface in the western US.** *Appl Geogr* 2019, **111**:102059.
- This paper is of high interest because it demonstrates how human exposure assessment, obtained with wildfire simulations of large fires and geospatial data, applying the fireshed concept, can be used to estimate required investments and identify priority areas for fuel management and safety measures.

46. Palaiologou P, Ager AA, Nielsen-Pincus M, Evers CR, Day MA: **Social vulnerability to large wildfires in the western USA.** *Landsc Urban Plann* 2019, **189**:99–116.
 ** This paper is of outstanding interest because it connects wildfire exposure and vulnerability models in different study areas in the United States and presents a solid literature review on the social attributes to consider in social vulnerability assessment to wildfires.
47. Dupire S, Curt T, Bigot S, Fréjaville T: **Vulnerability of forest ecosystems to fire in the French Alps.** *Eur J For Res* 2019, **138**:813–830.
48. Adagbasa EG, Adelabu SA, Okello TW: **Development of post-fire vegetation response-ability model in grassland mountainous ecosystem using GIS and remote sensing.** *ISPRS J Photogrammetry Remote Sens* 2020, **164**:173–183.
49. Sánchez-Pinillos M, De Cáceres M, Ameztegui A, Coll L: **Temporal dimension of forest vulnerability to fire along successional trajectories.** *J Environ Manag* 2019, **248**:109301.
50. Cutter SL, Boruff BJ, Shirley WL: **Social vulnerability to environmental hazards.** *Soc Sci Q* 2003, **84**:242–261.
51. Andersen LM, Sugg MM: **Geographic multi-criteria evaluation and validation: a case study of wildfire vulnerability in Western North Carolina, USA following the 2016 wildfires.** *Int J Disaster Risk Reduct* 2019, **39**:101123.
52. Paveglio TB, Edgeley CM, Stasiewicz AM: **Assessing influences on social vulnerability to wildfire using surveys, spatial data and wildfire simulations.** *J Environ Manag* 2018, **213**:425–439.
 * This work combines social data surveys, geospatial methods, and wildfire simulations to assess human vulnerability to wildfires, presenting a robust example of interaction among different systems.
53. Nielsen-Pincus M, Ager AA, Evers CR: **Wildfire exposure and community capacity to manage wildfire risk: a coupled bio-physical and sociological analysis of wildfire risk in 60 communities across the western United States.** *Fire* 2019, **2**:59.
54. Edgeley CM, Paveglio TB: **Exploring influences on intended evacuation behaviors during wildfire: what roles for pre-fire actions and event-based cues?** *Int J Disaster Risk Reduct* 2019, **37**:101182.
55. Strahan K, Whittaker J, Handmer J: **Self-evacuation archetypes in Australian bushfire.** *Int J Disaster Risk Reduct* 2018, **27**:307–316.
56. Steer K, Abebe E, Almashor M, Beloglazov A, Zhong X: **On the utility of shelters in wildfire evacuations.** *Fire Saf J* 2017, **94**:22–32.
57. Lindell MK, Perry RW: **The protective action decision model: theoretical modifications and additional evidence.** *Risk Anal* 2012, **32**:616–632.
58. McLennan J, Ryan B, Bearman C, Toh K: **Should we leave now? Behavioral factors in evacuation under wildfire threat.** *Fire Technol* 2019, **55**:487–516.
59. Zhao X, Lovreglio R, Kuligowski E, Nilsson D: **Using artificial intelligence for safe and effective wildfire evacuations.** *Fire Technol* 2020, <https://doi.org/10.1007/s10694-020-00979-x>.
60. Li D, Cova TJ, Dennison PE: **Setting wildfire evacuation triggers by coupling fire and traffic simulation models: a spatiotemporal GIS approach.** *Fire Technol* 2019, **55**:617–642.
61. Paveglio TB, Carroll MS, Stasiewicz AM, Williams DR, Becker DR: **Incorporating social diversity into wildfire management: proposing “Pathways” for fire adaptation.** *For Sci* 2018, **64**:515–532.
62. Billings MC, Carroll MS, Paveglio TB: **The “strings attached” to community difference and potential pathways to fire adaptiveness in the wildland urban interface.** *J For* 2021, **119**:13–27.
63. Dunn CJ, O'Connor CD, Abrams J, Thompson MP, Calkin DE, Johnston JD, Stratton R, Gilbertson-Day J: **Wildfire risk science facilitates adaptation of fire-prone social-ecological systems to the new fire reality.** *Environ Res Lett* 2020, **15**.
64. Ziccardi LG, Thiersch CR, Yanai AM, Fearnside PM, Ferreira-Filho PJ: **Forest fire risk indices and zoning of hazardous areas in Sorocaba, São Paulo state, Brazil.** *J For Res* 2020, **31**:581–590.
65. Wei Y, Thompson MP, Scott JH, O'Connor CD, Dunn CJ: **Designing operationally relevant daily large fire containment strategies using risk assessment results.** *Forests* 2019, **10**:1–23.
66. De Groeve T, Poljansek Karmen, Vernaccini L: *Index for risk management-INFORM*. 2014.
67. Thompson MP, Gannon BM, Caggiano MD, O'Connor CD, Brough A, Gilbertson-Day JW, Scott JH: **Prototyping a geospatial Atlas for wildfire planning and management.** *For-ests* 2020, **11**:1–17.
68. Castellnou M, Prat-Guitart N, Arilla E, Larrañaga A, Nebot E, Castellarnau X, Vendrell J, Pallàs J, Herrera J, Monturiol M, et al.: **Empowering strategic decision-making for wildfire management: avoiding the fear trap and creating a resilient landscape.** *Fire Ecol* 2019, **15**.
69. Dupuy J luc, Fargeon H, Martin-StPaul N, Pimont F, Ruffault J, Guijarro M, Hernando C, Madrigal J, Fernandes P: **Climate change impact on future wildfire danger and activity in southern Europe: a review.** *Ann For Sci* 2020, **77**.
70. Oliveira S, Gonçalves A, Zêzere JL: **Reassessing wildfire susceptibility and hazard for mainland Portugal.** *Sci Total Environ* 2021, **762**:143121.