



Institute for Advanced Studies
in Basic Sciences
Gava Zang, Zanjan, Iran

Regression with Linear Models

By Dr. Parvin Razzaghi (p.razzaghi@iasbs.ac.ir)

Winter 2024





Slides credit

- Professor:
 - Dr. Parvin Razzaghi
- Original Slides:
 - Amir-massoud Farahman
 - Eric Eaton
- Link:
 - <https://B2n.ir/n25709>
- University:
 - University of Toronto and Vector Institute
- Acknowledgment:
 - Credit for slides goes to many members of the ML Group at the U of T, and beyond, including (recent past): Roger Grosse, Murat Erdogdu, Richard Zemel, Juan Felipe Carrasquilla, Emad Andrews.
 - For any comments or suggestions, please contact: p.razzaghi@iasbs.ac.ir





Table of Contents

- Modular Approach to ML
- Regression
 - Linear Regression
 - Basic Expansion
 - Regularization
 - Probabilistic Interpretation of the Squared Error
 - Bias-variance tradeoff
 - Bayesian linear regression



Modular Approach ML Algorithm Design







Modular Approach ML Algorithm Design

- we will take a more **modular** approach:
 - define the problem
 - provide a training dataset
 - choose a **model** describing the relationships between variables of interest
 - define a **loss function** quantifying how bad the fit to the data is
 - (possibly) choose a **regularizer** saying how much we prefer different candidate models (or explanations of data), **before (prior to)** seeing the data
 - fit the model that minimizes the loss function and satisfy the constraint/penalty imposed by the regularizer, possibly using an **optimization algorithm**
- Mixing and matching these modular components gives us a lot of new ML methods.



Define the problem & provide a dataset



The Supervised Learning Setup


- Features
 - Living area
 - Number of bedrooms
 - ...
- Target variable
 - Price



Living area (feet ²)	#bedrooms	Price (1000\$s)
2104	3	400
1600	3	330
2400	3	369
1416	2	232
3000	4	540
⋮	⋮	⋮



The Supervised Learning Setup



Recall that in supervised learning:

- There is a target $t \in T$ (also called response, outcome, output, class)
- There are features $\mathbf{x} \in X$ (also called inputs or covariates)


The goal is to learn a function $f : X \rightarrow T$ such that

$$t \approx y = f(\mathbf{x}),$$

based on given data $D = \{(\mathbf{x}^{(i)}, t^{(i)}) \text{ for } i = 1, 2, \dots, N\}$.




choose a **model** describing the relationships between variables of interest?






Regression with Linear Models




"HEY, I DID A
REGRESSION."




"I WANTED A CURVED
LINE, SO I MADE ONE
WITH MATH."




"LOOK, IT'S
TAPERING OFF!"



"LOOK, IT'S GROWING
UNCONTROLLABLY!"



"I'M SOPHISTICATED, NOT
LIKE THOSE BUMBLING
POLYNOMIAL PEOPLE."



"I'M MAKING A
SCATTER PLOT BUT
I DON'T WANT TO."



Linear Regression – Model


- **Model:** In linear regression, we use linear functions of the inputs $\mathbf{x} = (x_1, \dots, x_D)$ to make predictions y of the target value t :

$$y = f(\mathbf{x}) = \sum_j w_j x_j + b$$


- y is the **prediction**
- \mathbf{w} is the **weights**
- b is the **bias** (or **intercept**) (do not confuse with the bias-variance tradeoff in the next lecture)
- \mathbf{w} and b together are the **parameters**
- We hope that our prediction is close to the target: $y \approx t$.



What is Linear? 1 Feature vs. D Features



- If we have only 1 feature:
 $y = wx + b$ where $w, x, b \in \mathbb{R}$
- y is linear in x




- If we have D features:
 $y = \mathbf{w}^T \mathbf{x} + b$ where $\mathbf{w}, \mathbf{x} \in \mathbb{R}^D, b \in \mathbb{R}$
- y is linear in \mathbf{x}

Relation between the prediction y and inputs \mathbf{x} is linear in both cases.



Weight Space vs. Data Space



Recall that:


$$y = f(x) = \sum_j w_j x_j + b$$



Linear Regression

We have a dataset $D = \{(x^{(i)}, t^{(i)})\}_{i=1}^N$ where:

- $x^{(i)} = (x_1^{(i)}, x_2^{(i)}, \dots, x_D^{(i)})^T \in \mathbb{R}^D$ are the inputs, e.g., age, height
- $t^{(i)} \in \mathbb{R}$ is the target or response, e.g., income
- predict $t^{(i)}$ with a linear function of $x^{(i)}$:



- $t^{(i)} \approx y^{(i)} = \mathbf{w}^T \mathbf{x}^{(i)} + b$
- Find the “best” line (\mathbf{w}, b) .
- **Q:** How should we find the **best** line?




Linear Regression – Loss Function

- How to quantify the quality of the fit to data?
- A **loss function** $\mathcal{L}(y, t)$ defines how bad it is if, for some input x , the algorithm predicts y , but the target is actually t .
- **Squared error loss function:**

$$\mathcal{L}(y, t) = \frac{1}{2} (y - t)^2$$

- $y - t$ is the **residual**, and we want to make its magnitude small
- The $\frac{1}{2}$ factor is just to make the calculations convenient.





Linear Regression – Loss Function

- **Cost function:** loss function averaged over all training examples

$$\mathcal{J}(\mathbf{w}, b) = \frac{1}{N} \sum_{i=1}^N \mathcal{L}(y^{(i)}, t^{(i)})$$

$$= \frac{1}{2N} \sum_{i=1}^N (y^{(i)} - t^{(i)})^2$$

$$= \frac{1}{2N} \sum_{i=1}^N (\mathbf{w}^T x^{(i)} + b - t^{(i)})^2$$



Linear Regression – Loss Function

- To find the best fit, we find a model (parameterized by its weights \mathbf{w} and b) that minimizes the cost:

$$\underset{(\mathbf{w}, b)}{\text{minimize}} \mathcal{J}(\mathbf{w}, b) = \frac{1}{N} \sum_{i=1}^N \mathcal{L}(y^{(i)}, t^{(i)})$$

- The terminology is not universal. Some might call “loss” **pointwise loss** and the “cost function” the **empirical loss** or **average loss**.



Vector Notation

- We can organize all the training examples into a **design matrix \mathbf{X}** with one row per training example, and all the targets into the **target vector \mathbf{t}** .

$$\mathbf{X} = \begin{bmatrix} \mathbf{x}^{(1)T} \\ \mathbf{x}^{(2)T} \\ \mathbf{x}^{(3)T} \end{bmatrix} = \begin{bmatrix} 8 & 0 & 3 & 0 \\ 6 & -1 & 5 & 3 \\ 2 & 5 & -2 & 8 \end{bmatrix}$$

One feature across
all training examples

One training
example (vector)

- Computing the predictions for the whole dataset:

$$\mathbf{X}\mathbf{w} + b\mathbf{1} = \begin{bmatrix} \mathbf{w}^T \mathbf{x}^{(1)} + b \\ \vdots \\ \mathbf{w}^T \mathbf{x}^{(N)} + b \end{bmatrix} = \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(N)} \end{bmatrix} = \mathbf{y}$$



Vectorization

- Computing the squared error cost across the whole dataset:

$$\mathbf{y} = \mathbf{X}\mathbf{w} + b\mathbf{1}$$

$$\mathcal{J} = \frac{1}{2N} \|\mathbf{y} - \mathbf{t}\|^2$$

- Note that sometimes we may use $\mathcal{J} = \frac{1}{2N} \|\mathbf{y} - \mathbf{t}\|^2$, without $\frac{1}{N}$ normalizer. That would correspond to the sum of losses, and not the average loss. That does not matter as the minimizer does not depend on N .



Vectorization

- We can also add a column of 1s to the design matrix, combine the bias and the weights, and conveniently write


$$X = \begin{bmatrix} 1 & [x^{(1)}]^T \\ 1 & [x^{(2)}]^T \\ 1 & \vdots \end{bmatrix} \in \mathbb{R}^{N \times D+1} \text{ and } w = \begin{bmatrix} b \\ w_1 \\ w_2 \\ \vdots \end{bmatrix} \in \mathbb{R}^{D+1}$$

Then, our predictions reduce to $\mathbf{y} = \mathbf{X}\mathbf{w}$.




Intuition on Minimizing MSE Loss

- Consider $x \in \mathbb{R}$ and $w \in \mathbb{R}$



$$w = 1$$


$$J(w, D)$$






Intuition on Minimizing MSE Loss

- Consider $x \in \mathbb{R}$ and $w \in \mathbb{R}$




$$w = 0.5$$






Intuition on Minimizing MSE Loss

- Consider $x \in \mathbb{R}$ and $w \in \mathbb{R}$



$w = 0.25$


$J(w, D)$






Intuition on Minimizing MSE Loss

- Consider $x \in \mathbb{R}$ and $w \in \mathbb{R}$





$J(w, D)$






Intuition on Minimizing MSE Loss

- **Convex** (“bowl shaped”) in general





Solving the Minimization Problem



- We defined a model (linear).
- We defined a loss and the cost function to be minimized.
- **Q:** How should we solve this minimization problem?



Solving the Minimization Problem

- Recall from your calculus class: minimum of a differentiable function (if it exists) occurs at a **critical point**, i.e., point where the derivative is zero.
- Multivariate generalization: set the partial derivatives to zero (or equivalently the gradient).
- We would like to find a point where the gradient is (close to) zero. How can we do it?
- Sometimes it is possible to directly find the parameters that make the gradient zero in a closed-form. We call this the **direct solution**.
- We may also use optimization techniques that iteratively get us closer to the solution. We will get back to this soon.



Direct Solution

- **Partial derivatives:** derivatives of a multivariate function with respect to (w.r.t.) one of its arguments.

$$\frac{\partial}{\partial x_1} f(x_1, x_2) = \lim_{h \rightarrow 0} \frac{f(x_1 + h, x_2) - f(x_1, x_2)}{h}$$



Direct Solution

- To compute, take the single variable derivatives, pretending the other arguments are constant.
- Example: partial derivatives of the prediction y with respect to weight w_j and bias b :

$$\frac{\partial y}{\partial w_j} = \frac{\partial}{\partial w_j} \left[\sum_{j'} w_{j'} x_{j'} + b \right] = x_j$$

$$\frac{\partial y}{\partial b} = \frac{\partial}{\partial b} \left[\sum_{j'} w_{j'} x_{j'} + b \right] = 1$$



Direct Solution

- **The derivative of loss:** We apply the chain rule: first we take the derivative of the loss L w.r.t. output y of the model, and then the derivative of the output y w.r.t. a parameter of the model such as w_j or b :

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial w_j} &= \frac{d\mathcal{L}}{dy} \frac{\partial y}{\partial w_j} \\ &= \frac{d}{dy} \left[\frac{1}{2} (y - t)^2 \right] \cdot x_j \\ &= (y - t) x_j \\ \frac{\partial \mathcal{L}}{\partial b} &= y - t\end{aligned}$$



Direct Solution

- Cost derivatives (average over data points):

$$\frac{\partial \mathcal{J}}{\partial w_j} = \frac{1}{N} \sum_{i=1}^N (y^{(i)} - t^{(i)}) x_j^{(i)}$$

$$\frac{\partial \mathcal{J}}{\partial b} = \frac{1}{N} \sum_{i=1}^N (y^{(i)} - t^{(i)})$$



Direct Solution

- Recall that the output y is a function of the parameters as $y = \mathbf{w}^T \mathbf{x}$.
- The minimum of the cost function must occur at a point where the partial derivatives are zero, i.e.,

$$\nabla_{\mathbf{w}} \mathcal{J} = 0 \iff \frac{\partial \mathcal{J}}{\partial w_j} = 0 \quad (\forall j), \quad \frac{\partial \mathcal{J}}{\partial b} = 0.$$

- If $\partial \mathcal{J} / \partial w_j \neq 0$, you could reduce the cost by changing w_j .



Direct Solution

If we follow this recipe, we get that we have to set the gradient of

$\mathcal{J} = \frac{1}{2N} \|\mathbf{y} - \mathbf{t}\|^2$, with $\mathbf{y} = \mathbf{X}\mathbf{w}$ (bias absorbed in \mathbf{X}) equal to zero.

We have

$$\mathcal{J} = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{t})^T (\mathbf{X}\mathbf{w} - \mathbf{t}),$$

So

$$\nabla_{\mathbf{w}} \mathcal{J} = \frac{1}{N} \mathbf{X}^T (\mathbf{X}\mathbf{w} - \mathbf{t}) = 0 \implies (\mathbf{X}^T \mathbf{X})\mathbf{w} = \mathbf{X}^T \mathbf{t}.$$

This is a **linear system of equations**.

Q: What are the dimensions of each component?



Direct Solution

- Assuming that $\mathbf{X}^T \mathbf{X}$ is invertible, the optimal weights are

$$\mathbf{w}^{LS} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{t}.$$


This solution is also called **Ordinary Least Squares (OLS)** solution.

At an arbitrary point \mathbf{x} , our prediction is $y = \mathbf{w}^{LS^T} \mathbf{x}$.

Q: What happens if $\mathbf{X}^T \mathbf{X}$ is not invertible?

Basis Expansion (Feature Mapping)


- The relation between the input and output may not be linear.



- We can still use linear regression by mapping the input feature to another space using **basis expansion** (or **feature mapping**) $\psi(\mathbf{x}) : \mathbb{R}^D \rightarrow \mathbb{R}^d$ and treat the mapped feature (in \mathbb{R}^d) as the input of a linear regression procedure.
- Let us see how it works when $x \in \mathbb{R}$ and we use polynomial feature mapping.



Polynomial Feature Mapping




- Fit the data using a degree- M polynomial function of the form:

$$y = w_0 + w_1x + w_2x^2 + \dots + w_Mx^M = \sum_{i=0}^M w_i x^i$$

- The feature mapping is $\psi(x) = [1, x, x^2, \dots, x^M]^T$.
- We can still use the linear regression framework with least squares loss to find \mathbf{w} since $y = \psi(x)^T \mathbf{w}$ is linear in w_0, w_1, \dots .
- In general, ψ can be any function. Another example: Fourier map
$$\psi = [1, \sin(2\pi x), \cos(2\pi x), \sin(4\pi x), \cos(4\pi x), \sin(6\pi x), \cos(6\pi x), \dots]^T.$$



Polynomial Feature Mapping with $M = 0$



Any questions?

