

Anchored Geodesic Analysis for Multivariate Extremes

Alberto Quaini* Chen Zhou

Erasmus University Rotterdam

Abstract

Extremal dependence is naturally described by the angular law of large multivariate observations. We introduce anchored geodesic component analysis (AGCA), a dimension-reduction method for extremal angular laws on the positive unit sphere. AGCA approximates angular variation by great subspheres constrained to pass through a chosen reference direction, with balanced complete dependence as the default anchor. Under a bounded sine-squared geodesic loss, the population and empirical problems reduce exactly to eigenanalysis of a second-moment matrix of anchored tangent departures. The resulting scores, loadings, residual risks and explained-variation summaries describe departures from the benchmark and remain well defined for face and near-axis extremes. Low-rank AGCA reconstructions also support tail simulation: bounded Lipschitz functionals and homogeneous tail scores, including portfolio capped excesses and value-at-risk, inherit explicit error bounds from the AGCA residual risk. We establish top- k consistency for oracle and rank-Pareto AGCA summaries and an oracle central limit theorem whose covariance is that of an independent sample from the limiting angular law. In daily equity-portfolio losses, AGCA finds concentrated benchmark-relative tail directions: ten components explain about 91% of anchored variation and approximate capped-excess and normalized value-at-risk summaries with about 1.25% average relative error.

Keywords: Multivariate extremes; geodesic analysis; angular measure; dimension reduction; tail simulation; portfolio tail risk.

JEL classification: C14; C38; C58; G11.

*Corresponding author: quaini@ese.eur.nl. We thank Nicola Gnecco for the valuable comments.

1 Introduction

Extremes increasingly arise as multivariate events. A storm is a joint profile over a region, a market crisis is a configuration of simultaneous tail losses, and an infrastructure failure may be visible only through several stressed measurements. For such data, the central question is extremal dependence: how do coordinates participate in large events, and can the relative pattern of participation be summarized in a small number of interpretable coordinates?

Classical multivariate extreme value theory formalizes this question through angular measures. After standardizing margins to a common tail scale, a large observation decomposes into radial size and angular direction; the limiting angular law records the relative contribution of the coordinates in the tail (De Haan and Ferreira, 2006; Resnick, 2007). This representation underlies peaks-over-threshold models and multivariate generalized Pareto limits (Rootzén and Tajvidi, 2006; Rootzén et al., 2018; Mourahib et al., 2025). It is also a natural starting point for dimension reduction, because the angular law is the object that remains after radial severity has been separated from extremal direction.

This paper develops anchored geodesic component analysis (AGCA) for that purpose. We represent standardized extreme directions on the Euclidean unit sphere, which preserves extremal rays in finite dimension while placing the angular component on a familiar Riemannian manifold. The default reference direction is the balanced complete-dependence anchor μ_0 in (2.1), corresponding to equal participation of all variables after marginal standardization. AGCA fits great subspheres through this anchor and measures reconstruction by a bounded sine-squared geodesic loss. The resulting scores, loadings, residual risks, and explained variation describe angular departures from balanced extremal co-movement.

The construction is related to principal geodesic analysis and principal nested spheres, which adapt principal component ideas to nonlinear spaces (e.g., Fletcher et al., 2004; Jung et al., 2012; Sommer et al., 2014; Tabaghi et al., 2024). Those methods typically build geodesic summaries around data-adaptive base points and use squared geodesic reconstruction loss. AGCA instead uses a reference direction tailored to extremal dependence and a bounded sine-squared geodesic

loss. In contrast to principal nested spheres, which allow nested small subspheres rather than only great subspheres (Jung et al., 2012), AGCA deliberately restricts to great subspheres through the anchor. The restriction makes the bounded sine-squared loss reduce spherical projection to ordinary orthogonal projection of bounded departures in the tangent hyperplane. Consequently, the population and empirical AGCA problems reduce exactly to eigendecomposition of an anchored second-moment matrix. In addition, since all standardized extremal rays are represented on the sphere, near-axis directions associated with asymptotic independence remain finite angular observations.

This exact spectral form is useful both computationally and statistically. It gives one eigen-solution for the fitted component space and for diagnostics such as anchored variation explained, residual geodesic risk, scores, loadings, and component-space stability. It also makes perturbation analysis transparent: error in the anchored second-moment matrix translates into eigenvalue error, residual-risk error, and, under an eigengap, projector error for the fitted component space. The loadings retain a direct extremal interpretation. Near μ_0 , loading differences determine how pairwise angular participation changes relative to balanced complete dependence.

Several neighboring literatures reduce the angular complexity, but they target different objects. Support and sparsity methods ask which variables, faces, subcones, or representative directions can become extreme together (Chautru, 2015; Goix et al., 2017; Meyer and Wintenberger, 2021). Extremal graphical models encode conditional independence and sparse Markov structure in tail limits (Engelke and Hitz, 2020). Tail dependence matrices and their decompositions summarize pairwise extremal association, while necessarily losing higher-order angular information (Cooley and Thibaud, 2019). Clustering methods for angular extremes group directions into representative regimes (Janßen and Wan, 2020; Fomichov and Ivanovs, 2023). Principal component methods for extremes seek low-dimensional linear subspaces close to extreme directions and study reconstruction risk, excess risk, and component selection (Drees and Sabourin, 2021; Drees, 2025; Butsch and Fasen-Hartmann, 2025; Cooley et al., 2026). Kernel, hyperplane, and compositional approaches provide further projections or transformed linear coordinates for angular samples (Avella Medina

et al., 2025; Wan, 2026; Lhaut et al., 2026). These methods are valuable when the target is an active face set, a sparse tail graph or direction set, a collection of representative regimes, a pairwise matrix, or a linear representation. They do not directly provide a benchmark-relative component analysis of the angular law itself. Several of these approaches impose sparsity, discrete-regime, or linear-subspace structure, whereas AGCA is designed to summarize the angular law itself without assuming sparsity and remains well defined for face and near-axis extremes.

The paper contributes at population, estimation, and inferential levels. At the population level, we define the anchored geodesic reconstruction problem, prove the exact projection identity, derive the eigensolution and loading interpretation, and characterize face- and axis-supported angular laws. Then we show how low-rank AGCA reconstructions can simulate bounded Lipschitz and homogeneous tail functionals, including portfolio capped excesses and Value-at-Risk. At the sample level, we give the rank-Pareto top- k construction used in applications, building on empirical angular measure estimation as in Einmahl et al. (1997, 2001); Einmahl and Segers (2009), and establish consistency and stability of scalar and component-space AGCA summaries. We also prove an oracle central limit theorem for the anchored top- k second-moment matrix. Under a second-order angular-bias condition, random threshold selection has the same first-order limit as an independent sample from the limiting angular law, yielding plug-in inference for the AGCA-specific eigenvalues, anchored explained variation, residual geodesic risk, and component projectors. Unlike the multivariate regular variation framework common to this line (Drees and Sabourin, 2021; Drees, 2025; Butsch and Fasen-Hartmann, 2025), the oracle AGCA CLT depends on angular convergence and a second-order angular-bias condition rather than full radial regular variation or tail-moment assumptions.

The empirical application studies daily equity-portfolio loss extremes. In a $d = 24$ Fama–French panel (French, 2026), the AGCA spectrum is concentrated: the first five components explain 76.3% of anchored variation and the first ten explain 91.1%. The loadings identify repeated cross-sectional contrasts in extreme portfolio-loss directions. The low-rank reconstruction also preserves the portfolio tail summaries motivated by the theory: by rank ten, the average relative error across




Figure 1: Sphere-level geometry in two controlled simulations. The left panel shows a low-dimensional angular law; the right panel shows a variable-specific near-axis regime. Each panel displays selected extreme directions, the canonical anchor and first AGC, and the Fréchet anchor and first AGC.

capped-excess and normalized value-at-risk summaries is about 1.25% over a broad portfolio class. A second daily anomaly portfolio panel from Open Source Asset Pricing (Chen and Zimmermann, 2026) gives the same qualitative message, with a less compressed but still stable anchored spectrum.

The Supplementary Material collects proofs, complementary theory, simulation analyses, and complementary empirical results. The simulations illustrate AGCA in designs that vary in complexity of the angular law and the number of near-axis directions; see Figure 1 for a simple visual preview. The AGCA methodology is implemented in the R package `AGCA4extremes` (<https://github.com/a91quaini/AGCA4extremes>). Data and replication code for the simulation and empirical analyses are available at <https://github.com/a91quaini/replicateAGCApaper>.

2 AGCA: population theory

We use the following notation throughout. For $x, y \in \mathbb{R}^d$, $\langle x, y \rangle = x^\top y$ and $\|x\|_2 = \langle x, x \rangle^{1/2}$. Let $\mathbb{S}^{d-1} := \{x \in \mathbb{R}^d : \|x\|_2 = 1\}$, and let $\mathbb{S}_+^{d-1}$ and $\mathbb{S}_{++}^{d-1}$ denote its intersections with the nonnegative orthant and the strictly positive orthant, respectively. For $x, y \in \mathbb{S}^{d-1}$,

$$d_g(x, y) := \arccos(\langle x, y \rangle)$$

denotes the spherical geodesic distance, and for $B \subset \mathbb{S}^{d-1}$, $d_g(x, B) := \inf_{b \in B} d_g(x, b)$. For $\mu \in \mathbb{S}^{d-1}$, $\mu^\perp := \{x \in \mathbb{R}^d : \langle x, \mu \rangle = 0\}$ denotes the tangent hyperplane to the sphere at μ , and for a linear subspace W , Π_W is the Euclidean orthogonal projector onto W . The matrix I is the identity, $\mathbf{1}$ is the vector of ones, and $\text{span}(\cdot)$, $\text{tr}(\cdot)$, $\text{rank}(\cdot)$, $\text{range}(\cdot)$, and $\text{diag}(\cdot)$ denote span, trace, rank, range, and diagonal matrix, respectively. The operator norm is $\|\cdot\|_{\text{op}}$, $|B|$ is the cardinality of a finite set B , and $a_+ := \max(a, 0)$ is the positive part of $a \in \mathbb{R}$. The indicator of an event A is $\mathbb{I}\{A\}$, $\mathcal{L}(Z)$ is the law of a random element Z , $\mathbb{P}$, $\mathbb{E}$, Var , and Cov denote probability, expectation, variance, and covariance, respectively, $\rightsquigarrow$ denotes weak convergence, and stochastic order notation such as $o_p(\cdot)$ has its usual meaning. For a bounded real-valued function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, let

$$\|f\|_\infty := \sup_{z \in \mathbb{R}^d} |f(z)|, \quad \text{Lip}(f) := \sup_{z \neq z'} \frac{|f(z) - f(z')|}{\|z - z'\|_2},$$

with the convention $\text{Lip}(f) = \infty$ if f is not Lipschitz.

2.1 Angular limits and the anchor

Let $Y = (Y_1, \dots, Y_d)$ have continuous marginal distribution functions $F_1, \dots, F_d$, and put the data on a common tail scale by the standard Pareto transform

$$X := ((1 - F_1(Y_1))^{-1}, \dots, (1 - F_d(Y_d))^{-1}).$$

Write $R := \|X\|_2$. For standardized X , the norm used for polar normalization only chooses one representative on each positive ray; it preserves component ratios and does not change the finite-dimensional tail measure (De Haan and Ferreira, 2006; Resnick, 2007, Chapter 6). We use the Euclidean representative because it places angular extremes on the ordinary unit sphere, where geodesics, tangent spaces, great subspheres, and orthogonal projections have their standard forms.

The population target is the limiting conditional law of the direction $X/\|X\|_2$ given large R . It records relative participation across variables in an extreme episode after marginal tail scales and radial size have been removed.

Assumption 1 (Euclidean-spherical angular limit). *There exists a random vector G taking values in $\mathbb{S}_+^{d-1}$ such that*

$$\mathcal{L}(X / \|X\|_2 \mid \|X\|_2 > r) \rightsquigarrow \mathcal{L}(G), \quad r \rightarrow \infty.$$

Assumption 1 is the Euclidean-spherical form of standard angular convergence in multivariate extremes. It is deliberately only directional, so it allows mass near the diagonal, coordinate faces, or axes. It is implied by Euclidean-polar regular variation as in Assumption 2 below, but is weaker because it does not require radial regular variation or a homogeneous tail measure on the cone.

AGCA is built relative to an interior reference direction $\mu \in \mathbb{S}_{++}^{d-1}$. The anchor fixes the origin of the score system. It may be specified from subject-matter considerations, chosen as a canonical benchmark, or estimated from the angular law. Our default choice is

$$\mu_0 := d^{-1/2} \mathbf{1}. \tag{2.1}$$

The direction μ_0 in (2.1) is the spherical image of the diagonal ray $t \mapsto t\mathbf{1}$. This is the balanced complete-dependence direction: after Pareto standardization, all components participate equally in the angular profile. With μ_0 , AGCA components are coordinates for asymmetric departures from balanced extremal co-movement. Because the canonical anchor does not depend on the sample, threshold, or fitted angular law, it can be used across data sets and thresholds, and the resulting loadings retain the interpretation of departures from balanced co-movement. The canonical anchor also has a distribution-free minimax justification: among all interior anchors, it uniquely maximizes the smallest cosine similarity with the positive sphere, equivalently minimizing the worst-case geodesic distance and worst-case gnomonic coordinate size; see Supplementary Proposition S9.

Data-adaptive anchors can improve empirical alignment, but they change the reference profile: the resulting components describe variation around an empirical angular benchmark rather than a fixed scientific one. This is analogous to the role of estimated base points in geodesic dimension-reduction methods (Fletcher et al., 2004; Huckemann and Ziezold, 2006; Jung et al., 2012; Dai and Müller, 2018). Two commonly used references are the spherical Fréchet anchor and the principal

anchor,

$$\mu_F \in \arg \min_{\mu \in \mathbb{S}_+^{d-1}} \mathbb{E}[d_g(G, \mu)^2], \quad \mu_P := v_1(\mathbb{E}[GG^\top]), \quad (2.2)$$

where $v_1(M)$ denotes the unit leading eigenvector of a matrix M , oriented to lie in $\mathbb{S}_{++}^{d-1}$ when possible. We use the Fréchet reference as an AGCA anchor when a chosen minimizer lies in $\mathbb{S}_{++}^{d-1}$; Supplementary Section S1.3 discusses these anchors and anchor-free alternatives.

2.2 Anchored coordinates and model spaces

For an interior anchor μ , all positive angular directions lie in the open hemisphere

$$\mathcal{H}_\mu := \{g \in \mathbb{S}^{d-1} : \langle \mu, g \rangle > 0\},$$

because $\langle \mu, g \rangle > 0$ for every $g \in \mathbb{S}_+^{d-1}$. Thus coordinate-face and near-axis extremes remain in a single anchored chart. For $g \in \mathcal{H}_\mu$, write $a_\mu(g) := \langle \mu, g \rangle$ for the anchor coordinate and $u_\mu(g) := (I - \mu\mu^\top)g \in \mu^\perp$ for the bounded tangent departure. Since $a_\mu(g) = \cos d_g(g, \mu)$, the distance $d_g(g, \mu) = \arccos a_\mu(g)$ measures how far g lies from the anchor along the sphere, while $u_\mu(g)$ records the departure from the anchor axis inside the tangent hyperplane. The decomposition

$$g = a_\mu(g)\mu + u_\mu(g), \quad a_\mu(g) = (1 - \|u_\mu(g)\|_2^2)^{1/2},$$

separates the anchored coefficient from the orthogonal departure. We also use the gnomonic coordinate centered at μ ,¹

$$v_\mu(g) := \frac{g}{\langle \mu, g \rangle} - \mu \in \mu^\perp, \quad \phi_\mu(v) := \frac{\mu + v}{(1 + \|v\|_2^2)^{1/2}}, \quad v \in \mu^\perp.$$

The coordinate $v_\mu(g)$ is the displacement obtained by projecting the ray through g onto the tangent affine hyperplane $\mu + \mu^\perp$, and ϕ_μ maps such tangent displacements back to the sphere. The bounded

¹Gnomonic coordinates project points from the sphere's center onto the tangent affine hyperplane at μ . Great subspheres through μ become linear subspaces in this chart.




Figure 2: Geometry of anchored geodesic component analysis. In the left panel, the open hemisphere $\mathcal{H}_\mu$ contains the positive Euclidean-spherical directions, and the anchored great subsphere $M_\mu(W)$ is the approximating set through the anchor. In the central panel, the gnomonic chart centered at μ sends the hemisphere $\mathcal{H}_\mu$ to the tangent affine hyperplane $\mu + \mu^\perp$, identified with $\mu^\perp$, and maps the anchored great subsphere $M_\mu(W)$ to the linear subspace W . The bounded departure $u_\mu(g)$ is a radial rescaling of the gnomonic coordinate $v_\mu(g)$. In the right panel, under the loss $\sin^2 d_g$, geodesic approximation of g by $M_\mu(W)$ on the sphere is the same as Euclidean orthogonal projection of $u_\mu(g)$ onto W in $\mu^\perp$.

and gnomonic coordinates satisfy

$$u_\mu(g) = \frac{v_\mu(g)}{(1 + \|v_\mu(g)\|_2^2)^{1/2}}, \quad v_\mu(g) = \frac{u_\mu(g)}{a_\mu(g)}.$$

Thus v_μ is the linear chart for anchored great subspheres, while u_μ is the bounded rescaling used for fitting. Near the edge of the hemisphere, $v_\mu(g)$ can be large, whereas $u_\mu(g)$ always has norm at most one and keeps the same tangent direction.

For a linear subspace $W \subset \mu^\perp$, define the anchored great-subsphere model

$$M_\mu(W) := \text{span}(\{\mu\} \cup W) \cap \mathcal{H}_\mu.$$

Thus $M_\mu(W)$ is the great subsphere through the anchor whose allowed directions away from the anchor are the tangent directions in W , restricted to the open hemisphere $\mathcal{H}_\mu$. The rank of the approximation is $\dim(W)$: rank zero is the anchor alone, and rank p allows p tangent directions through the anchor. The target is therefore a low-dimensional geodesic summary of angular variation around a fixed benchmark. Figure 2 shows the model set, the gnomonic chart, and the bounded projection geometry.

2.3 Population criterion and eigensolution

With the anchor μ fixed, the population fitting problem is to choose the tangent subspace $W \subset \mu^\perp$ that best reconstructs the limiting angular direction. Let G denote the limiting angular direction in Assumption 1. For a p -dimensional subspace $W \subset \mu^\perp$, AGCA uses the anchored geodesic risk

$$\mathcal{R}_\mu(W) := \mathbb{E}[\sin^2 d_g(G, M_\mu(W))], \quad d_g(g, M_\mu(W)) := \inf_{m \in M_\mu(W)} \arccos(\langle g, m \rangle). \quad (2.3)$$

Here, $d_g(g, M_\mu(W))$ denotes the shortest spherical distance from $g \in \mathbb{S}_+^{d-1}$ to the anchored great-subsphere model. The rank p is the dimension of the tangent subspace used to reconstruct the angular law. Equivalently, it is the number of anchored geodesic component directions retained.

The sine-squared geodesic residual is a central modeling choice. In principal geodesic analysis and principal nested spheres, reconstruction is usually measured by squared geodesic distance, often around a Fréchet mean, fitted base point, or recursively fitted small sphere (e.g., [Fletcher et al., 2004](#); [Jung et al., 2012](#); [Sommer et al., 2014](#); [Tabaghi et al., 2024](#)). That loss is the direct Riemannian analogue of Euclidean least squares, but it does not generally yield a quadratic projection problem; implementations typically require tangent-space approximations or nonlinear optimization over geodesics, base points, or spherical submanifolds. For the anchored positive-sphere problem we instead use $\sin^2 d_g$: on the admissible hemisphere it is bounded, monotone in geodesic residual distance, comparable to squared geodesic loss, and has the same zero-risk sets; see Supplementary Proposition S3. Its key advantage is exact linearization: the spherical residual becomes an ordinary Euclidean residual of the bounded departure $u_\mu(g)$, which leads to the eigendecomposition below.

Proposition 1 (Exact anchored projection identity). *For every linear subspace $W \subset \mu^\perp$ and every $g \in \mathbb{S}_+^{d-1}$,*

$$\sin^2 d_g(g, M_\mu(W)) = \|u_\mu(g) - \Pi_W u_\mu(g)\|_2^2. \quad (2.4)$$

Moreover, the unique geodesic projection of g onto $M_\mu(W)$ is

$$\Pi_{M_\mu(W)}(g) = \frac{a_\mu(g)\mu + \Pi_W u_\mu(g)}{(a_\mu(g)^2 + \|\Pi_W u_\mu(g)\|_2^2)^{1/2}}. \quad (2.5)$$

Proposition 1 uses standard spherical geometry to provide the exact linearization step for our fixed-anchor extremal-angular reconstruction problem, as illustrated in the right panel of Figure 2. For a fixed anchor, the bounded geodesic residual is exactly the Euclidean residual obtained by projecting $u_\mu(g)$ onto W . The formula for $\Pi_{M_\mu(W)}(g)$ then normalizes the retained anchored departure back to the sphere. The identity is conditional on the anchor being held fixed: it applies after choosing either the canonical anchor or a data-adaptive anchor, but it is not a joint optimization over the anchor and the component space.

Applying the identity to the angular limit G from Assumption 1 turns AGCA into a second-moment problem for bounded anchored departures. Define

$$U_\mu := u_\mu(G), \quad \Sigma_\mu := \mathbb{E}[U_\mu U_\mu^\top], \quad (2.6)$$

where Σ_μ acts on the tangent hyperplane $\mu^\perp$. The vector U_μ records how the angular direction departs from the fixed anchor after radial magnitude has been removed. Because U_μ is bounded, Σ_μ is finite even when the angular law has mass near coordinate faces or axes, as in asymptotically independent regimes. Thus the origin of $\mu^\perp$ is the anchor, and AGCA approximates benchmark-relative departures by linear subspaces through that origin. This boundary behavior is a distinguishing feature relative to many projection, graphical, and hyperplane summaries of extremes: Supplementary Section S1.2 shows that face- and axis-supported angular laws enter the same finite second-moment problem, with axis-dominant laws producing interpretable loading contrasts and symmetric axis laws producing tied eigenvalues.

Theorem 1 (Population eigensolution). *For every $p = 0, \dots, d - 1$ and every p -dimensional $W \subset \mu^\perp$, the anchored geodesic risk satisfies $\mathcal{R}_\mu(W) = \text{tr}((I - \Pi_W)\Sigma_\mu)$. Write the spectral*

decomposition of Σ_μ on $\mu^\perp$, with $b_1, \dots, b_{d-1}$ orthonormal in $\mu^\perp$, as

$$\Sigma_\mu = \sum_{j=1}^{d-1} \lambda_j b_j b_j^\top, \quad \lambda_1 \geq \dots \geq \lambda_{d-1} \geq 0. \quad (2.7)$$

For $p = 0, \dots, d-1$, define the population component space and spectral projector, and for $1 \leq p \leq d-2$, define the associated eigengap, by

$$\begin{aligned} W_{\mu,p} &:= \text{span}(b_1, \dots, b_p), & W_{\mu,0} &:= \{0\}, \\ P_{\mu,p} &:= \sum_{j=1}^p b_j b_j^\top, & P_{\mu,0} &:= 0, & \Delta_{\mu,p} &:= \lambda_p - \lambda_{p+1}, \quad 1 \leq p \leq d-2. \end{aligned} \quad (2.8)$$

Then $W_{\mu,p}$ minimizes $\mathcal{R}_\mu(W)$ among all p -dimensional anchored subspaces, and $\mathcal{R}_\mu(W_{\mu,p}) = \sum_{j=p+1}^{d-1} \lambda_j$.

Theorem 1 is the population PCA theorem for bounded anchored departures. It gives an exact spectral solution for a geodesic reconstruction problem on extremal angular laws: the leading eigenvectors of Σ_μ are the optimal anchored component directions, and the trailing eigensum is the residual geodesic risk.

2.4 Rank, scores, and loading interpretation

The total anchored variation is

$$\tau_\mu := \text{tr}(\Sigma_\mu) = \mathbb{E}[\|U_\mu\|_2^2] = \mathbb{E}[1 - \langle \mu, G \rangle^2]. \quad (2.9)$$

It is the rank-zero risk, namely the average sine-squared distance from G to the anchor. The eigenvalues λ_j decompose this variation exactly: λ_j is the amount of anchored geodesic variation captured by the j th component, and the trailing eigensum is the residual risk after retaining p components. For $p = 0, \dots, d-1$, the population rank- p residual risk and, when $\tau_\mu > 0$, the

proportion of anchored geodesic variation explained are

$$\rho_{\mu,p} := \sum_{j=p+1}^{d-1} \lambda_j, \quad \text{AVE}_{\mu,p} := (\tau_\mu)^{-1} \sum_{j=1}^p \lambda_j. \quad (2.10)$$

Here $P_{\mu,p}$ in (2.8) is the rank- p population spectral projector, and $\Delta_{\mu,p}$ is the eigengap separating the retained and discarded component spaces. This gives the same operational clarity as ordinary PCA, but the interpretation is tied to geodesic approximation of spherical extremal directions through the fixed anchor. The next corollary identifies r_μ as the exact anchored dimension: the smallest number of anchored component directions needed for zero population reconstruction risk.

Corollary 1 (Exact anchored rank). *Let $r_\mu := \text{rank}(\Sigma_\mu)$. There exists a p -dimensional subspace $W \subset \mu^\perp$ with $\mathcal{R}_\mu(W) = 0$ if and only if $p \geq r_\mu$. At the minimal dimension $p = r_\mu$, the exact component space is unique and equals $\text{range}(\Sigma_\mu)$.*

The scores and loadings have a direct interpretation as extremal contrasts. A loading eigenvector $b_\ell \in \mu^\perp$ is a tangent direction through the anchor, and the score of an angular direction g_i is its coordinate along that direction,

$$\alpha_{i\ell} := \langle b_\ell, u_\mu(g_i) \rangle. \quad (2.11)$$

Large positive or negative scores indicate strong movement away from the anchor along the corresponding loading, while scores near zero indicate little contribution from that contrast. The overall sign of a loading is conventional, but relative signs within a loading are meaningful once an orientation is fixed.

Given a rank p , the retained tangent departure is

$$\Pi_{W_{\mu,p}} u_\mu(g_i) = \sum_{\ell=1}^p \alpha_{i\ell} b_\ell. \quad (2.12)$$

Using (2.5), normalization back to the sphere gives

$$g_i^{(p)} = \frac{a_\mu(g_i)\mu + \sum_{\ell=1}^p \alpha_{i\ell} b_\ell}{\left(a_\mu(g_i)^2 + \|\sum_{\ell=1}^p \alpha_{i\ell} b_\ell\|_2^2\right)^{1/2}}. \quad (2.13)$$

This is the AGCA reconstruction of the angular profile; radial severity is not reconstructed.

Proposition 2 (Loadings as first-order log-ratio contrasts). *Let $\eta \in \mu^\perp$, and let $c_\eta(t) := \text{Exp}_\mu(t\eta)$ be the geodesic through μ with initial tangent direction η , where Exp_μ is the Riemannian exponential map on the unit sphere at μ . Then, for every pair of coordinate indices j, m ,*

$$\left. \frac{d}{dt} \right|_{t=0} \log \frac{c_{\eta,j}(t)}{c_{\eta,m}(t)} = \frac{\eta_j}{\mu_j} - \frac{\eta_m}{\mu_m}.$$

Proposition 2 gives the local ratio interpretation. Along a component direction η , the sign of $\eta_j/\mu_j - \eta_m/\mu_m$ determines whether positive scores increase or decrease the pairwise angular ratio g_j/g_m relative to the anchor ratio μ_j/μ_m . For the canonical anchor μ_0 , the baseline ratio is one and the derivative reduces to $d^{1/2}(\eta_j - \eta_m)$. Thus loading contrasts identify which variables drive benchmark-relative changes in angular participation. This is why the canonical anchor is the default in the simulations and empirical application: data-driven anchors may improve fit, but their loadings are less clearly interpretable because they describe departures from an empirical angular benchmark rather than from balanced complete dependence.

2.5 Low-rank reconstruction and tail simulation

The population eigensolution also defines a natural angular reconstruction. This is useful when AGCA is used not only as a descriptive summary, but also as a low-dimensional device for simulating extreme directions. The reconstruction should be understood as an angular approximation: AGCA controls how far the limiting direction moves on the sphere, while a separate radial tail model is needed to simulate the size of the extreme event.

Using the component space $W_{\mu,p}$ in (2.12), we define the rank- p reconstructed angular direction

as the geodesic projection of the angular limit G from Assumption 1 onto the AGCA model:

$$\tilde{G}_{\mu,p} := \Pi_{M_\mu(W_{\mu,p})}(G) = \frac{a_\mu(G)\mu + \Pi_{W_{\mu,p}}u_\mu(G)}{\left(a_\mu(G)^2 + \|\Pi_{W_{\mu,p}}u_\mu(G)\|_2^2\right)^{1/2}}. \quad (2.14)$$

Thus the reconstruction retains the projection of $u_\mu(G)$ onto the leading AGCA component space and renormalizes the result to a spherical direction. The next proposition shows that this unconstrained spherical reconstruction has an exact AGCA error and a corresponding Euclidean error bound.

Proposition 3 (AGCA angular reconstruction). *For $p = 0, \dots, d-1$,*

$$\sin^2 d_g(G, \tilde{G}_{\mu,p}) = \|u_\mu(G) - \Pi_{W_{\mu,p}}u_\mu(G)\|_2^2.$$

Consequently,

$$\mathbb{E}[\sin^2 d_g(G, \tilde{G}_{\mu,p})] = \rho_{\mu,p}, \quad \mathbb{E}\left[\|G - \tilde{G}_{\mu,p}\|_2^2\right] = \mathbb{E}\left[\frac{2 \sin^2 d_g(G, \tilde{G}_{\mu,p})}{1 + \cos d_g(G, \tilde{G}_{\mu,p})}\right] \leq 2\rho_{\mu,p}.$$

To turn an angular reconstruction into a tail simulator, the radial size must be generated separately and then attached to the reconstructed direction. Let $S > 0$ denote such a radial draw. If the same radius is attached to G and to $\tilde{G}_{\mu,p}$, the relative Euclidean error is purely angular and is controlled by the AGCA residual risk:

$$\mathbb{E}\left[\|SG - S\tilde{G}_{\mu,p}\|_2^2 / S^2\right] = \mathbb{E}\left[\|G - \tilde{G}_{\mu,p}\|_2^2\right] \leq 2\rho_{\mu,p}.$$

Thus AGCA controls the relative vector error once the radial size has been supplied, but it does not supply the radial tail law itself. For simulation of extremes one must combine the reconstructed angular law with an extreme-value model for the radius. We use the following standard Euclidean-polar regular variation condition for this purpose.

Assumption 2 (Euclidean-polar regular variation). *There exists $\alpha > 0$ such that, for a Pareto*

variable P_α independent of G and satisfying $\mathbb{P}[P_\alpha > x] = x^{-\alpha}$ for $x \geq 1$,

$$\mathcal{L}(R/r, X/R \mid R > r) \rightsquigarrow \mathcal{L}(P_\alpha, G), \quad r \rightarrow \infty.$$

Assumption 2 strengthens Assumption 1 by adding regular radial scaling. By the continuous mapping theorem, $\mathcal{L}(X/r \mid R > r) \rightsquigarrow \mathcal{L}(P_\alpha G)$, so the normalized rank- p AGCA tail-simulation target is $P_\alpha \tilde{G}_{\mu,p}$. Multiplying by the threshold gives $r P_\alpha \tilde{G}_{\mu,p}$, the corresponding simulator on the original Pareto scale.

In applications, however, the goal is often not to approximate the whole conditional law of extremes for its own sake, but to approximate tail functionals computed from that law. Examples include portfolio exceedance probabilities, value-at-risk, expected shortfall when the relevant tail moment exists, and stress scores such as the maximum loss over a collection of positions. The next two results cover complementary classes of such summaries. The first gives a uniform transport-type bound for bounded Lipschitz functionals of the simulated extreme vector. For probability laws on $\mathbb{R}^d$, write the bounded-Lipschitz distance as

$$d_{\text{BL}}(\mathcal{L}(A), \mathcal{L}(B)) := \sup_{\|f\|_\infty \leq 1, \text{Lip}(f) \leq 1} |\mathbb{E}[f(A)] - \mathbb{E}[f(B)]|.$$

Proposition 4 (Tail simulation under regular variation). *Suppose Assumption 2 holds, and let P_α be independent of G with $\mathbb{P}[P_\alpha > x] = x^{-\alpha}$, $x \geq 1$. Put*

$$\varepsilon_r := d_{\text{BL}}(\mathcal{L}(X/r \mid R > r), \mathcal{L}(P_\alpha G)). \quad (2.15)$$

Then $\varepsilon_r \rightarrow 0$. Moreover, for every $0 < \beta \leq 1$ with $\beta < \alpha$,

$$d_{\text{BL}}(\mathcal{L}(X/r \mid R > r), \mathcal{L}(P_\alpha \tilde{G}_{\mu,p})) \leq \varepsilon_r + 2^{1-\beta/2} \frac{\alpha}{\alpha - \beta} \rho_{\mu,p}^{\beta/2}.$$

The proposition gives a uniform bound over bounded Lipschitz test functionals. This is the

sense in which the low-rank AGCA simulator is controlled by the sum of two errors: the threshold error from replacing finite exceedances by the regular-variation limit, and the angular approximation error $\rho_{\mu,p}$. When $\alpha > 1$, one may take $\beta = 1$ and obtain the familiar $\sqrt{\rho_{\mu,p}}$ rate for the angular term. With exact standard Pareto marginal scaling, the radial index is $\alpha = 1$, so the statement uses any $\beta < 1$.

The reconstruction $\tilde{G}_{\mu,p}$ is an unconstrained spherical projection. Hence $M_\mu(W_{\mu,p}) \subset \mathcal{H}_\mu$ need not lie in $\mathbb{S}_+^{d-1}$, and simulated vectors $P_\alpha \tilde{G}_{\mu,p}$ need not remain in the positive cone. For positive-cone simulation, one should verify positivity on the relevant support, impose a constrained angular reconstruction, or model residual variation around the unconstrained AGCA fit. Supplementary Section S2.5 develops a positive post-projection that enforces $\mathbb{S}_+^{d-1}$ -valued simulated directions while adding an explicit correction term to the simulation bounds.

Many risk summaries are not bounded. For positively homogeneous scores, the Pareto radial excess can instead be separated from the angular profile. Let $h : \mathbb{R}^d \rightarrow [0, \infty)$ satisfy $h(cz) = ch(z)$ for $c \geq 0$. Suppose that, on $\mathbb{S}^{d-1}$, for constants $0 < H_h < \infty$ and $L_h < \infty$,

$$0 \leq h(s) \leq H_h, \quad |h(s) - h(t)| \leq L_h \|s - t\|_2.$$

Set $C_h := \mathbb{E}[h(G)^\alpha]$, $\tilde{C}_{h,p} := \mathbb{E}[h(\tilde{G}_{\mu,p})^\alpha]$, $\gamma := \min(\alpha, 1)$, and define

$$\eta_{r,h} := \sup_{x \geq H_h} \left| \frac{\mathbb{P}[h(X) > rx]}{\mathbb{P}[R > r]} - x^{-\alpha} C_h \right|, \quad K_{\alpha,h} := \begin{cases} L_h^\alpha, & 0 < \alpha \leq 1, \\ \alpha H_h^{\alpha-1} L_h, & \alpha > 1. \end{cases}$$

The next proposition is the homogeneous-score analogue of the bounded-Lipschitz bound: the finite-threshold error is inherited from Assumption 2, and the additional error comes from the AGCA angular approximation.

Proposition 5 (Homogeneous tail scores). *Suppose Assumption 2 holds. Then $\eta_{r,h} \rightarrow 0$ as $r \rightarrow \infty$.*

Moreover, for every $x \geq H_h$,

$$\mathbb{P}[h(P_\alpha G) > x] = x^{-\alpha} C_h, \quad \mathbb{P}[h(P_\alpha \tilde{G}_{\mu,p}) > x] = x^{-\alpha} \tilde{C}_{h,p}.$$

The angular constants satisfy $|C_h - \tilde{C}_{h,p}| \leq K_{\alpha,h}(2\rho_{\mu,p})^{\gamma/2}$. Consequently,

$$\sup_{x \geq H_h} \left| \frac{\mathbb{P}[h(X) > rx]}{\mathbb{P}[R > r]} - \mathbb{P}[h(P_\alpha \tilde{G}_{\mu,p}) > x] \right| \leq \eta_{r,h} + H_h^{-\alpha} K_{\alpha,h}(2\rho_{\mu,p})^{\gamma/2}.$$

Thus the two simulation bounds have the same structure: a first-order regular-variation error, ε_r in (2.15) or $\eta_{r,h}$, plus an AGCA angular approximation error. Assumption 2 ensures that the threshold terms vanish, but does not quantify their rates. Rates for either term would require second-order regular variation or another quantitative tail approximation condition.

The preceding bounds do not imply that the AGCA simulator can approximate $\mathbb{P}[P_\alpha G \in A]$ uniformly over all Borel sets A . Such a statement would require total-variation control of the law of $P_\alpha G$ by the law of $P_\alpha \tilde{G}_{\mu,p}$. Low-rank reconstruction changes the support of the angular law: $P_\alpha \tilde{G}_{\mu,p}$ is supported on the cone generated by $M_\mu(W_{\mu,p})$, whereas $P_\alpha G$ need not be.

Proposition 6 (Total-variation obstruction). *For P_α as above, let*

$$\mathcal{C}_{\mu,p} := \{z \in \mathbb{R}^d : \|z\|_2 \geq 1, z / \|z\|_2 \in M_\mu(W_{\mu,p})\}.$$

Then $\mathbb{P}[P_\alpha \tilde{G}_{\mu,p} \in \mathcal{C}_{\mu,p}] = 1$ and $\mathbb{P}[P_\alpha G \in \mathcal{C}_{\mu,p}] = \mathbb{P}[G \in M_\mu(W_{\mu,p})]$. Consequently,

$$\sup_A \left| \mathbb{P}[P_\alpha G \in A] - \mathbb{P}[P_\alpha \tilde{G}_{\mu,p} \in A] \right| \geq 1 - \mathbb{P}[G \in M_\mu(W_{\mu,p})],$$

where the supremum is over Borel $A \subset \mathbb{R}^d$.

If the angular law is full-dimensional and $p < d - 1$, the probability $\mathbb{P}[G \in M_\mu(W_{\mu,p})]$ is typically zero, so the total-variation discrepancy is maximal even when $\rho_{\mu,p}$ is small. This is not a defect of the projection formula; it is the usual support mismatch created by any exact low-

dimensional representation. Total-variation closeness is available only when the angular law is itself supported on the fitted model, or after adding an explicit residual/noise model around $M_\mu(W_{\mu,p})$. This low-dimensional support mismatch is separate from the positivity issue discussed above.

2.6 Portfolio functionals for tail simulation

We make the preceding bounds operational through portfolio scores. Let $z \in \mathbb{R}^d$ denote a vector of standardized portfolio losses and let $w \in \mathbb{R}^d$ collect portfolio positions. The portfolio loss score is $\pi_w(z) := w^\top z$, and we impose the budget constraint $\mathbf{1}^\top w = 1$. In applications below, the leverage constraint $\|w\|_1 \leq L$, for a positive constant L , gives uniform control over a portfolio class. We highlight two summaries that are useful for portfolio comparison and risk control. The first is a bounded score for comparing candidate portfolios under simulated tail scenarios. For $t \geq 0$, define the capped portfolio excess

$$f_{w,t,L}(z) := \min \{1, L^{-1}(\pi_w(z) - t)_+\}. \quad (2.16)$$

It records excess over a portfolio stress level, but caps the contribution of very large radial draws. Since $\|w\|_2 \leq \|w\|_1 \leq L$, it satisfies $\|f_{w,t,L}\|_\infty \leq 1$ and $\text{Lip}(f_{w,t,L}) \leq 1$, and is therefore covered directly by Proposition 4.

Corollary 2 (Capped portfolio excess). *Suppose Assumption 2 holds. Let $w \in \mathbb{R}^d$, $\mathbf{1}^\top w = 1$, $\|w\|_1 \leq L$ for some $L \geq 1$, $t \geq 0$, and let ε_r be as in (2.15). Then, for every $0 < \beta \leq 1$ with $\beta < \alpha$,*

$$\left| \mathbb{E}[f_{w,t,L}(X/r) \mid R > r] - \mathbb{E}[f_{w,t,L}(P_\alpha \tilde{G}_{\mu,p})] \right| \leq \varepsilon_r + 2^{1-\beta/2} \frac{\alpha}{\alpha - \beta} \rho_{\mu,p}^{\beta/2}.$$

The second summary is value-at-risk. For a scalar portfolio loss score Y , the upper-tail value-at-risk at tail probability u , $\text{VaR}_u(Y) := \inf\{q \in \mathbb{R} : \mathbb{P}[Y > q] \leq u\}$, is the high quantile exceeded with probability at most u . In the Pareto limit, VaR for the score π_w is driven by the unbounded

homogeneous score $(w^\top z)_+$. Define the true and reconstructed angular tail constants

$$C_w := \mathbb{E}[(w^\top G)_+^\alpha], \quad \tilde{C}_{w,p} := \mathbb{E}[(w^\top \tilde{G}_{\mu,p})_+^\alpha]. \quad (2.17)$$

For tail probability u , define the normalized Pareto-limit VaR values for the true and AGCA-reconstructed portfolio tails

$$q_w(u) := (C_w/u)^{1/\alpha}, \quad \tilde{q}_{w,p}(u) := (\tilde{C}_{w,p}/u)^{1/\alpha}. \quad (2.18)$$

Multiplying them by a high threshold r gives approximate VaR values on the original Pareto scale.

Corollary 3 (Portfolio tail constants and VaR). *Suppose Assumption 2 holds. Let $\pi_w(z) := w^\top z$, where $w \in \mathbb{R}^d$, $\mathbf{1}^\top w = 1$, and $\|w\|_1 \leq L$ for some $L \geq 1$. Then, for every $x \geq L$,*

$$\mathbb{P}[\pi_w(P_\alpha G) > x] = x^{-\alpha} C_w, \quad \mathbb{P}[\pi_w(P_\alpha \tilde{G}_{\mu,p}) > x] = x^{-\alpha} \tilde{C}_{w,p}.$$

For $0 < u < L^{-\alpha} \min(C_w, \tilde{C}_{w,p})$, the normalized VaR ratio satisfies $q_w(u)/\tilde{q}_{w,p}(u) = (C_w/\tilde{C}_{w,p})^{1/\alpha}$. If $\min(C_w, \tilde{C}_{w,p}) \geq c > 0$, then

$$|\log q_w(u) - \log \tilde{q}_{w,p}(u)| \leq \frac{1}{\alpha c} (2\rho_{\mu,p})^{\min(\alpha,1)/2} \begin{cases} L^\alpha, & 0 < \alpha \leq 1, \\ \alpha L^\alpha, & \alpha > 1. \end{cases}$$

Thus the capped excess gives a bounded simulation diagnostic, while VaR gives a scale-normalized tail risk summary. A related important measure is expected shortfall, $\text{ES}_u(Y) := \mathbb{E}[Y \mid Y > \text{VaR}_u(Y)]$, the conditional mean beyond the corresponding high quantile. For the Pareto limit considered here, this mean is finite only when $\alpha > 1$. In that case, the limiting expected shortfall equals $\alpha/(\alpha - 1)$ times the corresponding VaR, so the same relative angular error bound applies. For exact standard Pareto marginal scaling, $\alpha = 1$, the limiting expected shortfall is infinite, whereas VaR remains finite.

3 AGCA: estimation and inference

In data analysis we observe sample vectors $Y_i = (Y_{i1}, \dots, Y_{id})$, standardize the margins, select observations with large radial size, and apply the population eigensolution of Section 2 to the selected angular directions. The tail sample size k plays the usual peaks-over-threshold role: it balances extremeness against estimation variance. In asymptotic statements we use an intermediate sequence k_n , with $k_n \rightarrow \infty$ and $k_n/n \rightarrow 0$. We use hats with the superscript `orc` to denote empirical quantities computed from the true Pareto-standardized margins, while hats without the oracle superscript denote empirical quantities computed from rank-Pareto margins. The section separates the deterministic eigensolution, consistency under top- k selection and rank marginal standardization, oracle distributional inference, and the remaining first-order effect of estimating the margins.

3.1 Empirical construction and eigensolution

For a fixed rank p , anchor μ , and tail sample size k , we use two parallel sample constructions. For the oracle theory, let $X_1, \dots, X_n$ be independent copies of the Pareto-standardized vector, set $R_i := \|X_i\|_2$, $G_i := X_i/R_i$, and $U_{\mu,i} := u_\mu(G_i)$, and let $\mathcal{I}_n^{(k)}$ be the indices of the k largest values R_i . With unknown margins, a standard rank-Pareto transform is

$$\hat{X}_{ij} := \frac{n+1}{n+1 - \text{rank}(Y_{ij})}, \quad i = 1, \dots, n, \quad j = 1, \dots, d, \quad (3.1)$$

with ranks computed within each component. This is the standard empirical EVT device for removing marginal tail scales before estimating extremal dependence. Set $\hat{R}_i := \|\hat{X}_i\|_2$, $\hat{G}_i := \hat{X}_i/\hat{R}_i$, and $\hat{U}_{\mu,i} := u_\mu(\hat{G}_i)$, and let $\hat{\mathcal{I}}_n^{(k)}$ be the indices of the k largest values $\hat{R}_i$. In both constructions, radii select tail observations and directions are the angular profiles summarized by AGCA. The corresponding anchored second-moment matrices are

$$\hat{\Sigma}_{\mu,n}^{(k),\text{orc}} := \frac{1}{k} \sum_{i \in \mathcal{I}_n^{(k)}} U_{\mu,i} U_{\mu,i}^\top, \quad \hat{\Sigma}_{\mu,n}^{(k),\text{emp}} := \frac{1}{k} \sum_{i \in \hat{\mathcal{I}}_n^{(k)}} \hat{U}_{\mu,i} \hat{U}_{\mu,i}^\top. \quad (3.2)$$

Their eigenvectors represent departures from the fixed benchmark μ . The rank-Pareto eigensolution diagonalizes $\widehat{\Sigma}_{\mu,n}^{(k),\text{emp}}$ in the tangent hyperplane $\mu^\perp$. Explicitly, let $Q_\mu \in \mathbb{R}^{d \times (d-1)}$ have orthonormal columns spanning $\mu^\perp$, so that $Q_\mu^\top Q_\mu = I_{d-1}$ and $Q_\mu Q_\mu^\top = I - \mu\mu^\top$. Diagonalize the ordinary $(d-1) \times (d-1)$ coordinate matrix

$$Q_\mu^\top \widehat{\Sigma}_{\mu,n}^{(k),\text{emp}} Q_\mu = H \widehat{\Lambda} H^\top, \quad \widehat{\Lambda} = \text{diag}(\widehat{\lambda}_1, \dots, \widehat{\lambda}_{d-1}),$$

with $\widehat{\lambda}_1 \geq \dots \geq \widehat{\lambda}_{d-1} \geq 0$. If h_ℓ is the ℓ th column of H , the corresponding AGCA loading in the original coordinates is $\widehat{b}_\ell := Q_\mu h_\ell \in \mu^\perp$. For later use, write $\widehat{P}_{\mu,p} := \sum_{\ell=1}^p \widehat{b}_\ell \widehat{b}_\ell^\top$, with $\widehat{P}_{\mu,0} := 0$, for the empirical rank- p spectral projector. The empirical scores are the plug-in analogues of (2.11), $\widehat{\alpha}_{i\ell} := \langle \widehat{b}_\ell, \widehat{U}_{\mu,i} \rangle$, and the plug-in version of (2.13), equivalently (2.5), gives the rank- p reconstructions. The choice of orthonormal basis Q_μ only changes the coordinates of the calculation; the resulting subspaces in $\mu^\perp$ are the same, up to the usual sign and eigenspace nonuniqueness.

The next statement treats the selected directions as fixed, so the corresponding second moment and risk are written without hats.

Proposition 7 (Finite-sample eigensolution for selected directions). *Let $\mathcal{I} \subset \mathbb{N}$ be a nonempty finite index set and let $\{g_i : i \in \mathcal{I}\} \subset \mathbb{S}_+^{d-1}$ be selected angular directions. Define*

$$\Sigma_{\mu,\mathcal{I}} := \frac{1}{|\mathcal{I}|} \sum_{i \in \mathcal{I}} u_\mu(g_i) u_\mu(g_i)^\top, \quad \mathcal{R}_{\mu,\mathcal{I}}(W) := \frac{1}{|\mathcal{I}|} \sum_{i \in \mathcal{I}} \sin^2 d_g(g_i, M_\mu(W)), \quad W \subset \mu^\perp.$$

Then, for every linear subspace $W \subset \mu^\perp$, $\mathcal{R}_{\mu,\mathcal{I}}(W) = \text{tr}((I - \Pi_W) \Sigma_{\mu,\mathcal{I}})$. Consequently, if $b_{\mathcal{I},1}, \dots, b_{\mathcal{I},d-1}$ are orthonormal eigenvectors of $\Sigma_{\mu,\mathcal{I}}$ in $\mu^\perp$, with eigenvalues $\lambda_{\mathcal{I},1} \geq \dots \geq \lambda_{\mathcal{I},d-1}$, then $W_{\mu,\mathcal{I},p} := \text{span}(b_{\mathcal{I},1}, \dots, b_{\mathcal{I},p})$ minimizes $\mathcal{R}_{\mu,\mathcal{I}}(W)$ over all p -dimensional $W \subset \mu^\perp$, and the corresponding rank- p residual risk is $\mathcal{R}_{\mu,\mathcal{I}}(W_{\mu,\mathcal{I},p}) = \sum_{j=p+1}^{d-1} \lambda_{\mathcal{I},j}$.

The selected set $\mathcal{I}$ in Proposition 7 can be the top- k radial set or a rank-margin version of that set. The proposition is deterministic: conditional on those directions, empirical AGCA is the leading eigenspace of the second-moment operator formed from the selected directions, with only the usual

eigenspace nonuniqueness under ties.

3.2 Top- k consistency and marginal standardization

The following theorem gives the oracle top- k justification for $\widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}}$ in (3.2). The next theorem gives the corresponding primitive rank-Pareto consistency result.

Theorem 2 (Consistency under empirical intermediate thresholds). *Assume that R_1 has a continuous distribution and that Assumption 1 holds. Let $k_n \rightarrow \infty$ and $k_n/n \rightarrow 0$. Then $\left\| \widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} - \Sigma_\mu \right\|_{\text{op}} \rightarrow 0$ in probability. Consequently, the eigenvalues, anchored residual risks, and, under an eigengap, the rank- p AGCA projector obtained from the top k_n radial observations are consistent for their population counterparts.*

The main tail condition in Theorem 2 is Assumption 1: conditional on being far enough in the tail, the Euclidean spherical direction must approach a limiting angular law. This is a standard condition that underlies empirical angular-measure estimation in multivariate EVT. The continuity assumption on R_1 ensures that the top- k_n radial set is unambiguously defined: with probability one, no positive number of observations has the same radius as the empirical threshold. It is mild for continuous measurements and for continuous Pareto-standardized variables, and it does not exclude discrete angular regimes such as mass on coordinate axes.² The intermediate sequence k_n only requires increasingly many selected extremes with vanishing tail fraction k_n/n ; finite-sample choice remains a threshold-selection problem. In applications, this is normally handled through threshold paths and stability diagnostics. For AGCA the natural diagnostics are eigenvalues, anchored variation explained, residual risk, loadings, scores, and projector distances across a range of k .

Replacing the unknown marginal transforms by ranks affects the estimator in two ways: the direction of a fixed observation changes from G_i to $\widehat{G}_i$, and the top- k_n set based on $\widehat{R}_i$ need not equal

²The assumption could be relaxed by specifying a deterministic or randomized convention for observations with equal radii, or by replacing exact top- k_n selection with a threshold exceedance set and assuming that the number of observations at the boundary is $o_p(k_n)$. Such extensions are useful for heavily rounded, count-valued, or otherwise coarse data.

the oracle top- k_n set. The next theorem shows that both effects are negligible at the consistency scale under primitive tail conditions. Rank-based estimation of limiting angular (spectral) measures goes back to [Einmahl et al. \(1997, 2001\)](#); see also [Einmahl and Segers \(2009\)](#) and, for the multivariate rank-tail empirical process, [Einmahl et al. \(2012\)](#).

Theorem 3 (Rank-based consistency). *Assume that the margins $F_1, \dots, F_d$ and R_1 are continuous, and that Assumption 2 holds. Let $k_n \rightarrow \infty$ and $k_n/n \rightarrow 0$. Then $\left\| \widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \Sigma_\mu \right\|_{\text{op}} \rightarrow 0$ in probability. In particular, the eigenvalues, anchored residual risks, anchored variation explained when $\tau_\mu > 0$, and, under an eigengap, the rank- p AGCA projector obtained from the rank-Pareto top- k_n observations are consistent for their population counterparts.*

The proof, in Supplementary Section [S2.7](#), shows that the average rank-induced angular error over the relevant selected observations is $o_p(1)$, and that the symmetric difference between the oracle and rank-based selected sets is $o_p(k_n)$. The latter step is why Theorem 3 uses Assumption 2: angular convergence alone does not control how many radii lie in a shrinking relative band around the top- k_n threshold. The average form of the angular control is essential, because the rank transform distorts the top few order statistics of each margin at constant order.

3.3 Oracle distributional inference and stability diagnostics

The consistency result above is enough to justify the AGCA target, but it does not quantify sampling variation. For the oracle estimator, distributional inference is particularly simple because the anchored departure $u_\mu(G)$ is bounded. The only additional condition needed here is a second-order version of the angular limit at the k_n/n tail level.

With the oracle notation above, define

$$\bar{F}_R(r) := \mathbb{P}[R_1 > r], \quad H_\mu(r) := \mathbb{E}[u_\mu(G_1)u_\mu(G_1)^\top \mid R_1 > r],$$

and the angular-bias envelope

$$\eta_\mu(t) := \sup \left\{ \|H_\mu(r) - \Sigma_\mu\|_{\text{op}} : \bar{F}_R(r) \leq t \right\}, \quad 0 < t < 1.$$

Assumption 3 (Second-order angular bias). *As $n \rightarrow \infty$, $\sqrt{k_n} \eta_\mu(2k_n/n) \rightarrow 0$ for the intermediate sequence $k_n \rightarrow \infty$ with $k_n/n \rightarrow 0$.*

Assumption 3 assumes away the asymptotic bias. Such a condition is often assumed in classical extreme value statistics; see, e.g. (De Haan and Ferreira, 2006, Chapter 3). It places an upper bound for the intermediate sequence k_n .

Theorem 4 (Oracle CLT for top- k AGCA). *Assume the conditions of Theorem 2 and Assumption 3. Then*

$$\sqrt{k_n} \left(\widehat{\Sigma}_{\mu,n}^{(k_n), \text{orc}} - \Sigma_\mu \right) \rightsquigarrow Z,$$

where Z is a centered symmetric Gaussian matrix supported on $\mu^\perp \times \mu^\perp$. Moreover, for all symmetric $A, B \in \mathbb{R}^{d \times d}$, $\text{Cov}[\text{tr}(AZ), \text{tr}(BZ)] = \text{Cov}[U_\mu^\top A U_\mu, U_\mu^\top B U_\mu]$. Equivalently, given $U_{\mu,1}, \dots, U_{\mu,m}$ independent copies of U_μ , Z has the same law as the weak limit of

$$m^{-1/2} \sum_{\ell=1}^m (U_{\mu,\ell} U_{\mu,\ell}^\top - \Sigma_\mu).$$

Theorem 4 says that top- k_n selection is first-order free for the bounded anchored second moment: the random radial threshold has the same first-order covariance effect as the empirical tail count and cancels from the final covariance. This is the main inferential simplification created by using the bounded spherical departure $u_\mu(G)$. If the second-order bias is not negligible and $\sqrt{k_n}(H_\mu(r_n) - \Sigma_\mu)$ converges along the relevant deterministic threshold sequence, the same CLT has the corresponding deterministic bias shift.

Using the population projector and eigengap in (2.8), let $\widehat{\lambda}_j^{\text{orc}}$ and $\widehat{b}_j^{\text{orc}}$ denote the ordered

eigenvalues and associated eigenvectors of $\widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}}$, and define the corresponding oracle estimators

$$\widehat{\tau}_{\mu}^{\text{orc}} := \text{tr}(\widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}}), \quad \widehat{\text{AVE}}_{\mu,p}^{\text{orc}} := \frac{\sum_{j=1}^p \widehat{\lambda}_j^{\text{orc}}}{\widehat{\tau}_{\mu}^{\text{orc}}}, \quad \widehat{P}_{\mu,p}^{\text{orc}} := \sum_{j=1}^p \widehat{b}_j^{\text{orc}} (\widehat{b}_j^{\text{orc}})^{\top}.$$

Corollary 4 (Oracle scalar and spectral inference). *Under the assumptions of Theorem 4, the following limits hold.*

1. $\sqrt{k_n} (\widehat{\tau}_{\mu}^{\text{orc}} - \tau_{\mu}) \rightsquigarrow N(0, \text{Var}[\|U_{\mu}\|_2^2]).$

2. *If λ_j is a simple eigenvalue, then*

$$\sqrt{k_n} (\widehat{\lambda}_j^{\text{orc}} - \lambda_j) \rightsquigarrow N(0, \text{Var}[(b_j^{\top} U_{\mu})^2]).$$

The corresponding limits are jointly Gaussian over any fixed set of simple eigenvalues, with covariances $\text{Cov}[(b_j^{\top} U_{\mu})^2, (b_{\ell}^{\top} U_{\mu})^2]$.

3. *If $1 \leq p \leq d-2$, $\tau_{\mu} > 0$, and $\Delta_{\mu,p} > 0$, then*

$$\sqrt{k_n} (\widehat{\text{AVE}}_{\mu,p}^{\text{orc}} - \text{AVE}_{\mu,p}) \rightsquigarrow N(0, \sigma_{\mu,p}^2),$$

where $\sigma_{\mu,p}^2 := \text{Var}[\psi_{\mu,p}(U_{\mu})]$ and $\psi_{\mu,p}(u) := \tau_{\mu}^{-1} (u^{\top} P_{\mu,p} u - \text{AVE}_{\mu,p} \|u\|_2^2)$.

4. *If $1 \leq p \leq d-2$ and $\Delta_{\mu,p} > 0$, then*

$$\sqrt{k_n} (\widehat{P}_{\mu,p}^{\text{orc}} - P_{\mu,p}) \rightsquigarrow \mathcal{D}_{\mu,p}(Z), \quad \mathcal{D}_{\mu,p}(E) := \sum_{j=1}^p \sum_{m=p+1}^{d-1} \frac{b_m b_m^{\top} E b_j b_j^{\top} + b_j b_j^{\top} E b_m b_m^{\top}}{\lambda_j - \lambda_m}, \quad E = E^{\top}.$$

The eigengap condition is an identifiability condition: if $\lambda_p = \lambda_{p+1}$, the rank- p space is not unique, and small gaps allow large rotations of individual eigenvectors. Thus unstable loading vectors need not imply an unstable low-dimensional fitted subspace. For this reason, comparisons across thresholds, bootstrap samples, or anchors use spectral projectors whenever individual component orientation is not the target.

Corollary 5 (Plug-in variance for oracle explained variation). *Under the assumptions of Corollary 4, with $1 \leq p \leq d - 2$, $\tau_\mu > 0$, and $\Delta_{\mu,p} > 0$, define*

$$\hat{\psi}_{\mu,p,i}^{\text{orc}} := (\hat{\tau}_\mu^{\text{orc}})^{-1} \left[U_{\mu,i}^\top \hat{P}_{\mu,p}^{\text{orc}} U_{\mu,i} - \widehat{\text{AVE}}_{\mu,p}^{\text{orc}} \|U_{\mu,i}\|_2^2 \right]$$

and let $\hat{\sigma}_{\mu,p}^{2,\text{orc}}$ be the empirical variance of $\hat{\psi}_{\mu,p,i}^{\text{orc}}$ over $i \in \mathcal{I}_n^{(k_n)}$. Then $\hat{\sigma}_{\mu,p}^{2,\text{orc}} \rightarrow \sigma_{\mu,p}^2$ in probability.

Consequently,

$$\widehat{\text{AVE}}_{\mu,p}^{\text{orc}} \pm z_{1-\gamma/2} \frac{\hat{\sigma}_{\mu,p}^{\text{orc}}}{\sqrt{k_n}}$$

is an asymptotically valid oracle $(1 - \gamma)$ confidence interval for $\text{AVE}_{\mu,p}$, with z_q the q -quantile of $N(0, 1)$. The same plug-in principle estimates the asymptotic variances for $\hat{\tau}_\mu^{\text{orc}}$ and simple oracle eigenvalues.

3.4 Rank standardization beyond consistency

The preceding CLT applies to the oracle sample. The proof of Theorem 3 shows that rank-Pareto marginal standardization is negligible at the consistency scale, but this $o_p(1)$ control does not imply distributional equivalence at the $k_n^{-1/2}$ scale. With $h_\mu(g) := u_\mu(g)u_\mu(g)^\top$,

$$\sqrt{k_n} \left(\hat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \hat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} \right) = \frac{1}{\sqrt{k_n}} \sum_{i \in \mathcal{I}_n^{(k_n)}} \{h_\mu(\hat{G}_i) - h_\mu(G_i)\} + \frac{1}{\sqrt{k_n}} \left(\sum_{i \in \hat{\mathcal{I}}_n^{(k_n)}} - \sum_{i \in \mathcal{I}_n^{(k_n)}} \right) h_\mu(\hat{G}_i).$$

The first term is the angular perturbation induced by estimating the marginal Pareto scales; the second is the selection perturbation from recomputing the top- k_n set after rank standardization. Weighted empirical-tail approximations indicate that both terms can enter at the same $k_n^{-1/2}$ scale as the oracle fluctuation. Thus the oracle interval in Corollary 5 is a formal interval for the oracle estimator, not automatically for the rank-Pareto estimator.

A full rank-based CLT would add a Gaussian marginal-estimation correction to Theorem 4, under Euclidean-polar regular variation (Assumption 2), second-order marginal and joint-tail conditions, and differentiability of the limiting tail-measure functional under componentwise marginal scalings.

Supplementary Section S2.9 records the corresponding conditional delta-method theorem, but we do not derive primitive rank-tail empirical-process conditions here. Supplementary Section S3.8 illustrates the issue: oracle intervals are reasonably calibrated with oracle margins, whereas applying the same formula after rank-Pareto standardization is not uniformly reliable. The rank-standardized simulations and empirical analyses below therefore use bootstrap resampling, threshold paths, anchor checks, and projector distances as finite-sample stability diagnostics.

3.5 Plug-in reconstruction for tail simulation

The population simulation results in Section 2.5 use the rank- p component space through its orthogonal projector. For an angular direction $g \in \mathbb{S}_+^{d-1}$, the plug-in AGCA reconstruction uses the empirical projector $\hat{P}_{\mu,p}$ and is

$$\hat{g}_{\mu,p} := \frac{a_\mu(g)\mu + \hat{P}_{\mu,p}u_\mu(g)}{\sqrt{a_\mu(g)^2 + \left\| \hat{P}_{\mu,p}u_\mu(g) \right\|_2^2}}.$$

Applied to the selected empirical directions $\hat{G}_i$, this gives reconstructed angular profiles that can be combined with a fitted radial tail model to simulate extremes. For example, one draws an independent Pareto radial multiplier and attaches it to $\hat{G}_{\mu,p,i}$, or to a positive post-projected version if positivity is enforced as in Supplementary Section S2.5.

This plug-in step is controlled by the preceding consistency and oracle inference results. Theorem 2 gives consistency of the anchored second-moment matrix in the oracle case, and Theorem 3 gives the corresponding consistency when margins are rank-standardized. These results also give consistency of the eigenvalue residual-risk curve. Under an eigengap, the leading spectral projector is a continuous function of that matrix, so $\hat{P}_{\mu,p}$ is consistent; in the oracle case, Corollary 4 also gives its first-order distribution. The reconstruction map above is continuous in the projector on the anchored hemisphere. Remaining errors, such as radial tail-index estimation, threshold selection, and Monte Carlo error, belong to the radial EVT and simulation layers rather than to the AGCA

eigensolution.

4 Empirical study: portfolio-loss extremes

The empirical study examines whether daily equity-portfolio loss extremes have a low-dimensional angular structure and whether that structure is useful for portfolio tail summaries. The object is the angular law of tail directions obtained after marginal rank-Pareto standardization. Each extreme day is represented by a radius and a direction on the positive sphere; AGCA is then fit to the largest radii. A low-rank fit means that, conditional on a large joint portfolio-loss event, the cross-sectional pattern of standardized losses is well approximated by a small number of anchored geodesic coordinates. Supplementary Section S4 gives data-construction details and secondary diagnostics.

4.1 Data and tail construction

The data set consists of daily value-weighted Fama–French 2×3 portfolios (French, 2026). We use four size-based bivariate sorts: Size–Book-to-Market, Size–Operating Profitability, Size–Investment, and Size–Momentum. Each sort contributes six portfolios, giving $d = 24$ daily return series. We interpret losses as negative returns, transform each loss margin to the pseudo-Pareto scale, and form Euclidean radii and directions. The main AGCA fit uses the largest 5% radii from the complete daily panel beginning on July 2, 1973. A modern anomaly-sorted portfolio panel from Open Source Asset Pricing (Chen and Zimmermann, 2026) is used as a robustness check and is reported in Supplementary Section S4.1.

The analysis uses the canonical anchor $\mu_0 = d^{-1/2}\mathbf{1}$, the balanced complete-dependence direction. Loadings are therefore departures from a day on which all standardized portfolio losses participate equally in the extreme direction. This fixed anchor is used for the main interpretation; the Supplementary Material contains data-adaptive anchor checks. Whenever shaded bootstrap 95% diagnostic intervals (DIs) are shown below, they are conditional stability summaries: after the rank-Pareto transform and the top-5% radial sample are fixed, we resample the selected directions




Figure 3: Fama–French empirical summaries. Left: cumulative anchored variation explained at the main 5% radial threshold. Right: average relative errors for capped excess and normalized VaR over the portfolio evaluation class. Shaded bands are bootstrap 95% diagnostic intervals, computed as pointwise 2.5% and 97.5% conditional bootstrap percentiles.

with replacement, refit AGCA, and recompute the displayed summaries. These DIs do not include uncertainty from marginal rank estimation, threshold selection, or serial dependence in daily losses; they are used to assess stability of the fitted angular cloud, while threshold and anchor sensitivity are checked separately.

4.2 Effective dimension and portfolio tail summaries

The portfolio-functional check follows Section 2.6. We use the capped portfolio excess in (2.16), evaluated as $f_{w,2||w||_2,||w||_2}$, and the normalized Pareto-limit VaR summary in Corollary 3. With rank-Pareto margins the empirical Pareto index is $\alpha = 1$, so the VaR comparison is reported as the relative normalized VaR error $\exp\{|\log \tilde{C}_{w,p} - \log C_w|\} - 1$. We evaluate both summaries on the equal-weight portfolio, all block-equal portfolios, 250 random long-only portfolios, and 250 random limited-leverage portfolios with $||w||_1 \leq 1.5$. The rank-level curves average over this fixed evaluation set; Supplementary Figure S20 reports the class-level breakdown.

Figure 3 shows that the Fama–French portfolio tails have a concentrated anchored spectrum. The first three AGCs explain 64.6% of anchored variation, the first five explain 76.3%, the first eight explain 86.4%, and the first ten explain 91.1%. The same low-rank structure preserves the portfolio summaries in financially interpretable units: rank five gives mean relative error 3.4%




Figure 4: Canonical-anchor AGCA loadings for the Fama–French panel. Columns are portfolios, grouped by bivariate sort, and rows are the first five AGCA loading vectors. Red and blue denote opposite signed departures from the balanced complete-dependence anchor. Loading signs are oriented deterministically so that the largest absolute entry in each loading vector is positive.

for both capped excess and normalized VaR, rank eight reduces this to 1.9%, rank ten to 1.25%, and rank twelve to 0.84%. A 1.25% normalized VaR error means that the AGCA approximation changes the Pareto-limit VaR level by about 1.25%. The bootstrap DIs are narrow; for rank ten, the pointwise DI is approximately 1.14%–1.30%. Supplementary Figures S15 and S17 show that the effective-dimension conclusion is stable under data-adaptive anchor choices and over tail fractions from 0.5% to 10%.

4.3 Loading structure

Figure 4 gives the main loading display. The horizontal axis is the portfolio universe and the rows are AGCA coordinates. This orientation emphasizes that a loading is a cross-sectional contrast: it identifies which sorted portfolios are above or below the balanced extreme direction when the corresponding AGCA score is positive. The first few loadings are not single-asset spikes. They are structured contrasts across the four economically related 2×3 blocks, so the low-rank approximation is not simply discarding all but a few portfolios. Rather, AGCA is capturing repeated cross-sectional patterns in the directions of joint portfolio losses. In AGC1, the main pattern is a size contrast repeated across the bivariate sorts: the small-size portfolios tend to load negatively, while the large-size portfolios tend to load positively. A positive AGC1 score therefore corresponds to an

extreme loss direction tilted toward relatively larger standardized losses among the large-stock portfolios, after centering at balanced complete dependence.

The sign of a loading is arbitrary up to orientation. The canonical anchor fixes the interpretation of the coordinate origin, while the deterministic sign convention fixes display orientation. Data-driven anchors give similar scalar dimension diagnostics, but their loading vectors are less directly interpretable because the coordinate origin is then a sample-dependent tail direction rather than the balanced complete-dependence direction. Consequently, the empirical content is in the blockwise relative signs and magnitudes. Supplementary Figure S19 adds pointwise conditional bootstrap DIs for AGC1; most displayed signs are stable in this diagnostic sense, with the least stable entries concentrated near zero.

Because the unconstrained anchored subsphere may leave the positive orthant, the Supplementary Material also reports the positive post-projection. In this application its effect on the scalar functionals is small: the raw and post-projected curves are close, and the post-projection does not change the rank-level conclusion. Supplementary Section S4.1 reports the anomaly-panel robustness exercise.

The empirical conclusion is therefore twofold. First, the directions of daily portfolio-loss extremes have a concentrated angular structure: in the $d = 24$ panel, ten AGCs explain about 91% of anchored variation. Second, this angular reduction has practical content beyond the descriptive spectrum. It preserves bounded capped-excess summaries and normalized VaR tail constants across a broad set of portfolio weights, giving a direct empirical role for the reconstruction theory in Section 2.5.

5 Discussion

Relative to the extremal-PCA (Drees and Sabourin, 2021; Drees, 2025) and spherical-PCA literatures (Fletcher et al., 2004; Jung et al., 2012), AGCA occupies a deliberate niche. It trades some geometric flexibility for a fixed, interpretable benchmark and for an oracle CLT based on bounded angular

departures. The benchmark makes scores and loadings read as departures from balanced complete dependence, and the bounded tangent geometry makes the oracle AGCA CLT depend on angular convergence and a second-order angular-bias condition rather than radial regular variation or tail-moment assumptions. These features are hard to obtain jointly in the more flexible manifold or multivariate regular variation approaches. The restriction can be less parsimonious when the angular law has curved geometry, strong face structure, or several active regimes, but it gives AGCA its benchmark-relative interpretation, exact spectral solution, and clean inferential target.

References

- Avella Medina, M., R. A. Davis, and G. Samorodnitsky (2025). Insights into kernel pca with application to multivariate extremes. *SIAM Journal on Mathematics of Data Science* 7(2), 777–801.
- Butsch, L. and V. Fasen-Hartmann (2025). Estimation of the number of principal components in high-dimensional multivariate extremes. *Scandinavian Journal of Statistics* 52(4), 2270–2313.
- Chautru, E. (2015). Dimension reduction in multivariate extreme value analysis. *Electronic Journal of Statistics* 9(1), 383–418.
- Chen, A. Y. and T. Zimmermann (2026). Open Source Asset Pricing. <https://www.openassetpricing.com/>. Accessed July 7, 2026.
- Cooley, D., A. Sabourin, and T. Wixson (2026). Principal component analysis for multivariate extremes. *arXiv preprint arXiv:2606.07213*.
- Cooley, D. and E. Thibaud (2019). Decompositions of dependence for high-dimensional extremes. *Biometrika* 106(3), 587–604.
- Dai, X. and H.-G. Müller (2018). Principal component analysis for functional data on riemannian manifolds and spheres. *The Annals of Statistics* 46(6B), 3334–3361.

- De Haan, L. and A. Ferreira (2006). *Extreme value theory: an introduction*. Springer.
- Drees, H. (2025). Asymptotic behavior of principal component projections for multivariate extremes. *arXiv preprint arXiv:2503.22296*.
- Drees, H. and A. Sabourin (2021). Principal component analysis for multivariate extremes. *Electronic Journal of Statistics* 15(1), 908–943.
- Einmahl, J. H. J., L. De Haan, and V. I. Piterbarg (2001). Nonparametric estimation of the spectral measure of an extreme value distribution. *The Annals of Statistics* 29(5), 1401–1423.
- Einmahl, J. H. J., L. De Haan, and A. K. Sinha (1997). Estimating the spectral measure of an extreme value distribution. *Stochastic Processes and their Applications* 70(2), 143–171.
- Einmahl, J. H. J., A. Krajina, and J. Segers (2012). An M-estimator for tail dependence in arbitrary dimensions. *The Annals of Statistics* 40(3), 1764–1793.
- Einmahl, J. H. J. and J. Segers (2009). Maximum empirical likelihood estimation of the spectral measure of an extreme-value distribution. *The Annals of Statistics* 37(5B), 2953–2989.
- Engelke, S. and A. S. Hitz (2020). Graphical models for extremes. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 82(4), 871–932.
- Fletcher, P. T., C. Lu, S. M. Pizer, and S. Joshi (2004). Principal geodesic analysis for the study of nonlinear statistics of shape. *IEEE Transactions on Medical Imaging* 23(8), 995–1005.
- Fomichov, V. and J. Ivanovs (2023). Spherical clustering in detection of groups of concomitant extremes. *Biometrika* 110(1), 135–153.
- French, K. R. (2026). Kenneth R. French Data Library. https://mba.tuck.dartmouth.edu/pages/faculty/ken.french/data_library.html. Accessed July 7, 2026.
- Goix, N., A. Sabourin, and S. Cléménçon (2017). Sparse representation of multivariate extremes with applications to anomaly detection. *Journal of Multivariate Analysis* 161, 12–31.

- Huckemann, S. and H. Ziezold (2006). Principal component analysis for riemannian manifolds, with an application to triangular shape spaces. *Advances in Applied Probability* 38(2), 299–319.
- Janßen, A. and P. Wan (2020). k -means clustering of extremes. *Electronic Journal of Statistics* 14(1), 1211–1233.
- Jung, S., I. L. Dryden, and J. S. Marron (2012). Analysis of principal nested spheres. *Biometrika* 99(3), 551–568.
- Lhaut, S., H. Rootzén, and J. Segers (2026). Simulation of multivariate extremes: A wasserstein–aitchison gan approach. *Extremes* 29(2), 157–194.
- Meyer, N. and O. Wintenberger (2021). Sparse regular variation. *Advances in Applied Probability* 53(4), 1115–1148.
- Mourahib, A., A. Kiriliouk, and J. Segers (2025). Multivariate generalized pareto distributions along extreme directions. *Extremes* 28(2), 239–272.
- Resnick, S. I. (2007). *Heavy-tail phenomena: probabilistic and statistical modeling*. Springer.
- Rootzén, H., J. Segers, and J. L. Wadsworth (2018). Multivariate generalized pareto distributions: Parametrizations, representations, and properties. *Journal of Multivariate Analysis* 165, 117–131.
- Rootzén, H. and N. Tajvidi (2006). Multivariate generalized pareto distributions. *Bernoulli* 12(5), 917–930.
- Sommer, S., F. Lauze, and M. Nielsen (2014). Optimization over geodesics for exact principal geodesic analysis. *Advances in Computational Mathematics* 40(2), 283–313.
- Tabaghi, P., M. Khanzadeh, Y. Wang, and S. Mirarab (2024). Principal component analysis in space forms. *IEEE Transactions on Signal Processing* 72, 4428–4443.
- Wan, P. (2026). Characterizing extremal dependence on a hyperplane. *Biometrika* 113(2), asag015.

Supplementary Material for

Anchored Geodesic Analysis for Multivariate Extremes

S1 Complementary theory

This supplementary section collects theoretical material that supports the construction of AGCA. It records the geometry of the anchored coordinates, studies face- and axis-supported angular laws, and discusses the canonical anchor, data-adaptive anchors, and anchor-free spherical variants.

S1.1 Geometry of the anchored coordinates

This subsection collects auxiliary coordinate identities used in the proofs. It also records that the anchored coordinates agree with the Riemannian log map to first order near the anchor, while $u_\mu(g)$ remains bounded near the edge of the hemisphere.

Lemma S1 (Positive directions lie in every interior hemisphere). *If $\mu \in \mathbb{S}_{++}^{d-1}$, then*

$$\mathbb{S}_+^{d-1} \subset \mathcal{H}_\mu, \quad \mathcal{H}_\mu := \{g \in \mathbb{S}^{d-1} : \langle \mu, g \rangle > 0\}.$$

Proof. For $g \in \mathbb{S}_+^{d-1}$, all coordinates are nonnegative and at least one coordinate is positive. Since every coordinate of μ is positive, $\langle \mu, g \rangle > 0$. □

The lemma is the reason the anchored coordinates are globally valid on the positive angular sample space. For $g \in \mathcal{H}_\mu$, recall

$$a_\mu(g) := \langle \mu, g \rangle, \quad u_\mu(g) := (I - \mu\mu^\top)g, \quad v_\mu(g) := g / \langle \mu, g \rangle - \mu.$$

The inverse gnomonic map is

$$\phi_\mu(v) := \frac{\mu + v}{(1 + \|v\|_2^2)^{1/2}}, \quad v \in \mu^\perp.$$

Proposition S1 (Anchored coordinate and membership identities). *For every $g \in \mathbb{S}_+^{d-1}$,*

$$g = \phi_\mu(v_\mu(g)), \quad d_g(g, \mu) = \arccos(a_\mu(g)) = \arcsin \|u_\mu(g)\|_2 = \arctan \|v_\mu(g)\|_2.$$

For any linear $W \subset \mu^\perp$,

$$M_\mu(W) = \phi_\mu(W),$$

where $\phi_\mu(W) := \{\phi_\mu(w) : w \in W\}$. Consequently,

$$g \in M_\mu(W) \iff v_\mu(g) \in W \iff u_\mu(g) \in W.$$

Proof. For $g \in \mathbb{S}_+^{d-1}$, Lemma S1 gives $a_\mu(g) > 0$, and

$$g = a_\mu(g)\mu + u_\mu(g), \quad a_\mu(g)^2 + \|u_\mu(g)\|_2^2 = 1.$$

Since $d_g(g, \mu) = \arccos(a_\mu(g))$, this gives the first three distance identities. The last one follows from $v_\mu(g) = u_\mu(g)/a_\mu(g)$ and $\tan(d_g(g, \mu)) = \|u_\mu(g)\|_2/a_\mu(g)$. Since $v_\mu(g) = u_\mu(g)/a_\mu(g)$, the identity $g = \phi_\mu(v_\mu(g))$ follows from

$$\phi_\mu(v_\mu(g)) = \frac{\mu + u_\mu(g)/a_\mu(g)}{(1 + \|u_\mu(g)\|_2^2/a_\mu(g)^2)^{1/2}} = a_\mu(g)\mu + u_\mu(g) = g.$$

If $w \in W$, then $\phi_\mu(w) \in \mathbb{S}^{d-1}$, has positive inner product with μ , and belongs to $\text{span}(\{\mu\} \cup W)$, so $\phi_\mu(w) \in M_\mu(W)$. Conversely, if $h \in M_\mu(W)$, then $h = a(\mu + w)$ for some $a > 0$ and $w \in W$. Since $h \in \mathbb{S}^{d-1}$ and $w \perp \mu$, $a = (1 + \|w\|_2^2)^{-1/2}$, so $h = \phi_\mu(w)$. Hence $M_\mu(W) = \phi_\mu(W)$. The membership equivalences follow because $g \in M_\mu(W)$ if and only if $v_\mu(g) \in W$, and $u_\mu(g) =$

$$a_\mu(g)v_\mu(g).$$

□

These identities are the coordinate facts used by the AGCA criterion. The membership equivalence says that anchored great subspheres become linear subspaces in either u_μ - or v_μ -coordinates. Proposition 1 gives the additional residual identity that turns spherical reconstruction into the spectral second-moment problem in the main text.

Proposition S2 (Log-map and gnomonic coordinate formulas). *For $v \in \mu^\perp$,*

$$a_\mu(\phi_\mu(v)) = \frac{1}{(1 + \|v\|_2^2)^{1/2}}, \quad d_g(\phi_\mu(v), \mu) = \arctan \|v\|_2.$$

For $g \in \mathbb{S}_+^{d-1}$, let $\text{Log}_\mu(g)$ denote the Riemannian logarithm map on the unit sphere at μ . It satisfies

$$\text{Log}_\mu(g) = \frac{\arctan \|v_\mu(g)\|_2}{\|v_\mu(g)\|_2} v_\mu(g) = \frac{\arcsin \|u_\mu(g)\|_2}{\|u_\mu(g)\|_2} u_\mu(g),$$

with the prefactors interpreted by continuity as one at $g = \mu$. Hence

$$\text{Log}_\mu(g) = v_\mu(g) + O\{\|v_\mu(g)\|_2^3\} = u_\mu(g) + O\{\|u_\mu(g)\|_2^3\} \quad \text{as } g \rightarrow \mu.$$

Finally, for any linear $W \subset \mu^\perp$,

$$d_g(g, M_\mu(W)) = \arctan \left[\frac{\|v_\mu(g) - \Pi_W v_\mu(g)\|_2}{(1 + \|\Pi_W v_\mu(g)\|_2^2)^{1/2}} \right].$$

Proof. The identities for $a_\mu(\phi_\mu(v))$ and $d_g(\phi_\mu(v), \mu)$ follow from $\langle \mu, v \rangle = 0$, which gives

$$\langle \mu, \phi_\mu(v) \rangle = (1 + \|v\|_2^2)^{-1/2}.$$

The distance formula follows because $\arccos((1 + \|v\|_2^2)^{-1/2}) = \arctan \|v\|_2$. The log-map identity

is the standard unit-sphere formula

$$\text{Log}_\mu(g) = \frac{\theta}{\sin \theta}(g - \cos(\theta)\mu), \quad \theta = d_g(g, \mu),$$

together with $\cos \theta = a_\mu(g)$, $\sin \theta = \|u_\mu(g)\|_2$, and Proposition S1. The expansion follows from the Taylor series of $\arctan x/x$ and $\arcsin x/x$.

For the gnomonic residual, write $v = v_\mu(g)$, $v_W = \Pi_W v$, and $v_\perp = v - v_W$. Since $u_\mu(g) = v/(1 + \|v\|_2^2)^{1/2}$ and $\Pi_W u_\mu(g) = v_W/(1 + \|v\|_2^2)^{1/2}$, Proposition 1 gives

$$\sin d_g(g, M_\mu(W)) = \frac{\|v_\perp\|_2}{(1 + \|v_W\|_2^2 + \|v_\perp\|_2^2)^{1/2}}.$$

Since $d_g(g, M_\mu(W)) < \pi/2$,

$$d_g(g, M_\mu(W)) = \arcsin \left(\frac{\|v_\perp\|_2}{(1 + \|v_W\|_2^2 + \|v_\perp\|_2^2)^{1/2}} \right).$$

Using $\arcsin x = \arctan(x/(1 - x^2)^{1/2})$ gives the stated formula. $\square$

These formulas place the anchored coordinates relative to standard spherical geometry. The gnomonic map ϕ_μ gives a global coordinate map for the visible hemisphere, while the log-map expansion explains why both anchored coordinates behave like usual tangent coordinates near the anchor. Away from the anchor, the two coordinates differ: $v_\mu(g)$ keeps the exact gnomonic linearization, whereas $u_\mu(g)$ remains bounded and is therefore the coordinate used for the AGCA second moment.

The bounded loss $\sin^2 d_g$ is chosen to preserve this exact projection geometry, not as a generic replacement for squared geodesic distance. On the relevant hemisphere it is monotone in the geodesic residual, comparable to d_g^2 , and has the same zero-risk sets. Its advantage is that the residual becomes an ordinary Euclidean squared residual in the bounded coordinates $u_\mu(g)$, which is what leads to the eigensolution in the main text.

Proposition S3 (Squared geodesic loss and bounded geodesic loss). *For any linear subspace $W \subset \mu^\perp$ and any random $G \in \mathbb{S}_+^{d-1}$,*

$$(2/\pi)^2 d_g(G, M_\mu(W))^2 \leq \sin^2 d_g(G, M_\mu(W)) \leq d_g(G, M_\mu(W))^2$$

almost surely. Consequently, zero risk under $\sin^2 d_g$ is equivalent to exact geodesic representation in $M_\mu(W)$.

Proof. Because $\mu \in M_\mu(W)$ and $G \in \mathcal{H}_\mu$, $d_g(G, M_\mu(W)) < \pi/2$. The inequality follows from $(2/\pi)t \leq \sin t \leq t$ on $[0, \pi/2]$. The zero-risk statement follows because $\sin t = 0$ if and only if $t = 0$ on this interval. $\square$

S1.2 Face and axis-supported angular laws

This subsection studies how AGCA behaves when some variables have asymptotically independent extremes. At the angular level, such episodes may appear as mass on coordinate faces or near coordinate axes: one subset of variables dominates the extreme direction while the remaining coordinates are comparatively small. The goal here is to show what the AGCA population objects do once such angular directions are part of the limiting law G .

The results below make three simple points. First, face-supported angular laws enter AGCA through bounded anchored departures, whose span determines the component rank. Second, if one variable dominates the extreme direction, AGCA returns the corresponding contrast from the anchor toward that variable, and this conclusion is stable when the mass is only near the axis. Third, if all axes are represented symmetrically, there is no preferred one-dimensional contrast, so the spectrum is tied rather than artificially low-dimensional.

The first result makes precise the statement that coordinate faces do not cause a singularity for AGCA. The bounded coordinate $u_\mu(G)$ converts face-supported angular laws into ordinary bounded tangent vectors, and the population component space is exactly the span of those tangent departures. Thus asymptotically independent face support is compatible with the same eigendecomposition

used for jointly extreme angular laws.

Proposition S4 (Anchored span of face-supported laws). *Let $G \in \mathbb{S}_+^{d-1}$, let $U_\mu := u_\mu(G)$, and let $\Sigma_\mu := \mathbb{E}[U_\mu U_\mu^\top]$. Then $\text{range}(\Sigma_\mu) = \text{span}\{u_\mu(g) : g \in \text{supp}(G)\}$. Consequently, if G is supported on the coordinate face*

$$F_S := \{g \in \mathbb{S}_+^{d-1} : g_j = 0 \text{ for } j \notin S\}, \quad S \subset \{1, \dots, d\},$$

then $r_\mu = \text{rank}(\Sigma_\mu) \leq \min\{|S|, d-1\}$.

Proof. For any $x \in \mu^\perp$,

$$x^\top \Sigma_\mu x = \mathbb{E}[(x^\top U_\mu)^2].$$

Hence $x \in \ker(\Sigma_\mu)$ if and only if $x^\top U_\mu = 0$ almost surely. Since u_μ is continuous, this is equivalent to x being orthogonal to $u_\mu(g)$ for every $g \in \text{supp}(G)$. Thus $\ker(\Sigma_\mu) = \text{span}\{u_\mu(g) : g \in \text{supp}(G)\}^\perp$ inside $\mu^\perp$. Since Σ_μ is symmetric on $\mu^\perp$, this gives the range identity. If G is supported on F_S , then $G \in E_S := \text{span}\{e_j : j \in S\}$ almost surely, and $u_\mu(G) = (I - \mu\mu^\top)G$ lies in $(I - \mu\mu^\top)E_S$. This space has dimension at most $\min\{|S|, d-1\}$, so the same bound holds for $\text{rank}(\Sigma_\mu)$. $\square$

The next result gives the population loading for the simplest asymptotically independent episode: all angular mass is on one coordinate axis. In that case AGCA has a single nonzero population component. With the canonical anchor, the loading has one positive coordinate for the dominant variable and equal negative coordinates for the others, so it is exactly the contrast from balanced co-movement toward that axis. The same conclusion is stable for laws concentrated near, rather than exactly on, the axis.

Proposition S5 (Axis loading and near-axis stability). *Let e_j be the j th coordinate vector and let*

$$u_j := u_\mu(e_j) = e_j - \mu_j \mu, \quad \gamma_j := \|u_j\|_2^2 = 1 - \mu_j^2.$$

If $G = e_j$ almost surely, then $\Sigma_\mu = u_j u_j^\top$, $r_\mu = 1$, and the unique rank-one loading, up to sign, is $b_j^* := u_j / \|u_j\|_2$. For the canonical anchor $\mu_0 = d^{-1/2} \mathbf{1}$,

$$b_j^* = \frac{e_j - d^{-1} \mathbf{1}}{\sqrt{1 - 1/d}}.$$

More generally, suppose $\mathbb{E}[\|G - e_j\|_2^2] \leq \delta^2$. Then

$$\|\Sigma_\mu - u_j u_j^\top\|_{\text{op}} \leq 2\sqrt{\gamma_j} \delta + \delta^2.$$

If $\varepsilon_j := 2\sqrt{\gamma_j} \delta + \delta^2 < \gamma_j/2$, the leading rank-one population projector $P_{\mu,1}$ of Σ_μ is unique and satisfies

$$\|P_{\mu,1} - b_j^* b_j^{*\top}\|_{\text{op}} \leq \frac{2\varepsilon_j}{\gamma_j}.$$

Proof. If $G = e_j$ almost surely, then $U_\mu = u_j$ almost surely, so $\Sigma_\mu = u_j u_j^\top$. Since $\mu \in \mathbb{S}_{++}^{d-1}$, $\mu_j < 1$ for $d \geq 2$, and therefore $\gamma_j > 0$. The rank-one eigensolution and the canonical-anchor formula follow immediately from $u_{\mu_0}(e_j) = e_j - d^{-1} \mathbf{1}$. For the near-axis statement, write $V := u_\mu(G) - u_j = (I - \mu\mu^\top)(G - e_j)$. Since $I - \mu\mu^\top$ is an orthogonal projector, $\mathbb{E}[\|V\|_2^2] \leq \delta^2$. Also $\|\mathbb{E}V\|_2 \leq \mathbb{E}\|V\|_2 \leq \delta$. Expanding

$$\Sigma_\mu = \mathbb{E}[(u_j + V)(u_j + V)^\top] = u_j u_j^\top + u_j (\mathbb{E}V)^\top + (\mathbb{E}V) u_j^\top + \mathbb{E}[VV^\top]$$

gives

$$\|\Sigma_\mu - u_j u_j^\top\|_{\text{op}} \leq 2\|u_j\|_2 \|\mathbb{E}V\|_2 + \mathbb{E}[\|V\|_2^2] \leq 2\sqrt{\gamma_j} \delta + \delta^2.$$

The rank-one projector bound is the Davis–Kahan bound applied to the perturbation of $u_j u_j^\top$, whose leading eigengap is γ_j . $\square$

The final result records an important negative case. AGCA can represent axis-supported angular laws, but it does not force them to be low-dimensional. If a canonical-anchor angular law puts equal mass on every coordinate axis, every axis contrast is equally important. The population spectrum

is then completely tied on $\mu_0^\perp$, so individual loadings are not identified. This is the appropriate diagnostic: there is finite axis-supported angular structure, but no preferred one-dimensional component.

Proposition S6 (Symmetric axis support has no preferred loading). *Let $\mu_0 = d^{-1/2}\mathbf{1}$, and suppose $\mathbb{P}[G = e_j] = 1/d$ for $j = 1, \dots, d$. Then $\Sigma_{\mu_0} = \frac{1}{d} \left(I - \frac{\mathbf{1}\mathbf{1}^\top}{d} \right)$. Hence all $d - 1$ nonzero eigenvalues are equal to $1/d$, any orthonormal basis of $\mu_0^\perp$ is a population loading basis, and*

$$\text{AVE}_{\mu_0, p} = \frac{p}{d - 1}, \quad p = 0, \dots, d - 1.$$

Proof. For the canonical anchor, $u_{\mu_0}(e_j) = e_j - d^{-1}\mathbf{1}$. Therefore

$$\Sigma_{\mu_0} = \frac{1}{d} \sum_{j=1}^d (e_j - d^{-1}\mathbf{1})(e_j - d^{-1}\mathbf{1})^\top = \frac{1}{d} \left(I - \frac{\mathbf{1}\mathbf{1}^\top}{d} \right).$$

The matrix in parentheses is the orthogonal projector onto $\mu_0^\perp$. The eigenvalue and anchored-variation statements follow directly. □

S1.3 Anchor choice and anchor-free variants

Once the anchor is fixed, Theorem 1 identifies the optimal AGCA component spaces by a single eigendecomposition. The substantive modeling choice is therefore upstream: which benchmark should define the origin of the score system, and should one work in the anchored great-subspace family at all?

The anchor has two roles. First, it defines the zero-departure angular profile: a rank-zero anchored approximation replaces every angular observation by μ . Second, it fixes the origin of the component scores, so the loadings are contrasts describing how an extreme profile moves away from that benchmark. A data-adaptive anchor may reduce reconstruction loss, but the fitted contrasts then become relative to an estimated center or axis, which can weaken fixed-benchmark interpretation and add anchor-estimation variability.

Proposition S7 (Anchor-risk identities). *Let $G \in \mathbb{S}_+^{d-1}$, let $M := \mathbb{E}[GG^\top]$, and set $P_\mu := I - \mu\mu^\top$.*

Then

$$\Sigma_\mu = \mathbb{E}[u_\mu(G)u_\mu(G)^\top] = P_\mu M P_\mu$$

on $\mu^\perp$, and

$$\text{tr}(\Sigma_\mu) = 1 - \langle \mu, M\mu \rangle = \mathcal{R}_\mu(\{0\}) = \mathbb{E}[\sin^2 d_g(G, \mu)].$$

Proof. Since $u_\mu(G) = P_\mu G$, the first identity follows by taking expectations. Taking traces and using $\text{tr}(M) = \mathbb{E}[\|G\|_2^2] = 1$ gives $\text{tr}(\Sigma_\mu) = \text{tr}(P_\mu M) = 1 - \langle \mu, M\mu \rangle$. The remaining identities are the rank-zero case of the risk decomposition in Theorem 1. $\square$

These identities clarify what an empirical anchor can optimize. At rank zero, choosing an anchor to minimize AGCA risk is equivalent to choosing μ to maximize $\langle \mu, M\mu \rangle$, so the anchor moves toward the dominant angular direction of the law.

Proposition S8 (Anchor-free spherical eigensolution). *Let $G \in \mathbb{S}_+^{d-1}$, let $M := \mathbb{E}[GG^\top]$, and let $L \subset \mathbb{R}^d$ be a linear subspace with $\dim(L) = q$. Define the anchor-free bounded geodesic risk*

$$\mathcal{R}_{\text{af}}(L) := \mathbb{E}[\sin^2 d_g(G, \mathbb{S}^{d-1} \cap L)].$$

If $c_1, \dots, c_d$ are orthonormal eigenvectors of M , with eigenvalues $\alpha_1 \geq \dots \geq \alpha_d$, then $L_q^ := \text{span}(c_1, \dots, c_q)$ minimizes $\mathcal{R}_{\text{af}}(L)$ over all q -dimensional linear subspaces of $\mathbb{R}^d$, and $\mathcal{R}_{\text{af}}(L_q^*) = \sum_{j=q+1}^d \alpha_j$.*

Proof. For any $g \in \mathbb{S}^{d-1}$ and any linear subspace L ,

$$\sin^2 d_g(g, \mathbb{S}^{d-1} \cap L) = \|g - \Pi_L g\|_2^2 = 1 - \langle g, \Pi_L g \rangle,$$

where the projection onto the subspace is $\Pi_L g / \|\Pi_L g\|_2$ when $\Pi_L g \neq 0$; the identity also holds by

continuity if $\Pi_L g = 0$. Taking expectations,

$$\mathcal{R}_{\text{af}}(L) = 1 - \text{tr}(\Pi_L M).$$

Thus minimizing $\mathcal{R}_{\text{af}}(L)$ is equivalent to maximizing $\text{tr}(\Pi_L M)$ over q -dimensional L . The Ky Fan maximum principle gives the span of the leading q eigenvectors, and the minimized risk is the trailing eigensum. $\square$

Proposition S8 shows that the anchor-free problem is an exact benchmark-free alternative, not an approximation to fixed-anchor AGCA. The $q = 1$ version identifies the leading angular axis; when that axis is unique and can be oriented with positive coordinates, it also gives a loss-matched self-anchored variant.

Proposition S9 (Canonical anchor as maximal hemisphere margin). *For $\mu \in \mathbb{S}_{++}^{d-1}$, define the hemisphere margin, the smallest inner product between μ and any positive-sphere direction, by*

$$m_+(\mu) := \inf_{g \in \mathbb{S}_+^{d-1}} \langle \mu, g \rangle.$$

Then $m_+(\mu) = \min_j \mu_j$, and $\max_{\mu \in \mathbb{S}_{++}^{d-1}} m_+(\mu) = d^{-1/2}$, with unique maximizer μ_0 in (2.1). Moreover, the same anchor uniquely minimizes both the largest geodesic distance to the positive sphere and the largest gnomonic coordinate norm:

$$\arg \min_{\mu \in \mathbb{S}_{++}^{d-1}} \sup_{g \in \mathbb{S}_+^{d-1}} d_g(g, \mu) = \{\mu_0\}, \quad \arg \min_{\mu \in \mathbb{S}_{++}^{d-1}} \sup_{g \in \mathbb{S}_+^{d-1}} \|v_\mu(g)\|_2 = \{\mu_0\}.$$

Proof. For any $g \in \mathbb{S}_+^{d-1}$,

$$\langle \mu, g \rangle = \sum_{j=1}^d \mu_j g_j \geq \left(\min_j \mu_j \right) \sum_{j=1}^d g_j = \left(\min_j \mu_j \right) \|g\|_1 \geq \min_j \mu_j,$$

where the last inequality uses $g \in \mathbb{S}_+^{d-1}$, so $\|g\|_1 \geq \|g\|_2 = 1$. Equality is attained at e_j for an index minimizing μ_j . Hence $m_+(\mu) = \min_j \mu_j$. If $m = \min_j \mu_j$, then $1 = \sum_{j=1}^d \mu_j^2 \geq dm^2$, so

$m \leq d^{-1/2}$, with equality only when all coordinates are equal. The distance and gnomonic claims follow because

$$\sup_{g \in \mathbb{S}_+^{d-1}} d_g(g, \mu) = \arccos(m_+(\mu)), \quad \sup_{g \in \mathbb{S}_+^{d-1}} \|v_\mu(g)\|_2 = (m_+(\mu)^{-2} - 1)^{1/2}.$$

□

The canonical anchor is therefore a nonrandom, permutation-symmetric benchmark with maximal uniform angular margin over the positive sphere, not a sample center chosen to minimize empirical loss. Loss-matched and spherical Fréchet anchors are useful sensitivity analyses when no benchmark is privileged, but in samples they replace the fixed reference by an estimated one.

S2 Proofs

This section gives the proofs of the mathematical statements in the main text, in the order in which those statements appear.

S2.1 Anchored projection and population eigensolution

Proof of Proposition 1. Let $g \in \mathbb{S}_+^{d-1}$ and write $v := v_\mu(g) \in \mu^\perp$. By Proposition S1, $g = \phi_\mu(v)$ and $M_\mu(W) = \phi_\mu(W)$. Because $\arccos$ is decreasing, geodesic projection onto $M_\mu(W)$ is equivalent to maximizing $\langle g, \phi_\mu(w) \rangle$ over $w \in W$. Since $g = \phi_\mu(v)$,

$$\langle g, \phi_\mu(w) \rangle = \frac{1 + \langle v, w \rangle}{(1 + \|v\|_2^2)^{1/2}(1 + \|w\|_2^2)^{1/2}}.$$

Let $v_W := \Pi_W v$. Since $v - v_W \perp W$, $\langle v, w \rangle = \langle v_W, w \rangle$. By Cauchy–Schwarz in $\mathbb{R} \times W$,

$$1 + \langle v_W, w \rangle \leq (1 + \|v_W\|_2^2)^{1/2}(1 + \|w\|_2^2)^{1/2},$$

with equality if and only if $w = v_W$. Dividing by the denominator in the preceding display shows that $\langle g, \phi_\mu(w) \rangle$ is uniquely maximized over $w \in W$ at $w = v_W = \Pi_W v$. Hence the geodesic projection of g onto $M_\mu(W)$ is

$$\Pi_{M_\mu(W)}(g) = \phi_\mu(\Pi_W v).$$

At this maximizing point,

$$\cos d_g(g, M_\mu(W)) = \left(\frac{1 + \|\Pi_W v\|_2^2}{1 + \|v\|_2^2} \right)^{1/2}.$$

Therefore

$$\sin^2 d_g(g, M_\mu(W)) = \frac{\|v - \Pi_W v\|_2^2}{1 + \|v\|_2^2}.$$

Since

$$u_\mu(g) = \frac{v}{(1 + \|v\|_2^2)^{1/2}}, \quad \Pi_W u_\mu(g) = \frac{\Pi_W v}{(1 + \|v\|_2^2)^{1/2}},$$

the identity (2.4) follows.

Finally,

$$\phi_\mu(\Pi_W v) = \frac{\mu + \Pi_W v}{(1 + \|\Pi_W v\|_2^2)^{1/2}}.$$

Substituting $v = u_\mu(g)/a_\mu(g)$ gives

$$\phi_\mu(\Pi_W v) = \frac{a_\mu(g)\mu + \Pi_W u_\mu(g)}{(a_\mu(g)^2 + \|\Pi_W u_\mu(g)\|_2^2)^{1/2}},$$

which is (2.5). □

Proof of Theorem 1. By Proposition 1, $\mathcal{R}_\mu(W) = \mathbb{E}[\|U_\mu - \Pi_W U_\mu\|_2^2]$. Expanding the squared norm and using symmetry and idempotence of Π_W ,

$$\mathcal{R}_\mu(W) = \mathbb{E}[\|U_\mu\|_2^2] - \mathbb{E}[\langle U_\mu, \Pi_W U_\mu \rangle].$$

The second term is

$$\mathbb{E}[\langle U_\mu, \Pi_W U_\mu \rangle] = \text{tr}(\Pi_W \mathbb{E}[U_\mu U_\mu^\top]) = \text{tr}(\Pi_W \Sigma_\mu).$$

Since $\mathbb{E}[\|U_\mu\|_2^2] = \text{tr}(\Sigma_\mu)$, this proves $\mathcal{R}_\mu(W) = \text{tr}((I - \Pi_W)\Sigma_\mu)$.

The first term $\text{tr}(\Sigma_\mu)$ does not depend on W . Minimizing $\mathcal{R}_\mu(W)$ over p -dimensional subspaces is therefore equivalent to maximizing $\text{tr}(\Pi_W \Sigma_\mu)$. By the Ky Fan maximum principle (see, e.g., [Horn and Johnson, 2013](#)), the maximum is attained by the span of the eigenvectors corresponding to the largest p eigenvalues of Σ_μ . For $W_{\mu,p} := \text{span}(b_1, \dots, b_p)$,

$$\mathcal{R}_\mu(W_{\mu,p}) = \text{tr}(\Sigma_\mu) - \sum_{j=1}^p \lambda_j = \sum_{j=p+1}^{d-1} \lambda_j.$$

□

S2.2 Rank, scores, and loading interpretation

Proof of Corollary 1. By Proposition 1, $\mathcal{R}_\mu(W) = 0$ if and only if $U_\mu \in W$ almost surely. If this holds, then $\text{range}(\Sigma_\mu) \subset W$, so $\text{rank}(\Sigma_\mu) \leq \dim(W)$. Conversely, if y is orthogonal to $\text{range}(\Sigma_\mu)$, then $0 = \langle y, \Sigma_\mu y \rangle = \mathbb{E}[\langle y, U_\mu \rangle^2]$, so $\langle y, U_\mu \rangle = 0$ almost surely. Hence $U_\mu \in \text{range}(\Sigma_\mu)$ almost surely, and any $W \supset \text{range}(\Sigma_\mu)$ has zero risk. At the minimal dimension $p = \text{rank}(\Sigma_\mu)$, W must equal $\text{range}(\Sigma_\mu)$. □

Proof of Proposition 2. If $\eta = 0$, then $c_\eta(t) \equiv \mu$, and the derivative is zero. Suppose $\eta \neq 0$ and write $r := \|\eta\|_2$. On the unit sphere,

$$c_\eta(t) = \text{Exp}_\mu(t\eta) = \cos(tr)\mu + \sin(tr)\eta/r.$$

Thus $c_\eta(0) = \mu$ and $c'_\eta(0) = \eta$. Since $\mu_j, \mu_k > 0$, the map $t \mapsto \log\{c_{\eta,j}(t)/c_{\eta,k}(t)\}$ is differentiable

at $t = 0$, and the chain rule gives

$$\left. \frac{d}{dt} \right|_{t=0} \log \frac{c_{\eta,j}(t)}{c_{\eta,k}(t)} = \frac{c'_{\eta,j}(0)}{c_{\eta,j}(0)} - \frac{c'_{\eta,k}(0)}{c_{\eta,k}(0)} = \frac{\eta_j}{\mu_j} - \frac{\eta_k}{\mu_k}.$$

□

S2.3 Low-rank reconstruction and tail simulation

Proof of Proposition 3. By definition, $\tilde{G}_{\mu,p} = \Pi_{M_\mu(W_{\mu,p})}(G)$. Applying Proposition 1 with $g = G$ and $W = W_{\mu,p}$ gives

$$\sin^2 d_g(G, \tilde{G}_{\mu,p}) = \sin^2 d_g(G, M_\mu(W_{\mu,p})) = \|u_\mu(G) - \Pi_{W_{\mu,p}} u_\mu(G)\|_2^2.$$

Taking expectations and using (2.10) gives

$$\mathbb{E}[\sin^2 d_g(G, \tilde{G}_{\mu,p})] = \rho_{\mu,p} = \sum_{j=p+1}^{d-1} \lambda_j.$$

Finally, let $\theta = d_g(G, \tilde{G}_{\mu,p})$. Since $\mu \in M_\mu(W_{\mu,p})$ and $G \in \mathcal{H}_\mu$, the projection distance satisfies $\theta \leq d_g(G, \mu) < \pi/2$. Hence

$$\|G - \tilde{G}_{\mu,p}\|_2^2 = 2(1 - \cos \theta) = \frac{2}{1 + \cos \theta} \sin^2 \theta \leq 2 \sin^2 \theta.$$

Taking expectations gives the displayed equality for $\mathbb{E}[\|G - \tilde{G}_{\mu,p}\|_2^2]$, and the inequality follows from $\mathbb{E}[\sin^2 d_g(G, \tilde{G}_{\mu,p})] = \rho_{\mu,p}$. □

Proof of Proposition 4. Assumption 2 implies

$$\mathcal{L}(X/r \mid R > r) \rightsquigarrow \mathcal{L}(P_\alpha G),$$

where P_α is independent of G and $\mathbb{P}[P_\alpha > x] = x^{-\alpha}$, $x \geq 1$. Since bounded-Lipschitz distance

metrizes weak convergence on $\mathbb{R}^d$, $\varepsilon_r \rightarrow 0$.

It remains to bound $d_{\text{BL}}(\mathcal{L}(P_\alpha G), \mathcal{L}(P_\alpha \tilde{G}_{\mu,p}))$. Let

$$D_{\mu,p} := \|u_\mu(G) - \Pi_{W_{\mu,p}} u_\mu(G)\|_2.$$

By Proposition 3, $\mathbb{E}[D_{\mu,p}^2] = \rho_{\mu,p}$. If f satisfies $\|f\|_\infty \leq 1$ and $\text{Lip}(f) \leq 1$, then

$$\left| f(P_\alpha G) - f(P_\alpha \tilde{G}_{\mu,p}) \right| \leq \min\{2, P_\alpha \|G - \tilde{G}_{\mu,p}\|_2\}.$$

For $0 < \beta \leq 1$, $\min\{2, z\} \leq 2^{1-\beta} z^\beta$ for $z \geq 0$. Also, because $d_g(G, \tilde{G}_{\mu,p}) \leq d_g(G, \mu) < \pi/2$,

$$\|G - \tilde{G}_{\mu,p}\|_2^2 = 2\{1 - \cos d_g(G, \tilde{G}_{\mu,p})\} \leq 2 \sin^2 d_g(G, \tilde{G}_{\mu,p}) = 2D_{\mu,p}^2.$$

Hence

$$\left| \mathbb{E}[f(P_\alpha G)] - \mathbb{E}[f(P_\alpha \tilde{G}_{\mu,p})] \right| \leq 2^{1-\beta/2} \mathbb{E}[P_\alpha^\beta] \mathbb{E}[D_{\mu,p}^\beta].$$

The independence of P_α and G gives the factorization. Since $\beta < \alpha$,

$$\mathbb{E}[P_\alpha^\beta] = \frac{\alpha}{\alpha - \beta},$$

and Lyapunov's inequality gives

$$\mathbb{E}[D_{\mu,p}^\beta] \leq \{\mathbb{E}[D_{\mu,p}^2]\}^{\beta/2} = \rho_{\mu,p}^{\beta/2}.$$

Taking the supremum over the bounded-Lipschitz class yields

$$d_{\text{BL}}(\mathcal{L}(P_\alpha G), \mathcal{L}(P_\alpha \tilde{G}_{\mu,p})) \leq 2^{1-\beta/2} \frac{\alpha}{\alpha - \beta} \rho_{\mu,p}^{\beta/2}.$$

The stated inequality follows by the triangle inequality with ε_r . □

Proof of Proposition 5. Write $W := X/R$. Since $h(W) \leq H_h$, the event $\{h(X) > rx\}$ is contained in $\{R > r\}$ whenever $x \geq H_h$. Therefore

$$\frac{\mathbb{P}[h(X) > rx]}{\mathbb{P}[R > r]} = \mathbb{P}[(R/r)h(W) > x \mid R > r], \quad x \geq H_h.$$

By Assumption 2 and the continuous mapping theorem,

$$(R/r)h(W) \mid R > r \rightsquigarrow P_\alpha h(G).$$

Writing $A_h := h(G)$, independence of P_α and G gives, for $x \geq H_h$,

$$\mathbb{P}[P_\alpha h(G) > x] = \mathbb{E}[\mathbb{P}[P_\alpha > x/A_h \mid A_h]] = x^{-\alpha} \mathbb{E}[A_h^\alpha],$$

with the conditional probability interpreted as zero when $A_h = 0$. The limiting survival function is continuous on $[H_h, \infty)$. Uniform convergence on compact subintervals follows from weak convergence, and the remaining tail is controlled by tightness; hence $\eta_{r,h} \rightarrow 0$.

Similarly, with $\tilde{A}_{h,p} := h(\tilde{G}_{\mu,p})$,

$$\mathbb{P}[h(P_\alpha \tilde{G}_{\mu,p}) > x] = \mathbb{E}[\mathbb{P}[P_\alpha > x/\tilde{A}_{h,p} \mid \tilde{A}_{h,p}]] = x^{-\alpha} \mathbb{E}[\tilde{A}_{h,p}^\alpha], \quad x \geq H_h.$$

It remains to bound the angular constants. If $0 < \alpha \leq 1$, the map $a \mapsto a^\alpha$ is α -Hölder on $[0, \infty)$, so

$$\left| C_h - \tilde{C}_{h,p} \right| \leq \mathbb{E} \left[\left| h(G) - h(\tilde{G}_{\mu,p}) \right|^\alpha \right] \leq L_h^\alpha \{ \mathbb{E} [\|G - \tilde{G}_{\mu,p}\|_2^2] \}^{\alpha/2}.$$

If $\alpha > 1$, the map $a \mapsto a^\alpha$ is Lipschitz on $[0, H_h]$ with constant $\alpha H_h^{\alpha-1}$, and hence

$$\left| C_h - \tilde{C}_{h,p} \right| \leq \alpha H_h^{\alpha-1} L_h \mathbb{E} [\|G - \tilde{G}_{\mu,p}\|_2].$$

Using Proposition 3 in both cases gives

$$\left| C_h - \tilde{C}_{h,p} \right| \leq K_{\alpha,h} (2\rho_{\mu,p})^{\gamma/2}.$$

Finally, for $x \geq H_h$,

$$\left| \frac{\mathbb{P}[h(X) > rx]}{\mathbb{P}[R > r]} - \mathbb{P}[h(P_\alpha \tilde{G}_{\mu,p}) > x] \right| \leq \eta_{r,h} + x^{-\alpha} \left| C_h - \tilde{C}_{h,p} \right|.$$

Taking the supremum over $x \geq H_h$ gives the stated bound. $\square$

Proof of Proposition 6. By Proposition 3, $\tilde{G}_{\mu,p} \in M_\mu(W_{\mu,p})$ almost surely, and $\|P_\alpha \tilde{G}_{\mu,p}\|_2 = P_\alpha \geq 1$. Therefore $\mathbb{P}[P_\alpha \tilde{G}_{\mu,p} \in \mathcal{C}_{\mu,p}] = 1$. Similarly, $P_\alpha G \in \mathcal{C}_{\mu,p}$ if and only if $G \in M_\mu(W_{\mu,p})$, because $P_\alpha \geq 1$ and $(P_\alpha G) / \|P_\alpha G\|_2 = G$. Thus

$$\mathbb{P}[P_\alpha G \in \mathcal{C}_{\mu,p}] = \mathbb{P}[G \in M_\mu(W_{\mu,p})].$$

Taking $A = \mathcal{C}_{\mu,p}$ in the supremum over Borel sets gives the lower bound. $\square$

S2.4 Portfolio tail functionals

Proof of Corollary 2. The map $s \mapsto \min\{1, s_+\}$ is bounded by one and one-Lipschitz. Therefore

$$\|f_{w,t,L}\|_\infty \leq 1, \quad \text{Lip}(f_{w,t,L}) \leq \|w\|_2 / L \leq \|w\|_1 / L \leq 1.$$

The claim follows by applying Proposition 4 with $f = f_{w,t,L}$. $\square$

Proof of Corollary 3. Apply Proposition 5 to $h_w(z) := (w^\top z)_+$. This score is nonnegative and positively homogeneous. On $\mathbb{S}^{d-1}$,

$$h_w(s) \leq \|w\|_2 \leq \|w\|_1 \leq L, \quad |h_w(s) - h_w(t)| \leq |w^\top(s - t)| \leq L \|s - t\|_2.$$

The tail identities for all $x \geq L$ and the bound on $|C_w - \tilde{C}_{w,p}|$ therefore follow from the proposition, with

$$K_{\alpha, h_w} \leq \begin{cases} L^\alpha, & 0 < \alpha \leq 1, \\ \alpha L^\alpha, & \alpha > 1. \end{cases}$$

The VaR formulas follow by solving $x^{-\alpha}C = u$; the condition $u < L^{-\alpha} \min(C_w, \tilde{C}_{w,p})$ ensures that both resulting quantiles are at least L , where the tail identities apply. The ratio identity is immediate from the formulas. Finally, if $\min(C_w, \tilde{C}_{w,p}) \geq c$, then

$$|\log q_w(u) - \log \tilde{q}_{w,p}(u)| = \alpha^{-1} |\log C_w - \log \tilde{C}_{w,p}| \leq (\alpha c)^{-1} |C_w - \tilde{C}_{w,p}|,$$

and the angular-constant bound gives the stated logarithmic VaR bound. $\square$

S2.5 Positive post-projection for tail simulation

The population reconstruction $\tilde{G}_{\mu,p}$, defined in (2.14), is a unit vector in the fitted anchored great-subspace model

$$M_\mu(W_{\mu,p}) = \text{span}(\{\mu\} \cup W_{\mu,p}) \cap \mathcal{H}_\mu,$$

but this model itself need not be contained in the positive orthant. For simulation on the positive cone, one can enforce positivity after the AGCA reconstruction by applying a deterministic positive post-projection. This correction preserves the low-rank AGCA step as the primary approximation, but it no longer preserves exact membership in $M_\mu(W_{\mu,p})$. The point of the following results is to quantify precisely the additional price paid for this positivity correction.

For $z \in \mathbb{R}^d$, let z_+ denote the coordinatewise positive part. When $z_+ \neq 0$, define $\mathcal{P}_+(z) := z_+ / \|z_+\|_2$. Since $\tilde{G}_{\mu,p} \in \mathcal{H}_\mu$ and $\mu \in \mathbb{S}_{++}^{d-1}$, $(\tilde{G}_{\mu,p})_+ \neq 0$ almost surely. Indeed, if all coordinates of $\tilde{G}_{\mu,p}$ were nonpositive, then $\langle \mu, \tilde{G}_{\mu,p} \rangle \leq 0$, contradicting $\tilde{G}_{\mu,p} \in \mathcal{H}_\mu$. Thus the positive

post-projected reconstruction

$$\tilde{G}_{\mu,p}^+ := \mathcal{P}_+(\tilde{G}_{\mu,p}) = (\tilde{G}_{\mu,p})_+ / \left\| (\tilde{G}_{\mu,p})_+ \right\|_2$$

is well defined and takes values in $\mathbb{S}_+^{d-1}$. Define the positivity-correction cost

$$\delta_{\mu,p}^+ := \mathbb{E} \left[\left\| \tilde{G}_{\mu,p} - \tilde{G}_{\mu,p}^+ \right\|_2^2 \right], \quad \omega_{\mu,p}^+ := (2\rho_{\mu,p})^{1/2} + (\delta_{\mu,p}^+)^{1/2}.$$

The quantity $\omega_{\mu,p}^+$ is the L^2 angular error budget for positive simulation: the first term is the AGCA reconstruction error from Proposition 3, and the second term is the extra movement needed to return the reconstructed direction to the positive sphere.

Proposition S10 (Positive post-projected reconstruction). *For $p = 0, \dots, d-1$, $\tilde{G}_{\mu,p}^+ \in \mathbb{S}_+^{d-1}$ almost surely, and*

$$\mathbb{E} \left[\left\| G - \tilde{G}_{\mu,p}^+ \right\|_2^2 \right] \leq (\omega_{\mu,p}^+)^2.$$

Proof. The preceding paragraph proves that $\tilde{G}_{\mu,p}^+$ is well defined and belongs to $\mathbb{S}_+^{d-1}$. By Minkowski's inequality,

$$\sqrt{\mathbb{E} \left[\left\| G - \tilde{G}_{\mu,p}^+ \right\|_2^2 \right]} \leq \sqrt{\mathbb{E} \left[\left\| G - \tilde{G}_{\mu,p} \right\|_2^2 \right]} + \sqrt{\mathbb{E} \left[\left\| \tilde{G}_{\mu,p} - \tilde{G}_{\mu,p}^+ \right\|_2^2 \right]}.$$

The first term is bounded by $(2\rho_{\mu,p})^{1/2}$ by Proposition 3, and the second term is $(\delta_{\mu,p}^+)^{1/2}$ by definition. Squaring gives the final display. $\square$

Proposition S11 (Positive post-projected tail simulation). *Suppose Assumption 2 holds, and let ε_r be defined by (2.15). Then $\varepsilon_r \rightarrow 0$. Moreover, for every $0 < \beta \leq 1$ with $\beta < \alpha$,*

$$d_{\text{BL}} \left(\mathcal{L}(X/r \mid R > r), \mathcal{L}(P_\alpha \tilde{G}_{\mu,p}^+) \right) \leq \varepsilon_r + 2^{1-\beta} \frac{\alpha}{\alpha - \beta} (\omega_{\mu,p}^+)^{\beta}.$$

Proof. As in the proof of Proposition 4, $\varepsilon_r \rightarrow 0$, and it remains to bound the distance between

$\mathcal{L}(P_\alpha G)$ and $\mathcal{L}(P_\alpha \tilde{G}_{\mu,p}^+)$. If $\|f\|_\infty \leq 1$ and $\text{Lip}(f) \leq 1$, then

$$\left| f(P_\alpha G) - f(P_\alpha \tilde{G}_{\mu,p}^+) \right| \leq \min\{2, P_\alpha \|G - \tilde{G}_{\mu,p}^+\|_2\}.$$

For $0 < \beta \leq 1$, $\min\{2, z\} \leq 2^{1-\beta} z^\beta$. Since P_α is independent of the angular variables and $\beta < \alpha$,

$$\left| \mathbb{E}[f(P_\alpha G)] - \mathbb{E}[f(P_\alpha \tilde{G}_{\mu,p}^+)] \right| \leq 2^{1-\beta} \mathbb{E}[P_\alpha^\beta] \mathbb{E}[\|G - \tilde{G}_{\mu,p}^+\|_2^\beta] \leq 2^{1-\beta} \frac{\alpha}{\alpha - \beta} (\omega_{\mu,p}^+)^{\beta},$$

where the last step uses Lyapunov's inequality and Proposition S10. Taking the supremum over the bounded-Lipschitz class and applying the triangle inequality with ε_r gives the claim. $\square$

Proposition S12 (Positive post-projected homogeneous tail scores). *Suppose Assumption 2 holds, and let $h : \mathbb{R}^d \rightarrow [0, \infty)$ satisfy the conditions used in Proposition 5. Let C_h , $\eta_{r,h}$, γ , and $K_{\alpha,h}$ be the corresponding quantities from that setup, and define $\tilde{C}_{h,p}^+ := \mathbb{E}[h(\tilde{G}_{\mu,p}^+)^{\alpha}]$. Then $\eta_{r,h} \rightarrow 0$ as $r \rightarrow \infty$. Moreover, for every $x \geq H_h$,*

$$\mathbb{P}[h(P_\alpha \tilde{G}_{\mu,p}^+) > x] = x^{-\alpha} \tilde{C}_{h,p}^+, \quad \sup_{x \geq H_h} \left| \frac{\mathbb{P}[h(X) > rx]}{\mathbb{P}[R > r]} - \mathbb{P}[h(P_\alpha \tilde{G}_{\mu,p}^+) > x] \right| \leq \eta_{r,h} + H_h^{-\alpha} K_{\alpha,h} (\omega_{\mu,p}^+)^{\gamma}.$$

Proof. The convergence $\eta_{r,h} \rightarrow 0$ is the threshold part of Proposition 5. The tail identity follows from the independence of P_α and G , since $\tilde{G}_{\mu,p}^+$ is a measurable function of G :

$$\mathbb{P}[h(P_\alpha \tilde{G}_{\mu,p}^+) > x] = \mathbb{E} \left[\mathbb{P}[P_\alpha > x/h(\tilde{G}_{\mu,p}^+) \mid \tilde{G}_{\mu,p}^+] \right] = x^{-\alpha} \mathbb{E}[h(\tilde{G}_{\mu,p}^+)^{\alpha}],$$

with the conditional probability interpreted as zero when $h(\tilde{G}_{\mu,p}^+) = 0$. It remains to compare the angular constants. If $0 < \alpha \leq 1$, then

$$\left| C_h - \tilde{C}_{h,p}^+ \right| \leq L_h^\alpha \mathbb{E}[\|G - \tilde{G}_{\mu,p}^+\|_2^\alpha] \leq L_h^\alpha (\omega_{\mu,p}^+)^{\alpha}.$$

If $\alpha > 1$, the map $a \mapsto a^\alpha$ is Lipschitz on $[0, H_h]$ with constant $\alpha H_h^{\alpha-1}$, and hence

$$\left| C_h - \tilde{C}_{h,p}^+ \right| \leq \alpha H_h^{\alpha-1} L_h \mathbb{E}[\|G - \tilde{G}_{\mu,p}^+\|_2] \leq \alpha H_h^{\alpha-1} L_h \omega_{\mu,p}^+.$$

Together these bounds give

$$\left| C_h - \tilde{C}_{h,p}^+ \right| \leq K_{\alpha,h} (\omega_{\mu,p}^+)^{\gamma}.$$

The final display follows from the definition of $\eta_{r,h}$, the tail identity above, and $x^{-\alpha} \leq H_h^{-\alpha}$ for $x \geq H_h$. $\square$

The positive post-projection therefore solves the cone-membership problem without changing the radial model or the threshold errors ε_r and $\eta_{r,h}$. Its cost is explicit: the angular terms in the main simulation bounds are replaced by the combined budget $\omega_{\mu,p}^+$. It does not, however, remove the support mismatch in total variation. The simulator remains supported on a transformed low-dimensional set, so full-law approximation over all Borel events still requires either a correctly supported angular law or an additional residual model around the low-rank reconstruction.

S2.6 Empirical eigensolution and oracle-based consistency

Proof of Proposition 7. By Proposition 1,

$$\mathcal{R}_{\mu,\mathcal{I}}(W) = \frac{1}{|\mathcal{I}|} \sum_{i \in \mathcal{I}} \|u_\mu(g_i) - \Pi_W u_\mu(g_i)\|_2^2.$$

Expanding the squared norm and using symmetry and idempotence of Π_W gives

$$\mathcal{R}_{\mu,\mathcal{I}}(W) = \text{tr}(\Sigma_{\mu,\mathcal{I}}) - \text{tr}(\Pi_W \Sigma_{\mu,\mathcal{I}}) = \text{tr}((I - \Pi_W) \Sigma_{\mu,\mathcal{I}}).$$

The first term is fixed in W , so minimizing the empirical risk over p -dimensional subspaces is equivalent to maximizing $\text{tr}(\Pi_W \Sigma_{\mu,\mathcal{I}})$. The Ky Fan maximum principle gives a maximizer as the span of the leading p eigenvectors, and substituting that maximizer into the risk identity yields the

trailing eigensum. □

Proof of Theorem 2. The proof uses a deterministic-threshold sandwich. The random top- k_n set is trapped, with probability tending to one, between exceedance sets over two deterministic high thresholds whose expected sizes are slightly below and above k_n . Angular convergence for deterministic threshold exceedances can then be transferred to the random top- k_n selection.

We first prove this transfer at the level of empirical angular averages. Specifically, for every bounded continuous matrix-valued function h on $\mathbb{S}_+^{d-1}$, we show that

$$\frac{1}{k_n} \sum_{i \in \mathcal{I}_n^{(k_n)}} h(G_i) \rightarrow \mathbb{E}[h(G)] \quad \text{in probability.}$$

Taking $h(g) = u_\mu(g)u_\mu(g)^\top$, which is bounded and continuous because u_μ is linear and $g \in \mathbb{S}_+^{d-1}$, gives the consistency of $\widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}}$, and the remaining claims follow from standard eigenvalue and spectral-projector perturbation bounds.

Let B bound $\|h(g)\|_{\text{op}}$. For $0 < \varepsilon < 1$, choose deterministic thresholds u_n^- and u_n^+ such that

$$\mathbb{P}[R_1 > u_n^-] = (1 + \varepsilon)k_n/n, \quad \mathbb{P}[R_1 > u_n^+] = (1 - \varepsilon)k_n/n.$$

Because $k_n/n \rightarrow 0$, $u_n^\pm \rightarrow \infty$. Let $\mathcal{E}_n^\pm := \{i : R_i > u_n^\pm\}$. The binomial law of large numbers gives $|\mathcal{E}_n^-|/k_n \rightarrow 1 + \varepsilon$ and $|\mathcal{E}_n^+|/k_n \rightarrow 1 - \varepsilon$. With probability tending to one, $\mathcal{E}_n^+ \subset \mathcal{I}_n^{(k_n)} \subset \mathcal{E}_n^-$. By the angular convergence in Assumption 1 and a triangular-array law of large numbers for bounded variables,

$$\frac{1}{|\mathcal{E}_n^\pm|} \sum_{i \in \mathcal{E}_n^\pm} h(G_i) \rightarrow \mathbb{E}[h(G)] \quad \text{in probability.}$$

On the event $\mathcal{E}_n^+ \subset \mathcal{I}_n^{(k_n)} \subset \mathcal{E}_n^-$, write $A_I = k_n^{-1} \sum_{i \in \mathcal{I}_n^{(k_n)}} h(G_i)$ and $A_\pm = |\mathcal{E}_n^\pm|^{-1} \sum_{i \in \mathcal{E}_n^\pm} h(G_i)$.

Since $\|h\|_{\text{op}} \leq B$,

$$\|A_I - A_-\|_{\text{op}} \leq 2B \frac{|\mathcal{E}_n^-| - k_n}{|\mathcal{E}_n^-|}, \quad \|A_I - A_+\|_{\text{op}} \leq 2B \frac{k_n - |\mathcal{E}_n^+|}{k_n}.$$

The two fractions converge in probability to $\varepsilon/(1+\varepsilon)$ and ε , respectively. Hence the top- k_n average has all its subsequential probability limits within an $O(\varepsilon)$ neighborhood of $\mathbb{E}[h(G)]$. Letting $\varepsilon \downarrow 0$ proves the claim. $\square$

S2.7 Rank-based consistency

This subsection contains the rank-margin part of Section 3.2. Proposition S13 first transfers average angular accuracy and selected-set stability to covariance consistency. We then verify those two stability requirements from primitive assumptions and prove Theorem 3.

Proposition S13 (Stability under marginal standardization error). *With the oracle and rank-Pareto quantities in (3.2), suppose that*

$$\frac{1}{k_n} \sum_{i \in \mathcal{I}_n^{(k_n)} \cup \widehat{\mathcal{I}}_n^{(k_n)}} \left\| \widehat{G}_i - G_i \right\|_2 \rightarrow 0 \quad \text{in probability}$$

and $\left| \mathcal{I}_n^{(k_n)} \triangle \widehat{\mathcal{I}}_n^{(k_n)} \right| / k_n \rightarrow 0$ in probability, where $\triangle$ denotes symmetric difference. Then

$$\left\| \widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} \right\|_{\text{op}} \rightarrow 0 \quad \text{in probability.}$$

Hence, whenever the oracle top- k_n estimator is consistent, the rank-based estimator is consistent for the same Σ_μ .

Proof of Proposition S13. The map $g \mapsto u_\mu(g)$ is nonexpansive and $\|u_\mu(g)\|_2 \leq 1$. Hence

$$\left\| u_\mu(\widehat{g})u_\mu(\widehat{g})^\top - u_\mu(g)u_\mu(g)^\top \right\|_{\text{op}} \leq 2 \|\widehat{g} - g\|_2.$$

On the common selected set, the assumed average directional error gives

$$\frac{1}{k_n} \sum_{i \in \mathcal{I}_n^{(k_n)} \cap \widehat{\mathcal{I}}_n^{(k_n)}} \left\| u_\mu(\widehat{G}_i)u_\mu(\widehat{G}_i)^\top - u_\mu(G_i)u_\mu(G_i)^\top \right\|_{\text{op}} \leq \frac{2}{k_n} \sum_{i \in \mathcal{I}_n^{(k_n)} \cup \widehat{\mathcal{I}}_n^{(k_n)}} \left\| \widehat{G}_i - G_i \right\|_2 = o_p(1).$$

The symmetric difference of the selected sets contributes at most $2 \left| \mathcal{I}_n^{(k_n)} \Delta \widehat{\mathcal{I}}_n^{(k_n)} \right| / k_n$, because each summand has operator norm at most one. This proves the covariance transfer. $\square$

We now verify the two stability requirements under primitive assumptions. They are handled separately and under different assumptions. The average angular condition holds under continuity of the margins alone (Proposition S14); the argument is a weighted Glivenko–Cantelli ratio bound for the uniform empirical distribution function in the spirit of Wellner (1978) and Shorack and Wellner (1986, Chapter 10). The selected-set condition is where Assumption 2 enters, through an anti-concentration bound for the radii near the top- k_n threshold (Proposition S15).

Throughout this subsection the conditions of Theorem 3 are in force: the margins $F_1, \dots, F_d$ and the distribution of R_1 are continuous, Assumption 2 holds, $k_n \rightarrow \infty$, and $k_n/n \rightarrow 0$. Write $\bar{F}_R(r) := \mathbb{P}[R_1 > r]$. Under Assumption 2, $\bar{F}_R$ is regularly varying with index $-\alpha$; beyond the angular limit already used in Theorem 2, only this consequence of the assumption is used below. With the standard Pareto margins of Section 2.1 the index is necessarily $\alpha = 1$, since $R_1 \geq X_{11}$ gives $\bar{F}_R(r) \geq r^{-1}$ while $R_1 > r$ forces $\max_j X_{1j} > r/\sqrt{d}$ and hence $\bar{F}_R(r) \leq d^{3/2}r^{-1}$; we keep the general index α in the statements. Because the margins are continuous, the R_i are almost surely distinct, so $\mathcal{I}_n^{(k_n)}$ is almost surely unique. The $\widehat{R}_i$ take finitely many values and can tie with positive probability; fix any measurable tie-breaking rule under which $|\widehat{\mathcal{I}}_n^{(k_n)}| = k_n$ and $\widehat{\mathcal{I}}_n^{(k_n)}$ contains every index with $\widehat{R}_i$ strictly above its k_n -th largest value. The results below hold for every such rule. Write $\mathcal{U}_n := \mathcal{I}_n^{(k_n)} \cup \widehat{\mathcal{I}}_n^{(k_n)}$ for the union of the oracle and rank-based selected sets.

We use an exact representation of the rank transform. Fix a margin j and set $V_{ij} := 1 - F_j(Y_{ij})$, so that $V_{1j}, \dots, V_{nj}$ are i.i.d. standard uniform and $X_{ij} = 1/V_{ij}$, with large Y_{ij} corresponding to small V_{ij} . Let

$$\widehat{F}_{j,n}(v) := \frac{1}{n} \sum_{m=1}^n \mathbb{I}\{V_{mj} \leq v\}$$

be the empirical distribution function of these uniforms. Since ranks are almost surely unique,

$$n + 1 - \text{rank}(Y_{ij}) = 1 + \#\{m : Y_{mj} > Y_{ij}\} = 1 + \#\{m : V_{mj} < V_{ij}\} = n\widehat{F}_{j,n}(V_{ij}),$$

and therefore, exactly,

$$\frac{\widehat{X}_{ij}}{X_{ij}} = \frac{(n+1) V_{ij}}{n \widehat{F}_{j,n}(V_{ij})}. \quad (\text{S1})$$

The componentwise relative error of the rank transform is thus the ratio of a uniform empirical distribution function to the identity, evaluated at the observation's own uniform. The marginal step controls this ratio above a slowly shrinking level.

Lemma S2 (Ratio uniformity above a shrinking level). *Let $V_1, \dots, V_n$ be i.i.d. standard uniform with empirical distribution function $\widehat{F}_n$, and let $c_n \in (0, 1)$ satisfy $nc_n \rightarrow \infty$. Then*

$$\sup_{c_n \leq v \leq 1} \left| \frac{\widehat{F}_n(v)}{v} - 1 \right| \rightarrow 0 \quad \text{in probability.}$$

Proof. This is classical (Shorack and Wellner, 1986, Chapter 10); we include the short argument for completeness. Fix $\theta \in (0, 1)$ and put $v_\ell := c_n(1 + \theta)^\ell \wedge 1$ for $\ell = 0, 1, \dots, L_n$ with $L_n := \lceil \log(1/c_n) / \log(1 + \theta) \rceil$, so that the grid reaches 1. For $v \in [v_\ell, v_{\ell+1}]$, monotonicity gives

$$\frac{\widehat{F}_n(v)}{v} \leq \frac{\widehat{F}_n(v_{\ell+1})}{v_\ell} \leq (1 + \theta) \frac{\widehat{F}_n(v_{\ell+1})}{v_{\ell+1}}, \quad \frac{\widehat{F}_n(v)}{v} \geq \frac{1}{1 + \theta} \frac{\widehat{F}_n(v_\ell)}{v_\ell},$$

so if $|\widehat{F}_n(v_\ell)/v_\ell - 1| \leq \theta$ for every ℓ , the supremum in the statement is at most $(1 + \theta)^2 - 1 \leq 3\theta$. At a fixed v , $n\widehat{F}_n(v)$ is binomial with parameters n and v , and Bernstein's inequality gives, for $0 < \theta \leq 1$,

$$\mathbb{P} \left[\left| \widehat{F}_n(v) - v \right| > \theta v \right] \leq 2 \exp \left(- \frac{(\theta n v)^2}{2 n v + \frac{2}{3} \theta n v} \right) \leq 2 \exp \left(- \frac{3}{8} \theta^2 n v \right).$$

With $\beta_n := \frac{3}{8} \theta^2 n c_n \rightarrow \infty$ and $(1 + \theta)^\ell \geq 1 + \ell \theta$,

$$\sum_{\ell=0}^{L_n} \mathbb{P} \left[\left| \widehat{F}_n(v_\ell)/v_\ell - 1 \right| > \theta \right] \leq 2 \sum_{\ell \geq 0} e^{-\beta_n(1 + \ell \theta)} = \frac{2e^{-\beta_n}}{1 - e^{-\beta_n \theta}} \rightarrow 0.$$

Hence the supremum exceeds 3θ with probability tending to zero, for every $\theta \in (0, 1)$. $\square$

Fix the level $c_n := \sqrt{k_n}/n$, so that $nc_n = \sqrt{k_n} \rightarrow \infty$ and $nc_n = o(k_n)$, and define the bad set

$$J_n := \left\{ i \leq n : \min_{1 \leq j \leq d} V_{ij} < c_n \right\},$$

the observations having at least one coordinate among the roughly $\sqrt{k_n}$ most extreme of its margin.

Set

$$\varepsilon_n^* := \max_{1 \leq j \leq d} \sup_{c_n \leq v \leq 1} \left| \frac{n \hat{F}_{j,n}(v)}{(n+1)v} - 1 \right|.$$

Since $n\hat{F}_{j,n}(v)/((n+1)v) - 1 = \frac{n}{n+1}(\hat{F}_{j,n}(v)/v - 1) - \frac{1}{n+1}$, Lemma S2 applied to each margin and a union bound over the d margins give $\varepsilon_n^* \rightarrow 0$ in probability; each margin's uniforms are i.i.d., and no independence across margins is needed.

Lemma S3 (Componentwise accuracy off the bad set). *On the event $\{\varepsilon_n^* \leq \frac{1}{2}\}$, every $i \notin J_n$ satisfies*

$$\max_{1 \leq j \leq d} \left| \frac{\hat{X}_{ij}}{X_{ij}} - 1 \right| \leq 2\varepsilon_n^*, \quad \left| \hat{R}_i - R_i \right| \leq 2\varepsilon_n^* R_i, \quad \left\| \hat{G}_i - G_i \right\|_2 \leq 4\varepsilon_n^*.$$

Moreover $\mathbb{E} |J_n| \leq d\sqrt{k_n}$, so $|J_n|/k_n \rightarrow 0$ in probability.

Proof. For $i \notin J_n$ and every j , $V_{ij} \geq c_n$, so by the definition of ε_n^* the ratio $n\hat{F}_{j,n}(V_{ij})/((n+1)V_{ij})$ lies in $[1 - \varepsilon_n^*, 1 + \varepsilon_n^*]$. By (S1), $\hat{X}_{ij}/X_{ij}$ is the reciprocal of this ratio, so, using $\varepsilon_n^* \leq \frac{1}{2}$,

$$\left| \frac{\hat{X}_{ij}}{X_{ij}} - 1 \right| \leq \frac{\varepsilon_n^*}{1 - \varepsilon_n^*} \leq 2\varepsilon_n^*.$$

Consequently

$$\left\| \hat{X}_i - X_i \right\|_2^2 = \sum_{j=1}^d X_{ij}^2 \left(\frac{\hat{X}_{ij}}{X_{ij}} - 1 \right)^2 \leq (2\varepsilon_n^*)^2 R_i^2,$$

which gives $\left| \hat{R}_i - R_i \right| \leq \left\| \hat{X}_i - X_i \right\|_2 \leq 2\varepsilon_n^* R_i$ and, by the normalization inequality $\|a/\|a\|_2 - b/\|b\|_2\|_2 \leq 2\|a - b\|_2/\|a\|_2$, the bound $\left\| \hat{G}_i - G_i \right\|_2 \leq 4\varepsilon_n^*$. Finally $\mathbb{E} |J_n| \leq \sum_{j=1}^d n \mathbb{P}[V_{1j} < c_n] = dnc_n = d\sqrt{k_n}$, and Markov's inequality gives $|J_n|/k_n \rightarrow 0$ in probability since $\sqrt{k_n}/k_n \rightarrow 0$. $\square$

Proposition S14 (Average angular accuracy from continuity alone). *Let $F_1, \dots, F_d$ be continuous and $k_n \rightarrow \infty$, $k_n/n \rightarrow 0$. Then*

$$\frac{1}{k_n} \sum_{i \in \mathcal{U}_n} \left\| \hat{G}_i - G_i \right\|_2 \rightarrow 0 \quad \text{in probability.}$$

No assumption on the joint tail is used.

Proof. $|\mathcal{U}_n| \leq 2k_n$, and $\left\| \hat{G}_i - G_i \right\|_2 \leq 2$ always, being a difference of unit vectors. Splitting the sum over $\mathcal{U}_n$ at J_n and applying Lemma S3 on $\{\varepsilon_n^* \leq \frac{1}{2}\}$,

$$\frac{1}{k_n} \sum_{i \in \mathcal{U}_n} \left\| \hat{G}_i - G_i \right\|_2 \leq \frac{2|J_n|}{k_n} + \frac{2k_n}{k_n} \cdot 4\varepsilon_n^* = \frac{2|J_n|}{k_n} + 8\varepsilon_n^* \rightarrow 0 \quad \text{in probability.} \quad \square$$

The supremum analogue of the average angular condition fails. For the largest observation of margin j , $\text{rank}(Y_{ij}) = n$ gives $\hat{X}_{ij} = n + 1$, while X_{ij} is the reciprocal of the minimal uniform $V_{(1),j}$, and $nV_{(1),j}$ converges weakly to a standard exponential variable; hence $\hat{X}_{ij}/X_{ij}$ converges weakly to that exponential variable rather than to 1. The rank transform thus distorts the top few order statistics of each margin at constant order, and such observations are typically selected. Under the level $c_n = \sqrt{k_n}/n$ there are only $O_p(\sqrt{k_n})$ of them, which is why the averaged condition of Proposition S13 absorbs them while a uniform condition cannot. This is the reason Proposition S13 is stated in average form.

Lemma S4 (Threshold location and band count). *Define r_n by $\bar{F}_R(r_n) = k_n/n$, let $t_n := R_{n-k_n+1:n}$ be the k_n -th largest radius, and for $\delta \in (0, \frac{1}{2})$ let*

$$B_n(\delta) := \# \{i \leq n : t_n(1 - \delta) \leq R_i < t_n\}.$$

Then $t_n/r_n \rightarrow 1$ in probability, and for each fixed $\delta \in (0, \frac{1}{2})$,

$$\frac{B_n(\delta)}{k_n} \leq (1 - 2\delta)^{-\alpha} - 1 + o_p(1).$$

Proof. The first statement is the classical consistency of intermediate order statistics under regular variation; see [De Haan and Ferreira \(2006, Chapter 2\)](#). For the band count, let $N_n(r) := \#\{i \leq n : R_i > r\}$. For fixed $s > 0$, $\mathbb{E}[N_n(r_n s)]/k_n = \bar{F}_R(r_n s)/\bar{F}_R(r_n) \rightarrow s^{-\alpha}$ by the uniform convergence theorem for regularly varying functions ([Bingham et al., 1987](#), Theorem 1.5.2), and $\text{Var}[N_n(r_n s)] \leq \mathbb{E}[N_n(r_n s)]$, so Chebyshev's inequality gives $N_n(r_n s)/k_n \rightarrow s^{-\alpha}$ in probability. Since the distribution of R_1 is continuous, the R_i are almost surely distinct, so $\#\{i : R_i \geq t_n\} = k_n$. On the event $\{t_n \geq r_n(1 - \delta)\}$, whose probability tends to one by the first statement, every index counted by $B_n(\delta)$ has $R_i \geq t_n(1 - \delta) \geq r_n(1 - \delta)^2 \geq r_n(1 - 2\delta)$, and the same holds for the k_n indices with $R_i \geq t_n$; hence

$$B_n(\delta) \leq \#\{i : R_i \geq r_n(1 - 2\delta)\} - k_n = N_n(r_n(1 - 2\delta)) - k_n = k_n [(1 - 2\delta)^{-\alpha} - 1 + o_p(1)],$$

where the middle equality holds almost surely because the threshold $r_n(1 - 2\delta)$ is deterministic and the distribution of R_1 is continuous. $\square$

Proposition S15 (Selected-set stability under radial regular variation). *Under the conditions of Theorem 3, $|\mathcal{I}_n^{(k_n)} \triangle \widehat{\mathcal{I}}_n^{(k_n)}|/k_n \rightarrow 0$ in probability.*

Proof. Both sets have cardinality k_n , so $|\mathcal{I}_n^{(k_n)} \triangle \widehat{\mathcal{I}}_n^{(k_n)}| = 2|\mathcal{I}_n^{(k_n)} \setminus \widehat{\mathcal{I}}_n^{(k_n)}|$. Fix $\eta > 0$ and choose $\delta_0 \in (0, \frac{1}{2})$ with $2[(1 - 2\delta_0)^{-\alpha} - 1] < \eta/2$. Since $\varepsilon_n^* \rightarrow 0$ in probability, the event $\{4\varepsilon_n^* \leq \delta_0\}$ has probability tending to one; work on this event, on which in particular $\varepsilon_n^* \leq \frac{1}{2}$, so Lemma S3 applies.

Call i *good* if $i \notin J_n$.

Every good $i \in \mathcal{I}_n^{(k_n)}$ has $R_i \geq t_n$ and hence $\widehat{R}_i \geq (1 - 2\varepsilon_n^*)R_i \geq (1 - 2\varepsilon_n^*)t_n$. Therefore

$$|\mathcal{I}_n^{(k_n)} \setminus \widehat{\mathcal{I}}_n^{(k_n)}| \leq |J_n| + \#\left(\{i : \widehat{R}_i \geq (1 - 2\varepsilon_n^*)t_n\} \setminus \widehat{\mathcal{I}}_n^{(k_n)}\right).$$

Because $\widehat{\mathcal{I}}_n^{(k_n)}$ consists of k_n indices with the largest $\widehat{R}_i$, under any tie-breaking rule the number of indices at level $\geq (1 - 2\varepsilon_n^*)t_n$ that it fails to contain is at most $\widehat{N}_n - k_n$ when $\widehat{N}_n := \#\{i : \widehat{R}_i \geq$

$(1 - 2\varepsilon_n^*)t_n\}$ exceeds k_n , and zero otherwise. A good i with $\widehat{R}_i \geq (1 - 2\varepsilon_n^*)t_n$ has

$$R_i \geq \frac{\widehat{R}_i}{1 + 2\varepsilon_n^*} \geq t_n \frac{1 - 2\varepsilon_n^*}{1 + 2\varepsilon_n^*} \geq t_n(1 - 4\varepsilon_n^*),$$

so, using $\#\{i : R_i \geq t_n\} = k_n$ almost surely,

$$\widehat{N}_n \leq |J_n| + \#\{i : R_i \geq t_n(1 - 4\varepsilon_n^*)\} = |J_n| + k_n + B_n(4\varepsilon_n^*).$$

Combining the two displays,

$$\left| \mathcal{I}_n^{(k_n)} \triangle \widehat{\mathcal{I}}_n^{(k_n)} \right| \leq 2(2|J_n| + B_n(4\varepsilon_n^*)),$$

and on the working event $B_n(4\varepsilon_n^*) \leq B_n(\delta_0)$ by monotonicity of the band in its width. Lemma S4 and the choice of δ_0 give $\limsup_n \mathbb{P}[2B_n(4\varepsilon_n^*)/k_n > \eta/2] = 0$, while $4|J_n|/k_n \rightarrow 0$ in probability by Lemma S3. Hence $\mathbb{P}[\left| \mathcal{I}_n^{(k_n)} \triangle \widehat{\mathcal{I}}_n^{(k_n)} \right| / k_n > \eta] \rightarrow 0$. $\square$

Proof of Theorem 3. Propositions S14 and S15 verify the two conditions of Proposition S13, which yields $\left\| \widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} \right\|_{\text{op}} \rightarrow 0$ in probability. Since Assumption 2 implies Assumption 1, Theorem 2 gives $\left\| \widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} - \Sigma_\mu \right\|_{\text{op}} \rightarrow 0$ in probability, and the triangle inequality gives consistency of $\widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}}$. The spectral consequences follow from the same eigenvalue and spectral-projector perturbation bounds as in the proof of Theorem 2. $\square$

The radial step uses regular variation of $\bar{F}_R$ only through the vanishing band ratio

$$\lim_{\delta \downarrow 0} \limsup_{r \rightarrow \infty} \frac{\bar{F}_R(r(1 - \delta))}{\bar{F}_R(r)} = 1.$$

Under Assumption 1 alone, the standard Pareto margins give only the two-sided bound $r\bar{F}_R(r) \in [1, d^{3/2}]$, which permits $\bar{F}_R$ to fall by a fixed factor over relative bands of vanishing width; a fixed fraction of the top- k_n radii could then sit in such a band, and a rank perturbation of vanishing relative size could reshuffle a non-vanishing fraction of the selected set. We do not claim that

the selected-set condition fails under Assumption 1, only that its verification requires some radial anti-concentration, for which Assumption 2 is the natural primitive condition.

The proof gives more than $o_p(1)$: since $g \mapsto u_\mu(g)u_\mu(g)^\top$ is bounded by one in operator norm and 2-Lipschitz on $\mathbb{S}_+^{d-1}$, the bounds above combine to

$$\left\| \widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \widehat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} \right\|_{\text{op}} = O_p \left(\varepsilon_n^* + k_n^{-1/2} + \frac{B_n(4\varepsilon_n^*)}{k_n} \right),$$

with sharper weighted-approximation bounds for ε_n^* , of order $(nc_n)^{-1/2} = k_n^{-1/4}$ up to logarithmic factors, available from [Shorack and Wellner \(1986\)](#). We do not pursue rates here because the $k_n^{-1/2}$ scale is governed by the separate program of Section [S2.9](#).

S2.8 Oracle CLT and plug-in inference

Lemma S5 (Oracle tail empirical process). *Let $\mathcal{H}$ be the finite class consisting of the constant function $h_0(g) := 1$ and the coordinate functions of $h_\mu(g) := u_\mu(g)u_\mu(g)^\top$. For $x \in [1/2, 2]$, let $r_n(x)$ be a deterministic threshold with $\bar{F}_R(r_n(x)) = xk_n/n$, and define, for scalar $\psi \in \mathcal{H}$,*

$$\mathbb{G}_n(\psi, x) := \frac{1}{\sqrt{k_n}} \sum_{i=1}^n \{ \psi(G_i) \mathbb{I}\{R_i > r_n(x)\} - \mathbb{E}[\psi(G_1) \mathbb{I}\{R_1 > r_n(x)\}] \}.$$

Under the assumptions of Theorem 2, $\mathbb{G}_n \rightsquigarrow W$ in $\ell^\infty(\mathcal{H} \times [1/2, 2])$, where W is centered Gaussian with covariance

$$\text{Cov}[W(\psi, x), W(\varphi, x')] = \min(x, x') \mathbb{E}[\psi(G)\varphi(G)].$$

Proof. For fixed (ψ, x) and (φ, x') , put $y := \min(x, x')$. Since the two exceedance events intersect in $\{R_1 > r_n(y)\}$,

$$\frac{n}{k_n} \mathbb{E}[\psi(G_1)\varphi(G_1) \mathbb{I}\{R_1 > r_n(y)\}] = y \mathbb{E}[\psi(G_1)\varphi(G_1) \mid R_1 > r_n(y)] \rightarrow y \mathbb{E}[\psi(G)\varphi(G)].$$

The product of the centering terms is $O(k_n/n)$ after multiplication by n/k_n . Because all functions

in $\mathcal{H}$ are bounded, the row-wise Lindeberg condition is immediate, and the finite-dimensional convergence follows from the Lindeberg–Feller theorem and the Cramer–Wold device.

It remains only to control the index x . The class $\{(r, g) \mapsto \psi(g)\mathbb{I}\{r > r_n(x)\} : x \in [1/2, 2], \psi \in \mathcal{H}\}$ has a bounded envelope after the n/k_n row scaling and is generated by a finite class of bounded angular functions multiplied by a monotone one-dimensional threshold class. A regular grid in x gives brackets whose squared L_2 -size is proportional to the grid width, uniformly in n , and the envelope Lindeberg condition follows from $k_n/n \rightarrow 0$. The triangular-array bracketing central limit theorem therefore gives tightness. The limiting semimetric is proportional to $|x - x'|^{1/2}$ on each fixed angular coordinate, so the limit admits continuous sample paths in x . $\square$

Lemma S6 (Oracle random threshold expansion). *Let $R_{(1)} \geq \dots \geq R_{(n)}$ be the descending order statistics of $R_1, \dots, R_n$, and set $\hat{x}_n := \frac{n}{k_n} \bar{F}_R(R_{(k_n+1)})$. Under the assumptions of Theorem 2, $\hat{x}_n \rightarrow 1$ in probability and*

$$\sqrt{k_n}(\hat{x}_n - 1) = -\mathbb{G}_n(h_0, 1) + o_p(1).$$

Moreover, with probability tending to one, the oracle top- k_n set equals $\{i : R_i > r_n(\hat{x}_n)\}$.

Proof. The continuity of R_1 implies that the radii are almost surely distinct. Hence the set $\{i : R_i > R_{(k_n+1)}\}$ is exactly the top- k_n set. By the definition of $\hat{x}_n$, this set can be written as $\{i : R_i > r_n(\hat{x}_n)\}$, up to null events caused by flat parts of the survival function.

For fixed $\delta \in (0, 1/2)$, the event $\hat{x}_n > 1 + \delta$ is the event that at most k_n observations exceed the deterministic threshold $r_n(1 + \delta)$. The exceedance count is binomial with mean $(1 + \delta)k_n$, so Chebyshev’s inequality sends this probability to zero. The event $\hat{x}_n < 1 - \delta$ is handled in the same way. Thus $\hat{x}_n \rightarrow 1$ in probability. Let $N_n(x) := \sum_{i=1}^n \mathbb{I}\{R_i > r_n(x)\}$. On the event above, $N_n(\hat{x}_n) = k_n$, while $\mathbb{G}_n(h_0, x) = [N_n(x) - xk_n]/\sqrt{k_n}$. Therefore

$$\sqrt{k_n}(1 - \hat{x}_n) = \mathbb{G}_n(h_0, \hat{x}_n).$$

Lemma S5, its asymptotic equicontinuity in x , and $\hat{x}_n \rightarrow 1$ give $\mathbb{G}_n(h_0, \hat{x}_n) = \mathbb{G}_n(h_0, 1) + o_p(1)$, which proves the expansion. $\square$

Proof of Theorem 4. Let

$$S_n(x) := \frac{1}{k_n} \sum_{i=1}^n h_\mu(G_i) \mathbb{I}\{R_i > r_n(x)\}.$$

By Lemma S6, $\hat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} = S_n(\hat{x}_n)$ with probability tending to one. Also $\mathbb{E}[S_n(x)] = x H_\mu(r_n(x))$.

On $\{\hat{x}_n \in [1/2, 2]\}$,

$$\begin{aligned} \sqrt{k_n} (S_n(\hat{x}_n) - \Sigma_\mu) &= \mathbb{G}_n(h_\mu, \hat{x}_n) + \sqrt{k_n}(\hat{x}_n - 1)H_\mu(r_n(\hat{x}_n)) \\ &\quad + \sqrt{k_n}\{H_\mu(r_n(\hat{x}_n)) - \Sigma_\mu\}. \end{aligned}$$

Since $\bar{F}_R(r_n(\hat{x}_n)) \leq 2k_n/n$ with probability tending to one, the last term is bounded in operator norm by $\sqrt{k_n}\eta_\mu(2k_n/n) = o(1)$. The same bound gives $H_\mu(r_n(\hat{x}_n)) \rightarrow \Sigma_\mu$. Lemma S5 and asymptotic equicontinuity yield $\mathbb{G}_n(h_\mu, \hat{x}_n) = \mathbb{G}_n(h_\mu, 1) + o_p(1)$, while Lemma S6 gives

$$\sqrt{k_n}(\hat{x}_n - 1)H_\mu(r_n(\hat{x}_n)) = -\mathbb{G}_n(h_0, 1)\Sigma_\mu + o_p(1).$$

Thus, entrywise,

$$\sqrt{k_n} \left(\hat{\Sigma}_{\mu,n}^{(k_n),\text{orc}} - \Sigma_\mu \right) = \mathbb{G}_n(h_\mu, 1) - \mathbb{G}_n(h_0, 1)\Sigma_\mu + o_p(1) \rightsquigarrow Z.$$

It remains to identify the covariance. For symmetric A, B , write $W_A := \text{tr}(AW(h_\mu, 1))$.

Lemma S5 gives

$$\text{Cov}[W_A, W_B] = \mathbb{E}[(U_\mu^\top A U_\mu)(U_\mu^\top B U_\mu)], \quad \text{Cov}[W_A, W(h_0, 1)] = \mathbb{E}[U_\mu^\top A U_\mu] = \text{tr}(A\Sigma_\mu),$$

and $\text{Var}[W(h_0, 1)] = 1$. Therefore

$$\text{Cov}[\text{tr}(AZ), \text{tr}(BZ)] = \text{Cov}[U_\mu^\top A U_\mu, U_\mu^\top B U_\mu].$$

This is exactly the covariance of $\text{tr}(A(U_\mu U_\mu^\top - \Sigma_\mu))$ and $\text{tr}(B(U_\mu U_\mu^\top - \Sigma_\mu))$ for a single draw of U_μ , which gives the i.i.d. weak-limit representation. Since $U_\mu \in \mu^\perp$ almost surely, Z is supported on symmetric matrices acting on $\mu^\perp$. $\square$

Proof of Corollary 4. Write $E_n := \widehat{\Sigma}_{\mu,n}^{(k_n), \text{orc}} - \Sigma_\mu$. By Theorem 4 and fixed dimension, $\|E_n\|_{\text{op}} = O_p(k_n^{-1/2})$. The total anchored variation is linear:

$$\sqrt{k_n}(\widehat{\tau}_\mu^{\text{orc}} - \tau_\mu) = \text{tr}(\sqrt{k_n} E_n),$$

so the first limit follows from Theorem 4 with $A = I$, since $U_\mu^\top I U_\mu = \|U_\mu\|_2^2$.

For a simple eigenvalue λ_j , standard symmetric-matrix perturbation theory gives

$$\widehat{\lambda}_j^{\text{orc}} - \lambda_j = b_j^\top E_n b_j + O_p(\|E_n\|_{\text{op}}^2),$$

where the constant depends on the isolation gap around λ_j . Multiplying by $\sqrt{k_n}$ makes the remainder negligible, and the stated normal limit follows from Theorem 4 with $A = b_j b_j^\top$. Joint convergence over any fixed set of simple eigenvalues follows from joint convergence of the corresponding linear forms.

Assume now that $\Delta_{\mu,p} > 0$. Let $S_{\mu,p} := \sum_{j=1}^p \lambda_j$ and $\widehat{S}_{\mu,p}^{\text{orc}} := \sum_{j=1}^p \widehat{\lambda}_j^{\text{orc}}$. The map taking a symmetric matrix near Σ_μ to the sum of its leading p eigenvalues is differentiable at Σ_μ , with derivative $E \mapsto \text{tr}(P_{\mu,p} E)$. Hence

$$\sqrt{k_n}(\widehat{S}_{\mu,p}^{\text{orc}} - S_{\mu,p}) = \text{tr}(P_{\mu,p} \sqrt{k_n} E_n) + o_p(1).$$

Applying the delta method to $\text{AVE}_{\mu,p} = S_{\mu,p}/\tau_\mu$ gives the influence function

$$\tau_\mu^{-1} (U_\mu^\top P_{\mu,p} U_\mu - \text{AVE}_{\mu,p} \|U_\mu\|_2^2),$$

which proves the AVE limit and variance formula.

Finally, the spectral projector map is differentiable under the eigengap. Its derivative at Σ_μ in the direction of a symmetric matrix E is

$$\mathcal{D}_{\mu,p}(E) = \sum_{j=1}^p \sum_{m=p+1}^{d-1} \frac{b_m b_m^\top E b_j b_j^\top + b_j b_j^\top E b_m b_m^\top}{\lambda_j - \lambda_m}.$$

For completeness, this follows from the resolvent representation of the spectral projector and a first-order resolvent expansion around a contour separating $\{\lambda_1, \dots, \lambda_p\}$ from the rest of the spectrum. Applying the continuous mapping theorem to $\sqrt{k_n} E_n \rightsquigarrow Z$ gives the projector limit. $\square$

Proof of Corollary 5. The proof of Theorem 2 shows that oracle top- k_n averages of any fixed bounded continuous function of G_i converge in probability to the corresponding expectation under the limiting angular law. This applies to the functions needed for the first and second moments of

$$\psi_{\mu,p}(u) = \tau_\mu^{-1} (u^\top P_{\mu,p} u - \text{AVE}_{\mu,p} \|u\|_2^2).$$

By Theorem 2 and the eigengap assumption, $\widehat{\tau}_\mu^{\text{orc}} \rightarrow \tau_\mu$, $\widehat{\text{AVE}}_{\mu,p}^{\text{orc}} \rightarrow \text{AVE}_{\mu,p}$, and $\widehat{P}_{\mu,p}^{\text{orc}} \rightarrow P_{\mu,p}$ in probability. Since $\tau_\mu > 0$ and $\|u\|_2 \leq 1$, the map from (τ, AVE, P, u) to $\tau^{-1}(u^\top P u - \text{AVE} \|u\|_2^2)$ is uniformly Lipschitz on a neighborhood of the truth. The empirical first and second moments of $\widehat{\psi}_{\mu,p,i}^{\text{orc}}$ therefore converge to the corresponding moments of $\psi_{\mu,p}(U_\mu)$, and the empirical variance converges to $\sigma_{\mu,p}^2$. The confidence interval follows from Corollary 4 and Slutsky's theorem. The same argument applies to the plug-in variances for $\widehat{\tau}_\mu^{\text{orc}}$ and simple oracle eigenvalues. $\square$

Two boundary cases are worth recording. If the angular law is exactly rank p , so that $U_\mu \in W_{\mu,p}$ almost surely, then $P_{\mu,p} U_\mu = U_\mu$, $\text{AVE}_{\mu,p} = 1$, and the influence function in Corollary 4 is

identically zero. Hence $\sigma_{\mu,p}^2 = 0$, and the first-order normal interval for $\text{AVE}_{\mu,p}$ degenerates. Likewise, if $b_j^\top U_\mu = 0$ almost surely for an eigen-direction b_j , then the simple-eigenvalue variance $\text{Var}[(b_j^\top U_\mu)^2]$ is zero. These are boundary cases of the population angular law, not failures of the oracle CLT; they only say that the corresponding first-order fluctuation vanishes.

The same proof also gives explicit plug-in variance estimators for the other scalar limits in Corollary 4. If $\hat{v}_\tau^{\text{orc}}$ is the empirical variance of $\|U_{\mu,i}\|_2^2$ over $i \in \mathcal{I}_n^{(k_n)}$, then $\hat{v}_\tau^{\text{orc}} \rightarrow \text{Var}[\|U_\mu\|_2^2]$ in probability. If λ_j is simple and $\hat{v}_{\lambda_j}^{\text{orc}}$ is the empirical variance of $\{(\hat{b}_j^{\text{orc}})^\top U_{\mu,i}\}^2$ over $i \in \mathcal{I}_n^{(k_n)}$, then $\hat{v}_{\lambda_j}^{\text{orc}} \rightarrow \text{Var}[(b_j^\top U_\mu)^2]$ in probability. Joint covariances over any fixed set of simple eigenvalues are estimated analogously by the empirical covariances of these plug-in squared scores.

S2.9 A conditional rank-margin CLT

Rank-tail empirical-process expansions of this kind underlie rank-based inference for extreme-value copulas and stable tail-dependence functions (e.g., [Genest and Segers, 2009](#); [Einmahl et al., 2012](#)). The following result states the corresponding AGCA consequence in a high-level form. It isolates the part specific to AGCA: once the rank-Pareto tail empirical measure has a Gaussian expansion and the induced top- k_n angular functional is differentiable, the same spectral delta method used for the oracle CLT applies.

Let $E_+ := \mathbb{R}_+^d \setminus \{0\}$, $s(x) := x/\|x\|_2$, and $\varphi_\mu(x) := u_\mu(s(x))u_\mu(s(x))^\top$. For $q > 0$, write $A_q := \{x \in E_+ : \|x\|_2 > q\}$. If m is a Radon measure on E_+ and a threshold $q(m)$ can be chosen so that $m(A_{q(m)}) = 1$, define

$$\Phi_\mu(m) := \int_{A_{q(m)}} \varphi_\mu(x) dm(x).$$

Theorem S1 (Conditional rank-margin CLT). *Assume Assumption 2. Let $a_n := n/k_n$ and define the rank-tail empirical measure*

$$\hat{\nu}_n^{\text{rank}} := \frac{1}{k_n} \sum_{i=1}^n \delta_{\hat{X}_i/a_n}.$$

Suppose that there exist a Radon tail measure ν , a normed linear space $\mathbb{D}$ of signed measures, a

linear subspace $\mathbb{D}_0 \subset \mathbb{D}$, and a centered Gaussian element $\mathbb{W}_\nu^{\text{rank}} \in \mathbb{D}$, with $\mathbb{W}_\nu^{\text{rank}} \in \mathbb{D}_0$ almost surely, such that

$$\sqrt{k_n} (\hat{\nu}_n^{\text{rank}} - \nu) \rightsquigarrow \mathbb{W}_\nu^{\text{rank}} \quad \text{in } \mathbb{D}.$$

Assume further that $q(\hat{\nu}_n^{\text{rank}})$ can be chosen so that $A_{q(\hat{\nu}_n^{\text{rank}})}$ contains the rank-Pareto top- k_n observations with probability tending to one, that $q(\nu)$ satisfies $\nu(A_{q(\nu)}) = 1$ and $\nu(\partial A_{q(\nu)}) = 0$, and that Φ_μ is Hadamard differentiable at ν tangentially to $\mathbb{D}_0$, with continuous linear derivative $\dot{\Phi}_{\mu,\nu}$.

Then

$$\sqrt{k_n} (\hat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \Sigma_\mu) \rightsquigarrow Z_\mu^{\text{rank}} := \dot{\Phi}_{\mu,\nu}(\mathbb{W}_\nu^{\text{rank}}),$$

where Z_μ^{rank} is a centered Gaussian symmetric matrix. Consequently, if $1 \leq p \leq d-2$, $\tau_\mu > 0$, and $\Delta_{\mu,p} > 0$, then $\widehat{\text{AVE}}_{\mu,p}$ and $\hat{P}_{\mu,p}$, formed from $\hat{\Sigma}_{\mu,n}^{(k_n),\text{emp}}$, satisfy

$$\sqrt{k_n} (\widehat{\text{AVE}}_{\mu,p} - \text{AVE}_{\mu,p}) \rightsquigarrow \tau_\mu^{-1} \{ \text{tr}(P_{\mu,p} Z_\mu^{\text{rank}}) - \text{AVE}_{\mu,p} \text{tr}(Z_\mu^{\text{rank}}) \},$$

and

$$\sqrt{k_n} (\hat{P}_{\mu,p} - P_{\mu,p}) \rightsquigarrow \mathcal{D}_{\mu,p}(Z_\mu^{\text{rank}}),$$

with $\mathcal{D}_{\mu,p}$ as in Corollary 4. The same replacement of Z by Z_μ^{rank} gives the limits for total anchored variation and simple eigenvalues.

The theorem makes two high-level assumptions. For tail empirical measures with estimated Pareto margins, Gaussian expansions of the assumed type have been studied in the literature (Einmahl et al., 1997, 2001; Einmahl and Segers, 2009). The expansion assumption should be attainable along that route under suitable second-order marginal and joint-tail conditions. The differentiability assumption is more delicate: when the limiting angular law places mass on the axes, Φ_μ can respond non-smoothly to componentwise marginal scalings. The two assumptions together take care of the face and near-axis regimes that the bounded departures u_μ accommodate at the oracle level.

Proof. Put $\hat{m}_n := \hat{\nu}_n^{\text{rank}}$. By the assumed choice of $q(\hat{m}_n)$, the set $A_{q(\hat{m}_n)}$ contains exactly the

observations selected by the rank-Pareto top- k_n rule, with probability tending to one. Therefore

$$\Phi_\mu(\widehat{m}_n) = \frac{1}{k_n} \sum_{i \in \widehat{\mathcal{I}}_n^{(k_n)}} u_\mu(\widehat{G}_i) u_\mu(\widehat{G}_i)^\top = \widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}}$$

with probability tending to one. Under Assumption 2, the limiting tail measure has a polar representation whose conditional angular law on any radial exceedance set A_q is the angular law of G . Since $\nu(A_{q(\nu)}) = 1$, this gives

$$\Phi_\mu(\nu) = \int_{A_{q(\nu)}} \varphi_\mu(x) d\nu(x) = \mathbb{E}[u_\mu(G) u_\mu(G)^\top] = \Sigma_\mu.$$

The functional delta method applied to the assumed weak convergence of $\widehat{m}_n$ and to the Hadamard differentiability of Φ_μ gives

$$\sqrt{k_n} (\Phi_\mu(\widehat{m}_n) - \Phi_\mu(\nu)) \rightsquigarrow \dot{\Phi}_{\mu,\nu}(\mathbb{W}_\nu^{\text{rank}}).$$

The first two displayed identities identify the left-hand side with $\sqrt{k_n}(\widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \Sigma_\mu)$, except on an event whose probability tends to zero. Because $\dot{\Phi}_{\mu,\nu}$ is continuous and linear and $\mathbb{W}_\nu^{\text{rank}}$ is centered Gaussian, the limit Z_μ^{rank} is centered Gaussian.

The scalar and spectral conclusions are the same perturbation steps used in the proof of Corollary 4. The trace map is linear, simple eigenvalues are first-order differentiable with derivative $E \mapsto b_j^\top E b_j$, the explained-variation map has derivative

$$E \mapsto \tau_\mu^{-1} \{ \text{tr}(P_{\mu,p} E) - \text{AVE}_{\mu,p} \text{tr}(E) \},$$

and the rank- p spectral projector has derivative $\mathcal{D}_{\mu,p}$ under $\Delta_{\mu,p} > 0$. Applying the continuous mapping theorem to $\sqrt{k_n}(\widehat{\Sigma}_{\mu,n}^{(k_n),\text{emp}} - \Sigma_\mu) \rightsquigarrow Z_\mu^{\text{rank}}$ proves all displayed limits. $\square$

S3 Simulation analysis

This supplementary section provides a simulation analysis used to illustrate AGCA. To make the geometry visible, we begin with two three-dimensional designs. Model 1 is a low-dimensional angular law whose selected directions form a slightly bent cloud on the positive sphere. Model 2 adds a variable-specific near-axis regime, so that one coordinate can dominate the angular direction while remaining a finite point on $\mathbb{S}_+^2$. We then add a ten-dimensional design in which eight variables share a lower-rank angular mechanism and two variables have asymptotically independent near-axis regimes. The diagnostics below focus on visualization, explained variation, scores, loadings, anchor choice, bootstrap stability, finite-sample sensitivity, and robustness of the canonical-anchor summary beyond the visible three-dimensional case. For the ten-dimensional design, we also include a repeated-sampling coverage check for the oracle plug-in intervals, comparing the formal oracle-margin setting with the same interval formula applied after rank-Pareto marginal standardization.

Throughout, the canonical anchor is $\mu_0 = d^{-1/2}\mathbf{1}_d$ in the relevant dimension. We compare it with the empirical Fréchet and principal anchors in (2.2), computed from the selected directions. Bootstrap 95% diagnostic intervals (DIs) are conditional summaries: after fixing the selected angular directions, we resample those directions with replacement, refit AGCA at the same rank and anchor, and align loading signs to the main-sample loadings.

S3.1 Data-generating mechanisms

Let $Y_{\theta,q}$ be a q -variate logistic max-stable vector with unit Fréchet margins,

$$\mathbb{P}[Y_{\theta,q,1} \leq y_1, \dots, Y_{\theta,q,q} \leq y_q] = \exp \left[- \left(\sum_{j=1}^q y_j^{-1/\theta} \right)^\theta \right], \quad y_j > 0, \quad 0 < \theta < 1. \quad (\text{S1})$$

The corresponding logistic-Pareto block is

$$Z_{\theta,q,j} := (1 - \exp(-1/Y_{\theta,q,j}))^{-1}, \quad j = 1, \dots, q, \quad (\text{S2})$$

Design	n, k	Extremal mechanism	Near-axis part
Model 1	2400, 120	Trivariate logistic-Pareto block embedded through three positive rays. Two rays generate the dominant X_1 -versus- X_2 contrast; a weaker third ray bends the cloud through X_3 .	None by design.
Model 2	2400, 120	Bivariate logistic-Pareto block for X_1, X_2 , plus an independent Pareto source for X_3 .	X_3 near-axis regime.
10D model	10000, 500	Five-ray logistic-Pareto block on $X_1, \dots, X_8$ lying in a two-dimensional anchored surface, plus independent Pareto sources for X_9 and X_{10} . Margins are rank-Pareto standardized before thresholding.	X_9 and X_{10} near-axis regimes.

Table S1: Simulation designs. Selected extremes are the k largest Euclidean radii after the stated preprocessing, with directions normalized to the positive sphere.

so each margin is standard Pareto. For this specific model, $\eta_\mu(t) = O(t^\kappa)$ with $\kappa = \min\{1, (1 - \theta)/\theta\}$. Therefore, Assumption 3 holds whenever $k_n = o(n^{2\kappa/(2\kappa+1)})$; for $\theta \leq 1/2$ this is $k_n = o(n^{2/3})$, up to a logarithmic refinement at $\theta = 1/2$.

Let P denote an independent standard Pareto variable and let E_i collect independent standard exponential noise variables. The light-tailed noise keeps finite samples nonsingular, but is asymptotically negligible relative to the Pareto signals.

For the two three-dimensional designs, thresholding and angular normalization are applied directly to the generated vectors X_i ; only the ten-dimensional design uses the rank-Pareto preprocessing in (S6).

For Model 1, with $\theta_1 = 0.50$ and $\tau_1 = 0.30$,

$$X_i = A_1 Z_i + \tau_1 E_i, \quad Z_i = (Z_{i1}, Z_{i2}, Z_{i3})^\top \sim Z_{\theta_1, 3}. \quad (\text{S3})$$

Let

$$b_1 = (1, -1, 0)^\top / \sqrt{2}, \quad b_2 = (1, 1, -2)^\top / \sqrt{6}, \quad \mu_3 = 3^{-1/2} \mathbf{1}_3.$$

With $\rho = 0.25$, $\gamma = (1 + \rho)/2$, $\eta = 0.45$, and $\kappa = 0.45$, the ray matrix is

$$A_1 = (a_1, a_2, a_3), \quad a_1 = \begin{pmatrix} 1 \\ \rho \\ \gamma \end{pmatrix}, \quad a_2 = \begin{pmatrix} \rho \\ 1 \\ \gamma \end{pmatrix}, \quad a_3 = \kappa(\mu_3 - \eta b_2).$$

The first two rays create the dominant b_1 contrast and sit slightly below the canonical anchor in the X_3 direction. The third ray has larger relative X_3 participation, so the selected angular cloud is bent rather than horizontal, with μ_0 visually inside the main cloud.

For Model 2, with $\theta_2 = 0.25$ and $\tau_2 = 0.30$,

$$X_i = (Z_{i1}, Z_{i2}, P_{i3})^\top + \tau_2 E_i, \quad Z_i = (Z_{i1}, Z_{i2})^\top \sim Z_{\theta_2, 2}. \quad (\text{S4})$$

The independent Pareto source P_{i3} creates the variable-specific near-axis regime. In the displayed realization, 72 selected extremes are shared X_1 - X_2 episodes and 48 are X_3 -axis episodes.

For the ten-dimensional model, with $\theta_{10} = 0.45$ and $\tau_{10} = 0.25$, let $\mu_8 = 8^{-1/2} \mathbf{1}_8$ and

$$h_1 = \left(-1, -\frac{5}{7}, -\frac{3}{7}, -\frac{1}{7}, \frac{1}{7}, \frac{3}{7}, \frac{5}{7}, 1 \right)^\top, \quad h_2 = (1, 1, 1, 1, -1, -1, -1, -1)^\top.$$

Let $B = (b_1, b_2)$ be the Gram–Schmidt orthonormalization of (h_1, h_2) . For the rows $s_\ell^\top$, $\ell = 1, \dots, 5$, of

$$S = \begin{pmatrix} -0.42 & -0.30 & 0.00 & 0.28 & 0.48 \\ -0.26 & 0.30 & -0.36 & 0.24 & -0.02 \end{pmatrix}^\top,$$

define shared rays $a_\ell := (\mu_8 + B s_\ell) / \|\mu_8 + B s_\ell\|_2 \in \mathbb{S}_+^7$, $\ell = 1, \dots, 5$, and $A_8 = (a_1, \dots, a_5)$.

With $Z_i = (Z_{i1}, \dots, Z_{i5})^\top \sim Z_{\theta_{10}, 5}$, the raw vector is

$$X_i^* = (A_8 Z_i, P_{i9}, P_{i,10}) + \tau_{10} E_i \in \mathbb{R}^{10}. \quad (\text{S5})$$

The shared $X_1, \dots, X_8$ mechanism lies in a two-dimensional anchored surface before normalization,




Figure S1: Sphere-level geometry in the two three-dimensional simulations. The left panel shows Model 1 and the right panel shows Model 2. Each panel displays selected extreme directions, the canonical anchor and first AGC, and the Fréchet anchor and first AGC.

whereas P_{i9} and $P_{i,10}$ create two variable-specific near-axis regimes. As in empirical work with unknown margins, the AGCA fit is applied after the marginal rank-Pareto transform

$$\tilde{X}_{ij} = \frac{n+1}{n+1 - \text{rank}(X_{ij}^*, X_{1j}^*, \dots, X_{nj}^*)}, \quad j = 1, \dots, 10, \quad (\text{S6})$$

where ranks are increasing within each margin. The selected extremes are the $k = 500$ largest radii of the transformed vectors $\tilde{X}_i$.

S3.2 Sphere geometry

Figure S1 shows the geometric object fitted by AGCA. The points are the selected extreme directions. The canonical fit uses the fixed balanced anchor; the Fréchet fit recenters the geometry at the sample spherical center. Only the first AGC is shown for each anchor. In Model 1, the selected directions form a bent but still essentially one-dimensional angular cloud. The canonical and Fréchet anchors are close, and their first AGCs trace similar directions through the cloud. In Model 2, the first AGC runs from the balanced interior toward the X_3 -axis part of the sphere. This is the geometric sense in which AGCA handles asymptotic independence: a near-axis episode is a large but bounded anchored displacement, not a point outside the coordinate system.




Figure S2: Anchored variation explained in the three-dimensional simulations. Each panel compares the canonical, Fréchet, and principal anchors.

S3.3 Explained variation and scores

Figure S2 reports cumulative anchored variation explained. In Model 1, the first canonical AGC explains 82.0% of anchored variation and the rank-one residual risk is 0.0085. The lower percentage relative to an exactly rank-one construction is intentional: the cloud is bent by the third ray, but the leading direction is still sharply identified, with absolute alignment 0.999996 with the designed b_1 contrast. In Model 2, the first canonical AGC explains 93.3% of anchored variation and has absolute alignment 0.999977 with the X_3 -axis contrast.

Figure S3 gives the score representation of the same geometry. In Model 1, the selected extremes spread primarily along AGC 1, with a smaller second-coordinate spread caused by the X_3 bend. In Model 2, the first score separates shared X_1 - X_2 episodes from X_3 -axis episodes, while the second score records residual variation inside the shared block.

S3.4 Loadings

Figure S4 shows the canonical-anchor loadings. In Model 1, AGC 1 is the X_1 -versus- X_2 contrast, while AGC 2 captures the weaker X_3 bending direction. In Model 2, AGC 1 is positive for X_3 and negative for X_1, X_2 ; it separates the variable-specific near-axis extremes from the shared block.




Figure S3: Canonical-anchor scores for the selected extremes.




Figure S4: Canonical-anchor loadings. From left to right: Model 1 AGC 1, Model 1 AGC 2, Model 2 AGC 1, and Model 2 AGC 2. Bars show sample loadings, vertical intervals show bootstrap 95% diagnostic intervals (DIs), and dots show population loadings matched to the sample AGCs.

AGC 2 then describes the residual X_1 -versus- X_2 contrast. The bootstrap diagnostic intervals (DIs) are narrow relative to the leading loading magnitudes, so the displayed contrasts are stable in the selected samples.

S3.5 Anchor choice

The canonical anchor fixes the interpretation as departure from balanced complete dependence. Data-adaptive anchors can give similar scalar fits, but they change the zero point of the loading and score coordinates. Figure S5 reports the geodesic distances from the canonical anchor. In Model 1, the Fréchet and principal anchors are both about 0.078 radians from μ_0 . In Model 2, the Fréchet




Figure S5: Anchor sensitivity distances. Distances are spherical geodesic distances from the canonical anchor.




Figure S6: Fréchet-anchor loadings. From left to right: Model 1 AGC 1, Model 1 AGC 2, Model 2 AGC 1, and Model 2 AGC 2. Bars show sample loadings, vertical intervals show bootstrap 95% diagnostic intervals (DIs), and dots show population loadings matched to the sample AGCs under the Fréchet anchor.

anchor remains close (0.029), while the principal anchor moves farther (0.296) because the leading angular direction is pulled toward the X_3 axis. The sphere-level Fréchet fit is already displayed in Figure S1; Figure S6 gives the corresponding Fréchet-anchor loadings. The dominant directions are comparable to the canonical directions, but their interpretation is sample-centered rather than benchmark-relative. This distinction is the main reason the canonical anchor remains the default in the paper.




Figure S7: Finite-sample recovery diagnostics for the two three-dimensional simulations. Curves report medians with 10–90% bands across repeated samples at a fixed tail fraction.

S3.6 Finite-sample sensitivity

Figures S7 and S8 repeat both three-dimensional designs over increasing n , keeping the tail fraction fixed. The recovery figure contains two diagnostics. The loading-alignment curves compare the estimated leading loading with the known target contrast: the X_1 -versus- X_2 contrast in Model 1 and the X_3 -axis contrast in Model 2. The projector-distance curves compare the estimated leading one-dimensional AGCA space with the corresponding target space, so smaller values mean better subspace recovery. In both designs the median alignment is already high at the displayed sample sizes and the projector distance decreases as n grows; Model 2 is slightly more variable because the selected sample mixes a shared X_1 - X_2 regime with a separate near-axis X_3 regime.

Figure S8 reports the scalar fit diagnostics along the same finite-sample path. The anchored-variation curves show the fraction of total anchored variation explained by the leading AGC, while the residual-risk curves show the average squared anchored-coordinate error left after the rank-one




Figure S8: Finite-sample anchored variation and residual-risk diagnostics for the two three-dimensional simulations. Curves report medians with 10–90% bands across repeated samples at a fixed tail fraction.

fit. These quantities stabilize near the main-sample values used in the preceding displays: Model 1 has a deliberately nonzero residual because the angular cloud is slightly bent, whereas Model 2 has high rank-one explained variation because the dominant contrast separates shared X_1 - X_2 episodes from the X_3 -axis episodes. The bands shrink with n , confirming that the visual conclusions are not artifacts of one favorable selected sample.

S3.7 Ten-dimensional design

The ten-dimensional design checks that the preceding messages are not artifacts of a visible three-dimensional sphere. Variables $X_1, \dots, X_8$ share a lower-rank angular mechanism, while X_9 and X_{10} have asymptotically independent near-axis regimes. After the rank-Pareto transformation, the first two AGCs are expected to capture the two large variable-specific regimes, and the next components should recover the lower-amplitude shared variation among $X_1, \dots, X_8$. Figure S9




Figure S9: Ten-dimensional anchor diagnostics. Panel (A) compares cumulative anchored variation explained under the canonical, Fréchet, and principal anchors. Panel (B) reports the spherical distance from the canonical anchor to the data-adaptive anchors.

shows that the canonical, Fréchet, and principal anchors give qualitatively the same anchored-variation summary: the first few components dominate, and the ordering of the low-rank fit is stable across anchors. The data-driven anchors therefore do not overturn the canonical summary, even though two variables have asymptotically independent tails. The canonical anchor remains preferable for interpretation because its loadings are departures from balanced complete dependence rather than from an empirical center.

Figures S10 and S11 give the loading interpretation. The heatmap shows the leading canonical-anchor loadings: the first two components concentrate on X_9 and X_{10} , which are the designed near-axis regimes, while the following components show structured contrasts across $X_1, \dots, X_8$. The separate AGC 3 and AGC 4 bar plots make this lower-amplitude shared structure explicit. Thus AGCA does not ignore the shared block; it orders it after the larger near-axis departures because components are ranked by anchored variation.

Figure S12 reports the oracle alignment matrix. The target columns are known only because this is a simulation. The first two sample AGCs align with the two-dimensional axis-regime subspace generated by X_9 and X_{10} , up to rotation within that subspace. The third and fourth sample AGCs




Figure S10: Ten-dimensional canonical-anchor loading heatmap. The first two components concentrate on the designed near-axis regimes, while later components recover structured contrasts in the shared block $X_1, \dots, X_8$.




Figure S11: Ten-dimensional canonical-anchor bar loadings for AGC 3 and AGC 4. Bars show sample loadings, vertical intervals show bootstrap 95% diagnostic intervals (DIs), and dots show population loadings matched to the sample AGCs. These components recover the lower-amplitude shared angular variation among $X_1, \dots, X_8$.




Figure S12: Oracle alignment for the ten-dimensional simulation. Rows correspond to sample AGCs and columns to known design contrasts.




Figure S13: Ten-dimensional stability diagnostics. Panel (A) varies the number of selected radial extremes around the main threshold. Panel (B) shows bootstrap 95% diagnostic intervals (DIs) for canonical-anchor variation explained.

align with the two designed shared AGCs for $X_1, \dots, X_8$. The threshold and bootstrap diagnostics in Figure S13 then show that this interpretation is not driven by a narrow threshold choice: the explained-variation path and the rank-two residual risk are stable around the main threshold, and bootstrap DIs for cumulative variation remain narrow.

Margins	Target	Population value	Coverage	Bias	Mean length
Oracle	τ_μ	0.4637	0.941	0.0028	0.0577
Rank	τ_μ	0.4637	1.000	0.0005	0.0576
Oracle	λ_1	0.2576	0.969	0.0029	0.0266
Rank	λ_1	0.2576	1.000	-0.0003	0.0211
Oracle	λ_2	0.1942	0.952	0.0001	0.0423
Rank	λ_2	0.1942	1.000	-0.0000	0.0407
Oracle	$AVE_{\mu,2}$	0.9742	0.931	0.0005	0.0099
Rank	$AVE_{\mu,2}$	0.9742	0.878	-0.0017	0.0107

Table S2: Monte Carlo coverage of nominal 95% oracle plug-in intervals in the ten-dimensional design. Bias is the Monte Carlo mean estimate minus the population target. In the rank-margin rows, each sample is first rank-Pareto standardized and then the oracle plug-in standard error is applied deliberately as a shortcut.

S3.8 Oracle interval coverage in the ten-dimensional design

We also use the ten-dimensional design to check the finite-sample behavior of the oracle confidence intervals from Corollary 4 and Corollary 5 in the main text. The goal is not to replace the bootstrap diagnostics above, but to verify that the bounded-departure oracle variance formulas give reasonable repeated-sampling calibration in the same setting used for the geometric diagnostics. We run 5000 independent samples with $n = 10000$ and $k = 500$, using the canonical anchor and rank $p = 2$. The population targets are estimated once from an independent sample of size 500000, with the same tail fraction $k/n = 0.05$. Since the exact finite-sample marginal distributions of the noisy ten-dimensional generator are not available in closed form, the oracle-margin analysis uses an independent calibration sample of size 500000 to approximate the true Pareto marginal transforms, with Pareto tail extrapolation beyond the calibration tail. For comparison, we apply the same oracle interval formulas after the rank-Pareto marginal standardization used in the main simulation analysis. This second comparison is diagnostic: as discussed in Section 3.4, the oracle variance formula is not claimed to be a full rank-margin CLT.

Table S2 shows that the oracle intervals are reasonably calibrated for the formal oracle problem. Coverage is close to the nominal level for λ_2 , slightly conservative for λ_1 , and mildly below nominal for τ_μ and $AVE_{\mu,2}$. The biases are small relative to the interval lengths, so the remaining discrepancies are consistent with finite-threshold and plug-in variance effects rather than a breakdown of the bounded-departure CLT. The rank-margin comparison is informative but should be read

differently. For τ_μ , λ_1 , and λ_2 , the oracle formula applied to rank margins overcovers: the rank transformation fixes much of the marginal tail-scale variation across samples, while the oracle plug-in standard errors remain comparatively large. For $AVE_{\mu,2}$, however, the same shortcut undercovers, with coverage 0.878. This ratio functional is sensitive to the joint rank-margin correction in the numerator and denominator, and the result illustrates why the paper uses bootstrap and stability diagnostics for rank-standardized empirical analyses. At the same time, the oracle-margin results support the main inferential message: for this high-dimensional design with near-axis regimes, AGCA’s bounded spherical departures lead to usable first-order uncertainty quantification for the oracle spectral summaries.

Together, the simulations isolate the core geometric messages. In Model 1, AGCA summarizes a smooth, slightly bent angular law by a stable anchored direction plus a smaller residual direction. In Model 2, AGCA summarizes an asymptotically independent variable by a finite near-axis displacement from the balanced anchor. The same diagnostics used in the empirical application (explained variation, scores, loadings, anchor checks, bootstrap DIs, and threshold or finite-sample paths) are already visible in this low-dimensional setting and remain informative in the ten-dimensional design with two near-axis regimes.

S4 Portfolio empirical analysis: additional details

This supplementary section gives the data-construction details and secondary diagnostics for Section 4. All portfolio-return panels are converted to losses by multiplying returns by -1 . Each margin is then transformed to the pseudo-Pareto scale using (3.1), and AGCA is fit to Euclidean-spherical directions for the largest radii. Unless otherwise stated, the anchor is the canonical complete-dependence direction $\mu_0 = d^{-1/2}\mathbf{1}$.

S4.1 Data construction and anomaly-panel robustness design

The main Fama–French panel uses daily value-weighted 2×3 portfolios from the Kenneth R. French Data Library (French, 2026). The four included sorts are Size–Book-to-Market, Size–Operating Profitability, Size–Investment, and Size–Momentum. Each sort contributes six portfolios, corresponding to two size groups crossed with three characteristic groups. The retained complete panel has $d = 24$ assets and $n = 13340$ daily observations from July 2, 1973 through May 29, 2026. The main threshold keeps the largest $k = 667$ radii.

The robustness panel uses the Open Source Asset Pricing daily value-weighted anomaly-sorted quintile portfolio file (Chen and Zimmermann, 2026). We use six continuous signals related to liquidity and trading frictions: bid–ask spread, illiquidity, dollar volume, one-month zero trading, six-month zero trading, and price. Each signal contributes five quintile portfolios, giving $d = 30$ assets. The retained complete panel has $n = 12988$ daily observations from July 2, 1973 through December 31, 2024. The main threshold keeps the largest $k = 649$ radii.

Table S3: Portfolio blocks in the two empirical panels.

Panel	Block	Portfolios retained
FF	Size–Book-to-Market	six 2×3 value-weighted portfolios
FF	Size–Operating Profitability	six 2×3 value-weighted portfolios
FF	Size–Investment	six 2×3 value-weighted portfolios
FF	Size–Momentum	six 2×3 value-weighted portfolios
OSAP	Bid–ask spread	five value-weighted quintile portfolios
OSAP	Illiquidity	five value-weighted quintile portfolios
OSAP	Dollar volume	five value-weighted quintile portfolios
OSAP	Zero trading, one month	five value-weighted quintile portfolios
OSAP	Zero trading, six months	five value-weighted quintile portfolios
OSAP	Price	five value-weighted quintile portfolios

The Fama–French panel is the main application because it gives a compact and familiar portfolio-formation universe with clear economic blocks. The OSAP panel is deliberately different: it is a modern anomaly-sorted trading-friction panel, distinct from a standard size-characteristic panel. Its role is to test whether the portfolio-functional conclusions survive in another long daily portfolio universe. The robustness evidence is reported below in the cross-panel spectrum, OSAP loading

heatmap, and OSAP portfolio-functional diagnostics.

S4.2 AGCA spectrum diagnostics

Figure S14 reports the main canonical-anchor spectra for both portfolio panels. It is included here as the cross-panel version of the main-text Fama–French spectrum: the Fama–French panel is more compressed, while the OSAP anomaly panel gives the same qualitative low-rank message in a distinct portfolio universe.




Figure S14: Cumulative anchored variation explained for the Fama–French and OSAP portfolio panels. The line is the canonical-anchor fit at the main 5% radial threshold. Shaded bands are bootstrap 95% diagnostic intervals, computed as pointwise 2.5% and 97.5% conditional bootstrap percentiles after resampling the selected extreme directions and refitting AGCA.

Figure S15 reports anchor sensitivity for the effective-dimension summary. Besides the canonical anchor, we fit AGCA using the leading principal direction of the selected angular sample and the normalized selected-sample mean direction. These anchors are diagnostic references; the main text uses the canonical anchor. The canonical anchor keeps the interpretation of loadings as departures from balanced complete dependence.




Figure S15: Anchor sensitivity for cumulative anchored variation explained. Each panel reports the canonical, principal, and mean-anchor spectra at the main 5% radial threshold.

Figure S16 puts this diagnostic in geometric perspective by reporting the spherical distance from each data-driven anchor to the canonical anchor over the same threshold grid used below. The data-driven anchors move modestly but measurably away from balanced complete dependence, especially at smaller thresholds in the OSAP panel, so the near overlap in Figure S15 is not merely a numerical identity between anchor choices.




Figure S16: Spherical distance from the canonical anchor to the principal and selected-sample mean anchors over the threshold grid. Each panel reports distances for one portfolio panel.

The scalar spectra are insensitive to these anchor changes. In the Fama–French panel, rank ten explains about 91% of anchored variation under all three anchors. In the OSAP panel, the rank-twelve explained variation remains close to 90%. Thus the low-rank conclusion remains stable across anchor choices, even though the diagnostic anchors move away from the canonical anchor by nonzero spherical distances.

Figure S17 reports threshold sensitivity. For each panel we refit the canonical-anchor AGCA model over a grid of radial tail fractions from 0.5% to 10%, keeping only thresholds with more than $d + 2$ selected directions. The expected bias-variance tradeoff is visible: the smallest k curves are less smooth, while larger thresholds admit less extreme days. The main empirical message nevertheless persists. The Fama–French panel remains more compressed than the OSAP panel, and both retain a clearly ordered low-rank spectrum across the displayed threshold choices.




Figure S17: Threshold sensitivity for cumulative anchored variation explained. Each line corresponds to a different number k of selected radial extremes.

S4.3 Loading heatmaps and AGC1 stability

The main text reports the Fama–French loading heatmap. Figure S18 gives the corresponding OSAP display. In both heatmaps, assets are on the horizontal axis and AGCA loading vectors are the rows. Signs are oriented by the deterministic convention used in the implementation: within each loading vector, the entry with largest absolute magnitude is positive. This is the same orientation convention used in the main text.




Figure S18: Canonical-anchor AGCA loadings for the OSAP robustness panel. Columns are portfolios, grouped by signal, and rows are the first five AGCA loading vectors. Loading signs are oriented so that the largest absolute entry in each loading vector is positive.

The OSAP heatmap is useful mainly as a robustness diagnostic. Because the assets are quintile portfolios formed on liquidity and trading-friction signals, the loadings should be read as contrasts across trading-friction portfolios, with the AGCA interpretation tied to angular tail directions. The important comparison with the Fama–French panel is the preservation of both angular variation and the portfolio functionals reported below with a similar number of AGCA coordinates. AGC1 has a simple gradient interpretation for several OSAP signals: bid–ask spread, dollar volume, and price move from negative loadings in the lower quintiles to positive loadings in the upper quintiles. The first coordinate therefore captures a signal-sorted rotation of the extreme-loss direction, analogous to the size contrast in the Fama–French panel.




Figure S19: AGC1 loadings with pointwise conditional bootstrap 95% diagnostic intervals. Bars are the canonical-anchor AGC1 loadings at the main 5% radial threshold. DIs are 2.5% and 97.5% bootstrap percentiles after resampling the selected extreme directions and aligning bootstrap loading signs to the full-sample AGC1.

Figure S19 checks whether the AGC1 signs are stable under resampling of the selected extreme directions. The DIs are diagnostic rather than formal confidence intervals, because they condition on the selected tail sample and on the rank-Pareto marginal transformation. In the Fama–French panel, 23 of 24 DIs exclude zero. The stable signs confirm that AGC1 is a repeated size contrast across the four bivariate sorts; the only near-zero case is the big high book-to-market portfolio. In the OSAP panel, 22 of 30 DIs exclude zero. The most stable gradients occur for bid–ask spread, dollar volume, and price, while several middle illiquidity and zero-trading quintiles have DIs that cross zero. These crossings identify entries whose signs should be read as weak local features of AGC1 rather than robust loadings.

S4.4 Portfolio tail-functional diagnostics

The portfolio-functional experiment uses four classes of weights. The first is the equal-weight portfolio across all assets in the panel. The second contains block-equal portfolios, one per Fama–French sort or OSAP signal. The third contains 250 randomly generated long-only portfolios with weights summing to one. The fourth contains 250 randomly generated limited-leverage portfolios with $\mathbf{1}^T w = 1$ and $\|w\|_1 \leq 1.5$. These random portfolios are fixed across ranks within each data set.

The capped-excess functional is bounded and Lipschitz by construction. The VaR diagnostic uses the Pareto-limit tail constant, with Pareto index $\alpha = 1$, matching the rank-Pareto marginal scale. The main text reports rank-level average relative errors for the Fama–French panel. Figure S20 gives the Fama–French class-level breakdown, and Figure S21 adds the OSAP robustness panel on the same relative-error scale.




Figure S20: Fama–French relative portfolio-functional errors by portfolio class. The left display reports relative capped-excess error and the right display reports relative normalized VaR error.




Figure S21: OSAP relative portfolio tail-functional errors by AGCA rank. The left display reports relative capped-excess errors; the right display reports relative normalized VaR errors.

Table S4 reports selected-rank values on both the original error scales and the relative financial scale used in the main text.

Table S4: Mean portfolio-functional errors at selected AGCA ranks.

Panel	Rank	AVE	Capped abs.	Capped rel. (%)	Log VaR	VaR rel. (%)
FF	3	0.646	0.0129	5.36	0.0522	5.36
FF	5	0.763	0.0082	3.41	0.0335	3.41
FF	8	0.864	0.0046	1.91	0.0189	1.91
FF	10	0.911	0.0030	1.25	0.0124	1.25
FF	12	0.939	0.0020	0.84	0.0083	0.84
OSAP	3	0.571	0.0192	8.59	0.0824	8.59
OSAP	5	0.697	0.0131	5.85	0.0569	5.85
OSAP	8	0.822	0.0075	3.36	0.0331	3.36
OSAP	10	0.869	0.0054	2.44	0.0241	2.44
OSAP	12	0.904	0.0040	1.79	0.0177	1.79

Figure S22 reports the original absolute capped-excess and log-VaR scales, including the positive post-projection. The post-projected curves are close to the unconstrained AGCA curves, so enforcing positive directions for simulation does not change the rank-level conclusion in these data.




Figure S22: Original-scale portfolio tail-functional errors by AGCA rank. The left display reports absolute capped-excess error; the right display reports absolute log error in the normalized Pareto-limit VaR tail constant. Each display includes the raw AGCA reconstruction and the positive post-projection.

The rank-level averages hide some variation across portfolio classes. Figure S23 reports the OSAP class-level relative errors, complementing the Fama–French class-level display above. The class-level patterns are consistent with the main conclusion: the errors decline monotonically with rank for all classes, and the Fama–French panel achieves small functional errors at lower ranks than the OSAP panel.




Figure S23: OSAP relative portfolio-functional errors by portfolio class. The left display reports relative capped-excess error and the right display reports relative normalized VaR error.

Figure S24 checks whether the same data-driven anchors used in the spectral diagnostic materially reduce the portfolio-functional errors. The comparison uses the same selected extreme directions and the same portfolio weights, changing only the anchor used to fit the AGCA recon-

struction. The rank effect is much larger than the anchor effect: principal and mean anchors can move the curves slightly, but they do not materially improve the capped-excess or normalized VaR errors relative to the canonical anchor. The canonical anchor is therefore competitive for these scalar tail summaries while retaining the clean loading interpretation used in the main text.




Figure S24: Anchor sensitivity for relative portfolio tail-functional errors. Rows correspond to the Fama–French and OSAP panels; columns report capped-excess and normalized VaR errors.

References

- Bingham, N. H., C. M. Goldie, and J. L. Teugels (1987). *Regular Variation*. Cambridge University Press.
- Chen, A. Y. and T. Zimmermann (2026). Open Source Asset Pricing. <https://www.openassetpricing.com/>. Accessed July 7, 2026.
- De Haan, L. and A. Ferreira (2006). *Extreme value theory: an introduction*. Springer.

- Einmahl, J. H. J., L. De Haan, and V. I. Piterbarg (2001). Nonparametric estimation of the spectral measure of an extreme value distribution. *The Annals of Statistics* 29(5), 1401–1423.
- Einmahl, J. H. J., L. De Haan, and A. K. Sinha (1997). Estimating the spectral measure of an extreme value distribution. *Stochastic Processes and their Applications* 70(2), 143–171.
- Einmahl, J. H. J., A. Krajina, and J. Segers (2012). An M-estimator for tail dependence in arbitrary dimensions. *The Annals of Statistics* 40(3), 1764–1793.
- Einmahl, J. H. J. and J. Segers (2009). Maximum empirical likelihood estimation of the spectral measure of an extreme-value distribution. *The Annals of Statistics* 37(5B), 2953–2989.
- French, K. R. (2026). Kenneth R. French Data Library. https://mba.tuck.dartmouth.edu/pages/faculty/ken.french/data_library.html. Accessed July 7, 2026.
- Genest, C. and J. Segers (2009). Rank-based inference for bivariate extreme-value copulas. *The Annals of Statistics* 37(5B), 2990–3022.
- Horn, R. A. and C. R. Johnson (2013). *Matrix Analysis* (2 ed.). Cambridge University Press.
- Shorack, G. R. and J. A. Wellner (1986). *Empirical Processes with Applications to Statistics*. Wiley.
- Wellner, J. A. (1978). Limit theorems for the ratio of the empirical distribution function to the true distribution function. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* 45(1), 73–88.