

Optimal use of a black-box learner in semiparametric estimation

Yihong Gu
Harvard University

Abstract

Consider the partial linear model $Y = \mu_0(X) + \beta_0 \cdot T + \varepsilon$ and $T = \pi_0(X) + u$ in the *structure-agnostic* setting, where we are blind to the structure μ_0 and π_0 and estimate the nuisances by a black-box hypothesis class. The learnability of the class is characterized by the estimation error δ_s in the absence of model misspecification and its L_2 mis-specification error $\delta_{a,\mu}$ and $\delta_{a,\pi}$ for μ_0 and π_0 , respectively. We propose a novel estimator of the target linear coefficient $\theta_0 = \beta_0$ with error rate

$$\frac{1}{\sqrt{n}} + \delta_{a,\mu} \cdot \delta_{a,\pi} + [\delta_s]^2.$$

A matching lower bound is also established, implying that this rate is unimprovable. Compared with the product rate yielded by double machine learning (DML), our estimator removes the suboptimal term $\max(\delta_{a,\mu}, \delta_{a,\pi}) \cdot \delta_s$ at no extra cost or assumption.

Building on the underlying insights, which are neither tailored to the one-learner setting nor the partial linear model, we propose *Transductive Adversarial Moment-calibrated Editing (TAME)*, which locally edits debiasing weights induced by black-box regression estimates on the inference sample through adversarial conditional moment calibration. TAME can be combined with any initial black-box estimates and can strictly improve on DML guarantees when the nuisance difficulties are imbalanced. We discuss how to fully exploit the advantages introduced by TAME, including the gains from using two learners, the resulting under-smoothing principle for model selection, and extensions to other linear functional estimation problems.

Keywords: Adversarial estimation, Black-box model, Causal inference, Partial linear model, Structure-agnostic optimality

1 Introduction

Modern semiparametric estimation leverages flexible black-box models to estimate nuisance functions, raising the fundamental question: *How can we optimally transcribe the nuisance errors of generic black-box learners into the final estimation error of the target parameter?* The dominant paradigm, exemplified by Double/Debiased Machine Learning (DML) (Chernozhukov et al., 2017, 2018), plugs the resulting black-box estimates into a Neyman-orthogonal score, yielding a bound in which the nuisance estimation errors enter multiplicatively. While this multiplicative propagation of nuisance errors is known to be sub-optimal in *structure-aware* settings, for example, the nuisance functions are Hölder smooth (Robins et al., 2008) or sparse linear models (Zhang & Zhang, 2014; Bradic et al., 2022), it was recently shown by Gu et al. (2026) that the DML multiplicative error rate can also be *provably* improved in *structure-agnostic* settings with generic black-box models. However, the proposed method therein only answers the above question in the specific regime where the treatment function (or more generally, the Riesz representer for the target linear functional) cannot be consistently estimated, under which only the outcome function comes into play. Building upon and generalizing the insights from Gu et al. (2026), this paper answers the above question when both the outcome function and the treatment function can be estimated efficiently by one black-box learner, and based on this, proposes a post-prediction debiasing procedure that can be applied to arbitrary initial black-box estimates.

1.1 A canonical example and our new message

We begin by studying the partial linear model (Engle et al., 1986; Rice, 1986), one of the most canonical semiparametric problems and a setting in which the structure-aware literature, including either Hölder smooth class (Donald & Newey, 1994; Carroll et al., 1997; Ai & Chen, 2003; Fan & Huang, 2005; Chen, 2007; Robins et al., 2008) or sparse linear models (Zhang & Zhang, 2014; Van de Geer et al., 2014; Javanmard & Montanari, 2014; Cai & Guo, 2017; Javanmard & Montanari, 2018), offers substantial messages. Suppose we have $2n$ observations $\{(X_i, T_i, Y_i)\}_{i=1}^{2n}$ drawn i.i.d. from the model

$$\begin{aligned} Y &= \beta_0 \cdot T + \mu_0(X) + \varepsilon & \text{with} & \quad \mathbb{E}[\varepsilon|X, T] = 0, \\ T &= \pi_0(X) + u & \text{with} & \quad \mathbb{E}[u|X] = 0. \end{aligned} \tag{1.1}$$

Here $X \in \mathbb{R}^d$ is the *covariate*, $Y \in \mathbb{R}$ is the *response*, T is the particular variable of interest, and we refer to it as a *treatment* variable, following the terminology in causal inference. The goal is to estimate the linear coefficient $\theta_0 = \beta_0 \in \mathbb{R}$, while μ_0 and π_0 are *nuisance functions*. We write the sample size as $2n$ to accommodate sample splitting.

Motivated by modern statistical learning, we consider the *structure-agnostic* setting, under which we are blind to any structure of μ_0 and π_0 like additivity, Hölder smoothness, or (sparse) linearity, but know that they can be estimated well by a provided black-box machine learning (ML) model, represented as a hypothesis function class $\mathcal{G}$. Following the notations of statistical learning (Bartlett et al., 2005; Koltchinskii, 2006; Massart & Nédélec, 2006; Wainwright, 2019), we use three quantities ($\delta_\mu^{\text{appr}}, \delta_\pi^{\text{appr}}, \delta_n^{\text{stoc}}$) to characterize how the black-box model $\mathcal{G}$ can estimate the nuisance functions. Denote $\|f\|_2 = (\mathbb{E}[f^2(X)])^{1/2}$, we use $\delta_h^{\text{appr}} := \inf_{g \in \mathcal{G}} \|g - h_0\|_2$ to denote the approximation error with respect to the ground-truth functions $h \in \{\mu, \pi\}$. We let δ_n^{stoc} be the stochastic error of the function class $\mathcal{G}$ based on n i.i.d. observations. It will be formally defined later and can be interpreted as the L_2 regression error when the ground-truth regression function belongs to $\mathcal{G}$. For example, a Vapnik–Chervonenkis (VC) (Vapnik & Chervonenkis, 1971) class with VC dimension $D = o(n)$ has stochastic error $\delta_n^{\text{stoc}} \lesssim \sqrt{D \log(en/D)/n}$. Applying the standard empirical process arguments, running least squares over $\mathcal{G}$ can estimate a regression function h_0 at the error rate

$O(\delta_h^{\text{appr}} + \delta_n^{\text{stoc}})$. For example, we have

$$\|\hat{\pi} - \pi_0\|_2 \lesssim \delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}} \quad \text{for} \quad \hat{\pi} = \underset{g \in \mathcal{G}}{\operatorname{argmin}} \frac{1}{n} \sum_{i=1}^n [T_i - g(X_i)]^2.$$

Here, we explicitly disentangle δ^{appr} and δ^{stoc} in our formalization instead of using the combined L_2 estimation error. This is because they affect the estimation of β_0 differently. In particular, structure-aware methods can improve on DML by reducing the contribution of the stochastic error δ^{stoc} ; see Section 1.3 of Gu et al. (2026) for a detailed discussion.

This paper characterizes the best attainable performance for estimating the target $\theta_0 = \beta_0$ using a generic black-box ML class $\mathcal{G}$ with known error budgets $(\delta_\mu^{\text{appr}}, \delta_\pi^{\text{appr}}, \delta_n^{\text{stoc}})$. As a benchmark, DML first uses $\mathcal{G}$ to estimate the two nuisance functions either explicitly (Chernozhukov et al., 2017, 2018, 2022b; Foster & Syrgkanis, 2023) or implicitly (Chernozhukov et al., 2022a, 2026) on one sample split and plugs $(\hat{\mu}, \hat{\pi})$ into a doubly robust (Robins et al., 1994) estimator on another hold-out split, yielding

$$\begin{aligned} |\hat{\theta}_{\text{DML}} - \theta_0| &\lesssim n^{-1/2} + \|\hat{\mu} - \mu_0\|_2 \cdot \|\hat{\pi} - \pi_0\|_2 \\ &\lesssim n^{-1/2} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2 + \max\{\delta_\pi^{\text{appr}}, \delta_\mu^{\text{appr}}\} \cdot \delta_n^{\text{stoc}}. \end{aligned}$$

where the second inequality substitutes the approximation and stochastic errors into the L_2 estimation error.

Recently, Gu et al. (2026) showed that this rate can be improved for a generic black-box class $\mathcal{G}$. They proposed another estimator with error guarantees $n^{-1/2} + \delta_\mu^{\text{appr}} + [\delta_n^{\text{stoc}}]^2$, together with a matching lower bound in the regime of $\delta_\pi^{\text{appr}} \asymp 1$, under which one cannot consistently estimate the treatment regression function. Their method removes the $\delta_\pi^{\text{appr}} \cdot \delta_n^{\text{stoc}}$ at the cost of inflating the bias $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}}$ by δ_μ^{appr} , leaving the optimal rate unresolved when $\delta_\pi^{\text{appr}} = o(1)$.

This paper proposes a novel method that, for any given black-box class $\mathcal{G}$, attains

$$|\hat{\theta}_{\text{TAME-1}} - \theta_0| \lesssim n^{-1/2} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2 \quad (1.2)$$

A lower bound is also proved, showing that the above error rate is not improvable even if $\mathcal{G}$ is specialized as a sparse linear model with identity covariance. The one-learner analysis further reveals a calibration principle that is not specific to either one learner or the partial linear model. This principle underlies the general TAME procedure and the extensions developed later in the paper.

1.2 New contributions

Optimally using a learner in partial linear models. In Section 2, we focus on the canonical one-learner partial linear model introduced in Section 1.1. We present an estimator attaining the rate (1.2), explain its rationale, and establish the upper bound result to be proved in Section 5. A matching lower bound is also established by restricting $\mathcal{G}$ to a sparse linear model with identity covariance. These theoretical results yield the three implications, extending Gu et al. (2026) to a broader regime in the *structure-agnostic* setting.

- (a) Regardless of which nuisance function is harder to learn, one can eliminate all mixed terms that are first-order in stochastic error, i.e., $(\delta_\mu^{\text{appr}} + \delta_\pi^{\text{appr}}) \cdot \delta_n^{\text{stoc}}$, in the DML rate.
- (b) Consequently, asymptotic normality remains possible even if one nuisance function is only consistently estimated, provided the other is estimated well and appropriate under-smoothing is adopted.
- (c) In the one-learner regime, which is where most provable structure-aware improvements are proposed, structure agnosticism introduces no fundamental limit: the rate (1.2) is attainable for any black-box class $\mathcal{G}$ and remains unimprovable when $\mathcal{G}$ is restricted to a sparse linear class.

The estimator adopts two key ideas. Firstly, it searches for debiasing weights that remain close to those induced by the black-box regression estimates while approximately satisfying the adversarial moment constraints introduced by Gu et al. (2026). Second, when the outcome function is harder to estimate, it constructs the outcome-side debiasing weights by a cross-sample objective. The weights depend only on the covariates in the inference split, while their moment constraints are imposed using the other split. This preserves the required conditional independence and removes the mixed term $\delta_n^{\text{stoc}} \cdot \delta_\mu^{\text{appr}}$.

TAME: A generic fine-tuning debiased procedure. In Section 3, we go beyond the one-learner setting and develop *Transductive Adversarial Moment-calibrated Editing (TAME)*, a post-prediction debiasing procedure that can be applied to arbitrary initial black-box estimates. We establish a generic oracle-type inequality and then specialize it to the two-learner setting. When the nuisance difficulties are imbalanced and the nuisance contribution dominates $n^{-1/2}$, the two-learner version of TAME attains a strictly faster rate than the DML rate. Its optimal model selection uses the standard bias-and-variance tradeoff for the harder nuisance and under-smooths the easier one. We illustrate these with the neural network learners.

Extensions beyond the partial linear model. In Section 4, we extend the outcome-easier branch (O-branch) to general linear functional estimation and the treatment-easier branch (R-branch) to average treatment effect estimation. These results show that the principles underlying TAME are not limited to the partial linear model.

1.3 Related works

Many structure-aware estimation error results for partial linear models can be expressed in our one-learner setting under additional structural assumptions; see the translation of selected results in our language in Table 1. When the class $\mathcal{G} = \mathcal{G}_p^{\text{lin}}$ to be used is a linear model with p basis, so that $\delta_n^{\text{stoc}} \asymp (p/n)^{1/2}$, Donald & Newey (1994) show the joint least squares estimator $\hat{\beta}$ can estimate β_0 at the error rate $|\hat{\beta} - \beta_0| \lesssim n^{-1/2} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + n^{-1} \sqrt{p}$ provided $p = o(n)$. Robins et al. (2008) uses the higher-order influence function for estimating a more general nonlinear functional that reduces to β_0 in (1.1) and attains the same error without the requirement $p = o(n)$. These linear sieve methods exploit a known p -dimensional subspace and its projection operator, a structure that does not directly extend to the non-convex sparse linear model (Bradic et al., 2019). When $\mathcal{G} = \mathcal{G}_{p,s}^{\text{lin}}$ is a sparse linear model of ambient dimension p and sparsity level s , its stochastic error is upper bounded by $(s \log(p)/n)^{1/2}$. As a special case of the capped ℓ_1 approximate sparsity considered by Zhang & Zhang (2014), the debiased Lasso estimator therein can estimate β_0 at the error rate $n^{-1/2} + n^{-1} s_\mu \log(p)$ when $\mu_0 \in \mathcal{G}_{p,s_\mu}^{\text{lin}}$ is an exact s_μ -sparse linear model and the L_2 projection of π_0 into the full linear p basis, which is s_π -sparse, can at least be consistently estimated by Lasso, i.e., $s_\pi \log(p) = o(n)$. On the other hand, without assuming any sparsity conditions on μ_0 , the estimator in Bradic et al. (2022) can estimate β_0 at the error rate $n^{-1/2} + n^{-1} s_\pi \log(p)$ when $\pi_0 \in \mathcal{G}_{p,s_\pi}^{\text{lin}}$ is an exact s_π -sparse linear model. Our estimator recovers the above two sharp rates for partial linear model, which is the “worst-case” structure-aware models, and generalizes them in two aspects: we allow $\mathcal{G}$ to be any hypothesis class, and also give a tight dependence $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}}$ on the misspecification error; see Remark 2.3.

Our constructed estimator combines two different estimators, handling the cases where either the treatment or outcome function is harder to learn separately. The estimator targeted at a harder treatment function is related to the idea of covariate balancing (Hellerstein & Imbens, 1999; Imai & Ratkovic, 2014; Zubizarreta, 2015; Zhao, 2019; Ning et al., 2020; Athey et al., 2018; Fan et al., 2022), or more generally, augmented minimax linear estimation for a convex class (Hirshberg & Wager, 2021; Bruns-Smith et al., 2025; Kong, 2025). The other estimator, which deals with a harder outcome function, uses a cross-sample objective for carefully controlling conditional independence structure and shares a similar spirit with the high-dimensional debiasing constructions of Bradic et al. (2022); Wang & Shah (2024). Other provable im-

Selected reference	$\mathcal{G}$	Nuisance remainder	Regime
Donald & Newey (1994)	Linear	$\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + n^{-1/2} \cdot \delta_n^{\text{stoc}}$	$\delta_n^{\text{stoc}} = o(1)$
Robins et al. (2008)	Linear	$\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + n^{-1/2} \cdot \delta_n^{\text{stoc}}$	-
Zhang & Zhang (2014)	Sparse linear	$(\delta_n^{\text{stoc}})^2$	$\delta_\mu^{\text{appr}} = 0$
Bradic et al. (2022)	Sparse linear	$(\delta_n^{\text{stoc}})^2$	$\delta_\pi^{\text{appr}} = 0$
Chernozhukov et al. (2017)	Generic	$\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + (\delta_n^{\text{stoc}})^2 + (\delta_\mu^{\text{appr}} \vee \delta_\pi^{\text{appr}}) \cdot \delta_n^{\text{stoc}}$	-
Gu et al. (2026)	Generic	$\delta_\mu^{\text{appr}} + [\delta_n^{\text{stoc}}]^2$	-
This paper	Generic	$\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2$	-

Table 1: Nuisance remainders from selected results for the partial linear model expressed in our notation; the common $n^{-1/2}$ term and regularity conditions are suppressed. The sparse linear rows display the result for exact s -sparse linear models used in our estimation error comparison rather than the full generality of the cited papers.

provements over the DML benchmark fall broadly into two categories. The first is structure-aware: some methods exploit linear sieve structure (Robins et al., 2008, 2016; Liu et al., 2017), whereas others exploit sparse high-dimensional structure (Athey et al., 2018; Tan, 2020; Ning et al., 2020; Wang & Shah, 2024; Celentano & Wainwright, 2023). The second category relies on additional assumptions on the first-stage fits (Van der Laan, 2014; Bonvini et al., 2024; van der Laan et al., 2024). By contrast, our method uses only the learnability of the black-box class, as characterized by its approximation and stochastic error budgets, to attain provable improvement.

Notations. Denote $x \vee y = \max\{x, y\}$ and $x \wedge y = \min\{x, y\}$. In this paper, we use c to denote a universal constant that is related to the boundedness from below and above conditions. We use $a \lesssim b$, $b \gtrsim a$, or $a = O(b)$ if there exists some constant C depending on c such that $a \leq Cb$ for any $n \geq C$, and write $a \asymp b$ if $a \lesssim b$ and $a \gtrsim b$. Denote $a = o(b)$, $a \ll b$, $b \gg a$, if a and b changes with n in that $\lim_{n \rightarrow \infty} a/b = 0$. We use $\text{poly}(a) \in \mathbb{R}$ to denote a scalar that depends polynomially on a .

2 Optimal one-learner TAME for the partial linear model

In this section, we propose the one-learner version of TAME, referred to as TAME-1, for estimating the linear coefficient β_0 in the partial linear model (1.1). As a proof of concept for the generic TAME framework, TAME-1 attains the structure-agnostic optimal rate (1.2) in the one-learner setting.

Section 2.1 introduces the regularity condition and formally defines the approximation error and stochastic error for the hypothesis class $\mathcal{G}$. The method is then presented in Section 2.2, accompanied by the underlying rationale in Section 2.3. The upper bound and asymptotic normality results are established in Section 2.4, with a matching lower bound result in Section 2.5.

2.1 Setup

We first impose a standard regularity condition for the partial linear model (1.1). Condition 2.1 (a) bounds the coefficient, nuisance functions, and noises, while (b) ensures that the residual $u = T - \pi_0(X)$ has nondegenerate variance, which allows β_0 to be identifiable.

Condition 2.1 (Regularity conditions for partial linear model). *There exists a constant $c \geq 1$ such that:*

- (a) *The coefficient, nuisance functions, and noises are uniformly bounded: $|\beta_0| \leq c$, $\|\mu_0\|_\infty \vee \|\pi_0\|_\infty \leq c$, $|\varepsilon| \vee |u| \leq c$.*

(b) The treatment variable T cannot be fully explained by X , i.e., $\mathbb{E}[u^2] \geq 1/c$.

To formalize the stochastic error of a function class, we first define its localized population Rademacher complexity.

Definition 1 (Localized population Rademacher complexity). *For a given radius $\delta > 0$, function class $\mathcal{F}$ on $\mathcal{Z}$, and a probability distribution ν on $\mathcal{Z}$, define*

$$R_{n,\nu}(\delta; \mathcal{F}) = \mathbb{E}_{Z,R} \left[\sup_{h \in \mathcal{F}, \|h\|_{L_2(\nu)} \leq \delta} \left| \frac{1}{n} \sum_{i=1}^n R_i h(Z_i) \right| \right],$$

where $Z_1, \dots, Z_n \in \mathcal{Z}$ are n i.i.d. observations from ν , and $R_1, \dots, R_n$, independent of $Z_1, \dots, Z_n$, are i.i.d. Rademacher variables taking values in $\{-1, +1\}$ with equal probability.

We next use this complexity to define the stochastic error of a uniformly bounded hypothesis class.

Definition 2 (Hypothesis class with stochastic error $\bar{\delta}$). *Let $\mathcal{F}$ be a function class on domain $\mathcal{Z}$, and let ν be a probability distribution on $\mathcal{Z}$. Define*

$$\delta^{\text{stoc}}(\mathcal{F}, n, \nu) := \inf \left\{ \delta_s > \sqrt{\frac{\log(n)}{n}} : \sup_{\delta \geq \delta_s} \frac{R_{n,\nu}(\delta; \partial\mathcal{F})}{c \cdot \delta} \leq \delta_s \right\}, \quad (2.1)$$

where $\partial\mathcal{F} := \{f - \tilde{f} : f, \tilde{f} \in \mathcal{F}\}$. We say $\mathcal{F}$ is a uniformly bounded class with stochastic error upper bounded by $\bar{\delta}$, denoted by $\mathcal{F} \in \mathcal{HCS}(\bar{\delta}; n, \nu)$, if $\|f\|_\infty \leq c$ for any $f \in \mathcal{F}$ and $\delta^{\text{stoc}}(\mathcal{F}, n, \nu) \leq \bar{\delta}$.

Condition 2.2 (Hypothesis class $\mathcal{G}$). *Let ν_x be the distribution of X . Assume $\mathcal{G} \in \mathcal{HCS}(\delta_n^{\text{stoc}}; n, \nu_x)$. Moreover, we define the approximation errors, i.e., L_2 misspecification errors, as*

$$\delta_\mu^{\text{appr}} := \inf_{g \in \mathcal{G}} \|g - \mu_0\|_{L_2(\nu_x)} \quad \text{and} \quad \delta_\pi^{\text{appr}} = \inf_{g \in \mathcal{G}} \|g - \pi_0\|_{L_2(\nu_x)}. \quad (2.2)$$

Remark 2.1. *For ease of presentation, we adopt three simplifications: (1) the coefficient, nuisance functions, noises, and hypothesis classes are uniformly bounded; (2) the local Rademacher complexity satisfies the bounds for any $\delta \geq \delta_n^{\text{stoc}}$; and (3) the critical radius is at least $\sqrt{\log(n)/n}$. The boundedness assumptions in (1) can be replaced by suitable sub-Gaussian tail conditions. For (2), if $\bar{\mathcal{F}} = \partial\mathcal{F}$ is star-shaped, i.e., $\alpha f \in \bar{\mathcal{F}}$ for any $f \in \bar{\mathcal{F}}$ and $|\alpha| \leq 1$, then (2.1) reduces to the usual critical radius definition; see [Wainwright \(2019\)](#). Without star-shapedness, one may instead use the usual critical radius at the cost of slightly more complicated final bounds. (3) avoids repeatedly displaying negligible $\log(n)/n$ terms.*

2.2 The debiased estimator for one learner

For the observed data $\mathcal{D} = \{(X_i, T_i, Y_i)\}_{i=1}^{2n}$ with splits $\mathcal{D}_1 = \{(X_i, T_i, Y_i)\}_{i=1}^n$ and $\mathcal{D}_2 = \{(X_i, T_i, Y_i)\}_{i=n+1}^{2n}$, we first use $\mathcal{D}_2$ to obtain initial estimates. In particular, we run joint least squares for (β_0, μ_0) and least squares for π_0

$$(\hat{\beta}, \hat{\mu}) \in \underset{(\beta, g) \in [-c, c] \times \mathcal{G}}{\operatorname{argmin}} \frac{1}{n} \sum_{i=n+1}^{2n} (Y_i - \beta \cdot T_i - g(X_i))^2, \quad \hat{\pi} \in \underset{g \in \mathcal{G}}{\operatorname{argmin}} \frac{1}{n} \sum_{i=n+1}^{2n} (T_i - g(X_i))^2, \quad (\text{S0})$$

to get an initial estimate of (β_0, μ_0) and π_0 , respectively. Using the standard empirical process argument, we have $\|\hat{\mu} - \mu_0\|_2 + |\hat{\beta} - \beta_0| \lesssim \delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}}$ and $\|\hat{\pi} - \pi_0\|_2 \lesssim \delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}}$ with high probability.

Our estimator requires knowledge of the ordering between $(\delta_\mu^{\text{appr}}, \delta_\pi^{\text{appr}})$, and selects between the two estimators as follows:

$$\hat{\theta} = \begin{cases} \hat{\theta}_\mu \text{ in (S2-}\mu\text{)} & \text{if } \delta_\mu^{\text{appr}} \leq \delta_\pi^{\text{appr}}, \\ \hat{\theta}_\pi \text{ in (S2-}\pi\text{)} & \text{if } \delta_\mu^{\text{appr}} > \delta_\pi^{\text{appr}} \end{cases} \quad (2.3)$$

We let $\lambda > 0$ denote the tuning parameter in the minimax programs below; it should be tuned according to the full error budgets $(\delta_\mu^{\text{appr}}, \delta_\pi^{\text{appr}}, \delta_n^{\text{stoc}})$ to attain the optimal rate.

CASE 1: $\delta_\mu^{\text{appr}} \leq \delta_\pi^{\text{appr}}$. Let $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_n) \in \mathbb{R}^n$ with $\tilde{a}_i = [T_i - \hat{\pi}(X_i)] / [\{\frac{1}{n} \sum_{l=1}^n T_l(T_l - \hat{\pi}(X_l))\} \vee (2c)^{-1}]$ be the debiasing weights induced by the black-box estimates. The lower truncation keeps the denominator in $\tilde{a}_i$ bounded away from zero. Here we first use $\mathcal{D}_1$ to construct empirical debiasing weights $\hat{a} = (\hat{a}_1, \dots, \hat{a}_n)$ through the minimax program

$$\begin{aligned} \hat{a} \in \operatorname{argmin}_{a \in \mathbb{R}^n} \lambda \cdot Q(a, \tilde{a}) + \sup_{\beta \in \mathbb{R}, f \in \partial \mathcal{G}} \Phi_\mu(f, \beta, a) \quad \text{where } Q(v, v') = \frac{1}{n} \sum_{i=1}^n (v_i - v'_i)^2, \\ \Phi_\mu(f, \beta, a) = \left| \frac{1}{n} \sum_{i=1}^n (\beta \cdot T_i + f(X_i)) \cdot a_i - \beta \right| - \frac{1}{n} \sum_{i=1}^n f^2(X_i). \end{aligned} \quad (\text{S1-}\mu)$$

For definiteness, if the objective in (S1- μ) does not admit a finite minimizer, we set $\hat{a} = 0$. Conditional on the initial estimates from $\mathcal{D}_2$, $\hat{a}$ depends on $\mathcal{D}_1$ only through $\{(X_i, T_i)\}_{i=1}^n$, and not through $\{Y_i\}_{i=1}^n$. The estimate of $\theta_0 = \beta_0$ in this case is the following debiased estimator

$$\hat{\theta}_\mu = \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{\mu}(X_i)) \cdot \hat{a}_i. \quad (\text{S2-}\mu)$$

CASE 2: $\delta_\mu^{\text{appr}} > \delta_\pi^{\text{appr}}$. Let $\tilde{m} = (\tilde{m}_1, \dots, \tilde{m}_n) \in \mathbb{R}^n$ with $\tilde{m}_i = \hat{\mu}(X_i) + \hat{\beta} \cdot \hat{\pi}(X_i)$ be the debiasing weights induced by the black-box estimates in this case. We will use both $\mathcal{D}_1$ and $\mathcal{D}_2$ to construct empirical debiasing weights $\hat{m} = (\hat{m}_1, \dots, \hat{m}_n)$ on $\mathcal{D}_1$ through the following minimax program,

$$\begin{aligned} \hat{m} \in \operatorname{argmin}_{m \in \mathbb{R}^n} \lambda \cdot Q(m, \tilde{m}) + \sup_{f \in \partial \mathcal{G}} \Phi_\pi(f, m), \quad \text{where} \\ \Phi_\pi(f, m) = \left| \frac{1}{n} \sum_{i=n+1}^{2n} Y_i f(X_i) - \frac{1}{n} \sum_{i=1}^n m_i f(X_i) \right| - \frac{1}{n} \sum_{i=1}^n f^2(X_i) \end{aligned} \quad (\text{S1-}\pi)$$

Thus, $\hat{m}$ depends on all of $\mathcal{D}_2$ and, conditional on $\mathcal{D}_2$, on $\mathcal{D}_1$ only through the covariates $\{X_i\}_{i=1}^n$. This differs from (S1- μ), where $\hat{a}$ may also depend on the treatments $\{T_i\}_{i=1}^n$ in $\mathcal{D}_1$. This *cross-sample* construction preserves the conditional mean-zero structure needed for the treatment noises $\{u_i\}_{i=1}^n$. The estimate of $\theta_0 = \beta_0$ in this case is the following debiased estimator

$$\hat{\theta}_\pi = \left\{ (2c)^{-1} \vee \frac{1}{n} \sum_{i=1}^n [T_i - \hat{\pi}(X_i)]^2 \right\}^{-1} \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{m}_i) (T_i - \hat{\pi}(X_i)). \quad (\text{S2-}\pi)$$

2.3 Rationale

To keep the presentation concise in this subsection, the inequalities involving “ $\lesssim$ ” are understood to hold with high probability. Denote $\|v\|_n = (\frac{1}{n} \sum_{i=1}^n v_i^2)^{1/2}$ for an n -dimensional vector $v = (v_1, \dots, v_n) \in \mathbb{R}^n$. We identify a function f with its evaluation vector $f_{1:n} := (f(X_1), \dots, f(X_n))$ and accordingly write $\|f\|_n := \|f_{1:n}\|_n$.

We first explain the rationale behind the estimator $\hat{\theta}_\mu$ by considering the regime where the denominator in $\tilde{a}_i$ exceeds $1/(2c)$ so the lower bound truncation is inactive. This event occurs with high probability when $\hat{\pi}$ consistently estimates π_0 ; the formal analysis also covers the truncated case.

At a high level, it uses the adversarial moment-calibration criterion from [Gu et al. \(2026\)](#) to fine-tune the debiasing weights in a neighbourhood of the weights $\tilde{a}$ used by the standard DML estimator, so that the resulting estimator benefits from both objectives. To see this, when $\lambda \rightarrow \infty$, we will have $\hat{a}_i \rightarrow \tilde{a}_i = (T_i - \hat{\pi}(X_i))/\{\frac{1}{n} \sum_{l=1}^n T_l(T_l - \hat{\pi}(X_l))\}$, which coincides with the double machine learning estimator (3.5) when the weights $\hat{a}$ are plugged into (S2- μ); when $\lambda \rightarrow 0$, it is similar to the minimax empirical debiasing estimator in [Gu et al. \(2026\)](#), which is designed specifically to remove the first-order stochastic error at the cost of replacing $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}}$ by δ_μ^{appr} . In particular, our construction of the debiasing weights in (S1- μ) is akin to the following constrained formulation obtained via the Lagrange multiplier method

$$\begin{aligned} \hat{a} \in \operatorname{argmin}_{a \in \mathcal{A}} \frac{1}{n} \sum_{i=1}^n (a_i - \tilde{a}_i)^2 \quad \text{where} \\ \mathcal{A} = \left\{ a \in \mathbb{R}^n : \frac{1}{n} \sum_{i=1}^n a_i T_i = 1, \sup_{f \in \partial \mathcal{G}} \left| \frac{1}{n} \sum_{i=1}^n f(X_i) \cdot a_i \right| - \frac{1}{n} \sum_{i=1}^n f^2(X_i) \leq \tilde{C}(\delta_n^{\text{stoc}})^2 \right\} \end{aligned} \quad (2.4)$$

for a large enough constant $\tilde{C} = \text{poly}(c)$. The moment inequality in $\mathcal{A}$ can be viewed as a black-box analogue of a Dantzig-type ([Candes & Tao, 2007](#)) feasibility constraint.

Now, let us briefly explain why the weights in (2.4), plugged into (S2- μ) can yield the upper bound $n^{-1/2} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2$ when $\delta_\mu^{\text{appr}} \leq \delta_\pi^{\text{appr}}$. Let $\bar{D} = \frac{1}{n} \sum_{l=1}^n T_l(T_l - \pi_0(X_l))$. Since $\bar{D}$ is an average of i.i.d. random variables with mean $\mathbb{E}[u^2] \geq 1/c$, we have $\bar{D} \gtrsim 1$ for large n . Moreover, the ‘‘oracle weights’’ $\bar{a} \in \mathbb{R}^n$ with $\bar{a}_i = [T_i - \pi_0(X_i)]/\bar{D} = u_i/\bar{D}$ lie in the feasible set $\mathcal{A}$ in (2.4) with high probability; see the detailed justification in Section 2.2 of [Gu et al. \(2026\)](#). Thus, the feasible set $\mathcal{A}$ is non-empty with high probability, and its constraints apply to the solution $\hat{a}$.

We pick fixed $\bar{g}_\mu \in \mathcal{G}$ such that $\|\bar{g}_\mu - \mu_0\|_2 \leq 2\delta_\mu^{\text{appr}}$. It follows from the model (1.1) and the constraint $\frac{1}{n} \sum_{i=1}^n \hat{a}_i T_i = 1$ that

$$\begin{aligned} \hat{\theta}_\mu - \theta_0 &= \frac{1}{n} \sum_{i=1}^n (\varepsilon_i + \beta_0 \cdot T_i + \mu_0(X_i) - \hat{\mu}(X_i)) \cdot \hat{a}_i - \beta_0 \\ &= \underbrace{\frac{1}{n} \sum_{i=1}^n \varepsilon_i \hat{a}_i}_{\mathsf{T}_{\mu,n}} + \underbrace{\frac{1}{n} \sum_{i=1}^n (\mu_0 - \bar{g}_\mu)(X_i) \cdot \hat{a}_i}_{\mathsf{T}_{\mu,a}} + \underbrace{\frac{1}{n} \sum_{i=1}^n (\bar{g}_\mu - \hat{\mu})(X_i) \cdot \hat{a}_i}_{\mathsf{T}_{\mu,s}}. \end{aligned}$$

As shown below $\|\hat{a}\|_n \lesssim 1$. Moreover, conditioned on $\mathcal{D}_2$ and $\{(X_i, T_i)\}_{i=1}^n$, the weights $\hat{a}$ are fixed and ε_i are independent with conditional zero mean. Hence $|\mathsf{T}_{\mu,n}| \lesssim n^{-1/2}$.

The minimization of the empirical L_2 error between our weights $\hat{a}$ and the weights $\tilde{a}$ given by the black-box estimate $\hat{\pi} \in \mathcal{G}$ will help minimize the error in $\mathsf{T}_{\mu,a}$. To see this, by the minimization objective in (2.4), we have $\|\hat{a} - \tilde{a}\|_n \leq \|\tilde{a} - \bar{a}\|_n$. It is easy to verify that $\|\tilde{a} - \bar{a}\|_n \lesssim \|\hat{\pi} - \pi_0\|_2 \lesssim \delta_n^{\text{stoc}} + \delta_\pi^{\text{appr}}$. This means our debiasing weights $\hat{a}$ are also close to the ‘‘oracle weights’’ $\bar{a}$ in a similar manner to $\tilde{a}$: $\|\hat{a} - \bar{a}\|_n \leq \|\hat{a} - \tilde{a}\|_n + \|\tilde{a} - \bar{a}\|_n \leq 2\|\tilde{a} - \bar{a}\|_n \lesssim \delta_n^{\text{stoc}} + \delta_\pi^{\text{appr}}$. Since $\{u_i\}_{i=1}^n$ are independent zero-mean random variables conditioned on $\{X_i\}_{i=1}^n$, it thus follows from the triangle inequality and the Cauchy-Schwarz inequality that

$$|\mathsf{T}_{\mu,a}| \leq \left| \frac{1}{n} \sum_{i=1}^n (\mu_0 - \bar{g}_\mu)(X_i) \cdot u_i \right| / |\bar{D}| + \|\mu_0 - \bar{g}_\mu\|_n \cdot \|\hat{a} - \bar{a}\|_n$$

$$\lesssim \frac{1}{\sqrt{n}} + \delta_\mu^{\text{appr}} \cdot (\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}}) \lesssim \frac{1}{\sqrt{n}} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2,$$

where the last inequality follows from $\delta_\mu^{\text{appr}} \cdot \delta_n^{\text{stoc}} \leq [\delta_\mu^{\text{appr}}]^2 + [\delta_n^{\text{stoc}}]^2 \leq \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2$ as $\delta_\mu^{\text{appr}} \leq \delta_\pi^{\text{appr}}$.

The adversarial moment constraint with respect to $\partial\mathcal{G}$ in $\mathcal{A}$ will make the term $\mathsf{T}_{\mu,s}$ scale quadratically in $(\delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}})$, which explicitly removes the $\delta_n^{\text{stoc}} \cdot \delta_\pi^{\text{appr}}$ error in the DML product rate when $\delta_\pi^{\text{appr}} \gg \delta_\mu^{\text{appr}}$.

To see this, given $\hat{a} \in \mathcal{A}$, the following instance-dependent error bound holds

$$\forall g, \tilde{g} \in \mathcal{G}, \quad \left| \frac{1}{n} \sum_{i=1}^n (g - \tilde{g})(X_i) \cdot \hat{a}_i \right| \leq \|g - \tilde{g}\|_n^2 + \tilde{C}(\delta_n^{\text{stoc}})^2. \quad (2.5)$$

Applying this with $(g, \tilde{g}) = (\bar{g}_\mu, \hat{\mu})$ gives $|\mathsf{T}_{\mu,s}| \lesssim \|\hat{\mu} - \bar{g}_\mu\|_n^2 + (\delta_n^{\text{stoc}})^2 \lesssim [\delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}}]^2$. The squared approximation error $[\delta_\mu^{\text{appr}}]^2$ is absorbed into $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}}$ given the regime $\delta_\mu^{\text{appr}} \leq \delta_\pi^{\text{appr}}$ under consideration. Combining the error bounds on $\mathsf{T}_{\mu,n}$, $\mathsf{T}_{\mu,a}$, and $\mathsf{T}_{\mu,s}$ yields the stated rate.

Remark 2.2 (Combining two objectives). *For linear (Bruns-Smith et al., 2025) and sparse linear classes (Zhang & Zhang, 2014; Van de Geer et al., 2014), regression risk minimization can be equivalent to imposing the moment constraints with respect to the adopted hypothesis class, by the minimax duality (Hirshberg & Wager, 2021) and the KKT condition (Bickel et al., 2009), respectively. Such an equivalence need not hold for a generic, possibly non-convex hypothesis class $\mathcal{G}$. Our estimator thus combines the centered ridge penalty $Q(\hat{a}, \tilde{a})$ for deviating from the regression-induced weights with an adversarial moment constraint, rather than relying on either objective to imply the other.*

For CASE 2, we will also adopt the idea of using flexible weights to eliminate the potential $\delta_\mu^{\text{appr}} \cdot \delta_n^{\text{stoc}}$ error term, but an additional challenge occurs here: the outcome-side debiasing weights must be constructed without using the treatment residuals in the inference sample $\mathcal{D}_1$. Otherwise, the term involving u_i is no longer conditionally mean zero. At first glance, this requirement conflicts with using the outcomes Y in the inference sample $\mathcal{D}_1$ to impose the desired moment constraints.

To see why the construction in CASE 1 cannot be directly mirrored, let us consider the natural DML-form estimator $\hat{\theta} = \frac{1}{n} \sum_{i=1}^n (Y_i - m_i)(T_i - \hat{\pi}(X_i)) / \frac{1}{n} \sum_{i=1}^n T_i(T_i - \hat{\pi}(X_i))$ with the candidate debiasing weights $m = (m_1, \dots, m_n) \in \mathbb{R}^n$. Picking a fixed $\bar{g}_\pi \in \mathcal{G}$ such that $\|\bar{g}_\pi - \pi_0\|_2 \leq 2\delta_\pi^{\text{appr}}$ and expanding $\hat{\theta} - \theta_0$, we need to control the two terms below that share similar spirits to $\mathsf{T}_{\mu,s}$ and $\mathsf{T}_{\mu,n}$, respectively, in CASE 1:

$$\underbrace{\frac{1}{n} \sum_{i=1}^n (\mu_0(X_i) - m_i) (\bar{g}_\pi - \hat{\pi})(X_i)}_{\check{\mathsf{T}}_{\pi,s}} + \underbrace{\frac{1}{n} \sum_{i=1}^n (\mu_0(X_i) - m_i) \cdot u_i}_{\check{\mathsf{T}}_{\pi,n}}$$

One could calibrate m on $\mathcal{D}_1$ to control $\check{\mathsf{T}}_{\pi,s}$. A same-sample analogue would impose the moment constraints using the residualized outcomes $Y - \hat{\beta}T$. These residualized outcomes will depend on u_i through $T_i = \pi_0(X_i) + u_i$, making m depend on the treatment residuals $\{u_i\}_{i=1}^n$ in $\mathcal{D}_1$. Consequently, $\check{\mathsf{T}}_{\pi,n}$ would no longer be an average of conditionally zero-mean random variables.

Our construction avoids this conflict by changing both the target of the outcome-side values and the sample used for calibration. The optimized debiasing weight $\hat{m}_i$ will instead approximate $\bar{m}_i = \mathbb{E}[Y|X_i] = \mu_0(X_i) + \beta_0\pi_0(X_i)$, and the required moment constraints are learned from $\mathcal{D}_2$, while $\hat{m}$ depends on $\mathcal{D}_1$ only through its covariates $\{X_i\}_{i=1}^n$. The high-level ideas behind the design of the objective (S1- π) are twofold. First, the empirical L_2 norm $\|\bar{m} - \hat{m}\|_n$ should be small. This is done by minimizing $\|\hat{m} - \tilde{m}\|_n$ using the initial estimate $\tilde{m}_i = \hat{\beta} \cdot \hat{\pi}(X_i) + \hat{\mu}(X_i)$ satisfying $\|\tilde{m} - \bar{m}\|_n \lesssim \delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}}$. Second, minimizing $\sup_{f \in \partial\mathcal{G}} \Phi_\pi(f, m)$ w.r.t. m will impose the instance-dependent error bound $|\frac{1}{n} \sum_{i=1}^n (\bar{m}_i - \hat{m}_i)(g - \tilde{g})(X_i)| \lesssim \|g - \tilde{g}\|_n^2 + (\delta_n^{\text{stoc}})^2$ for any $g, \tilde{g} \in \mathcal{G}$, analogous to (2.5).

After separating the semiparametric efficient term and the standard higher-order terms arising from the estimated treatment regression and denominator, the three additional dominant terms are

$$\underbrace{\left| \frac{1}{n} \sum_{i=1}^n (\tilde{m}_i - \hat{m}_i) \cdot u_i \right|}_{\mathsf{T}_{\pi,n}} + \underbrace{\left| \frac{1}{n} \sum_{i=1}^n (\tilde{m}_i - \hat{m}_i) \cdot (\pi_0 - \bar{g}_\pi)(X_i) \right|}_{\mathsf{T}_{\pi,a}} + \underbrace{\left| \frac{1}{n} \sum_{i=1}^n (\tilde{m}_i - \hat{m}_i) \cdot (\bar{g}_\pi - \hat{\pi})(X_i) \right|}_{\mathsf{T}_{\pi,s}}.$$

Similarly, the minimization of $\|\hat{m} - \tilde{m}\|_n$ will make $|\mathsf{T}_{\pi,a}| \lesssim n^{-1/2} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2$, while the minimization of adversarial objective controls the term $|\mathsf{T}_{\pi,s}| \lesssim [\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}}]^2$, which explicitly eliminates the sub-optimal term $\delta_\mu^{\text{appr}} \cdot \delta_n^{\text{stoc}}$ in the DML rate. At the same time, the *cross-sample* construction of the weights $\{\hat{m}_i\}_{i=1}^n$ make them dependent on the split $\mathcal{D}_1$ only through $\{X_i\}_{i=1}^n$, thus $\mathsf{T}_{\pi,n}$ continues to be the sum of independent zero-mean random variables conditioned on fixed $(\{X_i\}_{i=1}^n, \mathcal{D}_2)$ in a similar spirit to $\mathsf{T}_{\mu,n}$ for $\hat{\theta}_\mu$.

2.4 Upper bound and asymptotic normality

Recall the partial linear model in (1.1), and define the first-order oracle term as

$$\hat{\varphi}_{\text{PLM}} = \frac{1}{n} \sum_{i=1}^n \varepsilon_i \cdot w_i \quad \text{with} \quad w_i = u_i / \mathbb{E}[u^2] \quad (2.6)$$

Under the additional conditional homoskedasticity assumption $\mathbb{E}[\varepsilon^2 \mid X, T] \equiv \sigma_\varepsilon^2$, this is the empirical average of the semiparametric efficient influence function.

The following theorem provides an oracle-type inequality for any given hyperparameter $\lambda \in (0, 1]$.

Theorem 2.1 (Oracle-type inequality). *Consider the partial linear model (1.1) satisfying Condition 2.1 and the black-box model class $\mathcal{G}$ satisfying Condition 2.2. Then there exists a constant $\tilde{C} = \text{poly}(\mathfrak{c})$, if $\delta_n^{\text{stoc}} \cdot \tilde{C} \leq 1$, and $t \in [1, n/\tilde{C}]$, then with probability at least $1 - 2e^{-t} - 2e^{-n(\delta_n^{\text{stoc}})^2}$, our estimator $\hat{\theta}$ in (2.3) with $\lambda \in [(\delta_n^{\text{stoc}})^2, 1]$ satisfies*

$$\tilde{C}^{-1} |\hat{\theta} - \theta_0 - \hat{\varphi}_{\text{PLM}}| \leq \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + (\delta_n^{\text{stoc}})^2 + \left(\lambda \cdot (\delta_{\max}^{\text{appr}})^2 + \frac{\delta_n^{\text{stoc}} \cdot \delta_{\min}^{\text{appr}}}{\sqrt{\lambda}} \right) + \text{Res},$$

where $\delta_{\max}^{\text{appr}} = \delta_\mu^{\text{appr}} \vee \delta_\pi^{\text{appr}}$, $\delta_{\min}^{\text{appr}} = \delta_\mu^{\text{appr}} \wedge \delta_\pi^{\text{appr}}$, and

$$\text{Res} := \sqrt{\frac{t}{n}} \cdot \left[\delta_{\max}^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}} \right]$$

is a high-order remainder: for fixed $t \geq 1$, $\text{Res} = o(n^{-1/2})$ as long as $\delta_{\max}^{\text{appr}} = o(1)$ and $\lambda \gg (\delta_n^{\text{stoc}})^2$.

Theorem 2.1 will be proved in Section 5. The two terms dependent on λ quantify the trade-off between keeping the edited weights close to the initial black-box estimates and enforcing the adversarial moment constraints. Choosing λ as in Corollary 2.2 balances the two terms and gives the desired error bound (1.2). Under the stronger conditions in the corollary, the same choice also yields asymptotic normality.

Corollary 2.2. *Under Condition 2.1 and Condition 2.2, for any $t \geq 1$, with probability at least $1 - 3e^{-t} - 2e^{-n(\delta_n^{\text{stoc}})^2}$,*

$$|\hat{\theta} - \theta_0| \lesssim \sqrt{\frac{t}{n}} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + (\delta_n^{\text{stoc}})^2 \quad \text{with} \quad \lambda = \left(\frac{\delta_n^{\text{stoc}}}{\delta_{\max}^{\text{appr}} + \delta_n^{\text{stoc}}} \right)^2.$$

Furthermore, if $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} = o(n^{-1/2})$, $\delta_n^{\text{stoc}} = o(n^{-1/4})$, and $\delta_{\max}^{\text{appr}} = o(1)$, then the same choice of λ gives $\hat{\theta} - \theta_0 = \hat{\varphi}_{\text{PLM}} + o_P(n^{-1/2})$.

Remark 2.3 (Relation to structure-aware results for sparse linear models). *As a special case, if μ_0 is a well-specified sparse linear model with sparsity level s_μ and $\mathcal{G}$ is the corresponding sparse linear class, then $\delta_n^{\text{stoc}} = \sqrt{s_\mu \log(p)/n}$ and $\delta_\mu^{\text{appr}} = 0$. Thus, the above error rate is instantiated as $n^{-1/2} + 0 \cdot \delta_\mu^{\text{appr}} + (\delta_n^{\text{stoc}})^2 \asymp n^{-1/2} + s_\mu \log(p)/n$, matching that given by the debiased-Lasso estimator (Zhang & Zhang, 2014; Van de Geer et al., 2014). On the other hand, if π_0 is a well-specified sparse linear model with sparsity level s_π , and $\mathcal{G}$ is the corresponding sparse linear class, then $\delta_\pi^{\text{appr}} = 0$ and $\delta_n^{\text{stoc}} = \sqrt{s_\pi \log(p)/n}$, and the error bound becomes $n^{-1/2} + s_\pi \log(p)/n$, which coincides with the rate in Bradic et al. (2022).*

Thus, Corollary 2.2 extends the previous structure-aware results for the partial linear model in two aspects. First, it applies to any black-box hypothesis class $\mathcal{G}$ with a controlled stochastic error budget, including classes that cannot be reduced to a well-conditioned sparse linear model via the ϵ -cover argument. Second, it explicitly gives the tight dependence $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}}$ on the approximation error budgets; Section 2.5 shows that this dependence is unimprovable.

When $\delta_\pi^{\text{appr}} \asymp 1$, the upper bound reduces to the result of Gu et al. (2026). When $\delta_\mu^{\text{appr}} \asymp 1$, it gives the symmetric rate

$$n^{-1/2} + \delta_\pi^{\text{appr}} + [\delta_n^{\text{stoc}}]^2.$$

Thus, even if μ_0 cannot be consistently estimated, β_0 can still be estimated at this rate when π_0 can be estimated by the hypothesis class $\mathcal{G}$ with approximation error δ_π^{appr} and stochastic error δ_n^{stoc} . Moreover, the bound improves continuously as the approximation error of the harder nuisance $\delta_{\max}^{\text{appr}} = \delta_\mu^{\text{appr}} \vee \delta_\pi^{\text{appr}}$ decreases from the hardest regime $\delta_{\max}^{\text{appr}} \asymp 1$.

Remark 2.4 (Inference). *Additionally assume that $\sigma_0 = \sqrt{\mathbb{E}[u^2 \varepsilon^2]}/\mathbb{E}[u^2] > 0$. Under the asymptotic normality conditions in Corollary 2.2, it is easy to verify that the following plug-in estimate of σ_0 , denoted as $\hat{\sigma}$, satisfies $|\hat{\sigma} - \sigma_0| = o_P(1)$:*

$$\frac{\sqrt{n}(\hat{\theta} - \theta_0)}{\hat{\sigma}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{with} \quad \hat{\sigma} = \frac{\left[\frac{1}{n} \sum_{i=1}^n (Y_i - \hat{\beta} \cdot T_i - \hat{\mu}(X_i))^2 (T_i - \hat{\pi}(X_i))^2 \right]^{1/2}}{\frac{1}{n} \sum_{i=1}^n (T_i - \hat{\pi}(X_i))^2}.$$

In this case, one can construct a confidence interval accordingly for inference.

2.5 Structure-aware lower bound

In this subsection, we establish a no-gap result between the structure-agnostic setting and the worst-case structure-aware setting under the same approximation and stochastic error budgets. Specifically, the error rate (1.2) is attainable for any black-box class $\mathcal{G}$ by Corollary 2.2, yet remains unavoidable for a sparse linear class with an orthonormal dictionary. Thus, relative to the worst-case structure-aware model with the same budgets, structure agnosticism incurs no additional loss in estimation error rate.

We consider the simplified setting where $X \sim \nu_d := \text{Unif}([0, 1]^d)$, and let $\mathcal{T} := \{f : \|f\|_{\infty, [0, 1]^d} \leq 3\}$. For a given pair of function classes $(\mathcal{F}_\mu, \mathcal{F}_\pi)$ for (μ_0, π_0) , define the associated family of distributions as

$$\begin{aligned} \mathcal{P}(\mathcal{F}_\mu, \mathcal{F}_\pi) = & \left\{ (X, T, Y) \sim \mathbb{P}_{\beta, v, \mu, \pi} \text{ where } \mu \in \mathcal{F}_\mu, \pi \in \mathcal{F}_\pi, (\beta, v) \in [-3, 3] \times [1/3, 3] : \right. \\ & X \sim \nu_d, \quad T = \pi(X) + u \text{ with } \mathbb{E}[u|X] = 0, \mathbb{E}[u^2|X] \equiv v, |T| \leq 3, \\ & \left. Y = T \cdot \beta + \mu(X) + \varepsilon \text{ with } \mathbb{E}[\varepsilon|X, T] = 0, |Y| \leq 3 \right\}. \end{aligned} \quad (2.7)$$

The target parameter is $\theta(\mathbb{P}) = \beta$ for $\mathbb{P} = \mathbb{P}_{\beta, v, \mu, \pi} \in \mathcal{P}(\mathcal{F}_\mu, \mathcal{F}_\pi)$. The restriction of the constant conditional variance function $\mathbb{E}[u^2|X] \equiv v$ is imposed to rule out the possibility that one can exploit additional structure in the treatment variance to attain a faster rate.

The following proposition is analogous to Proposition 3.1 in Gu et al. (2026): the approximation error product term follows similarly to the construction in Balakrishnan et al. (2023); Jin et al. (2025), while the lower bound of $s \log(p)/n$ uses the construction of sparse linear models in Gu et al. (2026).

Proposition 2.3. *For any $\delta_\mu^{\text{appr}}, \delta_\pi^{\text{appr}} \in [0, 1/3)$, and $p, s \in \mathbb{N}^+$ with $s \leq p$ and $s[\log(p/s) + 1] \leq n \leq p^{1/3}$, there exists a set of orthonormal bases $\{\phi_j(x)\}_{j=1}^p$ with $\mathbb{E}_{X \sim \nu_d}[\phi(X)\phi(X)^\top] = I_p$ for $\phi(X) = (\phi_1(X), \dots, \phi_p(X))^\top \in \mathbb{R}^p$ such that the function classes defined as*

$$\mathcal{F}_\mu = \mathcal{G}_{p,s} \cup \mathcal{B}(0, \delta_\mu^{\text{appr}}), \quad \text{and} \quad \mathcal{F}_\pi = \mathcal{G}_{p,s} \cup \mathcal{B}(0, \delta_\pi^{\text{appr}}),$$

with $\mathcal{G}_{p,s} := \{g(x) = \sum_{j=1}^p \alpha_j \phi_j(x) \in \mathcal{T}, \|\alpha\|_0 \leq s\}$ and $\mathcal{B}(g, \delta) = \{f \in \mathcal{T} : \|f - g\|_{L_2(\nu_d)} \leq \delta\}$ satisfies

$$\inf_{\hat{\theta}} \sup_{\mathbb{P} \in \mathcal{P}(\mathcal{F}_\mu, \mathcal{F}_\pi)} \mathbb{E}_{\mathbb{P}^{2n}} [|\hat{\theta} - \theta(\mathbb{P})|] \geq \frac{1}{211} \left[n^{-1/2} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + \frac{s \log(p)}{n} \right]. \quad (2.8)$$

Here the randomness in $\mathbb{E}_{\mathbb{P}^{2n}}$ is $2n$ i.i.d. observations from $\mathbb{P}$, and $\hat{\theta}$ is the function of the $2n$ observations.

Because $0 \in \mathcal{G}_{p,s}$, every function in $\mathcal{F}_\mu$ (resp., $\mathcal{F}_\pi$) can be approximated by functions in the class $\mathcal{G}_{p,s}$ with L_2 error at most δ_μ^{appr} (resp., δ_π^{appr}). Hence, Proposition 2.3 provides a lower bound in estimating the linear coefficient β_0 when both the outcome function and treatment function are approximately s -sparse linear models with misspecification error δ_μ^{appr} and δ_π^{appr} , respectively. Moreover, under the conditions of Proposition 2.3, i.e., $s \leq n \leq p^{1/3}$, the stochastic error of the class $\mathcal{G}_{p,s}$ satisfies $\delta_n^{\text{stoc}} \asymp \sqrt{s \log(p)/n}$; see for example, Lemma S1.1 in Gu et al. (2026). Hence Proposition 2.3 establishes the structure-agnostic optimality: no estimator can guarantee a uniformly smaller error rate than (1.2) using only the three budgets.

Combining the lower bound in Proposition 2.3 and the upper bound in Corollary 2.2 yields the promised no-gap claim. For any hypothesis class $\mathcal{G}$ with fixed approximation errors and stochastic error budgets $(\delta_\mu^{\text{appr}}, \delta_\pi^{\text{appr}}, \delta_n^{\text{stoc}})$, Corollary 2.2 gives the estimation error

$$\frac{1}{\sqrt{n}} + \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + (\delta_n^{\text{stoc}})^2,$$

while the above error rate is not improvable even when restricting $\mathcal{G}$ to be a sparse linear model $\mathcal{G}_{p,s}$ with $\delta_n^{\text{stoc}} \asymp \sqrt{s \log(p)/n}$ as in Proposition 2.3. Therefore, the optimal structure-agnostic rate matches the worst-case structure-aware rate under compatible error budgets. Hence, being agnostic to nuisance structure incurs no additional statistical cost. This extends the no-gap result of Gu et al. (2026), which focuses on the regime $\delta_\pi^{\text{appr}} \asymp 1$, to the general regime under which both δ_μ^{appr} and δ_π^{appr} may vary.

3 Transductive Adversarial Moment-calibrated Editing

Section 2 characterizes the best attainable performance for a generic black-box model and serves as a proof of concept for the general TAME framework. There, the same class $\mathcal{G}$ both produces the initial nuisance estimates and indexes the directions used to calibrate the edited debiasing weights. In this section, we decouple these two roles, allowing arbitrary initial black-box estimates to be calibrated against a separately chosen reference class $\mathcal{G}_0$.

The two cases in Section 2 then become two branches. The O-branch applies when the outcome regression function is easier to estimate and fine-tunes the treatment-side debiasing weights induced by the treatment estimate. The R-branch applies when the treatment regression function is easier to estimate and fine-tunes the outcome-side debiasing weights induced by the initial estimates.

Section 3.1 introduces the design principles and generic TAME algorithm. Section 3.2 establishes an oracle-type inequality for arbitrary initial estimates and a separately chosen calibration class. Section 3.3

specializes this result to two nuisance learners and compares TAME-2 with DML. Section 3.4 then uses neural network learners to illustrate the under-smoothing principle for model selection and derive explicit rate comparisons. The resulting bounds show that TAME-1 and TAME-2 can improve on the standard DML guarantee when the nuisance difficulties are imbalanced, and that TAME-2 can further improve on TAME-1.

3.1 Design principles and generic algorithm

Let $\mathcal{D}_1 = \{(X_i, T_i, Y_i)\}_{i=1}^n$ and $\mathcal{D}_2 = \{(X_i, T_i, Y_i)\}_{i=n+1}^{2n}$. TAME takes as input initial estimates $(\hat{\mu}, \hat{\beta}, \hat{\pi})$ that are independent of $\mathcal{D}_1$, together with a reference class $\mathcal{G}_0$ used to index the adversarial moment constraints. The reference class need not be the class used to construct the initial estimates, although the oracle-type inequality requires it to contain the estimated easier nuisance.

Given a determination of which nuisance is easier, TAME use the corresponding branch $b \in \{0, R\}$: the 0-branch (resp., R-branch) is intended for the case where the outcome (resp. treatment) regression function is estimated more accurately. Its central difference from DML is an additional post-prediction fine-tuning procedure: TAME retains the component associated with the easier nuisance but fine-tunes the score component (which we referred to as debiasing weights) induced by the harder nuisance, rather than using it directly as in DML. Specifically, the 0-branch retains the outcome estimate $\hat{\mu}$ and fine-tunes the treatment-side weights $\tilde{a}$, whereas the R-branch retains the treatment estimate $\hat{\pi}$ and fine-tunes the outcome-side weights $\tilde{m}$. The fine-tuning remains anchored at the corresponding black-box initial estimates while imposing adversarial moment constraints indexed by the reference class $\mathcal{G}_0$. Particularly, TAME is guided by a symbiosis of two criteria:

- (C1) (*Local editing*) In each branch, TAME starts from the same debiasing weights that DML would use for the harder nuisance. A centered ridge penalty discourages deviation from this initialization, so that the fine-tuning preserves good properties of the initial black-box estimate. In the two-learner specialization, this retains

$$(\text{approx. error for the easier nuisance}) \times (L_2 \text{ error for the harder nuisance}). \quad (3.1)$$

- (C2) (*Transductive adversarial moment calibration*) Rather than leaving the harder-side weights at their initial value, TAME edits them directly on the inference sample to approximately satisfy the moment constraints indexed by $\mathcal{G}_0$. The weights are represented by an unrestricted vector in $\mathbb{R}^n$, optimized directly on the inference sample rather than represented by a fitted function of controlled complexity. Its statistical behavior is instead governed by adversarial moment constraints indexed by $\mathcal{G}_0$. In the 0-branch, the edited weights use $\{(X_i, T_i)\}_{i=1}^n$, but not $\{Y_i\}_{i=1}^n$, from $\mathcal{D}_1$. In the R-branch, the moment constraints use the outcome $\{Y_i\}_{i=n+1}^{2n}$ in $\mathcal{D}_2$, while the edited values depend on $\mathcal{D}_1$ only through the covariates $\{X_i\}_{i=1}^n$. These constructions preserve the required conditional mean-zero relations. In the two-learner specialization, moment calibration replaces the DML propagation of nuisance errors, i.e., the product of the two nuisance L_2 errors, by the term in (3.1) plus

$$(L_2 \text{ error for the easier nuisance})^2.$$

The three parts of the name describe the procedure directly: it is *transductive*, mirroring transductive learning, because it directly optimizes observation-specific debiasing weights on the inference sample rather than learning a function intended to generalize to unseen observations, *adversarially moment-calibrated* because it enforces worst-case moments over the entire class $\partial\mathcal{G}_0$, and an *editing* procedure because the optimized weights remain close to the initial black-box-induced weights.

Algorithm 1 can be run with any initial estimates that are independent of $\mathcal{D}_1$. For the oracle-type inequality in Section 3.2, the reference class is required to contain the estimate of the easier nuisance: $\hat{\mu} \in \mathcal{G}_0$

Algorithm 1 TAME for partial linear model

1: **Input:** Data split: $\mathcal{D}_1 = \{(X_i, T_i, Y_i)\}_{i=1}^n$, $\mathcal{D}_2 = \{(X_i, T_i, Y_i)\}_{i=n+1}^{2n}$; $\hat{\mu}, \hat{\beta}, \hat{\pi}$ independent of $\mathcal{D}_1$, debiasing reference class $\mathcal{G}_0$; branch choice $b \in \{0, R\}$, hyper-parameter $\lambda > 0$.

2: **if** $b = 0$ **then** ▷ Branch 0: the outcome part is easier

3: Initialize debiasing weights $a \in \mathbb{R}^n$ with $a_i \leftarrow \tilde{a}_i := \left[\left\{ \frac{1}{n} \sum_{\ell=1}^n T_\ell \{T_\ell - \hat{\pi}(X_\ell)\} \right\} \vee \frac{1}{2c} \right]^{-1} \{T_i - \hat{\pi}(X_i)\}$.

4: Fine-tune weights a by adversarial estimation on $(\mathcal{G}_0, \mathcal{D}_1)$ and get fine-tuned weights $\hat{a}$:

$$\hat{a} \in \operatorname{argmin}_{a \in \mathbb{R}^n} \frac{\lambda}{n} \sum_{i=1}^n (a_i - \tilde{a}_i)^2 + \sup_{\beta \in \mathbb{R}, f \in \partial \mathcal{G}_0} \Phi_\mu(f, \beta, a),$$

where $\Phi_\mu(f, \beta, a) := \left| \frac{1}{n} \sum_{i=1}^n (\beta \cdot T_i + f(X_i)) \cdot a_i - \beta \right| - \frac{1}{n} \sum_{i=1}^n f^2(X_i)$.

5: Assign $\hat{\theta} \leftarrow \hat{\theta}_0 = \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{\mu}(X_i)) \cdot \hat{a}_i$.

6: **else** ▷ Branch R: the treatment part is easier

7: Initialize debiasing weights $m \in \mathbb{R}^n$ with $m_i \leftarrow \tilde{m}_i := \hat{\beta} \cdot \hat{\pi}(X_i) + \hat{\mu}(X_i)$.

8: Fine-tune weights m by adversarial estimation on $(\mathcal{G}_0, \mathcal{D}_1, \mathcal{D}_2)$ and get fine-tuned weights $\hat{m}$:

$$\hat{m} \in \operatorname{argmin}_{m \in \mathbb{R}^n} \frac{\lambda}{n} \sum_{i=1}^n (m_i - \tilde{m}_i)^2 + \sup_{f \in \partial \mathcal{G}_0} \Phi_\pi(f, m),$$

where $\Phi_\pi(f, m) := \left| \frac{1}{n} \sum_{i=n+1}^{2n} Y_i f(X_i) - \frac{1}{n} \sum_{i=1}^n m_i f(X_i) \right| - \frac{1}{n} \sum_{i=1}^n f^2(X_i)$.

9: Assign $\hat{\theta} \leftarrow \hat{\theta}_R = [(2c)^{-1} \vee \frac{1}{n} \sum_{i=1}^n (T_i - \hat{\pi}(X_i))^2]^{-1} \left[\frac{1}{n} \sum_{i=1}^n (Y_i - \hat{m}_i)(T_i - \hat{\pi}(X_i)) \right]$.

10: **end if**

11: **Output:** The final estimate $\hat{\theta}$.

in the 0-branch and $\hat{\pi} \in \mathcal{G}_0$ in the R-branch. When the initial estimates are obtained from (S0) and $\mathcal{G}_0 = \mathcal{G}$, the 0- and R-branches reduce to $\hat{\theta}_\mu$ and $\hat{\theta}_\pi$, respectively. Thus, TAME-1 is a special case of the generic algorithm.

3.2 A generic oracle-type inequality

Theorem 3.1 gives an oracle-type inequality for **Algorithm 1** with arbitrary initial estimates $(\hat{\mu}, \hat{\beta}, \hat{\pi})$ and a separately chosen reference class $\mathcal{G}_0$. The result separates the errors of the initial estimates from the approximation and stochastic errors associated with $\mathcal{G}_0$, thereby making explicit the contributions of local editing and adversarial moment calibration. For concision, we state the result under an oracle choice of λ ; see the full error with explicit dependence on $\lambda > 0$ in the proof.

Condition 3.1 (Reference class and initial estimates). *The following holds:*

(a) As in **Definition 2**, $\mathcal{G}_0 \in \mathcal{HCS}(\delta_n^{\text{stoc}}; n, \nu_x)$, where ν_x is the distribution of X .

(b) $\hat{\mu}, \hat{\pi}, \hat{\beta}$ are uniformly bounded, i.e., $\|\hat{\mu}\|_\infty \vee \|\hat{\pi}\|_\infty \vee |\hat{\beta}| \leq c$, and are independent of $\mathcal{D}_1$.

Theorem 3.1. Assume **Condition 2.1** and **Condition 3.1**. Recall $\hat{\varphi}_{\text{PLM}}$ defined in (2.6). Then there exists a constant $\tilde{C} = \text{poly}(c)$ such that if $\delta_n^{\text{stoc}} \leq 1/\tilde{C}$, then for any $t \in [1, n/\tilde{C}]$, with probability at least $1 - [2e^{-t} + \exp(-n(\delta_n^{\text{stoc}})^2)]$, the following holds:

(O) If $\hat{\mu} \in \mathcal{G}_0$ and $\|\hat{\pi} - \pi_0\|_2 \leq \delta_\pi$ for $\delta_\pi \in \mathbb{R}^+$, then **Algorithm 1** with branch $b = 0$ and $\lambda \asymp (\delta_n^{\text{stoc}}/\delta_\pi)^2$

returns $\hat{\theta}$ satisfying

$$\begin{aligned} \tilde{C}^{-1}|\hat{\theta} - \theta_0 - \hat{\varphi}_{\text{PLM}}| \leq \inf_{g \in \mathcal{G}_0} \left[\|g - \mu_0\|_2 \cdot \delta_\pi + \|g - \hat{\mu}\|_2^2 + [\delta_n^{\text{stoc}}]^2 \right. \\ \left. + \sqrt{\frac{t}{n}} \left\{ \delta_\pi + \|g - \mu_0\|_2 + (1 + \lambda) \sqrt{\frac{t}{n}} \right\} \right]. \end{aligned} \quad (3.2)$$

(R) If $\hat{\pi} \in \mathcal{G}_0$ and $\|\hat{\mu} - \mu_0\|_2 + \|\hat{\pi} - \pi_0\|_2 + |\hat{\beta} - \beta_0| \leq \delta_{\text{all}}$ with $\delta_{\text{all}} \in \mathbb{R}^+$, then Algorithm 1 with branch $b = \text{R}$ and $\lambda \asymp (\delta_n^{\text{stoc}}/\delta_{\text{all}})^2$ returns $\hat{\theta}$ satisfying

$$\begin{aligned} \tilde{C}^{-1}|\hat{\theta} - \theta_0 - \hat{\varphi}_{\text{PLM}}| \leq \inf_{g \in \mathcal{G}_0} \left[\|g - \pi_0\|_2 \cdot \delta_{\text{all}} + \|g - \hat{\pi}\|_2^2 + [\delta_n^{\text{stoc}} + \|\hat{\pi} - \pi_0\|_2]^2 \right. \\ \left. + \sqrt{\frac{t}{n}} \left\{ \delta_{\text{all}} + \|g - \pi_0\|_2 + (1 + \lambda) \sqrt{\frac{t}{n}} \right\} \right]. \end{aligned} \quad (3.3)$$

For both branches, the first blue term is the direct result of the criterion [C1]: it multiplies the approximation error of the reference class by the L_2 error of the initial estimate on the opposite side. The subsequent red quadratic terms arise from the adversarial moment calibration in criterion [C2]. The last gray term is of order $o(n^{-1/2})$ when consistent estimations of μ_0 and π_0 are attained and $\lambda = o(\sqrt{n})$. In the TAME-2 specialization below, the tuning hyperparameter $\lambda \leq 1$, so this additional requirement is automatic. Their roles will be more explicit in the next subsection, when we use two black-box models to estimate μ_0 and π_0 separately.

The error bound for one black-box learner in Corollary 2.2 is a special case of the above theorem by picking g in (3.2) (resp. (3.3)) as the near-best approximation of μ_0 (resp. π_0) in the class $\mathcal{G} = \mathcal{G}_0$.

3.3 TAME-2: a two-learner specialization

When separate black-box models $\mathcal{G}_\mu$ and $\mathcal{G}_\pi$ are used to estimate μ_0 and π_0 , respectively, a natural specialization of our TAME method, which we refer to as TAME-2 estimator, first obtains $(\hat{\mu}, \hat{\beta})$ and $\hat{\pi}$ by (joint) least squares. It then uses the branch 0 (resp. R) with reference hypothesis class $\mathcal{G}_0 \leftarrow \mathcal{G}_\mu$ (resp. $\mathcal{G}_0 \leftarrow \mathcal{G}_\pi$) when the estimation of the outcome (resp., treatment) function is more accurate.

In this subsection, we derive the resulting rate and compare it with the standard DML guarantee. TAME-2 matches the DML guarantee up to constants and improves it when the nuisance difficulties are imbalanced and the easier learner is appropriately under-smoothed.

We first formalize the approximation and stochastic error budgets of the two learners.

Condition 3.2. Let ν_x be the distribution of X . Assume $\mathcal{G}_\mu \in \mathcal{HCS}(\delta_\mu^{\text{stoc}}; n, \nu_x)$, $\mathcal{G}_\pi \in \mathcal{HCS}(\delta_\pi^{\text{stoc}}; n, \nu_x)$. Define

$$\delta_\mu^{\text{appr}} := \inf_{g \in \mathcal{G}_\mu} \|g - \mu_0\|_{L_2(\nu_x)} \quad \text{and} \quad \delta_\pi^{\text{appr}} := \inf_{g \in \mathcal{G}_\pi} \|g - \pi_0\|_{L_2(\nu_x)}.$$

Given the split $\mathcal{D}_1 \cup \mathcal{D}_2$, both our TAME-2 estimator and DML estimator use the following initial estimates obtained from $\mathcal{D}_2$:

$$(\hat{\beta}, \hat{\mu}) \in \underset{\substack{\beta \in [-c, c] \\ g \in \mathcal{G}_\mu}}{\text{argmin}} \frac{1}{n} \sum_{i=n+1}^{2n} (Y_i - \beta \cdot T_i - g(X_i))^2, \quad \hat{\pi} \in \underset{g \in \mathcal{G}_\pi}{\text{argmin}} \frac{1}{n} \sum_{i=n+1}^{2n} (T_i - g(X_i))^2. \quad (3.4)$$

It follows from the standard empirical process argument that, with high probability,

$$\|\hat{\mu} - \mu_0\|_2 + |\hat{\beta} - \beta_0| \lesssim \delta_\mu^{\text{appr}} + \delta_\mu^{\text{stoc}} =: \delta_\mu \quad \text{and} \quad \|\hat{\pi} - \pi_0\|_2 \lesssim \delta_\pi^{\text{appr}} + \delta_\pi^{\text{stoc}} =: \delta_\pi.$$

Using the initial estimates in (3.4), the standard DML estimator $\hat{\theta}_{\text{DML}}$ is:

$$\hat{\theta}_{\text{DML}} = \left\{ \frac{1}{n} \sum_{l=1}^n T_l(T_l - \hat{\pi}(X_l)) \right\}^{-1} \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{\mu}(X_i))(T_i - \hat{\pi}(X_i)). \quad (3.5)$$

Using the upper error budgets, define TAME-2 by the following branch and tuning choices:

$$\hat{\theta}_{\text{TAME-2}} = \begin{cases} \hat{\theta} \text{ from Algorithm 1 with } b = \mathbf{0}, \mathcal{G}_0 = \mathcal{G}_\mu, \lambda = (\delta_\mu^{\text{stoc}}/\delta_\pi)^2 & \text{if } \delta_\mu \leq \delta_\pi \\ \hat{\theta} \text{ from Algorithm 1 with } b = \mathbf{R}, \mathcal{G}_0 = \mathcal{G}_\pi, \lambda = (\delta_\pi^{\text{stoc}}/\delta_\mu)^2 & \text{if } \delta_\pi < \delta_\mu \end{cases}. \quad (3.6)$$

The following corollary compares the nuisance contributed remainders of $\hat{\theta}_{\text{DML}}$ and $\hat{\theta}_{\text{TAME-2}}$.

Corollary 3.2. *Under Condition 2.1 and Condition 3.2, suppose further $\delta_\mu + \delta_\pi = o(1)$. Then there exists $R_{\text{DML}}, R_{\text{TAME-2}}$, satisfying $R_{\text{DML}} \vee R_{\text{TAME-2}} = o(n^{-1/2})$, such that, with high probability*

$$\begin{aligned} |\hat{\theta}_{\text{DML}} - \theta_0 - \hat{\varphi}_{\text{PLM}}| &\lesssim \delta_\mu \cdot \delta_\pi + R_{\text{DML}}, \\ |\hat{\theta}_{\text{TAME-2}} - \theta_0 - \hat{\varphi}_{\text{PLM}}| &\lesssim [\delta_\mu^{\text{appr}} \cdot \delta_\pi + (\delta_\mu)^2] \wedge [\delta_\pi^{\text{appr}} \cdot \delta_\mu + (\delta_\pi)^2] + R_{\text{TAME-2}}. \end{aligned} \quad (3.7)$$

To interpret (3.7), suppose first that $\delta_\mu \leq \delta_\pi$, so the outcome regression is easier and TAME-2 uses the 0-branch. DML has nuisance remainder $\delta_\mu \cdot \delta_\pi$, whereas TAME-2 has $\delta_\mu^{\text{appr}} \delta_\pi + \delta_\mu^2$. Since

$$\frac{\text{TAME-2 nuisance error}}{\text{DML nuisance error}} = \frac{\delta_\mu^{\text{appr}} \delta_\pi + \delta_\mu^2}{\delta_\mu \delta_\pi} = \frac{\delta_\mu^{\text{appr}}}{\delta_\mu} + \frac{\delta_\mu}{\delta_\pi},$$

and $\delta_\mu^{\text{appr}} \leq \delta_\mu \leq \delta_\pi$, TAME-2 is no worse than DML up to constants. Furthermore, if $\delta_\mu \ll \delta_\pi$ and $\delta_\mu^{\text{appr}} \ll \delta_\mu^{\text{stoc}}$, then the TAME-2 nuisance remainder is $o(\delta_\mu \delta_\pi)$. The second condition $\delta_\mu^{\text{appr}} \ll \delta_\mu^{\text{stoc}}$ is precisely the under-smoothing requirement for the easier outcome part. The treatment-easier case is symmetric. If, in addition, $\delta_\mu \delta_\pi \gg n^{-1/2}$, this reduction yields a strictly faster overall estimation rate than DML. Section 3.4 makes the comparison explicit for neural network learners.

3.4 Application to the neural network learners via under-smoothing

In this subsection, we instantiate the black-box learners by neural networks and use them to illustrate the under-smoothing principle (Rice, 1986). Let r_{DML} , $r_{\text{TAME-1}}$, and $r_{\text{TAME-2}}$ denote the optimally tuned convergence rate for estimating θ_0 , with poly-logarithmic factors suppressed. We show that

$$r_{\text{DML}} \gtrsim r_{\text{TAME-1}} \gtrsim r_{\text{TAME-2}}. \quad (3.8)$$

The conditions under which these comparisons are strict are given after Corollary 3.3. We first introduce the neural network (NN) class and nuisance function class. Then we present the estimator with optimal model selection and the resulting rates.

Neural network class. We consider the fully connected deep neural network with ReLU activation $\text{ReLU}(\cdot) = \max\{0, \cdot\}$. For positive integers L and N , a *deep ReLU network with depth L width N* admits the form of

$$g(x) = T_{L+1} \circ \overline{\text{ReLU}}_L \circ T_L \circ \text{ReLU}_{L-1} \circ \cdots \circ T_2 \circ \overline{\text{ReLU}}_1 \circ T_1(x). \quad (3.9)$$

Here $T_l(z) = W_l z + b_l : \mathbb{R}^{d_{l-1}} \rightarrow \mathbb{R}^{d_l}$ with $l \in [L+1]$ is an affine map with weight matrix $W_l \in \mathbb{R}^{d_l \times d_{l-1}}$ and bias vector $b_l \in \mathbb{R}^{d_l}$, where $(d_0, d_1, \dots, d_L, d_{L+1}) = (d, N, \dots, N, 1)$, and $\overline{\text{ReLU}}_l : \mathbb{R}^{d_l} \rightarrow \mathbb{R}^{d_l}$ applies the ReLU activation $\text{ReLU}(\cdot)$ to each entry of a d_l -dimensional vector for each $l \in [L]$. We define the neural network class as follows.

Definition 3 (Deep ReLU network class). *Define the family of deep ReLU networks with a d -dimensional input, depth L , width N , and truncation level B by*

$$\mathcal{F}_{\text{nn}}(d, L, N, B) = \{\tilde{g}(\cdot) = \text{Tc}_B(g(\cdot)) : g(\cdot) \text{ in (3.9)}\},$$

where $\text{Tc}_B : \mathbb{R} \rightarrow \mathbb{R}$ is the truncation operator defined as $\text{Tc}_B(z) = \min\{|z|, B\} \cdot \text{sign}(z)$.

Nuisance function class. Here we introduce the concept of the hierarchical composition model (Bauer & Kohler, 2019; Schmidt-Hieber, 2020; Kohler & Langer, 2021), a low-dimensional function class that neural networks can efficiently estimate.

Definition 4 (Hierarchical composition model $\mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}, C)$). *Let $\ell, d \in \mathbb{N}^+$, $C > 0$, and let $\mathcal{O} \subset [1, \infty) \times \mathbb{N}^+$ be a non-empty finite set. Define $\mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}, C)$ (Kohler & Langer, 2021) recursively as follows. Let $\mathcal{F}_{\text{HCM}}(d, 0, \mathcal{O}, C) = \{h(x) = x_j, j \in [d]\}$, and for each $\ell \geq 1$,*

$$\begin{aligned} \mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}, C) = \{ & h : \mathbb{R}^d \rightarrow \mathbb{R} : h(x) = g(f_1(x), \dots, f_r(x)), \text{ where} \\ & g \in \mathcal{F}_{\text{HS}}(r, \beta, C) \text{ with } (\beta, r) \in \mathcal{O} \text{ and } f_i \in \mathcal{F}_{\text{HCM}}(d, \ell-1, \mathcal{O}, C)\}. \end{aligned}$$

Here $\mathcal{F}_{\text{HS}}(d, \beta, C)$ denotes the set of all the d -variate (β, C) -Hölder smooth functions in Gu et al. (2025).

The class $\mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}, C)$ consists of finite compositions of r -variate functions with Hölder smoothness β , where $(\beta, r) \in \mathcal{O}$. Define the smallest dimension-adjusted smoothness (Fan & Gu, 2024) $\gamma^* = \min_{(\beta, r) \in \mathcal{O}} (\beta/r)$. For example, if $f(x) = f_1(x_1) + f_2(f_3(x_2, x_3), f_4(x_1, x_4), x_5)$ and all functions have smoothness $\beta = 2$, then the hardest component is f_2 with $(\beta, r) = (2, 3)$, resulting in $\gamma^* = 2/3$.

If f_0 lies in $\mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}, C)$, then the approximation error (Fan et al., 2024) and stochastic error for a neural network class $\mathcal{F}_{\text{nn}}(d, L, N, B)$ can be represented by its statistical complexity $S = (NL)^2$ as

$$\delta^{\text{appr}} \lesssim \text{poly}(\log(S)) \cdot S^{-\gamma^*} \quad \text{and} \quad \delta^{\text{stoc}} \lesssim \text{poly}(\log(Sn)) \cdot \sqrt{\frac{S}{n}}. \quad (3.10)$$

See, for example, proof of Theorem 2.1 in Gu et al. (2025). Hence, a neural network estimator $\hat{f}$ can optimally estimate the target regression function $f_0 \in \mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}, C)$ by choosing $S \asymp n^{\frac{1}{2\gamma^*+1}} =: \mathbb{S}(\gamma^*)$ that balances $\delta^{\text{appr}} \asymp \delta^{\text{stoc}}$, resulting in a L_2 estimation error $\|\hat{f} - f_0\|_2 \lesssim \text{poly}(\log(n)) \cdot n^{-\frac{\gamma^*}{2\gamma^*+1}}$, which matches the minimax optimal L_2 estimation risk over $\mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}, C)$.

Model selection for optimal convergence rate. We next impose regularity conditions under which the approximation error and stochastic error budgets in (3.10) apply to both nuisance functions. Condition 3.3 specifies the HCM structure, boundedness on the data generating process, and the neural network architectures used by DML, TAME-1, and TAME-2 estimators.

Condition 3.3. *Recall the partial linear model in (1.1), there exists a constant $c_1 > 1$ such that:*

(a) *Let $\mathcal{O}_\mu$ and $\mathcal{O}_\pi$ be two sets of smoothness-dimension indices, $\ell \in \mathbb{N}^+$, $C_0 \in \mathbb{R}^+$ such that*

$$\max_{(\beta, r) \in \mathcal{O}_\mu \cup \mathcal{O}_\pi} \{\beta \vee r\} \vee \ell \vee C_0 \vee d \leq c_1,$$

$f_0 \in \mathcal{F}_{\text{HCM}}(d, \ell, \mathcal{O}_f, C_0)$ with $\gamma_f^ = \min_{(\beta, r) \in \mathcal{O}_f} (\beta/r)$ for $f \in \{\mu, \pi\}$. We assume $\gamma_\mu^* \wedge \gamma_\pi^* > 0$.*

- (b) The nuisance functions and variables are uniformly bounded: $\|X\|_\infty \leq c_1$, $|\varepsilon| \vee |u| \leq c_1$, $|\beta_0| \vee \|\mu_0\|_\infty \vee \|\pi_0\|_\infty \leq c_1$, and $\mathbb{E}[u^2] \geq c_1^{-1}$.
- (c) We use $\mathcal{G}_1 = \mathcal{F}_{\text{nn}}(d, L_1, N_1, B)$, $\mathcal{G}_\mu = \mathcal{F}_{\text{nn}}(d, L_\mu, N_\mu, B)$ and $\mathcal{G}_\pi = \mathcal{F}_{\text{nn}}(d, L_\pi, N_\pi, B)$ with same $B = c_1$ but different depth and width parameters (L_q, N_q) for $q \in \{1, \mu, \pi\}$ satisfying $L_q \wedge N_q \geq c_2 \log(n)$ for some constant c_2 dependent on c_1 . We also denote $S_q = (N_q L_q)^2$ for $c \in \{1, \mu, \pi\}$.

The following corollary calculates the convergence rate for different methods under optimal model selection by plugging the choice of S into the approximation error and stochastic error budgets (3.10): the inequality (3.12) establishes the rate ordering in (3.8).

Corollary 3.3. Under Condition 3.3, recall $\mathbb{S}(\gamma^*) = n^{\frac{1}{2\gamma^*+1}}$, consider the following three estimators,

- (a) $\hat{\theta}_{\text{DML}, \text{NN}} \leftarrow \hat{\theta}_{\text{DML}}$ in (3.5) with $\mathcal{G}_\mu, \mathcal{G}_\pi$ satisfying $S_\mu \asymp \mathbb{S}(\gamma_\mu^*)$ and $S_\pi \asymp \mathbb{S}(\gamma_\pi^*)$.
- (b) $\hat{\theta}_{\text{TAME-1}, \text{NN}} \leftarrow \hat{\theta}$ in (2.3) with $\mathcal{G} = \mathcal{G}_1$ satisfying $S_1 \asymp n^{\frac{1}{\gamma_\mu^* + \gamma_\pi^* + 1}}$ and the optimal choice of λ in Theorem 2.1. Here S_1 balances the polynomial upper bounds of $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}}$ and $(\delta_n^{\text{stoc}})^2$ as $S_1^{-(\gamma_\mu^* + \gamma_\pi^*)} \asymp S_1/n$.
- (c) $\hat{\theta}_{\text{TAME-2}, \text{NN}} \leftarrow \hat{\theta}_{\text{TAME-2}}$ in (3.6) with $\mathcal{G}_\mu$ and $\mathcal{G}_\pi$ satisfying the following hyper-parameter choice

$$\begin{cases} S_\pi \asymp \mathbb{S}(\gamma_\pi^*), S_\mu \asymp n^{\frac{\gamma_\mu^*+1}{2\gamma_\mu^*+1} \cdot \frac{1}{\gamma_\mu^*+1}} \gg \mathbb{S}(\gamma_\mu^*) & \text{if } \gamma_\mu^* > \gamma_\pi^* \\ S_\mu \asymp \mathbb{S}(\gamma_\mu^*), S_\pi \asymp \mathbb{S}(\gamma_\pi^*) & \text{if } \gamma_\mu^* = \gamma_\pi^* \\ S_\mu \asymp \mathbb{S}(\gamma_\mu^*), S_\pi \asymp n^{\frac{\gamma_\mu^*+1}{2\gamma_\mu^*+1} \cdot \frac{1}{\gamma_\pi^*+1}} \gg \mathbb{S}(\gamma_\pi^*) & \text{if } \gamma_\pi^* > \gamma_\mu^* \end{cases} \quad (3.11)$$

For any $\mathbf{M} \in \{\text{DML}, \text{TAME-1}, \text{TAME-2}\}$, $\hat{\theta}_{\mathbf{M}, \text{NN}}$ satisfies $|\hat{\theta}_{\mathbf{M}, \text{NN}} - \theta_0| \lesssim n^{-1/2} + \text{poly}(\log(n)) \cdot n^{-\alpha_{\mathbf{M}}}$ with high probability, where

$$\underbrace{\frac{\gamma_\mu^*}{2\gamma_\mu^*+1} + \frac{\gamma_\pi^*}{2\gamma_\pi^*+1}}_{=: \alpha_{\text{DML}}} \leq \underbrace{\frac{\gamma_\mu^* + \gamma_\pi^*}{\gamma_\mu^* + \gamma_\pi^* + 1}}_{=: \alpha_{\text{TAME-1}}} \leq \underbrace{\frac{s^*}{s^* + \{\gamma_\mu^* \wedge \gamma_\pi^*\} + 1}}_{=: \alpha_{\text{TAME-2}}} \quad (3.12)$$

and where $s^* = 2\gamma_\mu^* \gamma_\pi^* + \gamma_\mu^* + \gamma_\pi^*$, with both equalities in (3.12) holding if and only if $\gamma_\mu^* = \gamma_\pi^*$.

Remark 3.1 (Optimal model selection for DML). For DML estimator, the nuisance errors enter in the final error as $\delta_\mu \cdot \delta_\pi$ in Corollary 3.2, which means the model selections for $\mathcal{G}_\mu$ and $\mathcal{G}_\pi$ should match that for optimal estimations of μ_0 and π_0 , i.e., $\delta_\mu^{\text{appr}} \asymp \delta_\mu^{\text{stoc}}$ and $\delta_\pi^{\text{appr}} \asymp \delta_\pi^{\text{stoc}}$.

Remark 3.2 (Optimal model selection for TAME). For TAME-1, the nuisance error budgets enter in the final error as $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + (\delta_n^{\text{stoc}})^2$, which suggests the corresponding model selection rule $\delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} \asymp (\delta_n^{\text{stoc}})^2$. This choice under-smooths the easier nuisance, but over-smooths the harder nuisance. TAME-2 avoids the drawbacks introduced by the latter restriction by using separate neural networks. To see this, when $\gamma_\mu^* > \gamma_\pi^*$, i.e., the outcome is easier to estimate, Corollary 3.2 gives the oracle-type error bound $(\delta_\pi^{\text{appr}} + \delta_\pi^{\text{stoc}}) \cdot \delta_\mu^{\text{appr}} + \delta_\mu^2$, this suggests doing standard bias-variance tradeoff on the harder π_0 part $\delta_\pi^{\text{appr}} \asymp \delta_\pi^{\text{stoc}}$ and further under-smoothing on easier μ_0 part $\delta_\mu^{\text{appr}} \cdot \delta_\pi^* \asymp (\delta_\mu^{\text{stoc}})^2$ on top of the balanced $\delta_\pi^* := \delta_\pi^{\text{appr}} \asymp \delta_\pi^{\text{stoc}}$ as in the first line of (3.11). The mirror case $\gamma_\pi^* \geq \gamma_\mu^*$ follows similarly.

Remark 3.3 (The case where $\alpha_{\mathbf{M}}$ coincides). Under Condition 3.3, both inequalities in (3.12) are strict unless $\gamma_\mu^* = \gamma_\pi^*$. Thus, at the nonparametric rate level, TAME-1 strictly improves on DML, and TAME-2 strictly improves on TAME-1 when the nuisance difficulties are imbalanced. These exponent gains further yield strict improvements in the overall estimation error upper bound whenever the nonparametric term dominates $n^{-1/2}$.

4 Extensions beyond the partial linear model

This section extends TAME beyond the partial linear model. We discuss the extension of 0-branch to a broad class of linear functional estimation problems in [Section 4.1](#). The extension of R-branch for estimating average treatment effect (ATE) is discussed in [Section 4.2](#). With a slight abuse of notation, we adopt μ, θ with different semantic meanings than those for the partial linear model before.

4.1 The 0-branch for general linear functionals

Setup. Consider the following general model for $(X, T, Y) \in \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}$:

$$Y = \mu_0(X, T) + \varepsilon \quad \text{with} \quad \mu_0 \in \mathcal{H}_\mu, \quad \mathbb{E}[\varepsilon|X, T] = 0, \quad (4.1)$$

where $\mathcal{H}_\mu$ is a Hilbert space equipped with the inner product $\langle h_1, h_2 \rangle = \mathbb{E}_{\nu_z}[h_1(Z)h_2(Z)]$, and where ν_z is the distribution of $Z = (X, T) \in \mathbb{R}^d \times \mathbb{R}$. Given $2n$ observations $\mathcal{D} = \{(Z_i, Y_i)\}_{i=1}^{2n}$ drawn i.i.d. from the above model, the goal is to estimate the following linear functional

$$\theta_0 = \theta(\mu_0) \quad \text{with} \quad \theta(\mu) = \mathbb{E}[\psi(\mu, Z)] \quad (4.2)$$

for a known operator $\psi : \mathcal{H}_\mu \times \mathbb{R}^{d+1} \rightarrow \mathbb{R}$. We focus on the setting under which $\theta(\cdot)$ is linear and the map $\mu \mapsto \psi(\mu, \cdot)$ is Lipschitz from $L_2(\nu_z)$ to $L_2(\nu_z)$, as formalized below.

Condition 4.1 (Linearity and continuity). *There exists a constant $c \geq 1$ such that the following condition holds for $\theta(\cdot)$ on $\mathcal{H}_\mu$:*

- (a) For any $\mu, \tilde{\mu} \in \mathcal{H}_\mu$ and $\alpha \in \mathbb{R}$, $\theta(\alpha\mu + \tilde{\mu}) = \alpha \cdot \theta(\mu) + \theta(\tilde{\mu})$;
- (b) For any $\mu, \tilde{\mu} \in \mathcal{H}_\mu$, $\mathbb{E}[\|\psi(\mu, Z) - \psi(\tilde{\mu}, Z)\|^2] \leq c\mathbb{E}[\|\mu(Z) - \tilde{\mu}(Z)\|^2]$.

[Condition 4.1](#) implies that θ is a linear continuous functional. Indeed, $\theta(0) = 0$ by the linearity of $\theta(\cdot)$, and [Condition 4.1](#) (b) then implies that, for any $\mu \in \mathcal{H}_\mu$,

$$|\theta(\mu)| = |\mathbb{E}[\psi(\mu, Z) - \psi(0, Z)]| \leq \{\mathbb{E}[\|\psi(\mu, Z) - \psi(0, Z)\|^2]\}^{1/2} \leq c^{1/2}\|\mu\|_{L_2(\nu_z)}.$$

The Riesz representer theorem therefore yields a unique Riesz representer $r_0 \in \mathcal{H}_\mu$ for the linear functional $\theta(\cdot)$ such that

$$\forall \mu \in \mathcal{H}_\mu, \quad \theta(\mu) = \mathbb{E}[\psi(\mu, Z)] = \mathbb{E}[\mu(Z) \cdot r_0(Z)]. \quad (4.3)$$

The partial linear model in [Section 2](#) is an instance of the above general framework with $\mathcal{H}_\mu = \{\mu(X, T) = \beta \cdot T + f(X) : \beta \in \mathbb{R}, f \in L_2(\nu_x)\}$, $\psi(\mu, Z) = \beta$ and Riesz representer $r_0(X, T) = (T - \mathbb{E}[T|X])/\mathbb{E}[(T - \mathbb{E}[T|X])^2]$.

Method and main results. In a similar spirit to [Section 4](#) of [Gu et al. \(2026\)](#), [Algorithm 2](#) extends the 0-branch in [Algorithm 1](#) by initializing the debiasing weights with an arbitrary estimate $\hat{r}$ of the Riesz representer and then fine-tuning these weights against the reference class $\mathcal{G}_0$.

Algorithm 2 TAME for linear functional $\theta(\cdot)$, 0-branch

- 1: **Input:** $\mathcal{D}_1 = \{(Z_i, Y_i)\}_{i=1}^n$; $\hat{\mu}, \hat{r}$ independent of $\mathcal{D}_1$, debiasing reference class $\mathcal{G}_0$, hyper-parameter λ .
- 2: Initialize debiasing weights $a \in \mathbb{R}^n$ with $a_i \leftarrow \tilde{a}_i$ where $\tilde{a}_i = \hat{r}(Z_i)$.
- 3: Fine-tune weights a by adversarial estimation on $(\mathcal{G}_0, \mathcal{D}_1)$ and get fine-tuned weights $\hat{a}$:

$$\hat{a} \in \operatorname{argmin}_{a \in \mathbb{R}^n} \frac{\lambda}{n} \sum_{i=1}^n |a_i - \tilde{a}_i|^2 + \sup_{f \in \partial \mathcal{G}_0} \left| \frac{1}{n} \sum_{i=1}^n \{\psi(f, Z_i) - a_i \cdot f(Z_i)\} \right| - \frac{1}{n} \sum_{i=1}^n f^2(Z_i)$$

- 4: **Output:** The final estimate $\hat{\theta} \leftarrow \frac{1}{n} \sum_{i=1}^n \{\psi(\hat{\mu}, Z_i) + (Y_i - \hat{\mu}(Z_i)) \cdot \hat{a}_i\}$.
-

Remark 4.1 (Implementing the initial estimate). *The algorithm can be applied to any initial black-box estimates $\hat{\mu}, \hat{r}$ of the outcome function μ_0 and the Riesz representer r_0 . For example, $\hat{\mu}$ may be obtained by fitting Y on Z via least squares over a hypothesis class $\mathcal{G}_\mu$ on $\mathcal{D}_2 = \{(Z_i, Y_i)\}_{i=n+1}^{2n}$. The Riesz representer $\hat{r}$ may be estimated either by plugging in the estimated nuisance functions into a known analytic expression or directly via adversarial Riesz estimation as in [Chernozhukov et al. \(2026\)](#).*

The following conditions allow the 0-branch oracle-type inequality, i.e., (3.2) in [Theorem 3.1](#), to extend to the general linear functional θ .

Condition 4.2. *There exists a constant $c \geq 1$ such that:*

- (a) (Boundedness) $|\varepsilon| \leq c$, $\|r_0\|_\infty \vee \|\mu_0\|_\infty \leq c$, $\|\psi(\mu_0, \cdot)\|_\infty \leq c$.
- (b) (Function class $\mathcal{G}_0$) The reference class $\mathcal{G} \subset \mathcal{H}_\mu$ satisfies $\mathcal{G}_0 \in \mathcal{HCS}(\delta_n^{\text{stoc}}; n, \nu_z)$ and $\hat{\mu} \in \mathcal{G}_0$. Moreover, $\|\psi(f, \cdot)\|_\infty \leq c$ for any $f \in \partial\mathcal{G}_0$, and

$$\sup_{\delta \geq \delta_n^{\text{stoc}}} \frac{R_{n, \nu_z}(\delta; (\partial\mathcal{G}_0)^\psi)}{c \cdot \delta} \leq \delta_n^{\text{stoc}} \quad \text{for} \quad (\partial\mathcal{G}_0)^\psi := \{z \mapsto \psi(g - \tilde{g}, z) : g, \tilde{g} \in \mathcal{G}_0\}.$$

- (c) (Initial black-box estimate) $\hat{r}, \hat{\mu}$ are independent of $\mathcal{D}_1$ and uniformly bounded, i.e., $\|\hat{r}\|_\infty \vee \|\hat{\mu}\|_\infty \leq c$.

Theorem 4.1. *Under [Condition 4.1](#) and [Condition 4.2](#), if $\|\hat{r} - r_0\|_{L_2(\nu_z)} \leq \delta_r$ for $\delta_r \in \mathbb{R}^+$ and we choose $\lambda \asymp [\delta_n^{\text{stoc}}/(\delta_r + \delta_n^{\text{stoc}})]^2$, then for any $t \geq 1$, the estimate returned by [Algorithm 2](#) satisfies, with probability at least $1 - 2e^{-t} - e^{-n(\delta_n^{\text{stoc}})^2}$,*

$$\begin{aligned} \tilde{C}^{-1} |\hat{\theta} - \theta_0 - \hat{\varphi}_{\text{LF}, \theta}| &\leq \inf_{\bar{g} \in \mathcal{G}_0} (\delta_r + \delta_n^{\text{stoc}}) \cdot \|\bar{g} - \mu_0\|_2 + \|\bar{g} - \hat{\mu}\|_2^2 + (\delta_n^{\text{stoc}})^2 \\ &\quad + \left(\delta_r + \|\bar{g} - \mu_0\|_2 + \delta_n^{\text{stoc}} + \sqrt{\frac{t}{n}} \right) \cdot \sqrt{\frac{t}{n}} \end{aligned} \quad (4.4)$$

where $\tilde{C} = \text{poly}(c)$ and $\hat{\varphi}_{\text{LF}, \theta} = \frac{1}{n} \sum_{i=1}^n \psi(\mu_0; Z_i) - \theta_0 + \varepsilon_i r_0(Z_i)$ is the empirical average of an influence function for θ_0 . Under the additional condition $\mathbb{E}[\varepsilon^2 | Z] \equiv \sigma_\varepsilon^2$, this influence function is semi-parametrically efficient.

Remark 4.2 (Two-learner specialization). *Suppose $\hat{\mu}$ is an empirical least squares estimator over a hypothesis class $\mathcal{G}_\mu$ and we choose $\mathcal{G}_0 = \mathcal{G}_\mu$ in [Algorithm 2](#). Define $\delta_\mu^{\text{appr}} := \inf_{g \in \mathcal{G}_\mu} \|g - \mu_0\|_{L_2(\nu_z)}$, and $\delta_\mu := \delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}}$. If both $\hat{\mu}$ and $\hat{r}$ can consistently estimate μ_0 and r_0 , respectively, then (4.4) yields*

$$|\hat{\theta} - \theta_0 - \hat{\varphi}_{\text{LF}, \theta}| \lesssim \delta_r \cdot \delta_\mu^{\text{appr}} + (\delta_\mu)^2 + o(n^{-1/2}) \quad (4.5)$$

Recall that DML has the nuisance remainder $\delta_\mu \cdot \delta_r + o(n^{-1/2})$ ([Chernozhukov et al., 2018, 2026](#)). Thus, TAME is no worse up to constants when $\delta_\mu \leq \delta_r$. Moreover, when $\delta_\mu \ll \delta_r$, i.e., the outcome function is easier to estimate, the use of TAME estimator gives strict improvement over the DML nuisance remainder when under-smoothing on $\mathcal{G}_\mu$ is adopted, i.e., $\delta_\mu^{\text{appr}} \ll \delta_\mu \asymp \delta_n^{\text{stoc}}$. This yields a strictly faster overall rate whenever $\delta_\mu \cdot \delta_r \gg n^{-1/2}$.

Remark 4.3 (Asymptotic normality). *Continue with the two-learner specialization in [Remark 4.2](#), and suppose $\delta_r \cdot \delta_\mu^{\text{appr}} + (\delta_\mu)^2 = o(n^{-1/2})$. Define $\sigma_0 := [\text{Var}(\psi(\mu_0, Z_i) + \varepsilon \cdot r_0(Z))]^{1/2} > 0$, and the estimated influence values $\hat{\varphi}_i := \psi(\hat{\mu}, Z_i) - \hat{\theta} + (Y_i - \hat{\mu}(Z_i)) \cdot \hat{a}_i$. Then we have*

$$\frac{\sqrt{n}(\hat{\theta} - \theta_0)}{\hat{\sigma}} \xrightarrow{d} \mathcal{N}(0, 1) \quad \text{for} \quad \hat{\sigma} = \sqrt{\frac{1}{n} \sum_{i=1}^n \hat{\varphi}_i^2} \quad \text{satisfying} \quad |\hat{\sigma} - \sigma_0| = o_P(1).$$

4.2 The R-branch for ATE estimation

Let $\mathcal{D} = \{(X_i, T_i, Y_i)\}_{i=1}^{2n}$ be $2n$ i.i.d. observations generated from

$$\begin{aligned} T &= \pi_0(X) + u && \text{with } \mathbb{E}[u|X] = 0, T \in \{0, 1\}, \\ Y &= \mu_0(X, T) + \varepsilon && \text{with } \mathbb{E}[\varepsilon|X, T] = 0, \end{aligned} \quad (4.6)$$

where $X \in \mathcal{X} \subseteq \mathbb{R}^d$, and $\pi_0(x) = \mathbb{P}(T = 1|X = x)$ is the propensity score. The target is the linear functional $\theta_0 = \mathbb{E}[\mu_0(X, 1) - \mu_0(X, 0)]$. Under the usual consistency, unconfoundedness, and overlap assumptions, θ_0 equals the average treatment effect in the potential outcome framework

[Algorithm 3](#) is motivated by and extends the central construction of [Wang & Shah \(2024\)](#). Their debiased inverse propensity weighting (DIPW) estimator edits observation-specific outcome weights through cross-sample linear moment constraints under a sparse logistic propensity model. We replace these linear constraints with adversarial moment calibration over a generic reference class $\mathcal{G}_0$, allowing the same principle to be applied with generic black-box propensity learners. When $\hat{m}_i = \tilde{m}_i$, the estimator in Step 4 reduces algebraically to the usual AIPW estimator based on $(\hat{\mu}, \hat{\pi})$.

Algorithm 3 TAME for ATE, R-branch

- 1: **Input:** Data split: $\mathcal{D}_1 = \{(X_i, T_i, Y_i)\}_{i=1}^n$, $\mathcal{D}_2 = \{(X_i, T_i, Y_i)\}_{i=n+1}^{2n}$; $\hat{\mu}, \hat{\pi}$ independent of $\mathcal{D}_1$, debiasing reference class $\mathcal{G}_0$, hyper-parameter λ .
- 2: Initialize debiasing weights $m \in \mathbb{R}^n$ with $m_i \leftarrow \tilde{m}_i$ where $\tilde{m}_i = (1 - \hat{\pi}(X_i))\hat{\mu}(X_i, 1) + \hat{\pi}(X_i)\hat{\mu}(X_i, 0)$.
- 3: Fine-tune weights m by adversarial estimation on $(\mathcal{G}_0, \mathcal{D}_1, \mathcal{D}_2)$ and get fine-tuned weights $\hat{m}$:

$$\begin{aligned} \hat{m} \in \operatorname{argmin}_{m \in \mathbb{R}^n} & \frac{\lambda}{n} \sum_{i=1}^n |m_i - \tilde{m}_i|^2 + \sup_{g \in \mathcal{G}_0} \Phi_{\text{ATE}, \pi}(g, m) \quad \text{with} \\ \Phi_{\text{ATE}, \pi}(g, m) &= \left| \frac{1}{n} \sum_{i=n+1}^{2n} \hat{Y}_i \cdot \Delta_g(X_i) - \frac{1}{n} \sum_{i=1}^n m_i \cdot \Delta_g(X_i) \right| - \frac{1}{n} \sum_{i=1}^n (g - \hat{\pi})^2(X_i), \end{aligned}$$

where the pseudo-outcome $\hat{Y}_i$ and direction Δ_g are defined by $\hat{Y}_i = \frac{1 - \hat{\pi}(X_i)}{\hat{\pi}(X_i)} Y_i T_i + \frac{\hat{\pi}(X_i)}{1 - \hat{\pi}(X_i)} Y_i (1 - T_i)$ and $\Delta_g(X_i) = (g - \hat{\pi})(X_i) / [\hat{\pi}(X_i)(1 - \hat{\pi}(X_i))]$.

- 4: Assign $\hat{\theta} \leftarrow \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{m}_i) \left(\frac{T_i}{\hat{\pi}(X_i)} - \frac{1 - T_i}{1 - \hat{\pi}(X_i)} \right)$.
 - 5: **Output:** The final estimate $\hat{\theta}$.
-

The analysis of [Algorithm 3](#) uses two conditions. [Condition 4.3](#) imposes boundedness and overlap on the data-generating process, while [Condition 4.4](#) controls the reference class and the initial estimates.

Condition 4.3 (Boundedness and overlap). *There exists a constant $c \geq 2$ such that $\|\mu_0\|_\infty \vee |\varepsilon| \leq c$, and $\pi_0(x) \in [1/c, 1 - 1/c]$ for any $x \in \mathcal{X}$.*

Condition 4.4. *Let ν_x be the distribution of X , the following holds for a constant $c \geq 2$:*

- (a) *The reference class satisfies $\mathcal{G}_0 \in \mathcal{HCS}(\delta_n^{\text{stoc}}; n, \nu_x)$ and $\hat{\pi} \in \mathcal{G}_0$. Moreover, $g(x) \in [1/c, 1 - 1/c]$ for any $x \in \mathcal{X}$ and $g \in \mathcal{G}_0$.*
- (b) *The initial estimates $(\hat{\mu}, \hat{\pi})$ are independent of $\mathcal{D}_1 = \{(X_i, T_i, Y_i)\}_{i=1}^n$, and $\|\hat{\mu}\|_\infty \leq c$.*

[Theorem 4.2](#) gives the resulting oracle-type inequality for the estimator in [Algorithm 3](#).

Theorem 4.2. *Under [Condition 4.3](#) and [Condition 4.4](#), if the sum of first-stage estimation errors $\|\hat{\pi} - \pi_0\|_{L_2(\nu_x)} + \sup_{w \in \{0, 1\}} \|\hat{\mu}(\cdot, w) - \mu_0(\cdot, w)\|_{L_2(\nu_x)} \leq \delta_{\text{all}}$ for $\delta_{\text{all}} \in \mathbb{R}^+$ and we choose $\lambda \asymp [\delta_n^{\text{stoc}} / (\delta_{\text{all}} + \delta_n^{\text{stoc}})]^2$,*

then for any $t \in [1, n]$, the estimate $\hat{\theta}$ returned by [Algorithm 3](#) satisfies, with probability at least $1 - 3e^{-t} - e^{-n(\delta_n^{\text{stoc}})^2}$,

$$\begin{aligned} \tilde{C}^{-1}|\hat{\theta} - \theta_0 - \hat{\varphi}_{ATE}| &\leq \inf_{\bar{g} \in \mathcal{G}_0} (\delta_{all} + \delta_n^{\text{stoc}}) \cdot \|\bar{g} - \pi_0\|_2 + \|\bar{g} - \hat{\pi}\|_2^2 + \|\pi_0 - \hat{\pi}\|_2^2 + (\delta_n^{\text{stoc}})^2 \\ &\quad + \sqrt{\frac{t}{n}} \left[\delta_{all} + \delta_n^{\text{stoc}} + \|\bar{g} - \pi_0\|_2 + \|\hat{\pi} - \pi_0\|_2^{1/2} + \left(\frac{t}{n}\right)^{1/4} \right] \end{aligned} \quad (4.7)$$

where $\tilde{C} = \text{poly}(c)$ and $\hat{\varphi}_{ATE} = \frac{1}{n} \sum_{i=1}^n \{\mu_0(X_i, 1) - \mu_0(X_i, 0)\} + \varepsilon_i \cdot u_i / \{\pi_0(X_i) \cdot (1 - \pi_0(X_i))\} - \theta_0$ is the empirical average of the semiparametric efficient influence function for the ATE.

[Theorem 4.2](#) also yields direct analogues of [Remark 4.2](#) and [Remark 4.3](#). Suppose that $\hat{\pi}$ is an empirical least squares estimator over $\mathcal{G}_\pi = \mathcal{G}_0$, and define $\delta_\pi^{\text{appr}} := \inf_{g \in \mathcal{G}_\pi} \|g - \pi_0\|_{L_2(\nu_x)}$ and $\delta_\pi := \delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}}$. If the first stage errors are consistent, then (4.7) yields

$$|\hat{\theta} - \theta_0 - \hat{\varphi}_{ATE}| \lesssim \delta_{all} \cdot \delta_\pi^{\text{appr}} + \delta_\pi^2 + o(n^{-1/2}).$$

Consequently, asymptotic normality follows when the displayed nuisance remainder is $o(n^{-1/2})$.

5 Proof of Theorem 2.1

We first introduce notation used only in the proof of [Theorem 2.1](#). We let $\|f\|_2 = \sqrt{\mathbb{E}[f^2(X)]}$ and $\|f\|_n = \sqrt{\frac{1}{n} \sum_{i=1}^n f^2(X_i)}$, and adopt $\lesssim$ to omit the $\text{poly}(c)$ factors. The following proposition characterizes the estimation error of the initial estimate $(\hat{\beta}, \hat{\mu}, \hat{\pi})$, which is a restatement of Proposition S2.3 in [Gu et al. \(2026\)](#). The proof is standard; one can directly refer to the proof therein.

Proposition 5.1. *The following event occurs with probability at least $1 - e^{-n(\delta_n^{\text{stoc}})^2}$ for some constant $\tilde{C}_0 = \text{poly}(c)$,*

$$\mathcal{C}_0 := \left\{ |\hat{\beta} - \beta_0| + \|\hat{\mu} - \mu_0\|_2 \leq \tilde{C}_0 (\delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}}), \|\hat{\pi} - \pi_0\|_2 \leq \tilde{C}_0 (\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}}) \right\}.$$

Without loss of generality, we consider the case where $\delta_\mu^{\text{appr}} \geq n^{-100}$ and $\delta_\pi^{\text{appr}} \geq n^{-100}$. Otherwise, define $\bar{\delta}_\mu^{\text{appr}} \leftarrow \delta_\mu^{\text{appr}} + n^{-100}$ and $\bar{\delta}_\pi^{\text{appr}} \leftarrow \delta_\pi^{\text{appr}} + n^{-100}$, we can prove the corresponding branch-wise bound with $\bar{\delta}_h^{\text{appr}}$ in place of δ_h^{appr} , using the facts that (1) $\bar{\delta}_h^{\text{appr}}$ remains an upper bound on the corresponding approximation error; and (2) $\lambda \geq (\delta_n^{\text{stoc}})^2 \geq \log(n)/n$.

We will prove in the two subsequent subsections

$$\begin{aligned} |\hat{\theta}_\mu - \theta_0 - \hat{\varphi}_{PLM}| &\lesssim \delta^* + \lambda(\delta_\pi^{\text{appr}})^2 + \frac{\delta_n^{\text{stoc}} \cdot \delta_\mu^{\text{appr}}}{\sqrt{\lambda}} \quad \text{if } \delta_\mu^{\text{appr}} \leq \delta_\pi^{\text{appr}} \\ |\hat{\theta}_\pi - \theta_0 - \hat{\varphi}_{PLM}| &\lesssim \delta^* + \lambda(\delta_\mu^{\text{appr}})^2 + \frac{\delta_n^{\text{stoc}} \cdot \delta_\pi^{\text{appr}}}{\sqrt{\lambda}} \quad \text{if } \delta_\mu^{\text{appr}} > \delta_\pi^{\text{appr}} \end{aligned}$$

where $\delta^* = \delta_\pi^{\text{appr}} \cdot \delta_\mu^{\text{appr}} + (\delta_n^{\text{stoc}})^2 + \sqrt{\frac{t}{n}} \cdot \left[\delta_\mu^{\text{appr}} + \delta_\pi^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}} \right]$. Combining the separate error bounds with the estimator choice criterion (2.3) concludes the proof.

5.1 Proof of CASE 1

Denote $H_\mu(a) = \sup_{\beta, f \in \partial \mathcal{G}} \Phi_\mu(f, \beta, a)$ for any $a \in \mathbb{R}^n$ and $\bar{D} = \frac{1}{n} \sum_{l=1}^n T_l(T_l - \pi_0(X_l))$. Define $\bar{a} \in \mathbb{R}^n$ be such that $\bar{a}_i = (T_i - \pi_0(X_i))/\bar{D}$ if $\bar{D} \neq 0$ and $\bar{a}_i \equiv 0$ otherwise. With a slight abuse of notation, we will

also use $\|a - a'\|_n = \{\frac{1}{n} \sum_{i=1}^n (a_i - a'_i)^2\}^{1/2}$ for $a, a' \in \{\bar{a}, \tilde{a}, w, \hat{a}\} \subseteq \mathbb{R}^n$. We first pick $\bar{g} \in \mathcal{G}$ such that $\|\bar{g} - \mu_0\|_2 \leq 2\delta_\mu^{\text{appr}}$ by the definition of the approximation error in (2.2). Here, we first define the high-probability event used to derive our error rate, which follows directly from the standard empirical process arguments and concentration inequalities.

Proposition 5.2. *Under the event $\mathcal{C}_0$ in Proposition 5.1, assume $\delta_n^{\text{stoc}} \tilde{C}_0 \leq 1$ for some $\tilde{C}_0 = \text{poly}(c)$. For any $t \in [1, n]$, the following event occurs with probability at least $1 - e^{-n(\delta_n^{\text{stoc}})^2} - e^{-t}$ for some $\tilde{C}_1 = \text{poly}(c)$:*

$$\mathbf{H}_\mu(\bar{a}) \leq \tilde{C}_1 [\delta_n^{\text{stoc}}]^2, \quad (5.1)$$

$$\|\bar{g} - \hat{\mu}\|_n^2 \leq \tilde{C}_1 [(\delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}})^2 + (t/n)], \quad (5.2)$$

$$\sup_{i \in [n]} |\bar{a}_i - w_i| \leq \tilde{C}_1 \cdot (t/n)^{1/2}, \quad (5.3)$$

$$\|\bar{a} - \tilde{a}\|_n \leq \tilde{C}_1 \left[(t/n)^{1/2} + \delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}} \right], \quad (5.4)$$

$$\max \left\{ \sqrt{\frac{n}{t}} \cdot \left| \frac{1}{n} \sum_{i=1}^n (\mu_0 - \bar{g})(X_i) \cdot w_i \right|, \|\bar{g} - \mu_0\|_n \right\} \leq \tilde{C}_1 \left[\delta_\mu^{\text{appr}} + \sqrt{\frac{t}{n}} \right]. \quad (5.5)$$

Let $\mathcal{C}_1$ be the event in Proposition 5.2. The rest of the proof will proceed conditioned on the event $\mathcal{C}_0 \cap \mathcal{C}_1$. On $\mathcal{C}_1$, (5.1) implies $\bar{D} \neq 0$, since $\mathbf{H}_\mu(0) = \infty$. Given that $\tilde{a}$ is uniformly bounded by its construction and Condition 2.1, $\lambda \|\bar{a} - \tilde{a}\|_n^2 + \mathbf{H}_\mu(\bar{a}) < \infty$, thus $\bar{a}$ has finite objective value. Since $\mathbf{H}_\mu(a)$ is lower semi-continuous and $\lambda \|a - \tilde{a}\|_n^2$ is coercive, the objective in (S1- μ) has a finite minimum. Because $\hat{a}$ in (S1- μ) minimizes the objective therein, this means

$$\begin{aligned} \lambda \|\tilde{a} - \hat{a}\|_n^2 + \mathbf{H}_\mu(\hat{a}) &\leq \lambda \|\tilde{a} - \bar{a}\|_n^2 + \mathbf{H}_\mu(\bar{a}) \\ &\stackrel{(a)}{\lesssim} \lambda \left[(\delta_\pi^{\text{appr}})^2 + \frac{t}{n} \right] + (\delta_n^{\text{stoc}})^2. \end{aligned} \quad (5.6)$$

Here (a) follows from the error bound of $\mathbf{H}_\mu(\bar{a})$ (5.1), $\|\tilde{a} - \bar{a}\|_n$ (5.4) in Proposition 5.2, and the condition $\lambda \leq 1$ in the statement. Since $\mathbf{H}_\mu(\hat{a}) \geq \Phi_\mu(0, 0, \hat{a}) = 0$, $\|\hat{a} - \tilde{a}\|_n^2 \lesssim (\delta_\pi^{\text{appr}})^2 + \frac{t}{n} + \frac{(\delta_n^{\text{stoc}})^2}{\lambda}$. Recall the definition of w_i in (2.6), with a slight abuse of notation $\|\hat{a} - w\|_n = \sqrt{\frac{1}{n} \sum_{i=1}^n (\hat{a}_i - w_i)^2}$, combining the triangle inequality with the error bound of $\|\tilde{a} - \bar{a}\|_n$ (5.4) and $\|\bar{a} - w\|_\infty$ (5.3), we find

$$\|\hat{a} - w\|_n \leq \|\hat{a} - \tilde{a}\|_n + \|\tilde{a} - \bar{a}\|_n + \|\bar{a} - w\|_n \lesssim \delta_\pi^{\text{appr}} + \sqrt{\frac{t}{n}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}}. \quad (5.7)$$

On the other hand, combining the non-negativity of $\lambda \cdot \|\tilde{a} - \hat{a}\|_n^2$ and (5.6) also gives

$$\mathbf{H}_\mu(\hat{a}) \lesssim \lambda \left[(\delta_\pi^{\text{appr}})^2 + \frac{t}{n} \right] + (\delta_n^{\text{stoc}})^2. \quad (5.8)$$

Recall the definition of $\mathbf{H}_\mu(\hat{a})$ and $\Phi_\mu(f, \beta, a)$, this implies the following instance-dependent error bound by letting $\beta = 0$ in the supremum:

$$\forall g, \tilde{g} \in \mathcal{G}, \quad \left| \frac{1}{n} \sum_{i=1}^n (g - \tilde{g})(X_i) \cdot \hat{a}_i \right| \leq \|g - \tilde{g}\|_n^2 + \mathbf{H}_\mu(\hat{a}). \quad (5.9)$$

On the other hand, let $f = 0$ in the sup operator, we obtain $|\frac{1}{n} \sum_{i=1}^n T_i \hat{a}_i \cdot \beta - \beta| \leq \mathbf{H}_\mu(\hat{a})$ for any $\beta \in \mathbb{R}$, the finiteness of $\mathbf{H}_\mu(\hat{a})$ implies the normalization constraint $\frac{1}{n} \sum_{i=1}^n \hat{a}_i \cdot T_i = 1$. Recall our choice of fixed $\bar{g} \in \mathcal{G}$

satisfying $\|\bar{g} - \mu_0\|_2 \leq 2\delta_\mu^{\text{appr}}$, it follows from the definition of our estimator in (S2- μ) the model (1.1), and the normalization constraint $\frac{1}{n} \sum_{i=1}^n \hat{a}_i \cdot T_i = 1$ that

$$\hat{\theta}_\mu - \theta_0 \stackrel{(a)}{=} \underbrace{\frac{1}{n} \sum_{i=1}^n (\varepsilon_i + \mu_0(X_i) - \bar{g}(X_i)) \cdot \hat{a}_i}_{\mathsf{T}_1} + \underbrace{\frac{1}{n} \sum_{i=1}^n (\bar{g}(X_i) - \hat{\mu}(X_i)) \cdot \hat{a}_i}_{\mathsf{T}_2}.$$

We will handle T_1 and T_2 separately. For T_2 , we plug in the instance-dependent error bound (5.9) with $g \leftarrow \bar{g}$ and $\tilde{g} \leftarrow \hat{\mu}$ and obtain

$$\begin{aligned} |\mathsf{T}_2| &= \left| \frac{1}{n} \sum_{i=1}^n (\bar{g}(X_i) - \hat{\mu}(X_i)) \cdot \hat{a}_i \right| \lesssim \|\bar{g} - \hat{\mu}\|_n^2 + \left[(\delta_n^{\text{stoc}})^2 + \lambda \cdot \left((\delta_\pi^{\text{appr}})^2 + \frac{t}{n} \right) \right] \\ &\lesssim (\delta_n^{\text{stoc}})^2 + (\delta_\mu^{\text{appr}})^2 + \frac{t}{n} + \lambda \cdot (\delta_\pi^{\text{appr}})^2 \end{aligned}$$

by the error bound of $\|\hat{\mu} - \bar{g}\|_n$ in (5.2), the error bound of $\mathsf{H}_\mu(\hat{a})$ in (5.8), and the fact that $\lambda \leq 1$.

On the other hand, we will further decompose T_1 into

$$\begin{aligned} \mathsf{T}_1 &= \frac{1}{n} \sum_{i=1}^n \varepsilon_i w_i + \frac{1}{n} \sum_{i=1}^n \varepsilon_i (\hat{a}_i - w_i) + \frac{1}{n} \sum_{i=1}^n (\mu_0(X_i) - \bar{g}(X_i)) \cdot (\hat{a}_i - w_i) \\ &\quad + \frac{1}{n} \sum_{i=1}^n (\mu_0(X_i) - \bar{g}(X_i)) \cdot w_i. \end{aligned}$$

We will use the following concentration result: it follows from the fact that $\{\varepsilon_i\}_{i=1}^n$ are zero-mean random variables conditioned on fixed $\mathcal{D}_2$ and $\{(X_i, T_i)\}_{i=1}^n$, under which $\{\hat{a}_i - w_i\}_{i=1}^n$ are fixed because our construction of the weights only depends on $\mathcal{D}_2$ and $\{(X_i, T_i)\}_{i=1}^n$ in the first split $\mathcal{D}_1$.

Lemma 5.3. *Under the setting of Theorem 2.1, assume further that $\|\hat{a} - w\|_n \leq \delta$. There exists some constant $\tilde{C}_1 = \text{poly}(c)$ such that the following event occurs with probability at least $1 - e^{-t}$ for any $t \geq 1$:*

$$\left| \frac{1}{n} \sum_{i=1}^n \varepsilon_i (\hat{a}_i - w_i) \right| \leq \tilde{C}_1 \delta \cdot \sqrt{\frac{t}{n}}$$

Let $\mathcal{C}_2$ be the event in Lemma 5.3. We proceed with the rest of the proof conditioned on the event $\mathcal{C}_0 \cap \mathcal{C}_1 \cap \mathcal{C}_2$. It then follows from the triangle inequality, Cauchy-Schwarz inequality, the error bound of $\|\hat{a} - w\|_n$ in (5.7) and the error bounds in (5.5) and Lemma 5.3 that

$$\begin{aligned} \left| \mathsf{T}_1 - \frac{1}{n} \sum_{i=1}^n \varepsilon_i w_i \right| &\lesssim \|\bar{g} - \mu_0\|_n \cdot \|\hat{a} - w\|_n + \sqrt{\frac{t}{n}} \cdot \left[\delta_\pi^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \delta_\mu^{\text{appr}} + \sqrt{\frac{t}{n}} \right] \\ &\lesssim \left[\sqrt{\frac{t}{n}} + \delta_\mu^{\text{appr}} \right] \cdot \left[\delta_\pi^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}} \right] \end{aligned}$$

Putting all the pieces together, under the intersection of events in Proposition 5.2, Proposition 5.1, and Lemma 5.3 that occurs with probability at least $1 - 2(e^{-n(\delta_n^{\text{stoc}})^2} + e^{-t})$ by the union bound, we have

$$\begin{aligned} \left| \hat{\theta}_\mu - \theta_0 - \hat{\varphi}_{\text{PLM}} \right| &\leq |\mathsf{T}_2| + \left| \mathsf{T}_1 - \frac{1}{n} \sum_{i=1}^n \varepsilon_i w_i \right| \\ &\lesssim \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + (\delta_n^{\text{stoc}})^2 + \lambda (\delta_\pi^{\text{appr}})^2 + \frac{\delta_\mu^{\text{appr}} \cdot \delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}} \left(\delta_\pi^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}} \right) \end{aligned}$$

by noting $\delta_\mu^{\text{appr}} \leq \delta_\pi^{\text{appr}}$ and $\lambda \leq 1$. This concludes the proof for CASE 1. $\square$

5.2 Proof of CASE 2

Denote $m_0(x) = \mathbb{E}[Y|X = x] = \beta_0 \cdot \pi_0(x) + \mu_0(x)$ and $\xi := Y - m_0(X) = \beta_0 u + \varepsilon$. Let $\tilde{m}(x) = \hat{\beta} \cdot \hat{\pi}(x) + \hat{\mu}(x)$. With a slight abuse of notation, we will also denote $\|m - f\|_n = \{\frac{1}{n} \sum_{i=1}^n |m_i - f(X_i)|^2\}^{1/2}$ for $m \in \{\tilde{m}, \hat{m}\}$ and $f \in L_2(\nu_x)$ and use $\tilde{m} \in \mathbb{R}^n$ or $\tilde{m} \in L_2(\nu_x)$ exchangeably based on the context.

We also pick $\bar{g} \in \mathcal{G}$ such that $\|\bar{g} - \pi_0\|_2 \leq 2\delta_\pi^{\text{appr}}$. The proof is based on the following instance-dependent error bounds (5.10) – (5.12) and concentration results (5.13) – (5.17). The concentration results are all standard and follow directly from the Bernstein-type inequality.

Proposition 5.4. Denote $\mathbb{P}_{n,l}[f] = \frac{1}{n} \sum_{i=1+(l-1) \cdot n}^{l \cdot n} f(X_i, T_i, Y_i)$ be the empirical average operator on the split $\mathcal{D}_l$ and $\mathbb{P}[f] = \mathbb{E}[f(X, T, Y)]$, and define $\delta_\pi := \delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}} + (t/n)^{1/2}$ only for the proof of Theorem 2.1. The following event occurs with probability at least $1 - e^{-n(\delta_n^{\text{stoc}})^2} - e^{-t}$ for some $\tilde{C}_1 = \text{poly}(c)$:

$$\forall f \in \partial\mathcal{G}, l \in \{1, 2\} \quad |\mathbb{P}_{n,l}[f(X) \cdot \xi]| \leq \tilde{C}_1 [\|f\|_2 \cdot \delta_n^{\text{stoc}} + (\delta_n^{\text{stoc}})^2] \quad (5.10)$$

$$|(\mathbb{P}_{n,l} - \mathbb{P})[m_0(X)f(X)]| \leq \tilde{C}_1 [\|f\|_2 \cdot \delta_n^{\text{stoc}} + (\delta_n^{\text{stoc}})^2] \quad (5.11)$$

$$\|f\|_2 - \|f\|_n \leq \tilde{C}_1 \delta_n^{\text{stoc}} \quad (5.12)$$

$$\|\tilde{m} - m_0\|_n \leq \tilde{C}_1 [\delta_\pi + \delta_\mu^{\text{appr}}] \quad (5.13)$$

$$\|\bar{g} - \pi_0\|_n \leq \tilde{C}_1 [\delta_\pi^{\text{appr}} + (t/n)^{1/2}] \quad (5.14)$$

$$|\mathbb{P}_{n,1}[(\hat{\pi} - \pi_0)(X) \cdot \varepsilon]| + |\mathbb{P}_{n,1}[(\hat{\pi} - \pi_0)(X) \cdot u]| \leq \tilde{C}_1 \delta_\pi \cdot (t/n)^{1/2} \quad (5.15)$$

$$\|\pi_0 - \hat{\pi}\|_n^2 \leq \tilde{C}_1 \delta_\pi^2 \quad (5.16)$$

$$|\mathbb{P}_{n,1}[(T - \hat{\pi}(X))^2] - \mathbb{E}[u^2]| \leq \tilde{C}_1 [\delta_\pi^2 + (t/n)^{1/2}] \quad (5.17)$$

Let $C_1 = \text{poly}(c)$ be the constant $\tilde{C}_1$ in Proposition 5.4, and the event in Proposition 5.4 be $\mathcal{C}_1$. The rest of the proof proceeds conditioned on the event $\mathcal{C}_0 \cap \mathcal{C}_1$. Denote $H_\pi(m) = \sup_{f \in \partial\mathcal{G}} \Phi_\pi(f, m)$ and $\bar{m} = (m_0(X_1), \dots, m_0(X_n)) \in \mathbb{R}^n$. We first claim that under the inequalities (5.10) – (5.12),

$$H_\pi(\bar{m}) \lesssim (\delta_n^{\text{stoc}})^2, \quad (5.18)$$

whose proof can be found at the end of this section. Given $\hat{m}$ minimizes the objective in (S1- π) and $H_\pi(\hat{m}) \geq \Phi_\pi(0, \hat{m}) = 0$, we obtain $\lambda \cdot \|\hat{m} - \tilde{m}\|_n^2 \leq H_\pi(\bar{m}) + \lambda \cdot \|\bar{m} - \tilde{m}\|_n^2$. Combining it with the triangle inequality further yields

$$\begin{aligned} \|\hat{m} - m_0\|_n &\leq \|\hat{m} - \tilde{m}\|_n + \|\tilde{m} - m_0\|_n \\ &\leq 2\|\tilde{m} - m_0\|_n + \sqrt{\frac{H_\pi(\bar{m})}{\lambda}} \lesssim \delta_\mu^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}}, \end{aligned} \quad (5.19)$$

where the last inequality follows from the error bound (5.13) of $\|m_0 - \tilde{m}\|_n$ and the facts that $\lambda \leq 1$ and $\delta_\pi^{\text{appr}} \leq \delta_\mu^{\text{appr}}$. We further argue the following instance-dependent error bound by the inequalities (5.10) – (5.12), the proof is also deferred to the end of the section.

$$\forall f \in \partial\mathcal{G}, \quad \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \hat{m}_i) \cdot f(X_i) \right| \lesssim \|f\|_2^2 + (\delta_n^{\text{stoc}})^2 + \lambda(\delta_\mu^{\text{appr}})^2 + \frac{t}{n} \quad (5.20)$$

We assume without loss of generality that $2c \cdot C_1 \left\{ (\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}})^2 + \sqrt{\frac{t}{n}} \right\} \leq 1$, otherwise the error rate is trivial because $\hat{\theta}_\pi$ is bounded. This is because the denominator in $\hat{\theta}_\pi$ is bounded from below. At the same time, as $\|\hat{m}\|_n \leq \|\hat{m} - m_0\|_n + \|m_0\|_n \lesssim 1$ provided $\lambda \geq (\delta_n^{\text{stoc}})^2$, the numerator is also bounded by applying the triangle inequality and Cauchy-Schwarz inequality.

Denote $\widehat{D} = \frac{1}{n} \sum_{i=1}^n [T_i - \widehat{\pi}(X_i)]^2$, note $|\widehat{D} - \mathbb{E}[u^2]| \leq C_1 \{(\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}})^2 + (t/n)^{1/2}\}$ by (5.17). As long as $2c \cdot C_1 \{(\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}})^2 + (t/n)^{1/2}\} \leq 1$, $\widehat{D} \geq 1/(2c)$ hence the truncation in (S2- π) is inactive and its denominator equals $\widehat{D}$. By our constructed estimator (S2- π) and the model (1.1), we decompose

$$\frac{1}{n} \sum_{i=1}^n [T_i - \widehat{\pi}(X_i)]^2 [\widehat{\theta}_\pi - \theta_0] = \frac{1}{n} \sum_{i=1}^n \{\varepsilon_i + m_0(X_i) - \widehat{m}_i - (\pi_0(X_i) - \widehat{\pi}(X_i))\beta_0\} \{u_i + \pi_0(X_i) - \widehat{\pi}(X_i)\}$$

Recall our choice of fixed $\bar{g} \in \mathcal{G}$ such that $\|\bar{g} - \pi_0\|_2 \leq 2\delta_\pi^{\text{appr}}$, it further follows from the triangle inequality that

$$\begin{aligned} |\widehat{\theta}_\pi - \theta_0 - \widehat{\varphi}_{\text{PLM}}| &\leq \left| \frac{1}{\mathbb{E}[u^2]} - \frac{1}{\widehat{D}} \right| \cdot \left| \frac{1}{n} \sum_{i=1}^n \varepsilon_i u_i \right| + \frac{1}{|\widehat{D}|} \cdot \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i) \cdot u_i \right| \\ &\quad + \frac{|\beta_0|}{|\widehat{D}|} \frac{1}{n} \sum_{i=1}^n |\widehat{\pi}(X_i) - \pi_0(X_i)|^2 + \frac{|\beta_0|}{|\widehat{D}|} \cdot \left| \frac{1}{n} \sum_{i=1}^n (\pi_0(X_i) - \widehat{\pi}(X_i)) \cdot u_i \right| \\ &\quad + \frac{1}{|\widehat{D}|} \cdot \left| \frac{1}{n} \sum_{i=1}^n (\pi_0 - \widehat{\pi})(X_i) \varepsilon_i \right| + \frac{1}{|\widehat{D}|} \cdot \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i)(\pi_0 - \widehat{\pi})(X_i) \right| \end{aligned}$$

It then follows from the error bounds (5.15), (5.16), (5.17) and the fact that $|\widehat{D}|^{-1} + |\beta_0| \lesssim 1$ by the above discussion and Condition 2.1 that

$$\begin{aligned} |\widehat{\theta}_\pi - \theta_0 - \widehat{\varphi}_{\text{PLM}}| &\lesssim \left[(\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}})^2 + \sqrt{\frac{t}{n}} \right] \cdot \left| \frac{1}{n} \sum_{i=1}^n \varepsilon_i u_i \right| + \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i) \cdot u_i \right| \\ &\quad + (\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}})^2 + \frac{t}{n} + \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i)(\pi_0 - \bar{g})(X_i) \right| \\ &\quad + \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i)(\bar{g}(X_i) - \widehat{\pi}(X_i)) \right|. \end{aligned}$$

We will further use the following concentration result, which is due to our construction of the cross-sample debiasing weights $\{\widehat{m}_i\}_{i=1}^n$ that depend on $\mathcal{D}_2$ and only $\{X_i\}_{i=1}^n$ in the first sample $\mathcal{D}_1$. This will make $\{u_i \cdot (m_0(X_i) - \widehat{m}_i)\}_{i=1}^n$ be independent zero-mean random variables given fixed $\mathcal{D}_2, \{X_i\}_{i=1}^n$.

Lemma 5.5. *Under the setting of Theorem 2.1, assume further that $\|\widehat{m} - m_0\|_n \leq \delta$, then with probability at least $1 - e^{-t}$,*

$$\left| \frac{1}{n} \sum_{i=1}^n \varepsilon_i \cdot u_i \right| \leq \widetilde{C}_1 \sqrt{\frac{t}{n}} \quad \text{and} \quad \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i) \cdot u_i \right| \leq \widetilde{C}_1 \cdot \delta \sqrt{\frac{t}{n}}.$$

Let $\mathcal{C}_2$ be the event in Lemma 5.5. We proceed with the rest of the proof conditioned on the event $\mathcal{C}_0 \cap \mathcal{C}_1 \cap \mathcal{C}_2$, which occurs with probability at least $1 - 2(e^{-n(\delta_n^{\text{stoc}})^2} + e^{-t})$ by the union bound. Substituting the error bound in Lemma 5.5 with $\delta \leftarrow \delta_\mu^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}}$ by (5.19), we arrive at

$$|\widehat{\theta}_\pi - \theta_0 - \widehat{\varphi}_{\text{PLM}}| \lesssim (\delta_\pi^{\text{appr}} + \delta_n^{\text{stoc}})^2 + \frac{t}{n} + \sqrt{\frac{t}{n}} \cdot \left(\delta_\mu^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \delta_n^{\text{stoc}} \right) + \mathsf{T}_1 + \mathsf{T}_2. \quad (5.21)$$

where $\mathsf{T}_1 = \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i)(\pi_0 - \bar{g})(X_i) \right|$ and $\mathsf{T}_2 = \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i)(\bar{g}(X_i) - \widehat{\pi}(X_i)) \right|$. On the one hand, it follows from the Cauchy-Schwarz inequality and the error bound for $\|\widehat{m} - m_0\|_n$ in (5.19) and the error bound for $\|\bar{g} - \pi_0\|_n$ in (5.14) that

$$\mathsf{T}_1 \leq \|\widehat{m} - m_0\|_n \cdot \|\bar{g} - \pi_0\|_n \lesssim \left[\delta_\pi^{\text{appr}} + \sqrt{\frac{t}{n}} \right] \cdot \left[\delta_\mu^{\text{appr}} + \frac{\delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \sqrt{\frac{t}{n}} \right].$$

On the other hand, applying the instance-dependent error bound (5.20) with $f = \bar{g} - \hat{\pi} \in \partial\mathcal{G}$ and the fact that $\|\hat{\pi} - \bar{g}\|_2 \lesssim \|\hat{\pi} - \pi_0\|_2 + \|\pi_0 - \bar{g}\|_2 \lesssim \delta_n^{\text{stoc}} + \delta_\pi^{\text{appr}}$ under the event in Proposition 5.1 yield

$$\mathsf{T}_2 \lesssim \|\bar{g} - \hat{\pi}\|_2^2 + (\delta_n^{\text{stoc}})^2 + \lambda(\delta_\mu^{\text{appr}})^2 + t/n \lesssim (\delta_\pi^{\text{appr}})^2 + (\delta_n^{\text{stoc}})^2 + \lambda(\delta_\mu^{\text{appr}})^2 + t/n.$$

Now putting the two inequalities back into (5.21) and keeping in mind that we are considering the case where $\delta_\pi^{\text{appr}} \leq \delta_\mu^{\text{appr}}$ and $\lambda \leq 1$, we can conclude that, with the high-order residual term $\text{Res} = (t/n)^{1/2} \cdot \{\delta_\mu^{\text{appr}} + \delta_n^{\text{stoc}}/\sqrt{\lambda} + (t/n)^{1/2}\}$,

$$\left| \hat{\theta}_\pi - \theta_0 - \hat{\varphi}_{\text{PLM}} \right| \lesssim \delta_\mu^{\text{appr}} \cdot \delta_\pi^{\text{appr}} + (\delta_n^{\text{stoc}})^2 + \lambda(\delta_\mu^{\text{appr}})^2 + \frac{\delta_\pi^{\text{appr}} \cdot \delta_n^{\text{stoc}}}{\sqrt{\lambda}} + \text{Res}.$$

This concludes the proof for CASE 2. $\square$

Proof of (5.18). To this end, it follows from the triangle inequality that, for any $f = g - \tilde{g} \in \partial\mathcal{G}$,

$$\begin{aligned} \Phi_\pi(f, \bar{m}) &= \left| \frac{1}{n} \sum_{i=n+1}^{2n} Y_i f(X_i) - \frac{1}{n} \sum_{i=1}^n m_0(X_i) f(X_i) \right| - \|f\|_n^2 \\ &\leq \left| \frac{1}{n} \sum_{i=n+1}^{2n} \xi_i f(X_i) \right| + \left| \frac{1}{n} \sum_{i=n+1}^{2n} m_0(X_i) f(X_i) - \mathbb{E}[m_0(X) f(X)] \right| \\ &\quad + \left| \mathbb{E}[m_0(X) f(X)] - \frac{1}{n} \sum_{i=1}^n m_0(X_i) f(X_i) \right| - \|f\|_n^2 \\ &\stackrel{(a)}{\leq} 3C_1 \{ \|f\|_2 \cdot \delta_n^{\text{stoc}} + (\delta_n^{\text{stoc}})^2 \} - \|f\|_n^2 \\ &\stackrel{(b)}{\leq} 3C_1 \{ (\|f\|_n + C_1 \delta_n^{\text{stoc}}) \cdot \delta_n^{\text{stoc}} + (\delta_n^{\text{stoc}})^2 \} - \|f\|_n^2 \\ &\stackrel{(c)}{\leq} \|f\|_n^2 + \frac{(3C_1)^2}{4} (\delta_n^{\text{stoc}})^2 + 3C_1(C_1 + 1) \cdot (\delta_n^{\text{stoc}})^2 - \|f\|_n^2 \lesssim (\delta_n^{\text{stoc}})^2 \end{aligned}$$

where we apply (5.10) and (5.11) in (a), (5.12) in (b), and use a variant of Young's inequality $xy \leq x^2 + \frac{y^2}{4}$ with $x = \|f\|_n$ and $y = (3C_1)\delta_n^{\text{stoc}}$ in (c). Taking the supremum on both sides concludes the proof. $\square$

Proof of the inequality (5.20). By the optimality of $\hat{m}$, and after dropping the non-negative term $\lambda\|\hat{m} - \tilde{m}\|_n^2$, we find

$$\mathsf{H}_\pi(\hat{m}) \leq \mathsf{H}_\pi(\bar{m}) + \lambda \cdot \|m_0 - \tilde{m}\|_n^2 \lesssim (\delta_n^{\text{stoc}})^2 + \lambda \left[(\delta_\mu^{\text{appr}})^2 + \frac{t}{n} \right] \quad (5.22)$$

where again we plug in the upper bound (5.13) for $\|\tilde{m} - m_0\|_n$ and use the facts $\delta_\pi^{\text{appr}} \leq \delta_\mu^{\text{appr}}$, $\lambda \leq 1$ to simplify the presentation. It follows from the triangle inequality that, for any $f = g - \tilde{g} \in \partial\mathcal{G}$,

$$\begin{aligned} \left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \hat{m}_i) f(X_i) \right| &\leq \left| \frac{1}{n} \sum_{i=n+1}^{2n} Y_i f(X_i) - \frac{1}{n} \sum_{i=1}^n \hat{m}_i f(X_i) \right| \\ &\quad + \left| \frac{1}{n} \sum_{i=n+1}^{2n} Y_i f(X_i) - \frac{1}{n} \sum_{i=1}^n m_0(X_i) f(X_i) \right| \\ &\stackrel{(a)}{=} \Phi_\pi(f, \hat{m}) + \|f\|_n^2 \\ &\quad + \left| \frac{1}{n} \sum_{i=n+1}^{2n} (m_0(X_i) + \xi_i) \cdot f(X_i) - \frac{1}{n} \sum_{i=1}^n m_0(X_i) f(X_i) \right| \end{aligned}$$

$$\begin{aligned}
&\stackrel{(b)}{\leq} \mathbf{H}_\pi(\widehat{m}) + \|f\|_n^2 + \left| \frac{1}{n} \sum_{i=n+1}^{2n} \xi_i f(X_i) \right| \\
&\quad + \sum_{l \in \{1,2\}} \left| \frac{1}{n} \sum_{i=1+(l-1)n}^{ln} m_0(X_i) \cdot f(X_i) - \mathbb{E}[m_0(X)f(X)] \right|
\end{aligned}$$

Here (a) follows from the definition of Φ_π , (b) follows from the definition $\mathbf{H}_\pi(m) = \sup_{f \in \partial \mathcal{G}} \Phi_\pi(f, m)$ and the triangle inequality. Now we plug in the error bound (5.22) for $\mathbf{H}_\pi(\widehat{m})$ as well as the instance-dependent error bounds (5.10) and (5.11),

$$\begin{aligned}
\left| \frac{1}{n} \sum_{i=1}^n (m_0(X_i) - \widehat{m}_i) f(X_i) \right| &\lesssim \|f\|_n^2 + (\delta_n^{\text{stoc}})^2 + \lambda \left[(\delta_\mu^{\text{appr}})^2 + \frac{t}{n} \right] + \|f\|_2 \cdot \delta_n^{\text{stoc}} + (\delta_n^{\text{stoc}})^2 \\
&\stackrel{(a)}{\lesssim} \|f\|_2^2 + (\delta_n^{\text{stoc}})^2 + \lambda \left[(\delta_\mu^{\text{appr}})^2 + \frac{t}{n} \right],
\end{aligned}$$

where (a) follows from the equivalence between L_2 norm and its empirical counterparts in (5.12) and Young's inequality. This concludes the proof for (5.20). $\square$

References

- Ai, C. & Chen, X. (2003). Efficient estimation of models with conditional moment restrictions containing unknown functions. *Econometrica*, 71(6), 1795–1843.
- Athey, S., Imbens, G. W., & Wager, S. (2018). Approximate residual balancing: debiased inference of average treatment effects in high dimensions. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80(4), 597–623.
- Balakrishnan, S., Kennedy, E. H., & Wasserman, L. (2023). The fundamental limits of structure-agnostic functional estimation. *arXiv preprint arXiv:2305.04116*.
- Bartlett, P., Bousquet, O., & Mendelson, S. (2005). Local rademacher complexities. *Annals of Statistics*, 33(4), 1497–1537.
- Bauer, B. & Kohler, M. (2019). On deep learning as a remedy for the curse of dimensionality in nonparametric regression. *The Annals of Statistics*, 47(4), 2261–2285.
- Bickel, P. J., Ritov, Y., & Tsybakov, A. B. (2009). Simultaneous analysis of Lasso and Dantzig selector. *The Annals of Statistics*, 37(4), 1705 – 1732.
- Bonvini, M., Kennedy, E. H., Dukes, O., & Balakrishnan, S. (2024). Doubly-robust inference and optimality in structure-agnostic models with smoothness. *arXiv preprint arXiv:2405.08525*.
- Bradic, J., Chernozhukov, V., Newey, W. K., & Zhu, Y. (2019). Minimax semiparametric learning with approximate sparsity. *arXiv preprint arXiv:1912.12213*.
- Bradic, J., Fan, J., & Zhu, Y. (2022). Testability of high-dimensional linear models with nonsparse structures. *Annals of statistics*, 50(2), 615.
- Bruns-Smith, D., Dukes, O., Feller, A., & Ogburn, E. L. (2025). Augmented balancing weights as linear regression. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, (pp. qkaf019).

- Cai, T. T. & Guo, Z. (2017). Confidence intervals for high-dimensional linear regression: Minimax rates and adaptivity. *The Annals of Statistics*, 45(2), 615 – 646.
- Candes, E. & Tao, T. (2007). The dantzig selector: Statistical estimation when p is much larger than n . *The Annals of Statistics*, (pp. 2313–2351).
- Carroll, R. J., Fan, J., Gijbels, I., & Wand, M. P. (1997). Generalized partially linear single-index models. *Journal of the American Statistical Association*, 92(438), 477–489.
- Celentano, M. & Wainwright, M. J. (2023). Challenges of the inconsistency regime: Novel debiasing methods for missing data models. *arXiv preprint arXiv:2309.01362*.
- Chen, X. (2007). Large sample sieve estimation of semi-nonparametric models. *Handbook of econometrics*, 6, 5549–5632.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., & Newey, W. (2017). Double/debiased/neyman machine learning of treatment effects. *American Economic Review*, 107(5), 261–265.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., & Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1), C1–C68.
- Chernozhukov, V., Newey, W. K., & Singh, R. (2022a). Automatic debiased machine learning of causal and structural effects. *Econometrica*, 90(3), 967–1027.
- Chernozhukov, V., Newey, W. K., & Singh, R. (2022b). Debiased machine learning of global and local parameters using regularized riesz representers. *The Econometrics Journal*, 25(3), 576–601.
- Chernozhukov, V., Newey, W. K., Singh, R., & Syrgkanis, V. (2026). Adversarial estimation of riesz representers. *Journal of the American Statistical Association*, (pp. 1–12).
- Donald, S. G. & Newey, W. K. (1994). Series estimation of semilinear models. *Journal of Multivariate Analysis*, 50(1), 30–40.
- Engle, R. F., Granger, C. W., Rice, J., & Weiss, A. (1986). Semiparametric estimates of the relation between weather and electricity sales. *Journal of the American statistical Association*, 81(394), 310–320.
- Fan, J. & Gu, Y. (2024). Factor augmented sparse throughput deep relu neural networks for high dimensional regression. *Journal of the American Statistical Association*, 119(548), 2680–2694.
- Fan, J., Gu, Y., & Zhou, W.-X. (2024). How do noise tails impact on deep ReLU networks? *The Annals of Statistics*, 52(4), 1845 – 1871.
- Fan, J. & Huang, T. (2005). Profile likelihood inferences on semiparametric varying-coefficient partially linear models. *Bernoulli*, 11(6), 1031–1057.
- Fan, J., Imai, K., Lee, I., Liu, H., Ning, Y., & Yang, X. (2022). Optimal covariate balancing conditions in propensity score estimation. *Journal of Business & Economic Statistics*, 41(1), 97–110.
- Foster, D. J. & Syrgkanis, V. (2023). Orthogonal statistical learning. *The Annals of Statistics*, 51(3), 879–908.
- Gu, Y., Fang, C., Bühlmann, P., & Fan, J. (2025). Causality pursuit from heterogeneous environments via neural adversarial invariance learning. *The Annals of Statistics*, 53(5), 2230 – 2257.

- Gu, Y., Yin, Q., Cai, T., & Fan, J. (2026). Optimally taming biases in black-box models for efficient semiparametric estimation. *arXiv preprint arXiv:2606.06368*.
- Hellerstein, J. K. & Imbens, G. W. (1999). Imposing moment restrictions from auxiliary data by weighting. *Review of Economics and Statistics*, 81(1), 1–14.
- Hirshberg, D. A. & Wager, S. (2021). Augmented minimax linear estimation. *The Annals of Statistics*, 49(6), 3206–3227.
- Imai, K. & Ratkovic, M. (2014). Covariate balancing propensity score. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 76(1), 243–263.
- Javanmard, A. & Montanari, A. (2014). Confidence intervals and hypothesis testing for high-dimensional regression. *Journal of Machine Learning Research*, 15(82), 2869–2909.
- Javanmard, A. & Montanari, A. (2018). Debiasing the lasso: Optimal sample size for Gaussian designs. *The Annals of Statistics*, 46(6A), 2593 – 2622.
- Jin, J., Mackey, L., & Syrgkanis, V. (2025). It’s hard to be normal: The impact of noise on structure-agnostic estimation. *arXiv preprint arXiv:2507.02275*.
- Kohler, M. & Langer, S. (2021). On the rate of convergence of fully connected deep neural network regression estimates. *The Annals of Statistics*, 49(4), 2231–2249.
- Koltchinskii, V. (2006). Local Rademacher complexities and oracle inequalities in risk minimization. *The Annals of Statistics*, 34(6), 2593 – 2656.
- Kong, J. (2025). On the asymptotics of the minimax linear estimator. *arXiv preprint arXiv:2510.16661*.
- Liu, L., Mukherjee, R., Newey, W. K., & Robins, J. M. (2017). Semiparametric efficient empirical higher order influence function estimators. *arXiv preprint arXiv:1705.07577*.
- Massart, P. & Nédélec, É. (2006). Risk bounds for statistical learning. *Ann. Statist.*, 34(1), 2326–2366.
- Ning, Y., Sida, P., & Imai, K. (2020). Robust estimation of causal effects via a high-dimensional covariate balancing propensity score. *Biometrika*, 107(3), 533–554.
- Rice, J. (1986). Convergence rates for partially splined models. *Statistics & probability letters*, 4(4), 203–208.
- Robins, J., Li, L., Tchetgen, E., van der Vaart, A., et al. (2008). Higher order influence functions and minimax estimation of nonlinear functionals. In *Probability and statistics: essays in honor of David A. Freedman*, volume 2 (pp. 335–422). Institute of Mathematical Statistics.
- Robins, J., Li, L., Tchetgen, E. T., & van der Vaart, A. (2016). Technical report: Higher order influence functions and minimax estimation of nonlinear functionals. *arXiv preprint arXiv:1601.05820*.
- Robins, J. M., Rotnitzky, A., & Zhao, L. P. (1994). Estimation of regression coefficients when some regressors are not always observed. *Journal of the American statistical Association*, 89(427), 846–866.
- Schmidt-Hieber, J. (2020). Nonparametric regression using deep neural networks with relu activation function (with discussion). *The Annals of Statistics*, 48(4), 1875–1921.
- Tan, Z. (2020). Model-assisted inference for treatment effects using regularized calibrated estimation with high-dimensional data. *Annals of Statistics*, 48(2), 811–837.

- Van de Geer, S., Bühlmann, P., Ritov, Y., & Dezeure, R. (2014). On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics*, 42(3), 1166–1202.
- van der Laan, L., Luedtke, A., & Carone, M. (2024). Doubly robust inference via calibration. *arXiv preprint arXiv:2411.02771*.
- Van der Laan, M. J. (2014). Targeted estimation of nuisance parameters to obtain valid statistical inference. *The international journal of biostatistics*, 10(1), 29–57.
- Vapnik, V. & Chervonenkis, A. Y. (1971). On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability & Its Applications*, 16(2), 264–280.
- Wainwright, M. J. (2019). *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press.
- Wang, Y. & Shah, R. D. (2024). Debiased inverse propensity score weighting for estimation of average treatment effects with high-dimensional confounders. *The Annals of Statistics*, 52(5), 1978 – 2003.
- Zhang, C.-H. & Zhang, S. S. (2014). Confidence intervals for low dimensional parameters in high dimensional linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(1), 217–242.
- Zhao, Q. (2019). Covariate balancing propensity score by tailored loss functions. *The Annals of Statistics*, 47(2), 965 – 993.
- Zubizarreta, J. R. (2015). Stable weights that balance covariates for estimation with incomplete outcome data. *Journal of the American Statistical Association*, 110(511), 910–922.