

The Convergence Rate of Stochastic Tracking with Application to Optimal Execution

Marcel Nutz* Moritz Voss†

September 1, 2026

Abstract

We study the quadratic tracking problem of a general stochastic target process with absolutely continuous controls, with and without terminal constraint. We derive explicit, non-asymptotic upper bounds in terms of a Besov-type modulus of the target. These bounds yield sharp explicit rates that specialize to the square-root order for semimartingale targets. We then apply these results to a generalized Obizhaeva–Wang execution model with random terminal inventory. We first develop a Hilbert-space approach to characterize its optimal strategy, which includes jumps. To avoid such trading spikes, one regularizes the problem by a quadratic trading-rate penalty with coefficient ε . We then show that the regularized optimal execution cost—and therefore the excess price impact cost of the regularized optimal strategy—converges at the sharp rate $O(\sqrt{\varepsilon})$. Since the regularized optimal strategy is not available in closed form, we further construct a nearly optimal strategy which is readily implementable and shares the same approximation rate.

Mathematics Subject Classification (2020): 93E20, 91G80, 60H30.

JEL Classification: G11, C61.

Keywords: Stochastic optimal tracking, convergence rate, optimal execution, transient price impact.

Contents

1	Introduction	2
2	Unconstrained optimal tracking	5
3	Optimal tracking with terminal state constraint	16
4	An Obizhaeva–Wang model and its solution	24
5	Regularized execution and its approximation rate	32

*Departments of Statistics and Mathematics, Columbia University, mnutz@columbia.edu. Research supported by NSF Grants DMS-2106056, DMS-2407074, DMS-2606731.

†Department of Mathematics, University of California Los Angeles, voss@math.ucla.edu.

1 Introduction

We consider stochastic optimal tracking problems of the form

$$\min_u \mathbb{E} \left[\frac{1}{2} \int_0^T (X_t^u - \xi_t)^2 dt + \frac{\kappa}{2} \int_0^T u_t^2 dt \right], \quad (1.1)$$

in which a controller chooses the velocity $u = (u_t)_{0 \leq t \leq T}$ of an absolutely continuous state process

$$X_t^u = x + \int_0^t u_s ds, \quad 0 \leq t \leq T,$$

so as to remain close to a prescribed progressively measurable target process $\xi = (\xi_t)_{0 \leq t \leq T}$, while penalizing rapid adjustments through the quadratic cost parameter $\kappa > 0$. In the constrained version, the controller is additionally required to satisfy

$$X_T^u = \Xi_T \quad \mathbb{P}\text{-a.s.}$$

for a given random target position Ξ_T at time T . Our main interest is the small-friction regime $\kappa \downarrow 0$, in which we investigate how fast the optimal tracking cost vanishes as adjustment costs disappear. In our optimal execution application, the running target ξ will be the inventory of the optimal execution strategy in a generalized Obizhaeva–Wang model. The constrained tracking problem will quantify the cost of replacing this strategy, which generally has infinite variation and includes block trades, by a smoother strategy while preserving the random terminal position Ξ_T .

For fixed $\kappa > 0$, semi-explicit solutions for both the unconstrained and constrained versions of (1.1) were derived by Bank, Soner, and Voß [7] in terms of certain kernels. These formulas provide useful insights into the stochastic tracking problem. It is, however, not straightforward to derive general properties, such as asymptotic bounds that can be read from regularity properties of the target (or at least we have not succeeded in doing so).

Our first main contribution is to derive constructive, non-asymptotic upper bounds for the value function (1.1). For the unconstrained problem, the bound is expressed in terms of an exponentially weighted average of the squared L^2 time-translation modulus of the target and immediately yields a generic $O(\sqrt{\kappa})$ rate whenever the associated seminorm in (2.7) is finite. We show that this includes all square-integrable càdlàg semimartingales, as well as certain non-semimartingale targets such as fractional Brownian motion with Hurst parameter $H > 1/2$. In general, a lower bound shows that whenever the modulus behaves approximately like a power law, it fully determines the convergence rate up to multiplicative constants. We show that the generic square-root rate is sharp for continuous martingales and for certain deterministic targets with jumps. By contrast, for fractional Brownian motion with $H < 1/2$, the time-translation seminorm is infinite and (1.1) vanishes at the slower rate κ^H .

For the constrained problem with prescribed terminal state Ξ_T , the upper bound is additionally governed by the short-horizon mismatch between the running target ξ and the terminal-target martingale

$$\Xi_t := \mathbb{E}[\Xi_T \mid \mathcal{F}_t], \quad 0 \leq t \leq T,$$

as well as by the local cost of learning Ξ_T near maturity. The latter is related to the reachability criterion identified in [7] as the necessary and sufficient admissibility condition for

a random terminal position. Again, the typical rate is $O(\sqrt{\kappa})$ in the most relevant cases, and sharpness is established in benchmark examples. Importantly, all our upper bounds are constructive. In the unconstrained problem, the bound follows from a suboptimal state X^u that exponentially filters the target at relaxation scale $\sqrt{\kappa}$. In the constrained problem, this tracker is concatenated with a terminal bridge that enforces the random terminal position. This construction will also be used to produce an explicit, implementable execution strategy in the second part of the paper.

Indeed, our second main contribution concerns optimal execution with a stochastic terminal target. The model is motivated by the optimal execution problem of a central risk book (CRB) as in Nutz, Webster, and Zhao [19]. A CRB receives random order flow while unwinding its inventory, so that the total quantity to be executed is learned over time and the terminal constraint is inherently stochastic. If the order flow is truth-telling (cf. [19]), the time- t inventory is precisely Ξ_t . We study a generalized Obizhaeva–Wang model [20] of transient price impact with deterministic time-varying resilience and depth, random terminal position Ξ_T , and càdlàg semimartingale execution strategies. The central financial question is the cost of suppressing the block trades and infinite variation of the optimal singular strategy by imposing absolute continuity and a quadratic penalty on the trading rate. This question was first posed by Ulrich Horst in connection with the regularized model that was used in [19].

We first solve the unregularized execution problem explicitly by a direct Hilbert-space argument. This yields the unique optimal càdlàg semimartingale strategy without relying on the backward stochastic differential equation (BSDE) techniques, and the corresponding mathematical restrictions, employed in previous works (see also below). The optimal inventory is driven by the terminal-target martingale $(\Xi_t)_{0 \leq t \leq T}$ and therefore typically has infinite variation, in addition to block trades. To produce an absolutely continuous optimal execution strategy, the problem is often regularized by adding an instantaneous quadratic penalty on the trading rate (as in, for example, Graewe and Horst [17]), forcing execution strategies to avoid spikes in trading activity that would leak too much information to the market.

Our Hilbert-space formulation provides the connection between the execution and tracking parts of the paper. Writing $J_0(Q)$ for the execution cost of an inventory strategy Q in the (unregularized) price impact model and Q^0 for the optimal inventory, the excess cost of Q is precisely the squared distance between the corresponding impact processes Y^Q and Y^{Q^0} in the Hilbert space $\mathcal{H}$ that we introduce:

$$J_0(Q) - J_0(Q^0) = \|Y^Q - Y^{Q^0}\|_{\mathcal{H}}^2.$$

By the impact dynamics of the Obizhaeva–Wang model, the right-hand side is further bounded by a constant times $\mathbb{E} \int_0^T |Q_t - Q_t^0|^2 dt$. As a result, the approximation of the regularized execution problem reduces to constrained tracking of the unregularized optimizer Q^0 by an absolutely continuous strategy Q with terminal position $Q_T = Q_T^0$.

We verify that Q^0 has a finite squared L^2 time-translation seminorm and satisfies a uniform L^2 bound relative to the terminal-target martingale. These estimates allow us to specialize the constructive tracking mechanism developed in the first part of the paper. The resulting strategy Q^ε exponentially tracks Q^0 with relaxation scale $\sqrt{\varepsilon}$ until the deterministic time $T - \sqrt{\varepsilon}$, and then switches to a terminal bridge enforcing $Q_T^\varepsilon = \Xi_T$. The feedback form (5.20) of Q^ε is simple, explicit, and implementable; it does not require solving the less tractable regularized optimization problem.

We derive non-asymptotic bounds showing that the excess price impact cost $J_0(Q^\varepsilon) - J_0(Q^0)$ is of order $O(\sqrt{\varepsilon})$. Thus, Q^ε is an implementable, nearly optimal alternative to the generally unknown regularized optimizer. Indeed, it approximates $J_0(Q^0)$ at the same rate as the true optimizer: we also show that the difference between the regularized cost of Q^ε and the unregularized value function is of order $O(\sqrt{\varepsilon})$, which is a stronger result and implies that the difference between the value functions of the regularized and unregularized problems is of order $O(\sqrt{\varepsilon})$. Finally, we show that this rate is sharp in the benchmark case of constant model parameters and deterministic terminal position.

Our paper sits at the intersection of two strands of the literature. The first concerns optimal tracking of a general target process with absolutely continuous controls and a random terminal state constraint, as introduced by Bank, Soner, and Voß [7], whose setup we adopt. Bank and Voß [8] study a more general framework with stochastic coefficients. Related examples of targeting problems and terminal-state constraints appear in Ankirchner and Kruse [5], Graewe and Horst [17], and Dolinsky, Gottesman, and Gurel-Gurevich [13], among others. These works, however, do not study the optimal tracking of a general stochastic target process. Compared with [7], our contribution differs in emphasis: rather than seeking exact solution formulas, we derive non-asymptotic bounds tied directly to the roughness of the target and, in the constrained problem, to the information flow associated with the random terminal position. Cai, Rosenbaum, and Tankov derive asymptotic lower bounds for a class of tracking objectives and control types in [9] and construct asymptotically optimal feedback strategies for many examples in the companion paper [10]. Their framework is restricted to continuous Itô-semimartingale targets and does not include random terminal state constraints, and therefore does not apply directly to our problem.

The second strand concerns transient price impact and optimal execution. Variants of the Obizhaeva–Wang model [20] with deterministic time-varying illiquidity parameters were analyzed, among others, by Bank and Fruth [6], Alfonsi and Acevedo [4], and Fruth, Schöneborn, and Urusov [15]. Extensions with stochastic liquidity include Fruth, Schöneborn, and Urusov [16] and Ackermann, Kruse, and Urusov [2, 1, 3]. In contrast to these works, we derive the optimal strategy for our model through a direct Hilbert-space argument rather than through BSDEs. See also Webster [21] and Cartea, Jaimungal, and Penalva [11] for extensive references.

The regularized problem with quadratic instantaneous costs is related to the optimal liquidation problems studied by Graewe and Horst [17], Nutz, Webster, and Zhao [19], and Chen, Horst, and Tran [12]. The latter obtain an explicit solution in the benchmark case of constant coefficients and a deterministic terminal position. Following [19], we cover deterministic time-varying coefficients and a general stochastic terminal target. The work of Horst and Kivman [18] is comparable to our asymptotic analysis; they study the convergence of the optimal execution strategy in the optimal liquidation framework of [17] when the instantaneous cost parameter vanishes. The convergence result is qualitative, it holds in probability and uniformly in time over time intervals bounded away from the initial and terminal times. At $t = 0$ and $t = T$, this convergence fails as the limiting strategy has jumps. They show that the limiting càdlàg semimartingale strategy solves an optimal liquidation problem of Obizhaeva–Wang-type as in [1] with a deterministic liquidation constraint, but the speed of convergence is not considered. Our focus is on random terminal targets and quantitative results, both for the optimizer and the directly implementable strategy that we propose.

In a different direction, related asymptotic questions are also considered by Dolinsky and Dolinsky [14], who study exponential-utility indifference prices of European options under a liquidation constraint in a regime of small linear price impact and high risk aversion.

The remainder of the paper is organized as follows. Section 2 studies the asymptotics of the unconstrained tracking problem, and Section 3 treats the version with a random terminal state constraint. In Section 4, we introduce and explicitly solve the unregularized Obizhaeva–Wang execution problem. Finally, Section 5 combines the Hilbert-space structure of the execution problem with the tracking mechanisms from Sections 2 and 3 to derive an explicit absolutely continuous execution strategy with non-asymptotic cost guarantees, as well as the sharp square-root convergence rate for the regularized value.

2 Unconstrained optimal tracking

Fix $T > 0$, a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$ satisfying the usual conditions, and an initial position $x \in \mathbb{R}$. For any process u in the class of admissible controls

$$\mathcal{U} = \left\{ u = (u_t)_{0 \leq t \leq T} : u \text{ is progressively measurable and } \mathbb{E} \int_0^T u_t^2 dt < \infty \right\}, \quad (2.1)$$

we define the associated state process

$$X_t^u = x + \int_0^t u_s ds, \quad 0 \leq t \leq T. \quad (2.2)$$

Throughout, we fix a progressively measurable target process $\xi = (\xi_t)_{0 \leq t \leq T}$ satisfying

$$\xi_0 \in L^2(\mathbb{P}), \quad \mathbb{E} \int_0^T \xi_t^2 dt < \infty, \quad (2.3)$$

and consider the quadratic tracking objective

$$J_\kappa(u) = \mathbb{E} \left[\frac{1}{2} \int_0^T (X_t^u - \xi_t)^2 dt + \frac{\kappa}{2} \int_0^T u_t^2 dt \right], \quad \kappa \geq 0. \quad (2.4)$$

The associated value is

$$v(\kappa) := \inf_{u \in \mathcal{U}} J_\kappa(u). \quad (2.5)$$

Our aim is to quantify the behavior of $v(\kappa)$ as $\kappa \downarrow 0$, which clearly depends on the regularity of the target ξ . Our key insight is that the relevant roughness can be quantified by the following (squared) L^2 time-translation modulus of ξ :

$$\omega_\xi(h) := \mathbb{E} \int_0^T |\xi_t - \xi_{(t-h)^+}|^2 dt, \quad h > 0, \quad (2.6)$$

where $(t-h)^+ := \max\{t-h, 0\}$. By definition,

$$\begin{aligned} \omega_\xi(h) &= \mathbb{E} \int_0^h |\xi_t - \xi_0|^2 dt + \mathbb{E} \int_h^T |\xi_t - \xi_{t-h}|^2 dt, \quad 0 < h \leq T, \\ \omega_\xi(h) &= \omega_\xi(T) = \mathbb{E} \int_0^T |\xi_t - \xi_0|^2 dt, \quad h \geq T, \end{aligned}$$

and $\lim_{h \downarrow 0} \omega_\xi(h) = 0$. In particular, note that $\omega_\xi(h) < \infty$ for all $h > 0$, for any target process ξ satisfying (2.3). Moreover, we introduce

$$[\xi]_{\text{tr}}^2 := \sup_{h>0} \frac{\omega_\xi(h)}{h} \quad (2.7)$$

and refer to it as the squared L^2 time-translation seminorm of ξ .

The following theorem provides a general upper bound on the value (2.5) in terms of an exponentially weighted average of the time-translation modulus $\omega_\xi(\cdot)$ of ξ .

Theorem 2.1. *For every $\kappa > 0$,*

$$v(\kappa) \leq \sqrt{\kappa} \mathbb{E}[(x - \xi_0)^2] + \frac{2}{\sqrt{\kappa}} \int_0^\infty e^{-h/\sqrt{\kappa}} \omega_\xi(h) dh. \quad (2.8)$$

Proof. The idea of the proof is to construct a control $u \in \mathcal{U}$ such that $J_\kappa(u)$ is bounded by the right-hand side of (2.8). It turns out that this is achieved by a myopic control which steers X_t^u toward ξ_t at every instant $t \in [0, T]$ at a suitably chosen rate.

Specifically, set $a = \sqrt{\kappa}$, consider the progressively measurable process X defined by

$$X_t = x + \frac{1}{a} \int_0^t (\xi_s - X_s) ds, \quad 0 \leq t \leq T, \quad (2.9)$$

and define

$$u_t := \frac{1}{a}(\xi_t - X_t), \quad 0 \leq t \leq T.$$

The linear equation (2.9) has the explicit solution

$$X_t = e^{-t/a} x + \frac{1}{a} \int_0^t e^{-(t-s)/a} \xi_s ds. \quad (2.10)$$

By Jensen's inequality and (2.3), we have $X \in L^2(\Omega \times [0, T])$, hence $u \in \mathcal{U}$ and $X = X^u$.

Consider the error

$$e_t := X_t - \xi_t.$$

Since $u_t = -a^{-1}e_t$ and $a^2 = \kappa$, the tracking objective in (2.4) can be written as

$$J_\kappa(u) = \mathbb{E} \int_0^T e_t^2 dt. \quad (2.11)$$

Moreover, let $\mu_a(dh) = a^{-1}e^{-h/a} dh$, a probability measure on $[0, \infty)$. By the change of variables $h = t - s$ in (2.10), we can write

$$X_t = e^{-t/a} x + \int_0^t \xi_{t-h} \mu_a(dh).$$

Since $\int_t^\infty \mu_a(dh) = e^{-t/a}$, we obtain

$$\begin{aligned} e_t &= e^{-t/a} x + \int_0^t \xi_{t-h} \mu_a(dh) - \int_0^{+\infty} \xi_t \mu_a(dh) \\ &= e^{-t/a} (x - \xi_0) - \int_0^\infty (\xi_t - \xi_{(t-h)^+}) \mu_a(dh). \end{aligned} \quad (2.12)$$

Hence, by $(r + s)^2 \leq 2r^2 + 2s^2$ and Jensen's inequality,

$$|e_t|^2 \leq 2e^{-2t/a}(x - \xi_0)^2 + 2 \int_0^\infty |\xi_t - \xi_{(t-h)+}|^2 \mu_a(dh).$$

Integrating over $t \in [0, T]$, taking expectations, and applying Fubini's theorem yield

$$\begin{aligned} \mathbb{E} \int_0^T e_t^2 dt &\leq 2\mathbb{E}[(x - \xi_0)^2] \int_0^T e^{-2t/a} dt + 2 \int_0^\infty \omega_\xi(h) \mu_a(dh) \\ &\leq a \mathbb{E}[(x - \xi_0)^2] + \frac{2}{a} \int_0^\infty e^{-h/a} \omega_\xi(h) dh. \end{aligned}$$

Combining this estimate with (2.11) and $v(\kappa) \leq J_\kappa(u)$ proves (2.8). $\square$

Remark 2.2. If $x = \xi_0$ almost surely, the proof of Theorem 2.1 gives the sharper estimate

$$v(\kappa) \leq \frac{1}{\sqrt{\kappa}} \int_0^\infty e^{-h/\sqrt{\kappa}} \omega_\xi(h) dh. \quad (2.13)$$

Indeed, in that case, the initial displacement term vanishes, and Jensen's inequality can be applied directly in (2.12) to omit the factor two.

The next corollary shows that the L^2 time-translation seminorm of ξ in (2.7) provides the natural constant associated with the generic rate $\sqrt{\kappa}$ in the limiting behavior of $v(\kappa)$. We write $\Gamma(\cdot)$ for the usual Gamma function.

Corollary 2.3. *Suppose that, for some constants $L > 0$ and $\alpha > 0$,*

$$\omega_\xi(h) \leq Lh^\alpha, \quad h > 0. \quad (2.14)$$

Then, for every $\kappa > 0$,

$$v(\kappa) \leq \sqrt{\kappa} \mathbb{E}[(x - \xi_0)^2] + 2L \kappa^{\alpha/2} \Gamma(\alpha + 1). \quad (2.15)$$

Consequently, for $0 < \kappa \leq 1$,

$$v(\kappa) \leq \kappa^{(\alpha \wedge 1)/2} \left(\mathbb{E}[(x - \xi_0)^2] + 2L \Gamma(\alpha + 1) \right). \quad (2.16)$$

In particular, if $[\xi]_{\text{tr}}^2 < \infty$, it holds that

$$v(\kappa) \leq \sqrt{\kappa} \left(\mathbb{E}[(x - \xi_0)^2] + 2[\xi]_{\text{tr}}^2 \right) \quad (2.17)$$

for all $\kappa > 0$. Finally, if $x = \xi_0$ a.s., we have the sharper estimate

$$v(\kappa) \leq L \kappa^{\alpha/2} \Gamma(\alpha + 1), \quad \kappa > 0. \quad (2.18)$$

Proof. Insert (2.14) into (2.8). With the change of variables $h = \sqrt{\kappa} r$,

$$\frac{1}{\sqrt{\kappa}} \int_0^\infty e^{-h/\sqrt{\kappa}} h^\alpha dh = \kappa^{\alpha/2} \int_0^\infty e^{-r} r^\alpha dr = \Gamma(\alpha + 1) \kappa^{\alpha/2}.$$

This proves (2.15). For $0 < \kappa \leq 1$, the two terms in (2.15) are both bounded by the right-hand side of (2.16), depending on whether $\alpha < 1$ or $\alpha \geq 1$. Moreover, (2.17) is the special case with $L = [\xi]_{\text{tr}}^2$ and $\alpha = 1$. Lastly, if $x = \xi_0$ a.s., the sharper estimate in (2.13) from Remark 2.2 applies, and the same change of variables from above yields (2.18). $\square$

The preceding upper bounds admit a complementary lower bound in terms of the same time-translation modulus, showing that under mild scale regularity of ω_ξ , the modulus determines the convergence rate up to multiplicative constants.

Proposition 2.4 (Lower bound). *For every $0 < \kappa \leq T^2/2$,*

$$v(\kappa) \geq \frac{1}{12} \omega_\xi(\sqrt{2\kappa}) - \frac{\sqrt{2}}{4} \sqrt{\kappa} \mathbb{E}[(x - \xi_0)^2]. \quad (2.19)$$

In particular, if $x = \xi_0$ almost surely, then

$$\frac{1}{12} \omega_\xi(\sqrt{2\kappa}) \leq v(\kappa) \leq \frac{1}{\sqrt{\kappa}} \int_0^\infty e^{-h/\sqrt{\kappa}} \omega_\xi(h) dh, \quad 0 < \kappa \leq T^2/2. \quad (2.20)$$

If $x = \xi_0$ almost surely and

$$ch^\alpha \leq \omega_\xi(h) \leq Ch^\alpha, \quad 0 < h \leq T \quad (2.21)$$

for some $c, C, \alpha > 0$, then

$$\frac{c2^{\alpha/2}}{12} \kappa^{\alpha/2} \leq v(\kappa) \leq CT(\alpha+1)\kappa^{\alpha/2}, \quad 0 < \kappa \leq T^2/2 \quad (2.22)$$

and thus

$$v(\kappa) \asymp \kappa^{\alpha/2} \asymp \omega_\xi(\sqrt{\kappa}), \quad \kappa \downarrow 0.$$

Proof. Fix $u \in \mathcal{U}$, and write

$$X_t := x + \int_0^t u_s ds, \quad e_t := X_t - \xi_t, \quad A := \mathbb{E} \int_0^T e_t^2 dt, \quad B := \mathbb{E} \int_0^T u_t^2 dt,$$

so that $J_\kappa(u) = \frac{1}{2}(A + \kappa B)$. Fix $0 < h \leq T$. Since

$$\xi_t - \xi_{(t-h)^+} = (X_t - X_{(t-h)^+}) - e_t + e_{(t-h)^+},$$

the inequality $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ gives

$$\omega_\xi(h) \leq 3 \mathbb{E} \int_0^T |X_t - X_{(t-h)^+}|^2 dt + 3A + 3 \mathbb{E} \int_0^T e_{(t-h)^+}^2 dt.$$

Because $e_0 = x - \xi_0$, a change of variables yields

$$\mathbb{E} \int_0^T e_{(t-h)^+}^2 dt = h \mathbb{E}[(x - \xi_0)^2] + \mathbb{E} \int_0^{T-h} e_t^2 dt \leq h \mathbb{E}[(x - \xi_0)^2] + A.$$

Moreover, by Cauchy–Schwarz and Fubini,

$$\mathbb{E} \int_0^T |X_t - X_{(t-h)^+}|^2 dt = \mathbb{E} \int_0^T \left| \int_{(t-h)^+}^t u_s ds \right|^2 dt \leq h \mathbb{E} \int_0^T \int_{(t-h)^+}^t u_s^2 ds dt \leq h^2 B.$$

Consequently,

$$\omega_\xi(h) \leq 6A + 3h^2 B + 3h \mathbb{E}[(x - \xi_0)^2]. \quad (2.23)$$

Now let $0 < \kappa \leq T^2/2$ and choose $h = \sqrt{2\kappa}$. Then $h \leq T$, and

$$6A + 3h^2B = 6A + 6\kappa B = 12J_\kappa(u).$$

Thus, (2.23) implies

$$J_\kappa(u) \geq \frac{1}{12} \omega_\xi(\sqrt{2\kappa}) - \frac{\sqrt{2}}{4} \sqrt{\kappa} \mathbb{E}[(x - \xi_0)^2].$$

Taking the infimum over $u \in \mathcal{U}$ proves (2.19). If $x = \xi_0$ almost surely, the lower bound in (2.20) follows immediately, while the upper bound is (2.13) from Remark 2.2.

Finally, assume (2.21). The lower bound in (2.22) follows from (2.19). Since $\omega_\xi(h) = \omega_\xi(T)$ for $h \geq T$, the upper bound in (2.21) extends to all $h > 0$. Hence (2.18), with $L = C$, gives the upper bound in (2.22). The final equivalences follow from (2.21). $\square$

2.1 Examples and special cases

We first show that every square-integrable càdlàg semimartingale target process ξ has a finite L^2 time-translation seminorm, as defined in (2.7), and thus has the generic $\sqrt{\kappa}$ rate established in (2.17).

Proposition 2.5 (Semimartingale target). *Suppose*

$$\xi_t = \xi_0 + M_t + A_t, \quad 0 \leq t \leq T,$$

where M is a square-integrable càdlàg martingale with $M_0 = 0$ and A is an adapted càdlàg finite-variation process with $A_0 = 0$, such that

$$\mathbb{E}[\langle M \rangle_T] + \mathbb{E}[|A|_T^2] < \infty.$$

Then, for every $0 < h \leq T$,

$$\omega_\xi(h) \leq 2h \mathbb{E}[\langle M \rangle_T] + 2h \mathbb{E}[|A|_T^2]. \quad (2.24)$$

In particular, $[\xi]_{\text{tr}}^2 < \infty$, and therefore $v(\kappa) = O(\sqrt{\kappa})$.

Proof. Write

$$\xi_t - \xi_{(t-h)+} = (M_t - M_{(t-h)+}) + (A_t - A_{(t-h)+}).$$

Hence

$$\omega_\xi(h) \leq 2\omega_M(h) + 2\omega_A(h),$$

where

$$\omega_M(h) := \mathbb{E} \int_0^T |M_t - M_{(t-h)+}|^2 dt, \quad \omega_A(h) := \mathbb{E} \int_0^T |A_t - A_{(t-h)+}|^2 dt.$$

For the martingale part, using Fubini's theorem and the properties of the predictable quadratic variation $\langle M \rangle$ of M yields

$$\begin{aligned} \omega_M(h) &= \mathbb{E} \int_0^h M_t^2 dt + \mathbb{E} \int_h^T (M_t - M_{t-h})^2 dt \\ &= \mathbb{E} \int_0^h \langle M \rangle_t dt + \mathbb{E} \int_h^T (\langle M \rangle_t - \langle M \rangle_{t-h}) dt \\ &= \mathbb{E} \int_{T-h}^T \langle M \rangle_t dt \leq h \mathbb{E}[\langle M \rangle_T]. \end{aligned} \quad (2.25)$$

For the finite-variation part, let $V_t := |A|_t$ and note that

$$|A_t - A_{(t-h)^+}| \leq \int_{((t-h)^+, t]} dV_s =: D_t^{(h)}.$$

Moreover, since $D_t^{(h)} \leq V_T$, we obtain via Fubini's theorem that

$$\begin{aligned} \int_0^T |A_t - A_{(t-h)^+}|^2 dt &\leq V_T \int_0^T D_t^{(h)} dt = V_T \int_0^T \int_{((t-h)^+, t]} dV_s dt \\ &= V_T \int_0^T \int_s^{(s+h) \wedge T} dt dV_s \leq h V_T^2. \end{aligned} \quad (2.26)$$

Taking expectations and applying once more Fubini's theorem yields $\omega_A(h) \leq h \mathbb{E}[|A|_T^2]$. Combining the two estimates proves (2.24). Lastly, since $\omega_\xi(h) = \omega_\xi(T)$ for $h \geq T$, this also implies $[\xi]_{\text{tr}}^2 < \infty$, and thus $v(\kappa) = O(\sqrt{\kappa})$ by virtue of (2.17). $\square$

Note that the bounds of Theorem 2.1 and Corollary 2.3 are one-sided. To see that the $\sqrt{\kappa}$ rate for semimartingale target processes (Proposition 2.5) is sharp in specific cases, it is convenient to use the exact formula of Bank et al. [7, Theorem 3.1, formula (8)] for fixed κ . This formula simplifies substantially in the two regimes considered below: continuous martingale targets and deterministic finite-variation targets.

Proposition 2.6 (Continuous martingale target). *Assume that*

$$\xi_t = \xi_0 + M_t, \quad 0 \leq t \leq T,$$

where $\xi_0 \in \mathbb{R}$ is deterministic and M is a continuous square-integrable martingale with $M_0 = 0$. Then

$$v(\kappa) = \frac{\sqrt{\kappa}}{2} \tanh\left(\frac{T}{\sqrt{\kappa}}\right) (x - \xi_0)^2 + \frac{1}{2} \mathbb{E} \int_0^T \sqrt{\kappa} \tanh\left(\frac{T-t}{\sqrt{\kappa}}\right) d\langle M \rangle_t. \quad (2.27)$$

In particular,

$$v(\kappa) = \frac{\sqrt{\kappa}}{2} \left((x - \xi_0)^2 + \mathbb{E}[\langle M \rangle_T] \right) + o(\sqrt{\kappa}), \quad \kappa \downarrow 0, \quad (2.28)$$

so that the $\sqrt{\kappa}$ rate is sharp whenever ξ is not identically equal to x .

Proof. Continuous adapted processes are predictable, so the exact unconstrained value formula of Bank et al. [7, Theorem 3.1, formula (8)] applies, and since ξ is a martingale, the formula simplifies to the expression in (2.27). Hence

$$\frac{2v(\kappa)}{\sqrt{\kappa}} = \tanh\left(\frac{T}{\sqrt{\kappa}}\right) (x - \xi_0)^2 + \mathbb{E} \int_0^T \tanh\left(\frac{T-t}{\sqrt{\kappa}}\right) d\langle M \rangle_t.$$

Since $\tanh(T/\sqrt{\kappa}) \rightarrow 1$ as $\kappa \downarrow 0$, the first term converges to $(x - \xi_0)^2$. For the second term, since $0 \leq \tanh((T-t)/\sqrt{\kappa}) \leq 1$ for all $0 \leq t \leq T$, $\tanh((T-t)/\sqrt{\kappa}) \rightarrow 1$ for all $t < T$, and $\langle M \rangle$ is continuous, dominated convergence yields

$$\mathbb{E} \int_0^T \tanh\left(\frac{T-t}{\sqrt{\kappa}}\right) d\langle M \rangle_t \longrightarrow \mathbb{E} \int_0^T d\langle M \rangle_t = \mathbb{E}[\langle M \rangle_T]$$

for $\kappa \downarrow 0$ and we conclude (2.28). $\square$

Remark 2.7 (Brownian motion target). If $M = W$ is a standard Brownian motion, then $\langle M \rangle_t = t$ for all $0 \leq t \leq T$, and (2.27) reduces to

$$\begin{aligned} v(\kappa) &= \frac{\sqrt{\kappa}}{2} \tanh\left(\frac{T}{\sqrt{\kappa}}\right) (x - \xi_0)^2 + \frac{\kappa}{2} \log\left(\cosh\left(\frac{T}{\sqrt{\kappa}}\right)\right) \\ &= \frac{\sqrt{\kappa}}{2} \tanh\left(\frac{T}{\sqrt{\kappa}}\right) (x - \xi_0)^2 + \frac{\sqrt{\kappa} T}{2} - \frac{\kappa}{2} \log 2 + \frac{\kappa}{2} \log(1 + e^{-2T/\sqrt{\kappa}}). \end{aligned}$$

Hence, using the fact that $\tanh(T/\sqrt{\kappa}) = 1 + O(e^{-2T/\sqrt{\kappa}})$ and $\log(1 + e^{-2T/\sqrt{\kappa}}) = O(e^{-2T/\sqrt{\kappa}})$ for $\kappa \downarrow 0$, we obtain

$$v(\kappa) = \frac{\sqrt{\kappa}}{2} ((x - \xi_0)^2 + T) - \frac{\kappa}{2} \log 2 + O\left(\sqrt{\kappa} e^{-2T/\sqrt{\kappa}}\right), \quad \kappa \downarrow 0.$$

In particular,

$$v(\kappa) = \frac{\sqrt{\kappa}}{2} ((x - \xi_0)^2 + T) + O(\kappa), \quad \kappa \downarrow 0,$$

showing most transparently that the $\sqrt{\kappa}$ upper bound has the correct order.

For the next example, it is notationally convenient to introduce the Sobolev space

$$\begin{aligned} H^1(0, T) = \left\{ g \in L^2(0, T) : g \text{ admits a representative } \tilde{g} \text{ with } \tilde{g}(t) = \tilde{g}(s) + \int_s^t \dot{g}(r) dr \right. \\ \left. \text{for all } s, t \in [0, T], \text{ for some } \dot{g} \in L^2(0, T) \right\}. \end{aligned}$$

Proposition 2.8 (Deterministic targets with finitely many jumps). *Assume that the target process is deterministic and of the form*

$$\xi_t = \xi_0 + g(t) + \sum_{j=1}^N \Delta_j \mathbf{1}_{\{t \geq t_j\}}, \quad 0 \leq t \leq T, \quad (2.29)$$

where $N \geq 0$, $g \in H^1(0, T)$, $g(0) = 0$, and the jump times satisfy $0 < t_1 < \dots < t_N < T$. Then

$$v(\kappa) = \sqrt{\kappa} \left(\frac{1}{2} (x - \xi_0)^2 + \frac{1}{4} \sum_{j=1}^N \Delta_j^2 \right) + o(\sqrt{\kappa}), \quad \kappa \downarrow 0. \quad (2.30)$$

In particular, whenever $x \neq \xi_0$ or at least one jump is nonzero, the generic $\sqrt{\kappa}$ rate cannot be improved.

Proof. Set $a := \sqrt{\kappa}$. Since ξ differs at most at the deterministic times $t_1, \dots, t_N$ from a predictable deterministic process, the exact unconstrained value formula of Bank et al. [7, Theorem 3.1, formula (8)] applies after changing ξ on a Lebesgue-null set. Because the target process is deterministic, this formula simplifies to

$$v(\kappa) = \frac{a}{2} \tanh\left(\frac{T}{a}\right) (x - \hat{\xi}_0^\kappa)^2 + \frac{1}{2} \int_0^T (\xi_t - \hat{\xi}_t^\kappa)^2 dt, \quad (2.31)$$

where

$$\hat{\xi}_t^\kappa := \int_t^T \xi_u K_a(t, u) du, \quad K_a(t, u) := \frac{\cosh((T-u)/a)}{a \sinh((T-t)/a)}, \quad 0 \leq t \leq u < T.$$

Note that for all $0 \leq t < T$, the kernel $K_a(t, \cdot)$ integrates to one over $[t, T]$. Let us decompose $\xi = g + J$ with

$$J(t) := \xi_0 + \sum_{j=1}^N \Delta_j \mathbf{1}_{\{t \geq t_j\}}, \quad 0 \leq t \leq T.$$

By linearity, $\hat{\xi}^\kappa = \hat{g}^\kappa + \hat{J}^\kappa$, where $\hat{g}_t^\kappa := \int_t^T g(u) K_a(t, u) du$ and $\hat{J}_t^\kappa := \int_t^T J(u) K_a(t, u) du$.

For the H^1 part, using $g(t) - g(u) = -\int_t^u \dot{g}(r) dr$ for $u > t$ and

$$\int_r^T K_a(t, u) du = \frac{\sinh((T-r)/a)}{\sinh((T-t)/a)} \leq e^{-(r-t)/a}, \quad t \leq r < T,$$

we obtain with Fubini's theorem that

$$|g(t) - \hat{g}_t^\kappa| = \left| \int_t^T (g(t) - g(u)) K_a(t, u) du \right| \leq \int_t^T |\dot{g}(r)| e^{-(r-t)/a} dr.$$

Writing

$$\int_t^T |\dot{g}(r)| e^{-(r-t)/a} dr = \int_{\mathbb{R}} k_a(t-r) |\dot{g}(r)| dr = (k_a * |\dot{g}|)(t)$$

with $k_a(s) := e^{s/a} \mathbf{1}_{(-\infty, 0]}(s)$ (and extending $\dot{g}$ by zero outside $[0, T]$), Young's convolution inequality gives

$$\|g - \hat{g}^\kappa\|_{L^2(0, T)} \leq \|k_a * |\dot{g}|\|_{L^2(\mathbb{R})} \leq \|k_a\|_{L^1(\mathbb{R})} \|\dot{g}\|_{L^2(0, T)} \leq a \|\dot{g}\|_{L^2(0, T)},$$

and thus

$$\int_0^T (g(t) - \hat{g}_t^\kappa)^2 dt = O(a^2). \quad (2.32)$$

Similarly,

$$|\hat{g}_0^\kappa| = \left| \int_0^T g(u) K_a(0, u) du \right| \leq \int_0^T |\dot{g}(r)| e^{-r/a} dr \leq \sqrt{\frac{a}{2}} \|\dot{g}\|_{L^2(0, T)},$$

where we used Cauchy-Schwarz in the last inequality, and hence

$$\hat{g}_0^\kappa = O(a^{1/2}). \quad (2.33)$$

For the pure jump part, a direct computation gives

$$\hat{J}_t^\kappa = \int_t^T J(u) K_a(t, u) du = \xi_0 + \sum_{j: t \geq t_j} \Delta_j + \sum_{j: t < t_j} \Delta_j \int_{t_j}^T K_a(t, u) du,$$

and therefore

$$\hat{J}_t^\kappa - J(t) = \sum_{j=1}^N \Delta_j q_j^a(t) \mathbf{1}_{\{t < t_j\}}, \quad q_j^a(t) := \frac{\sinh((T-t_j)/a)}{\sinh((T-t)/a)}.$$

Consequently,

$$\int_0^T (\hat{J}_t^\kappa - J(t))^2 dt = \sum_{j=1}^N \Delta_j^2 \int_0^{t_j} (q_j^a(t))^2 dt + 2 \sum_{1 \leq j < k \leq N} \Delta_j \Delta_k \int_0^{t_j} q_j^a(t) q_k^a(t) dt. \quad (2.34)$$

Next, since

$$q_j^a(t) = e^{-(t_j-t)/a} \frac{1 - e^{-2(T-t_j)/a}}{1 - e^{-2(T-t)/a}}, \quad 0 \leq t < t_j < T,$$

and

$$\frac{1}{1 - e^{-2(T-t)/a}} = 1 + \frac{e^{-2(T-t)/a}}{1 - e^{-2(T-t)/a}} = 1 + O(e^{-2(T-t)/a}),$$

we get

$$\frac{1 - e^{-2(T-t_j)/a}}{1 - e^{-2(T-t)/a}} = (1 - e^{-2(T-t_j)/a}) (1 + O(e^{-2(T-t)/a})) = 1 + O(e^{-2(T-t_j)/a})$$

for all $0 \leq t \leq t_j$, because $2(T - t_j) \leq 2(T - t)$. This yields

$$q_j^a(t) = e^{-(t_j-t)/a} (1 + O(e^{-2(T-t_j)/a}))$$

uniformly on $[0, t_j]$ and thus

$$\begin{aligned} \int_0^{t_j} (q_j^a(t))^2 dt &= (1 + O(e^{-2(T-t_j)/a})) \int_0^{t_j} e^{-2(t_j-t)/a} dt \\ &= \frac{a}{2} (1 - e^{-2t_j/a}) (1 + O(e^{-2(T-t_j)/a})) = \frac{a}{2} + o(a), \quad a \downarrow 0. \end{aligned} \quad (2.35)$$

Similarly, for $j < k$ with $t_j < t_k$, we obtain

$$\int_0^{t_j} q_j^a(t) q_k^a(t) dt = e^{-(t_k-t_j)/a} (1 + O(e^{-c/a})) \int_0^{t_j} e^{-2(t_j-t)/a} dt = o(a), \quad (2.36)$$

where $c > 0$ is some constant. Plugging (2.35) and (2.36) back into (2.34) yields

$$\int_0^T (\hat{J}_t^\kappa - J(t))^2 dt = \frac{a}{2} \sum_{j=1}^N \Delta_j^2 + o(a). \quad (2.37)$$

Moreover, note that

$$\hat{J}_0^\kappa - \xi_0 = \sum_{j=1}^N \Delta_j \frac{\sinh((T - t_j)/a)}{\sinh(T/a)} = o(1), \quad a \downarrow 0.$$

Hence, together with (2.33), we can write

$$\hat{\xi}_0^\kappa = \hat{g}_0^\kappa + \hat{J}_0^\kappa - \xi_0 + \xi_0 = \xi_0 + o(1), \quad a \downarrow 0. \quad (2.38)$$

Finally, we can put everything together. Regarding the second term in (2.31), using $\xi - \hat{\xi}^\kappa = (J - \hat{J}^\kappa) + (g - \hat{g}^\kappa)$, we obtain

$$\begin{aligned} \int_0^T (\xi_t - \hat{\xi}_t^\kappa)^2 dt &= \int_0^T (J(t) - \hat{J}_t^\kappa)^2 dt + \int_0^T (g(t) - \hat{g}_t^\kappa)^2 dt \\ &\quad + 2 \int_0^T (J(t) - \hat{J}_t^\kappa)(g(t) - \hat{g}_t^\kappa) dt \frac{a}{2} \sum_{j=1}^N \Delta_j^2 + o(a) \end{aligned} \quad (2.39)$$

due to (2.37), (2.32), and the Cauchy–Schwarz estimate

$$\left| \int_0^T (J(t) - \hat{J}_t^\kappa)(g(t) - \hat{g}_t^\kappa) dt \right| \leq \|J - \hat{J}^\kappa\|_{L^2(0,T)} \|g - \hat{g}^\kappa\|_{L^2(0,T)} = O(a^{3/2}) = o(a), \quad (2.40)$$

since $\|J - \hat{J}^\kappa\|_{L^2(0,T)} = O(a^{1/2})$ and $\|g - \hat{g}^\kappa\|_{L^2(0,T)} = O(a)$. In particular, we observe that the jump component determines the leading $O(a)$ error in (2.39), while the smoother g -component contributes only $O(a^2)$, and its interaction with the jump component is $O(a^{3/2})$. Regarding the first term in (2.31), we can write

$$\frac{a}{2} \tanh\left(\frac{T}{a}\right) (x - \hat{\xi}_0^\kappa)^2 = \frac{a}{2} (x - \xi_0)^2 + o(a), \quad (2.41)$$

because $\tanh(T/a) = 1 + o(1)$ and $(x - \hat{\xi}_0^\kappa)^2 = (x - \xi_0)^2 + o(1)$ by (2.38) as $a \downarrow 0$, as well as the fact that $a o(1) = o(a)$. Thus,

$$v(\kappa) = \frac{a}{2} \tanh\left(\frac{T}{a}\right) (x - \hat{\xi}_0^\kappa)^2 + \frac{1}{2} \int_0^T (\xi_t - \hat{\xi}_t^\kappa)^2 dt = \frac{a}{2} (x - \xi_0)^2 + \frac{a}{4} \sum_{j=1}^N \Delta_j^2 + o(a),$$

which is (2.30). $\square$

Remark 2.9. In the setting of Proposition 2.8, if there are no jumps and $x = \xi_0$, then the choice $u_t = \dot{g}(t)$ for a deterministic, absolutely continuous target process $\xi_t = \xi_0 + g(t)$ with $g \in H^1(0, T)$ tracks the target in (2.4) exactly, and thus

$$v(\kappa) \leq \frac{\kappa}{2} \int_0^T \dot{\xi}_t^2 dt.$$

In particular, $v(\kappa) = O(\kappa)$. Thus, once the initial mismatch is removed, the $\sqrt{\kappa}$ contribution in Proposition 2.8 is produced entirely by the jumps.

Our final example is concerned with the case where the target process follows a fractional Brownian motion with Hurst parameter $H \in (0, 1)$. It is well known that the fractional Brownian motion is a semimartingale with respect to its natural filtration only in the Brownian case $H = 0.5$. Hence, when $H \neq 0.5$, it serves as a natural example of a non-semimartingale target process which satisfies (2.3).

We first observe that the fractional Brownian motion has finite L^2 time-translation seminorm, as defined in (2.7), only if $H \geq 0.5$. Then, we show that the limiting behavior of $v(\kappa)$ is governed by the Hurst parameter H through the scaling parameter κ^H .

Lemma 2.10. *Let $B^H = (B_t^H)_{0 \leq t \leq T}$ be a fractional Brownian motion with Hurst parameter $H \in (0, 1)$, considered in its natural filtration. Its L^2 time-translation modulus is given by*

$$\omega_{B^H}(h) = Th^{2H} - \frac{2H}{2H+1}h^{2H+1}, \quad 0 < h \leq T. \quad (2.42)$$

Moreover,

$$[B^H]_{\text{tr}}^2 = \sup_{h>0} \frac{\omega_{B^H}(h)}{h} = \begin{cases} \infty, & 0 < H < \frac{1}{2}, \\ T, & H = \frac{1}{2}, \\ \frac{T^{2H}}{2H} \left(1 - \frac{1}{4H^2}\right)^{2H-1}, & H > \frac{1}{2}. \end{cases} \quad (2.43)$$

Proof. Let $0 < h \leq T$. Since $B_0^H = 0$ and $\mathbb{E}|B_t^H - B_s^H|^2 = |t - s|^{2H}$, we obtain

$$\begin{aligned} \omega_{B^H}(h) &= \mathbb{E} \int_0^h |B_t^H|^2 dt + \mathbb{E} \int_h^T |B_t^H - B_{t-h}^H|^2 dt \\ &= \int_0^h t^{2H} dt + (T-h)h^{2H} = Th^{2H} - \frac{2H}{2H+1}h^{2H+1}. \end{aligned}$$

For $h \geq T$, we get

$$\omega_{B^H}(h) = \omega_{B^H}(T) = \frac{T^{2H+1}}{2H+1}.$$

The claim (2.43) then follows from direct computations. $\square$

Proposition 2.11 (Fractional Brownian motion). *Let $B^H = (B_t^H)_{0 \leq t \leq T}$ be fractional Brownian motion with Hurst parameter $H \in (0, 1)$, considered in its natural filtration. Let*

$$\xi_t = B_t^H, \quad 0 \leq t \leq T, \quad \text{and} \quad x = 0.$$

Then

$$v(\kappa) \asymp \kappa^H, \quad \kappa \downarrow 0.$$

More explicitly, for $0 < \kappa \leq T^2/2$,

$$\frac{\kappa^H}{12} \left(T^{2H} - \frac{2H}{2H+1} 2^{H+\frac{1}{2}} \sqrt{\kappa} \right) \leq v(\kappa) \leq T\Gamma(2H+1)\kappa^H. \quad (2.44)$$

Proof. By Lemma 2.10, the time-translation modulus of $\xi = B^H$ satisfies

$$\omega_\xi(h) \leq Th^{2H}, \quad h > 0.$$

Indeed, this follows directly from (2.42) for $0 < h \leq T$, whereas for $h \geq T$,

$$\omega_\xi(h) = \frac{T^{2H+1}}{2H+1} \leq Th^{2H}.$$

Thus, (2.14) holds with $L = T$ and $\alpha = 2H$. Since $x = \xi_0 = 0$, the sharper estimate (2.18) yields

$$v(\kappa) \leq T\Gamma(2H+1)\kappa^H, \quad \kappa > 0.$$

For the lower bound, let $0 < \kappa \leq T^2/2$. Since $x = \xi_0$, Proposition 2.4 gives

$$v(\kappa) \geq \frac{1}{12} \omega_\xi(\sqrt{2\kappa}).$$

Moreover, $\sqrt{2\kappa} \leq T$, so (2.42) implies

$$\begin{aligned} v(\kappa) &\geq \frac{1}{12} \left(T(2\kappa)^H - \frac{2H}{2H+1} (2\kappa)^{H+\frac{1}{2}} \right) \\ &= \frac{\kappa^H}{12} \left(T 2^H - \frac{2H}{2H+1} 2^{H+\frac{1}{2}} \sqrt{\kappa} \right). \end{aligned}$$

This proves (2.44). Since the expression in parentheses in the lower bound converges to $T 2^H > 0$ as $\kappa \downarrow 0$, the two bounds also yield $v(\kappa) \asymp \kappa^H$. $\square$

3 Optimal tracking with terminal state constraint

In this section, we study the asymptotic behavior of the optimal tracking problem from Section 2 with an additional $\mathbb{P}$ -almost sure state constraint at terminal time T .

Recall the admissible class (2.1), the state dynamics of $X^u = (X_t^u)_{0 \leq t \leq T}$ in (2.2) and, for a given progressively measurable target process ξ satisfying (2.3), the tracking objective $J_\kappa(u)$ in (2.4). In addition, we let $\Xi_T \in L^2(\mathcal{F}_T)$ denote a predetermined terminal position and define its càdlàg square-integrable martingale $\Xi = (\Xi_t)_{0 \leq t \leq T}$ via

$$\Xi_t := \mathbb{E}[\Xi_T \mid \mathcal{F}_t], \quad 0 \leq t \leq T.$$

We assume that

$$\int_0^T \frac{d\mathbb{E}[\Xi_t^2]}{T-t} < \infty, \quad (3.1)$$

introduce the constrained admissible set

$$\mathcal{U}^c := \left\{ u \in \mathcal{U} : X_T^u = x + \int_0^T u_s ds = \Xi_T \quad \mathbb{P}\text{-a.s.} \right\} \quad (3.2)$$

and consider the value of the constrained optimization problem

$$v^c(\kappa) := \inf_{u \in \mathcal{U}^c} J_\kappa(u). \quad (3.3)$$

The integral in (3.1) is understood in the Stieltjes sense, with the convention that any atom of $d\mathbb{E}[\Xi_t^2]$ at T contributes $+\infty$. The condition (3.1) was derived in Bank et al. [7] as the relevant reachability criterion; indeed, [7, Lemma 5.4] proves that (3.1) is equivalent to $\mathcal{U}^c \neq \emptyset$, so that it characterizes the terminal positions that can be reached by absolutely continuous state processes with square-integrable controls. We note that (3.1) rules out a terminal jump of Ξ , implies $\Xi_T \in \mathcal{F}_{T-}$, and may be interpreted as a condition on the rate at which the terminal target is revealed as $t \uparrow T$.

As in the previous section, our aim is to quantify the behavior of $v^c(\kappa)$ as $\kappa \downarrow 0$. The relevant objects are the following: For $0 < a \leq T/2$, define

$$\Omega(a) := a \mathbb{E}[(x - \xi_0)^2] + \frac{2}{a} \int_0^\infty e^{-h/a} \omega_\xi(h) dh, \quad (3.4)$$

$$\Theta(a) := \mathbb{E} \int_{T-2a}^T |\xi_t - \Xi_t|^2 dt, \quad (3.5)$$

and

$$\Lambda(a) := \int_{T-2a}^T \frac{d\mathbb{E}[\Xi_t^2]}{T-t}. \quad (3.6)$$

The term $\Omega(a)$ is the familiar right-hand side of (2.8) evaluated at scale a , while $\Theta(a)$ measures the short-horizon mismatch between the running target ξ and the terminal martingale Ξ , and $\Lambda(a)$ records the local cost of learning Ξ_T near maturity. Note that, under our assumptions, we have $\Theta(a) < \infty$ and $\Lambda(a) < \infty$.

The next theorem is the counterpart to Theorem 2.1, providing a general upper bound on the constrained value (3.3).

Theorem 3.1. *Let ξ be progressively measurable and satisfy (2.3), and let $\Xi_T \in L^2(\mathcal{F}_T)$ satisfy (3.1). Then, for every $\kappa \in (0, T^2/4]$, with $a := \sqrt{\kappa}$,*

$$v^c(\kappa) \leq 6(\Omega(a) + \Theta(a) + a^2\Lambda(a)). \quad (3.7)$$

Proof. Fix $\kappa \in (0, T^2/4]$ and set $a := \sqrt{\kappa}$. The idea of the proof is to construct a control $u \in \mathcal{U}^c$ such that $J_\kappa(u)$ is bounded by the right-hand side of (3.7). It turns out that this is achieved by the concatenation of a myopic control targeting ξ up to some suitable time $s \in (0, T)$, and then an asymptotically singular control forcing X_t^u toward the terminal value Ξ_T by time T .

The myopic phase. On $[0, T-a]$, let Y solve

$$Y_t = x + \frac{1}{a} \int_0^t (\xi_r - Y_r) dr, \quad 0 \leq t \leq T-a,$$

and put $e_t := Y_t - \xi_t$. Exactly as in the proof of Theorem 2.1, we have

$$\mathbb{E} \int_0^{T-a} e_t^2 dt \leq \Omega(a).$$

Moreover, introducing $u_t^{(1)} := a^{-1}(\xi_t - Y_t) \in L^2(\Omega \times [0, T-a])$, we obtain

$$\frac{1}{2} \int_0^{T-a} (Y_t - \xi_t)^2 dt + \frac{\kappa}{2} \int_0^{T-a} (u_t^{(1)})^2 dt = \int_0^{T-a} e_t^2 dt,$$

and for every $s \in [T-2a, T-a]$, we can conclude that

$$\mathbb{E} \left[\frac{1}{2} \int_0^s (Y_t - \xi_t)^2 dt + \frac{\kappa}{2} \int_0^s (u_t^{(1)})^2 dt \right] \leq \Omega(a). \quad (3.8)$$

Now, choose $s \in [T-2a, T-a]$ such that

$$\mathbb{E}[e_s^2 + |\Xi_s - \xi_s|^2] \leq \frac{1}{a} \int_{T-2a}^{T-a} \mathbb{E}[e_t^2 + |\Xi_t - \xi_t|^2] dt.$$

Since

$$\int_{T-2a}^{T-a} \mathbb{E}[e_t^2] dt \leq \Omega(a), \quad \int_{T-2a}^{T-a} \mathbb{E}[|\Xi_t - \xi_t|^2] dt \leq \Theta(a),$$

it follows that

$$\mathbb{E}[e_s^2 + |\Xi_s - \xi_s|^2] \leq \frac{\Omega(a) + \Theta(a)}{a}. \quad (3.9)$$

We also introduce the constant

$$B := \mathbb{E}[|\Xi_s - Y_s|^2] \quad (3.10)$$

and note that

$$B = \mathbb{E}[|\Xi_s - \xi_s + \xi_s - Y_s|^2] \leq 2 \mathbb{E}[e_s^2 + |\Xi_s - \xi_s|^2] \leq \frac{2}{a} (\Omega(a) + \Theta(a)). \quad (3.11)$$

The terminal bridge. On $[s, T)$, define X by

$$dX_t = \frac{\Xi_t - X_t}{T - t} dt, \quad X_s = Y_s.$$

Its explicit solution is given by

$$X_t = \frac{T - t}{T - s} Y_s + (T - t) \int_s^t \frac{\Xi_r}{(T - r)^2} dr, \quad s \leq t < T. \quad (3.12)$$

We first argue that $\lim_{t \uparrow T} X_t = \Xi_T$ $\mathbb{P}$ -a.s. To this end, note that due to the martingale convergence theorem, we have

$$\lim_{t \uparrow T} \Xi_t = \mathbb{E}[\Xi_T \mid \mathcal{F}_{T-}] = \Xi_T \quad \mathbb{P}\text{-a.s.}, \quad (3.13)$$

because (3.1) implies $\Xi_T \in \mathcal{F}_{T-}$. Moreover, we can write

$$X_t - \Xi_T = \frac{T - t}{T - s} (Y_s - \Xi_T) + (T - t) \int_s^t \frac{\Xi_r - \Xi_T}{(T - r)^2} dr, \quad s \leq t < T, \quad (3.14)$$

since

$$(T - t) \int_s^t \frac{1}{(T - r)^2} dr = 1 - \frac{T - t}{T - s}.$$

The first term in (3.14) converges to zero as $t \uparrow T$. For the second term, fix an outcome $\omega \in \Omega$ for which $\Xi_t(\omega) \rightarrow \Xi_T(\omega)$ for $t \uparrow T$. Given $\varepsilon > 0$, choose $s < u < T$ such that

$$|\Xi_r(\omega) - \Xi_T(\omega)| \leq \varepsilon, \quad u \leq r < T.$$

Then, for $t > u$, we obtain the estimate

$$(T - t) \int_s^t \frac{|\Xi_r - \Xi_T|}{(T - r)^2} dr \leq (T - t) \int_s^u \frac{|\Xi_r - \Xi_T|}{(T - r)^2} dr + \varepsilon (T - t) \int_u^t \frac{1}{(T - r)^2} dr.$$

The first term tends to zero as $t \uparrow T$, while the second term is bounded by ε . Hence, $\limsup_{t \uparrow T} |X_t - \Xi_T| \leq \varepsilon$ in (3.14). Since $\varepsilon > 0$ is arbitrary, we get that $\lim_{t \uparrow T} X_t = \Xi_T$ $\mathbb{P}$ -a.s.

Next, we define

$$u_t^{(2)} := \frac{\Xi_t - X_t}{T - t}, \quad s \leq t < T.$$

Using integration by parts in (3.12) yields

$$X_t = \frac{T-t}{T-s} Y_s + \Xi_t - \frac{T-t}{T-s} \Xi_s - (T-t) \int_{(s,t]} \frac{1}{T-r} d\Xi_r, \quad s \leq t < T,$$

and hence

$$\Xi_t - X_t = \frac{T-t}{T-s} (\Xi_s - Y_s) + (T-t) \int_{(s,t]} \frac{1}{T-r} d\Xi_r, \quad s \leq t < T, \quad (3.15)$$

so that we obtain the representation

$$u_t^{(2)} = \frac{\Xi_s - Y_s}{T-s} + \int_{(s,t]} \frac{1}{T-r} d\Xi_r, \quad s \leq t < T. \quad (3.16)$$

By the martingale isometry, Fubini's theorem, and (3.1), we have

$$\mathbb{E} \int_s^T \left(\int_{(s,t]} \frac{1}{T-r} d\Xi_r \right)^2 dt = \int_{(s,T]} \frac{d\mathbb{E}[\Xi_r^2]}{T-r} \leq \Lambda(a) < \infty. \quad (3.17)$$

Hence, we can conclude that $u^{(2)} \in L^2(\Omega \times [s, T])$. In particular, note that the concatenated control

$$u_t := u_t^{(1)} \mathbf{1}_{[0,s)}(t) + u_t^{(2)} \mathbf{1}_{[s,T)}(t)$$

belongs to $\mathcal{U}^c$.

The total cost. We start with estimating the cost on $[s, T]$, where we recall $s \in [T-2a, T-a]$. First,

$$|X_t - \xi_t|^2 \leq 2|X_t - \Xi_t|^2 + 2|\Xi_t - \xi_t|^2.$$

Moreover, for $s \leq t < T$, using (3.15) and the constant B defined in (3.10), we obtain

$$\begin{aligned} \mathbb{E}[|\Xi_t - X_t|^2] &\leq 2 \frac{(T-t)^2}{(T-s)^2} B + 2(T-t)^2 \mathbb{E} \left[\left(\int_{(s,t]} \frac{1}{T-r} d\Xi_r \right)^2 \right] \\ &\leq 2 \frac{(T-t)^2}{(T-s)^2} B + 2(T-t) \int_{(s,t]} \frac{1}{T-r} d\mathbb{E}[\Xi_r^2] \\ &\leq 2 \frac{(T-t)^2}{(T-s)^2} B + 2(T-t) \Lambda(a), \end{aligned}$$

where we used the fact that $r \leq t$, which implies $\frac{T-t}{T-r} \leq 1$ and hence $(\frac{T-t}{T-r})^2 \leq \frac{T-t}{T-r}$.

Consequently,

$$\begin{aligned}
\mathbb{E} \int_s^T |X_t - \xi_t|^2 dt &\leq 2 \mathbb{E} \int_s^T |X_t - \Xi_t|^2 dt + 2 \Theta(a) \\
&\leq 2 \int_s^T \left(2 \frac{(T-t)^2}{(T-s)^2} B + 2(T-t)\Lambda(a) \right) dt + 2 \Theta(a) \\
&= \frac{4}{3}(T-s)B + 2(T-s)^2\Lambda(a) + 2\Theta(a) \\
&\leq \frac{8a}{3}B + 8a^2\Lambda(a) + 2\Theta(a),
\end{aligned}$$

because $T-s \leq 2a$. Hence,

$$\frac{1}{2} \mathbb{E} \int_s^T |X_t - \xi_t|^2 dt \leq \frac{4a}{3}B + 4a^2\Lambda(a) + \Theta(a). \quad (3.18)$$

Next, using (3.16) and (3.17) together with the definition of B , we get the upper bound

$$\begin{aligned}
\mathbb{E} \int_s^T (u_t^{(2)})^2 dt &\leq \frac{2}{T-s}B + 2 \mathbb{E} \int_s^T \left(\int_{(s,t]} \frac{1}{T-r} d\Xi_r \right)^2 dt \\
&\leq \frac{2}{a}B + 2\Lambda(a),
\end{aligned}$$

since $T-s \geq a$. Thus, we have

$$\frac{\kappa}{2} \mathbb{E} \int_s^T (u_t^{(2)})^2 dt \leq aB + a^2\Lambda(a). \quad (3.19)$$

Combining (3.18) and (3.19), the cost on the terminal interval $[s, T]$ is bounded by

$$\frac{7a}{3}B + 5a^2\Lambda(a) + \Theta(a).$$

Using (3.11), this yields

$$\mathbb{E} \left[\frac{1}{2} \int_s^T (X_t - \xi_t)^2 dt + \frac{\kappa}{2} \int_s^T (u_t^{(2)})^2 dt \right] \leq \frac{14}{3}\Omega(a) + \frac{17}{3}\Theta(a) + 5a^2\Lambda(a).$$

Finally, adding the contribution from (3.8) on $[0, s]$, we obtain

$$J_\kappa(u) \leq \frac{17}{3}\Omega(a) + \frac{17}{3}\Theta(a) + 5a^2\Lambda(a) \leq 6(\Omega(a) + \Theta(a) + a^2\Lambda(a)).$$

Since the constructed control satisfies $u \in \mathcal{U}^c$, this proves (3.7). $\square$

Remark 3.2. The bound (3.7) separates the constrained cost into three pieces: the same contribution $\Omega(a)$ as in the unconstrained problem, the terminal compatibility defect $\Theta(a)$, and the learning term $a^2\Lambda(a)$. Under the general reachability assumption (3.1), we have $\Lambda(a) \rightarrow 0$ as $a \downarrow 0$, and hence, in general,

$$a^2\Lambda(a) = o(a) = o(\sqrt{\kappa}) \quad \text{when } a = \sqrt{\kappa}.$$

Thus, at the generic $\sqrt{\kappa}$ scale, the new effect of the terminal constraint $X_T^u = \Xi_T$ $\mathbb{P}$ -a.s. is governed by the short-horizon mismatch between the running target ξ and the terminal-target martingale Ξ measured by $\Theta(a)$.

The next corollary is analogous to Corollary 2.3 in the unconstrained case, and again establishes $\sqrt{\kappa}$ as the generic rate.

Corollary 3.3. *Under the assumptions of Theorem 3.1, assume that there exist constants $L_1, L_2 \geq 0$ and exponents $\alpha, \beta > 0$ such that*

$$\omega_\xi(h) \leq L_1 h^\alpha, \quad h > 0, \quad (3.20)$$

and

$$\Theta(a) \leq L_2 a^\beta, \quad 0 < a \leq T/2. \quad (3.21)$$

Then, for every $\kappa \in (0, T^2/4]$,

$$v^c(\kappa) \leq 6\sqrt{\kappa} \mathbb{E}[(x - \xi_0)^2] + 12L_1 \Gamma(\alpha + 1) \kappa^{\alpha/2} + 6L_2 \kappa^{\beta/2} + 6\kappa \Lambda(\sqrt{\kappa}). \quad (3.22)$$

Consequently,

$$v^c(\kappa) = O\left(\kappa^{(1 \wedge \alpha \wedge \beta)/2}\right), \quad \kappa \downarrow 0,$$

and in particular $v^c(\kappa) = O(\sqrt{\kappa})$ if $\alpha, \beta \geq 1$.

If, in addition, $x = \xi_0$ almost surely, then the sharper estimate

$$v^c(\kappa) \leq 6L_1 \Gamma(\alpha + 1) \kappa^{\alpha/2} + 6L_2 \kappa^{\beta/2} + 6\kappa \Lambda(\sqrt{\kappa}) \quad (3.23)$$

holds. In particular,

$$v^c(\kappa) = O\left(\kappa^{(\alpha \wedge \beta \wedge 2)/2}\right), \quad \kappa \downarrow 0,$$

and $v^c(\kappa) = o(\sqrt{\kappa})$ if $\alpha, \beta > 1$.

Proof. From (3.4) and (3.20), with the change of variables $h = ar$,

$$\frac{1}{a} \int_0^\infty e^{-h/a} \omega_\xi(h) dh \leq L_1 a^\alpha \int_0^\infty e^{-r} r^\alpha dr = L_1 \Gamma(\alpha + 1) a^\alpha.$$

Therefore

$$\Omega(a) \leq a \mathbb{E}[(x - \xi_0)^2] + 2L_1 \Gamma(\alpha + 1) a^\alpha.$$

Inserting this estimate and (3.21) into (3.7), with $a = \sqrt{\kappa}$, gives (3.22).

By (3.1), we have $\Lambda(a) \rightarrow 0$ as $a \downarrow 0$. Hence

$$\kappa \Lambda(\sqrt{\kappa}) = o(\kappa) = O\left(\kappa^{(1 \wedge \alpha \wedge \beta)/2}\right), \quad \kappa \downarrow 0,$$

and the first asymptotic estimate follows.

If $x = \xi_0$ a.s., the initial displacement term in the myopic phase vanishes. More precisely, in the proof of Theorem 3.1, using the sharper estimate from Remark 2.2 gives the same bound as (3.7) with $\Omega(a)$ replaced by

$$\Omega_0(a) := \frac{1}{a} \int_0^\infty e^{-h/a} \omega_\xi(h) dh.$$

The estimate above gives

$$\Omega_0(a) \leq L_1 \Gamma(\alpha + 1) a^\alpha.$$

Substituting this into (3.7), again with $a = \sqrt{\kappa}$, proves (3.23), and the remaining claims follow. $\square$

Remark 3.4 (Lower bound). Since the unconstrained value function v is trivially a lower bound for the constrained value function v^c , the lower bounds in Proposition 2.4 immediately extend to the constrained case.

3.1 Examples and special cases

We first record a direct consequence of Theorem 3.1 for target processes ξ with finite L^2 time-translation seminorm, which includes square-integrable semimartingales by Proposition 2.5. Then, we isolate two complementary benchmarks: a mismatch between a running target ξ and a deterministic terminal position $\Xi_T = z \in \mathbb{R}$, which is controlled by $\Theta(a)$, and the exact continuous-martingale case furnished by [7, Theorem 3.2, formula (10)] with sharp rate $\sqrt{\kappa}$.

Corollary 3.5. *In the setting of Theorem 3.1, assume that $[\xi]_{\text{tr}}^2 < \infty$. Then, for every $\kappa \in (0, T^2/4]$,*

$$v^c(\kappa) \leq 6\sqrt{\kappa} \left(\mathbb{E}[(x - \xi_0)^2] + 2[\xi]_{\text{tr}}^2 \right) + 6\Theta(\sqrt{\kappa}) + 6\kappa \Lambda(\sqrt{\kappa}). \quad (3.24)$$

In particular, if

$$\Theta(a) = O(a), \quad a \downarrow 0, \quad (3.25)$$

then

$$v^c(\kappa) = O(\sqrt{\kappa}), \quad \kappa \downarrow 0.$$

Proof. By the definition of the seminorm $[\xi]_{\text{tr}}^2$ in (2.7), we have $\omega_\xi(h) \leq [\xi]_{\text{tr}}^2 h$ for all $h > 0$. Hence (3.4) yields

$$\Omega(a) \leq a \left(\mathbb{E}[(x - \xi_0)^2] + 2[\xi]_{\text{tr}}^2 \right), \quad 0 < a \leq T/2.$$

Insert this estimate into Theorem 3.1 with $a = \sqrt{\kappa}$ to obtain (3.24). If (3.25) holds, then $\Theta(\sqrt{\kappa}) = O(\sqrt{\kappa})$, and Remark 3.2 yields $\kappa \Lambda(\sqrt{\kappa}) = o(\sqrt{\kappa})$. Therefore $v^c(\kappa) = O(\sqrt{\kappa})$. $\square$

Remark 3.6. A simple but quite general sufficient condition for (3.25) is the local uniform L^2 -boundedness

$$\sup_{t \in [T-\delta, T]} \mathbb{E}|\xi_t - \Xi_t|^2 < \infty \quad (3.26)$$

for some $\delta > 0$. Indeed, for $0 < a \leq \delta/2$,

$$\Theta(a) = \mathbb{E} \int_{T-2a}^T |\xi_t - \Xi_t|^2 dt \leq 2a \sup_{t \in [T-\delta, T]} \mathbb{E}|\xi_t - \Xi_t|^2.$$

In particular, observe that exact agreement of ξ and Ξ on a terminal time interval is not needed for the generic $\sqrt{\kappa}$ rate.

Example 3.7 (Deterministic terminal target). Let $\Xi_T = z \in \mathbb{R}$ be deterministic. Then $\Xi_t \equiv z$ for all $t \in [0, T]$, so

$$\Lambda(a) = 0, \quad \Theta(a) = \mathbb{E} \int_{T-2a}^T |\xi_t - z|^2 dt.$$

Hence Theorem 3.1 reduces to

$$v^c(\kappa) \leq 6(\Omega(a) + \Theta(a)), \quad a = \sqrt{\kappa}.$$

Assume, moreover, that ξ is deterministic, piecewise $H^1(0, T)$, and continuous at T with $\xi_T = z$. Then $[\xi]_{\text{tr}}^2 < \infty$ by Proposition 2.5 and

$$\Omega(a) \leq a \left((x - \xi_0)^2 + 2[\xi]_{\text{tr}}^2 \right).$$

If, in addition, $\xi \in H^1(T - \delta, T)$ for some $\delta > 0$, then for $a \leq \delta/2$, applying Jensen yields

$$\Theta(a) = \int_{T-2a}^T |\xi_t - z|^2 dt \leq \int_{T-2a}^T (T-t) \left(\int_t^T |\dot{\xi}_s|^2 ds \right) dt \leq 2a^2 \int_{T-2a}^T |\dot{\xi}_t|^2 dt = o(a^2),$$

since $\dot{\xi} \in L^2(T - \delta, T)$. Consequently,

$$v^c(\kappa) = O(\sqrt{\kappa}), \quad \kappa \downarrow 0.$$

If one further has $\xi \in H^1(0, T)$ and $x = \xi_0$, then the control $u_t = \dot{\xi}_t$ belongs to $\mathcal{U}^c$ and yields the sharper upper bound

$$v^c(\kappa) \leq \frac{\kappa}{2} \int_0^T \dot{\xi}_t^2 dt.$$

Example 3.8 (Continuous martingale running and terminal targets). Assume that

$$\xi_t = \Xi_t = \Xi_0 + M_t, \quad 0 \leq t \leq T,$$

where $\Xi_0 \in \mathbb{R}$ is deterministic, M is a continuous square-integrable martingale with $M_0 = 0$, and (3.1) holds. In this case, the exact constrained value formula of Bank et al. [7, Theorem 3.2, formula (10)] yields

$$v^c(\kappa) = \frac{\sqrt{\kappa}}{2} \coth\left(\frac{T}{\sqrt{\kappa}}\right) (x - \Xi_0)^2 + \frac{1}{2} \mathbb{E} \int_0^T \sqrt{\kappa} \coth\left(\frac{T-t}{\sqrt{\kappa}}\right) d\langle M \rangle_t. \quad (3.27)$$

Then

$$v^c(\kappa) = \frac{\sqrt{\kappa}}{2} \left((x - \Xi_0)^2 + \mathbb{E}[\langle M \rangle_T] \right) + o(\sqrt{\kappa}), \quad \kappa \downarrow 0, \quad (3.28)$$

so that the generic $\sqrt{\kappa}$ rate is sharp whenever ξ is not identically equal to x .

To justify (3.28), first note that $\coth(T/\sqrt{\kappa}) = 1 + o(1)$ as $\kappa \downarrow 0$ and hence

$$\sqrt{\kappa} \coth\left(\frac{T}{\sqrt{\kappa}}\right) = \sqrt{\kappa} + o(\sqrt{\kappa}), \quad \kappa \downarrow 0. \quad (3.29)$$

Next, for $0 \leq t < T$,

$$0 \leq \sqrt{\kappa} \coth\left(\frac{T-t}{\sqrt{\kappa}}\right) - \sqrt{\kappa} = \frac{2\sqrt{\kappa}}{e^{2(T-t)/\sqrt{\kappa}} - 1} \leq \frac{\kappa}{T-t},$$

because $e^{2x} - 1 \geq 2x$ for $x \geq 0$. Since $d\mathbb{E}[\Xi_t^2] = d\mathbb{E}[\langle M \rangle_t]$, the reachability condition (3.1) gives

$$\int_0^T \frac{d\mathbb{E}[\langle M \rangle_t]}{T-t} < \infty.$$

Therefore

$$0 \leq \mathbb{E} \int_0^T \left(\sqrt{\kappa} \coth\left(\frac{T-t}{\sqrt{\kappa}}\right) - \sqrt{\kappa} \right) d\langle M \rangle_t \leq \kappa \int_0^T \frac{d\mathbb{E}[\langle M \rangle_t]}{T-t} = o(\sqrt{\kappa}),$$

and hence

$$\begin{aligned} \mathbb{E} \int_0^T \sqrt{\kappa} \coth\left(\frac{T-t}{\sqrt{\kappa}}\right) d\langle M \rangle_t &= \mathbb{E} \int_0^T \left(\sqrt{\kappa} \coth\left(\frac{T-t}{\sqrt{\kappa}}\right) - \sqrt{\kappa} \right) d\langle M \rangle_t + \sqrt{\kappa} \mathbb{E}[\langle M \rangle_T] \\ &= \sqrt{\kappa} \mathbb{E}[\langle M \rangle_T] + o(\sqrt{\kappa}), \end{aligned}$$

which proves (3.28) together with (3.29).

4 An Obizhaeva–Wang model and its solution

In the remaining two sections, we apply our general tracking results to the optimal execution problem that motivated it. Specifically, we consider a generalized Obizhaeva–Wang model of transient price impact with a random target position at the terminal time and semimartingale execution strategies. In this section, we first solve the resulting singular stochastic control problem using a Hilbert-space approach. Then, in Section 5, we introduce a regularized version of the problem with instantaneous quadratic costs imposed on the trading rate, and study its small-cost asymptotics. We will show that the asymptotic analysis reduces to a constrained tracking problem of the optimal unregularized strategy with terminal state constraint. Consequently, the theory from Section 3 will allow us to quantify the asymptotic error rate when approximating the optimal singular solution in our original Obizhaeva–Wang model with an absolutely continuous regularized strategy.

We fix the time horizon $T \in (0, \infty)$ and let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ be a filtered probability space satisfying the usual assumptions. By convention, $\mathcal{F}_{0-}$ is the augmentation of the trivial σ -field. Let $\beta, \lambda : [0, T] \rightarrow (0, \infty)$ be deterministic functions with the following properties.

Assumption 4.1. (i) β is càdlàg and of finite variation.

(ii) λ is absolutely continuous and bounded away from zero, and its a.e. derivative admits a càdlàg version $\dot{\lambda}$ of finite variation. We set $\lambda_{0-} := \lambda_0$ and define

$$\gamma_t := \log(\lambda_t), \quad 0 \leq t \leq T.$$

Then, γ is also absolutely continuous and its a.e. derivative admits the càdlàg finite-variation version

$$\dot{\gamma}_t := \frac{\dot{\lambda}_t}{\lambda_t}, \quad 0 \leq t \leq T. \quad (4.1)$$

(iii) $2\beta + \dot{\gamma}$ is bounded away from zero, i.e., there exists $c > 0$ such that $2\beta_t + \dot{\gamma}_t > c$ for every $t \in [0, T]$.

We recall that (iii) ensures absence of price manipulation in the model. For later use, we also define the measurable functions

$$\eta_t := \frac{\beta_t + \dot{\gamma}_t}{2\beta_t + \dot{\gamma}_t}, \quad \theta_t := \int_0^t \frac{(\beta_s + \dot{\gamma}_s)^2}{\lambda_s(2\beta_s + \dot{\gamma}_s)} ds, \quad 0 \leq t \leq T. \quad (4.2)$$

Note that, under Assumption 4.1, the functions $\beta, \lambda, \dot{\lambda}, \dot{\gamma}$ are all bounded on $[0, T]$. Moreover, the function η is càdlàg and of finite variation on $[0, T]$ (in particular, it is bounded), and $\theta, \dot{\theta}$ are bounded on $[0, T]$.

Let $\mathcal{S}^2$ denote the set of càdlàg semimartingales S with $\mathbb{E}[\sup_{0 \leq t \leq T} |S_t|^2] < \infty$ and let $\Xi_T \in L^2(\mathcal{F}_T)$ be the given target position at the terminal time T . The set of admissible execution strategies is given by

$$\mathcal{Q} := \{Q \in \mathcal{S}^2 : Q_{0-} = 0, Q_T = \Xi_T\}$$

with Q_t representing the cumulative number of shares bought or sold by time t . For any strategy $Q \in \mathcal{Q}$, we define the impact process $Y = Y^Q$ as the unique solution of

$$dY_t = -\beta_t Y_t dt + \lambda_t dQ_t, \quad Y_{0-} = y, \quad (4.3)$$

where $y \in \mathbb{R}$ is the given initial impact and is fixed throughout. The impact process Y is interpreted as the deviation process, driven by Q , from an unaffected price process $S \in \mathcal{S}^2$, in the sense that marginal orders are effectively executed at the price $S_- + Y_-$. The two illiquidity parameters in the model, the resilience β and the depth (or push factor) λ , are allowed to vary deterministically over time, similar to [15].

Given the unaffected asset price process S , we are interested in minimizing the execution cost,

$$\inf_{Q \in \mathcal{Q}} \mathbb{E} \left[\int_0^T Y_{t-} dQ_t + \frac{1}{2} [Y, Q]_T - \int_0^T Q_{t-} dS_t \right], \quad (4.4)$$

subject to the terminal constraint $Q_T = \Xi_T$ specifying the total quantity to be executed. Throughout, note that the integral $\int_a^b := \int_{[a,b]}$ is defined over the closed interval, and the quadratic covariation includes jumps at zero. Assuming that S is a sufficiently regular martingale such that the last integral in (4.4) is a martingale, the optimization reduces to

$$\inf_{Q \in \mathcal{Q}} \mathbb{E} \left[\int_0^T Y_{t-} dQ_t + \frac{1}{2} [Y, Q]_T \right]. \quad (4.5)$$

This is the form of the problem typically used in the literature, and we take this formulation as our primitive. Observe that, despite the absence of S , the problem is inherently stochastic due to the random terminal condition Ξ_T . This differs from standard optimal execution problems, where the order size is known upfront. The random target position can be thought of as arising from a stochastic order flow, which is anticipated and executed over time, akin to the setup studied in Nutz, Webster, and Zhao [19]. Its presence also necessitates the consideration of general càdlàg semimartingale execution strategies $Q \in \mathcal{Q}$. A detailed discussion and derivation of the cost functionals in (4.4) and (4.5) can be found in [1].

The next theorem summarizes the optimal execution strategy for (4.5).

Theorem 4.2. *Suppose Assumption 4.1 is satisfied. Let $\Xi_t := \mathbb{E}[\Xi_T \mid \mathcal{F}_t]$ for $t \in \{0-\} \cup [0, T]$, and define the square-integrable martingale $M = (M_t)_{t \in \{0-\} \cup [0, T]}$ by*

$$M_{0-} := \frac{\lambda_T}{1 + \lambda_T \theta_T} \left(\Xi_{0-} + \frac{y}{\lambda_0} \right), \quad (4.6)$$

$$M_t := M_{0-} + \int_0^t \frac{\lambda_T}{1 + \lambda_T (\theta_T - \theta_s)} d\Xi_s, \quad t \in [0, T]. \quad (4.7)$$

Equivalently, for $t \in [0, T]$,

$$M_t = \frac{\lambda_T}{1 + \lambda_T \theta_T} \left(\Xi_0 + \frac{y}{\lambda_0} \right) + \int_{(0, t]} \frac{\lambda_T}{1 + \lambda_T (\theta_T - \theta_s)} d\Xi_s. \quad (4.8)$$

Then

$$Y_t^* := \eta_t M_t \quad \text{for } t \in [0, T), \quad Y_T^* := M_T \quad (4.9)$$

is the price impact process of the unique optimal strategy for (4.5). The optimal strategy is the càdlàg semimartingale $Q^* \in \mathcal{Q}$ satisfying $Q_{0-}^* = 0$, $Q_T^* = \Xi_T$, and

$$\begin{aligned} Q_t^* &= \left(\Xi_{0-} + \frac{y}{\lambda_0} \right) \left[\frac{\lambda_T}{1 + \lambda_T \theta_T} \left(\frac{\eta_t}{\lambda_t} + \theta_t \right) - 1 \right] + \Xi_{0-} \\ &\quad + \int_0^t \frac{\lambda_T}{1 + \lambda_T (\theta_T - \theta_s)} \left(\frac{\eta_t}{\lambda_t} + \theta_t - \theta_s \right) d\Xi_s, \quad 0 \leq t < T. \end{aligned} \quad (4.10)$$

Its initial and terminal block trades are given by

$$\begin{aligned} \Delta Q_0^* &= \frac{\eta_0}{\lambda_0} \frac{\lambda_T}{1 + \lambda_T \theta_T} \left(\Xi_0 + \frac{y}{\lambda_0} \right) - \frac{y}{\lambda_0}, \\ \Delta Q_T^* &= \frac{1}{\lambda_T} \left(\frac{\beta_{T-}}{2\beta_{T-} + \dot{\gamma}_{T-}} M_T + \eta_{T-} \Delta M_T \right). \end{aligned} \quad (4.11)$$

Finally, the optimal execution cost (4.5) is given by

$$\frac{1}{2} \left(\mathbb{E} \left[\int_0^T \dot{\theta}_t M_t^2 dt \right] + \frac{\mathbb{E}[M_T^2]}{\lambda_T} - \frac{y^2}{\lambda_0} \right). \quad (4.12)$$

Remark 4.3 (Special cases). Our Theorem 4.2 generalizes some existing results in the literature. In the case where the terminal position $\Xi_T = x > 0$ is deterministic, equations (4.10)–(4.12) retrieve the optimal strategy and cost derived in Fruth et al. [15, Theorem 8.4]. In the case where $\beta > 0$ and $\lambda > 0$ are constants, our formulas (4.10)–(4.11) reduce to the ones derived in Ackermann et al. [3, Example 5.3] for the special case of a Brownian filtration setting. In particular, this includes the original formulas of Obizhaeva and Wang [20, Proposition 2] with deterministic target position and $y = 0$.

Remark 4.4. The optimal solution in Theorem 4.2 has some intuitive properties. First, note that the optimal strategy Q^* is characterized by making the controlled deviation process Y^* in (4.9) a martingale on $(0, T)$ up to scaling by the deterministic function η , which is in turn determined by the time-varying illiquidity parameters β and λ . This generalizes an observation from [20] for the deterministic Obizhaeva–Wang model with constant β, λ . There, the optimal strategy keeps the deviation process flat over $(0, T)$ by offsetting incurred price impact λ against resilience β . Here, loosely speaking, any execution strategy $Q \in \mathcal{Q}$ that induces a drift in the rescaled deviation process Y^Q/η in (4.3) cannot be optimal. It can similarly be interpreted as optimally counterbalancing, now in expectation, the illiquidity effects while trading toward the random target Ξ_T . Second, as also observed in [3, Example 5.3], the optimal execution strategy on $(0, T)$ in (4.10) decomposes into two additive parts: (i) the optimal deterministic (not necessarily absolutely continuous) execution strategy for

known order sizes, where the unknown terminal position Ξ_T is simply replaced by its expected value $\Xi_{0-} = \mathbb{E}[\Xi_T \mid \mathcal{F}_{0-}]$; (ii) random fluctuations around the deterministic strategy, driven by the càdlàg terminal-target martingale Ξ , which incorporates updates about the terminal position Ξ_T . In particular, note that jumps in the martingale Ξ during the execution interval $(0, T)$ trigger block trades in the execution strategy Q^* .

4.1 A Hilbert-space proof of Theorem 4.2

The rest of the section is devoted to proving Theorem 4.2. As observed in [15, Lemma 8.6], the equation for the transient price impact (4.3) can be inverted pathwise. More precisely, fixing $Q_{0-} = 0$ and $Y_{0-} = y$, one has

$$dQ_t = \frac{\beta_t}{\lambda_t} Y_t dt + \frac{1}{\lambda_t} dY_t, \quad Q_{0-} = 0, \quad (4.13)$$

and hence, using integration by parts, the representation

$$Q_t = \frac{Y_t}{\lambda_t} - \frac{y}{\lambda_0} + \int_0^t \frac{\beta_s + \dot{\gamma}_s}{\lambda_s} Y_s ds, \quad 0 \leq t \leq T. \quad (4.14)$$

It will be beneficial to cast our control problem (4.5) purely in terms of the impact process Y .

Lemma 4.5. *Let $Q \in \mathcal{Q}$ and let Y be the associated impact process. Then*

$$\begin{aligned} \mathbb{E} \left[\int_0^T Y_{t-} dQ_t + \frac{1}{2} [Y, Q]_T \right] &= \mathbb{E} \left[\frac{1}{2} \left(\frac{Y_T^2}{\lambda_T} - \frac{y^2}{\lambda_0} \right) + \int_0^T \frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} Y_t^2 dt \right] \\ &= \langle Y, Y \rangle_{\mathcal{H}} - \frac{y^2}{2\lambda_0}, \end{aligned} \quad (4.15)$$

where

$$\langle Y, Z \rangle_{\mathcal{H}} := \mathbb{E} \left[\int_0^T \frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} Y_t Z_t dt + \frac{1}{2\lambda_T} Y_T Z_T \right]. \quad (4.16)$$

Moreover, the terminal constraint $Q_T = \Xi_T$ is equivalent to

$$\frac{Y_T}{\lambda_T} - \frac{y}{\lambda_0} + \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} Y_t dt = \Xi_T. \quad (4.17)$$

Proof. Using (4.13), the identity $Y_{t-} = Y_t$ dt -a.e., Itô's formula, and integration by parts yields

$$\begin{aligned} \int_0^T Y_{t-} dQ_t + \frac{1}{2} [Y, Q]_T &= \int_0^T Y_{t-} \left(\frac{1}{\lambda_t} dY_t + \frac{\beta_t}{\lambda_t} Y_t dt \right) + \frac{1}{2} \int_0^T \frac{1}{\lambda_t} d[Y, Y]_t \\ &= \int_0^T \frac{1}{\lambda_t} \left(Y_{t-} dY_t + \frac{1}{2} d[Y, Y]_t \right) + \int_0^T \frac{\beta_t}{\lambda_t} Y_t^2 dt \\ &= \frac{1}{2} \int_0^T \frac{1}{\lambda_t} d(Y_t^2) + \int_0^T \frac{\beta_t}{\lambda_t} Y_t^2 dt \\ &= \frac{1}{2} \left(\frac{Y_T^2}{\lambda_T} - \frac{y^2}{\lambda_0} - \int_0^T Y_t^2 d\left(\frac{1}{\lambda_t}\right) \right) + \int_0^T \frac{\beta_t}{\lambda_t} Y_t^2 dt \\ &= \frac{1}{2} \left(\frac{Y_T^2}{\lambda_T} - \frac{y^2}{\lambda_0} \right) + \int_0^T \frac{\dot{\lambda}_t}{2\lambda_t^2} Y_t^2 dt + \int_0^T \frac{\beta_t}{\lambda_t} Y_t^2 dt, \end{aligned}$$

which is (4.15). The terminal constraint is (4.14) at time T . $\square$

Let $\mathcal{R}$ be the progressive σ -field on $\Omega \times [0, T]$ and define the finite measures

$$\mu(dt) := \frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} dt + \frac{1}{2\lambda_T} \delta_T(dt), \quad dR(\omega, t) := \mathbb{P}(d\omega) \mu(dt).$$

Then

$$\mathcal{H} := L^2(\Omega \times [0, T], \mathcal{R}, R)$$

is the Hilbert space of all progressively measurable processes $Y = (Y_t)_{t \in [0, T]}$ with finite norm

$$\|Y\|_{\mathcal{H}} := \sqrt{\langle Y, Y \rangle_{\mathcal{H}}} \quad (4.18)$$

induced by the inner product defined in (4.16). We also introduce the map $G : \mathcal{H} \rightarrow L^2(\mathcal{F}_T)$ defined as

$$G(Y) := \frac{Y_T}{\lambda_T} + \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} Y_t dt, \quad Y \in \mathcal{H}.$$

The next result shows that our control problem in (4.5) can be seen as a simple projection in the Hilbert space $\mathcal{H}$ onto a suitable subspace, which is determined by the terminal constraint through the map G .

Proposition 4.6. *The map $G : \mathcal{H} \rightarrow L^2(\mathcal{F}_T)$ is linear, continuous, and*

$$\mathcal{Y} := \{Y \in \mathcal{H} : G(Y) = \Xi_T + \frac{y}{\lambda_0}\}$$

is a nonempty closed affine subspace of $\mathcal{H}$. Consequently, the problem

$$\inf_{Y \in \mathcal{Y}} \langle Y, Y \rangle_{\mathcal{H}} \quad (4.19)$$

admits a unique minimizer $Y^ \in \mathcal{Y}$. Writing*

$$\mathcal{J} := \ker G = \left\{ J \in \mathcal{H} : \frac{J_T}{\lambda_T} = - \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} J_t dt \right\},$$

this minimizer is characterized by the first-order condition

$$\langle Y^*, J \rangle_{\mathcal{H}} = 0, \quad J \in \mathcal{J}. \quad (4.20)$$

Proof. Clearly, the map G is linear. For $Y \in \mathcal{H}$, Cauchy–Schwarz gives

$$\begin{aligned} \mathbb{E} \left| \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} Y_t dt \right|^2 &= \mathbb{E} \left| \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\sqrt{\lambda_t} \sqrt{2\beta_t + \dot{\gamma}_t}} \frac{\sqrt{2\beta_t + \dot{\gamma}_t}}{\sqrt{\lambda_t}} Y_t dt \right|^2 \\ &\leq 2\theta_T \mathbb{E} \left[\int_0^T \frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} Y_t^2 dt \right] \leq 2\theta_T \|Y\|_{\mathcal{H}}^2. \end{aligned}$$

Moreover,

$$\mathbb{E} \left[\left| \frac{Y_T}{\lambda_T} \right|^2 \right] \leq \frac{2}{\lambda_T} \|Y\|_{\mathcal{H}}^2.$$

Hence, $\|G(Y)\|_{L^2(\mathcal{F}_T)} \leq c\|Y\|_{\mathcal{H}}$ for some constant $c > 0$, and G is continuous. Therefore, $\mathcal{Y} = G^{-1}(\Xi_T + y/\lambda_0)$ is a closed affine subspace of $\mathcal{H}$. It is nonempty because the element

$$Y_t^\circ := \mathbf{1}_{\{T\}}(t)\lambda_T \left(\Xi_T + \frac{y}{\lambda_0} \right), \quad 0 \leq t \leq T,$$

belongs to $\mathcal{Y}$. Consequently, the projection theorem in Hilbert spaces yields the existence of a unique minimizer $Y^* \in \mathcal{Y}$, and the usual characterization of projections onto affine subspaces gives the condition in (4.20). $\square$

The next step is to derive an explicit formula for Y^* from Proposition 4.6.

Proposition 4.7. *Suppose Assumption 4.1 is satisfied. Let M be given by (4.6)–(4.7). Then M is a square-integrable càdlàg martingale, and the unique minimizer from Proposition 4.6 is given by the square-integrable càdlàg semimartingale*

$$Y_t^* = \eta_t M_t, \quad t \in [0, T], \quad Y_T^* = M_T. \quad (4.21)$$

Proof. First, observe that (4.7) defines a square-integrable martingale on $[0, T]$. Indeed, the deterministic integrand

$$\frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_t)}, \quad 0 \leq t \leq T,$$

is bounded since $0 \leq \theta_T - \theta_t \leq \theta_T$ by the definition of θ in (4.2). Moreover, since η is deterministic, bounded, càdlàg, and of finite variation by Assumption 4.1, the process Y^* defined in (4.21) is càdlàg and a square-integrable semimartingale.

Next, we argue that Y^* is an element of $\mathcal{Y}$. First, since

$$\frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} (Y_t^*)^2 = \frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} (\eta_t M_t)^2 = \frac{1}{2} \dot{\theta}_t M_t^2 \quad \text{a.e.,}$$

we obtain that

$$\|Y^*\|_{\mathcal{H}}^2 = \frac{1}{2} \mathbb{E} \left[\int_0^T \dot{\theta}_t M_t^2 dt \right] + \frac{1}{2\lambda_T} \mathbb{E}[M_T^2] < \infty,$$

because $\dot{\theta}$ is bounded by Assumption 4.1, and hence $Y^* \in \mathcal{H}$.

Regarding the condition $G(Y^*) = \Xi_T + y/\lambda_0$, we introduce the function

$$a_t := \frac{1 + \lambda_T(\theta_T - \theta_t)}{\lambda_T} = \frac{1}{\lambda_T} + \theta_T - \theta_t, \quad 0 \leq t \leq T.$$

Clearly, a is continuous and of finite variation with $da_t = -\dot{\theta}_t dt$, and (4.7) is equivalent to

$$a_t dM_t = d\Xi_t.$$

Thus, integration by parts yields

$$d(a_t M_t) = a_t dM_t + M_t da_t = d\Xi_t - \dot{\theta}_t M_t dt. \quad (4.22)$$

Due to the definition of M_{0-} in (4.6), we have $a_0 M_{0-} = \Xi_{0-} + y/\lambda_0$, and hence, by integrating (4.22), we obtain for every $t \in [0, T]$ the representation

$$\Xi_t = \left(\frac{1}{\lambda_T} + \theta_T - \theta_t \right) M_t - \frac{y}{\lambda_0} + \int_0^t \dot{\theta}_s M_s ds. \quad (4.23)$$

In particular, at $t = T$, it holds that

$$\Xi_T = \frac{M_T}{\lambda_T} - \frac{y}{\lambda_0} + \int_0^T \dot{\theta}_s M_s ds, \quad (4.24)$$

which implies

$$\begin{aligned} G(Y^*) &= \frac{Y_T^*}{\lambda_T} + \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} Y_t^* dt = \frac{M_T}{\lambda_T} + \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} \eta_t M_t dt \\ &= \frac{M_T}{\lambda_T} + \int_0^T \dot{\theta}_s M_s ds = \Xi_T + \frac{y}{\lambda_0}, \end{aligned} \quad (4.25)$$

so $Y^* \in \mathcal{Y}$.

It remains to verify the first-order condition in (4.20). To this end, let $J \in \mathcal{J}$. By the representation of Y^* in (4.21), we get

$$\langle Y^*, J \rangle_{\mathcal{H}} = \frac{1}{2} \mathbb{E} \left[\int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} M_t J_t dt + \frac{M_T J_T}{\lambda_T} \right]. \quad (4.26)$$

Since $J \in \ker G$, it satisfies

$$\frac{J_T}{\lambda_T} = - \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} J_t dt.$$

Plugging this back into (4.26) yields

$$\begin{aligned} \langle Y^*, J \rangle_{\mathcal{H}} &= \frac{1}{2} \mathbb{E} \left[\int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} M_t J_t dt - M_T \int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} J_t dt \right] \\ &= \frac{1}{2} \mathbb{E} \left[\int_0^T \frac{\beta_t + \dot{\gamma}_t}{\lambda_t} (M_t - \mathbb{E}[M_T | \mathcal{F}_t]) J_t dt \right] = 0, \end{aligned}$$

because M is a martingale. Thus (4.20) holds, so Proposition 4.6 identifies Y^* in (4.21) as the unique minimizer of (4.19). $\square$

We are now ready to prove Theorem 4.2.

Proof of Theorem 4.2. By Proposition 4.7, the process $Y^* \in \mathcal{Y}$ in (4.21) represents the unique minimizer of (4.19). Moreover, formula (4.14) defines the associated càdlàg semimartingale inventory process Q^* given by

$$Q_t^* = \frac{Y_t^*}{\lambda_t} - \frac{y}{\lambda_0} + \int_0^t \frac{\beta_s + \dot{\gamma}_s}{\lambda_s} Y_s^* ds = \begin{cases} \frac{\eta_t M_t}{\lambda_t} - \frac{y}{\lambda_0} + \int_0^t \dot{\theta}_s M_s ds, & 0 \leq t < T, \\ \frac{M_T}{\lambda_T} - \frac{y}{\lambda_0} + \int_0^T \dot{\theta}_s M_s ds, & t = T, \end{cases} \quad (4.27)$$

with $Q_{0-}^* = 0$. Observe that $Q^* \in \mathcal{S}^2$ by Proposition 4.7. Furthermore, since $Y^* \in \mathcal{Y}$, the condition in (4.25) yields $Q_T^* = \Xi_T$. Thus $Q^* \in \mathcal{Q}$.

Let $Q \in \mathcal{Q}$ be arbitrary and let Y be its impact process. By Lemma 4.5,

$$\mathbb{E} \left[\int_0^T Y_{t-} dQ_t + \frac{1}{2} [Y, Q]_T \right] = \langle Y, Y \rangle_{\mathcal{H}} - \frac{y^2}{2\lambda_0} \geq \langle Y^*, Y^* \rangle_{\mathcal{H}} - \frac{y^2}{2\lambda_0},$$

because Y^* minimizes $\langle Y, Y \rangle_{\mathcal{H}}$ over $\mathcal{Y}$. Hence Q^* is optimal.

To prove uniqueness, let $\tilde{Q} \in \mathcal{Q}$ be another optimizer and $\tilde{Y}$ its impact process. Then $\tilde{Y}$ is another minimizer in $\mathcal{Y}$, so $\tilde{Y} = Y^*$ in $\mathcal{H}$. In particular, $\tilde{Y}_T = Y_T^*$ a.s. and $\tilde{Y}_t = Y_t^*$ for $\mathbb{P} \otimes dt$ -a.e. $(\omega, t) \in \Omega \times [0, T]$. Since both processes are càdlàg, they are indistinguishable. Then (4.14) implies that $\tilde{Q}$ and Q^* are also indistinguishable.

To justify the representation of the optimal inventory Q^* in (4.10), we first insert (4.7) into (4.27) to obtain

$$\begin{aligned} Q_t^* &= \frac{\eta_t}{\lambda_t} \frac{\lambda_T}{1 + \lambda_T \theta_T} \left(\Xi_{0-} + \frac{y}{\lambda_0} \right) + \frac{\eta_t}{\lambda_t} \int_0^t \frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_s)} d\Xi_s \\ &\quad - \frac{y}{\lambda_0} + \int_0^t \dot{\theta}_s \frac{\lambda_T}{1 + \lambda_T \theta_T} \left(\Xi_{0-} + \frac{y}{\lambda_0} \right) ds + \int_0^t \dot{\theta}_s \left(\int_0^s \frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_r)} d\Xi_r \right) ds \\ &= \left(\Xi_{0-} + \frac{y}{\lambda_0} \right) \left(\frac{\eta_t}{\lambda_t} \frac{\lambda_T}{1 + \lambda_T \theta_T} + \frac{\lambda_T}{1 + \lambda_T \theta_T} \theta_t - 1 \right) + \Xi_{0-} \\ &\quad + \frac{\eta_t}{\lambda_t} \int_0^t \frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_s)} d\Xi_s + \int_0^t \left(\int_0^s \frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_r)} d\Xi_r \right) d\theta_s, \quad 0 \leq t < T. \end{aligned}$$

Performing integration by parts in the last integral (note that θ is absolutely continuous) yields

$$\int_0^t \left(\int_0^s \frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_r)} d\Xi_r \right) d\theta_s = \theta_t \int_0^t \frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_s)} d\Xi_s - \int_0^t \theta_s \frac{\lambda_T}{1 + \lambda_T(\theta_T - \theta_s)} d\Xi_s.$$

Inserting this back into the previous equation results in (4.10).

For the initial and final block trades in (4.11), we note that

$$\Delta Q_0^* = \frac{1}{\lambda_0} \Delta Y_0^* = \frac{1}{\lambda_0} (Y_0^* - Y_{0-}^*) = \frac{1}{\lambda_0} (\eta_0 M_0 - y)$$

and

$$\Delta Q_T^* = \frac{1}{\lambda_T} \Delta Y_T^* = \frac{1}{\lambda_T} (Y_T^* - Y_{T-}^*) = \frac{1}{\lambda_T} (M_T - \eta_{T-} M_{T-}) = \frac{1}{\lambda_T} (M_T(1 - \eta_{T-}) + \eta_{T-} \Delta M_T).$$

Finally, for the optimal cost in (4.12), we use Lemma 4.5 and obtain

$$\begin{aligned} \mathbb{E} \left[\int_0^T Y_{t-}^* dQ_t^* + \frac{1}{2} [Y^*, Q^*]_T \right] &= \langle Y^*, Y^* \rangle_{\mathcal{H}} - \frac{y^2}{2\lambda_0} \\ &= \mathbb{E} \left[\int_0^T \frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} (Y_t^*)^2 dt + \frac{1}{2\lambda_T} (Y_T^*)^2 \right] - \frac{y^2}{2\lambda_0} \\ &= \frac{1}{2} \mathbb{E} \left[\int_0^T \dot{\theta}_t M_t^2 dt \right] + \frac{1}{2\lambda_T} \mathbb{E} [M_T^2] - \frac{y^2}{2\lambda_0}, \end{aligned}$$

which completes the proof. $\square$

5 Regularized execution and its approximation rate

As seen in Theorem 4.2, the optimal strategy for the unregularized Obizhaeva–Wang model naturally has jumps. While block trades at the initial and terminal times arise even in the deterministic case, the terminal-target martingale $(\Xi_t)_{0 \leq t \leq T}$ in our setting typically also leads to non-zero quadratic variation during the execution interval $(0, T)$. A common approach to smooth the strategy is to use a regularized version of the model where an instantaneous quadratic cost penalizes fast trading and in particular imposes absolute continuity of the execution strategy. If we consider the instantaneous cost as an auxiliary regularizer to produce better-behaved strategies, this regularity comes at a price, namely, additional execution cost.

While the standard approach would be to use the regularized optimal strategy as execution strategy, that optimizer is not explicitly available in our setting with stochastic terminal condition and time-varying coefficients. Below, we construct a *simple, explicit, nearly optimal* strategy Q^ε which achieves a similar cost as the optimizer. Specifically, our main result in Theorem 5.7 gives a non-asymptotic bound on the additional execution cost, which turns out to be of order $\sqrt{\varepsilon}$ for both strategies. We further show in Proposition 5.9 that this order is sharp even in the benchmark deterministic case.

Let J_0 denote the objective of the unregularized problem in (4.5) from Section 4, which can be rewritten as

$$J_0(Q) := \langle Y^Q, Y^Q \rangle_{\mathcal{H}} - \frac{y^2}{2\lambda_0}$$

according to Lemma 4.5. We also introduce the corresponding value

$$V(0) := \inf_{Q \in \mathcal{Q}} J_0(Q) = J_0(Q^0) \quad (5.1)$$

where $Q^0 := Q^* \in \mathcal{Q}$ denotes the unique optimizer from Theorem 4.2 with price impact process $Y^0 := Y^{Q^0}$.

Next, we introduce the regularized problem. Let $\varepsilon > 0$ and let $\mathcal{Q}_{\text{ac}} \subset \mathcal{Q}$ denote the subclass of absolutely continuous strategies Q of the form

$$Q_t = \int_0^t u_s ds, \quad 0 \leq t \leq T,$$

with progressively measurable u satisfying $\mathbb{E} \int_0^T u_t^2 dt < \infty$, and $Q_T = \Xi_T \in L^2(\mathcal{F}_T)$. For any $Q \in \mathcal{Q}_{\text{ac}}$ with impact process Y^Q given by (4.3), we introduce the objective J_ε by adding an instantaneous quadratic cost,

$$J_\varepsilon(Q) := J_0(Q) + \varepsilon \mathbb{E} \int_0^T \dot{Q}_t^2 dt. \quad (5.2)$$

We further denote the corresponding value by

$$V(\varepsilon) := \inf_{Q \in \mathcal{Q}_{\text{ac}}} J_\varepsilon(Q). \quad (5.3)$$

Throughout this section, Assumption 4.1 is in force. Moreover, for the square-integrable terminal-target martingale $\Xi_t = \mathbb{E}[\Xi_T \mid \mathcal{F}_t]$, $t \in [0, T]$, we assume that the reachability condition in (3.1) from Section 3 is satisfied and denote its value by

$$\Lambda_\Xi := \int_0^T \frac{d\mathbb{E}[\Xi_t^2]}{T-t} < \infty. \quad (5.4)$$

Remark 5.1. Under (5.4), for every $\varepsilon > 0$, the infimum in (5.3) is attained by a unique strategy $Q^{*,\varepsilon} \in \mathcal{Q}_{ac}$. Indeed, identifying an absolutely continuous strategy with its trading rate, the feasible set is a nonempty closed affine subset of $L^2(\Omega \times [0, T], \mathcal{R}, \mathbb{P} \otimes dt)$, while J_ε is weakly lower semicontinuous, coercive, and strictly convex.

We equip square-integrable càdlàg processes on $[0, T]$ with the norm

$$\|X\|_*^2 := \mathbb{E} \left[\int_0^T X_t^2 dt + X_T^2 \right] \quad (5.5)$$

and set

$$C_{\mathcal{H}}^2 := \frac{1}{2} \max \left\{ \left\| \frac{2\beta + \dot{\gamma}}{\lambda} \right\|_{L^\infty(0,T)}, \frac{1}{\lambda_T} \right\}, \quad (5.6)$$

so that

$$\|X\|_{\mathcal{H}} \leq C_{\mathcal{H}} \|X\|_* \quad (5.7)$$

for every square-integrable càdlàg process X , where we recall that $\|\cdot\|_{\mathcal{H}}$ is the norm in (4.18).

We start by recording the following simple Lipschitz property.

Lemma 5.2. *Let $Q, \tilde{Q} \in \mathcal{Q}$, and let $Y^Q, Y^{\tilde{Q}}$ be the associated impact processes following (4.3). Then*

$$\|Y^Q - Y^{\tilde{Q}}\|_* \leq L_Y \|Q - \tilde{Q}\|_*, \quad (5.8)$$

where

$$L_Y := \|\lambda\|_{L^\infty(0,T)} \sqrt{2 + T(T+2)\|\beta + \dot{\gamma}\|_{L^\infty(0,T)}^2}. \quad (5.9)$$

In particular, the map $Q \mapsto Y^Q$ is affine and Lipschitz on $(\mathcal{Q}, \|\cdot\|_*)$.

Proof. Let

$$R_t := Q_t - \tilde{Q}_t, \quad Z_t := Y_t^Q - Y_t^{\tilde{Q}}, \quad B_t := \int_0^t \beta_s ds, \quad 0 \leq t \leq T,$$

and introduce the constants

$$L_\lambda := \|\lambda\|_{L^\infty(0,T)}, \quad L_b := \|\beta + \dot{\gamma}\|_{L^\infty(0,T)}.$$

Then $R_{0-} = 0$, $Z_{0-} = 0$, and $Z = (Z_t)_{0 \leq t \leq T}$ satisfies

$$dZ_t = -\beta_t Z_t dt + \lambda_t dR_t, \quad 0 \leq t \leq T,$$

with explicit solution

$$Z_t = e^{-B_t} \int_0^t e^{B_s} \lambda_s dR_s \quad 0 \leq t \leq T.$$

Since $e^B \lambda$ is continuous and of finite variation, integration by parts yields

$$Z_t = \lambda_t R_t - \int_0^t e^{-(B_t - B_s)} \lambda_s (\beta_s + \dot{\gamma}_s) R_s ds.$$

As $\beta \geq 0$, the exponential factor is bounded by one, and we get the upper bound

$$|Z_t|^2 \leq 2L_\lambda^2 |R_t|^2 + 2L_\lambda^2 L_b^2 \left(\int_0^t |R_s| ds \right)^2.$$

Next, using Cauchy–Schwarz and Fubini yields

$$\begin{aligned} \int_0^T \left(\int_0^t |R_s| ds \right)^2 dt &\leq \int_0^T t \int_0^t R_s^2 ds dt \\ &= \frac{1}{2} \int_0^T (T^2 - s^2) R_s^2 ds \leq \frac{T^2}{2} \int_0^T R_s^2 ds. \end{aligned}$$

Therefore

$$\mathbb{E} \int_0^T Z_t^2 dt \leq L_\lambda^2 (2 + T^2 L_b^2) \mathbb{E} \int_0^T R_t^2 dt.$$

Likewise,

$$|Z_T|^2 \leq 2L_\lambda^2 |R_T|^2 + 2L_\lambda^2 L_b^2 T \int_0^T R_s^2 ds.$$

Adding the two bounds yields (5.8). $\square$

The next lemma shows that the difference $V(\varepsilon) - V(0)$ can be controlled by the value of the constrained tracking problem of the optimal unregularized inventory $Q^0 \in \mathcal{Q}$ by an absolutely continuous strategy $Q \in \mathcal{Q}_{ac}$ satisfying $Q_T = \Xi_T$.

Lemma 5.3. *Let*

$$C_J := C_{\mathcal{H}}^2 L_Y^2. \quad (5.10)$$

Then, for every $Q \in \mathcal{Q}$,

$$0 \leq J_0(Q) - J_0(Q^0) = \|Y^Q - Y^0\|_{\mathcal{H}}^2 \leq C_J \mathbb{E} \int_0^T (Q_t - Q_t^0)^2 dt. \quad (5.11)$$

Consequently, for every $Q \in \mathcal{Q}_{ac}$ and every $\varepsilon > 0$,

$$0 \leq V(\varepsilon) - V(0) \leq C_J \mathbb{E} \int_0^T (Q_t - Q_t^0)^2 dt + \varepsilon \mathbb{E} \int_0^T \dot{Q}_t^2 dt. \quad (5.12)$$

Proof. Let $Q \in \mathcal{Q}$. Since $Q_T = \Xi_T = Q_T^0$, the constraint identity (4.17) applied to Y^Q and Y^0 shows that

$$G(Y^Q - Y^0) = 0,$$

that is, $Y^Q - Y^0 \in \mathcal{J}$. Therefore, the first-order condition (4.20) gives

$$\langle Y^0, Y^Q - Y^0 \rangle_{\mathcal{H}} = 0.$$

It follows that

$$J_0(Q) - J_0(Q^0) = \|Y^Q\|_{\mathcal{H}}^2 - \|Y^0\|_{\mathcal{H}}^2 = \|Y^Q - Y^0\|_{\mathcal{H}}^2.$$

Moreover, using (5.7), Lemma 5.2, and again $Q_T = Q_T^0$, we obtain

$$\|Y^Q - Y^0\|_{\mathcal{H}}^2 \leq C_{\mathcal{H}}^2 \|Y^Q - Y^0\|_*^2 \leq C_J \|Q - Q^0\|_*^2 = C_J \mathbb{E} \int_0^T (Q_t - Q_t^0)^2 dt,$$

which proves (5.11) for every $Q \in \mathcal{Q}$.

Finally, if $Q \in \mathcal{Q}_{\text{ac}}$, then

$$V(\varepsilon) - V(0) \leq J_\varepsilon(Q) - J_0(Q^0) = J_0(Q) - J_0(Q^0) + \varepsilon \mathbb{E} \int_0^T \dot{Q}_t^2 dt.$$

Combining this with (5.11) yields (5.12). $\square$

We now collect some properties of the unregularized optimal strategy Q^0 from Theorem 4.2 to verify that it fits our framework developed in Section 3. Recall the definition of the processes $\eta = (\eta_t)_{0 \leq t \leq T}$ and $\theta = (\theta_t)_{0 \leq t \leq T}$ in (4.2), as well as the L^2 time-translation modulus and associated seminorm from (2.6) and (2.7).

Lemma 5.4. *Define*

$$\alpha_t := \frac{\eta_t}{\lambda_t}, \quad t \in [0, T].$$

Then α has finite variation on $[0, T)$, and we set

$$A_\alpha := \|\alpha\|_{L^\infty(0, T)}, \quad V_\alpha := |\alpha|_{T-}, \quad B_\theta := \|\dot{\theta}\|_{L^\infty(0, T)}, \quad m_2 := \mathbb{E}[M_T^2],$$

where $|\alpha|_{T-}$ denotes the total variation of α on $[0, T)$. Moreover, defining the constant

$$C_0 := 2A_\alpha^2 m_2 + \frac{2y^2}{\lambda_0^2},$$

we have

$$\mathbb{E}[(Q_0^0)^2] \leq C_0. \quad (5.13)$$

The L^2 time-translation modulus of Q^0 satisfies

$$\omega_{Q^0}(h) \leq 3m_2(A_\alpha^2 + 4V_\alpha^2 + T^2 B_\theta^2)h, \quad 0 < h \leq T, \quad (5.14)$$

and

$$[Q^0]_{\text{tr}}^2 \leq 3m_2(A_\alpha^2 + 4V_\alpha^2 + T^2 B_\theta^2). \quad (5.15)$$

Finally,

$$K_0 := \sup_{t \in [0, T)} \mathbb{E}|Q_t^0 - \Xi_t|^2 \leq 6m_2(A_\alpha^2 + T^2 B_\theta^2) + \frac{6y^2}{\lambda_0^2} + 2\mathbb{E}[\Xi_T^2]. \quad (5.16)$$

Proof. By Assumption 4.1, η has finite variation on $[0, T)$ and λ^{-1} is absolutely continuous. Hence, $\alpha = \eta/\lambda$ has finite variation on $[0, T)$ and we have $A_\alpha < \infty$, as well as $V_\alpha < \infty$. Moreover, $B_\theta < \infty$, again by Assumption 4.1, and $m_2 < \infty$ by Proposition 4.7.

Next, recall the representation of Q^0 from (4.27) given by

$$Q_t^0 = \alpha_t M_t - \frac{y}{\lambda_0} + \int_0^t \dot{\theta}_s M_s ds, \quad 0 \leq t < T. \quad (5.17)$$

At $t = 0$, we have

$$Q_0^0 = \alpha_0 M_0 - \frac{y}{\lambda_0},$$

and therefore

$$\mathbb{E}[(Q_0^0)^2] \leq 2A_\alpha^2 \mathbb{E}[M_0^2] + \frac{2y^2}{\lambda_0^2} \leq 2A_\alpha^2 m_2 + \frac{2y^2}{\lambda_0^2},$$

which proves (5.13).

Fix $0 < h \leq T$ and write $s := (t - h)^+$. For $t < T$, using (5.17) gives

$$Q_t^0 - Q_s^0 = \alpha_t(M_t - M_s) + (\alpha_t - \alpha_s)M_s + \int_s^t \dot{\theta}_r M_r dr.$$

Hence, the estimate $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ yields

$$\begin{aligned} \omega_{Q^0}(h) &= \mathbb{E} \int_0^T |Q_t^0 - Q_s^0|^2 dt \\ &\leq 3A_\alpha^2 \mathbb{E} \int_0^T |M_t - M_s|^2 dt + 3 \mathbb{E} \left[\sup_{u \in [0, T)} M_u^2 \right] \int_0^T |\alpha_t - \alpha_s|^2 dt \\ &\quad + 3B_\theta^2 \mathbb{E} \int_0^T \left(\int_s^t |M_r| dr \right)^2 dt. \end{aligned}$$

For the martingale term, due to the computations in (2.25), we have

$$\mathbb{E} \int_0^T |M_t - M_s|^2 dt \leq h \mathbb{E}[\langle M \rangle_T] \leq h m_2.$$

For the finite variation term, performing the same computations as in (2.26) gives

$$\int_0^T |\alpha_t - \alpha_s|^2 dt \leq h V_\alpha^2.$$

Moreover, using Doob's inequality, we have

$$\mathbb{E} \left[\sup_{u \in [0, T)} M_u^2 \right] \leq 4m_2.$$

Finally, Cauchy–Schwarz and Fubini imply, again similar to (2.26), the bound

$$\int_0^T \left(\int_s^t |M_r| dr \right)^2 dt \leq h \int_0^T \int_s^t M_r^2 dr dt \leq h^2 \int_0^T M_r^2 dr.$$

Taking expectations and using $h \leq T$ as well as $\mathbb{E}[M_r^2] \leq m_2$, we get

$$\mathbb{E} \int_0^T \left(\int_s^t |M_r| dr \right)^2 dt \leq T^2 h m_2.$$

Combining the three estimates proves (5.14). Since $\omega_{Q^0}(h) = \omega_{Q^0}(T)$ for every $h \geq T$, (5.15) follows immediately.

For the claim in (5.16), using once more the representation in (5.17) as well as $(a+b+c)^2 \leq 3(a^2 + b^2 + c^2)$ gives

$$|Q_t^0|^2 \leq 3A_\alpha^2 M_t^2 + \frac{3y^2}{\lambda_0^2} + 3B_\theta^2 T \int_0^T M_s^2 ds, \quad t < T.$$

Taking expectations and using $\mathbb{E}[M_t^2] \leq m_2$ yields

$$\sup_{t \in [0, T)} \mathbb{E}|Q_t^0|^2 \leq 3m_2(A_\alpha^2 + T^2 B_\theta^2) + \frac{3y^2}{\lambda_0^2}.$$

Lastly, since $\sup_{t \in [0, T]} \mathbb{E}[\Xi_t^2] \leq \mathbb{E}[\Xi_T^2]$, we conclude that

$$K_0 \leq 2 \sup_{t \in [0, T)} \mathbb{E}|Q_t^0|^2 + 2 \sup_{t \in [0, T]} \mathbb{E}[\Xi_t^2],$$

which is (5.16). $\square$

The next result exhibits an absolutely continuous execution strategy Q^ε which is based on the unregularized optimizer Q^0 , whose explicit form was derived in Theorem 4.2. Following the construction in the proof of Theorem 3.1, Q^ε tracks Q^0 through an exponential filter with relaxation scale $\sqrt{\varepsilon}$ and, over the final interval of that length, switches to a terminal bridge enforcing $Q_T^\varepsilon = \Xi_T$. The resulting feedback rule is explicit and does not require solving the regularized stochastic control problem.

Lemma 5.5 (Explicit execution strategy). *Fix $0 < \varepsilon \leq T^2/4$. Define*

$$\bar{Q}_t^\varepsilon := \frac{1}{\sqrt{\varepsilon}} \int_0^t e^{-(t-r)/\sqrt{\varepsilon}} Q_r^0 dr, \quad 0 \leq t < T, \quad (5.18)$$

and

$$Q_t^\varepsilon := \begin{cases} \bar{Q}_t^\varepsilon, & 0 \leq t \leq T - \sqrt{\varepsilon}, \\ \frac{T-t}{\sqrt{\varepsilon}} \bar{Q}_{T-\sqrt{\varepsilon}}^\varepsilon + (T-t) \int_{T-\sqrt{\varepsilon}}^t \frac{\Xi_r}{(T-r)^2} dr, & T - \sqrt{\varepsilon} < t < T, \\ \Xi_T, & t = T. \end{cases} \quad (5.19)$$

Then $Q^\varepsilon \in \mathcal{Q}_{\text{ac}}$. Its trading rate is given by the feedback rule

$$\dot{Q}_t^\varepsilon = \begin{cases} \frac{Q_t^0 - Q_t^\varepsilon}{\sqrt{\varepsilon}}, & 0 \leq t < T - \sqrt{\varepsilon}, \\ \frac{\Xi_t - Q_t^\varepsilon}{T-t}, & T - \sqrt{\varepsilon} \leq t < T, \end{cases} \quad \mathbb{P} \otimes dt\text{-a.e.} \quad (5.20)$$

Define

$$\tilde{C}_{\text{tr}} := 2(C_0 + 2[Q^0]_{\text{tr}}^2 + 8K_0 + 10\mathbb{E}[\Xi_T^2]) \quad (5.21)$$

and

$$C_{\text{tr}} := \tilde{C}_{\text{tr}} + 2T\Lambda_\Xi. \quad (5.22)$$

Then

$$\mathbb{E} \left[\int_0^T (Q_t^\varepsilon - Q_t^0)^2 dt + \varepsilon \int_0^T (\dot{Q}_t^\varepsilon)^2 dt \right] \leq \tilde{C}_{\text{tr}} \sqrt{\varepsilon} + 4\Lambda_\Xi \varepsilon \leq C_{\text{tr}} \sqrt{\varepsilon}. \quad (5.23)$$

Proof. For brevity, set $a := \sqrt{\varepsilon}$ and $s := T - \sqrt{\varepsilon}$. On $[0, s]$, differentiating (5.18) gives

$$\dot{Q}_t^\varepsilon = \frac{Q_t^0 - Q_t^\varepsilon}{a}.$$

Thus, the calculation in the proof of Theorem 2.1, applied with target Q^0 and initial position zero, together with (5.13) and (5.15), yields

$$\mathbb{E} \left[\frac{1}{2} \int_0^s (Q_t^\varepsilon - Q_t^0)^2 dt + \frac{a^2}{2} \int_0^s (\dot{Q}_t^\varepsilon)^2 dt \right] \leq a (C_0 + 2[Q^0]_{\text{tr}}^2). \quad (5.24)$$

Set

$$B_\varepsilon := \mathbb{E} [|\Xi_s - \bar{Q}_s^\varepsilon|^2].$$

By Jensen's inequality,

$$\mathbb{E} [|\bar{Q}_s^\varepsilon|^2] \leq \frac{1}{a} \int_0^s e^{-(s-r)/a} \mathbb{E} [|\bar{Q}_r^\varepsilon|^2] dr \leq \sup_{t \in [0, T]} \mathbb{E} [|\bar{Q}_t^\varepsilon|^2] \leq 2K_0 + 2\mathbb{E} [\Xi_T^2],$$

where the last inequality follows from $|\bar{Q}_t^\varepsilon|^2 \leq 2|Q_t^0 - \Xi_t|^2 + 2|\Xi_t|^2$. Consequently,

$$B_\varepsilon \leq 2\mathbb{E} [\Xi_s^2] + 2\mathbb{E} [|\bar{Q}_s^\varepsilon|^2] \leq 4K_0 + 6\mathbb{E} [\Xi_T^2]. \quad (5.25)$$

On $[s, T]$, the second branch of (5.19) solves the second feedback equation in (5.20). Integration by parts gives

$$\dot{Q}_t^\varepsilon = \frac{\Xi_s - \bar{Q}_s^\varepsilon}{a} + N_t, \quad N_t := \int_{(s, t]} \frac{1}{T-r} d\Xi_r, \quad s \leq t < T. \quad (5.26)$$

The process N is progressively measurable, and the martingale isometry and Fubini's theorem yield

$$\mathbb{E} \int_s^T N_t^2 dt = \int_{(s, T]} \frac{d\mathbb{E} [\Xi_r^2]}{T-r} \leq \Lambda_\Xi. \quad (5.27)$$

Together with (5.24) and (5.25), this shows that the trading rate in (5.20) is progressively measurable and square-integrable. Moreover, the argument following (3.14) shows that $Q_t^\varepsilon \rightarrow \Xi_T$ almost surely as $t \uparrow T$. Hence $Q^\varepsilon \in \mathcal{Q}_{\text{ac}}$.

For $s \leq t < T$, (5.20) and (5.26) imply

$$\mathbb{E} [|\Xi_t - Q_t^\varepsilon|^2] \leq 2 \frac{(T-t)^2}{a^2} B_\varepsilon + 2(T-t) \Lambda_\Xi.$$

Indeed, for $r \leq t$, $(T-t)^2/(T-r)^2 \leq (T-t)/(T-r)$, and the second term follows from the martingale isometry. Integrating over $[s, T]$ and using the definition of K_0 , we obtain

$$\frac{1}{2} \mathbb{E} \int_s^T (Q_t^\varepsilon - Q_t^0)^2 dt \leq \frac{2a}{3} B_\varepsilon + a^2 \Lambda_\Xi + aK_0. \quad (5.28)$$

Similarly, (5.26) and (5.27) give

$$\frac{a^2}{2} \mathbb{E} \int_s^T (\dot{Q}_t^\varepsilon)^2 dt \leq aB_\varepsilon + a^2 \Lambda_\Xi. \quad (5.29)$$

Combining (5.28) and (5.29) with (5.25) yields

$$\begin{aligned} \mathbb{E} \left[\frac{1}{2} \int_s^T (Q_t^\varepsilon - Q_t^0)^2 dt + \frac{a^2}{2} \int_s^T (\dot{Q}_t^\varepsilon)^2 dt \right] &\leq \frac{5a}{3} B_\varepsilon + aK_0 + 2a^2 \Lambda_\Xi \\ &\leq a(8K_0 + 10\mathbb{E}[\Xi_T^2]) + 2a^2 \Lambda_\Xi. \end{aligned}$$

Combining this estimate with (5.24) and multiplying by two gives

$$\mathbb{E} \left[\int_0^T (Q_t^\varepsilon - Q_t^0)^2 dt + a^2 \int_0^T (\dot{Q}_t^\varepsilon)^2 dt \right] \leq \tilde{C}_{\text{tr}} a + 4a^2 \Lambda_\Xi.$$

Since $a = \sqrt{\varepsilon}$ and $a \leq T/2$, this proves (5.23). The finiteness of the constants follows from Lemma 5.4 and (5.4). $\square$

Remark 5.6 (On Q^ε). The above construction of Q^ε deviates slightly from the proof of Theorem 3.1. There, under the general assumptions of the constrained tracking problem, the switching time is selected by an averaging argument from the interval $[T - 2\sqrt{\varepsilon}, T - \sqrt{\varepsilon}]$. Here, the additional uniform estimate $\sup_{t < T} \mathbb{E}|Q_t^0 - \Xi_t|^2 < \infty$ from Lemma 5.4 allows us to use the deterministic switching time $T - \sqrt{\varepsilon}$ and hence make the resulting strategy (5.19) fully explicit.

An example is shown in Figure 1. We remark that the slope of the terminal bridge could be changed by modifying the switching time to $T - c\sqrt{\varepsilon}$ for some $c > 0$.

We can now state our main theorem. Its bounds in (5.30) and (5.31) apply to the explicit strategy Q^ε from (5.19), not merely to the optimal value, and show that the excess impact cost $J_0(Q^\varepsilon) - J_0(Q^0)$ and the total regularized cost above $V(0)$ are both $O(\sqrt{\varepsilon})$.

Theorem 5.7 (Main result). *For every $0 < \varepsilon \leq T^2/4$, the explicit strategy $Q^\varepsilon \in \mathcal{Q}_{\text{ac}}$ from (5.19) satisfies*

$$0 \leq J_0(Q^\varepsilon) - J_0(Q^0) \leq C_J(\tilde{C}_{\text{tr}}\sqrt{\varepsilon} + 4\Lambda_\Xi\varepsilon), \quad (5.30)$$

$$0 \leq J_\varepsilon(Q^\varepsilon) - V(0) \leq (1 + C_J)(\tilde{C}_{\text{tr}}\sqrt{\varepsilon} + 4\Lambda_\Xi\varepsilon). \quad (5.31)$$

As a consequence, the value functions satisfy

$$0 \leq V(\varepsilon) - V(0) \leq (1 + C_J)(\tilde{C}_{\text{tr}}\sqrt{\varepsilon} + 4\Lambda_\Xi\varepsilon) \quad (5.32)$$

and in particular

$$V(\varepsilon) - V(0) = O(\sqrt{\varepsilon}), \quad \varepsilon \downarrow 0.$$

Proof. Let Q^ε be the explicit strategy from (5.19). Since

$$\mathbb{E} \int_0^T (Q_t^\varepsilon - Q_t^0)^2 dt \leq \mathbb{E} \left[\int_0^T (Q_t^\varepsilon - Q_t^0)^2 dt + \varepsilon \int_0^T (\dot{Q}_t^\varepsilon)^2 dt \right],$$

Lemmas 5.3 and 5.5 give

$$0 \leq J_0(Q^\varepsilon) - J_0(Q^0) \leq C_J(\tilde{C}_{\text{tr}}\sqrt{\varepsilon} + 4\Lambda_\Xi\varepsilon),$$

which is (5.30). Adding the regularization term and noting that likewise

$$\varepsilon \mathbb{E} \left[\int_0^T (\dot{Q}_t^\varepsilon)^2 dt \right] \leq \mathbb{E} \left[\int_0^T (Q_t^\varepsilon - Q_t^0)^2 dt + \varepsilon \int_0^T (\dot{Q}_t^\varepsilon)^2 dt \right]$$

yields (5.31). Since $V(\varepsilon) \leq J_\varepsilon(Q^\varepsilon)$, (5.32) follows. $\square$

Recall that convergence of Q^ε to Q^0 in L^2 was already shown in (5.23). The following complements that result with the convergence of the true optimizer $Q^{*,\varepsilon}$.

Corollary 5.8 (Convergence of the regularized optimizer). *Let $Q^{*,\varepsilon}$ be the regularized optimizer from Remark 5.1, and let $Y^{*,\varepsilon} := Y^{Q^{*,\varepsilon}}$. Define*

$$\mathfrak{C}_\mathcal{H} := \inf_{t \in [0, T]} \frac{2\beta_t + \dot{\gamma}_t}{2\lambda_t} > 0 \quad \text{and} \quad C_{\text{inv}} := \frac{2\|\lambda^{-1}\|_{L^\infty(0, T)}^2 + T^2 \left\| \frac{\beta + \dot{\gamma}}{\lambda} \right\|_{L^\infty(0, T)}^2}{\mathfrak{C}_\mathcal{H}}.$$

Then, for every $0 < \varepsilon \leq T^2/4$,

$$0 \leq J_0(Q^{*,\varepsilon}) - J_0(Q^0) = \|Y^{*,\varepsilon} - Y^0\|_\mathcal{H}^2 \leq (1 + C_J)(\tilde{C}_{\text{tr}}\sqrt{\varepsilon} + 4\Lambda_\Xi\varepsilon), \quad (5.33)$$

and

$$\mathbb{E} \int_0^T |Q_t^{*,\varepsilon} - Q_t^0|^2 dt \leq C_{\text{inv}}(1 + C_J)(\tilde{C}_{\text{tr}}\sqrt{\varepsilon} + 4\Lambda_\Xi\varepsilon). \quad (5.34)$$

In particular,

$$J_0(Q^{*,\varepsilon}) - J_0(Q^0) = O(\sqrt{\varepsilon}), \quad \mathbb{E} \int_0^T |Q_t^{*,\varepsilon} - Q_t^0|^2 dt = O(\sqrt{\varepsilon}), \quad \varepsilon \downarrow 0.$$

Proof. Since $Q^{*,\varepsilon}$ minimizes J_ε over $\mathcal{Q}_{\text{ac}}$, we have

$$J_0(Q^{*,\varepsilon}) - J_0(Q^0) \leq J_\varepsilon(Q^{*,\varepsilon}) - J_0(Q^0) = V(\varepsilon) - V(0).$$

The identity in (5.11) and the value bound (5.32) therefore yield (5.33).

It remains to control the inventory difference. More generally, let $Q \in \mathcal{Q}$, and set

$$R_t := Q_t - Q_t^0, \quad Z_t := Y_t^Q - Y_t^0.$$

Subtracting the two instances of (4.14) gives

$$R_t = \frac{Z_t}{\lambda_t} + \int_0^t \frac{\beta_s + \dot{\gamma}_s}{\lambda_s} Z_s ds, \quad 0 \leq t \leq T.$$

Consequently, by Cauchy–Schwarz and Fubini,

$$\begin{aligned} \mathbb{E} \int_0^T R_t^2 dt &\leq 2\|\lambda^{-1}\|_{L^\infty(0, T)}^2 \mathbb{E} \int_0^T Z_t^2 dt \\ &\quad + 2 \left\| \frac{\beta + \dot{\gamma}}{\lambda} \right\|_{L^\infty(0, T)}^2 \mathbb{E} \int_0^T t \int_0^t Z_s^2 ds dt \\ &\leq \left(2\|\lambda^{-1}\|_{L^\infty(0, T)}^2 + T^2 \left\| \frac{\beta + \dot{\gamma}}{\lambda} \right\|_{L^\infty(0, T)}^2 \right) \mathbb{E} \int_0^T Z_t^2 dt. \end{aligned}$$

Moreover, by the definition of the $\mathcal{H}$ -norm,

$$\mathbb{E} \int_0^T Z_t^2 dt \leq \frac{1}{\underline{c}_{\mathcal{H}}} \|Z\|_{\mathcal{H}}^2.$$

Thus,

$$\mathbb{E} \int_0^T (Q_t - Q_t^0)^2 dt \leq C_{\text{inv}} \|Y^Q - Y^0\|_{\mathcal{H}}^2.$$

Applying this estimate to $Q = Q^{*,\varepsilon}$ and using (5.33) proves (5.34). $\square$

Our final result shows that the generic rate $\sqrt{\varepsilon}$ is sharp in the benchmark case where $\beta, \lambda > 0$ are constants, the terminal target position Ξ_T is deterministic, and $y = 0$.

Proposition 5.9 (Sharpness). *Suppose $y = 0$, that $\beta, \lambda > 0$ are constants, and that the target position Ξ_T is deterministic. Then,*

$$V(\varepsilon) = \frac{\lambda \Xi_T^2}{\beta T + 2} + \frac{2\sqrt{\beta\lambda} \Xi_T^2}{(\beta T + 2)^2} \sqrt{\varepsilon} + O(\varepsilon), \quad \varepsilon \downarrow 0.$$

Consequently, if $\Xi_T \neq 0$, then

$$V(\varepsilon) - V(0) \sim \frac{2\sqrt{\beta\lambda} \Xi_T^2}{(\beta T + 2)^2} \sqrt{\varepsilon}, \quad \varepsilon \downarrow 0,$$

and hence the rate $O(\sqrt{\varepsilon})$ is sharp.

Proof. Set $D := \beta T + 2$. We first consider the unregularized case $\varepsilon = 0$. Under the stated assumptions, this is the original Obizhaeva–Wang model [20]. By [20, Proposition 2], the optimal strategy is

$$dQ_t^0 = \frac{\Xi_T}{D} \delta_0(dt) + \frac{\beta \Xi_T}{D} dt + \frac{\Xi_T}{D} \delta_T(dt), \quad 0 \leq t \leq T,$$

and its cost is

$$V(0) = \frac{\lambda \Xi_T^2}{D}. \tag{5.35}$$

We next use the explicit solution of the regularized problem (5.3) obtained by Chen, Horst, and Tran [12, Theorem 2.1]. Define

$$k_\varepsilon := \sqrt{\beta^2 + \frac{\beta\lambda}{\varepsilon}}, \quad K_\varepsilon := \frac{k_\varepsilon T}{2},$$

and

$$a_\varepsilon := \frac{\lambda}{\varepsilon} \sinh(K_\varepsilon), \quad b_\varepsilon := \beta k_\varepsilon \cosh(K_\varepsilon) + k_\varepsilon^2 \sinh(K_\varepsilon), \quad c_\varepsilon := \frac{\lambda}{\varepsilon}.$$

Then the optimal inventory and trading rate are

$$Q_t^{*,\varepsilon} = \Xi_T \frac{a_\varepsilon + b_\varepsilon t + c_\varepsilon \sinh(k_\varepsilon(t - T/2))}{2a_\varepsilon + b_\varepsilon T}, \quad 0 \leq t \leq T,$$

and

$$u_t^{*,\varepsilon} = d_\varepsilon + \frac{\Xi_T c_\varepsilon k_\varepsilon}{2a_\varepsilon + b_\varepsilon T} \cosh(k_\varepsilon(t - T/2)), \quad 0 \leq t < T,$$

where

$$d_\varepsilon := \frac{\Xi_T b_\varepsilon}{2a_\varepsilon + b_\varepsilon T}. \quad (5.36)$$

The optimal value is not stated explicitly in [12], but it follows from their proof. Indeed, [12, Appendix 6.1] gives the first-order condition

$$2\varepsilon u_t^{*,\varepsilon} + \lambda \int_0^T e^{-\beta|t-s|} u_s^{*,\varepsilon} ds + \mu_\varepsilon = 0$$

for some Lagrange multiplier μ_ε , together with

$$\mu_\varepsilon = -\frac{2\varepsilon k_\varepsilon^2}{\beta^2} d_\varepsilon.$$

For an absolutely continuous strategy with trading rate u , the objective in the present constant-coefficient setting can be written as

$$J_\varepsilon(Q) = \mathbb{E} \left[\frac{\lambda}{2} \int_0^T \int_0^T e^{-\beta|t-s|} u_t u_s ds dt + \varepsilon \int_0^T u_t^2 dt \right].$$

Multiplying the first-order condition by $u_t^{*,\varepsilon}$, integrating over $[0, T]$, and using $\int_0^T u_t^{*,\varepsilon} dt = \Xi_T$, we therefore obtain

$$V(\varepsilon) = J_\varepsilon(Q^{*,\varepsilon}) = -\frac{\mu_\varepsilon \Xi_T}{2} = \frac{\varepsilon k_\varepsilon^2}{\beta^2} d_\varepsilon \Xi_T. \quad (5.37)$$

We now expand d_ε . From (5.36),

$$d_\varepsilon = \Xi_T \frac{k_\varepsilon^2 + \beta k_\varepsilon \coth(K_\varepsilon)}{T(k_\varepsilon^2 + \beta k_\varepsilon \coth(K_\varepsilon)) + \frac{2}{\beta}(k_\varepsilon^2 - \beta^2)}. \quad (5.38)$$

Since $k_\varepsilon \rightarrow \infty$ and $\coth(K_\varepsilon) = 1 + O(e^{-k_\varepsilon T})$, we obtain

$$k_\varepsilon^2 + \beta k_\varepsilon \coth(K_\varepsilon) = k_\varepsilon^2 + \beta k_\varepsilon + O(k_\varepsilon^{-2}),$$

and

$$T(k_\varepsilon^2 + \beta k_\varepsilon \coth(K_\varepsilon)) + \frac{2}{\beta}(k_\varepsilon^2 - \beta^2) = \frac{D}{\beta} k_\varepsilon^2 + \beta T k_\varepsilon - 2\beta + O(k_\varepsilon^{-2}).$$

Hence

$$d_\varepsilon = \frac{\beta \Xi_T}{D} \frac{1 + \frac{\beta}{k_\varepsilon} + O(k_\varepsilon^{-2})}{1 + \frac{\beta^2 T}{D k_\varepsilon} + O(k_\varepsilon^{-2})}.$$

Using $(1+x)^{-1} = 1 - x + O(x^2)$ and $1 - \beta T/D = 2/D$, we conclude that

$$d_\varepsilon = \frac{\beta \Xi_T}{D} + \frac{2\beta^2 \Xi_T}{D^2} \frac{1}{k_\varepsilon} + O(k_\varepsilon^{-2}), \quad \varepsilon \downarrow 0. \quad (5.39)$$




Figure 1: Comparison of the unregularized optimal strategy Q^0 , the explicit strategy Q^ε from (5.19), and the regularized optimal strategy $Q^{*,\varepsilon}$ in the constant-coefficient benchmark with $T = 1$, $\beta = \lambda = 1$, $y = 0$, and deterministic terminal target $\Xi_T = 1$. The colored dotted and dashed lines show Q^ε and $Q^{*,\varepsilon}$, respectively, for $\varepsilon \in \{0.01, 0.0025\}$.

The remaining factor in (5.37) satisfies

$$\frac{\varepsilon k_\varepsilon^2}{\beta^2} = \frac{\lambda}{\beta} + \varepsilon. \quad (5.40)$$

Moreover,

$$\frac{1}{k_\varepsilon} = \sqrt{\frac{\varepsilon}{\beta\lambda}} \left(1 + \frac{\beta\varepsilon}{\lambda}\right)^{-1/2} = \sqrt{\frac{\varepsilon}{\beta\lambda}} + O(\varepsilon^{3/2}),$$

and therefore $\frac{1}{k_\varepsilon} = O(\varepsilon)$. Substituting (5.39) and (5.40) into (5.37) gives

$$\begin{aligned} V(\varepsilon) &= \left(\frac{\lambda}{\beta} + \varepsilon\right) \left(\frac{\beta\Xi_T}{D} + \frac{2\beta^2\Xi_T}{D^2} \frac{1}{k_\varepsilon} + O(k_\varepsilon^{-2})\right) \Xi_T \\ &= \frac{\lambda\Xi_T^2}{D} + \frac{2\lambda\beta\Xi_T^2}{D^2} \frac{1}{k_\varepsilon} + O(\varepsilon) = \frac{\lambda\Xi_T^2}{D} + \frac{2\sqrt{\beta\lambda}\Xi_T^2}{D^2} \sqrt{\varepsilon} + O(\varepsilon). \end{aligned}$$

Together with (5.35), this proves the claimed expansion. If $\Xi_T \neq 0$, the coefficient of $\sqrt{\varepsilon}$ is strictly positive, so the asserted asymptotic equivalence and sharpness follow. $\square$

Figure 1 illustrates the two regularizations in the constant-coefficient benchmark of Proposition 5.9. As ε decreases, both the simplified strategy Q^ε and the regularized optimizer $Q^{*,\varepsilon}$ approach the singular unregularized optimizer Q^0 .

References

- [1] J. Ackermann, T. Kruse, and M. Urusov. Càdlàg semimartingale strategies for optimal trade execution in stochastic order book models. *Finance Stoch.*, 25(4):757–810, 2021.

- [2] J. Ackermann, T. Kruse, and M. Urusov. Optimal trade execution in an order book model with stochastic liquidity parameters. *SIAM J. Financial Math.*, 12(2):788–822, 2021.
- [3] J. Ackermann, T. Kruse, and M. Urusov. Reducing Obizhaeva–Wang-type trade execution problems to LQ stochastic control problems. *Finance Stoch.*, 28(3):813–863, 2024.
- [4] A. Alfonsi and J. I. Acevedo. Optimal execution and price manipulations in time-varying limit order books. *Appl. Math. Finance*, 21(3):201–237, 2014.
- [5] S. Ankirchner and T. Kruse. Optimal position targeting with stochastic linear–quadratic costs. *Banach Center Publ.*, 104(1):9–24, 2015.
- [6] P. Bank and A. Fruth. Optimal order scheduling for deterministic liquidity patterns. *SIAM J. Financial Math.*, 5(1):137–152, 2014.
- [7] P. Bank, H. M. Soner, and M. Voß. Hedging with temporary price impact. *Math. Financ. Econ.*, 11(2):215–239, 2017.
- [8] P. Bank and M. Voß. Linear quadratic stochastic control problems with stochastic terminal constraint. *SIAM J. Control Optim.*, 56(2):672–699, 2018.
- [9] J. Cai, M. Rosenbaum, and P. Tankov. Asymptotic lower bounds for optimal tracking: A linear programming approach. *Ann. Appl. Probab.*, 27(4):2455–2514, 2017.
- [10] J. Cai, M. Rosenbaum, and P. Tankov. Asymptotic optimal tracking: Feedback strategies. *Stochastics*, 89(6–7):943–966, 2017.
- [11] Á. Cartea, S. Jaimungal, and J. Penalva. *Algorithmic and High-Frequency Trading*. Cambridge University Press, 2015.
- [12] Y. Chen, U. Horst, and H. H. Tran. Optimal trade execution strategy and implementation with deterministic market impact parameters. *Appl. Math. Finance*, 32(1):1–29, 2025.
- [13] Y. Dolinsky, B. Gottesman, and O. Gurel-Gurevich. A note on costs minimization with stochastic target constraints. *Electron. Commun. Probab.*, 25(11):1–12, 2020.
- [14] L. Dolinsky and Y. Dolinsky. Optimal liquidation with high risk aversion and small linear price impact. *Decis. Econ. Finance*, 47(1):183–198, 2024.
- [15] A. Fruth, T. Schöneborn, and M. Urusov. Optimal trade execution and price manipulation in order books with time-varying liquidity. *Math. Finance*, 24(4):651–695, 2014.
- [16] A. Fruth, T. Schöneborn, and M. Urusov. Optimal trade execution in order books with stochastic liquidity. *Math. Finance*, 29(2):507–541, 2019.
- [17] P. Graewe and U. Horst. Optimal trade execution with instantaneous price impact and stochastic resilience. *SIAM J. Control Optim.*, 55(6):3707–3725, 2017.
- [18] U. Horst and E. Kivman. Optimal trade execution under small market impact and portfolio liquidation with semimartingale strategies. *Finance Stoch.*, 28(3):759–812, 2024.
- [19] M. Nutz, K. Webster, and L. Zhao. Unwinding Stochastic Order Flow: When to Warehouse Trades. *Math. Finance*, 36(3):500–527, 2026.
- [20] A. A. Obizhaeva and J. Wang. Optimal trading strategy and supply/demand dynamics. *J. Financial Mark.*, 16(1):1–32, 2013.
- [21] K. Webster. *Handbook of Price Impact Modeling*. CRC Press, Boca Raton, FL, 2023.