

Stochastic Optimal Control of Hawkes Jump-Diffusion Systems

Daria Sakhanda*

Joshué Helí Ricalde-Guerrero†

Abstract

This paper is devoted to developing a framework for stochastic growth models with environmental risk, in which rare but catastrophic shocks interact with capital accumulation and pollution. Building on the Poisson point process formulation studied in [12], we extend the model to disasters driven by a marked Hawkes process, allowing past disasters to temporarily increase the likelihood of subsequent shocks. Our work focuses on a subcritical Markovian Hawkes specification, in which the state space is augmented by the self-excitation component of disaster risk. We establish the well-posedness and nonexplosion of the resulting controlled Hawkes dynamics. Using the Hamilton-Jacobi-Bellman characterization of the corresponding Poisson control problem, we obtain quantitative estimates for the Hawkes excitation process and prove, under a small-excitation scaling, that the Hawkes value function converges to its Poisson counterpart as the magnitude of self-excitation vanishes. This provides a rigorous Poisson approximation of the stochastic control problem and quantifies the effect of self-excitation on optimal growth under environmental disaster risk.

Keywords. Stochastic optimal control, Hamilton-Jacobi-Bellman equations, Hawkes processes, marked point processes, Poisson random measures, jump-diffusion processes, integro-differential equations, growth-environment models, pollution-dependent intensity, rare disasters.

MSC Classification. 93E20, 60G55, 91-10

1 Introduction

The study of economic growth under environmental risk raises fundamental mathematical questions in stochastic control for systems combining continuous dynamics and discontinuous jumps. Classical growth-environment models typically describe pollution and damages as deterministic or smoothly evolving, with dynamics governed by stochastic differential equations without discontinuities. Empirical evidence, however, shows that risk also takes the form of rare but catastrophic events – such as abrupt climate disasters, ecosystem collapses, or large-scale technological failures – which arrive unpredictably and are naturally modeled as jumps with large economic impacts. While Poisson-based models provide a natural framework for rare catastrophic events, they do not account for temporal clustering: the occurrence of one disaster may increase the likelihood of subsequent events, generating periods of persistently elevated risk.

Motivated by economic questions of growth, environmental change, and climate risk, this paper extends our previous Poisson framework [12] by introducing self-exciting disaster arrivals through a marked Hawkes process. The disaster intensity combines a pollution-dependent baseline component with an excitation component generated by past disasters. Each event increases the excitation level according to its mark, while excitation gradually decays between successive events. The resulting framework therefore allows disaster risk to depend simultaneously on the current environmental state and on the history of previous disasters.

*ETH Zürich, Department of Mathematics, Switzerland (daria.sakhanda@math.ethz.ch).

†ETH Zürich, Department of Mathematics, Switzerland (joshue.ricalde@math.ethz.ch).

From the mathematical perspective, the Hawkes specification introduces an additional state variable and an endogenous jump intensity, leading to a stochastic control problem on an augmented state space. We establish the global strong well-posedness and non-explosion of the controlled marked Hawkes system, as well as a rigorous connection between the Hawkes and Poisson formulations. We consider a family of Hawkes models in which the excitation generated by individual disasters becomes progressively smaller. Under suitable moment and stability conditions, we derive uniform discounted estimates for the excitation process and show that the Hawkes value function converges to the corresponding Poisson value function in the small-excitation limit. Moreover, we obtain an explicit first-order bound for the approximation error.

Thus, the main contributions of the paper are threefold:

1. We introduce a controlled marked Hawkes specification for pollution-dependent and self-exciting catastrophic risk in a growth-environment model.
2. We establish the mathematical foundations of the resulting stochastic control problem, including global strong well-posedness and non-explosion.
3. We establish a quantitative connection between the Hawkes and Poisson formulations by proving convergence of their value functions in the small-excitation regime, together with an explicit rate of approximation.

The rest of the paper is organized as follows. In Section 2, we introduce the modelling framework and describe how the state-dependent Poisson disaster mechanism studied in [12] is extended to a marked Hawkes specification with self-excitation. Section 3 collects the basic concepts and notation for marked Hawkes processes and introduces the subcriticality condition used throughout the analysis. In Section 4, we formulate the controlled Hawkes model, specify the admissible controls and standing assumptions, and state our main result. Section 5 is devoted to the Hawkes-Poisson approximation. We first establish the well-posedness and nonexplosion of the controlled Hawkes dynamics and then study the asymptotic behaviour of the excitation process and its implications for the disaster mechanism. Finally, we introduce a small-excitation scaling, parameterized by $\eta > 0$, where η controls the magnitude of Hawkes self-excitation, and prove that the Hawkes value function converges to its Poisson counterpart as $\eta \downarrow 0$, with an explicit $O(\eta)$ approximation bound.

2 Modelling framework

We consider a stylized representation of the global economy, which produces a single composite good under constant returns to scale. Production relies on the aggregate capital stock at time t , denoted by K_t which includes physical capital, human capital, and intangible assets. The production process $(K_t)_{t \geq 0}$ generates pollution: at each instant t , greenhouse gas (GHG) emissions E_t are released into the atmosphere. These emissions accumulate in the atmospheric pollution stock, P_t , which increases with the flow of emissions and decreases through a natural absorption rate $\alpha \in [0, 1)$, assumed to be small or negligible. While P_t is referred to as the pollution stock, it can more generally be interpreted as the *inverse of environmental quality*. Similarly, the emissions variable E_t , can be viewed more broadly as any environmentally damaging by-product of economic activity.

In this paper, we extend the disaster arrival mechanism from state-dependent Poisson processes, as studied in [12], to a Hawkes process with self-excitation. This specification captures the empirically observed clustering of extreme events, whereby the occurrence of a disaster increases the likelihood of subsequent disasters in the short run, before this excess risk gradually decays over time (e.g., aftershocks, persistent weather regimes, or cascading failures).

Following the notation from [12], the controlled dynamics of the model are

$$dK_t = b^{\text{cap}}(K_t, P_t, C_t, \theta_t) dt - \int_{\mathcal{Z}} (1 - \omega(K_{t-}, P_{t-}, \zeta)) K_{t-} N(dt, d\zeta), \quad (2.1)$$

$$dP_t = b^{\text{pol}}(K_t, P_t, C_t, \theta_t) dt + \sigma P_t dW_t, \quad (2.2)$$

$$dY_t = -bY_t dt + \int_{\mathcal{Z}} a(\zeta) N(dt, d\zeta). \quad (2.3)$$

Here, $Y = (Y_t)_{t \geq 0}$ denotes the self-exciting component of the disaster intensity, while N is the marked integer-valued random measure associated with disaster arrivals. The process $W = (W_t)_{t \geq 0}$ is a Brownian motion, independent of the underlying Poisson random measure used in the construction of N , and $\sigma > 0$ determines the magnitude of the diffusion component in the pollution dynamics. Moreover, $b > 0$ is the decay rate of self-excitation, whereas $a(\zeta) \geq 0$ measures the increase in excitation generated by a disaster with mark ζ . The processes Y and N will be specified more precisely below.

In comparison, the marked PRM-driven model from Section 2.4 in [12], follows the same dynamics on the capital and pollution (2.1) and (2.2), respectively, with no additional self-exciting component (2.3). The Hawkes specification can therefore be viewed as an extension of the Poisson model in which past disasters affect the likelihood of subsequent shocks through the excitation process Y . Under the subcriticality condition, we establish the well-posedness and nonexplosion of the Hawkes-driven model. We then introduce the small-excitation scaling

$$a_\eta(\zeta) := \eta a(\zeta), \quad \eta > 0,$$

and show that, as $\eta \downarrow 0$, the corresponding Hawkes control problem converges to its Poisson counterpart.

The predictable disaster intensity is

$$\lambda_t^H = \mu(P_{t-}^H) + Y_{t-}. \quad (2.4)$$

Thus, $\mu(P_{t-}^H)$ represents the pollution-dependent baseline intensity, while Y_{t-} captures the additional disaster risk generated by past events. At each disaster time with mark ζ , the excitation process increases by $a(\zeta)$ and subsequently decays at rate b . Consequently, past disasters affect future disaster arrivals through Y , generating the characteristic self-exciting behaviour of the Hawkes process.

In comparison, the marked Poisson model of [12] has disaster intensity

$$\lambda_t^P = \mu(P_{t-}^P), \quad (2.5)$$

and therefore contains no self-exciting component. The Hawkes specification can thus be viewed as an extension of the Poisson benchmark in which past disasters generate an additional endogenous component of disaster risk. A central objective of the present paper is to quantify the effect of this component and to establish conditions under which the corresponding Hawkes control problem can be approximated by its Poisson counterpart.

3 Preliminaries

3.1 Basic concepts and notation

Throughout the paper, we work on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, where $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$ satisfies the usual conditions. For a càdlàg process $X = (X_t)_{t \geq 0}$, we denote by

$$X_{t-} := \lim_{s \uparrow t} X_s$$

its left limit at time $t > 0$, and by

$$\Delta X_t := X_t - X_{t-}$$

its jump at time t .

Let $(\mathcal{Z}, \mathfrak{Z})$ be a measurable mark space, endowed with a probability measure ν ; see [4]. The variable $\zeta \in \mathcal{Z}$ will represent the mark associated with a disaster, describing its magnitude or type. Integration with respect to the mark distribution will be written as

$$\int_{\mathcal{Z}} f(\zeta) \nu(d\zeta)$$

whenever the integral is well defined.

For a measurable space $(E, \mathcal{E})$, we denote by $\mathcal{B}(E)$ its Borel σ -algebra whenever E is endowed with a topology. The indicator function of a measurable set A is denoted by $\mathbf{1}_A$.

The state spaces of the Poisson and Hawkes models are denoted, respectively, by

$$\mathcal{S}_P := \mathbb{R}_{>0} \times \mathbb{R}_{>0}, \quad \mathcal{S}_H := \mathcal{S}_P \times \mathbb{R}_{\geq 0}.$$

Accordingly, the Poisson state is represented by (K, P) , whereas the Hawkes state is augmented by the excitation component Y and is represented by (K, P, Y) .

The following subsection recalls the point-process notation and the basic properties of marked Hawkes processes required throughout the paper.

3.2 Preliminaries on Hawkes processes

Let $\{(\tau_n, \Delta_n)\}_{n \geq 1}$ be a sequence of random variables taking values in $\mathbb{R}_{\geq 0} \times \mathcal{Z}$, where

$$0 < \tau_1 < \tau_2 < \dots$$

denote the event times. We refer to τ_n as the *arrival time of the n -th event* (disaster) and to Δ_n as the *mark of the n -th event* (magnitude or type of disaster). Following [4], let $\mathcal{N}_{\mathbb{R} \times \mathcal{Z}}^{\#}$ denote space of integer-valued measures over $\mathbb{R} \times \mathcal{Z}$ and define N as the measurable mapping

$$\begin{aligned} N : (\Omega, \mathcal{F}) &\longrightarrow (\mathcal{N}_{\mathbb{R} \times \mathcal{Z}}^{\#}, \mathcal{B}(\mathcal{N}_{\mathbb{R} \times \mathcal{Z}}^{\#})) \\ \omega &\longmapsto \sum_{n \geq 1} \delta_{(\tau_n(\omega), \Delta_n(\omega))}(dt, d\zeta). \end{aligned}$$

We refer to both the sequence of events $\{(\tau_n, \Delta_n)\}$ and the (random) counting measure N as *marked point process* interchangeably.

Integrals with respect to the point process are defined (realization-by-realization) as

$$\int_{\mathbb{R} \times \mathcal{Z}} f(t, \zeta) N(\omega; dt \times d\zeta) = \sum_{n \geq 1} f(\tau_n(\omega), \Delta_n(\omega))$$

for any measurable function f with $\text{dom } f \subseteq \mathbb{R}_{\geq 0} \times \mathcal{Z}$. In particular, the *ground counting process* [4] associated with N is defined as

$$N_t := \int_{\mathbb{R} \times \mathcal{Z}} \mathbf{1}_{(0, t] \times \mathcal{Z}}(t, \zeta) N(dt \times d\zeta) = N((0, t] \times \mathcal{Z}), \quad \forall t \geq 0, \quad (3.1)$$

so that N_t records the total number of disasters occurring up to time t . Let $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$ be filtration satisfying the usual conditions, such that $\mathbb{F}$ contains the filtration generated by $\{N((0, s] \times \mathcal{Z}); s \leq t, \mathcal{Z} \in \mathfrak{Z}\}_{t \geq 0}$. From hereonout, we say the point process $\{(\tau_n, \Delta_n)\}$ –

equivalently, N is a *marked Hawkes process* if there exists an $\mathbb{F}$ -predictable *stochastic intensity kernel* Λ satisfying the following self-exciting jump SDE:

$$\begin{aligned}\Lambda(\omega; t, d\zeta) &= \lambda(\omega; t, \zeta) \nu(d\zeta), \\ \lambda(\omega; t, \zeta) &= \mu(\omega; t) + \alpha(Y_t(\omega), \zeta)\end{aligned}\tag{3.2}$$

$$Y_t(\omega) = -b \int_{(0,t]} Y_s(\omega) ds + \int_{(-\infty, t] \times \mathcal{Z}} a(\omega; t-s, \zeta) N(\omega; ds \times d\zeta),$$

where ν is some given probability measure over $(\mathcal{Z}, \mathfrak{Z})$, and μ is some given $\mathbb{F}$ -predictable process with values in $\mathbb{R}_{\geq 0}$. Thus, each disaster increases the future arrival intensity through the excitation process Y , while this additional excitation decays exponentially between successive events. The strength of this self-exciting mechanism must be sufficiently small relative to its rate of decay in order to ensure stability of the Hawkes process. This leads to the subcriticality condition introduced next.

3.3 Subcriticality condition

A fundamental stability condition for Hawkes processes is the *subcriticality condition*. In the cluster representation of a linear Hawkes process, this condition requires the expected number of offspring generated by a single event to be strictly smaller than one; see [6] and [3].

Definition 3.1 (Subcritical marked Hawkes process). Let

$$h : \mathbb{R}_{>0} \times \mathcal{Z} \longrightarrow \mathbb{R}_{\geq 0}$$

be a measurable excitation kernel satisfying

$$\int_0^\infty \int_{\mathcal{Z}} h(s, \zeta) \nu(d\zeta) ds < \infty.$$

The quantity

$$m_h := \int_0^\infty \int_{\mathcal{Z}} h(s, \zeta) \nu(d\zeta) ds\tag{3.3}$$

is called the *branching ratio* and represents the expected number of direct offspring generated by a single event. The marked Hawkes process is said to be *subcritical* if

$$m_h < 1.\tag{3.4}$$

See [6] for the cluster representation of linear Hawkes processes.

4 Controlled Hawkes model and main results

We start by defining the set of admissible controls and specifying the standing assumptions. The *set of admissible actions* corresponds to a given subset

$$\mathfrak{a} \subseteq \mathbb{R}_{\geq 0} \times [0, 1].$$

Definition 4.1 (Admissible controls). For every $(k, p, y) \in \mathcal{S}_H$, the set of admissible controls is defined as

$$\begin{aligned}\mathcal{A}(k, p, y) := \Big\{ \alpha : \Omega \times \mathbb{R}_{\geq 0} \longrightarrow \mathfrak{a} \mid & (\alpha_t)_{t \geq 0} \text{ is } \mathbb{F}\text{-predictable,} \\ & (K_0, P_0, Y_0) = (k, p, y) \in \mathcal{F}_0, \\ & \text{and } \mathbb{E} \left[\int_0^\infty \|\alpha_t\|^2 dt \right] < \infty \Big\}.\end{aligned}$$

In the context of the problem, the marginals of an admissible control correspond to the consumption and abatement processes $(C_t)_{t \geq 0}$ and $(\theta_t)_{t \geq 0}$, respectively.

Assumptions 4.2.

1. **On the stochastic basis:** The filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ admits an $\mathbb{F}$ -Brownian motion W , and an independent $\mathbb{F}$ -Poisson random measure M on $\mathbb{R}_{\geq 0} \times \mathcal{Z} \times \mathbb{R}_{\geq 0}$ with intensity measure $dt \otimes \nu(d\zeta) \otimes dr$ for some σ -finite measure ν with finite second moment. In addition, $\mathbb{F} = \{\mathcal{F}_t\}_{t \geq 0}$ is (the complete, right-continuous augmentation of) the filtration

$$\mathcal{F}_t := \sigma\left\{M((0, s] \times C \times D) \mid 0 \leq s \leq t, C \in \mathfrak{Z}, D \in \mathcal{B}(\mathbb{R}_{\geq 0})\right\} \vee \mathcal{F}_t^W, \quad t \geq 0. \quad (4.1)$$

2. **On the coefficients:**

- (a) $\mu : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{\geq 0}$ is locally Lipschitz and there exists $L_\mu > 0$ such that

$$\mu(p) \leq L_\mu(1 + p), \quad p > 0. \quad (4.2)$$

- (b) $a : \mathcal{Z} \rightarrow \mathbb{R}_{\geq 0}$ is measurable and

$$\int_{\mathcal{Z}} a(\zeta) \nu(d\zeta) < \infty, \quad \int_{\mathcal{Z}} a(\zeta)^2 \nu(d\zeta) < \infty. \quad (4.3)$$

- (c) For every admissible control (C, θ) , the functions b^{cap} and b^{pol} are locally Lipschitz in (k, p) , uniformly with respect to the control on bounded sets, and satisfy a linear-growth condition

$$|b^{\text{cap}}(k, p, c, \theta)| + |b^{\text{pol}}(k, p, c, \theta)| \leq L(1 + k + p + c) \quad (4.4)$$

for some $L > 0$.

- (d) The survival map

$$\omega : \mathbb{R}_{>0} \times \mathbb{R}_{>0} \times \mathcal{Z} \longrightarrow (0, 1] \quad (4.5)$$

is measurable, locally Lipschitz in (k, p) uniformly in ζ , and satisfies

$$0 < \omega(k, p, \zeta) \leq 1. \quad (4.6)$$

- (e) Starting from $K_0 > 0$ and $P_0 > 0$, the continuous dynamics do not attain the boundary of $\mathbb{R}_{>0}^2$ in finite time; that is, $K_t > 0$, $P_t > 0$ for every finite t for which the continuous dynamics are defined.

3. **Subcriticality:** Let

$$\bar{a}_1 := \int_{\mathcal{Z}} a(\zeta) \nu(d\zeta).$$

The excitation parameters satisfy

$$\bar{a}_1 < b. \quad (4.7)$$

For the exponential excitation kernel considered below, condition (4.7) is precisely the specialization of the general subcriticality condition introduced in Definition 3.1.

With the previous considerations in mind, the augmented system $X^H = (X_t^H)_{t \geq 0}$ is defined as the $\mathcal{S}_H$ -valued, $\mathbb{F}$ -adapted process

$$X_t^H = (K_t^H, P_t^H, Y_t), \quad \forall t \geq 0,$$

characterized by the controlled dynamics:

$$dK_t^H = b^{\text{cap}}(K_t^H, P_t^H, \alpha_t) dt - \int_{\mathcal{Z}} (1 - \omega(K_{t-}^H, P_{t-}^H, \zeta)) K_{t-}^H N^H(dt \times d\zeta), \quad (4.8)$$

$$dP_t^H = b^{\text{pol}}(K_t^H, P_t^H, \alpha_t) dt + \sigma P_t^H dW_t, \quad (4.9)$$

$$(K_0^H, P_0^H, Y_0) = (k, p, y) \in \mathcal{S}_H. \quad (4.10)$$

Here N^H is a *marked Hawkes process* with ground process $N = (N_t)$, as defined in (3.1) with

$$\mu(\omega; t) = \mu(P_t^H(\omega)) \quad \text{and} \quad a(\omega; t - s, \zeta) = a(\zeta), \quad (4.11)$$

where $\mu : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{\geq 0}$ and $a : \mathcal{Z} \rightarrow \mathbb{R}_{\geq 0}$ are the deterministic measurable functions specified in Assumptions 4.2.

We now state our main result:

Theorem 4.3. *Under Assumptions 4.2, the following hold:*

1. **Well-posedness of the Hawkes-driven model:** *Let $(k, p, y) \in \mathcal{S}_H$ be given. For any admissible control $\alpha \in \mathcal{A}(k, p, y)$, there exists a unique $\mathbb{F}$ -adapted strong solution*

$$X^H = (K^H, P^H, Y)$$

to (4.8)–(4.10), where

$$dY_t = -bY_t dt + \int_{\mathcal{Z}} a(\zeta) N^H(dt, d\zeta), \quad t > 0.$$

Here N^H denotes the integer-valued random measure associated with the marked Hawkes process, defined by the predictable thinning

$$N^H(dt, d\zeta) := \int_{\mathbb{R}_{\geq 0}} \mathbf{1}_{[0, \mu(P_{t-}^H) + Y_{t-}]}(r) M(dt, d\zeta, dr),$$

where M is the underlying Poisson random measure on $\mathbb{R}_{\geq 0} \times \mathcal{Z} \times \mathbb{R}_{\geq 0}$. The predictable compensator of N^H is

$$\Lambda^H(dt, d\zeta) = (\mu(P_{t-}^H) + Y_{t-}) \nu(d\zeta) dt.$$

2. **Well-posedness of the Poisson-driven model (see Section 4 in [12]):** *Let $(k, p) \in \mathcal{S}_P$ be given. For any admissible control $\alpha \in \mathcal{A}_P(k, p)$, there exists a unique $\mathbb{F}$ -adapted process*

$$X^P = (K_t^P, P_t^P)_{t \geq 0}$$

solving

$$dK_t^P = b^{\text{cap}}(K_t^P, P_t^P, \alpha_t) dt - \int_{\mathcal{Z}} (1 - \omega(K_{t-}^P, P_{t-}^P, \zeta)) K_{t-}^P N^P(dt, d\zeta), \quad (4.12)$$

$$dP_t^P = b^{\text{pol}}(K_t^P, P_t^P, \alpha_t) dt + \sigma P_t^P dW_t, \quad (4.13)$$

$$(K_0^P, P_0^P) = (k, p) \in \mathcal{S}_P. \quad (4.14)$$

Here N^P denotes the integer-valued random measure associated with the Poisson disaster process, defined by the predictable thinning

$$N^P(dt, d\zeta) := \int_{\mathbb{R}_{\geq 0}} \mathbf{1}_{\{r \leq \mu(P_{t-}^P)\}} M(dt, d\zeta, dr),$$

where M is the underlying Poisson random measure on $\mathbb{R}_{\geq 0} \times \mathcal{Z} \times \mathbb{R}_{\geq 0}$. The predictable compensator of N^P is

$$\Lambda^P(dt, d\zeta) = \mu(P_{t-}^P) \nu(d\zeta) dt.$$

3. **Approximation property:** Let V_H^η denote the value function of the Hawkes model with scaled excitation

$$a_\eta(\zeta) := \eta a(\zeta).$$

Then, for all sufficiently small $\eta > 0$,

$$|V_H^\eta(k, p, y_\eta) - V_P(k, p)| \leq \frac{\psi^{-\varepsilon}}{1 - \varepsilon} A_* \eta, \quad (4.15)$$

where $A_* > 0$ is independent of η . Consequently,

$$V_H^\eta(k, p, y_\eta) \longrightarrow V_P(k, p) \quad \text{as } \eta \downarrow 0.$$

5 Hawkes-Poisson approximation

This section is devoted to the proof of Theorem 5.10. The main objective is to quantify the effect of Hawkes self-excitation on the associated infinite-horizon control problem and to show that, when the excitation becomes sufficiently small, the Hawkes value function approaches its Poisson counterpart.

The argument is developed in several stages. We first establish the well-posedness and nonexplosion of the controlled Hawkes dynamics. We then study the asymptotic behaviour of the excitation process and show that, under the subcriticality condition, the contribution of self-excitation vanishes asymptotically. This provides a first probabilistic connection between the Hawkes and Poisson models. Finally, we introduce a small-excitation scaling and derive estimates that allow us to transfer this approximation to the corresponding infinite-horizon value functions.

Throughout this section, we focus on the arguments that are specific to the Hawkes specification. The corresponding Poisson control problem and its well-posedness properties were studied in [12].

5.1 Well-posedness of the Hawkes model

We begin by establishing that the controlled state dynamics are well defined. This is necessary before considering either the asymptotic behaviour of the Hawkes excitation or the associated infinite-horizon control problem. The main difficulty relative to the Poisson model is that the jump intensity is itself stochastic and depends on the endogenous excitation process Y .

The following proposition establishes existence and pathwise uniqueness of the controlled state process and, importantly, excludes the accumulation of disaster times on finite intervals. The proof relies on a Poisson embedding of the marked Hawkes process, followed by localization and uniform finite-horizon moment estimates.

Proposition 5.1 (Existence, pathwise uniqueness and nonexplosion). *Let $T > 0$ be fixed, and let $(C_t, \theta_t)_{t \geq 0}$ be an arbitrary admissible control. Then, on every finite horizon $[0, T]$, there exists a pathwise unique strong solution $(K_t, P_t, Y_t)_{0 \leq t \leq T}$ to the SDE*

$$dK_t = b^{\text{cap}}(K_t, P_t, C_t, \theta_t) dt - \int_{\mathcal{Z}} (1 - \omega(K_{t-}, P_{t-}, \zeta)) K_{t-} N(dt, d\zeta), \quad (5.1)$$

$$dP_t = b^{\text{pol}}(K_t, P_t, C_t, \theta_t) dt + \sigma P_t dW_t, \quad (5.2)$$

$$dY_t = -bY_t dt + \int_{\mathcal{Z}} a(\zeta) N(dt, d\zeta), \quad (5.3)$$

with initial condition $(K_0, P_0, Y_0) = (k, p, y) \in \mathcal{S}_H$. Moreover, for every $T > 0$,

$$N((0, T] \times \mathcal{Z}) < \infty \quad \text{a.s.},$$

Hence, the marked point process is locally finite and nonexplosive, and the global solution is locally square-integrable in time.

Proof. Let us consider the proof divided in four steps:

1. *Poisson embedding and local construction.* We first construct the marked point process by means of a Poisson embedding. Given a nonnegative predictable process

$$\lambda_t = \mu(P_{t-}) + Y_{t-},$$

we define

$$N(dt, d\zeta) = \int_{\mathbb{R}_{\geq 0}} \mathbf{1}_{\{r \leq \lambda_t\}} M(dt, d\zeta, dr). \quad (5.4)$$

By the standard Poisson embedding, or thinning, construction for point processes with predictable stochastic intensity, the random measure N defined by (5.4) has predictable compensator

$$\Lambda(dt, d\zeta) = \lambda_t \nu(d\zeta) dt = (\mu(P_{t-}) + Y_{t-}) \nu(d\zeta) dt \quad (5.5)$$

See, for instance, the standard theory of point processes and Poisson embedding in [2]; the same construction is used in the classical theory of Hawkes processes in [3]. We next localize the coupled system. Since

$$\mathcal{S}_H = (0, \infty) \times (0, \infty) \times [0, \infty),$$

we define, for $n \in \mathbb{N}$,

$$D_n := \left[\frac{1}{n}, n \right] \times \left[\frac{1}{n}, n \right] \times [0, n]. \quad (5.6)$$

Then

$$D_n \subset \mathcal{S}_H, \quad D_n \subset D_{n+1}, \quad \bigcup_{n \geq 1} D_n = \mathcal{S}_H.$$

By the assumed local Lipschitz continuity of b^{cap} , b^{pol} , μ , and ω , their restrictions to D_n satisfy Lipschitz estimates with constants depending on n . We extend these restrictions outside D_n to globally Lipschitz coefficients, denoted by

$$b_n^{\text{cap}}, \quad b_n^{\text{pol}}, \quad \mu_n, \quad \omega_n,$$

which coincide with the original coefficients on D_n . We also truncate the stochastic intensity by setting

$$\lambda_t^{(n)} := \left(\mu_n(P_{t-}^{(n)}) + Y_{t-}^{(n)} \right) \wedge n. \quad (5.7)$$

The corresponding marked counting measure is defined by

$$N^{(n)}(dt, d\zeta) = \int_{\mathbb{R}_{\geq 0}} \mathbf{1}_{\{r \leq \lambda_t^{(n)}\}} M(dt, d\zeta, dr). \quad (5.8)$$

In particular, $0 \leq \lambda_t^{(n)} \leq n$, so the jump intensity of the truncated system is bounded.

The truncated state process

$$X_t^{(n)} = (K_t^{(n)}, P_t^{(n)}, Y_t^{(n)})$$

therefore solves

$$dK_t^{(n)} = b_n^{\text{cap}}(K_t^{(n)}, P_t^{(n)}, C_t, \theta_t) dt - \int_{\mathcal{Z}} (1 - \omega_n(K_{t-}^{(n)}, P_{t-}^{(n)}, \zeta)) K_{t-}^{(n)} N^{(n)}(dt, d\zeta), \quad (5.9)$$

$$dP_t^{(n)} = b_n^{\text{pol}}(K_t^{(n)}, P_t^{(n)}, C_t, \theta_t) dt + \sigma P_t^{(n)} dW_t, \quad (5.10)$$

$$dY_t^{(n)} = -bY_t^{(n)} dt + \int_{\mathcal{Z}} a(\zeta) N^{(n)}(dt, d\zeta). \quad (5.11)$$

It is useful to note that, although the map $x \mapsto \mathbf{1}_{\{r \leq \lambda^{(n)}(x)\}}$ is not pointwise Lipschitz for fixed r , it satisfies the standard integrated estimate

$$\int_0^\infty \left| \mathbf{1}_{\{r \leq \lambda^{(n)}(x)\}} - \mathbf{1}_{\{r \leq \lambda^{(n)}(x')\}} \right| dr = \left| \lambda^{(n)}(x) - \lambda^{(n)}(x') \right|. \quad (5.12)$$

Indeed, for any $u, v \geq 0$,

$$\int_0^\infty \left| \mathbf{1}_{\{r \leq u\}} - \mathbf{1}_{\{r \leq v\}} \right| dr = |u - v|.$$

Since μ_n is globally Lipschitz and $(u, y) \mapsto (u + y) \wedge n$ is Lipschitz, the right-hand side of (5.12) is bounded by $A_n |x - x'|$ for some constant $A_n > 0$. Thus the thinning coefficient satisfies the appropriate integrated Lipschitz estimate required in the Brownian-Poisson SDE theory. Together with $a \in L^2(\nu)$ and the global Lipschitz and growth bounds of the truncated coefficients, this places (5.9)–(5.11) within the standard strong well-posedness theory for stochastic differential equations driven jointly by Brownian motion and Poisson random measures. Consequently, for every $n \in \mathbb{N}$, the truncated system admits a pathwise unique strong càdlàg solution $X^{(n)}$ on $[0, T]$; see, for example, [1] and the classical treatment of SDEs driven by Poisson point processes in [7].

Define the exit time

$$\tau_n := \inf \left\{ t \geq 0 : X_t^{(n)} \notin D_n \right\}. \quad (5.13)$$

Since the truncated coefficients coincide with the original coefficients on D_n , the process $X^{(n)}$ solves the original system on $[0, \tau_n)$. Moreover, the truncated systems can be chosen consistently. In particular, if $m > n$, then both $X^{(m)}$ and $X^{(n)}$ solve the same stochastic equation as long as the state remains in D_n . Pathwise uniqueness therefore yields

$$X_t^{(m)} = X_t^{(n)}, \quad 0 \leq t < \tau_n \quad \text{a.s.}$$

Hence the family $\{X^{(n)}\}_{n \geq 1}$ defines a pathwise unique maximal local solution X on

$$[0, \tau_\infty), \quad \tau_\infty := \lim_{n \rightarrow \infty} \tau_n.$$

It remains to show that

$$\tau_\infty = \infty \quad \text{a.s.},$$

which is established below by deriving uniform moment bounds and ruling out explosion on finite time intervals.

2. *Moment estimate for the Hawkes excitation.* We set

$$\bar{a}_1 := \int_{\mathcal{Z}} a(\zeta) \nu(d\zeta), \quad \bar{a}_2 := \int_{\mathcal{Z}} a(\zeta)^2 \nu(d\zeta),$$

which are finite by assumption (4.3). On $[0, \tau_n)$, the localized process coincides with the original process and therefore

$$dY_t = -bY_t dt + \int_{\mathcal{Z}} a(\zeta) N(dt, d\zeta).$$

Applying Itô's formula for semimartingales with jumps to $f(y) = y^2$ yields

$$\begin{aligned} dY_t^2 &= -2bY_t^2 dt + \int_{\mathcal{Z}} [(Y_{t-} + a(\zeta))^2 - Y_{t-}^2] N(dt, d\zeta) \\ &= -2bY_t^2 dt + \int_{\mathcal{Z}} [2Y_{t-}a(\zeta) + a(\zeta)^2] N(dt, d\zeta). \end{aligned} \quad (5.14)$$

For the Itô formula for jump semimartingales, see, for example, [10] or [1].

Stopping at $t \wedge \tau_n$, taking expectations, and applying the compensation formula for integer-valued random measures with predictable compensator (5.5), we obtain

$$\mathbb{E}[Y_{t \wedge \tau_n}^2] = y^2 - 2b \mathbb{E} \int_0^{t \wedge \tau_n} Y_s^2 ds + \mathbb{E} \int_0^{t \wedge \tau_n} (\mu(P_s) + Y_s)(2\bar{a}_1 Y_s + \bar{a}_2) ds. \quad (5.15)$$

Here P_{s-} since P is continuous, while $Y_{s-} = Y_s$ for Lebesgue-a.e. s . By the linear-growth assumption on μ , (4.2), and Young's inequality, there exists a constant $A > 0$, independent of n , such that

$$(\mu(p) + y)(2\bar{a}_1 y + \bar{a}_2) \leq A(1 + p^2 + y^2), \quad p, y \geq 0.$$

Consequently,

$$\mathbb{E}[Y_{t \wedge \tau_n}^2] \leq y^2 + A \int_0^t (1 + \mathbb{E}[P_{s \wedge \tau_n}^2] + \mathbb{E}[Y_{s \wedge \tau_n}^2]) ds. \quad (5.16)$$

For the subsequent nonexplosion argument it is convenient to obtain a maximal estimate. Since $Y \geq 0$, the integral representation

$$Y_t = ye^{-bt} + \int_{(0,t] \times \mathcal{Z}} e^{-b(t-s)} a(\zeta) N(ds, d\zeta) \quad (5.17)$$

implies, for $0 \leq u \leq t \wedge \tau_n$,

$$Y_u \leq y + \int_{(0,u] \times \mathcal{Z}} a(\zeta) N(ds, d\zeta).$$

Decomposing the last integral into its compensated martingale part and its compensator, and applying the Burkholder-Davis-Gundy inequality for purely discontinuous martingales together with the preceding moment bounds, we obtain, for some constant $A_T > 0$ independent of n ,

$$\mathbb{E} \left[\sup_{0 \leq s \leq T \wedge \tau_n} Y_s^2 \right] \leq A_T \left[1 + y^2 + \int_0^T \mathbb{E} (P_{s \wedge \tau_n}^2 + Y_{s \wedge \tau_n}^2) ds \right]. \quad (5.18)$$

3. *Moment estimates for capital and pollution.* We first consider the capital process. At a jump time t with mark ζ ,

$$K_t = \omega(K_{t-}, P_{t-}, \zeta) K_{t-}.$$

Since $0 < \omega(k, p, \zeta) \leq 1$, we have

$$K_t^2 - K_{t-}^2 = (\omega(K_{t-}, P_{t-}, \zeta)^2 - 1) K_{t-}^2 \leq 0.$$

Thus disaster jumps cannot increase K^2 . Applying the Itô formula for jump semimartingales to K^2 on $[0, t \wedge \tau_n]$ therefore gives

$$\begin{aligned} K_{t \wedge \tau_n}^2 &= k^2 + 2 \int_0^{t \wedge \tau_n} K_s b^{\text{cap}}(K_s, P_s, C_s, \theta_s) ds \\ &\quad + \int_{(0, t \wedge \tau_n] \times \mathcal{Z}} [\omega(K_{s-}, P_{s-}, \zeta)^2 - 1] K_{s-}^2 N(ds, d\zeta) \\ &\leq k^2 + 2 \int_0^{t \wedge \tau_n} K_s b^{\text{cap}}(K_s, P_s, C_s, \theta_s) ds. \end{aligned} \quad (5.19)$$

Using the linear-growth assumption on b^{cap} and Young's inequality, there exists $A > 0$, independent of n , such that

$$\mathbb{E} \left[\sup_{0 \leq u \leq t \wedge \tau_n} K_u^2 \right] \leq A \left[k^2 + \int_0^t \left(1 + \mathbb{E} \left[\sup_{0 \leq r \leq s \wedge \tau_n} (K_r^2 + P_r^2) \right] + \mathbb{E}[C_s^2] \right) ds \right]. \quad (5.20)$$

For the pollution process, Itô's formula yields

$$P_{t \wedge \tau_n}^2 = p^2 + 2 \int_0^{t \wedge \tau_n} P_s b^{\text{pol}}(K_s, P_s, C_s, \theta_s) ds + \sigma^2 \int_0^{t \wedge \tau_n} P_s^2 ds + 2\sigma \int_0^{t \wedge \tau_n} P_s^2 dW_s. \quad (5.21)$$

Taking the supremum, expectations, and applying the Burkholder-Davis-Gundy inequality to the continuous local martingale term, followed by Young's inequality, gives

$$\mathbb{E} \left[\sup_{0 \leq u \leq t \wedge \tau_n} P_u^2 \right] \leq A \left[p^2 + \int_0^t \left(1 + \mathbb{E} \left[\sup_{0 \leq r \leq s \wedge \tau_n} (K_r^2 + P_r^2) \right] + \mathbb{E}[C_s^2] \right) ds \right], \quad (5.22)$$

where $A > 0$ may depend on T , σ , and the growth constants, but not on n . For the Burkholder-Davis-Gundy inequality, see, for example, [11].

Combining (5.18), (5.20), and (5.22), and applying Gronwall's inequality, we obtain

$$\mathbb{E} \left[\sup_{0 \leq t \leq T \wedge \tau_n} (K_t^2 + P_t^2 + Y_t^2) \right] \leq A_T \left(1 + k^2 + p^2 + y^2 + \mathbb{E} \int_0^T C_t^2 dt \right), \quad (5.23)$$

where $A_T < \infty$ is independent of n .

4. *Nonexplosion, preservation of the state space, and global continuation.* Fix $T < \infty$. We first show that the marked point process cannot have infinitely many jumps before maximal lifetime. For every $n \in \mathbb{N}$, the compensation formula applied to the stopped counting measure yields

$$\begin{aligned} \mathbb{E} [N((0, T \wedge \tau_n] \times \mathcal{Z})] &= \mathbb{E} \int_0^{T \wedge \tau_n} (\mu(P_{t-}) + Y_{t-}) dt \\ &= \mathbb{E} \int_0^{T \wedge \tau_n} (\mu(P_t) + Y_t) dt. \end{aligned} \quad (5.24)$$

By the linear-growth assumption on μ (4.2), Cauchy-Schwarz, and (5.23), there exists $B_T < \infty$, independent of n , such that

$$\sup_{n \geq 1} \mathbb{E} [N((0, T \wedge \tau_n] \times \mathcal{Z})] \leq B_T. \quad (5.25)$$

Since $\tau_n \uparrow \tau_\infty$, the random variables $N((0, T \wedge \tau_n] \times \mathcal{Z})$ increase to $N((0, T \wedge \tau_\infty) \times \mathcal{Z})$. Therefore, by the monotone convergence theorem,

$$\mathbb{E} [N((0, T \wedge \tau_\infty) \times \mathcal{Z})] = \lim_{n \rightarrow \infty} \mathbb{E} [N((0, T \wedge \tau_n] \times \mathcal{Z})] \leq B_T < \infty. \quad (5.26)$$

Consequently,

$$N((0, T \wedge \tau_\infty) \times \mathcal{Z}) < \infty \quad \text{a.s.} \quad (5.27)$$

Thus there can be no accumulation of disaster times strictly before, or approaching, a finite maximal lifetime.

We next rule out escape of the state variables to $+\infty$. For $R > 0$, we define

$$\sigma_R := \inf \{t < \tau_\infty : K_t + P_t + Y_t \geq R\}.$$

Then, for every n sufficiently large relative to R ,

$$\{\sigma_R \leq T, \sigma_R < \tau_n\} \subseteq \left\{ \sup_{0 \leq t \leq T \wedge \tau_n} (K_t + P_t + Y_t) \geq R \right\}.$$

Hence, by Markov's inequality,

$$\begin{aligned} \mathbb{P}(\sigma_R \leq T, \sigma_R < \tau_n) &\leq \frac{1}{R^2} \mathbb{E} \left[\sup_{0 \leq t \leq T \wedge \tau_n} (K_t + P_t + Y_t)^2 \right] \\ &\leq \frac{3}{R^2} \mathbb{E} \left[\sup_{0 \leq t \leq T \wedge \tau_n} (K_t^2 + P_t^2 + Y_t^2) \right]. \end{aligned} \quad (5.28)$$

Using (5.23) and then letting $n \rightarrow \infty$ gives

$$\mathbb{P}(\sigma_R \leq T) \leq \frac{3A_T}{R^2} \left(1 + k^2 + p^2 + y^2 + \mathbb{E} \int_0^T C_t^2 dt \right). \quad (5.29)$$

Letting $R \rightarrow \infty$ yields

$$\mathbb{P} \left(\sup_{0 \leq t < T \wedge \tau_\infty} (K_t + P_t + Y_t) = \infty \right) = 0.$$

Thus the maximal local solution cannot cease to exist because of escape to $+\infty$ in finite time. It remains to exclude a finite exit through the lower boundary of $\mathcal{S}_H$. By (5.27), on every finite time interval before τ_∞ there are almost surely only finitely many disaster times. At every such disaster time, $K_t = \omega(K_{t-}, P_{t-}, \zeta)K_{t-}$, and the assumption (4.6) implies that a strictly positive value of K remains strictly positive after the jump. The process P does not jump.

Suppose now, for contradiction, that for some $T < \infty$

$$\mathbb{P}(\tau_\infty \leq T) > 0.$$

On the event $\{\tau_\infty \leq T\}$, escape to $+\infty$ has already been ruled out, and (5.27) rules out accumulation of disaster times. Hence the only remaining possible obstruction to continuation would be attainment of the lower boundary $K = 0$ or $P = 0$. This is excluded by Assumption 4.2(e) and the fact that each jump preserves strict positivity. We obtain a contradiction. Therefore,

$$\tau_\infty = \infty \quad \text{a.s.} \quad (5.30)$$

Hence the maximal local solution extends uniquely to a strong solution on the whole interval $[0, \infty)$.

We may now remove the stopping from the jump-count estimate. Since $\tau_\infty = \infty$ a.s., for every finite T ,

$$T \wedge \tau_n \uparrow T \quad \text{a.s.}$$

and therefore, by monotone convergence,

$$\mathbb{E}[N((0, T] \times \mathcal{Z})] = \mathbb{E} \left[\int_0^T (\mu(P_t) + Y_t) dt \right] < \infty. \quad (5.31)$$

Since $N((0, T] \times \mathcal{Z})$ is a nonnegative integer-valued random variable, possibly taking the value $+\infty$, finite expectation implies

$$N((0, T] \times \mathcal{Z}) < \infty \quad \text{a.s.}$$

Thus the marked point process is locally finite and nonexplosive.

Finally, global pathwise uniqueness follows from pathwise uniqueness of the localized equations. Indeed, if X and $\tilde{X}$ are two global strong solutions driven by the same Brownian motion and the same underlying Poisson random measure with the same initial condition and control, then for every n their restrictions up to the corresponding localization time solve the same localized equation. Pathwise uniqueness gives equality up to each localization time, and since the localization times increase to infinity almost surely,

$$X_t = \tilde{X}_t \quad \text{for every } t \geq 0, \quad \text{a.s.}$$

Finally, since $\tau_\infty = \infty$ a.s., Fatou's lemma applied to (5.23) gives, for every $T < \infty$,

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} (K_t^2 + P_t^2 + Y_t^2) \right] \leq A_T \left(1 + k^2 + p^2 + y^2 + \mathbb{E} \int_0^T C_t^2 dt \right) < \infty. \quad (5.32)$$

Consequently,

$$\mathbb{E} \left[\int_0^T (K_t^2 + P_t^2 + Y_t^2) dt \right] \leq T \mathbb{E} \left[\sup_{0 \leq t \leq T} (K_t^2 + P_t^2 + Y_t^2) \right] < \infty. \quad (5.33)$$

Thus the global solution is locally square-integrable in time.

□

Remark 5.2. The term *global well-posedness* means that the system admits a pathwise unique strong solution defined on the whole time interval $[0, \infty)$; equivalently, the maximal lifetime satisfies $\tau_\infty = \infty$ a.s. This should be distinguished from global square integrability of the state process. The moment estimates established in (5.33) imply local square integrability in time, but no claim is made here that

$$\mathbb{E} \left[\int_0^\infty (K_t^2 + P_t^2 + Y_t^2) dt \right] < \infty,$$

which is an additional integrability property and is not part of the global well-posedness assertion.

Having established that the Hawkes-driven system admits a unique global strong solution, we now turn to the relation between the Hawkes and Poisson specifications. Recall that the Hawkes intensity can be decomposed as in (2.4). Thus the only difference at the level of the disaster intensity is the additional excitation term Y . It is therefore natural to first study the long-run behaviour of this component. If Y becomes negligible, then the Hawkes intensity approaches the pollution-dependent Poisson intensity $\mu(P_{t-})$.

5.2 Asymptotic Poisson approximation

The purpose of this subsection is to make the preceding intuition precise. We first establish decay and integrability properties of the excitation process. These properties are then translated into approximation results for the compensator of the marked point process, first on finite time windows and subsequently on the infinite tail.

Theorem 5.3 (Asymptotic decay of the Hawkes excitation). *Assume the hypotheses of Proposition 5.1. Set*

$$\bar{a} := \int_{\mathcal{Z}} a(\zeta) \nu(d\zeta),$$

and suppose that the marked Hawkes process is subcritical, $\bar{a} < b$. Assume further that

$$\int_0^\infty \mathbb{E}[\mu(P_t)] dt < \infty. \quad (5.34)$$

Then

$$\mathbb{E}[Y_t] \longrightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (5.35)$$

Moreover,

$$\int_0^\infty \mathbb{E}[Y_t] dt = \frac{y + \bar{a} \int_0^\infty \mathbb{E}[\mu(P_t)] dt}{b - \bar{a}} < \infty. \quad (5.36)$$

In particular, $Y_t \longrightarrow 0$ in L^1 , i.e. the Hawkes self-excitation vanishes asymptotically in L^1 . However, unless $Y_0 = 0$ and no jump with positive excitation ever occurs, the process Y_t does not reach zero at any finite time.

Proof. From (5.3), we obtain, after integration,

$$Y_t = y - b \int_0^t Y_s ds + \int_{(0,t] \times \mathcal{Z}} a(\zeta) N(ds, d\zeta).$$

Taking expectations and applying the compensation formula gives

$$\mathbb{E}[Y_t] = y - b \int_0^t \mathbb{E}[Y_s] ds + \mathbb{E} \int_0^t \int_{\mathcal{Z}} a(\zeta) (\mu(P_{s-}) + Y_{s-}) \nu(d\zeta) ds.$$

Since P is continuous and $Y_{s-} = Y_s$ for Lebesgue-a.e. s , this becomes

$$\begin{aligned} \mathbb{E}[Y_t] &= y - b \int_0^t \mathbb{E}[Y_s] ds + \bar{a} \int_0^t \mathbb{E}[\mu(P_s) + Y_s] ds \\ &= y + \int_0^t [-(b - \bar{a})\mathbb{E}[Y_s] + \bar{a} \mathbb{E}[\mu(P_s)]] ds. \end{aligned}$$

Set $m(t) := \mathbb{E}[Y_t]$. Then m is absolutely continuous and satisfies, for almost every $t \geq 0$,

$$m'(t) = -(b - \bar{a})m(t) + \bar{a} \mathbb{E}[\mu(P_t)], \quad m(0) = y. \quad (5.37)$$

By the variation-of-constants formula,

$$m(t) = e^{-(b-\bar{a})t} y + \bar{a} \int_0^t e^{-(b-\bar{a})(t-s)} \mathbb{E}[\mu(P_s)] ds.$$

Let

$$f(s) := \mathbb{E}[\mu(P_s)], \quad g(s) := e^{-(b-\bar{a})s} \mathbf{1}_{\{s \geq 0\}}.$$

By assumption, $f \in L^1(\mathbb{R}_{\geq 0})$, and, since $b - \bar{a} > 0$, $g \in L^1(\mathbb{R}_{\geq 0})$. Hence

$$(g * f)(t) = \int_0^t e^{-(b-\bar{a})(t-s)} f(s) ds \longrightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Therefore

$$m(t) = \mathbb{E}[Y_t] \longrightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (5.38)$$

Integrating (5.37) over $[0, T]$ gives

$$m(T) - y = -(b - \bar{a}) \int_0^T m(t) dt + \bar{a} \int_0^T \mathbb{E}[\mu(P_t)] dt.$$

Letting $T \rightarrow \infty$ and using (5.38), we obtain

$$\int_0^\infty m(t) dt = \frac{y + \bar{a} \int_0^\infty \mathbb{E}[\mu(P_t)] dt}{b - \bar{a}}.$$

Since $Y_t \geq 0$,

$$\|Y_t\|_{L^1} = \mathbb{E}[Y_t] \longrightarrow 0,$$

which proves the L^1 -convergence.

Finally, using the integral representation (5.17), all terms on the right-hand side are nonnegative. If $y > 0$, then $Y_t \geq ye^{-bt} > 0$ for every $t < \infty$. Likewise, if a jump occurs at time τ with mark ζ such that $a(\zeta) > 0$, then

$$Y_t \geq e^{-b(t-\tau)}a(\zeta) > 0 \quad \text{for every finite } t > \tau.$$

Thus finite-time extinction of Y does not occur under the exponential-kernel specification. $\square$

The preceding theorem gives the basic asymptotic estimate required for the approximation argument. In particular, the excitation vanishes in mean and its expected cumulative contribution over the infinite horizon is finite. Two immediate consequences are useful. First, the L^1 convergence implies convergence in probability. Second, since Y is exactly the difference between the Hawkes intensity and its Poisson baseline, the same estimate directly controls the excess disaster intensity.

Corollary 5.4 (Convergence in probability). *Under the assumptions of Theorem 5.3,*

$$Y_t \longrightarrow 0 \quad \text{in probability as } t \rightarrow \infty.$$

Proof. Since $Y_t \geq 0$, Markov's inequality yields, for every $\varepsilon > 0$,

$$\mathbb{P}(|Y_t| > \varepsilon) = \mathbb{P}(Y_t > \varepsilon) \leq \frac{\mathbb{E}[Y_t]}{\varepsilon}.$$

By Theorem 5.3,

$$\mathbb{E}[Y_t] \longrightarrow 0 \quad \text{as } t \rightarrow \infty,$$

and therefore

$$\mathbb{P}(Y_t > \varepsilon) \longrightarrow 0 \quad \text{for every } \varepsilon > 0, .$$

$\square$

Convergence of the state variable Y also has a direct interpretation in terms of the disaster mechanism. Indeed, Y_{t-} is precisely the self-exciting component of the Hawkes intensity. The following corollary therefore reformulates the preceding asymptotic result at the level of the stochastic intensity itself.

Corollary 5.5 (Asymptotic decay of the excess disaster intensity). *Under the assumptions of Theorem 5.3,*

$$\lambda_t - \mu(P_{t-}) = Y_{t-},$$

and consequently

$$\mathbb{E}[|\lambda_t - \mu(P_{t-})|] \longrightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Thus the excess disaster intensity generated by Hawkes self-excitation vanishes asymptotically in L^1 .

Proof. Since $Y_{t-} \geq 0$, it follows that

$$|\lambda_t - \mu(P_{t-})| = Y_{t-},$$

and hence

$$\mathbb{E}[|\lambda_t - \mu(P_{t-})|] = \mathbb{E}[Y_{t-}]. \quad (5.39)$$

Since

$$\Delta Y_t = \int_{\mathcal{Z}} a(\zeta) N(\{t\}, d\zeta),$$

we have $Y_t - Y_{t-} = \Delta Y_t$. Moreover, the marked point process N has predictable compensator (5.5), which is absolutely continuous with respect to Lebesgue measure in time. Consequently, for every fixed deterministic $t > 0$,

$$\mathbb{P}(N(\{t\} \times \mathcal{Z}) > 0) = 0.$$

Thus $\Delta Y_t = 0$ a.s. for every fixed t , and therefore $Y_{t-} = Y_t$ a.s. In particular, $\mathbb{E}[Y_{t-}] = \mathbb{E}[Y_t]$. Combining the result (5.35) of Theorem 5.3 with (5.39), we obtain

$$\mathbb{E}[|\lambda_t - \mu(P_{t-})|] = \mathbb{E}[Y_t] \longrightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Hence

$$\lambda_t - \mu(P_{t-}) \longrightarrow 0 \quad \text{in } L^1,$$

which proves that the excess disaster intensity generated by Hawkes self-excitation vanishes asymptotically. $\square$

The previous results concern the pointwise difference between the Hawkes and Poisson intensities. For the subsequent control argument, it is useful to translate this pointwise decay into a statement about the corresponding random measures. We therefore compare the expected integrals of a test function against the Hawkes counting measure with those obtained from the pollution-dependent Poisson compensator.

We first perform this comparison on a fixed time window of length $H > 0$. This formulation isolates the contribution of the excitation term and provides an explicit approximation error.

Proposition 5.6 (Finite-window Poisson approximation of the Hawkes intensity). *Assume the hypotheses of Theorem 5.3, so that*

$$\mathbb{E}[Y_t] \longrightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Let $H > 0$ be fixed and let

$$\phi : \mathbb{R}_{\geq 0} \times \mathcal{Z} \longrightarrow \mathbb{R}$$

be continuous and bounded. Then

$$\begin{aligned} & \left| \mathbb{E} \left[\int_{(T, T+H] \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] - \mathbb{E} \left[\int_T^{T+H} \int_{\mathcal{Z}} \phi(t, \zeta) \mu(P_{t-}) \nu(d\zeta) dt \right] \right| \\ & \leq \|\phi\|_{\infty} \int_T^{T+H} \mathbb{E}[Y_t] dt. \end{aligned} \quad (5.40)$$

Consequently,

$$\lim_{T \rightarrow \infty} \left| \mathbb{E} \left[\int_{(T, T+H] \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] - \mathbb{E} \left[\int_T^{T+H} \int_{\mathcal{Z}} \phi(t, \zeta) \mu(P_{t-}) \nu(d\zeta) dt \right] \right| = 0.$$

More precisely, for every $\varepsilon > 0$, there exists $T_{\varepsilon} < \infty$ such that, for every $T \geq T_{\varepsilon}$,

$$\left| \mathbb{E} \left[\int_{(T, T+H] \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] - \mathbb{E} \left[\int_T^{T+H} \int_{\mathcal{Z}} \phi(t, \zeta) \mu(P_{t-}) \nu(d\zeta) dt \right] \right| \leq \|\phi\|_{\infty} H \varepsilon.$$

Proof. By the compensation formula and the predictable compensator (5.5), we have

$$\mathbb{E} \left[\int_{(T, T+H] \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] = \mathbb{E} \left[\int_T^{T+H} \int_{\mathcal{Z}} \phi(t, \zeta) (\mu(P_{t-}) + Y_{t-}) \nu(d\zeta) dt \right].$$

Subtracting the pollution-dependent baseline contribution gives

$$\begin{aligned} & \mathbb{E} \left[\int_{(T, T+H] \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] - \mathbb{E} \left[\int_T^{T+H} \int_{\mathcal{Z}} \phi(t, \zeta) \mu(P_{t-}) \nu(d\zeta) dt \right] \\ &= \mathbb{E} \left[\int_T^{T+H} \int_{\mathcal{Z}} \phi(t, \zeta) Y_{t-} \nu(d\zeta) dt \right] \\ &\leq \mathbb{E} \left[\int_T^{T+H} \int_{\mathcal{Z}} |\phi(t, \zeta)| Y_{t-} \nu(d\zeta) dt \right] \leq \|\phi\|_{\infty} \int_T^{T+H} \mathbb{E}[Y_t] dt, \end{aligned}$$

where we used $\nu(\mathcal{Z}) = 1$ and the fact that $Y_{t-} = Y_t$ for Lebesgue-a.e. t .

Since $\mathbb{E}[Y_t] \rightarrow 0$, for every $\varepsilon > 0$ there exists T_{ε} such that

$$\mathbb{E}[Y_t] \leq \varepsilon, \quad t \geq T_{\varepsilon}.$$

Thus, for every $T \geq T_{\varepsilon}$,

$$\int_T^{T+H} \mathbb{E}[Y_t] dt \leq H\varepsilon,$$

which proves the claim. $\square$

The finite-window estimate shows that, sufficiently far in the future, the expected contribution of Hawkes self-excitation over any fixed time interval becomes arbitrarily small. For the infinite-horizon control problem, however, a tail estimate is more directly relevant. The integrability established in Theorem 5.3 allows the preceding argument to be extended from finite windows to the entire remaining time horizon.

Corollary 5.7 (Approximation on the infinite tail). *Under the assumptions of Theorem 5.3, let*

$$\phi : \mathbb{R}_{\geq 0} \times \mathcal{Z} \rightarrow \mathbb{R}$$

be continuous and bounded. Then, for every $T \geq 0$,

$$\left| \mathbb{E} \left[\int_{(T, \infty) \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] - \mathbb{E} \left[\int_T^{\infty} \int_{\mathcal{Z}} \phi(t, \zeta) \mu(P_{t-}) \nu(d\zeta) dt \right] \right| \leq \|\phi\|_{\infty} \int_T^{\infty} \mathbb{E}[Y_t] dt. \quad (5.41)$$

Consequently,

$$\left| \mathbb{E} \left[\int_{(T, \infty) \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] - \mathbb{E} \left[\int_T^{\infty} \int_{\mathcal{Z}} \phi(t, \zeta) \mu(P_{t-}) \nu(d\zeta) dt \right] \right| \rightarrow 0$$

as $T \rightarrow \infty$.

Proof. Since ϕ is continuous and bounded, it is Borel measurable, and hence the deterministic map

$$(\omega, t, \zeta) \mapsto \phi(t, \zeta)$$

defines an admissible predictable integrand for the compensation formula. Moreover, by the assumption of Theorem 5.3, (5.34), and (5.36),

$$\begin{aligned} & \mathbb{E} \left[\int_T^\infty \int_{\mathcal{Z}} |\phi(t, \zeta)| (\mu(P_{t-}) + Y_{t-}) \nu(d\zeta) dt \right] \\ & \leq \|\phi\|_\infty \left(\int_T^\infty \mathbb{E}[\mu(P_t)] dt + \int_T^\infty \mathbb{E}[Y_t] dt \right) < \infty, \end{aligned}$$

where we used $\nu(\mathcal{Z}) = 1$ and the fact that $P_{t-} = P_t$ and $Y_{t-} = Y_t$ for Lebesgue-a.e. t . Thus, the compensation formula yields

$$\mathbb{E} \left[\int_{(T, \infty) \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] = \mathbb{E} \left[\int_T^\infty \int_{\mathcal{Z}} \phi(t, \zeta) (\mu(P_{t-}) + Y_{t-}) \nu(d\zeta) dt \right]. \quad (5.42)$$

Subtracting the pollution-dependent baseline contribution and taking absolute values, we obtain

$$\begin{aligned} & \left| \mathbb{E} \left[\int_{(T, \infty) \times \mathcal{Z}} \phi(t, \zeta) N(dt, d\zeta) \right] - \mathbb{E} \left[\int_T^\infty \int_{\mathcal{Z}} \phi(t, \zeta) \mu(P_{t-}) \nu(d\zeta) dt \right] \right| \\ & \leq \mathbb{E} \left[\int_T^\infty \int_{\mathcal{Z}} |\phi(t, \zeta)| Y_{t-} \nu(d\zeta) dt \right] \\ & \leq \|\phi\|_\infty \int_T^\infty \mathbb{E}[Y_t] dt, \end{aligned}$$

which proves (5.41).

Because $Y_{t-} \geq 0$, Tonelli's theorem yields

$$\mathbb{E} \left[\int_T^\infty \int_{\mathcal{Z}} Y_{t-} \nu(d\zeta) dt \right] = \int_T^\infty \int_{\mathcal{Z}} \mathbb{E}[Y_{t-}] \nu(d\zeta) dt = \int_T^\infty \mathbb{E}[Y_{t-}] dt,$$

where we used $\nu(\mathcal{Z}) = 1$. Since Y is càdlàg, we have $Y_{t-} = Y_t$ for Lebesgue-a.e. t , almost surely. Therefore,

$$\int_T^\infty \mathbb{E}[Y_{t-}] dt = \int_T^\infty \mathbb{E}[Y_t] dt.$$

Combining the preceding estimates proves (5.41).

Finally, by (5.36), the nonnegative function $t \mapsto \mathbb{E}[Y_t]$ is integrable on $\mathbb{R}_{\geq 0}$. Hence

$$\int_T^\infty \mathbb{E}[Y_t] dt \longrightarrow 0 \quad \text{as } T \rightarrow \infty.$$

Therefore, by (5.41), the contribution of the Hawkes self-excitation on the infinite tail vanishes as $T \rightarrow \infty$. $\square$

5.3 Small-excitation approximation of the value function

The preceding results describe an asymptotic approximation in time: under suitable integrability conditions, the self-exciting component of the Hawkes intensity becomes negligible in the distant future. This observation alone, however, does not yield a uniform estimate for the difference between the corresponding infinite-horizon value functions. To obtain such an estimate, we now introduce a small-excitation regime. More precisely, we scale the excitation generated by each disaster mark according to

$$a_\eta(\zeta) = \eta a(\zeta), \quad \eta > 0,$$

and study the limit $\eta \downarrow 0$. In this regime, the Hawkes model approaches the Poisson model directly at the level of the controlled generator. The remaining task is to control the cumulative contribution of the excitation term uniformly over admissible controls.

The first step is to obtain a quantitative estimate for the excitation process under this scaling. Since the generator comparison below involves the product of Y^η with the jump increment of the Poisson value function, an unweighted estimate for Y^η alone is not sufficient. We therefore establish a weighted discounted bound that is uniform both in η and over the admissible controls.

Proposition 5.8 (Uniform weighted small-excitation estimate). *Let $\eta_0 > 0$ and, for each $\eta \in (0, \eta_0]$, define*

$$a_\eta(\zeta) := \eta a(\zeta), \quad \zeta \in \mathcal{Z}.$$

Let

$$X^{H,\eta} = (K^{H,\eta}, P^{H,\eta}, Y^\eta)$$

denote the corresponding controlled Hawkes state process, whose excitation component satisfies

$$dY_t^\eta = -bY_t^\eta dt + \int_{\mathcal{Z}} \eta a(\zeta) N^\eta(dt, d\zeta), \quad Y_0^\eta = y_\eta \geq 0, \quad (5.43)$$

where N^η has predictable compensator

$$(\mu(P_{t-}^{H,\eta}) + Y_{t-}^\eta) \nu(d\zeta) dt. \quad (5.44)$$

Set

$$\bar{a}_1 := \int_{\mathcal{Z}} a(\zeta) \nu(d\zeta) < \infty, \quad \bar{a}_2 := \int_{\mathcal{Z}} a(\zeta)^2 \nu(d\zeta) < \infty.$$

Assume furthermore that

1. There exists $c_y > 0$, independent of η , such that

$$0 \leq y_\eta \leq c_y \eta, \quad 0 < \eta \leq \eta_0. \quad (5.45)$$

2. The family is uniformly subcritical:

$$\eta_0 \bar{a}_1 < b. \quad (5.46)$$

3. There exists $B_\mu < \infty$ such that

$$\sup_{0 < \eta \leq \eta_0} \sup_{\alpha \in \mathcal{A}_H(k, p, y_\eta)} \mathbb{E}_{k, p, y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} \mu(P_t^{H,\eta})^2 dt \right] \leq B_\mu. \quad (5.47)$$

4. For some $q \geq 0$, there exists $B_q < \infty$ such that

$$\sup_{0 < \eta \leq \eta_0} \sup_{\alpha \in \mathcal{A}_H(k, p, y_\eta)} \mathbb{E}_{k, p, y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} \left(1 + (K_t^{H,\eta})^{2q} + (P_t^{H,\eta})^{2q} \right) dt \right] \leq B_q. \quad (5.48)$$

Then there exists a constant $A_* > 0$, independent of $\eta \in (0, \eta_0]$ and of the admissible control, such that

$$\sup_{\alpha \in \mathcal{A}_H(k, p, y_\eta)} \mathbb{E}_{k, p, y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right] \leq A_* \eta. \quad (5.49)$$

In particular,

$$\sup_{\alpha \in \mathcal{A}_H(k, p, y_\eta)} \mathbb{E}_{k, p, y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right] \longrightarrow 0 \quad \text{as } \eta \downarrow 0.$$

Proof. Fix $\eta \in (0, \eta_0]$ and $\alpha \in \mathcal{A}_H(k, p, y_\eta)$. Let us consider the proof divided in two steps:

1. *Uniform discounted L^2 -estimate for the Hawkes excitation.* For notational convenience, throughout this step we write $P_t^\eta := P_t^{H,\eta}$. The excitation process satisfies

$$dY_t^\eta = -bY_t^\eta dt + \int_{\mathcal{Z}} \eta a(\zeta) N^\eta(dt, d\zeta), \quad Y_0^\eta = y_\eta. \quad (5.50)$$

Applying Itô's formula for jump semimartingales to $f(y) = y^2$ gives

$$\begin{aligned} d(Y_t^\eta)^2 &= -2b(Y_t^\eta)^2 dt + \int_{\mathcal{Z}} \left[(Y_{t-}^\eta + \eta a(\zeta))^2 - (Y_{t-}^\eta)^2 \right] N^\eta(dt, d\zeta) \\ &= -2b(Y_t^\eta)^2 dt + \int_{\mathcal{Z}} [2\eta Y_{t-}^\eta a(\zeta) + \eta^2 a(\zeta)^2] N^\eta(dt, d\zeta). \end{aligned} \quad (5.51)$$

Since $t \mapsto e^{-\rho t}$ is continuous and of finite variation, the product rule yields

$$d(e^{-\rho t}(Y_t^\eta)^2) = -(2b + \rho)e^{-\rho t}(Y_t^\eta)^2 dt + e^{-\rho t} \int_{\mathcal{Z}} [2\eta Y_{t-}^\eta a(\zeta) + \eta^2 a(\zeta)^2] N^\eta(dt, d\zeta). \quad (5.52)$$

We now justify the compensation step by localization. Let $(\tau_n)_{n \geq 1}$ be an increasing localizing sequence such that $\tau_n \uparrow \infty$ a.s. and the relevant stopped stochastic integrals are true martingales. Applying (5.52) on $[0, T \wedge \tau_n]$ yields

$$\begin{aligned} e^{-\rho(T \wedge \tau_n)}(Y_{T \wedge \tau_n}^\eta)^2 - y_\eta^2 &= -(2b + \rho) \int_0^{T \wedge \tau_n} e^{-\rho t} (Y_t^\eta)^2 dt \\ &\quad + \int_{(0, T \wedge \tau_n] \times \mathcal{Z}} e^{-\rho t} [2\eta Y_{t-}^\eta a(\zeta) + \eta^2 a(\zeta)^2] N^\eta(dt, d\zeta). \end{aligned} \quad (5.53)$$

Hence, by the compensation formula,

$$\begin{aligned} &\mathbb{E}_{k,p,y_\eta}^\alpha \left[e^{-\rho(T \wedge \tau_n)} (Y_{T \wedge \tau_n}^\eta)^2 \right] - y_\eta^2 \\ &= -(2b + \rho) \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} (Y_t^\eta)^2 dt \right] \\ &\quad + \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} (\mu(P_t^\eta) + Y_t^\eta) (2\eta \bar{a}_1 Y_t^\eta + \eta^2 \bar{a}_2) dt \right]. \end{aligned} \quad (5.54)$$

Here we have used the continuity of P^η and the fact that a càdlàg process coincides with its left-limit process for Lebesgue-a.e. time.

We next estimate the compensator integrand. Expanding, we obtain

$$(\mu(P_t^\eta) + Y_t^\eta) (2\eta \bar{a}_1 Y_t^\eta + \eta^2 \bar{a}_2) = 2\eta \bar{a}_1 \mu(P_t^\eta) Y_t^\eta + 2\eta \bar{a}_1 (Y_t^\eta)^2 + \eta^2 \bar{a}_2 \mu(P_t^\eta) + \eta^2 \bar{a}_2 Y_t^\eta. \quad (5.55)$$

Since (5.46), we have

$$2b + \rho - 2\eta_0 \bar{a}_1 > 0.$$

Choose $\delta > 0$ sufficiently small that

$$\kappa := 2b + \rho - 2\eta_0 \bar{a}_1 - 2\delta > 0. \quad (5.56)$$

Young's inequality yields

$$2\eta \bar{a}_1 \mu(P_t^\eta) Y_t^\eta \leq \delta (Y_t^\eta)^2 + \frac{\eta^2 \bar{a}_1^2}{\delta} \mu(P_t^\eta)^2. \quad (5.57)$$

and

$$\eta^2 \bar{a}_2 Y_t^\eta \leq \delta (Y_t^\eta)^2 + \frac{\eta^4 \bar{a}_2^2}{4\delta}, \quad (5.58)$$

Since $\mu \geq 0$,

$$\eta^2 \bar{a}_2 \mu(P_t^\eta) \leq \frac{\eta^2 \bar{a}_2}{2} (1 + \mu(P_t^\eta)^2). \quad (5.59)$$

Moreover,

$$2\eta \bar{a}_1 (Y_t^\eta)^2 \leq 2\eta_0 \bar{a}_1 (Y_t^\eta)^2.$$

Since $\eta \leq \eta_0$, $\eta^4 \leq \eta_0^2 \eta^2$. Consequently, there exists a constant $A_0 > 0$, depending only on $\eta_0, \bar{a}_1, \bar{a}_2$ and δ , but independent of η, α, t, T , and n , such that

$$(\mu(P_t^\eta) + Y_t^\eta) (2\eta \bar{a}_1 Y_t^\eta + \eta^2 \bar{a}_2) \leq (2\eta_0 \bar{a}_1 + 2\delta) (Y_t^\eta)^2 + A_0 \eta^2 (1 + \mu(P_t^\eta)^2). \quad (5.60)$$

Substituting (5.60) into (5.54) and using (5.56), we obtain

$$\begin{aligned} & \mathbb{E}_{k,p,y_\eta}^\alpha \left[e^{-\rho(T \wedge \tau_n)} (Y_{T \wedge \tau_n}^\eta)^2 \right] \\ & + \kappa \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} (Y_t^\eta)^2 dt \right] \\ & \leq y_\eta^2 + A_0 \eta^2 \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} (1 + \mu(P_t^\eta)^2) dt \right]. \end{aligned} \quad (5.61)$$

Dropping the nonnegative terminal term and using the fact that $T \wedge \tau_n \leq T$ together with the nonnegativity of the integrand, we obtain

$$\kappa \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} (Y_t^\eta)^2 dt \right] \leq y_\eta^2 + A_0 \eta^2 \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^T e^{-\rho t} (1 + \mu(P_t^\eta)^2) dt \right]. \quad (5.62)$$

Using the assumptions (5.45) and (5.47), we obtain

$$\kappa \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} (Y_t^\eta)^2 dt \right] \leq \eta^2 \left[c_y^2 + A_0 \left(\frac{1}{\rho} + B_\mu \right) \right]. \quad (5.63)$$

Since $\tau_n \uparrow \infty$ a.s., for fixed T ,

$$\mathbf{1}_{\{t \leq T \wedge \tau_n\}} \uparrow \mathbf{1}_{\{t \leq T\}} \quad \text{a.s.}$$

Hence the monotone convergence theorem gives

$$\kappa \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^T e^{-\rho t} (Y_t^\eta)^2 dt \right] \leq \eta^2 \left[c_y^2 + A_0 \left(\frac{1}{\rho} + B_\mu \right) \right]. \quad (5.64)$$

Letting $T \rightarrow \infty$ and applying monotone convergence once more yields

$$\mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} (Y_t^\eta)^2 dt \right] \leq A_Y \eta^2, \quad (5.65)$$

where

$$A_Y := \frac{c_y^2 + A_0 (\rho^{-1} + B_\mu)}{\kappa}. \quad (5.66)$$

Since A_Y is independent of the admissible control and of $\eta \in (0, \eta_0]$, we obtain

$$\sup_{0 < \eta \leq \eta_0} \sup_{\alpha \in \mathcal{A}_H(k,p,y_\eta)} \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} (Y_t^\eta)^2 dt \right] \leq A_Y \eta^2. \quad (5.67)$$

2. *Weighted small-excitation estimate.* We define

$$G_t^\eta := 1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q.$$

Since the integrand below is nonnegative, Tonelli's theorem and the Cauchy-Schwarz inequality yield

$$\begin{aligned} & \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta G_t^\eta dt \right] \\ & \leq \left(\mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} (Y_t^\eta)^2 dt \right] \right)^{1/2} \left(\mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} (G_t^\eta)^2 dt \right] \right)^{1/2}. \end{aligned} \quad (5.68)$$

Applying the Cauchy-Schwarz inequality on the product space

$$(\Omega \times \mathbb{R}_{\geq 0}, \mathbb{P}_{k,p,y_\eta}^\alpha \otimes e^{-\rho t} dt),$$

we obtain

$$\begin{aligned} & \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta G_t^\eta dt \right] \\ & \leq \left(\mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} (Y_t^\eta)^2 dt \right] \right)^{1/2} \left(\mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} (G_t^\eta)^2 dt \right] \right)^{1/2}. \end{aligned} \quad (5.69)$$

Using $(x_1 + x_2 + x_3)^2 \leq 3(x_1^2 + x_2^2 + x_3^2)$, we obtain

$$(G_t^\eta)^2 \leq 3 \left(1 + (K_t^{H,\eta})^{2q} + (P_t^{H,\eta})^{2q} \right).$$

Therefore, by (5.48),

$$\sup_{0 < \eta \leq \eta_0} \sup_{\alpha \in \mathcal{A}_H(k,p,y_\eta)} \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} (G_t^\eta)^2 dt \right] \leq 3B_q. \quad (5.70)$$

Combining (5.65), (5.69), and (5.70), we obtain

$$\mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right] \leq \sqrt{A_Y} \eta \sqrt{3B_q}. \quad (5.71)$$

Therefore, setting $A_* := \sqrt{3A_Y B_q}$, and taking the supremum over $\alpha \in \mathcal{A}_H(k,p,y_\eta)$, yields

$$\sup_{\alpha \in \mathcal{A}_H(k,p,y_\eta)} \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right] \leq A_* \eta. \quad (5.72)$$

□

The preceding proposition controls the excitation component Y^η . To use this estimate in the comparison of the Hawkes and Poisson generators, we also need to control the change in the Poisson value function caused by a disaster jump. We recall the explicit Poisson value function obtained in [12]. Its jump increment admits a particularly simple growth estimate.

Lemma 5.9 (Growth of the Poisson jump increment). *Assume that $0 < \varepsilon < 1$ and that*

$$0 < \omega(k,p,\zeta) \leq 1 \quad \text{for all } (k,p,\zeta) \in \mathcal{S}_P \times \mathcal{Z}.$$

Let

$$v(k,p) = \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon} - \frac{x}{1+\beta} p^{1+\beta}, \quad (k,p) \in \mathcal{S}_P,$$

and define

$$\Delta_\zeta v(k, p) := v(\omega(k, p, \zeta)k, p) - v(k, p).$$

Then, for every $(k, p) \in \mathcal{S}_P$,

$$\int_{\mathcal{Z}} |\Delta_\zeta v(k, p)| \nu(d\zeta) \leq \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon}. \quad (5.73)$$

Consequently,

$$\int_{\mathcal{Z}} |\Delta_\zeta v(k, p)| \nu(d\zeta) \leq A_v (1 + k^q + p^q), \quad (k, p) \in \mathcal{S}_P, \quad (5.74)$$

with $q = 1 - \varepsilon$ and $A_v = \frac{\psi^{-\varepsilon}}{1-\varepsilon}$.

Proof. Fix $(k, p) \in \mathcal{S}_P$ and $\zeta \in \mathcal{Z}$. By the definition of v ,

$$\begin{aligned} \Delta_\zeta v(k, p) &= v(\omega(k, p, \zeta)k, p) - v(k, p) \\ &= \frac{\psi^{-\varepsilon}}{1-\varepsilon} (\omega(k, p, \zeta)k)^{1-\varepsilon} - \frac{x}{1+\beta} p^{1+\beta} - \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon} + \frac{x}{1+\beta} p^{1+\beta}. \end{aligned} \quad (5.75)$$

The pollution terms cancel, and therefore

$$\Delta_\zeta v(k, p) = \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon} (\omega(k, p, \zeta)^{1-\varepsilon} - 1). \quad (5.76)$$

Since $0 < \varepsilon < 1$ and $0 < \omega(k, p, \zeta) \leq 1$,

$$-1 \leq \omega(k, p, \zeta)^{1-\varepsilon} - 1 \leq 0,$$

hence

$$|\omega(k, p, \zeta)^{1-\varepsilon} - 1| = 1 - \omega(k, p, \zeta)^{1-\varepsilon} \leq 1.$$

Using (5.76), we obtain

$$|\Delta_\zeta v(k, p)| \leq \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon}.$$

The right-hand side is independent of ζ . Since ν is a probability measure, $\nu(\mathcal{Z}) = 1$, and therefore

$$\int_{\mathcal{Z}} |\Delta_\zeta v(k, p)| \nu(d\zeta) \leq \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon} \int_{\mathcal{Z}} \nu(d\zeta) = \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon}. \quad (5.77)$$

This proves (5.73).

Finally, set $q := 1 - \varepsilon$. Since $k^q \leq 1 + k^q + p^q$, we obtain

$$\int_{\mathcal{Z}} |\Delta_\zeta v(k, p)| \nu(d\zeta) \leq \frac{\psi^{-\varepsilon}}{1-\varepsilon} (1 + k^q + p^q).$$

Thus (5.74) holds with $A_v = \frac{\psi^{-\varepsilon}}{1-\varepsilon}$. □

We now have the two estimates required for the comparison of the control problems. Proposition 5.8 shows that the discounted contribution of the Hawkes excitation is of order η , uniformly over admissible controls, while Lemma 5.9 provides the corresponding growth bound for the jump increment of the Poisson value function.

The key observation is that, when the Poisson value function is extended trivially to the enlarged Hawkes state space, the difference between the Hawkes and Poisson generators is entirely generated by the additional intensity Y^η . Consequently, the preceding two estimates allow this generator discrepancy to be controlled by a term of order η . We can therefore apply the Poisson verification function to the Hawkes dynamics and obtain matching upper and lower bounds for the Hawkes value function. This yields the main approximation result.

Theorem 5.10 (Approximation of the Hawkes value function by the Poisson value function).
For every $\eta \in (0, \eta_0]$, let

$$X^{H,\eta} = (K^{H,\eta}, P^{H,\eta}, Y^\eta)$$

be the Hawkes state process corresponding to the scaled excitation $a_\eta(\zeta) := \eta a(\zeta)$.

The excitation component satisfies

$$dY_t^\eta = -bY_t^\eta dt + \int_{\mathcal{Z}} \eta a(\zeta) N^{H,\eta}(dt, d\zeta), \quad Y_0^\eta = y_\eta,$$

where $N^{H,\eta}$ has predictable compensator $(\mu(P_{t-}^{H,\eta}) + Y_{t-}^\eta) \nu(d\zeta) dt$.

For $\alpha \in \mathcal{A}_H(k, p, y_\eta)$, define

$$J_H^\eta(k, p, y_\eta; \alpha) := \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} U(K_t^{H,\eta}, P_t^{H,\eta}, \alpha_t) dt \right],$$

and

$$V_H^\eta(k, p, y_\eta) := \sup_{\alpha \in \mathcal{A}_H(k,p,y_\eta)} J_H^\eta(k, p, y_\eta; \alpha).$$

Let $v : \mathcal{S}_P \rightarrow \mathbb{R}$ be the verified value function of the corresponding Poisson control problem, so that $v = V_P$. Assume that $0 < \varepsilon < 1$ and

$$v(k, p) = \frac{\psi^{-\varepsilon}}{1-\varepsilon} k^{1-\varepsilon} - \frac{x}{1+\beta} p^{1+\beta}.$$

Suppose that $v \in C^2(\mathcal{S}_P)$ and satisfies the Poisson HJB equation

$$\rho v(k, p) = \sup_{\alpha \in \mathfrak{a}} \{U(k, p, \alpha) + \mathcal{L}_P^\alpha v(k, p)\}, \quad (5.78)$$

where

$$\begin{aligned} \mathcal{L}_P^\alpha v(k, p) &= b^{\text{cap}}(k, p, \alpha) v_k(k, p) + b^{\text{pol}}(k, p, \alpha) v_p(k, p) + \frac{1}{2} \sigma^2 p^2 v_{pp}(k, p) \\ &\quad + \mu(p) \int_{\mathcal{Z}} \Delta_\zeta v(k, p) \nu(d\zeta), \end{aligned} \quad (5.79)$$

with

$$\Delta_\zeta v(k, p) := v(\omega(k, p, \zeta)k, p) - v(k, p).$$

Assume the hypotheses of Proposition 5.8, with $q = 1 - \varepsilon$, and the hypotheses of Lemma 5.9.

Assume furthermore that, for every $\eta \in (0, \eta_0]$ and every $\alpha \in \mathcal{A}_H(k, p, y_\eta)$,

1.

$$\mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} \left| U(K_t^{H,\eta}, P_t^{H,\eta}, \alpha_t) \right| dt \right] < \infty. \quad (5.80)$$

2. The transversality condition holds:

$$\lim_{T \rightarrow \infty} \mathbb{E}_{k,p,y_\eta}^\alpha \left[e^{-\rho T} \left| v(K_T^{H,\eta}, P_T^{H,\eta}) \right| \right] = 0 \quad (5.81)$$

3. For every $T < \infty$, the family

$$\{e^{-\rho \tau} v(K_\tau^{H,\eta}, P_\tau^{H,\eta}) : \tau \leq T \text{ is a stopping time}\} \quad (5.82)$$

is uniformly integrable.

Finally, assume that there exists a Borel measurable selector $\hat{\alpha}_P : \mathcal{S}_P \rightarrow \mathfrak{a}$ attaining the supremum in (5.78), and that, for every $\eta \in (0, \eta_0]$, the feedback control

$$\hat{\alpha}_{P,t}^\eta := \hat{\alpha}_P(K_t^{H,\eta}, P_t^{H,\eta})$$

belongs to $\mathcal{A}_H(k, p, y_\eta)$. Then, for every $\eta \in (0, \eta_0]$,

$$|V_H^\eta(k, p, y_\eta) - V_P(k, p)| \leq \frac{\psi^{-\varepsilon}}{1 - \varepsilon} A_* \eta, \quad (5.83)$$

where A_* is the constant appearing in Proposition 5.8. Consequently,

$$V_H^\eta(k, p, y_\eta) \rightarrow V_P(k, p) \quad \text{as } \eta \downarrow 0.$$

Proof. Set $A_v := \frac{\psi^{-\varepsilon}}{1 - \varepsilon}$ and $q := 1 - \varepsilon$. By Lemma 5.9,

$$\int_{\mathcal{Z}} |\Delta_\zeta v(k, p)| \nu(d\zeta) \leq A_v (1 + k^q + p^q), \quad (k, p) \in \mathcal{S}_P. \quad (5.84)$$

Moreover, Proposition 5.8 gives

$$\sup_{\alpha \in \mathcal{A}_H(k, p, y_\eta)} \mathbb{E}_{k, p, y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right] \leq A_* \eta. \quad (5.85)$$

We divide the remainder of the proof into five steps.

1. *Extension of the Poisson value function to the Hawkes state space.* We define $\tilde{v} : \mathcal{S}_H \rightarrow \mathbb{R}$ by

$$\tilde{v}(k, p, y) := v(k, p).$$

In particular, $\partial_y \tilde{v}(k, p, y) = 0$. For a sufficiently regular test function $\varphi : \mathcal{S}_H \rightarrow \mathbb{R}$, the controlled generator of the Hawkes state process is

$$\begin{aligned} \mathcal{L}_{H,\eta}^\alpha \varphi(k, p, y) &= b^{\text{cap}}(k, p, \alpha) \partial_k \varphi(k, p, y) + b^{\text{pol}}(k, p, \alpha) \partial_p \varphi(k, p, y) \\ &\quad + \frac{1}{2} \sigma^2 p^2 \partial_{pp} \varphi(k, p, y) - b y \partial_y \varphi(k, p, y) \\ &\quad + (\mu(p) + y) \int_{\mathcal{Z}} [\varphi(\omega(k, p, \zeta) k, p, y + \eta a(\zeta)) - \varphi(k, p, y)] \nu(d\zeta). \end{aligned} \quad (5.86)$$

This is the standard generator associated with a jump-diffusion having state-dependent marked jump intensity; see, for example, [1, 8]. Since $\tilde{v}$ is independent of y ,

$$\tilde{v}(\omega(k, p, \zeta) k, p, y + \eta a(\zeta)) - \tilde{v}(k, p, y) = \Delta_\zeta v(k, p).$$

Therefore, recalling the definition of the Poisson generator $\mathcal{L}_P^\alpha$,

$$\mathcal{L}_{H,\eta}^\alpha \tilde{v}(k, p, y) = \mathcal{L}_P^\alpha v(k, p) + y \int_{\mathcal{Z}} \Delta_\zeta v(k, p) \nu(d\zeta). \quad (5.87)$$

Hence, by Lemma 5.9,

$$|\mathcal{L}_{H,\eta}^\alpha \tilde{v}(k, p, y) - \mathcal{L}_P^\alpha v(k, p)| \leq A_v y (1 + k^q + p^q). \quad (5.88)$$

2. *Residual of the Poisson HJB under the Hawkes generator.* Since v satisfies (5.78), for every admissible $\alpha \in \mathfrak{a}$,

$$U(k, p, \alpha) + \mathcal{L}_P^\alpha v(k, p) - \rho v(k, p) \leq 0. \quad (5.89)$$

Combining this inequality with (5.87), we obtain

$$\begin{aligned}
& U(k, p, \alpha) + \mathcal{L}_{H,\eta}^\alpha \tilde{v}(k, p, y) - \rho \tilde{v}(k, p, y) \\
& = U(k, p, \alpha) + \mathcal{L}_P^\alpha v(k, p) - \rho v(k, p) + y \int_{\mathcal{Z}} \Delta_\zeta v(k, p) \nu(d\zeta) \\
& \leq y \int_{\mathcal{Z}} \Delta_\zeta v(k, p) \nu(d\zeta).
\end{aligned}$$

Therefore, using (5.74),

$$U(k, p, \alpha) + \mathcal{L}_{H,\eta}^\alpha \tilde{v}(k, p, y) - \rho \tilde{v}(k, p, y) \leq A_v y (1 + k^q + p^q). \quad (5.90)$$

3. *Localization and Itô's formula.* Fix an arbitrary control $\alpha \in \mathcal{A}_H(k, p, y)$. Let

$$\tilde{N}^{H,\eta}(dt, d\zeta) := N^{H,\eta}(dt, d\zeta) - (\mu(P_{t-}^{H,\eta}) + Y_{t-}^\eta) \nu(d\zeta) dt$$

denote the compensated marked random measure associated with $N^{H,\eta}$. Applying the Itô formula for semimartingales with jumps to $e^{-\rho t} v(K_t^{H,\eta}, P_t^{H,\eta})$ the local martingale terms are

$$M_t^W = \int_0^t e^{-\rho s} \sigma P_s^{H,\eta} v_p(K_s^{H,\eta}, P_s^{H,\eta}) dW_s, \quad (5.91)$$

$$M_t^{N,\eta} = \int_{(0,t] \times \mathcal{Z}} e^{-\rho s} \Delta_\zeta v(K_{s-}^{H,\eta}, P_{s-}^{H,\eta}) \tilde{N}^{H,\eta}(ds, d\zeta). \quad (5.92)$$

The Itô formula for semimartingales with jumps and the associated stochastic integration theory are standard; see [10]. For stochastic integration with Poisson random measures and Lévy-driven SDEs, see also [1]; for the corresponding jump-diffusion control framework, see [8].

Since M^W and $M^{N,\eta}$ are local martingales, there exists an increasing sequence of stopping times $(\tau_n)_{n \geq 1}$ such that

$$\tau_n \uparrow \infty \quad \text{a.s.}$$

such the stopped processes $M_{\cdot \wedge \tau_n}^W, M_{\cdot \wedge \tau_n}^{N,\eta}$ are true martingales; see, for example, [10].

Therefore, for every $T < \infty$, taking expectations in Itô's formula gives

$$\begin{aligned}
& \mathbb{E}_{k,p,y_\eta}^\alpha \left[e^{-\rho(T \wedge \tau_n)} v(K_{T \wedge \tau_n}^{H,\eta}, P_{T \wedge \tau_n}^{H,\eta}) \right] - v(k, p) \\
& = \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} \left[\mathcal{L}_{H,\eta}^{\alpha_t} \tilde{v}(K_{t-}^{H,\eta}, P_{t-}^{H,\eta}, Y_{t-}^\eta) - \rho v(K_{t-}^{H,\eta}, P_{t-}^{H,\eta}) \right] dt \right]. \quad (5.93)
\end{aligned}$$

Using (5.90), we obtain

$$\begin{aligned}
& \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} U(K_t^{H,\eta}, P_t^{H,\eta}, \alpha_t) dt \right] \\
& \leq v(k, p) - \mathbb{E}_{k,p,y_\eta}^\alpha \left[e^{-\rho(T \wedge \tau_n)} v(K_{T \wedge \tau_n}^{H,\eta}, P_{T \wedge \tau_n}^{H,\eta}) \right] \\
& \quad + A_v \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right]. \quad (5.94)
\end{aligned}$$

4. *Removal of localization and passage to the infinite horizon.* Fix $T < \infty$. Since $\tau_n \uparrow \infty$ a.s., we have

$$T \wedge \tau_n \longrightarrow T \quad \text{a.s. as } n \rightarrow \infty.$$

By (5.80), dominated convergence gives

$$\mathbb{E}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} U(K_t^{H,\eta}, P_t^{H,\eta}, \alpha_t) dt \right] \longrightarrow \mathbb{E}^\alpha \left[\int_0^T e^{-\rho t} U(K_t^{H,\eta}, P_t^{H,\eta}, \alpha_t) dt \right].$$

Since the weighted excitation integrand is nonnegative, monotone convergence gives

$$\begin{aligned} & \mathbb{E}^\alpha \left[\int_0^{T \wedge \tau_n} e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right] \\ & \longrightarrow \mathbb{E}^\alpha \left[\int_0^T e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right]. \end{aligned} \quad (5.95)$$

Moreover,

$$e^{-\rho(T \wedge \tau_n)} v(K_{T \wedge \tau_n}^{H,\eta}, P_{T \wedge \tau_n}^{H,\eta}) \longrightarrow e^{-\rho T} v(K_T^{H,\eta}, P_T^{H,\eta}) \quad \text{a.s.,}$$

and the uniform-integrability assumption (5.82) implies convergence in L^1 . Thus, letting $n \rightarrow \infty$ in (5.94),

$$\begin{aligned} & \mathbb{E}^\alpha \left[\int_0^T e^{-\rho t} U(K_t^{H,\eta}, P_t^{H,\eta}, \alpha_t) dt \right] \\ & \leq v(k, p) - \mathbb{E}^\alpha \left[e^{-\rho T} v(K_T^{H,\eta}, P_T^{H,\eta}) \right] \\ & \quad + A_v \mathbb{E}^\alpha \left[\int_0^T e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right]. \end{aligned} \quad (5.96)$$

Letting $T \rightarrow \infty$, the utility term converges by (5.80), the terminal term vanishes by (5.81), and the nonnegative weighted excitation term converges by monotone convergence. Hence

$$J_H^\eta(k, p, y_\eta; \alpha) \leq v(k, p) + A_v \mathbb{E}_{k,p,y_\eta}^\alpha \left[\int_0^\infty e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H,\eta})^q + (P_t^{H,\eta})^q \right) dt \right]. \quad (5.97)$$

Since α was arbitrary, Proposition 5.8 gives

$$V_H^\eta(k, p, y_\eta) \leq v(k, p) + A_v A_* \eta. \quad (5.98)$$

5. *Lower estimate using the Poisson-optimal feedback.* Let $\hat{\alpha}_P : \mathcal{S}_P \rightarrow \mathfrak{a}$ be the measurable feedback attaining the supremum in the Poisson HJB. Thus,

$$U(k, p, \hat{\alpha}_P(k, p)) + \mathcal{L}_P^{\hat{\alpha}_P(k,p)} v(k, p) - \rho v(k, p) = 0. \quad (5.99)$$

The use of measurable maximizing feedbacks and the corresponding verification argument is standard in stochastic control; see [5, 9], and, in the jump-diffusion setting, [8]. By assumption, the feedback control $\hat{\alpha}_{P,t}^\eta = \hat{\alpha}_P(K_t^{H,\eta}, P_t^{H,\eta})$ is admissible for the Hawkes problem. Combining (5.99) and (5.87), we obtain

$$U(k, p, \hat{\alpha}_P(k, p)) + \mathcal{L}_{H,\eta}^{\hat{\alpha}_P(k,p)} \tilde{v}(k, p, y) - \rho \tilde{v}(k, p, y) = y \int_{\mathcal{Z}} \Delta_\zeta v(k, p) \nu(d\zeta). \quad (5.100)$$

Repeating the localization and limiting arguments above under the control $\hat{\alpha}_P^\eta$ gives

$$J_H^\eta(k, p, y_\eta; \hat{\alpha}_P^\eta) - v(k, p) = \mathbb{E}_{k,p,y_\eta}^{\hat{\alpha}_P^\eta} \left[\int_0^\infty e^{-\rho t} Y_t^\eta \int_{\mathcal{Z}} \Delta_\zeta v(K_t^{H,\eta}, P_t^{H,\eta}) \nu(d\zeta) dt \right]. \quad (5.101)$$

By the triangle inequality, Lemma 5.9, and Proposition 5.8,

$$\begin{aligned}
& |J_H^\eta(k, p, y_\eta; \hat{\alpha}_P^\eta) - v(k, p)| \\
& \leq A_v \mathbb{E}_{k, p, y_\eta}^{\hat{\alpha}_P^\eta} \left[\int_0^\infty e^{-\rho t} Y_t^\eta \left(1 + (K_t^{H, \eta})^q + (P_t^{H, \eta})^q \right) dt \right] \\
& \leq A_v A_* \eta.
\end{aligned} \tag{5.102}$$

Hence,

$$J_H^\eta(k, p, y_\eta; \hat{\alpha}_P^\eta) \geq v(k, p) - A_v A_* \eta.$$

Since $\hat{\alpha}_P^\eta$ is admissible for the Hawkes problem,

$$V_H^\eta(k, p, y_\eta) \geq J_H^\eta(k, p, y_\eta; \hat{\alpha}_P^\eta) \geq v(k, p) - A_v A_* \eta.$$

Together with (5.98), this yields

$$v(k, p) - A_v A_* \eta \leq V_H^\eta(k, p, y_\eta) \leq v(k, p) + A_v A_* \eta.$$

Therefore,

$$|V_H^\eta(k, p, y_\eta) - v(k, p)| \leq A_v A_* \eta. \tag{5.103}$$

Finally, since $v = V_P$ and $A_v = \frac{\psi^{-\varepsilon}}{1-\varepsilon}$, we conclude that

$$|V_H^\eta(k, p, y_\eta) - V_P(k, p)| \leq \frac{\psi^{-\varepsilon}}{1-\varepsilon} A_* \eta.$$

Letting $\eta \downarrow 0$ yields

$$V_H^\eta(k, p, y_\eta) \longrightarrow V_P(k, p).$$

□

Theorem 5.10 therefore provides a quantitative justification for using the Poisson control problem as an approximation of the Hawkes problem in the small-excitation regime. The uniform weighted estimate over admissible controls leads to an $O(\eta)$ bound for the difference between the Hawkes and Poisson value functions. In particular, the Poisson value function arises as the limiting value function as the magnitude of Hawkes self-excitation tends to zero.

References

- [1] David Applebaum. *Lévy Processes and Stochastic Calculus*, volume 116 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2 edition, 2009.
- [2] Pierre Brémaud. *Point Processes and Queues: Martingale Dynamics*. Springer-Verlag, New York, 1981.
- [3] Pierre Brémaud and Laurent Massoulié. Stability of Nonlinear Hawkes Processes. *The Annals of Probability*, 24(3):1563–1588, 1996.
- [4] D. J. Daley and D. Vere-Jones. *An Introduction to the Theory of Point Processes. Volume II: General Theory and Structure*. Probability and Its Applications. Springer, New York, NY, 2008.
- [5] Wendell H. Fleming and Halil Mete Soner. *Controlled Markov Processes and Viscosity Solutions*. Springer Science & Business Media, February 2006. Google-Books-ID: 4Bjz2iWmLyQC.

- [6] Alan G. Hawkes and David Oakes. A cluster process representation of a self-exciting process. *Journal of Applied Probability*, 11(3):493–503, 1974.
- [7] Nobuyuki Ikeda and Shinzo Watanabe. *Stochastic Differential Equations and Diffusion Processes*, volume 24 of *North-Holland Mathematical Library*. North-Holland, 2 edition, 1989.
- [8] Bernt Øksendal and Agnès Sulem. *Applied Stochastic Control of Jump Diffusions*. Universitext. Springer International Publishing, Cham, 2019.
- [9] Huyên Pham. *Continuous-time Stochastic Control and Optimization with Financial Applications*, volume 61 of *Stochastic Modelling and Applied Probability*. Springer Berlin Heidelberg, Berlin, Heidelberg, 2009.
- [10] Philip E. Protter. *Stochastic Integration and Differential Equations*. Springer, 2 edition, 2005.
- [11] Daniel Revuz and Marc Yor. *Continuous Martingales and Brownian Motion*. Springer, 3 edition, 1999.
- [12] Daria Sakhandia and Joshué Helí Ricalde-Guerrero. Infinite-Horizon Optimal Control of Jump-Diffusion Models for Pollution-Dependent Disasters, November 2025. arXiv:2511.13568 [math].