

Wasserstein-Barycentric Interaction Fields for Spatial Factor Models: Evidence from Language-Model Representations

Marcus Gawronsky, Chun-Sung Huang
Department of Finance and Tax, University of Cape Town

August 2026

Abstract

Spatial return models take the interaction matrix as given and leave feedback uninterpreted. We construct a bandwidth-free field from firms' language-model article-embedding distributions using target-anchored Wasserstein barycentric reconstruction. A quadratic exposure-adjustment problem maps feedback into a peer-misalignment penalty ratio. For 52 firms, the field, frozen from 2018–2022 news, yields a 2023–2026 penalty ratio of 3.46 (95% interval [2.89, 4.17]) and higher conditional quasi-likelihood than equal-weighted peer support or RBF weighting of the same distances. Joint penalty ratios for the barycentric and news co-mention fields are 2.33 and 0.86; boundary-calibrated tests reject both exclusions ($p = 0.0005$ each).

Keywords: Spatial Exposure Adjustment; Barycentric Interaction Fields; Wasserstein Barycentric Reconstruction; Language-Model Representations; Spatial Autoregression

JEL classification: G12; G11; C21; C58

Introduction

Spatial asset-pricing models organize cross-sectional return dependence through an interaction matrix and estimate its strength by spatial quasi-likelihood. The researcher typically supplies W from geography, industry, supply chains, or news-derived networks and estimates a spatial coefficient ρ conditional on that choice (Fernandez, 2011; Kou et al., 2018; Ge et al., 2023). This approach is powerful once W has been specified, but the spatial equation begins after the economically relevant relations among firms have already been chosen. It therefore leaves two questions outside the model: where the interaction field comes from, and what ρ measures beyond return dependence conditional on that field.

Existing constructions encode economic location in different ways. Point-based spatial models locate a firm in a geographic or characteristic space, whereas network models represent the firm as a node connected by observed links. Text-based finance extends the latter approach by extracting co-coverage and co-mention relations that balance-sheet classifications can miss (Scherbina and Schlusche, 2013; Schwenkler and Zheng, 2020; Ge et al., 2023). Other work summarizes text as a predictive feature, a named factor, or a learned graph (Ben-Rephael et al., 2019; Cong et al., 2024; Son and Lee, 2022). These approaches establish the relevance of text, but an adjacency still leaves primitive what constitutes a link and why its cardinal weight should measure economic interaction.

This paper moves one level upstream of the conventional spatial specification by changing the mathematical object used to represent a firm. A diversified firm does not occupy only one economic location: its products, technologies, supply chains, and information form a

footprint across many positions. Two firms can share a centroid while having very different footprints, much as two restaurant chains with the same average store coordinates can cover different regions. We therefore represent each firm by a probability measure over economic-information positions rather than by one point or averaged text score. A point-valued firm is the special case in which the entire footprint is concentrated at one position; a genuine distribution also retains spread, multimodality, and internal composition.

A language model maps each article to a numerical position, so a firm’s corpus forms an empirical distribution in the embedding space. Once firms are measures, proximity must compare entire footprints rather than only their centroids. In the Kantorovich formulation, quadratic optimal transport considers all feasible couplings of two distributions and selects the one with the smallest average squared displacement. Here the ground cost is squared displacement in a language-model embedding space, so Wasserstein distance measures the least information-space displacement needed to align one observed footprint with another. This least-cost formulation gives the geometry an economic interpretation while making no claim of physical reallocation or literal spatial arbitrage.

Pairwise transport nevertheless does not yet produce an interaction field. A distance says how far two distributions are separated; it does not say how several peers jointly represent a fixed target firm. For each target, optimal transport first aligns its articles separately with the articles of every candidate peer. Holding those target-specific alignments fixed, target-anchored Wasserstein barycentric reconstruction chooses nonnegative unit-sum weights that combine the aligned peer clouds to approximate the target cloud. These distributional spanning weights answer which combination of peers jointly represents the firm, rather than only which single firm lies nearest to it. Repeating the reconstruction across targets produces W^b , the directed, row-stochastic barycentric interaction field. Its simplex and leave-one-out restrictions give nonnegative unit row sums and a zero diagonal without a conventional kernel-bandwidth choice.

The paper’s primary conceptual contribution is to turn this target-anchored Wasserstein barycentric reconstruction into an admissible field for spatial exposure adjustment. Section 4 formalizes the fixed-target construction and its boundary with the unrestricted Wasserstein barycentre; the latter appears only in Section 5 as a representation of cross-sectional dispersion.

Given this field, the paper next asks what the spatial coefficient means. Each firm has a stand-alone exposure ξ_i implied by its own characteristics, and a quadratic adjustment problem balances departure from ξ_i against misalignment with peer exposures. Solving that problem yields a spatial equilibrium in the peer-adjusted exposures B_i with feedback coefficient

$$\rho = \frac{\lambda}{1 + \lambda},$$

where λ is the penalty on peer misalignment relative to the penalty on departing from the stand-alone exposure. This spatial closure derives the lag in exposures rather than assuming it in returns and maps nonnegative adjustment intensity exactly to $0 \leq \rho < 1$. A subsequent return bridge connects the exposure equilibrium to observed returns.

The same closure permits several admissible fields to enter one adjustment problem, each with its own coefficient and adjustment index. The barycentric interaction field and a persistent news co-mention field can therefore represent separate channels rather than rival estimates of one privileged network. A supporting attenuation result then asks how much characteristic-implied latent exposure dispersion survives peer adjustment under an explicit maintained transfer restriction. Related research uses distribution-valued firm characteristics to derive pairwise covariance restrictions and to study portfolio-risk

bounds and allocation (Gawronsky and Huang, 2026a,b). The present paper works at the intervening cross-sectional level: it constructs the field relating firms and traces how stand-alone exposures propagate through that field.

Figure 1 separates what is observed or constructed from what is latent and model-implied, and from what is estimated.




Figure 1: From text geometry to model-implied adjustment. Text distributions construct the frozen barycentric interaction field W^b . In the latent model, firm-specific information generates stand-alone exposure ξ_i , which adjustment transforms into peer-adjusted exposure B_i . The return bridge connects exposure to returns; conditional on the frozen field, QMLE yields a working-model estimate of ρ and hence the adjustment index $\lambda = \rho/(1 - \rho)$. Solid arrows denote construction or estimation; dashed arrows denote model relations.

In the empirical application, we constructed W^b from 2018–2022 text for 52 firms and froze it before estimating the return equation on 885 aligned trading days from 2023 through the incomplete 2026 period. The pooled QMLE implies the working-model adjustment index $\hat{\lambda} = 3.46$. Within the quadratic representation, the fitted penalty on peer misalignment is therefore 3.46 times the penalty on departing from stand-alone exposure. When the persistent news co-mention field enters jointly, the estimated indices are $\hat{\lambda}_B = 2.33$ for the barycentric field and $\hat{\lambda}_N = 0.86$ for the news-link field. Each field improves conditional fit once the other is included.

These estimates have a deliberately conditional interpretation. Freezing the fields before the return window removes mechanical same-sample feedback, but it does not make the text geometry exogenous to omitted industries, technologies, attention, or reporting selection. QMLE estimates working-model return coefficients conditional on the specified fields and return quasi-likelihood; it does not separately identify the peer-adjusted exposures or adjustment costs. Only under the maintained return bridge and quadratic closure does $\hat{\lambda}$ inherit the model’s adjustment interpretation. Accordingly, $\hat{\lambda}$ is a working-model adjustment index, not an observed managerial cost, a geometry-invariant structural parameter, or a causal peer effect.

Section 1 positions the contribution relative to spatial finance, network adjustment, text-based measurement, and optimal transport. Sections 2 and 3 define stand-alone exposures and derive the spatial closure, while Section 4 constructs the barycentric interaction

field and Section 5 derives the attenuation result. Section 6 states the return bridge, identification boundary, data, and estimation design; Section 7 reports the evidence, and Sections 8 and 9 discuss scope and implications.

1 Related Literature

Spatial asset pricing asks how local interaction modifies factor-based pricing once the researcher supplies an interaction matrix W . Across geographic and financial applications, W organizes local dependence before its strength is estimated (Fernandez, 2011; Kou et al., 2018; Ge et al., 2023). In the spatial CAPM and spatial APT of Kou et al. (2018), assets occupy point-valued locations and inverse geographic distance supplies the weights. The resulting spatial multiplier accumulates direct and higher-order interactions, but the asset representation and the rule that converts it into W remain exogenous to the pricing model.

Linear-quadratic network games address a different part of this problem. Costly local complementarity yields best responses summarized by a network resolvent, providing an economic route from individual objectives to aggregate propagation conditional on a given graph (Ballester et al., 2006). The adjustment model in this paper applies that logic one layer below returns: firms trade off departures from their stand-alone factor exposures against misalignment among peer-adjusted exposures, and the spatial coefficient records the relative intensity of that adjustment. This interpretation explains why interaction may arise, but it leaves open which firms should be peers and how their weights should be determined.

The news-implied-network literature supplies one influential answer by replacing geographic location with an observed informational edge. Beginning with economic linkages inferred from news and their relation to return predictability, this line uses sentence-level co-mentions to construct firm networks that trace contagion and aggregate risk (Scherbina and Schlusche, 2013; Schwenkler and Zheng, 2020). Ge et al. (2023) places such a co-mention matrix inside a spatial factor model, and subsequent work studies whether auxiliary network information improves covariance estimation (Ge et al., 2026). These studies expand the meaning of economic proximity beyond literal distance, but the observed or count-weighted graph remains the empirical primitive that supplies W .

A broader text-finance literature usually maps information into a first-order object such as an attention measure, a textual factor, or an embedding-based input to return or stochastic-discount-factor estimation (Ben-Rephael et al., 2019; Cong et al., 2024; Wang et al., 2025). Related learned-network methods map connectivity directly into factor exposures or priced network factors (Son and Lee, 2022; Uddin et al., 2024). Together, these approaches establish that text and learned representations carry economically relevant predictive and pricing information. The distinct question here is not whether text predicts a scalar outcome, but whether the full cross-section of within-firm information can construct the peer field along which exposures adjust.

That question changes the primitive representation of a firm. A diversified firm need not occupy one geographic or characteristic point; its articles can instead be represented by a probability distribution C_i over positions in a maintained information space. This representation preserves dispersion, multimodality, and other differences that a centroid or single embedding suppresses. When every C_i degenerates to a point mass, Wasserstein distance reduces to the underlying point distance, so point-based spatial logic is nested at the level of the representation. For genuinely distribution-valued firms, however, the graph becomes an output of the distributional geometry rather than its starting point.

Quadratic Wasserstein transport gives that geometry economic content beyond a

generic similarity metric. Its primal problem finds the least aggregate squared displacement required to reallocate one distribution into another. Unlike the W_1 Kantorovich–Rubinstein case, the quadratic W_2 dual is not a single Lipschitz price schedule, so it does not carry the same direct spatial-arbitrage interpretation. In this application, the ground cost is displacement in a maintained embedding space rather than a monetary shipping cost, so the construction neither prices literal transport nor tests for semantic arbitrage. It measures least semantic displacement conditional on the chosen representation and ground metric.

Pairwise transport nevertheless stops short of an interaction field. A distance answers how much reallocation separates two firms, and a transport plan establishes article-level correspondence, but neither determines how several candidate firms should jointly represent one target. Turning pairwise distances directly into weights would require an additional kernel, bandwidth, or nearest-neighbour rule. The central abstraction of this paper is instead *target-anchored Wasserstein barycentric reconstruction*. For each target firm, quadratic transport first aligns its articles separately with those of every candidate firm. Holding those target-specific correspondences fixed, a convex simplex step selects the nonnegative unit-sum distributional spanning weights that best reconstruct the target cloud from its aligned peers. The fitted rows form the *barycentric interaction field* W^b .

The closest geometric reference is the unrestricted Wasserstein barycentre, which selects a free centre distribution for several input laws (Agueh and Carlier, 2011). The present construction instead holds the target and pairwise alignments fixed and estimates target-specific distributional spanning weights. Section 4 states this boundary formally. The distinction is economically consequential: a large W_{ij}^b records firm j ’s conditional usefulness in reconstructing target i from the investable universe, not merely small pairwise distance. Because that usefulness is target anchored, the barycentric interaction field can be directed even though pairwise Wasserstein distance is symmetric. Its direction records reconstruction relevance, not causal influence.

At neighboring levels of aggregation, related studies use distribution-valued firm characteristics for different finance questions. Gawronsky and Huang (2026a) studies pairwise covariance envelopes implied by distances between characteristic laws, whereas Gawronsky and Huang (2026b) studies portfolio-risk bounds and allocation from distributional structure. The present paper occupies the intermediate, multi-firm level: it turns target-specific correspondences into a cross-sectional field and studies the propagation of stand-alone exposures through that field. Its field construction and adjustment model are stated independently of the pairwise and portfolio results, which locate the contribution without supplying a premise for it.

Constructing the field from text also preserves the conditioning requirement of spatial inference. Classical spatial-autoregressive methods condition on a known W , whereas estimating W from the same outcomes used in the spatial lag creates mechanical reflection (Anselin, 1988; LeSage and Pace, 2009; Kelejian and Prucha, 2010). Freezing every text-derived field before the return-evaluation window removes that same-sample feedback. It does not identify causal peer effects when omitted industries, technologies, attention, or selection jointly influence text and returns.

The possibility of several admissible fields produces the final change in the literature’s empirical question. Estimating candidate matrices separately asks “which W wins?” but cannot distinguish redundant descriptions of the same relations from distinct channels of exposure adjustment. The multi-field quadratic model instead places the barycentric interaction field beside a conventional news-link field and assigns each its own coefficient. Separate-network specifications become boundary cases of the joint model, and the estimand

becomes how adjustment divides across channels, including the case in which one field absorbs the other. With the source of the field, the exposure-adjustment mechanism, and the return bridge kept distinct, the next section introduces the stand-alone and peer-adjusted exposures linked by that mechanism.

2 Economic Environment and Stand-Alone Exposures

Begin with the familiar finite-dimensional factor model, in which an exposure is a vector of K factor loadings in $\mathbb{R}^K$. Firm i belongs to a finite universe of n firms and has a population *characteristic law* C_i on an embedding space (Ω, d_Ω) , which summarizes the distribution of its information. In the application, observed articles produce the empirical law $\hat{C}_i$ as a proxy for C_i . Think of the firm's *stand-alone exposure* ξ_i as the factor-loading vector implied by its own information before any peer adjustment. Peer adjustment maps ξ_i into the latent, model-implied *peer-adjusted exposure* B_i , and a maintained factor bridge then links B_i to observed centered excess returns. The economic sequence is therefore characteristic law, stand-alone exposure, peer-adjusted exposure, and return.

The firm's economic object is an exposure, not a return. The quadratic criterion introduced in the next section represents, in reduced form, the costs of operational, financing, or portfolio reconfiguration. It is an as-if adjustment problem and does not require firms literally to choose factor loadings each period. Keeping ξ_i and B_i separate lets the model ask how much of the exposure that enters returns reflects the firm's own information and how much reflects alignment with other firms.

Observed information is not itself an exposure: embedding locations describe an information distribution, whereas factor loadings measure sensitivity to common shocks. A transmission map is therefore needed to connect the distribution-valued characteristic law to stand-alone exposure in factor-loading units. Formally, draw $X_i \sim C_i$ and let U_i collect idiosyncratic transmission randomness. A common measurable map T converts the information draw and transmission shock into the exposure space:

$$\xi_i = T(X_i, U_i). \quad (1)$$

Together, C_i , U_i , and T generate the stand-alone exposure ξ_i . The characteristic law is measured in embedding-space units, whereas ξ_i is measured in factor-exposure units. The map T , its latent inputs, and the cross-firm coupling of the resulting stand-alone exposures are not identified from the return panel. No peer response has yet entered Equation (1).

Moving from ξ_i to B_i requires a peer average and a relative adjustment weight. Let W be the peer matrix that describes how each firm weights the other firms, and let $\lambda \geq 0$ be the dimensionless weight on peer alignment relative to the unit cost of departing from ξ_i . Because peer alignment is averaging over an economic neighborhood rather than forming a signed contrast, each row must use nonnegative weights, exclude the firm itself, and sum to one.

Definition 1 (Row-stochastic, zero-diagonal interaction matrix). An interaction matrix is any $W \in \mathbb{R}^{n \times n}$ with $W_{ij} \geq 0$ for all i and j , $W_{ii} = 0$ for all i , and $\sum_j W_{ij} = 1$ for all i . A firm never interacts with itself, weights every other firm nonnegatively, and spends exactly one unit of interaction weight on the remaining firms.

The definition makes $\sum_j W_{ij} B_j$ a peer-weighted average in the same factor-exposure units as B_i . It restricts the economic role of each row but does not select its weights. Later, Wasserstein geometry will align the empirical characteristic laws, and simplex weights will

form a target-specific barycentric interaction field. That field will be represented by a matrix satisfying the definition above; Section 4 supplies the formal construction. For now, the economic environment takes W as a fixed admissible peer matrix.

In the finite-dimensional case, ξ_i and B_i are ordinary vectors of K factor loadings, and peer adjustment operates coordinate by coordinate. To cover either finitely or countably many risk directions in one statement, we now let the exposures take values in a real separable Hilbert space $\mathcal{H}$ that generalizes $\mathbb{R}^K$. Let $F_t \in \mathcal{H}$ be a centered common factor innovation with covariance operator Γ , and let e_t^i be a centered idiosyncratic return component. The maintained return bridge specifies how the peer-adjusted exposure enters centered excess returns:

$$\tilde{r}_t^i = \langle B_i, F_t \rangle_{\mathcal{H}} + e_t^i. \quad (2)$$

Here B_i has loading units, F_t has factor-innovation units, and their inner product has return units. This bridge requires the maintained conditions that e_t^i is square-integrable, orthogonal to F_t , and has zero cross-firm covariance.

The environment now contains the stand-alone exposure ξ , the peer-adjusted exposure B , an admissible peer matrix W , and the relative adjustment weight λ . The equilibrium is solved pointwise for each realization of ξ . Consequently, B is random whenever ξ is random, even when W and λ are fixed. The next section asks whether a transparent firm-level objective maps stand-alone exposures into a unique peer-adjustment equilibrium.

3 Exposure Adjustment and Spatial Closure

How does peer adjustment transform the stand-alone exposures ξ into peer-adjusted exposures B ? We answer with an as-if reduced-form adjustment criterion that summarizes costly operational, financing, or portfolio reconfiguration. The criterion balances fidelity to the firm's stand-alone exposure against alignment with its peer-weighted exposure. We call the resulting link *spatial closure*: the objective yields, rather than assumes, a spatial autoregression in exposures.

To formalize this trade-off, fix an interaction matrix W satisfying Definition 1 and an adjustment intensity $\lambda \geq 0$. For a fixed peer profile b_{-i} , let $a \in \mathcal{H}$ denote firm i 's candidate exposure, let ξ_i denote its stand-alone exposure, and let $b_j \in \mathcal{H}$ denote the peer exposures. The adjustment criterion is the following quadratic problem.

Definition 2 (Quadratic adjustment problem). Holding the profile b_{-i} of peer exposures fixed, firm i chooses $a \in \mathcal{H}$ to minimize

$$\frac{1}{2} \|a - \xi_i\|^2 + \frac{\lambda}{2} \sum_j W_{ij} \|a - b_j\|^2. \quad (3)$$

A profile b is an equilibrium when every b_i minimizes (3) at b_{-i} .

Both terms are measured in squared exposure units. The first penalizes departure from the stand-alone exposure, whereas the second penalizes disagreement with the peer profile. Row stochasticity keeps the scale of the peer penalty comparable across firms, and λ measures its weight relative to the stand-alone-exposure penalty. The quadratic form and fixed W are maintained inputs to the model.

To characterize equilibrium, first ask when each firm's candidate exposure minimizes its criterion. Unit row sums collect the quadratic terms and yield a condition that is both necessary and sufficient.

Lemma 1 (Adjustment equilibrium condition). *For $\lambda \geq 0$ and row-stochastic W , a profile b is an equilibrium of Definition 2 if and only if*

$$b_i - \xi_i + \lambda \sum_j W_{ij}(b_i - b_j) = 0 \quad \text{for every } i. \quad (4)$$

Economically, (4) balances displacement from the stand-alone exposure against the weighted gap from peer exposures. Mathematically, completing the square isolates a linear term whose coefficient is the left-hand side of (4). A minimizer forces that coefficient to vanish; once it does, the objective gap at any alternative a is $\frac{1}{2}(1 + \lambda)\|a - b_i\|^2$. The condition is therefore necessary and sufficient in any real inner-product space, without a finite-dimensional differentiability argument. This lemma characterizes equilibrium for a fixed admissible W .

To close the firm-level conditions as a simultaneous system, use $\sum_j W_{ij} = 1$ in (4) and rearrange.

Theorem 1 (Spatial closure). *Under the hypotheses of Lemma 1, every equilibrium satisfies the Hilbert-valued spatial autoregression*

$$B = \rho WB + (1 - \rho)\xi, \quad \rho = \frac{\lambda}{1 + \lambda}. \quad (5)$$

Equation (5) is the spatial closure: the familiar spatial lag follows from the two adjustment penalties rather than entering as an assumed return equation. For each firm, peer-adjusted exposure is a convex balance between its peer average and its stand-alone exposure. Accordingly, ρ is a model-implied index of relative adjustment, not an identified causal share attributable to peers. The map $\lambda \mapsto \lambda/(1 + \lambda)$ is a strictly increasing bijection from nonnegative adjustment intensity to $0 \leq \rho < 1$. Consequently, the relative penalty weight can be recovered from a spatial coefficient by

$$\lambda = \frac{\rho}{1 - \rho}. \quad (6)$$

A value $\rho = 1/2$ gives the peer average and the stand-alone exposure equal weight, while larger values place more weight on peers. At $\rho = 0$, peer-adjusted exposure equals stand-alone exposure, whereas values approaching one place progressively greater model-implied weight on peer alignment.

To express the simultaneous system as a unique reduced form, the spatial multiplier must exist. We impose the following standard sufficient stability condition.

Hypothesis 1 (Stability condition). The pair (ρ, W) satisfies $\|\rho W\| < 1$ in the ℓ^∞ -induced operator norm. That norm is the largest absolute row sum, so the nonnegative unit rows of Definition 1 give $\|W\| = 1$ and $0 \leq \rho < 1$ is sufficient.

Economically, the condition keeps iterated peer feedback anchored by the stand-alone exposures. Mathematically, rearranging (5) and applying the convergent Neumann series for $(I - \rho W)^{-1}$ gives the following unique equilibrium.

Theorem 2 (Reduced form). *If in addition $\|\rho W\| < 1$, the equilibrium is unique and*

$$B = (1 - \rho)(I - \rho W)^{-1}\xi. \quad (7)$$

The multiplier aggregates direct and iterated peer adjustment, while the factor $1 - \rho$ preserves the scale of the stand-alone exposures. This is an equilibrium result for exposures; its observed-return interpretation requires the projection argument in Section 6.

To connect spatial closure to the pairwise exposure model of Gawronsky and Huang (2026a), consider the zero-feedback benchmark.

Corollary 1 (Zero-feedback nesting). *At $\rho = 0$, $B = \xi$.*

The equality is exact rather than limiting: when peer alignment receives zero weight, peer-adjusted exposure equals stand-alone exposure. The benchmark therefore recovers the exposure object used by their pairwise covariance restriction without reproducing that theory.

3.1 Joint Adjustment Across Two Interaction Fields

Nothing in the adjustment criterion requires the researcher to settle on one definition of a peer. Distributional similarity and explicit news links may each carry distinct conditional information about peer relevance. The economic question is therefore how peer-adjusted exposure balances both peer averages against the firm's stand-alone exposure. The two-field objective represents that trade-off directly.

Let W^B and W^N be two matrices satisfying Definition 1. Let $\lambda_B, \lambda_N \geq 0$ be the relative penalty weights on misalignment with each. The labels anticipate the two fields the application supplies; the statements below use nothing beyond admissibility of each matrix.

Definition 3 (Two-field adjustment problem). Holding the profile b_{-i} fixed, firm i chooses $a \in \mathcal{H}$ to minimize

$$\frac{1}{2} \|a - \xi_i\|^2 + \frac{\lambda_B}{2} \sum_j W_{ij}^B \|a - b_j\|^2 + \frac{\lambda_N}{2} \sum_j W_{ij}^N \|a - b_j\|^2. \quad (8)$$

A profile b is an equilibrium when every b_i minimizes (8) at b_{-i} .

All three terms carry squared exposure units, so λ_B and λ_N are dimensionless penalty weights relative to the stand-alone term. Setting either intensity to zero returns Definition 2 at the other field.

To connect the two-field objective to the one-field closure, write $\lambda = \lambda_B + \lambda_N$ and, when $\lambda > 0$, $\theta = \lambda_B/\lambda$, and define the mixture $W(\theta) = \theta W^B + (1 - \theta)W^N$. The two peer penalties in (8) then equal the one-field penalty in (3) at total intensity λ and matrix $W(\theta)$. Before applying the one-field results, the next result verifies that this mixture remains an admissible peer matrix.

Corollary 2 (Mixture nesting). *For $0 \leq \theta \leq 1$ and W^B, W^N satisfying Definition 1, the mixture $W(\theta)$ satisfies Definition 1. The endpoints $\theta = 1$ and $\theta = 0$ return W^B and W^N .*

Economically, the mixture allocates one unit of peer weight across two peer definitions while preserving the peer-average interpretation. Nonnegativity and the zero diagonal are preserved entrywise, and the row sums are a convex combination of two unit row sums. Thus the one-field results apply unchanged.

With admissibility established, the joint-field result answers how total model-implied adjustment divides across the two channels.

Theorem 3 (Two-field spatial closure). *For $\lambda_B, \lambda_N \geq 0$ and admissible W^B, W^N , every equilibrium of Definition 3 satisfies*

$$B = \rho_B W^B B + \rho_N W^N B + (1 - \rho_B - \rho_N) \xi, \quad \rho_k = \frac{\lambda_k}{1 + \lambda_B + \lambda_N}, \quad (9)$$

for $k \in \{B, N\}$.

Equation (9) makes the channel allocation explicit: $\rho_B + \rho_N$ is the total model-implied peer-adjustment weight, while ρ_B and ρ_N assign that total to the two fields. The remaining weight $1 - \rho_B - \rho_N$ anchors peer-adjusted exposure to the firm's stand-alone exposure. When $\lambda_N > 0$, equivalently $\rho_N > 0$, the ratio $\lambda_B/\lambda_N = \rho_B/\rho_N$ describes how peer adjustment divides between the channels. To obtain these coefficients, apply Theorem 1 at $(\lambda, W(\theta))$ and expand $B = \rho W(\theta)B + (1 - \rho)\xi$ with $\rho = \lambda/(1 + \lambda)$.

To establish stability of the joint system and recover each channel's relative penalty weight, combine the coefficient levels in the following result.

Proposition 1 (Two-field stability and channel inversion). *Under the hypotheses of Theorem 3, $\rho_B + \rho_N = \lambda/(1 + \lambda) < 1$, and $\|\rho_B W^B + \rho_N W^N\| \leq \rho_B + \rho_N$ in the ℓ^∞ -induced operator norm, so Hypothesis 1 holds and the reduced form of Theorem 2 applies at the mixture. Each intensity is recovered from the coefficients by*

$$\lambda_k = \frac{\rho_k}{1 - \rho_B - \rho_N}, \quad k \in \{B, N\}. \quad (10)$$

Equation (10) generalizes (6): each channel's adjustment index is its own spatial coefficient measured against the weight the firm still places on its stand-alone exposure. The norm bound follows from Corollary 2 and $\|W(\theta)\| \leq 1$. For the inversion, $1 - \rho_B - \rho_N = (1 + \lambda)^{-1}$, so dividing ρ_k by this common stand-alone weight returns λ_k .

The closure argument is complete conditional on an admissible interaction field. Where does that field come from when peer relevance must be inferred from distributions of firm text? The next section develops a *target-anchored Wasserstein barycentric reconstruction*: candidate firms are aligned to a fixed target, and their distributional spanning weights form the *barycentric interaction field* W^b . Each row is admissible, and the computation does not require a conventional kernel bandwidth.

4 Constructing the Barycentric Interaction Field

The preceding section shows how an admissible peer matrix enters the exposure-adjustment model. We now ask what should populate that matrix when each firm is represented by a distribution of text rather than by a single characteristic. For a target firm, the economic question is which combination of other firms best represents its information footprint within the investable universe. A pairwise distance can identify proximity, but it cannot determine how several candidate firms jointly represent the target.

A firm's article embeddings define an empirical characteristic law $\hat{C}_i$; its population counterpart is C_i . Quadratic Wasserstein distance aligns two article clouds by minimizing their average squared displacement in embedding units. That pairwise alignment supplies correspondence, after which one common set of peer weights can be chosen across all aligned article positions.

Consider a three-firm universe with target A and candidate peers B and C . After separately aligning the article positions of B and C with those of A , suppose the best common convex approximation assigns coefficient 0.60 to the aligned positions of B and 0.40 to those of C at every matched index. In the order (A, B, C) , the resulting target row is $W_A^b = (0, 0.60, 0.40)$. The 60–40 weights are coordinates on aligned peer positions, not probabilities of drawing whole articles from B or C .

This coordinate problem differs from an unrestricted Wasserstein barycentre. An unrestricted barycentre holds input laws and barycentre weights a_j fixed and optimizes over a new centre distribution Q through an objective such as $\inf_Q \sum_{j \neq i} a_j W_2^2(C_j, Q)$ (Agueh and Carlier, 2011). The present construction instead fixes the target C_i and, in

the empirical implementation, fixes its pairwise transport alignments before optimizing over simplex peer coordinates w_i . Target anchoring therefore narrows the claim from constructing a new centre to finding coordinates for an existing firm, while retaining the barycentric geometry.

The following definition formalizes this two-stage construction.

Definition 4 (Target-anchored Wasserstein barycentric reconstruction). Write the equally sized empirical clouds as $\hat{C}_i = M^{-1} \sum_{m=1}^M \delta_{x_{im}}$. Let $\mathcal{A}_M$ be the finite set of permutations of the M support indices. For every ordered pair $i \neq j$, first fix one optimal balanced assignment under the implementation’s predetermined deterministic selection rule,

$$\pi_{ij} \in \operatorname{argmin}_{\pi \in \mathcal{A}_M} \frac{1}{M} \sum_{m=1}^M \|x_{im} - x_{j, \pi(m)}\|_2^2, \quad y_{ijm} = x_{j, \pi_{ij}(m)}. \quad (11)$$

Conditional on these target-specific assignments, define

$$w_i \in \operatorname{argmin}_{w \in \Delta_{-i}} \frac{1}{M} \sum_{m=1}^M \left\| x_{im} - \sum_{j \neq i} w_j y_{ijm} \right\|_2^2, \quad \Delta_{-i} = \{w : w_j \geq 0, \sum_{j \neq i} w_j = 1\}. \quad (12)$$

The output is the barycentric interaction field $W_{ij}^b = (w_i)_j$ for $j \neq i$ and $W_{ii}^b = 0$.

Equation (11) fixes a transport alignment for each ordered target–candidate pair, conditional on the empirical clouds and their ground metric. Equation (12) then selects the target’s simplex coordinates by minimizing reconstruction loss across all aligned candidates jointly. Because the second objective is joint, a row of W^b is not obtained by applying a scalar transformation separately to each pairwise distance.

Every other eligible firm enters the candidate set, self-links are excluded, and the assignments are target specific. The second stage uses squared reconstruction loss and simplex-normalized weights, while equal cloud size is part of the implementation. These features, including the rule that selects among tied optimal assignments, are maintained design choices rather than consequences of Wasserstein distance alone.

Geometrically, the coefficients in w_i are barycentric-type coordinates on the aligned peer positions. Economically, we call them approximate distributional spanning weights within the investable universe: the objective value records how well the aligned convex span represents the target’s information footprint. A small residual supports approximate semantic substitutability of that peer combination for the target within the maintained representation, ground metric, and candidate universe; the simplex constraint does not assert exact spanning. The weights do not identify causal influence, a tradable replicating portfolio, or a literal arbitrage relation.

Because the target determines both the alignments and the reconstruction problem, W^b is generally asymmetric even though pairwise Wasserstein distance is symmetric. A large W_{ij}^b need not imply a large W_{ji}^b : the direction records reconstruction relevance for firm i .

Figure 2 shows how article-level correspondence becomes firm-level reconstruction relevance.




Figure 2: Target-anchored reconstruction of one row of W^b . For target firm i , separate target-to-peer transport assignments establish article-level correspondence. Holding those assignments fixed, simplex coordinates combine the aligned peer clouds to reconstruct the target. The fitted coordinates form a nonnegative unit-sum row with zero self-weight.

Read the figure from left to right: transport establishes correspondence, simplex reconstruction selects joint coordinates, and the fitted row becomes the target's peer average. The simplex constraint and leave-one-out policy imply, row by row,

$$W_{ij}^b \geq 0, \quad W_{ii}^b = 0, \quad \sum_j W_{ij}^b = 1.$$

Thus the construction satisfies Definition 1 without a separate row-normalization step. It avoids a conventional kernel-bandwidth choice, not researcher choices about the representation, candidate set, or reconstruction rule. The formal norm implication needed for the spatial multiplier is stated in the appendix as Proposition 4; in the main argument, the economic point is that every empirical row has the admissible peer-average interpretation.

4.1 Comparators

The three comparators vary one margin at a time: cardinal coordinates within a selected support, the rule that maps transport geometry into weights, and the information object that defines a relation between firms.

The equal-active-support comparator holds fixed the active peer set selected by W^b and changes only the cardinal magnitudes within that set. For each row, a peer is active when $W_{ij}^b > 10^{-8}$, the numerical tolerance fixed in the empirical producer. Writing $A_i = \{j \neq i : W_{ij}^b > 10^{-8}\}$, the equal-active-support matrix is

$$W_{ij}^{\text{eq}} = \frac{\mathbf{1}\{j \in A_i\}}{|A_i|}.$$

It has exactly the same numerical support as W^b but weights every active peer equally. Its economic question is whether fitted distributional spanning weights improve conditional spatial fit beyond selecting the active peer set.

The RBF diffusion comparator holds fixed the empirical characteristic laws, their pairwise quadratic Wasserstein distances, and the candidate universe, but changes how that geometry becomes a peer row. It replaces joint target reconstruction with a separate

proximity weight for each target–candidate pair. For $D_{ij} = W_2(\hat{C}_i, \hat{C}_j)$ and bandwidth $h > 0$, it sets

$$W_{ij}^h = \frac{\mathbf{1}\{j \neq i\} \exp(-D_{ij}^2/h)}{\sum_{k \neq i} \exp(-D_{ik}^2/h)}.$$

The application sets h to the median off-diagonal value of D_{ij}^2 . The economic question is whether peer dependence is organized by pairwise proximity alone or by the joint reconstruction of a fixed target. This dense operator also makes the conventional bandwidth choice explicit, and the appendix documents its construction.

The persistent news co-mention comparator holds fixed the firm universe, the 2018–2022 pre-evaluation timing, and the row-stochastic econometric role, but changes the information object and the link rule. The matrix W^{news} links firms that are the only two tagged names in an article and co-occur in at least two calendar years during that period. Total co-mention counts are symmetrized and row normalized. Its economic question is whether interaction is organized by approximate distributional spanning or by persistent shared news coverage. Because it is admissible in the sense of Definition 1, it can also enter Definition 3 as a separate adjustment field rather than only as an alternative to W^b . Section 7 uses it both ways: first as a comparator estimated on its own, then as the second channel of the joint model, where the two roles are the boundary and the interior of one nested family.

The barycentric field establishes empirical peers from observable characteristic laws. It does not yet determine what separation between those laws implies for latent risk exposures. The next section states that characteristic-to-exposure restriction and asks how much of the implied exposure dispersion survives equilibrium peer adjustment.

5 Cross-Sectional Wasserstein Dispersion and Spatial Attenuation

The dispersion analysis is the only part of the paper that uses a free-centre problem. Unlike the fixed-target reconstruction in Section 4, the unrestricted Wasserstein barycentre problem below selects a centre distribution Q that summarizes heterogeneity across firms.

How much characteristic-implied exposure heterogeneity survives peer adjustment? The conditional answer is economically simple: characteristic dispersion places a floor on stand-alone exposure dispersion, and adjustment through an admissible peer field cannot erase more than a coefficient-dependent share of that floor. This is a supporting closure result for the field constructed in Section 4, not a second field construction or an empirical calibration of the carrier restrictions introduced below.

Fix cross-sectional weights $q = (q_1, \dots, q_N)$ in the unit simplex. Assume that the laws considered below have finite second moments. The simplex restriction makes the weights nonnegative with unit sum, so they determine each firm’s contribution without changing the scale of the aggregate comparison. Finite second moments make the quadratic transport costs below well defined. Economically, $\mathcal{D}_q(C_1, \dots, C_N)$ is the least weighted separation compatible with all observed firm-level characteristic laws. It is therefore a conservative measure of cross-sectional information heterogeneity rather than the dispersion generated by one selected matching. Formally, for characteristic laws on (Ω, d_Ω) and $(X_1, \dots, X_N) \sim \gamma$, define

$$\mathcal{D}_q(C_1, \dots, C_N) = \inf_{\gamma \in \Pi} \mathbb{E}_\gamma \left[\sum_{i < j} q_i q_j d_\Omega(X_i, X_j)^2 \right],$$

with Π the joint laws on Ω^N whose i th marginal is C_i . The infimum asks how close the laws could be under their most favourable common coupling. Its units are squared embedding distance, and its value depends on the chosen ground metric and weights.

The Hilbert-space restriction begins to matter for the centre representation: for any laws $R_1, \dots, R_N$ on a Hilbert space, the same functional satisfies $\mathcal{D}_q(R_1, \dots, R_N) = \inf_Q \sum_i q_i W_2^2(R_i, Q)$ (Agueh and Carlier, 2011; Gawronsky and Huang, 2026b). Here Q is free and summarizes cross-sectional dispersion. Its minimizer is the unrestricted Wasserstein barycentre of the firm laws under weights q .

For two firms, the aggregate functional reduces to the familiar pairwise Wasserstein distance.

Theorem 4 (Pairwise nesting). *For $N = 2$,*

$$\mathcal{D}_{(q_1, q_2)}(C_1, C_2) = q_1 q_2 W_2^2(C_1, C_2). \quad (13)$$

For two marginals, Π is exactly the set of couplings of C_1, C_2 . The objective is therefore $q_1 q_2$ times the quadratic transport cost, so taking the infimum gives Equation (13).

Thus the geometric object nests the squared Wasserstein term used to derive the pairwise covariance envelope in Gawronsky and Huang (2026a). The equality establishes geometric nesting only; it does not import that paper's covariance or coupling assumptions.

The Wasserstein dispersion measure is in embedding units, whereas stand-alone exposures are in factor-loading units, so an explicit transfer restriction is needed before the two can be compared. Write $Z_i = \Gamma^{1/2} \xi_i$ and $P_i = \mathcal{L}(Z_i)$ for the stand-alone exposure laws in covariance coordinates. The transformation by $\Gamma^{1/2}$ places exposure differences in the risk coordinates used by the dispersion certificate. Let $t : \Omega \rightarrow \mathcal{H}$ be a common measurable carrier in those coordinates. Commonness supplies one benchmark across firms, and measurability makes the carrier image of each information draw a valid random exposure. For some $L > 0$, impose the noncollapse condition at the point where characteristic distance must become exposure distance:

$$\|t(x) - t(y)\| \geq L^{-1} d_\Omega(x, y) \quad \text{for every } x, y \in \Omega.$$

This lower-distance restriction prevents the common carrier from erasing separation between distinct information states. No upper-distance bound enters the result. For each firm, next allow synchronous slack $\tau_i \geq 0$:

$$\|\Gamma^{1/2} T(X_i, U_i) - t(X_i)\| \leq \tau_i \quad \text{almost surely.}$$

The same draw X_i appears on both sides because the bound must couple firm i 's actual stand-alone exposure to the carrier image of that information state. The slack permits firm-specific transmission noise of at most τ_i in risk-coordinate norm. Together, noncollapse and synchronous slack state the entire characteristic-to-exposure restriction used below. They do not restrict W or the peer-adjustment mechanism.

Writing $\tau_q = (\sum_i q_i \tau_i^2)^{1/2}$ for weighted root-mean-square slack and $[a]_+ = \max\{a, 0\}$, we maintain the following characteristic-to-exposure transfer inequality:

$$\sqrt{\mathcal{D}_q(P_1, \dots, P_N)} \geq \left[L^{-1} \sqrt{\mathcal{D}_q(C_1, \dots, C_N)} - \tau_q \right]_+. \quad (14)$$

The root-mean-square aggregation matches the cross-sectional weights, while the positive part sets the lower floor to zero when transmission slack absorbs the carrier-adjusted separation. This restriction is stated here in full; Gawronsky and Huang (2026b) studies its portfolio implications.

Observable separation is informative whenever its carrier-adjusted magnitude exceeds the root-mean-square transmission slack. At zero slack (14) becomes $\mathcal{D}_q(P_1, \dots, P_N) \geq L^{-2} \mathcal{D}_q(C_1, \dots, C_N)$. The carrier, L , and τ_i are maintained inputs rather than estimated objects, so (14) is a conditional restriction and is not numerically calibrated here.

5.1 Spatial attenuation under peer adjustment

The transfer inequality stops at stand-alone exposure dispersion; attenuation enters only to ask what the spatial closure does to that floor. The reduced form in Theorem 2 maps stand-alone exposure ξ into peer-adjusted exposure B through the normalized spatial multiplier:

$$S_\rho = (1 - \rho)(I - \rho W)^{-1}, \quad B = S_\rho \xi. \quad (15)$$

The restriction $0 \leq \rho < 1$ makes the Neumann expansion of S_ρ a convex mixture of current and iterated peer averages. Nonnegativity and unit row sums make each application of W an average, which is the property needed for Jensen's inequality below. Because W is directed, however, equal cross-sectional weights need not be preserved by peer averaging. Let π instead be a strictly positive stationary distribution, so $\pi^\top W = \pi^\top$. Economically, π_i measures firm i 's long-run influence under repeated peer averaging. Stationarity makes the aggregate mean invariant to W , while strict positivity keeps every firm in the comparison and makes the weighted quadratic norm nondegenerate. For any exposure profile z , define

$$V_\pi(z) = \sum_i \pi_i \|z_i - \bar{z}_\pi\|^2, \quad \bar{z}_\pi = \sum_i \pi_i z_i. \quad (16)$$

Under these restrictions, peer adjustment cannot increase cross-sectional dispersion. It also cannot eliminate more than a coefficient-dependent share: at least $\{(1 - \rho)/(1 + \rho)\}^2$ of stand-alone dispersion remains. The finite, nonempty cross-section in the theorem keeps the weighted mean and sums well defined. The next result formalizes the bracket without requiring W to be symmetric.

Theorem 5 (Spatial attenuation). *Let W be nonnegative and row stochastic, let π be a strictly positive stationary distribution, and let $0 \leq \rho < 1$. For every finite nonempty cross-section z ,*

$$\left(\frac{1 - \rho}{1 + \rho}\right)^2 V_\pi(z) \leq V_\pi(S_\rho z) \leq V_\pi(z). \quad (17)$$

To establish the upper bound, let $f = z - \bar{z}_\pi \mathbf{1}$ and define $\|f\|_\pi^2 = V_\pi(z)$. Stationarity makes Wf centered whenever f is, while nonnegative unit rows permit Jensen's inequality:

$$\sum_i \pi_i \|(Wf)_i\|^2 \leq \sum_{i,j} \pi_i W_{ij} \|f_j\|^2 = \sum_j \pi_j \|f_j\|^2.$$

The final equality uses stationarity, so W contracts the centered norm and hence so does every W^k . Because $0 \leq \rho < 1$, the Neumann mixture $S_\rho = (1 - \rho) \sum_{k \geq 0} \rho^k W^k$ is a convex combination of these contractions. This proves the upper bound. For the lower bound, the centered profile $g = S_\rho f$ satisfies $(1 - \rho)f = (I - \rho W)g$. The triangle inequality and the contraction $\|Wg\|_\pi \leq \|g\|_\pi$ give

$$(1 - \rho)\|f\|_\pi \leq \|g\|_\pi + \rho\|Wg\|_\pi \leq (1 + \rho)\|g\|_\pi.$$

Rearranging and squaring gives the lower bound in (17).

At $\rho = 0$, the two bounds coincide and peer adjustment leaves dispersion unchanged. For every admissible ρ , peer-adjusted exposures cannot be more dispersed than stand-alone exposures, while the guaranteed retained fraction $\{(1 - \rho)/(1 + \rho)\}^2$ declines with ρ . Evaluating this fraction at the fitted working-model spatial-feedback coefficient $\hat{\rho}$ gives a descriptive plug-in lower envelope, not the exact amount of attenuation or a calibration of t , L , or τ_i . It has a structural interpretation as a retained-share bound only under the return-projection conditions stated in the next section. The bracket needs neither symmetry nor an additional spectral-gap restriction, but it need not be sharp for the fitted operator.

The transfer and attenuation bounds can now be composed because they measure dispersion in the same risk coordinates. Set $Y_i = \Gamma^{1/2} B_i$, the risk-coordinate form of peer-adjusted exposure, so linearity gives $Y = S_\rho Z$. Then set the dispersion weights $q = \pi$. This is not an arbitrary aggregation choice: the stationary weights both preserve the mean under the directed operator and determine each firm's contribution to the characteristic dispersion floor.

Corollary 3 (Spatial-dispersion certificate). *Under the carrier and slack conditions stated above and the hypotheses of Theorem 5,*

$$\mathbb{E} V_\pi(Y) \geq \left(\frac{1 - \rho}{1 + \rho}\right)^2 \left[L^{-1} \sqrt{\mathcal{D}_\pi(C_1, \dots, C_N)} - \tau_\pi \right]_+^2. \quad (18)$$

The complete chain is visible in one display. The identity $V_\pi(Z) = \sum_{i < j} \pi_i \pi_j \|Z_i - Z_j\|^2$ links weighted variance to the multimarginal objective, and the actual joint law of Z is one feasible coupling. Therefore

$$\begin{aligned} \mathbb{E} V_\pi(Y) &\geq \left(\frac{1 - \rho}{1 + \rho}\right)^2 \mathbb{E} V_\pi(Z) \\ &\geq \left(\frac{1 - \rho}{1 + \rho}\right)^2 \mathcal{D}_\pi(P_1, \dots, P_N) \\ &\geq \left(\frac{1 - \rho}{1 + \rho}\right)^2 \left[L^{-1} \sqrt{\mathcal{D}_\pi(C_1, \dots, C_N)} - \tau_\pi \right]_+^2. \end{aligned}$$

The first step applies attenuation realization by realization, the second uses feasibility of the actual stand-alone exposure coupling, and the third squares the transfer inequality. This ordering shows how observed information heterogeneity becomes a stand-alone exposure floor and then a peer-adjusted exposure floor under the common adjustment coefficient ρ . Information separation, carrier distortion, transmission noise, and peer adjustment each enter once with a known direction.

The certificate is expressed in the latent risk coordinates of peer-adjusted exposure, whereas the application observes scalar returns. The next section states the return projection needed to preserve W and the common adjustment coefficient ρ , then distinguishes that structural parameter from the working-model adjustment index estimated by QMLE.

6 Return Bridge, Empirical Design, and Data

6.1 Empirical questions and return bridge

The empirical analysis follows four successive questions. First, does the barycentric interaction field organize conditional cross-sectional return dependence? This is the existence question; it concerns the return dependence associated with a predetermined field,

not whether semantic similarity causes returns. Second, does target-anchored Wasserstein barycentric reconstruction add information beyond pairwise RBF proximity computed from the same Wasserstein distances and beyond equal weighting of the selected peer support? This mechanism question distinguishes joint representability from pairwise distance decay and cardinal reconstruction weights from peer selection. Third, does the barycentric interaction field remain informative beside a persistent co-mention field? This distinctness question motivates estimating the two fields jointly. Finally, do the comparisons remain stable when representation width, model capacity, model family, or information vintage changes? These last exercises diagnose sensitivity to the measurement design; they are not additional hypotheses about economic behavior.

The theory concerns latent peer-adjusted exposure vectors, whereas the data contain scalar returns. The empirical bridge therefore has two steps. First, scalar projection preserves the spatial operator and the structural coefficient for the systematic return component. Second, the application estimates an observable return equation under a spherical working quasi-likelihood. Only the first step follows algebraically from the exposure model; interpreting the QMLE coefficient as the structural adjustment parameter requires the additional restriction stated below.

The theory's stand-alone exposure ξ_i , peer-adjusted exposure B_i , transmission map T , and risk-coordinate law P_i are latent. The application observes empirical text laws $\hat{C}_i$ and returns r_t and constructs W^b before the return-evaluation window. The following result shows that the equilibrium derivation survives when an exposure is viewed along one scalar risk direction.

Proposition 2 (Scalar projection). *For every $h \in \mathcal{H}$, the scalar field $\langle B_i, h \rangle$ satisfies the ordinary spatial autoregression with the same W and ρ , with reduced form $(1 - \rho)(I - \rho W)^{-1}$ applied to $\langle \xi_i, h \rangle$.*

By linearity of the inner product,

$$\langle B_i, h \rangle = \rho \sum_j W_{ij} \langle B_j, h \rangle + (1 - \rho) \langle \xi_i, h \rangle.$$

Stacking preserves the same W and ρ ; applying the already-established inverse in Theorem 2 gives the reduced form stated in the proposition.

To state the exact return implication, define the systematic return component and the projected stand-alone exposure by $s_t^i = \langle B_i, F_t \rangle_{\mathcal{H}}$ and $z_t^i = \langle \xi_i, F_t \rangle_{\mathcal{H}}$, and stack them across firms. Conditional on the realized factor direction F_t , scalar projection gives $s_t = \rho W s_t + (1 - \rho) z_t$. Since centered excess returns satisfy $\tilde{r}_t = s_t + e_t$, substitution, not an additional stochastic assumption, yields

$$\tilde{r}_t = \rho W \tilde{r}_t + x_t^{\text{model}}, \quad x_t^{\text{model}} = (1 - \rho) z_t + (I - \rho W) e_t. \quad (19)$$

Equation (19) is the return equation implied by the exposure model. Even when the components of e_t are cross-sectionally uncorrelated, filtering them by $I - \rho W$ and adding the projected stand-alone exposure generally makes x_t^{model} nonspherical and correlated across firms. Allowing an intercept to absorb centering and possible mean components gives the empirical notation

$$r_t = \rho W r_t + x_t. \quad (20)$$

When $x_t = x_t^{\text{model}}$, the ρ in (20) is the structural coefficient inherited from the exposure model. The QMLE instead replaces that innovation with a working specification and targets a potentially different coefficient.

6.2 Estimand and identification

Three coefficients must remain distinct. The structural ρ governs peer adjustment of latent exposures in Section 3 and lies on the model’s nonnegative branch $0 \leq \rho < 1$ fixed by Hypothesis 1; on this branch, the structural ratio is $\lambda = \rho/(1 - \rho)$. For the concentrated working log-likelihood $\ell_t^Q(\rho; W)$ introduced below, the population target is the pseudo-true coefficient

$$\rho^* \in \operatorname{argmax}_{|\rho| < 1} \mathbb{E}[\ell_t^Q(\rho; W)].$$

The sample coefficient $\hat{\rho}$ is a finite-sample maximizer of the corresponding pooled working objective for a supplied W . Thus ρ is a structural exposure-adjustment parameter, ρ^* is the best population approximation within the working likelihood, and $\hat{\rho}$ is its sample estimate.

Scalar projection establishes the algebraic bridge from latent exposures to systematic returns; it does not identify the structural adjustment parameter from the observed return panel. For ρ^* to equal the structural ρ , the expected derivative of the working objective—the population quasi-score—must have its unique zero at the structural value despite the richer innovation in (19). This return-bridge restriction is neither implied by projection nor established by the empirical design.

Absent that restriction, QMLE describes conditional spatial dependence for a specified geometry and working likelihood. Then $\lambda^* = \rho^*/(1 - \rho^*)$ is the population working-model adjustment index, and $\hat{\lambda} = \hat{\rho}/(1 - \hat{\rho})$ is its sample analogue. It is not the structural cost ratio, a causal peer effect, an observed cost, or a separately identified latent exposure parameter. The return panel also cannot identify the carrier parameters, the transmission map, individual adjustment costs, or individual peer-adjusted exposures. A negative statistical solution would describe negative conditional spatial dependence under the working likelihood. It would lie outside the peer-alignment environment of Section 2, rather than imply a negative adjustment cost.

6.3 QMLE and stationary-bootstrap inference

For a fixed W , quasi-maximum likelihood estimation (QMLE) replaces the innovation in (20) with a common intercept and a spherical disturbance:

$$r_t = \alpha \mathbf{1} + \rho W r_t + \varepsilon_t, \quad \varepsilon_t \stackrel{\text{work}}{\sim} \text{iid}(0, \sigma^2 I).$$

This mean-zero, homoskedastic, cross-sectionally spherical innovation defines a working quasi-likelihood; it is not implied by (19) or by the text construction. Concentrating out α and σ^2 leaves the spatial Jacobian term $T \log |I - \rho W|$, which accounts for the simultaneous mapping from spatial innovations to observed returns. The numerical objective is searched over the stable symmetric region $|\rho| < 1$, allowing the working likelihood to diagnose negative as well as positive conditional spatial dependence.

The pooled single-field analysis applies this same estimator to the barycentric interaction field, its RBF and equal-active-support comparators, and the persistent co-mention field. For descriptive persistence, only the coefficient and nuisance parameters are re-estimated in four non-overlapping annual samples; every geometry remains frozen. Inference uses 2,000 joint-date stationary-bootstrap refits with expected block length 21. Each resampled date retains the full cross-section, while the blocks allow for temporal dependence. Reported 95 per cent intervals are empirical quantiles of the bootstrap distribution. The QMLE

and bootstrap are therefore tools for answering the empirical questions, not separate hypotheses.

6.4 Comparator and joint-field design

The single-field comparators isolate different parts of the mechanism defined in Section 4.1. Equal weighting on W^b 's active support holds peer selection fixed and removes only the fitted cardinal reconstruction weights. The RBF field holds fixed the empirical laws, pairwise quadratic Wasserstein distances, and candidate universe, but replaces target-anchored Wasserstein barycentric reconstruction with a separate median-bandwidth distance-decay weight for each pair. The first comparison asks whether the fitted coordinate magnitudes add to peer selection; the second asks whether joint reconstruction adds to pairwise proximity. Because the RBF and barycentric fields are supplied non-nested geometries, their quasi-log-likelihood comparison is descriptive rather than a generic formal QLR test.

The persistent co-mention matrix changes the information object and link rule while retaining the same pre-evaluation window and row-stochastic econometric role. Its single-field fit and overlap diagnostics describe how the two constructions compare in isolation. The distinctness question is sharper: does the barycentric interaction field retain conditional return content once the co-mention field enters the same model? At the structural level, scalar projection applies to (9) exactly as it does to the one-field equilibrium because Proposition 2 uses only linearity of the inner product and the cross-sectional operator. Substituting as before gives

$$r_t = \rho_B W^b r_t + \rho_N W^{\text{news}} r_t + x_t. \quad (21)$$

Estimation replaces x_t with the same spherical working disturbance as in the one-field likelihood. The fitted ρ_B and ρ_N , and their transformed indices, therefore have the same pseudo-true interpretation unless the return bridge holds for both channels.

We enter the two fixed matrices unchanged and estimate their coefficients jointly rather than orthogonalizing one field against the other. Residualization would generally produce negative entries and rows that do not sum to one, destroying the admissibility required by Definition 1 and the peer-average interpretation of the peer-adjusted exposure. The joint specification preserves both admissible fields while asking whether each is needed conditional on the other.

Because $I - \rho_B W^b - \rho_N W^{\text{news}} = I - (\rho_B + \rho_N)W(\theta)$, the concentrated objective is the one-field objective evaluated at the mixture; no separate Jacobian is required. Restricting $0 \leq \theta \leq 1$ and $0 \leq \rho_B + \rho_N < 1$ imposes exactly $\lambda_B, \lambda_N \geq 0$, as required by the quadratic model.

Two consequences matter for interpretation. First, the single-field fits are the $\theta = 1$ and $\theta = 0$ boundaries of the same family. The joint quasi-log-likelihood therefore cannot fall below either boundary, and the reported gains compare specifications within one objective rather than across estimators. Both boundary fits are re-estimated here rather than carried over from Table 2. Second, a coefficient pinned at a boundary would indicate that the corresponding channel is unnecessary within the working model; the share of bootstrap refits at each boundary records this possibility.

The joint model uses the same 2,000 joint-date stationary-bootstrap refits and expected block length 21 as the pooled analysis. Both nested boundaries are refit on the same draws, making their intervals comparable with the joint intervals. Alongside the marginal intervals, the bootstrap correlation between channel coefficients measures how separately

the sample estimates them: a value near -1 would indicate that the sample estimates their sum more sharply than their division.

The quasi-likelihood-ratio statistic is reserved for these nested joint-field boundaries. The pooled comparisons use $QLR_k = 2\{\sup_{\rho_B, \rho_N \geq 0} \ell^Q(\rho_B, \rho_N) - \sup_{\rho_k=0} \ell^Q(\rho_B, \rho_N)\}$ for $H_{0,k} : \rho_k = 0$ against the nonnegative-channel alternative, with both suprema taken over the stationary parameter region. Because each null lies on the boundary of the working quasi-likelihood, its reference distribution is simulated under the corresponding restricted single-field fit rather than taken from a chi-square law. The restricted-null residual simulation, spatial inverse mapping, refitting schedule, and continuous grid-cell refinement are documented in Appendix B.4.

6.5 Data, timing, and representation construction

We constructed the barycentric interaction, RBF, equal-support, and persistent co-mention fields from information ending in 2022 and froze every matrix. We then aligned firms with compatible return histories from 3 January 2023 through 15 July 2026 and estimated conditional spatial dependence over 885 common dates. The 2023–2025 folds cover complete calendar years, whereas 2026 ends at the last available observation. No evaluation return enters the text geometry or the comparator matrices.

Within each common date, the coefficient is estimated from the co-movement between a firm’s return and the fixed weighted return of its peers, pooled under one common coefficient. Predetermination removes the direct same-sample reflection that would arise if W were estimated from these returns. It does not make text exogenous: industries, technologies, attention, and news selection can jointly shape the clouds and returns. Comparisons across frozen matrices therefore ask which predetermined geometry better organizes conditional spatial dependence, not which peer links cause returns.

We assembled firm text from Nasdaq’s public per-symbol archive of syndicated news (Nasdaq, Inc., 2026). We selected this source because its ticker-indexed retrieval, dated article bodies, and public access support one reproducible collection rule across the pre-specified universe. The choice prioritizes auditable corpus construction rather than a claim that Nasdaq is comprehensive relative to alternative news sources, and we do not treat its ticker assignment as manually validated article-level entity annotation. We constructed daily simple returns as Yahoo Finance adjusted-close ratios minus one, using the pinned `yfinance` client (Yahoo Finance, 2026; Aroussi, 2026). These are raw rather than excess returns: no risk-free series was subtracted, and the intercept in the empirical equation accommodates centering and possible mean components without removing the return-bridge restriction. Per-firm return files were inner-joined on date, and dates with a missing return were removed rather than interpolated or filled with zeros.

The source universe is a prespecified Nasdaq-100-based frame, admitted by an annual raw-coverage screen requiring at least 64 raw articles per firm in each calendar year from 2018 through 2022. The annual floor ensured that every retained firm had raw text support throughout the construction window; it did not make the frame representative of listed firms. The estimation sample is the intersection of firms with a frozen embedding cloud and a compatible return history, producing 52 firms. The news frame contains one additional firm: WBA has an embedding cloud but no compatible local return series and was dropped before return alignment. This distinction explains why the news-frame count in Table 1 exceeds the return-panel count by one.

The estimation sample is a fixed 52-firm intersection observed on 885 common evaluation dates from 3 January 2023 through 15 July 2026. This large-cap intersection excludes firms without compatible evaluation histories and is therefore survivor-conditioned. Two restric-

tions bind the reading of every estimate below. The coverage screen selects continuously and heavily covered large-cap names, so the cross-section is not representative of listed firms. Moreover, because articles were truncated to balanced clouds, article counts carry no information about a firm’s true news volume. Section 8 states what each restriction costs the interpretation.

Within the 2018–2022 window, we deterministically ordered eligible articles by URL hash and retained 128 articles per firm over the pooled period. The common cloud size put every empirical law on the support size required by the balanced assignment in (11); the annual coverage floor and the pooled cap therefore serve different purposes. This maintained truncation rule makes the geometry comparable across firms but deliberately removes raw news-volume information and need not preserve raw annual article shares.

An embedding model maps an article body to a fixed-dimensional numerical vector; its training is designed to place semantically related texts nearer in the representation space (Zhang et al., 2025). We encoded each retained article with the fixed, full-width Qwen3-Embedding-8B model and row-normalized its 4,096-coordinate vector. Rather than average a firm’s vectors into one point, we treated its balanced article cloud as an equal-mass empirical probability distribution. Wasserstein geometry supplied target-specific alignments between those distributions. The target-anchored Wasserstein barycentric reconstruction used those alignments to produce the barycentric interaction field W^b . The article cloud is a proxy for C_i ; it does not observe T , ξ_i , B_i , or P_i .

The representation exercises address design sensitivity rather than new economic hypotheses. For output width, we recomputed the Qwen3-8B geometry at 1,024, 256, and 64 coordinates beside its 4,096-coordinate primary representation. For model capacity, the fixed-width comparison placed Qwen3-Embedding-8B and Qwen3-Embedding-4B at 1,024 coordinates; their native 4,096- and 2,560-coordinate comparison changes capacity and width jointly. The 1,024-coordinate BGE-large-v1.5 arm changes model family and pretraining jointly, so it is an external sensitivity rather than an identified architecture effect. The production Qwen encoder postdates the 2018–2022 article window, so the vintage diagnostic asks whether the comparison is unusually sensitive to an encoder’s information cutoff. Finally, three matched 320-coordinate EttaX encoders held architecture, optimization recipe, compute budget, and training-token budget fixed while varying the Wikipedia snapshot: V0 used 20 December 2017, V1 used 20 December 2020, and V3 used 1 August 2026. V3 deliberately postdates both the article-construction and return-evaluation windows, so it is a negative control rather than a valid point-in-time encoder. Because EttaX capacity differs materially from the production Qwen encoders, the vintage comparisons remain descriptive sensitivities rather than identified vintage effects. Every representation arm uses the same article inputs through 2022, the same 2023–2026 return dates, and one common joint-date bootstrap schedule. The contrasts are paired within resample and corrected for multiplicity within their declared families; they are sensitivity diagnostics rather than a representation-selection test.

Sector	n	Articles	Volatility	W_2 within	W_2 cross
Communication Services	7	128.0	0.0225	1.052	1.069
Consumer Cyclical	11	128.0	0.0276	1.033	1.055
Consumer Defensive	5	128.0	0.0182	1.009	1.061
Financial Services	1	128.0	0.0272	—	1.066
Healthcare	9	128.0	0.0226	1.024	1.072
Industrials	2	128.0	0.0380	0.955	1.081
Technology	18	128.0	0.0288	0.999	1.057
All sectors	53	128.0	0.0261	1.013	1.063

Table 1: Per-sector composition of the news frame and its balanced-cloud sample: firm count, mean article count per firm, mean daily return volatility, and mean within- versus cross-sector rooted W_2 distance under the frozen Qwen3-Embedding-8B geometry. Firm counts cover the whole frame, while volatility covers only the firms carrying a return series, so the two need not agree; the estimation panel is the latter set. The Articles column reports the constant balanced-cloud size after per-firm truncation, not raw frame coverage, and therefore carries no information about a firm’s true news volume. Volatility is daily and unannualised. A dash marks a statistic undefined for that sector, such as within-sector distance for a singleton. Authors’ calculations.

Table 1 records the news frame’s industry composition and its return and text dispersion. Two features matter for what follows. Sector sizes are heavily unbalanced, with Technology holding roughly a third of the frame and two sectors holding fewer than three firms, so peer sets built from the text geometry inherit that imbalance rather than correct it. Mean within-sector W_2 distance is also below the corresponding cross-sector distance in every sector holding more than one firm. The frozen geometry therefore contains industry structure before any return is used. This composition contextualizes the resulting peer sets rather than validating them: the matched-sparsity null in Section 7 holds each row’s peer count fixed, not its sector composition.

Before turning to estimation, Figure 3 illustrates the structure of the constructed field to clarify its interpretation; it is not evidence for the return hypotheses. The heat map reads by row: each row is a target firm, each column is a candidate source, and color marks the five largest actual coefficients in that target’s row without renormalizing them. The upper panel aligns with the source columns and reports each source’s total incoming mass across all target rows. The two right-hand panels align with target rows and report the effective source count and the share of each complete row captured by the five displayed cells.




Figure 3: Illustrative barycentric interaction field from the target-anchored Wasserstein barycentric reconstruction. Each row is the actual leave-one-out solution $W_{i\cdot}^b$ to (12) over the frozen 2018–2022 article clouds: weights are nonnegative, sum to one, and assign zero self-weight. The heat map shows each target’s five largest *actual* coefficients without renormalizing them; the remaining positive coefficients still belong to W^b but are not colored. The upper and right-hand diagnostics are calculated from the complete field.

The full-field diagnostics show that both concentration within rows and total incoming mass across rows vary across firms; the five colored entries should not be mistaken for the complete operator. As an illustrative reading, AMD’s row combines firms from complementary chip design, equipment-supply, and memory segments. This example makes one target-anchored reconstruction tangible, but it neither establishes economic substitutability nor tests whether the field organizes returns. Thus W^b is a barycentric interaction field rather than a product-link map, and its effective source count is a description of sourcing breadth rather than evidence about an economic mechanism.

The next section answers existence, mechanism, and distinctness in that order before reporting descriptive annual persistence. Appendix B.2 addresses the fourth question as a design-sensitivity diagnostic.

7 Results

7.1 Conditional return dependence and the reconstruction mechanism

The first empirical question is whether the barycentric interaction field organizes conditional cross-sectional return dependence in the subsequent 2023–2026 panel. Table 2 reports separate working-QMLE estimates for three headline fields and one support-fixed diagnostic, all frozen before that return panel. The headline field W^b is the output of target-anchored Wasserstein barycentric reconstruction. Within the working QMLE, $\hat{\lambda}$ is a dimensionless adjustment index and $\hat{\rho}$ is its equivalent spatial feedback coefficient. The benchmark $\lambda = 0$ removes conditional peer alignment from the working model.

Table 2: Pooled 2023–2026 model-implied adjustment estimates. The 95 per cent intervals use 2,000 joint-date stationary-bootstrap refits, and $\hat{\rho} = \hat{\lambda}/(1 + \hat{\lambda})$. The final column reports the maximized conditional quasi-log-likelihood on the common return sample.

Interaction field	$\hat{\lambda}$	95% interval	Implied $\hat{\rho}$	ℓ^Q
Barycentric field (W^b)	3.46	[2.89, 4.17]	0.776	110 994.7
RBF–Wasserstein diffusion	2.65	[2.13, 3.30]	0.726	108 580.9
Persistent news co-mentions	1.54	[1.26, 1.92]	0.606	110 278.5
Equal active support	2.91	[2.39, 3.55]	0.744	108 987.4

The barycentric interaction field organizes substantial conditional dependence within the working model. Its pooled adjustment index is $\hat{\lambda} = 3.46$, with interval [2.89, 4.17]. Because both terms in the quadratic adjustment problem have squared-exposure units, this index is a unit-free ratio: the fitted objective places about three and a half times as much weight on misalignment between a firm’s peer-adjusted exposure and its barycentric peer average as on departure from its stand-alone exposure. The equivalent feedback coefficient is $\hat{\rho} = 0.776$. The bootstrap interval excludes the zero-feedback benchmark but spans a meaningful range of relative weights, so the sign is more precisely estimated than the magnitude. These are conditional working-model quantities, not geometry-invariant structural or causal effects.

Given that result, the mechanism question is whether target-anchored Wasserstein barycentric reconstruction adds information beyond equal weighting of the selected peer support and pairwise RBF proximity computed from the same Wasserstein distances. The equal-active-support row retains every peer selected by W^b but replaces the fitted distributional spanning weights with equal weights. Its adjustment estimate remains positive, so peer selection itself carries conditional signal. Its lower conditional quasi-log-likelihood indicates that support alone does not reproduce the fit obtained from the fitted coordinates. The RBF–Wasserstein row holds fixed the firm distributions, pairwise transport distances, and candidate universe, but replaces target-anchored joint reconstruction with a separate distance-decay weight for each pair. This field also has a positive adjustment estimate, so pairwise proximity organizes some conditional dependence. Its lower conditional fit provides evidence consistent with joint representability carrying information beyond pairwise distance decay in this sample. These likelihood rankings compare non-nested working models and are therefore mechanism diagnostics, not formal likelihood-ratio tests. The news-link row also makes clear why feedback magnitude is not a fit ranking: it attains higher conditional likelihood than the RBF field with a smaller adjustment index. Appendix B.2 treats model family, capacity, output width, and encoder

vintage as design-sensitivity diagnostics for the barycentric–RBF comparison.

7.2 Are the barycentric and news-link fields distinct?

The two fields choose substantially different peers but produce related, non-interchangeable induced return signals. Table 3 separates this conclusion into peer selection, cardinal weighting, and the resulting peer-return series. Support and induced-signal agreement are evaluated against matched-random nulls: row sparsity creates a floor for peer overlap, while common cross-sectional return variation creates one for induced-signal correlation.

Table 3: Agreement between the frozen barycentric interaction field and the persistent co-mention field. The null draws 1000 fixed-seed baskets carrying W^b ’s own row sparsities; p is the share of null draws reaching the observed value. Rows with no mechanical floor carry no null.

Stage	Statistic	Observed	Null	p
Peer sets	Directed share of W^b peers	0.579	0.531	0.001
	Rank-matched share	0.637	—	—
	Jaccard overlap	0.508	—	—
Weights	Off-diagonal correlation	0.669	—	—
	Cosine on the support union	0.726	—	—
	Rank correlation on common edges	0.467	—	—
Induced field	Correlation of Wr_t	0.898	0.700	0.001
	Coefficient of determination	0.806	—	—

Peer selection provides the clearest evidence of distinctness. The barycentric rows are the denser of the two, carrying 38.8 active peers on average against 27.1 for the co-mention graph, and 0.579 of the barycentric peers are also co-mention peers. A random basket with the same row sparsities reaches 0.531, so the observed excess is about five percentage points and exceeds the matched-random benchmark at $p = 0.001$. The null comparison therefore establishes above-chance overlap, not equivalence of the peer sets.

The cardinal weights also show only partial agreement. Their off-diagonal correlation is 0.669, while the rank correlation over the edges the two constructions share—an average of 22.3 per row—is lower, at 0.467.

The induced peer-return series are more similar than the underlying supports or weights, but this comparison is also the one most in need of a null. The two spatial lags correlate at 0.898 over the evaluation panel, so one accounts for 0.806 of the other’s variation. Taken alone, that number would overstate agreement because any two row-stochastic averages of a strongly co-moving cross-section are correlated by construction: a random basket at W^b ’s sparsity already reaches 0.700 against the same comparator. The observed value exceeds that floor at $p = 0.001$, and the per-firm correlations range from 0.553 to 0.973. Thus common return variation makes the fields look more alike after they are applied to returns than they do as peer maps. The observed excess over the matched-random floor nevertheless leaves room for distinct conditional content, which the joint model evaluates next.

7.3 Does the barycentric field remain informative beside a persistent co-mention field?

Joint estimation reallocates working-model feedback across the two fields rather than materially increasing its total. The barycentric-only boundary gives $\hat{\rho} = 0.776$, whereas the joint estimate is $\hat{\rho}_B + \hat{\rho}_N = 0.761$. The 95 per cent interval for the joint total, $[0.729, 0.793]$, contains the barycentric-only point estimate. Thus the news-link field absorbs part of the conditional dependence attributed to W^b when that field is estimated alone, while total feedback remains similar.

Table 4 estimates (21) on the same pooled panel and with the same two frozen matrices. It treats them as channels in one conditional return specification rather than as rival structural mechanisms. Within this working model, each channel has the adjustment index $\hat{\lambda}_k = \hat{\rho}_k / (1 - \hat{\rho}_B - \hat{\rho}_N)$, and the relevant null for each is $\rho_k = 0$: the corresponding field adds no conditional fit once the other is present.

Table 4: Pooled 2023–2026 joint two-field estimates. The 95 per cent intervals use the same 2,000 joint-date stationary-bootstrap refits as Table 2. The single-field rows are the $\theta = 1$ and $\theta = 0$ boundaries of the same nested family, re-estimated and re-resampled here on those same draws, and the quasi-log-likelihood column is measured against the joint fit.

Channel	$\hat{\rho}$	95% interval	$\hat{\lambda}$	95% interval	$\Delta\ell^Q$
Barycentric (W^b)	0.556	[0.521, 0.588]	2.33	[2.00, 2.70]	
News-link (W^{news})	0.205	[0.164, 0.250]	0.86	[0.61, 1.18]	
Total	0.761	[0.729, 0.793]			
<i>Nested single-field boundaries</i>					
W^b only ($\theta = 1$)	0.776	[0.743, 0.807]	3.46	[2.89, 4.17]	−169.8
W^{news} only ($\theta = 0$)	0.606	[0.557, 0.657]	1.54	[1.26, 1.92]	−886.0

The channel split is economically asymmetric. The barycentric channel has the adjustment index $\hat{\lambda}_B = 2.33$, with interval $[2.00, 2.70]$. The corresponding news-link index is $\hat{\lambda}_N = 0.86$, with interval $[0.61, 1.18]$. As dimensionless penalties relative to the stand-alone exposure term, these estimates assign more than twice as much weight to misalignment with the barycentric peer average as to misalignment among firms reported alongside one another. Both intervals exclude zero, and no bootstrap refit is pinned at either boundary.

Figure 4 describes the likelihood and bootstrap geometry behind this channel uncertainty; it does not supply a calibrated confidence region.




Figure 4: Joint two-field likelihood and bootstrap geometry. The horizontal axis is barycentric-channel feedback ρ_B , and the vertical axis is news-link feedback ρ_N . The left panel places the joint estimate inside the admissible triangle, with both single-field fits on the axes and the stability boundary $\rho_B + \rho_N = 1$ marked. The right panel is the same profiled surface zoomed to the bootstrap cloud, with a descriptive two-parameter quasi-log-likelihood contour. Neither the contour nor the cloud is a calibrated confidence region. The cloud’s diagonal tilt records the trade-off between channels, while its dispersion in both directions shows the sampling variation in their allocation.

The two channel coefficients have bootstrap correlation -0.683 . That pattern is expected given the induced-field correlation of 0.898 : when the two peer-return signals move together, one coefficient can partly offset the other across refits. The draws nevertheless vary in both directions rather than collapsing onto a one-dimensional ridge. The figure therefore indicates that total feedback is estimated more precisely than its allocation across fields, without calibrating a joint confidence set.

The incremental-fit decisions instead come from boundary-calibrated QLR tests. Because each nonnegative-channel null lies on the boundary, the reference distribution is simulated under the corresponding restricted single-field fit. For $H_{0,N} : \rho_N = 0$, the observed statistic is $QLR_N = 339.5$, with restricted-null $p = 0.0005$. For $H_{0,B} : \rho_B = 0$, the corresponding values are $QLR_B = 1772.1$ and $p = 0.0005$. Both restrictions are rejected under the null-imposed calibration, so each field improves conditional fit once the other is included. Across 885 common dates, the raw quasi-log-likelihood gains over the barycentric-only and news-only boundaries are, respectively, 169.8 and 886.0 . The larger loss from dropping the barycentric field is consistent with the asymmetric channel indices; it remains a conditional-fit comparison rather than a structural or causal decomposition.

7.4 Descriptive persistence under a frozen field

The conditional association persists descriptively across all four annual evaluation samples. Table 5 keeps the 2018–2022 barycentric interaction field fixed and re-estimates only the common coefficient and nuisance parameters within each non-overlapping return sample. The annual refits are reported without a cross-year equality statistic or joint calibration, so this exercise describes persistence rather than testing coefficient constancy.

Table 5: Annual model-implied adjustment under the frozen barycentric interaction field. The 2026 period ends on 15 July.

Evaluation year	$\hat{\lambda}$	95% interval	Implied $\hat{\rho}$
2023	3.30	[2.73, 3.88]	0.768
2024	2.89	[2.42, 3.36]	0.743
2025	3.94	[2.46, 5.69]	0.798
2026	3.62	[2.83, 4.60]	0.784

The annual adjustment indices span 2.89– 3.94. All four bootstrap intervals remain above the zero-feedback benchmark, although their widths vary, especially in 2025; the annual evidence therefore supports persistence but not equality of magnitudes. The 2026 estimate uses only 133 dates, so its precision is not directly comparable with that of a complete year.

Table 9 in the appendix reports joint annual estimates under the same frozen matrices; the exercise is descriptive rather than a formal constancy test. The next section sets out the economic and empirical limits on that interpretation.

8 Discussion and Limitations

Within its prespecified information geometry and working return model, the paper establishes a bounded positive implication. The target-anchored Wasserstein barycentric reconstruction maps distribution-valued information positions into a predetermined, row-stochastic barycentric interaction field, and the quadratic closure gives an economic scale to the relation between stand-alone and peer-adjusted exposure in Theorem 1. In the return panel, this field and the persistent news co-mention field select largely different peers but generate correlated peer-return series. Estimating them jointly leaves total model-implied adjustment close to its single-field level and reallocates it across the two channels, a result that separate one-field fits conceal.

The first limit is identification. The quadratic criterion is an as-if reduced-form representation of exposure adjustment, not a claim that managers observe the barycentric interaction field or literally solve the displayed optimization problem. Constructing every field from information ending in 2022 and freezing it before the 2023–2026 return panel prevents evaluation returns from mechanically determining their own peer weights. That predetermination does not make text exogenous: industries, technologies, investor attention, and state-dependent reporting selection can jointly shape the information positions and returns. The likelihood gains and channel decomposition therefore measure conditional field fit rather than causal peer effects; isolating peer transmission would require an exclusion restriction or shock design. The full distinction between the structural adjustment parameter and the pseudo-true target of QMLE remains the identification boundary in Section 6.

The second limit concerns measurement and design. The central field contribution depends on a fixed representation and ground metric for quadratic Wasserstein transport, balanced article clouds, and the target-specific alignment and simplex restrictions of the target-anchored Wasserstein barycentric reconstruction. Balancing the clouds makes firms comparable but removes news volume as a source of information, while changing the representation, ground metric, or reconstruction design can alter the barycentric interaction field and its conditional fit. The implementation uses a predetermined deterministic rule to select one optimal pairwise assignment, but some assignments are tied; the paper does not establish that alternative optimal selections would leave W^b unchanged. Cross-geometry

likelihood comparisons are consequently evidence about competing measurement designs, not geometry-free rankings. Prespecified alternatives using weighted or unbalanced clouds, other ground metrics, and held-out evaluation windows would show which features of the field survive those measurement choices.

Measurement also bounds the dispersion and multi-field conclusions. The common carrier, its noncollapse constant L , and the transmission radii τ_i in Equation (14) remain latent and are not estimated by the return QMLE. The attenuation bracket therefore limits the share of implied stand-alone exposure dispersion that peer adjustment can erase but does not determine its level, so Corollary 3 remains a conditional restriction rather than a measured quantity. The two fields’ peer-return series correlate at 0.898, and the bootstrap correlation between $\hat{\rho}_B$ and $\hat{\rho}_N$ is -0.683 . Both channel intervals exclude zero in this sample, but more nearly collinear fields could leave their division weakly determined even when total adjustment remains precise. The split should therefore be read from the joint region in Figure 4; extensions to additional fields should report the corresponding joint uncertainty and induced-signal collinearity.

The final limit is external validity. The evidence comes from a large-cap, survivor-conditioned intersection of 52 firms with complete return histories, rather than a representative cross-section of listed firms. A broader or changing universe would alter each target’s feasible peer set and could change the field’s stationary weighting. The timing is also specific: the barycentric interaction field uses 2018–2022 text, the return evaluation begins in 2023, and the 2026 window ends on 15 July, making that final period unsuitable for like-for-like comparison with complete calendar years. Reconstructing fields in successive ex ante windows and extending the sample to firms with shorter histories would test whether the conditional patterns persist across firm entry, market segments, and information regimes.

9 Conclusion

Firms occupy distribution-valued information positions rather than single economic locations. This paper turns target-anchored Wasserstein barycentric reconstruction into a cross-sectional interaction field for those positions. For each target, the fitted distributional spanning weights combine aligned peer positions; repeated across firms, they form the admissible, directed barycentric interaction field W^b , with nonnegative unit-sum rows and a zero diagonal.

Placing that field inside the quadratic exposure-adjustment problem yields spatial closure: peer-adjusted exposure balances its stand-alone counterpart against the peer average and obeys the familiar spatial autoregression. Within this closure, $\rho = \lambda/(1 + \lambda)$ records the relative intensity of peer alignment, so spatial feedback acquires an exposure-adjustment interpretation rather than remaining an unexplained coefficient conditional on a supplied matrix.

In the 2023–2026 return panel, the barycentric interaction field (constructed without using those evaluation returns) organizes conditional cross-sectional return dependence and delivers stronger conditional fit than pairwise RBF proximity constructed from the same Wasserstein distances. It selects a distinct peer structure and remains incrementally informative when a persistent field of explicit news co-mention links enters the same specification. Joint estimation reallocates model-implied adjustment between these fields rather than materially increasing total feedback.

Methodologically, the paper provides a bridge from distribution-valued representation to spatial econometrics: it converts target-specific joint representability in Wasserstein space

into an admissible field for spatial quasi-likelihood. As Section 6 explains, the estimates are conditional working-model quantities, not causal peer effects or geometry-invariant structural adjustment parameters. Spatial econometrics therefore need not begin after the researcher has supplied W : it can begin one level upstream, by constructing the field from firms’ distribution-valued information positions.

Data availability statement

Firm news records were obtained from Nasdaq’s public ticker-indexed news archive (Nasdaq, Inc., 2026). Daily adjusted-price histories were obtained from Yahoo Finance (Yahoo Finance, 2026) through the pinned `yfinance` client (Aroussi, 2026). Article vectors were produced with `qwen/qwen3-embedding-8b` through OpenRouter; the embedding model is documented by Zhang et al. (2025). The provider-hosted articles, model endpoint, and market data remain subject to their providers’ terms and are not redistributed. A citable public release containing the versioned code, environment specifications, derived interaction fields, estimation outputs, and scripts needed to regenerate the manuscript’s tables and figures will be deposited upon acceptance; the archive DOI will be added to the accepted manuscript.

Funding

The authors received no financial support for the research, authorship, or publication of this article.

Disclosure statement

The authors report no potential conflict of interest.

References

- Martial Agueh and Guillaume Carlier. Barycenters in the Wasserstein Space. *SIAM Journal on Mathematical Analysis*, 43(2):904–924, 2011. URL <https://doi.org/10.1137/100805741>.
- Luc Anselin. *Spatial Econometrics: Methods and Models*, volume 4. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1988. URL <https://www.springer.com/gp/book/9789024737352>.
- Ran Aroussi. Yfinance 1.5.1: Download market data from Yahoo! Finance’s API, jun 2026. URL <https://pypi.org/project/yfinance/1.5.1/>.
- Coralio Ballester, Antoni Calvó-Armengol, and Yves Zenou. Who’s Who in Networks: Wanted: The Key Player. *Econometrica*, 74(5):1403–1417, 2006. URL <https://doi.org/10.1111/j.1468-0262.2006.00709.x>.
- Azi Ben-Rephael, Bruce I. Carlin, Zhi Da, and Ryan D. Israelsen. Information Consumption and Asset Pricing, 2019. URL <https://ssrn.com/abstract=3002339>.

- Lin William Cong, Tengyuan Liang, Xiao Zhang, and Wu Zhu. Textual Factors: A Scalable, Interpretable, and Data-Driven Approach to Analyzing Unstructured Information, 2024. URL <https://www.nber.org/papers/w33168>.
- Viviana Fernandez. Spatial Linkages in International Financial Markets. *Quantitative Finance*, 11(2):237–245, feb 2011. URL <http://www.tandfonline.com/doi/abs/10.1080/14697680903127403>.
- Marcus Gawronsky and Chun-Sung Huang. Systematic Covariance Envelopes from Wasserstein Geometry: Evidence from Language-Model Representations. jul 2026a.
- Marcus Gawronsky and Chun-Sung Huang. Portfolio Risk Bounds without a Return Covariance: Leveraging Distributional Fields from Language-Model Representations. jul 2026b.
- Shuyi Ge, Shaoran Li, and Oliver Linton. News-implied Linkages and Local Dependency in the Equity Market. *Journal of Econometrics*, 235(2):779–815, 2023.
- Shuyi Ge, Shaoran Li, Oliver Linton, Weiguang Liu, and Wen Su. Should We Augment Large Covariance Matrix Estimation with Auxiliary Network Information?, 2026.
- Harry H. Kelejian and Ingmar R. Prucha. Specification and Estimation of Spatial Autoregressive Models with Autoregressive and Heteroskedastic Disturbances. *Journal of Econometrics*, 157(1):53–67, jul 2010. URL <https://www.sciencedirect.com/science/article/abs/pii/S0304407609002784>.
- Steven Kou, Xianhua Peng, and Haowen Zhong. Asset Pricing with Spatial Interaction. *Management Science*, 64(5):2083–2101, may 2018. URL <https://pubsonline.informs.org/doi/10.1287/mnsc.2016.2627>.
- James P. LeSage and Robert Kelley Pace. *Introduction to Spatial Econometrics*. Chapman and Hall/CRC, Boca Raton, Florida, may 2009. URL <https://www.routledge.com/Introduction-to-Spatial-Econometrics/LeSage-Pace/p/book/9781032477749>.
- Nasdaq, Inc. Nasdaq ticker-indexed news headlines [dataset], 2026. URL <https://www.nasdaq.com/market-activity/quotes/news-headlines>.
- Anna Scherbina and Bernd Schlusche. Economic {linkages} {inferred} from {news} {stories} and the {predictability} of {stock} {returns}. *SSRN Electronic Journal*, 2013. URL <http://www.ssrn.com/abstract=2363436>.
- Gustavo Schwenkler and Hannan Zheng. The Network of Firms Implied by the News. Technical Report 108, European Systemic Risk Board, 2020. URL <https://hdl.handle.net/10419/244260>.
- Bumho Son and Jaewook Lee. Graph-based Multi-Factor Asset Pricing Model. *Finance Research Letters*, 44:102032, 2022. URL <https://doi.org/10.1016/j.frl.2021.102032>.
- Ajim Uddin, Xinyuan Tao, and Dantong Yu. The Network Factor of Equity Pricing: A Signed Graph Laplacian Approach. *Journal of Financial Econometrics*, 22(5):1616–1655, 2024. URL <https://doi.org/10.1093/jjfinec/nbae010>.
- Shunyao Wang, Ming Cheng, and Christina Dan Wang. Newsnet--sdf: Stochastic Discount Factor Estimation with Pre-trained Language-Model News Embeddings via Adversarial Networks. *China Digital Finance Conference*, pages 3–21, 2025. URL <https://arxiv.org/abs/2505.06864>.

Shitao Xiao, Zheng Liu, Peitian Zhang, Niklas Muennighoff, Defu Lian, and Jian-Yun Nie. C-pack: Packed Resources For General Chinese Embeddings. *Proceedings of the 47th International ACM SIGIR Conference on Research and Development in Information Retrieval*, pages 641–649, 2024. URL <https://dl.acm.org/doi/10.1145/3626772.3657878>.

Yahoo Finance. Yahoo Finance historical market data [dataset], 2026. URL <https://finance.yahoo.com/>.

Yanzhao Zhang, Mingxin Li, Dingkun Long, Xin Zhang, Huan Lin, Baosong Yang, Pengjun Xie, An Yang, Dayiheng Liu, Junyang Lin, Fei Huang, and Jingren Zhou. Qwen3 Embedding: Advancing Text Embedding and Reranking Through Foundation Models. Technical Report 2506.05176, arXiv, 2025. URL <https://arxiv.org/abs/2506.05176>.

A Stability and spectral diagnostics for the barycentric field

Two questions organize this appendix: what makes the constructed field a stable peer-averaging system, and how close is the fitted model to the strong-interaction boundary? We first state the maintained conditions and derive their implications for peer averaging, stationary weighting, and the spatial multiplier. We then report the fitted residual, Dobrushin, and finite- ρ diagnostics before characterizing the rank-one boundary. These results certify the constructed field, although their matrix implications apply to any row-stochastic interaction matrix that satisfies the stated conditions.

A.1 Maintained operator conditions

The first question is which matrix properties are needed for this interpretation. The main argument requires a nonnegative row-stochastic interaction matrix W , a strictly positive stationary probability vector π , and a coefficient ρ that satisfies the stability condition in Hypothesis 1. Row stochasticity defines the directed peer average, the stationary vector supplies the weighting in Theorem 5, and the stability condition makes the spatial multiplier well defined. Let $P = \mathbf{1}\pi^\top$. All matrix norms here are maximum absolute row-sum norms.

In the application, $W = W^b$ is the barycentric interaction field obtained from target-anchored reconstruction. The spectral results operate on that fitted matrix.

The boundary analysis also requires a rate at which repeated peer averaging approaches P . The following certificate states that condition.

Definition 5 (Geometric Perron certificate). A row-stochastic matrix W has a geometric Perron certificate (π, C, q) if $0 \leq q < 1$ and

$$\|W^k - P\| \leq Cq^k \quad \text{for every } k \geq 0.$$

A.2 Implications for peer averaging and the multiplier

The next question is whether the maintained conditions produce coherent long-run peer weights. Row stochasticity makes Wx a peer average, while the stability condition makes $(I - \rho W)^{-1}$, and hence the equilibrium mapping from stand-alone exposures, well defined. The geometric certificate adds the limiting result needed for the boundary analysis.

Theorem 6 (Geometric certificate implies the Perron limit). *If W has a geometric Perron certificate (π, C, q) , then $W^k \rightarrow P$ as $k \rightarrow \infty$.*

Because $0 \leq q < 1$, $Cq^k \rightarrow 0$, so the defining inequality yields $W^k \rightarrow P$.

The Perron limit has two consequences for the economic aggregation in the main text. First, repeated peer averaging replaces any initial profile by its π -weighted cross-sectional mean.

Theorem 7 (Stationary interaction centrality). *Under the Perron limit, $W^k x \rightarrow (\pi^\top x)\mathbf{1}$ for every vector x .*

Multiplying $W^k \rightarrow P = \mathbf{1}\pi^\top$ by a fixed vector x gives $W^k x \rightarrow Px = (\pi^\top x)\mathbf{1}$.

Second, the weights that define this long-run mean are invariant under one application of the operator.

Theorem 8 (Stationarity). *Under the Perron limit, $\pi^\top W = \pi^\top$.*

Since $W^{k+1} = W^k W$, the Perron limit gives $W^{k+1} \rightarrow PW$, while the same limit with $k+1$ gives $W^{k+1} \rightarrow P$. Uniqueness of limits implies $PW = P$, and $P = \mathbf{1}\pi^\top$ then gives $\pi^\top W = \pi^\top$.

Together, the two results make π_i firm i 's long-run influence and justify the stationary centring in Theorem 5.

A.3 Fitted operator and finite- ρ diagnostics

The empirical diagnostic question is whether the fitted field and coefficient satisfy these conditions and whether the boundary closely approximates the empirical model. The fitted 52×52 operator is row stochastic to a maximum residual of 0.000 000 00. Because its entries are nonnegative and its rows sum to one, its maximum absolute row-sum norm is one. The pooled estimate $\hat{\rho} = 0.776$ therefore satisfies the stability condition and makes the spatial multiplier well defined. Power iteration also yields a strictly positive stationary distribution with minimum mass 0.0105 and stationarity residual 3.38×10^{-13} . These fitted checks support, respectively, the peer average, the equilibrium mapping from stand-alone exposures, and the stationary weighting used in the main text.

The Dobrushin coefficient $\delta(W) = \frac{1}{2} \max_{i,j} \sum_m |W_{im} - W_{jm}|$ measures the largest difference between any two firms' peer-weight profiles. A value below one means that all row pairs retain some overlap and supplies the auditable envelope $C = 2$ and $q = \delta(W)$: every row of W^k lies within $2\delta(W)^k$ of the stationary distribution in ℓ_1 distance. For the fitted operator, the second power is strictly positive with minimum entry 0.001 630, and the Dobrushin coefficient is 0.862. Thus the fitted operator has a geometric Perron certificate: its iterated peer profiles converge even though the operator is directed.

The Perron limit describes the boundary $\rho \uparrow 1$, whereas the empirical application uses a finite working-model coefficient. An exact decomposition separates the rank-one part of the multiplier from the remaining cross-sectional variation at any admissible ρ . Write $N = W - P$. Row stochasticity, stationarity, and unit mass imply $WP = PW = P^2 = P$. Whenever the displayed inverses exist and $\rho \neq 1$, the Leontief multiplier splits exactly as

$$(I - \rho W)^{-1} = (1 - \rho)^{-1}P + (I - \rho N)^{-1}(I - P). \quad (22)$$

The error made by replacing the normalized multiplier with its limiting projection is therefore explicit rather than asymptotic:

$$(1 - \rho)(I - \rho W)^{-1} - P = (1 - \rho)(I - \rho N)^{-1}(I - P). \quad (23)$$

Proposition 3 (Finite- ρ resolvent bound). *If $\|\rho(W - P)\| < 1$, then*

$$\|(1 - \rho)(I - \rho W)^{-1} - P\| \leq |1 - \rho| \{1 - \|\rho(W - P)\|\}^{-1} \|I - P\|.$$

The proposition provides an auditable upper bound on the distance between the finite- ρ normalized multiplier and its limiting projection. Taking norms in (23) and applying the Neumann bound

$$\|(I - \rho N)^{-1}\| \leq \{1 - \|\rho N\|\}^{-1},$$

gives the displayed bound.

At the pooled fitted coefficient, the exact normalized-resolvent error is 0.675, against a Dobrushin bound of 1.353 and a Neumann-region bound of 3.786. The exact error remains materially above zero, so the rank-one limit is not a close approximation to the fitted normalized multiplier in this norm. The two larger bounds are valid but conservative envelopes. Thus the fitted coefficient lies inside the stable region, but the boundary approximation is not the empirical working model.

A.4 The rank-one boundary

The boundary question is what remains of cross-sectional variation as peer adjustment approaches its strong-interaction limit. The boundary result follows in two steps. First, convergence of repeated peer averaging makes the normalized spatial multiplier converge to the stationary projection.

Theorem 9 (Resolvent limit from the power limit). *Let W be row stochastic and suppose $W^k \rightarrow P$. Then $(1 - \rho)(I - \rho W)^{-1} \rightarrow P$ as $\rho \uparrow 1$.*

This result converts the power limit into a statement about spatial feedback. The normalized Neumann series is

$$S_\rho = (1 - \rho)(I - \rho W)^{-1} = (1 - \rho) \sum_{k \geq 0} \rho^k W^k.$$

For any $\varepsilon > 0$, choose K so that $\|W^k - P\| < \varepsilon$ for $k \geq K$. Since $P = (1 - \rho) \sum_{k \geq 0} \rho^k P$, split the difference into its finite head and tail:

$$\|S_\rho - P\| \leq (1 - \rho) \sum_{k < K} \rho^k \|W^k - P\| + (1 - \rho) \sum_{k \geq K} \rho^k \|W^k - P\|.$$

The finite head vanishes as $\rho \uparrow 1$, while the tail is at most ε ; the uniform bound $\|W^k - P\| \leq 2$ justifies the split. Hence $S_\rho \rightarrow P$.

Second, substituting that multiplier limit into the spatial covariance removes all cross-sectional directions except the common one.

Theorem 10 (Rank-one collapse of the rescaled spatial covariance). *Let $\Sigma_{\text{SAR}} = (I - \rho W)^{-1} V (I - \rho W)^{-\top}$. Under the Perron limit,*

$$(1 - \rho)^2 \Sigma_{\text{SAR}} \longrightarrow (\pi^\top V \pi) \mathbf{1} \mathbf{1}^\top \quad \text{as } \rho \uparrow 1.$$

The proof follows by continuity of matrix multiplication. Set $S_\rho = (1 - \rho)(I - \rho W)^{-1} \rightarrow P$. Continuity of matrix multiplication gives $S_\rho V S_\rho^\top \rightarrow P V P^\top$. Since $P = \mathbf{1} \pi^\top$,

$$P V P^\top = (\pi^\top V \pi) \mathbf{1} \mathbf{1}^\top.$$

Theorem 10 is the $\rho \uparrow 1$ endpoint of the attenuation result: at the boundary, spatial feedback removes cross-sectional dispersion entirely and the covariance becomes rank one. The finite- ρ remainder shows why this endpoint is useful as a theoretical boundary but not as an approximation to the fitted working model.

B Supporting Interaction-Field Results and Implementation

The main analysis leaves four supporting questions. The first two ask whether the distinction between joint reconstruction and pairwise proximity survives a conventional distance-to-weight map and alternative text representations. The third asks whether the pooled division between the barycentric and news-link fields recurs across annual samples. The fourth verifies the estimation and operator conditions used in the main text. This appendix answers those questions in order and then records the reproducibility and formal-verification boundaries.

The firms used to construct the primary barycentric interaction field are listed in Table 6.

Table 6: Canonical empirical roster: ticker symbol, firm name, and sector label from Nasdaq summary metadata for the firms in the canonical priced W_2 geometry, sorted by sector then symbol. Authors’ calculations.

Symbol	Name	Sector
CMCSA	Comcast Corporation	Communication Services
EA	Electronic Arts Inc.	Communication Services
GOOG	Alphabet Inc.	Communication Services
MTCH	Match Group Inc.	Communication Services
NFLX	Netflix Inc.	Communication Services
SIRI	Sirius XM Holdings Inc.	Communication Services
TMUS	T-Mobile US Inc.	Communication Services
EXPE	Expedia Group Inc.	Consumer Cyclical
HAS	Hasbro Inc.	Consumer Cyclical
JD	JD.com Inc.	Consumer Cyclical
LULU	Lululemon Athletica Inc.	Consumer Cyclical
MAR	Marriott International Inc.	Consumer Cyclical
MAT	Mattel Inc.	Consumer Cyclical
MELI	MercadoLibre Inc.	Consumer Cyclical
ORLY	O’Reilly Automotive Inc.	Consumer Cyclical
SBUX	Starbucks Corporation	Consumer Cyclical
ULTA	Ulta Beauty Inc.	Consumer Cyclical
WYNN	Wynn Resorts Limited	Consumer Cyclical
COST	Costco Wholesale Corporation	Consumer Defensive
DLTR	Dollar Tree Inc.	Consumer Defensive
KDP	Keurig Dr Pepper Inc.	Consumer Defensive
KHC	The Kraft Heinz Company	Consumer Defensive
PEP	PepsiCo Inc.	Consumer Defensive
PYPL	PayPal Holdings Inc.	Financial Services
ALGN	Align Technology Inc.	Healthcare
AMGN	Amgen Inc.	Healthcare
AZN	AstraZeneca PLC	Healthcare
BIIB	Biogen Inc.	Healthcare
GILD	Gilead Sciences Inc.	Healthcare
ISRG	Intuitive Surgical Inc.	Healthcare
REGN	Regeneron Pharmaceuticals Inc.	Healthcare
VRTX	Vertex Pharmaceuticals Incorporated	Healthcare

Symbol	Name	Sector
AAL	American Airlines Group Inc.	Industrials
UAL	United Airlines Holdings Inc.	Industrials
ADP	Automatic Data Processing Inc.	Technology
AMAT	Applied Materials Inc.	Technology
AMD	Advanced Micro Devices Inc.	Technology
AVGO	Broadcom Inc.	Technology
CSCO	Cisco Systems Inc.	Technology
ENPH	Enphase Energy Inc.	Technology
FTNT	Fortinet Inc.	Technology
INTC	Intel Corporation	Technology
INTU	Intuit Inc.	Technology
LRCX	Lam Research Corporation	Technology
MRVL	Marvell Technology Inc.	Technology
MU	Micron Technology Inc.	Technology
NVDA	NVIDIA Corporation	Technology
NXPI	NXP Semiconductors N.V.	Technology
PANW	Palo Alto Networks Inc.	Technology
QCOM	QUALCOMM Incorporated	Technology
WDAY	Workday Inc.	Technology
ZS	Zscaler Inc.	Technology

B.1 Diffusion-Kernel Construction Benchmark

The first question is whether the fitted relation reflects target-anchored joint reconstruction or only pairwise proximity. The RBF comparator defined in Section 4.1 retains the pairwise quadratic Wasserstein distances but changes how those distances become peer weights. Its implementation uses the median off-diagonal squared distance as the bandwidth, excludes self-links, and row-normalizes the resulting dense matrix. The pooled comparison estimates this operator beside the barycentric interaction field and the persistent news-link field on the same return panel and bootstrap schedule. Because the empirical laws, candidate universe, and Wasserstein distances remain fixed, the contrast isolates the distance-to-weight rule. The RBF operator weights each source separately by its distance from the target, whereas W^b selects simplex coordinates jointly to reconstruct that fixed target. The comparison is therefore a diagnostic of pairwise proximity versus joint barycentric reconstruction, not a generic model-selection exercise.

B.2 Embedding-Representation Sensitivity

The second question is whether the field comparison is specific to one text representation. The sensitivity roster separates four design margins: output width, model capacity, model family, and information vintage. An encoder maps each article into a vector, whose dimension determines the coordinates entering both the pairwise W_2 distances and the target-anchored reconstruction.

The output-width comparison uses Qwen3-Embedding 8B at full width and at 1,024, 256, and 64 coordinates. Qwen’s Matryoshka training is designed to keep leading coordinate prefixes informative, so the shorter vectors test how much compression the interaction geometry tolerates. The common 1,024-coordinate Qwen3-Embedding 8B and 4B rows then vary model capacity while holding output width fixed. Their full-width comparison changes

capacity and output width together, so it does not isolate capacity. The 1,024-coordinate BGE-large-v1.5 row changes model family and pretraining relative to Qwen3-Embedding 8B, making it an external sensitivity check rather than an identified training-mechanism effect (Zhang et al., 2025; Xiao et al., 2024).

The information-vintage comparison uses the three 320-coordinate EttaX encoders. EttaX V0 and V1 use Wikipedia snapshots from 20 December 2017 and 20 December 2020, respectively. The V3 snapshot, dated 1 August 2026, postdates both the article-construction and return-evaluation windows. It is therefore a deliberately post-window negative control, not a point-in-time specification. The three encoders are matched on architecture, optimization recipe, compute budget, and training-token budget; only the Wikipedia snapshot changes. Their capacity differs materially from the production Qwen encoders, so the vintage rows remain descriptive sensitivities rather than identified vintage effects.

Every row in Table 7 uses the same article inputs through 2022, the same 2023–2026 return dates, and one common joint-date bootstrap schedule. Pairing each contrast within the same resample removes common bootstrap variation, while the Holm correction accounts for testing several representations in each family. The table reports direct differences in ρ for the barycentric field and changes in its ρ gap relative to RBF. For representation r , write $g_r = \rho_r^b - \rho_r^h$. The two candidate-minus-reference estimands are $\rho_c^b - \rho_r^b$ and $g_c - g_r$; negative values indicate, respectively, weaker fitted feedback for the barycentric field and a smaller advantage over the matching RBF operator. The persistent co-mention graph is embedding-free and is therefore reported once. The equal-active-support comparator replaces the fitted coordinate magnitudes with uniform weights on the same selected support, so it asks whether the barycentric coordinate magnitudes improve conditional spatial fit beyond peer selection.

The contrast table states one joint equivalence rule for the vintage rows and reports its two benchmark-calibrated bounds. A pair meets the rule only when the adjusted intervals for both estimands lie strictly inside their corresponding bounds. These post-specified thresholds organize a descriptive sensitivity exercise; they are not prespecified confirmatory equivalence margins.

Encoder	Dim.	Interaction matrix	$\hat{\rho}$ [95% CI]	$\hat{\lambda}$	Log-likelihood
Qwen3-8B full (primary)	4096	Barycentric W^b	0.776 [0.743, 0.807]	3.46	110994.7
		RBF–Wasserstein W^h	0.726 [0.680, 0.768]	2.65	108580.9
Qwen3-4B full	2560	Barycentric W^b	0.779 [0.746, 0.809]	3.53	110909.4
		RBF–Wasserstein W^h	0.726 [0.679, 0.768]	2.64	108564.9
Qwen3-8B@1024	1024	Barycentric W^b	0.776 [0.743, 0.807]	3.47	110963.8
		RBF–Wasserstein W^h	0.726 [0.680, 0.768]	2.65	108583.9
Qwen3-4B@1024	1024	Barycentric W^b	0.777 [0.743, 0.808]	3.49	110932.3
		RBF–Wasserstein W^h	0.726 [0.680, 0.768]	2.65	108580.4
BGE-large-v1.5 full	1024	Barycentric W^b	0.772 [0.738, 0.804]	3.38	110611.1
		RBF–Wasserstein W^h	0.725 [0.678, 0.767]	2.63	108562.7
Qwen3-8B@256	256	Barycentric W^b	0.771 [0.737, 0.803]	3.36	110881.2
		RBF–Wasserstein W^h	0.728 [0.683, 0.769]	2.68	108603.7
Qwen3-8B@64	64	Barycentric W^b	0.749 [0.713, 0.784]	2.98	110726.6
		RBF–Wasserstein W^h	0.732 [0.689, 0.772]	2.73	108669.8
EttaX V0	320	Barycentric W^b	0.756 [0.719, 0.792]	3.10	109561.5
		RBF–Wasserstein W^h	0.720 [0.672, 0.764]	2.58	108501.7
EttaX V1	320	Barycentric W^b	0.757 [0.720, 0.793]	3.11	109684.4
		RBF–Wasserstein W^h	0.721 [0.673, 0.764]	2.58	108505.4
EttaX V3	320	Barycentric W^b	0.756 [0.719, 0.793]	3.10	109571.8
		RBF–Wasserstein W^h	0.721 [0.673, 0.764]	2.58	108503.5
Embedding-free	–	Persistent news W^{news}	0.606 [0.557, 0.657]	1.54	110278.5
Primary support	–	Equal active support	0.744 [0.705, 0.780]	2.91	108987.4

Table 7: Pooled 2023–2026 representation sensitivity of the single-field spatial horse race across seven base representations and three matched EttaX encoder vintages. Each encoder row compares the barycentric interaction field W^b , obtained by target-anchored Wasserstein-barycentric reconstruction, with the median-bandwidth RBF transform of its matching squared- W_2 matrix. Intervals use the same 2,000 joint-date stationary bootstrap on the common 885-date schedule. $\hat{\lambda} = \hat{\rho}/(1 - \hat{\rho})$ is the implied adjustment index; log-likelihood is descriptive. Persistent news co-mentions are embedding-free; equal active support is shown once for the primary barycentric interaction field. These are paired conditional-fit sensitivities, not a representation-selection test.

Contrast ($c - r$)	Estimate	95% CI	Raw p	Holm p	Bound	Joint equiv.
<i>Panel A: direct barycentric-ρ contrasts</i>						
Qwen3-4B full – Qwen3-8B full	0.003	[0.002, 0.004]	0.001	0.007	–	–
Qwen3-4B@1024 – Qwen3-8B@1024	0.001	[-0.001, 0.003]	0.247	0.494	–	–
Qwen3-8B@1024 – Qwen3-8B full	0.000	[-0.001, 0.001]	0.624	0.624	–	–
Qwen3-8B@256 – Qwen3-8B@1024	-0.005	[-0.007, -0.004]	0.001	0.007	–	–
Qwen3-8B@64 – Qwen3-8B@256	-0.022	[-0.025, -0.018]	0.001	0.007	–	–
BGE-large-v1.5 full – Qwen3-8B@1024	-0.004	[-0.007, -0.002]	0.001	0.007	–	–
BGE-large-v1.5 full – Qwen3-4B@1024	-0.005	[-0.008, -0.003]	0.001	0.007	–	–
EttaX V1 – EttaX V0	0.001	[-0.001, 0.002]	0.504	1.000	0.007	Yes
EttaX V3 – EttaX V0	-0.000	[-0.003, 0.002]	0.875	1.000	0.007	Yes
EttaX V3 – EttaX V1	-0.001	[-0.003, 0.001]	0.570	1.000	0.007	Yes
<i>Panel B: barycentric-minus-RBF gap contrasts</i>						
Qwen3-4B full – Qwen3-8B full	0.004	[0.003, 0.004]	0.001	0.007	–	–
Qwen3-4B@1024 – Qwen3-8B@1024	0.001	[-0.001, 0.003]	0.273	0.546	–	–
Qwen3-8B@1024 – Qwen3-8B full	0.000	[-0.001, 0.001]	0.786	0.786	–	–
Qwen3-8B@256 – Qwen3-8B@1024	-0.007	[-0.010, -0.006]	0.001	0.007	–	–
Qwen3-8B@64 – Qwen3-8B@256	-0.026	[-0.030, -0.022]	0.001	0.007	–	–
BGE-large-v1.5 full – Qwen3-8B@1024	-0.003	[-0.004, -0.001]	0.003	0.009	–	–
BGE-large-v1.5 full – Qwen3-4B@1024	-0.004	[-0.006, -0.001]	0.001	0.007	–	–
EttaX V1 – EttaX V0	-0.000	[-0.001, 0.002]	0.982	1.000	0.004	Yes
EttaX V3 – EttaX V0	-0.001	[-0.003, 0.001]	0.669	1.000	0.004	Yes
EttaX V3 – EttaX V1	-0.001	[-0.003, 0.002]	0.672	1.000	0.004	Yes

Table 8: Paired common-schedule QMLE contrasts. Panel A estimates $\rho_c^b - \rho_r^b$; Panel B estimates $(\rho_c^b - \rho_c^h) - (\rho_r^b - \rho_r^h)$. Raw p -values are two-sided add-one sign-tail probabilities, and Holm adjustment is applied separately within each metric’s seven base and three vintage contrasts. Vintage equivalence requires both intervals to lie strictly inside their respective BGE-large-minus-Qwen3-8B@1024 benchmark-calibrated, post-specified bounds; the decision shown is joint. No contrast is causal or an encoder-superiority claim.

The intervals for moderate Qwen3-8B truncation from full width to 1,024 coordinates contain zero for both paired estimands. The subsequent 1,024-to-256 and 256-to-64 steps yield lower fitted feedback for the barycentric field and smaller gaps relative to RBF after Holm adjustment. The full-width 4B cell raises both estimands, whereas the fixed-width 4B comparison includes zero; the native-width comparison therefore cannot be read as a pure capacity effect. BGE-large lowers both estimands relative to Qwen3-8B@1,024, but the comparison does not isolate architecture.

All three EttaX pairs meet the table’s joint equivalence rule, including both contrasts involving the post-window V3 negative control. This stability shows that the reported comparison is not uniquely tied to the two earlier EttaX snapshots; it does not make V3 a valid point-in-time encoder. Log-likelihood remains descriptive throughout.

B.3 Annual Joint-Field Persistence

Does the pooled division between barycentric and news-link fields recur in shorter samples? Table 9 re-estimates both coefficients within each non-overlapping annual return sample while holding the two 2018–2022 matrices fixed.

Table 9: Annual joint estimates under the frozen barycentric and news-link fields. The 2026 period ends on 15 July.

Evaluation year	$\hat{\rho}_B$	$\hat{\rho}_N$	$\hat{\lambda}_B$	$\hat{\lambda}_N$
2023	0.600	0.158	2.47	0.65
2024	0.567	0.161	2.09	0.59
2025	0.567	0.218	2.63	1.01
2026	0.476	0.286	2.00	1.20

Both channel coefficients remain positive in every year, and the barycentric estimate is larger throughout. The gap narrows in the incomplete 2026 period, which uses only 133 dates and is not directly comparable with a complete-year estimate. The table therefore documents descriptive persistence, not structural constancy or a formal test of equality across years.

B.4 Joint-Field Estimation and Restricted-Null Simulation

The fourth question concerns implementation, boundary inference, and field admissibility. The main text states the estimand and inference design; this subsection begins with the details needed to reproduce them. The pooled sample uses the fixed base seed, and each successive annual fold adds one to that seed in period order. Within a period, the resulting stationary-bootstrap bank is shared by the joint fit and both single-field boundaries, so corresponding intervals compare the same joint-date resamples. These interval draws preserve the cross-section observed on each resampled date. Numerical-Hessian standard errors from the one-field QMLE are retained only as diagnostics; all reported intervals use bootstrap quantiles.

The two-field search profiles the mixture weight and total feedback on a 401 by 1000 grid. The maximum is then refined continuously within its grid cell, and every bootstrap draw receives the same refinement. Consequently, reported coefficients are not restricted to grid nodes and their intervals do not inherit the grid spacing. On the grid, the Jacobian term is evaluated from the eigenvalue spectrum of each field mixture, which amortizes the calculation across candidate values of total feedback. The continuous refinement instead evaluates each candidate by direct LU factorization, which is cheaper than computing a new eigendecomposition at every isolated point. Tests verify that the two determinant routes agree to floating-point tolerance. This agreement is a numerical cross-check; the routes do not define different estimators.

The QLR calibration is used only for the nested restrictions $\rho_B = 0$ and $\rho_N = 0$, under which the joint model reduces to one of its single-field boundaries. It is not applied to the non-nested comparison between the RBF and barycentric fields. For each nested null, the procedure first recovers the residual series from the corresponding restricted fit. For null draw b , each asset receives a stationary-block path from the shared bank by offsetting the path index by asset and wrapping cyclically. Cycling uses each precomputed path equally often across the null draws. The asset-specific offsets preserve each series' temporal blocks while removing the excluded field's contemporaneous residual alignment under the null. The resampled innovations are mapped back through the restricted spatial inverse, after which both the joint and restricted models are refitted with the same continuous refinement. The reported probability is $(1 + \#\{QLR_k^* \geq QLR_k\})/(B + 1)$ over the null draws. Here, B is the number of null draws. This restricted-null simulation accommodates the nonnegative-channel boundary; no chi-square reference is used.

The remaining checks establish that the constructed field and the joint specification retain the peer-average interpretation required by the adjustment model.

B.5 Compatibility of the Barycentric Interaction Field

The target-anchored Wasserstein barycentric reconstruction returns one simplex row for each firm, and the main text uses the resulting W^b as an economically admissible peer-average field. The required algebra follows directly from the leave-one-out simplex rather than from an empirical normalization: nonnegativity and unit row sums come from Δ_{-i} , the zero diagonal comes from excluding the target, and the ℓ^∞ norm is the maximum absolute row sum.

Proposition 4 (SAR compatibility). *Every operator satisfying Definition 4 is entrywise nonnegative, row-stochastic, and zero-diagonal. In the induced ℓ^∞ operator norm, $\|W^b\| \leq 1$; hence $|\rho| < 1$ implies $\|\rho W^b\| < 1$ and is sufficient for the SAR resolvent.*

The proposition verifies that the constructed field can enter the spatial multiplier without further normalization. It concerns the fitted distributional spanning weights and their matrix properties.

B.6 Why the Two Fields Are Not Orthogonalized

The two supplied fields enter jointly because each must remain a nonnegative, unit-sum peer-average operator if its fitted coefficient is to retain the model’s economic interpretation. Orthogonalizing one matrix against the other would sacrifice that admissibility for algebraic separation. A natural residualization would replace W^b by

$$W^b - \frac{\langle W^b, W^{\text{news}} \rangle_F}{\|W^{\text{news}}\|_F^2} W^{\text{news}}.$$

The Frobenius residual is generally neither entrywise nonnegative nor row-stochastic, so it fails Definition 1 and $\sum_j W_{ij} B_j$ ceases to be a peer-weighted average in exposure units. The construction would therefore buy statistical separation by discarding the economic interpretation that Theorem 3 supplies, and its coefficient would not map to an adjustment intensity. Residualizing the induced signals $W^b r_t$ and $W^{\text{news}} r_t$ instead preserves return units but occurs only after the contemporaneous r_t has entered both signals. It therefore does not produce an alternative pair of predetermined admissible fields for the spatial likelihood. The joint model is used in preference to both alternatives because it leaves each supplied field admissible and estimates their coefficients within the same spatial equilibrium.

C Formal Verification Status

The generated claim manifest records the scope and hypotheses of each machine-checked declaration. It distinguishes algebraic results from assumptions about the empirical operator and from statistical interpretation of the return estimates.

- Proposition 4: Machine-checked for the exact finite operator contract in the pinned Lean 4 / mathlib v4.31.0 workspace. This verifies SAR compatibility, not the economic validity of the text-derived weights.

- Definition 5: Formalized as the exact quantitative premise consumed by the convergence bridge. The declaration does not claim that a parquet operator satisfies the premise and does not formalize general Perron–Frobenius theory.
- Theorem 6: Machine-checked under standard axioms only. This verifies that a quantitative contraction certificate is sufficient for the Perron limit; the empirical float64 certificate remains a separate numerical witness.
- Proposition 3: Machine-checked under standard axioms only in the maximum absolute row-sum matrix norm. It is a finite-rho algebraic bound, not a statistical confidence interval.
- Theorem 7: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace, conditional on the assumed Perron–Frobenius limit hypothesis.
- Theorem 8: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace, conditional on the assumed Perron–Frobenius limit hypothesis.
- Theorem 10: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace, conditional on the single assumed Perron–Frobenius power-limit hypothesis. The resolvent limit it previously also assumed is now derived from that hypothesis rather than assumed alongside it.
- Theorem 9: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace under standard axioms only. The derivation is elementary linear algebra via an absorbing-idempotent resolvent split; it introduces no Perron–Frobenius content of its own, no Abel/Tauberian summation, and no spectral theory.
- Lemma 1: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace by a completing-square identity rather than a differentiability argument, so the equivalence holds in any real inner-product exposure space.
- Theorem 1: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace. The spatial equation is derived from the stated adjustment problem; it is not imposed as a reduced form, and the interaction weights themselves are a maintained input.
- Corollary 2: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace. Admissibility of the mixture is what lets every one-field result transfer; it says nothing about whether either field is economically correct.
- Theorem 3: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace by reduction to the one-field closure at the intensity-weighted mixture. The two-channel equation is derived from the stated adjustment problem; both fields remain maintained inputs, and the result does not establish that they are separately identified in any sample.
- Proposition 1: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace. The inversion is an identity of the quadratic model; reading a fitted coefficient through it presumes the return-bridge restriction stated in the identification section.
- Theorem 2: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace for Hilbert-valued exposures under the displayed norm gate.

- Proposition 2: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace for scalar projections of the systematic exposure equation. Its return-QMLE interpretation remains conditional on the maintained innovation specification and is not causal.
- Corollary 1: Machine-checked unconditionally in the pinned Lean 4 / mathlib v4.31.0 workspace: at zero feedback, peer-adjusted exposure equals stand-alone exposure exactly. This verifies the exposure identity only; it does not import the related paper’s transmission or covariance assumptions.
- Theorem 4: Machine-checked as an exact equality in the pinned Lean 4 / mathlib v4.31.0 workspace, with no optimal-coupling existence theorem required. It identifies the pairwise quadratic Wasserstein term as the two-firm case of the multi-firm dispersion functional.
- Theorem 5: Machine-checked in the pinned Lean 4 / mathlib v4.31.0 workspace for a nonnegative row-stochastic matrix centered at strictly positive stationary probability weights. No reversibility, symmetry, or spectral premise is used; the upper factor of one is exact and the lower factor is a universal worst case.

Together, these records close the algebraic audit trail without extending formal verification to empirical identification, representation validity, or causal interpretation; those remain governed by the statistical design and limitations stated in the paper.