

A note on markets with semi-static trading strategies*

Miklós Rásonyi[†]

September 1, 2026

Abstract

We investigate arbitrage in a discrete-time financial market model where, in addition to finitely many dynamically traded assets, there are also static options to choose from. We introduce the concept of *small cones* of random variables and present a sufficient condition for the attainable positions in the market to be closed in probability.

A fundamental theorem of asset pricing is shown in the present context. Utility maximization will also be considered. We will provide economically meaningful examples of infinite dimensional small cones to demonstrate the pertinence of our approach.

Keywords: arbitrage; static options; fundamental theorem of asset pricing; utility maximization

JEL Classification: D52, G11, G12

1 Introduction

In [1] a discrete-time model of a financial market was considered where, in addition to dynamically traded assets one could invest into static options as well: the investor could buy these at time 0 and receive their payoff at time T . See the bibliography of [1] for further literature about such models.

The study [1] drew attention to the mathematical difficulties associated with such models. Investigations in arbitrage theory are concerned with certain infinite-dimensional cones in topological vector spaces: the cone of attainable positions in the given market. It is pivotal for this theory that the cone of superhedgeable claims should form a *closed* set.

In this respect, [1] conveyed an intimidating message: if the set of available options coincides with all contingent claims that have finite price under a given pricing measure then the set of superhedgeable claims may *fail* to be closed, see Remark 3.2 below for a detailed discussion.

In this note we point out that, if the cone of available options is *small* in a technical sense to be defined, closedness does hold in numerous economically meaningful settings.

*The author gratefully acknowledges the support of the National Research, Development and Innovation Office (NKFIH) through grant K 143529.

[†]HUN-REN Alfréd Rényi Institute of Mathematics, Budapest, Hungary; rasonyi@renyi.hu

In Section 2 we investigate the closedness of cones of random variables and related technicalities. In Section 3 we recall a market with finitely many dynamically traded assets and static options. Absence of arbitrage is characterized in Theorem 3.5 under the hypothesis that the available set of options is a finite sum of small cones. Section 4 treats utility maximization in the same model. Section 5 presents examples of small cones. Some conclusions are drawn in Remark 5.10.

2 Cones of random variables

For $x \in \mathbb{R}$, the notation x^+/x^- refers to positive/negative parts. Let $(\Omega, \mathcal{F}, P)$ be the probability space we are working on, expectation is denoted by $E[\cdot]$. We denote by L^0 the vector space of all (almost sure equivalence classes of) $\mathbb{R}$ -valued random variables, equipped with the topology of convergence in probability. L^∞ will stand for the set of almost surely bounded elements in L^0 , equipped with the essential supremum norm $\|\cdot\|_\infty$. L_+^0 (resp. L_+^∞) refers to the set of non-negative elements in L^0 (resp. L^∞). For $H \subset L^0$, its closure will be denoted $\bar{H}$. For a probability $P' \ll P$ we denote by $E_{P'}[\cdot]$ the corresponding expectation and by $L^1(P')$ the usual Banach-space of P' -integrable random variables.

A subset $C \subset L^0$ is called a *cone* if, for all $\xi_1, \xi_2 \in C$ and $\lambda_1, \lambda_2 \geq 0$, also $\lambda_1 \xi_1 + \lambda_2 \xi_2 \in C$. For a subset $H \subset L^0$, $\mathcal{C}(H)$ denotes the cone generated by H . For $H_1, H_2 \subset L^0$ we denote $H_1 + H_2 := \{h_1 + h_2 : h_1 \in H_1, h_2 \in H_2\}$, $H_1 - H_2$ is then self-explanatory. A subset $H \subset L^0$ is called *bounded* if

$$\sup_{\xi \in H} P(|\xi| \geq n) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We recall two beautiful theorems that are crucial for the present paper.

Lemma 2.1. *If $H \subset L^0$ is convex and bounded then there is $R \sim P$ with $dR/dP \in L^\infty$ and $\sup_{h \in H} E_R[|h|] < \infty$.*

Proof. In [9] this is proved under the supplementary condition $0 \in H$. To see the general case, pick some $h_0 \in H$ and consider $\tilde{H} := \{h - h_0 : h \in H\}$ which is also convex, bounded and contains 0. By the result of [9], there is $\tilde{R}$ with $d\tilde{R}/dP$ bounded and $\sup_{h \in \tilde{H}} E_{\tilde{R}}[|h|] < \infty$. Now define $dR/d\tilde{R} := e^{-|h_0|}/E[e^{-|h_0|}] \leq M_0 < \infty$. Clearly, dR/dP is bounded and

$$\sup_{h \in H} E_R[|h|] \leq \sup_{h \in \tilde{H}} E_R[|h| + |h_0|] \leq M_0 \sup_{h \in \tilde{H}} E_{\tilde{R}}[|h|] + E_R[|h_0|] < \infty,$$

by the choice of $\tilde{R}$ and R . □

Lemma 2.2. *Let $H \subset L^0$ be bounded and convex. Let $\xi_n \in H$, $n \in \mathbb{N}$. Then there exists a subsequence n_k and $\xi_* \in L^0$ such that*

$$\frac{\xi_{n_1} + \dots + \xi_{n_k}}{k} \rightarrow \xi_*$$

almost surely.

Proof. Apply Lemma 2.1. Since H is bounded in $L^1(R)$, Theorem 1 of [4] implies the statement. □

We introduce a rather technical concept which, however, will prove to be mathematically convenient.

Definition 2.3. A set $L \subset L^0$ is called a small cone if there exists a convex, closed and bounded set H with $0 \notin H$ such that $L = \mathcal{C}(H)$. Such an H is called a generator of L .

It is a well-known fact that, in a finite-dimensional Euclidean space, a cone generated by a closed, bounded set away from the origin is closed (see Corollary 9.6.1 of [7]). In the current, infinite dimensional setting, small cones have similar, favourable closedness properties.

Lemma 2.4. Every small cone is closed. If L is a small cone and $C \subset L^0$ is an arbitrary closed cone with $C \cap (-L) = \{0\}$ then $C + L$ is also closed.

Proof. The first statement follows from the second with the choice $C = \{0\}$ so we only prove the latter. Let H be a generator of L as in Definition 2.3 and let R be as given by Lemma 2.1 for this choice of H , set $S := \sup_{h \in H} E_R[|h|] < \infty$.

Let $c_n + \alpha_n h_n \rightarrow \zeta$ in probability as $n \rightarrow \infty$, where $c_n \in C$, $h_n \in H$, $\alpha_n \geq 0$. If $\sup_n \alpha_n < \infty$ then there is a subsequence n_k such that both $\alpha_{n_k} \rightarrow \alpha_* \geq 0$, $k \rightarrow \infty$ and $\sum_{j=1}^k h_{n_j}/k \rightarrow h_* \in H$ hold almost surely, by Lemma 2.2. Then for every ε there is $N(\varepsilon) \geq 1$ such that for $j \geq N(\varepsilon)$, $|\alpha_{n_j} - \alpha_*| \leq \varepsilon$. For $k \geq N(\varepsilon) + 1$,

$$\begin{aligned} & \left| \frac{1}{k} \sum_{j=1}^k \alpha_{n_j} h_{n_j} - \alpha_* h_* \right| \\ & \leq \left| \frac{1}{k} \sum_{j=1}^k \alpha_{n_j} h_{n_j} - \frac{1}{k} \sum_{j=1}^k \alpha_* h_{n_j} \right| + \left| \frac{1}{k} \sum_{j=1}^k \alpha_* h_{n_j} - \alpha_* h_* \right| \\ & \leq \left| \frac{1}{k} \sum_{j=N(\varepsilon)+1}^k \alpha_{n_j} h_{n_j} - \frac{1}{k} \sum_{j=N(\varepsilon)+1}^k \alpha_* h_{n_j} \right| + \frac{1}{k} \left| \sum_{j=1}^{N(\varepsilon)} \alpha_{n_j} h_{n_j} \right| \\ & + \frac{\alpha_*}{k} \left| \sum_{j=1}^{N(\varepsilon)} h_{n_j} \right| + \alpha_* \left| \frac{1}{k} \sum_{j=1}^k h_{n_j} - h_* \right|. \end{aligned}$$

For fixed ε , the third and fourth terms tend to 0 in probability as $k \rightarrow \infty$. As to the second term,

$$E_R \left[\left| \sum_{j=1}^{N(\varepsilon)} \alpha_{n_j} h_{n_j} \right| \right] / k \leq N(\varepsilon) S \sup_n \alpha_n / k,$$

so this term also tends to 0 in probability when $k \rightarrow \infty$ (keeping ε fixed). As to the first term,

$$\begin{aligned} & E_R \left[\left| \sum_{j=N(\varepsilon)+1}^k \alpha_{n_j} h_{n_j} - \sum_{j=N(\varepsilon)+1}^k \alpha_* h_{n_j} \right| \right] \\ & \leq \sum_{j=N(\varepsilon)+1}^k |\alpha_{n_j} - \alpha_*| E_R[|h_{n_j}|] \leq \varepsilon S k \end{aligned}$$

so by Markov's inequality, for every $\eta > 0$,

$$P\left(\left|\frac{1}{k}\sum_{j=N(\varepsilon)+1}^k \alpha_{n_j} h_{n_j} - \frac{1}{k}\sum_{j=N(\varepsilon)+1}^k \alpha_* h_{n_j}\right| \geq \eta\right) \leq \varepsilon S/\eta,$$

which tends to 0 as $\varepsilon \rightarrow 0$. It follows from these arguments that $\frac{1}{k}\sum_{j=1}^k \alpha_{n_j} h_{n_j}$ tends to $\alpha_* h_*$ in probability. Since

$$\frac{1}{k}\sum_{j=1}^k c_{n_j} + \frac{1}{k}\sum_{j=1}^k \alpha_{n_j} h_{n_j} \rightarrow \zeta, \quad k \rightarrow \infty,$$

necessarily

$$\frac{1}{k}\sum_{j=1}^k c_{n_j} \rightarrow c_*,$$

in probability for some $c_* \in C$, since C is closed. But then $\zeta = c_* + \alpha_* h_* \in C + L$.

In the case $\sup_n \alpha_n = \infty$ there is a subsequence n_k with $0 < \alpha_{n_k} \rightarrow \infty, k \rightarrow \infty$ and $\sum_{j=1}^k h_{n_j}/k \rightarrow h_*$ almost surely for some $h_* \in H$. Since

$$\frac{c_{n_k}}{\alpha_{n_k}} + h_{n_k} \rightarrow 0, \quad k \rightarrow \infty,$$

also

$$\frac{1}{k}\sum_{j=1}^k \frac{c_{n_j}}{\alpha_{n_j}} + \frac{1}{k}\sum_{j=1}^k h_{n_j} \rightarrow 0, \quad k \rightarrow \infty,$$

thus $\frac{1}{k}\sum_{j=1}^k \frac{c_{n_j}}{\alpha_{n_j}} \rightarrow -h_* \in C$ in probability, as $k \rightarrow \infty$. But then

$$-h_* \in C \cap (-H) \subset C \cap (-L) = \{0\},$$

which contradicts $0 \notin H$. We conclude that $\sup_n \alpha_n = \infty$ is not possible hence $C + L$ is closed. $\square$

Remark 2.5. Clearly, not all cones are small. One may consider $L_+^\infty \subset L^0$. This is generated by the convex set $H := \{g \in L_+^0 : \|g\|_\infty \leq 1\}$ which is closed and bounded in L^0 (but contains 0). However, L_+^∞ is clearly not closed in L^0 so it is not a small cone.

A subspace $L \subset L^0$ is never a small cone. Indeed, arguing by contradiction, let H be a generator of L . Take an arbitrary $h_0 \in H$. If L is a subspace then also $-h_0 \in L$ hence $h_0 = -\alpha h_1$ for some $h_1 \in H$ and $\alpha \geq 0$. Then, by convexity of H , $\frac{h_0 + \alpha h_1}{1 + \alpha} = 0 \in H$, which contradicts Definition 2.3.

3 Market with dynamically traded assets and static options

Let $\mathcal{F}_t, t = 0, \dots, T$ be a discrete-time filtration on the given probability space, $\mathcal{F}_0$ is the trivial sigma-algebra. Let $X_t, t = 0, \dots, T$ be an adapted $\mathbb{R}^d$ -valued process describing the discounted prices of d dynamically traded assets. We denote by $\langle \cdot, \cdot \rangle$ the standard scalar product in $\mathbb{R}^d$.

A trading strategy is a $\mathbb{R}^d$ -valued process $\phi_t, t = 1, \dots, T$ such that ϕ_t is $\mathcal{F}_{t-1}$ -measurable, the set of all such strategies is denoted Φ . We further assume that a set of zero-price static options $\mathcal{Y} \subset L^0$ is given. These are not liquid, an investor with initial capital 0 can choose one element $h \in \mathcal{Y}$ at time 0 and then his/her portfolio value at time $t = 0, \dots, T$ will be

$$V(\phi, h) := h + \sum_{j=1}^t \langle \phi_j, X_j - X_{j-1} \rangle.$$

We will also write $W_t(\phi) := \sum_{j=1}^t \langle \phi_j, X_j - X_{j-1} \rangle, 0 \leq t \leq T$.

Denote

$$K_0 := \{V(\phi, 0) : \phi \in \Phi\} = \{W_T(\phi) : \phi \in \Phi\},$$

the set of portfolio values in the dynamically traded assets. Then $K_0 + \mathcal{Y}$ describes all attainable positions from 0 initial capital.

We say that **NA** holds is $K_0 \cap L_+^0 = \{0\}$. We say that **NA**($\mathcal{Y}$) holds if

$$(K_0 + \mathcal{Y}) \cap L_+^0 = \{0\}.$$

It is well-known that, under **NA**, the set $K_0 - L_+^0$ is closed in L^0 , see Theorem 6.9.2 of [2].

$\mathcal{M}$ is the set of probabilities $Q \sim P$ on $(\Omega, \mathcal{F})$ such that $X_t, t = 0, \dots, T$ is a Q -martingale with respect to the given filtration. Define

$$\mathcal{M}_\infty = \{Q \in \mathcal{M} : dQ/dP \in L^\infty\}.$$

Analogously,

$$\mathcal{M}(\mathcal{Y}) := \{Q \in \mathcal{M} : \text{for all } Y \in \mathcal{Y}, Y \in L^1(Q) \text{ and } E_Q[Y] \leq 0\}$$

and $\mathcal{M}_\infty(\mathcal{Y}) := \{Q \in \mathcal{M}(\mathcal{Y}) : dQ/dP \in L^\infty\}$.

Clearly, **NA**($\mathcal{Y}$) implies **NA** and the latter is equivalent to $\mathcal{M}_\infty \neq \emptyset$, see Theorem 6.1.1 of [2]. It is clear that $\mathcal{M}(\mathcal{Y}) \neq \emptyset \Rightarrow \mathbf{NA}(\mathcal{Y})$. One may cherish the hope that **NA**($\mathcal{Y}$) is equivalent to $\mathcal{M}(\mathcal{Y}) \neq \emptyset$. The situation, however, is not so simple.

Example 3.1. Let $d = T = 1$, $X_0 = 0$, X_1 a standard Cauchy random variable, $\mathcal{F} := \mathcal{F}_1 = \sigma(X_1)$; let $\mathcal{Y}$ be the set of zero-mean random variables. If $Q \sim P$ and $E_Q[Y] \leq 0$ for all $Y \in \mathcal{Y}$ then necessarily $Q = P$. But $P \notin \mathcal{M}(\mathcal{Y})$ since X_1 is not P -integrable. Hence $\mathcal{M}(\mathcal{Y}) = \emptyset$. We show that, nevertheless, **NA**($\mathcal{Y}$) holds.

Suppose we had $wX_1 + Y \geq 0$ for some $Y \in \mathcal{Y}$ and $w > 0$. Then $X_1 \geq -Y/w$ so X_1^- would be integrable which is nonsense. $w < 0$ leads to a similar contradiction. Hence $w = 0$ but then $E_P[Y] = 0$ implies $Y = 0$ a.s.

This example shows that we need additional hypotheses on $\mathcal{Y}$ for the equivalence

$$\mathcal{M}(\mathcal{Y}) \neq \emptyset \iff \mathbf{NA}(\mathcal{Y})$$

to hold. Note that a proof would probably require the use of the Kreps-Yan separation theorem which needs, roughly speaking, that the cone $K_0 + \mathcal{Y} - L_+^0$ is closed.

Remark 3.2. We briefly review certain results of [1] in order to put our achievements in context. In that paper $P \in \mathcal{M}$ was assumed and, instead of K_0 , the set

$$\tilde{K}_0 := \{W_T(\phi) : (W_t(\phi))_{0 \leq t \leq T} \text{ is a supermartingale}\} \subset K_0$$

was considered. The authors took $\mathcal{Y} = L^1(\Omega, \sigma(X_T), P)$ (the set of integrable $\sigma(X_T)$ -measurable random variables). They exhibited (in the case $d = 1, T = 2$) a bounded price process for which $\tilde{K}_0 + \mathcal{Y} - L_+^0$ failed to be closed in L^0 (this is not surprising, $\mathcal{Y}$ not being closed in L^0 either), even $(\tilde{K}_0 + \mathcal{Y} - L_+^0) \cap L^1(P)$ failed to be closed in $L^1(P)$. The sequences of random variables they constructed had further specific properties that we do not list here.

The proof of [1] actually shows that $(K_0 + \mathcal{Y} - L_+^0) \cap L^1(P)$ is not closed either. Indeed, in the last step of the proof in Section 3 of [1], one takes $0 \leq g \leq u + v$ with $u \in \tilde{K}_0, v \in \mathcal{Y}$. We point out that even if we take u from the larger set K_0 , necessarily $-v \leq g - v \leq u$ and, since v is integrable, it follows that $u \in \tilde{K}_0$ (a martingale transform with terminal value bounded from below by an integrable random variable is, in fact, a martingale, see Theorems 1 and 2 of [3]) hence the proof of [1] goes through with $\tilde{K}_0$ replaced by K_0 .

There is, however, a conceptual difference between [1] and the present paper. We regard $\mathcal{Y}$ as a family of options available at 0 cost while the choice in [1] is not (and cannot be) interpreted in this way.

Our purpose in the present paper is to provide conditions on $\mathcal{Y}$ that imply closedness of $K_0 + \mathcal{Y} - L_+^0$ in meaningful cases. The set $\mathcal{Y}$ in Remark 3.2 was rather large so choosing smaller sets might lead to success. We propose the following hypothesis.

Assumption 3.3. $\mathcal{Y} = \sum_{j=1}^m C_j$ where each C_j is a small cone and

$$\left(K_0 + \sum_{j=1}^n C_j \right) \cap (-C_{n+1}) = \{0\}$$

for all $0 \leq n \leq m - 1$.

Remark 3.4. Recall that K_0 is a vector space so in the case $m = 1, K_0 \cap (-C_1) = \{0\}$ is equivalent to $K_0 \cap C_1 = \{0\}$ which means that the non-zero static options should not be replicable from 0 initial capital. Hence this is a sort of no-redundancy hypothesis.

Our main result comes next: a fundamental theorem of asset pricing in the present setting.

Theorem 3.5. Let $\mathcal{Y} \subset L^0$ be arbitrary. Consider the following statements.

1. $\mathbf{NA}(\mathcal{Y})$ holds.
2. For each $Y \in \mathcal{Y}$ there is $Q(Y) \in \mathcal{M}_\infty$ such that Y is $Q(Y)$ -integrable and

$$E_{Q(Y)}[Y] \leq 0.$$

3. $\mathcal{M}_\infty(\mathcal{Y}) \neq \emptyset$.
4. The set $K_0 + \mathcal{Y} - L_+^0$ is closed.

Then $3. \rightarrow 2. \leftrightarrow 1.$ always. If Assumption 3.3 holds then $1. \rightarrow 4.$ and also $3. \leftrightarrow 2. \leftrightarrow 1.$

Proof. Suppose that 2. holds and $V(\phi, Y) \geq 0$ almost surely for some $(\phi, Y) \in \mathcal{A}$. Notice that $W_T(\phi) \geq -h$ in this case and, Y being $Q(Y)$ -integrable and $(X_t)_{0 \leq t \leq T}$

being a $Q(Y)$ -martingale, the $Q(Y)$ -martingale transform $(W_t(\phi))_{0 \leq t \leq T}$ has a final value that is above a $Q(Y)$ -integrable random variable hence $(W_t(\phi))_{0 \leq t \leq T}$ is a true $Q(Y)$ -martingale by Theorems 1 and 2 of [3]. Then $E_{Q(Y)}[W_T(\phi)] = 0$ and $E_{Q(Y)}[Y] \leq 0$. Consequently, $E_{Q(Y)}[V(\phi, Y)] \leq 0$ and $V(\phi, Y) = 0$ almost surely: $\mathbf{NA}(\mathcal{Y})$ holds.

Suppose that 1. is true. Let $Y \in \mathcal{Y}$ be arbitrary. Define $dP'/dP := e^{-|Y|}/E[e^{-|Y|}]$, clearly $Y \in L^1(P')$. $\mathbf{NA}(\mathcal{Y})$ implies $\mathbf{NA}$ but, equivalently, also $\mathbf{NA}$ under the probability P' . Hence Theorem 6.1.1 implies the existence of $Q_0 \in \mathcal{M}$ with $dQ_0/dP' \in L^\infty$. If $Y \in K_0$ then $E_{Q_0}[Y] = 0$ (since $Y = W_T(\phi)$ for some ϕ and it is integrable so $W_t(\phi)$, $t = 0, \dots, T$ is a Q_0 -martingale). $dQ_0/dP \in L^\infty$ by construction.

In the case $Y \notin K_0$, $L := \{\alpha Y : \alpha \geq 0\}$ is a small cone with $K_0 \cap (-L) = \{0\}$. We claim that also $(K_0 - L_+^0) \cap (-L) = \{0\}$. Indeed, if $k_0 - l = -\alpha Y$ for some $k_0 \in K_0$, $l \in L$, $\alpha \geq 0$ then $k_0 + \alpha Y = l \in L_+^0$ so $l = 0$ by $\mathbf{NA}(\mathcal{Y})$ and $k_0 = -\alpha Y$ which necessitates $\alpha = 0$. $K_0 - L_+^0$ is closed by Theorem 6.9.2 of [2]. Now by Lemma 2.4 (or by direct argument), $K_0 - L_+^0 + L$ is a closed cone in L^0 .

Theorem 5.2.3 of [2] (with the choice $p = 1$, $C := (K_0 - L_+^0 + L) \cap L^1(Q_0)$ and dual pair $E = L^1(Q_0)$, $E' = L^\infty$) provides a probability $Q(Y) \sim P$ with $dQ(Y)/dQ_0 \in L^\infty$ such that $Q(Y) \in \mathcal{M}_\infty$ and also $E_{Q(Y)}[Y] \leq 0$: 2. follows.

3. trivially implies 2. Now let Assumption 3.3 be in force and assume 1. which implies $\mathbf{NA}$ so $K_0 - L_+^0$ is closed by Theorem 6.9.2 of [2].

Apply Lemma 2.4 inductively. Assume that $K_0 - L_+^0 + \sum_{j=1}^n C_j$ is closed for some $0 \leq n \leq m-1$. Take $\xi = k - l + \sum_{j=1}^n c_j \in K_0 - L_+^0 + \sum_{j=1}^n C_j$. If we had $\xi \in -C_{n+1}$ then $l = k + \sum_{j=1}^n c_j - \xi \in (K_0 + \sum_{j=1}^{n+1} C_j) \cap L_+^0$ so $\mathbf{NA}(\mathcal{Y})$ implies $l = 0$ and then $\xi \in (K_0 + \sum_{j=1}^n C_j) \cap (-C_{n+1}) = \{0\}$, by assumption. As C_{n+1} is a small cone, Lemma 2.4 implies (with the choice $C = K_0 + \sum_{j=1}^n C_j$, $L = C_{n+1}$) that $K_0 + \sum_{j=1}^{n+1} C_j$ is also a closed cone. Finally, $K_0 - L_+^0 + \sum_{j=1}^m C_j$ is shown to be closed and 4. follows.

Now we prove that, under Assumption 3.3, 1. implies 3. We have already seen that $\mathbf{NA}(\mathcal{Y})$ implies 4. Let H_j generate C_j as in Definition 2.3. Lemma 2.1 provides $R_j \sim P$, $j = 1, \dots, m$ with dR_j/dR_{j-1} , $j = 2, \dots, m$ bounded and also dR_1/dP bounded, such that $\sup_{h \in H_j} E_{R_j}[|h|] < \infty$. It follows that $E_{R_m}[|\xi|] < \infty$ for all $\xi \in \sum_{j=1}^m C_j$. Now define

$$dR/dR_m := \exp \left(\sum_{t=0}^T |X_t| \right) / E_{R_m} \left[\exp \left(\sum_{t=0}^T |X_t| \right) \right].$$

Notice that $\langle \zeta, X_t - X_{t-1} \rangle$ are R -integrable for each $1 \leq t \leq T$ and for each $\mathcal{F}_{t-1}$ -measurable d -dimensional bounded random variable ζ .

Theorem 5.2.3 of [2] (with the choice $p = 1$, $C := (K_0 - L_+^0 + \mathcal{Y}) \cap L^1(R)$ and dual pair $E = L^1(R)$, $E' = L^\infty$) provides a probability $Q_* \sim P$ such that $E_{Q_*}[c] \leq 0$ for all $c \in (K_0 - L_+^0 + \mathcal{Y}) \cap L^1(R)$ and dQ_*/dR is bounded (hence dQ_*/dP is also bounded). By the discussion above, $(K_0 - L_+^0 + \mathcal{Y}) \cap L^1(R)$ contains all random variables of the form $\langle \zeta, X_t - X_{t-1} \rangle$ which implies $Q_* \in \mathcal{M}_\infty$. The construction of Q_* ensures that $E_{Q_*}[Y] \leq 0$ for all $Y \in \mathcal{Y}$, 3. follows. $\square$

As already mentioned above, the equivalence 1. $\leftrightarrow$ 3. can be considered as a “right” fundamental theorem of asset pricing for a market with both liquid assets and static options. It asserts that there is a “uniform” pricing measure $Q_* \in \mathcal{M}_\infty(\mathcal{Y})$ which works for all the static options at the same time. Without Assumption 3.3, we

can guarantee only that for each $Y \in \mathcal{Y}$ there is *some* pricing measure $Q(Y) \in \mathcal{M}_\infty$ which may depend on Y : this is the content of 2. in Theorem 3.5.

4 Utility maximization with semi-static strategies

In the discussion following Theorem 1.3 of [1] the authors point out that the failure of the closedness of final outcomes in a market with semistatic trading strategies makes utility maximization a delicate issue in these models. An important consequence of 4. in Theorem 3.5 is that, under Assumption 3.3 and $\mathbf{NA}(\mathcal{Y})$, it is possible to prove the existence of optimal strategies for a utility maximizer. Define $\mathcal{A} := \Phi \times \mathcal{Y}$.

Theorem 4.1. *Let Assumption 3.3 and $\mathbf{NA}(\mathcal{Y})$ hold. Let $u : \mathbb{R} \rightarrow \mathbb{R}$ be non-decreasing, concave and bounded from above. Then there exist $(\phi^*, h^*) \in \mathcal{A}$ such that*

$$\sup_{(\phi, h) \in \mathcal{A}} E[u(V(\phi, h))] = E[u(V(\phi^*, h^*))],$$

that is, the utility maximization problem with dynamic assets and static options is well-posed.

Proof. We follow ideas from Proposition 4.3 of [6]. Let $(\phi_n, h_n), n \in \mathbb{N}$ be a sequence such that

$$\sup_{(\phi, h) \in \mathcal{A}} E[u(V(\phi, h))] = \lim_{n \rightarrow \infty} E[u(V(\phi_n, h_n))]$$

and $E[u(V(\phi_n, h_n))] > -\infty$ for all n . u being bounded from above, necessarily

$$\sup_n E[u^-(V(\phi_n, h_n))] < \infty. \quad (1)$$

The case of constant u is trivial, otherwise there exist $c_0, c_1 > 0$ such that $u(x) \leq -c_0|x| + c_1$ for all $x \leq 0$, see Lemma 4.1 of [6] for this simple fact. Now (1) implies

$$\sup_n E[V^-(\phi_n, h_n)] < \infty.$$

Proposition 2 of [5] implies that there is $\tilde{P} \sim P$ with bounded $d\tilde{P}/dP$ such that $V(\phi_n, h_n)$ are $\tilde{P}$ -integrable for all n .

Take $Q \in \mathcal{M}(\mathcal{Y})$ with $dQ/d\tilde{P}$ bounded, as constructed in Theorem 3.5 (note that $\mathbf{NA}(\mathcal{Y})$ is true under P iff it is true under $\tilde{P}$). Hence $V(\phi_n, h_n)$ are Q -integrable and $\sup_n E_Q[V^-(\phi_n, h_n)] < \infty$. As $E_Q[Z] \leq 0$ for each $Z \in (K_0 + \mathcal{Y} - L_+^0) \cap L^1(\tilde{P})$ and $V(\phi_n, h_n) \in L^1(\tilde{P})$ for all n , it follows that also

$$\sup_n E_Q[V^+(\phi_n, h_n)] \leq \sup_n E_Q[V^-(\phi_n, h_n)] < \infty$$

so, finally, $\sup_n E_Q[|V(\phi_n, h_n)|] < \infty$.

Applying Theorem 1 of [4] in $L^1(Q)$, for some subsequence $n_k, k \in \mathbb{N}$, the Césaro means of $V(\phi_{n_k}, h_{n_k})$ converge to some random variable X almost surely. By 4. in Theorem 3.5, $X \in K_0 + \mathcal{Y} - L_+^0$ so $X \leq V(\phi^*, h^*)$ for some $(\phi^*, h^*) \in \mathcal{A}$. By

concavity of u and the limsup Fatou lemma,

$$\begin{aligned}
E[u(V(\phi^*, h^*))] &\geq E[u(X)] \geq \lim_{k \rightarrow \infty} E \left[u \left(V \left(\frac{1}{k} \sum_{l=1}^k \phi_{n_l}, \frac{1}{k} \sum_{l=1}^k h_{n_l} \right) \right) \right] \\
&\geq \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{l=1}^k E[u(V(\phi_{n_l}, h_{n_l}))] = \lim_{k \rightarrow \infty} E[u(V(\phi_{n_k}, h_{n_k}))] \\
&= \sup_{(\phi, h) \in \mathcal{A}} E[u(V(\phi, h))],
\end{aligned}$$

and the result follows since

$$E[u(V(\phi^*, h^*))] \leq \sup_{(\phi, h) \in \mathcal{A}} E[u(V(\phi, h))],$$

trivially. $\square$

5 Examples

In this section we present several examples of small cones. For $H \subset L^0$, $\text{conv}(H)$ denotes the set of finite convex combinations of elements in H .

Example 5.1. Let $\xi_n, n \in \mathbb{N}$ be a sequence of zero-mean independent random variables such that $J := \sup_n E[\xi_n^2] < \infty$. Let $a_n, n \in \mathbb{N}$ be a sequence of constants satisfying $a_* \leq a_n \leq a_\dagger, n \in \mathbb{N}$ for some $0 < a_* \leq a_\dagger$. Let $H := \{\xi_n + a_n, n \in \mathbb{N}\}$.

For each $h \in \text{conv}(H)$, $h = \sum_{j=1}^k \lambda_j (\xi_j + a_j)$ for some k and convex weights $\lambda_1, \dots, \lambda_k$ so $\text{var}(h) \leq J \sum_{j=1}^k \lambda_j^2 \leq J$ and $E[|h|^2] \leq J + a_\dagger^2$. It follows that $\text{conv}(H)$ is uniformly integrable.

Clearly, $E[h] \geq a_*$ for all $h \in \text{conv}(H)$. If $h_n \in H$ converge almost surely to some $\hat{h}$ also $E[\hat{h}] \geq a_*$ by uniform integrability. This implies $0 \notin H' := \overline{\text{conv}(H)}$. Also note that, for all $h \in \text{conv}(H)$,

$$P(|h| \geq l) \leq E[h^2]/l^2 \leq (J + a_\dagger^2)/l^2 \rightarrow 0$$

as $l \rightarrow \infty$, so $\text{conv}(H)$ and hence $H' := \overline{\text{conv}(H)}$ are bounded in L^0 . We conclude that $\mathcal{C}(H')$ is a small cone.

Such an example is not artificial at all: the set of payoffs in $\mathcal{C}(H')$ are similar to those available in the Arbitrage Pricing Theory model of [8] which can be used to describe one-step *large financial markets*, see also [5, 6].

Example 5.2. In a similar vein, if $H = \{\xi_n, n \in \mathbb{N}\}$ are independent with standard Cauchy distribution then all elements of $H' := \overline{\text{conv}(H)}$ have this same distribution so H' is a closed, bounded convex set not containing 0. $\mathcal{C}(H')$ is thus a small cone.

Example 5.3. Let $\xi_i \in L^0, i = 1, \dots, N$ be linearly independent. Then

$$\mathcal{C}(\{\xi_i, i = 1, \dots, N\})$$

is a small cone. Indeed, $\text{conv}(\{\xi_i, i = 1, \dots, N\})$ is closed in this case (see e.g. Theorem 19.1 of [7]) and it does not contain 0 by linear independence.

In the rest of this section, we enter the setting of Section 3. For simplicity, we assume that the number of dynamically traded assets is $d = 1$ and we will only look at the case $m = 1$ in Assumption 3.3. In the two examples below we provide infinite dimensional small cones with clear financial interpretations.

Example 5.4. Let $0 < K_b < K_\#$ and consider the call options

$$\{(X_T - K)^+ : K_b \leq K \leq K_\#\}.$$

Take a mapping $Q : [K_b, K_\#] \rightarrow \mathcal{M}$ such that

$$D := \sup_K E_{Q(K)}[(X_T - K)^+] < \infty,$$

that is, the prices of the available call options should be bounded from above: a rather natural assumption.

Define

$$H_1 := \{(X_T - K)^+ - E_{Q(K)}[(X_T - K)^+] : K_b \leq K \leq K_\#\},$$

set $\tilde{H}_1 := \overline{\text{conv}(H_1)}$ and $L_1 := \mathcal{C}(\tilde{H}_1)$.

Proposition 5.5. Assume that $X_T \geq 0$ is unbounded but P -integrable. The set $\tilde{H}_1$ is bounded, does not contain 0 hence L_1 is a small cone.

Proof. Since $-D \leq h \leq X_T$ for all $h \in H_1$ this is also true for all $h \in \text{conv}(H_1)$ so $\tilde{H}_1$ is bounded.

It is easy to construct an auxiliary probability $P^\natural \ll P$ with bounded $dP^\natural/dP$ such that

$$E_{P^\natural}[(X_T - K_\#)^+] \geq D + 1.$$

Indeed, $p := P(X_T - K_\# > D + 1) > 0$. Define $dP^\natural/dP := 1_{\{X_T > D+1\}}/p$.

It follows that, for each $K \in [K_b, K_\#]$,

$$E_{P^\natural}[(X_T - K)^+] \geq E_{P^\natural}[(X_T - K_\#)^+] \geq D + 1 \geq E_{Q(K)}[(X_T - K)^+] + 1,$$

hence $E_{P^\natural}[h] \geq 1$ for all $h \in H$, even for all $h \in \text{conv}(H)$. Now taking $h_n \in \text{conv}(H)$, $h_n \rightarrow h_*$ in L^0 , the limsup Fatou lemma implies $E_{P^\natural}[h_*] \geq 1$ since $E_P[X_T] < \infty$, $dP^\natural/dP$ is bounded and hence X_T is a $P^\natural$ -integrable majorant of the sequence h_n . $\square$

Proposition 5.6. Assume that the $\mathcal{F}_{T-1}$ -conditional law of X_T has almost surely support $[0, \infty)$. Then $K_0 \cap L_1 = \{0\}$.

Proof. We argue by contradiction. If $K_0 \cap L_1 \neq \{0\}$ then there is $h_* \in K_0 \cap \tilde{H}_1$. Let $\lambda_1^n, \dots, \lambda_{k(n)}^n$ be convex weights and assume that

$$h_n := \sum_{j=1}^{k(n)} \lambda_j^n [(X_T - K_j^n)^+ - E_{Q(K_j^n)}[(X_T - K_j^n)^+]] \rightarrow h_*$$

almost surely as $n \rightarrow \infty$, where $K_j^n \in [K_b, K_\#]$, $j = 1, \dots, k(n)$. If one had $h_* \in K_0$ then also $X_T - X_0 - h_* \in K_0$. Notice that

$$K_\# + D - X_0 \geq X_T - h_n - X_0 \geq -X_0, \quad n \in \mathbb{N}$$

so $X_T - X_0 - h_*$ is a bounded random variable.

Since $X_T - X_0 - h_*$ is in K_0 , $X_T - X_0 - h_* = \xi_{T-1} + \phi_{T-1}(X_T - X_{T-1})$ holds for some $\mathcal{F}_{T-1}$ -measurable ξ_{T-1}, ϕ_{T-1} . By our hypothesis on the conditional support, necessarily $\phi_{T-1} = 0$. In other words, $X_T - X_0 - h_*$ and hence also $X_T - h_*$ are $\mathcal{F}_{T-1}$ -measurable.

We will show that this is not possible. On the event $\{X_T \geq K_\# \}$,

$$X_T - h_n = c_n := \sum_{j=1}^{k(n)} \lambda_j^n [K_j^n + E_{Q(K_j^n)}[(X_T - K_j^n)^+]]$$

so $c_n \rightarrow c_*, n \rightarrow \infty$ for some constant c_* . It follows that

$$P(X_T - h_* = c_* | \mathcal{F}_{T-1}) \geq P(X_T \geq K_\# | \mathcal{F}_{T-1}) > 0$$

almost surely by our hypothesis on the conditional support of X_T . On the other hand, on the event $\{X_T \leq K_b/2\}$,

$$h_n = - \sum_{j=1}^{k(n)} \lambda_j^n E_{Q(K_j^n)}[(X_T - K_j^n)^+] \rightarrow d_*, n \rightarrow \infty$$

for some constant d_* . Hence on this event $X_T - h_* \leq K_b/2 - d_*$ holds. Clearly, $c_* \geq K_b - d_*$ hence $X_T - h_* < c_*$ on $\{X_T \leq K_b/2\}$. Notice that

$$P(X_T \leq K_b/2 | \mathcal{F}_{T-1}) > 0$$

almost surely but this contradicts that the conditional law of $X_T - h_*$ with respect to $\mathcal{F}_{T-1}$ is degenerate. $\square$

Example 5.7. Consider now the *put options* $\{(K - X_T)^+ : K_b \leq K \leq K_\#\}$. Take a mapping $Q : [K_b, K_\#] \rightarrow \mathcal{M}$ such that $q := \inf_K E_{Q(K)}[(K - X_T)^+] > 0$ (that is, the prices of the put options cannot be arbitrarily small, a rather natural assumption).

Define

$$H_2 := \{(K - X_T)^+ - E_{Q(K)}[(K - X_T)^+] : K_b \leq K \leq K_\#\},$$

set $\tilde{H}_2 := \overline{\text{conv}(H)}$ and $L_2 := \mathcal{C}(\tilde{H}_2)$.

Proposition 5.8. *If $X_T \geq 0$ is unbounded then L_2 is a small cone.*

Proof. As $-K_\# \leq h \leq K_\#$ for all $h \in H_2$, $\tilde{H}_2$ is bounded. Notice that $\delta := P(X_T \geq K_\#)$ is positive. On the event $\{X_T \geq K_\#\}$,

$$(K - X_T)^+ - E_{Q(K)}[(K - X_T)^+] \leq -E_{Q(K)}[(K - X_T)^+] \leq -q,$$

for all K . Hence for any $h \in \text{conv}(H_2)$, $P(h \leq -q) \geq \delta$. If $h_n \in \text{conv}(H_2) \rightarrow h_*$ in probability as $n \rightarrow \infty$, necessarily $h_* \neq 0$. It follows that $0 \notin \tilde{H}_2$ and $\mathcal{C}(\tilde{H}_2)$ is a small cone. $\square$

Proposition 5.9. *Assume that the $\mathcal{F}_{T-1}$ -conditional law of X_T has almost surely support $[0, \infty)$. Then $K_0 \cap L_2 = \{0\}$.*

Proof. As in the proof of Proposition 5.6, suppose that

$$h_n := \sum_{j=1}^{k(n)} \lambda_j^n [(K_j^n - X_T)^+ - E_{Q(K_j^n)}[(K_j^n - X_T)^+]] \rightarrow h_*.$$

Clearly, h_* is a bounded random variable hence, as before, h_* must be $\mathcal{F}_{T-1}$ -measurable. We will show that this cannot be the case.

Arguing on the event $\{X_T \geq K_\sharp\}$,

$$- \sum_{j=1}^{k(n)} \lambda_j^n E_{Q(K_j^n)}[(K_j^n - X_T)^+] \rightarrow d_*, \quad n \rightarrow \infty$$

for some constant d_* . Hence

$$P(h_* = d_* | \mathcal{F}_{T-1}) \geq P(X_T \geq K_\sharp | \mathcal{F}_{T-1}) > 0 \text{ a.s.}$$

At the same time, on the event $\{X_T \leq K_\flat/2\}$, $h_* \geq K_\flat/2 + d_*$ and

$$P(X_T \leq K_\flat/2 | \mathcal{F}_{T-1}) > 0 \text{ a.s.}$$

This contradicts that h_* has a degenerate conditional law with respect to $\mathcal{F}_{T-1}$. $\square$

Remark 5.10. We have thus shown in Propositions 5.5, 5.6, 5.8 and 5.9 that choosing $\mathcal{Y}$ as the convex closure of either call or put options, the conditions of Theorems 3.5 and 4.1 are met.

Our work suggests that, in markets with both dynamical assets and static options, a satisfactory arbitrage theory can be built up provided that the set of options is not too large, as described by the concept of small cones.

Extending Theorem 4.1 to unbounded utilities requires more involved arguments and will be done elsewhere. The use of small cones in continuous-time models is an exciting direction of future research that could be pursued.

References

- [1] B. Acciaio, M. Larsson, and W. Schachermayer. The space of outcomes of semi-static trading strategies need not be closed. *Finance Stoch.*, 21:741–751, 2017.
- [2] F. Delbaen, and W. Schachermayer. *The mathematics of arbitrage*. Springer, Berlin, Heidelberg, 2006.
- [3] J. Jacod and A. N. Shiryaev. Local martingales and the fundamental asset pricing theorems in the discrete-time case. *Fin. Stoch.*, 2:259–273, 1998.
- [4] J. Komlós. A generalization of a problem of Steinhaus. *Acta Mathematica Academiae Scientiarum Hungaricae*, 18:217–229, 1967.
- [5] M. Rásonyi. Equivalent martingale measures for large financial markets in discrete time. *Mathematical Methods of Operations Research*, 58:401–415, 2003.
- [6] M. Rásonyi. Maximizing expected utility in the Arbitrage Pricing Model. *Journal of Mathematical Analysis and Applications*, 454:127–143, 2017.

- [7] R. T. Rockafellar. *Convex analysis*. Princeton, N.J., 1970.
- [8] S. A. Ross. The arbitrage theory of capital asset pricing. *J. Econom. Theory*, 13:341–360, 1976.
- [9] J.-A. Yan. Caractérisation d’une classe d’ensembles convexes de L^1 ou H^1 . *Séminaire Proba. XIV*. Lecture Notes in Mathematics, vol. 784, 220–222. Springer, Berlin, Heidelberg, New York, 1980.