

Importance Sampling Enhanced with the COS Method for the Portfolio Risk Allocation

Fang Fang^{*}

Xiaoyu Shen[†]

Qinling Wang[‡]

September 1, 2026

Abstract

We study the calibration of importance-sampling proposals for rare portfolio losses and obligor-level tail-risk contributions in multi-factor credit models. Cross-entropy importance sampling approximates the common-state distribution conditional on a tail event, but its standard implementation (CEIS) estimates the required moments from binary tail indicators and can be unstable when few pilot paths reach an extreme threshold. We propose ISCOS, which replaces each indicator by a filtered COS approximation of $q_x(u) = \mathbb{P}(L \geq x \mid U = u)$. Conditional on the common state, defaults are independent and the portfolio-loss characteristic function is available in closed form, so these smooth calibration weights can be computed without an additional default simulation. We derive Gaussian proposals for the systematic factors and Gaussian–inverse-Gamma product proposals for Student t -copulas. With exact conditional probabilities, the ISCOS raw-moment estimator is the Rao–Blackwellisation of CEIS and has no larger covariance matrix. If the filtered COS error is uniformly $O(K^{-p})$, the induced proposal parameters inherit the same order, yielding a fixed-threshold calibration error of $O_{\mathbb{P}}(M_0^{-1/2}) + O(K^{-p})$. On an eleven-factor benchmark, exact conditional block convolution confirms convergence of the Gaussian COS-calibrated moments. Under matched simulation budgets, ISCOS gives higher event-weight effective sample sizes and shorter average nominal pointwise intervals than CEIS in the reported Gaussian and Student t -copula configurations, particularly for tail contributions, while end-to-end costs remain comparable. The Gaussian comparison also includes the paper-aligned Glasserman’s proposal.

Keywords. Credit portfolio; importance sampling; cross-entropy method; Fourier–cosine method; conditional Monte Carlo; rare events; Gaussian copula; Student t -copula; risk allocation.

^{*}FF Quant Advisory B.V., 3531 WR Utrecht, the Netherlands (fang.fang@ffquant.nl) and Delft Institute of Applied Mathematics, Delft University of Technology, 2628 CD Delft, the Netherlands (f.fang@tudelft.nl).

[†]FF Quant Advisory B.V., 3531 WR Utrecht, the Netherlands (xiaoyu.shen@ffquant.nl).

[‡]Delft Institute of Applied Mathematics, Delft University of Technology, 2628 CD Delft, the Netherlands (q.wang-7@tudelft.nl).

Contents

1	Introduction	3
1.1	Related Literature and Methodological Position	3
1.2	Contributions and Paper Organisation	4
2	Credit-Portfolio Models and Tail-Risk Quantities	5
2.1	Factor-copula loss model	5
2.2	Conditional loss distribution	6
2.3	Tail-risk quantities and allocations	7
3	Cross-Entropy Importance Sampling in Factor Space	7
3.1	The factor distribution behind a tail loss	8
3.2	Cross-entropy as a weighted fit	8
3.3	Gaussian proposal for the systematic factors	9
3.4	Student t -copula product proposal	9
3.5	Pilot reweighting and likelihood ratios	11
4	Estimating the Cross-Entropy Proposal	12
4.1	One pilot sample, two weights	12
4.2	Computing the smooth COS weights	13
4.3	Practical calibration	13
5	Error Analysis for ISCOS Calibration	14
5.1	Exact conditional weighting and Rao–Blackwellisation	15
5.2	A uniform COS error for the conditional tail probability	17
5.3	Propagation of the COS error to the CE parameters	19
5.4	Finite-COS covariance and the combined calibration error	21
5.5	Threshold estimation and scope of the result	24
6	Numerical Experiments: Gaussian Copula	25
6.1	Benchmark portfolio and matched experimental design	25
6.2	Accuracy of the filtered COS calibration	27
6.3	Proposal-calibration diagnostics	29
6.4	Risk contributions and nominal pointwise intervals	30
6.5	Computational cost and limitations	31
7	Numerical Experiments: Student t-Copula	33
7.1	Portfolio, Product Proposal, and Reproducible Setup	34
7.2	Proposal Calibration	35
7.3	Risk Contributions and Nominal Pointwise Intervals	36
7.4	Computational Cost	39
7.5	Summary and Limitations of the Student t Experiment	40
8	Conclusion	40

1 Introduction

Credit-portfolio risk measurement is a rare-event problem. For a portfolio loss

$$L = \sum_{n=1}^N l_n Y_n,$$

Value-at-Risk, threshold-based expected shortfall, and their obligor-level allocations depend on events such as $\{L \geq x\}$ and $\{L = x\}$. At a confidence level of 99.9%, an ordinary Monte Carlo sample of size M contains only about $M/1000$ tail observations on average, while the exact-level event used for a VaR contribution may be much rarer. Accurate allocation can therefore require substantially more simulation than estimation of an unconditional portfolio quantity.

Importance sampling makes these events more frequent under the simulation law and corrects the result by a likelihood ratio. In factor-copula credit models, a natural strategy changes the distribution of the common factors and then exponentially twists the conditional Bernoulli defaults. Its performance depends strongly on the factor proposal. A proposal that is too mild produces few useful tail observations; one that is too concentrated can generate unstable likelihood ratios and may fail the moment conditions needed for standard error estimates.

This paper studies how that proposal should be calibrated. For a fixed loss threshold x , the natural target is the distribution of the common state U conditional on $A_x = \{L \geq x\}$, whose density is proportional to

$$f_U(u)q_x(u), \quad q_x(u) = \mathbb{P}(L \geq x \mid U = u).$$

The cross-entropy projection of this distribution onto a Gaussian family is therefore determined by tail-conditional factor moments. Standard cross-entropy importance sampling (CEIS) estimates those moments from the binary indicators $\mathbf{1}_{\{L \geq x\}}$. At an extreme threshold, most pilot observations receive zero weight, making the fitted covariance noisy or nearly singular.

We propose ISCOS, which replaces the indicator by a finite COS approximation $q_{x,K}(U)$ of the conditional tail probability. Conditional on the common state, obligor defaults are independent and the conditional loss characteristic function is available analytically. The COS method can thus integrate out the idiosyncratic default simulation at the calibration stage. Each pilot state is weighted by its chance of producing a tail loss, rather than by the outcome of one conditional Bernoulli draw. CEIS and ISCOS then use the same weighted proposal fit and the same production importance-sampling estimator. ISCOS is therefore a proposal-calibration method, not a direct COS estimator of the final risk contributions.

1.1 Related Literature and Methodological Position

Gaussian factor-copula models are standard in portfolio credit risk, while Student t -copulas provide a heavy-tailed extension with tail dependence; see, for example, [5, 4, 2]. Marginal VaR and expected-shortfall contributions can be represented as conditional expected obligor losses and, under suitable conditions, provide full allocations of the portfolio quantity [6, 4].

Glasserman and Li [5] developed a two-stage rare-event simulation method that combines a change of measure for the common factors with conditional default twisting. Glasserman [4] extended this framework to marginal risk contributions and studied factor mean shifts, covariance changes, and related asymptotic approximations. Algorithms 6.1 and 6.2 use the paper-aligned Gaussian proposal $\mathcal{N}(\mu_x^{\text{GL}}, I)$, where the mean is selected by a saddle-point optimisation. This mean-only proposal provides the Glasserman benchmark in our Gaussian experiment; the Hessian covariance extension is not used in that comparison.

The cross-entropy method instead projects the formal zero-variance law onto a tractable family by minimising Kullback–Leibler divergence [7, 1]. For an exponential family, this is a sufficient-statistic moment-matching problem. Cross-entropy therefore gives a global description of the tail-conditioned factor distribution, but its usual indicator-based implementation remains vulnerable to a small number of pilot tail hits.

The COS method recovers distributional quantities from characteristic functions using Fourier–cosine coefficients [3]. Filtered discrete COS expansions, needed to control oscillations near jumps, are analysed in [8]. ISCOS uses this machinery only to approximate $q_x(u)$ during calibration. The final CVaR and CES contributions are still estimated from independent importance-sampling runs with the full factor and conditional-default likelihood ratios.

1.2 Contributions and Paper Organisation

The paper makes four contributions.

- (i) We formulate cross-entropy proposal construction in the low-dimensional common-state space. For the Gaussian component,

$$\mu_x^* = \mathbb{E}[Z \mid L \geq x], \quad \Sigma_x^* = \text{Cov}(Z \mid L \geq x).$$

For the Student t -copula, we derive a Gaussian–inverse-Gamma product proposal; the scale component matches the tail-conditional sufficient statistics $\log W$ and W^{-1} .

- (ii) We introduce ISCOS calibration based on clipped COS weights $q_{x,K}(U)$. The construction handles the left limit required by a discrete non-strict tail event and supports filtering, batching, exact grouping of homogeneous obligors, and covariance regularisation.
- (iii) We separate the statistical and numerical errors. With the exact conditional probability, the ISCOS raw-moment estimator is the Rao–Blackwellisation of its CEIS counterpart and has a no-larger covariance matrix. If

$$\sup_u |q_{x,K}(u) - q_x(u)| = O(K^{-p}),$$

then the induced proposal parameters inherit the same order, and the total fixed-threshold calibration error is

$$O_{\mathbb{P}}(M_0^{-1/2}) + O(K^{-p}).$$

- (iv) We report Gaussian and Student t -copula experiments on the same eleven-factor portfolio. For the Gaussian model, exact conditional block convolution validates the COS-calibrated moments, followed by a matched-budget comparison of the Glasserman mean-shift proposal $\mathcal{N}(\mu_x^{\text{GL}}, I)$, CEIS, and ISCOS. For the Student t -copula, CEIS- t and ISCOS- t fit the same Gaussian-inverse-Gamma family from a shared pilot. In both reported configurations, ISCOS gives higher event-weight effective sample sizes and shorter average nominal pointwise intervals than CEIS, especially for tail contributions, while the additional calibration cost remains small relative to production simulation. These numerical findings are descriptive and are not presented as a universal variance ordering.

The remainder of the paper introduces the portfolio models and risk-allocation conventions, derives the population cross-entropy proposals, presents the CEIS and ISCOS estimators, and analyses their calibration errors. The final two numerical sections treat the Gaussian and Student t -copula models before the paper concludes.

Declaration on the use of generative AI ChatGPT was used during the preparation of this manuscript to assist with language editing and to support parts of the research code. Anything generated by AI was reviewed by authors.

2 Credit-Portfolio Models and Tail-Risk Quantities

We consider a portfolio observed over a fixed time horizon. Loss-given-default amounts are deterministic, while dependence across obligors is generated by a small number of common factors. The Gaussian and Student t specifications used below differ in their common state, but both make defaults conditionally independent. This is the property that later allows us to evaluate conditional tail probabilities with the COS method.

2.1 Factor-copula loss model

For obligor n , let X_n be a latent creditworthiness variable, ξ_n its default threshold, and $l_n > 0$ its loss-at-default. We use the lower-tail convention

$$Y_n = \mathbf{1}_{\{X_n \leq \xi_n\}}, \quad L = \sum_{n=1}^N l_n Y_n, \quad (2.1)$$

so that $p_n = \mathbb{P}(X_n \leq \xi_n)$ is the unconditional default probability. The portfolio loss has finite support in $[0, L_{\max}]$, where $L_{\max} = \sum_{n=1}^N l_n$.

Let $Z \sim \mathcal{N}_d(0, I_d)$ denote the systematic factors and let $\varepsilon_1, \dots, \varepsilon_N$ be independent standard normal idiosyncratic factors, independent of Z . Obligor n has loading vector $\beta_n \in \mathbb{R}^d$, with $\|\beta_n\|_2^2 < 1$, and idiosyncratic loading

$$b_n = \sqrt{1 - \|\beta_n\|_2^2}.$$

Gaussian copula. The latent variable is

$$X_n = \beta_n^\top Z + b_n \varepsilon_n, \quad \xi_n = \Phi^{-1}(p_n),$$

where Φ is the standard normal CDF. Conditional on $Z = z$, the default probability is

$$p_n(z) = \Phi\left(\frac{\xi_n - \beta_n^\top z}{b_n}\right). \quad (2.2)$$

Student t -copula. Let $V \sim \chi_\nu^2$ be independent of Z and the idiosyncratic factors, and set

$$W = \frac{\nu}{V} \sim \text{InvGamma}\left(\frac{\nu}{2}, \frac{\nu}{2}\right).$$

We use the shape–scale parameterisation

$$f_W(w; \alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} w^{-\alpha-1} \exp\left(-\frac{\beta}{w}\right), \quad w > 0. \quad (2.3)$$

The common scale W multiplies both the systematic and idiosyncratic parts:

$$X_n = \sqrt{W} \left(\beta_n^\top Z + b_n \varepsilon_n \right). \quad (2.4)$$

Then X_n has a standard Student t_ν marginal, $\xi_n = T_\nu^{-1}(p_n)$, and

$$p_n(z, w) = \Phi\left(\frac{\xi_n / \sqrt{w} - \beta_n^\top z}{b_n}\right). \quad (2.5)$$

It is convenient to write the common state as

$$U = \begin{cases} Z, & \text{Gaussian copula,} \\ (Z, W), & \text{Student } t\text{-copula.} \end{cases} \quad (2.6)$$

In either model, we write $p_n(u) = \mathbb{P}(Y_n = 1 \mid U = u)$ for the corresponding conditional default probability.

2.2 Conditional loss distribution

Conditional on $U = u$, the indicators $Y_1, \dots, Y_N$ are independent Bernoulli variables with success probabilities $p_1(u), \dots, p_N(u)$. Hence the conditional characteristic function of the portfolio loss is

$$\varphi_{L|u}(\omega) = \mathbb{E}\left[e^{i\omega L} \mid U = u\right] = \prod_{n=1}^N \left[1 + p_n(u) \left(e^{i\omega l_n} - 1\right)\right]. \quad (2.7)$$

This closed-form expression is the main model input used by ISCOS.

For a loss threshold x , define

$$\mathcal{A}_x = \{L \geq x\}, \quad q_x(u) = \mathbb{P}(L \geq x \mid U = u). \quad (2.8)$$

Because L is discrete,

$$q_x(u) = 1 - F_{L|u}(x^-),$$

where $F_{L|u}$ is the conditional CDF. Averaging over the common state gives

$$\mathbb{P}(L \geq x) = \mathbb{E}[q_x(U)].$$

The COS approximation developed later is applied to (2.7) to evaluate $q_x(u)$.

2.3 Tail-risk quantities and allocations

Let $F_L(x) = \mathbb{P}(L \leq x)$. For $\alpha \in (0, 1)$,

$$\text{VaR}_\alpha = \inf\{x \in \mathbb{R} : F_L(x) \geq \alpha\}. \quad (2.9)$$

For a discrete loss distribution, the probability mass at VaR_α need not vanish; in particular,

$$\mathbb{P}(L > \text{VaR}_\alpha) \leq 1 - \alpha \leq \mathbb{P}(L \geq \text{VaR}_\alpha).$$

We therefore use the threshold-based convention

$$\text{ES}_\alpha = \mathbb{E}[L \mid L \geq \text{VaR}_\alpha].$$

The obligor-level contribution to VaR and the conditional expected-shortfall contribution are defined by

$$\text{CVaR}_{n,\alpha} = \mathbb{E}[l_n Y_n \mid L = \text{VaR}_\alpha], \quad (2.10)$$

$$\text{CES}_{n,\alpha} = \mathbb{E}[l_n Y_n \mid L \geq \text{VaR}_\alpha]. \quad (2.11)$$

Since $L = \sum_n l_n Y_n$, these quantities satisfy

$$\sum_{n=1}^N \text{CVaR}_{n,\alpha} = \text{VaR}_\alpha, \quad \sum_{n=1}^N \text{CES}_{n,\alpha} = \text{ES}_\alpha.$$

In smooth models these conditional allocations agree with the familiar Euler contributions. For a finite discrete portfolio, however, VaR need not be differentiable with respect to every exposure, so we take the conditional expectations above as the primary definitions.

When the exact level event is too sparse, it can be replaced by the local band

$$\mathcal{B}_{\alpha,h} = \{|L - \text{VaR}_\alpha| \leq h\}, \quad h \geq 0. \quad (2.12)$$

The corresponding local VaR contribution is obtained by conditioning on $\mathcal{B}_{\alpha,h}$ instead of $\{L = \text{VaR}_\alpha\}$. For an integer-valued portfolio with an attainable VaR, one may take $h = 0$.

The importance-sampling construction below therefore depends on the chosen copula only through the density of U and the conditional probabilities $p_n(u)$; the COS method supplies the smooth tail weight $q_x(u)$.

3 Cross-Entropy Importance Sampling in Factor Space

We now choose an importance-sampling proposal for the common state U . The idea is straightforward: factor realisations that are often followed by a large portfolio loss should be sampled more frequently. The conditional tail probability

$$q_x(u) = \mathbb{P}(L \geq x \mid U = u)$$

measures exactly how strongly a factor state u is associated with the tail event. This section uses that quantity to define the ideal, population-level proposal. Section 4 explains how the proposal is estimated in practice.

3.1 The factor distribution behind a tail loss

Fix a threshold x and write

$$\mathcal{A}_x = \{L \geq x\}, \quad \kappa_x = \mathbb{P}(\mathcal{A}_x) > 0. \quad (3.1)$$

Bayes' rule gives the density of the common state conditional on this event:

$$\pi_{U,x}^*(u) = \frac{f_U(u)q_x(u)}{\kappa_x} = f_{U|\mathcal{A}_x}(u). \quad (3.2)$$

Thus, a factor state receives more mass whenever it makes the loss event more likely. Equivalently, for any integrable statistic T ,

$$\mathbb{E}[T(U) | \mathcal{A}_x] = \frac{\mathbb{E}[T(U)q_x(U)]}{\mathbb{E}[q_x(U)]}. \quad (3.3)$$

This identity is the basis of all proposal parameters derived below.

The density in (3.2) is the factor marginal of the usual zero-variance distribution for estimating κ_x . It is also a zero-variance proposal for the conditional Monte Carlo representation $\kappa_x = \mathbb{E}[q_x(U)]$. It is not directly available, however, because both its normalising constant and the function q_x are unknown in closed form. We therefore approximate it within a tractable family. The same construction can be used for an exact-level event $\{L = x\}$ or a local band by replacing q_x with the corresponding conditional probability.

The word “optimal” should be read in this specific sense. The target in (3.2) is natural for conditional Monte Carlo, and its cross-entropy projection is optimal within the chosen proposal family. This does not imply that the fitted proposal minimises the variance of every downstream risk estimator.

3.2 Cross-entropy as a weighted fit

Let $\mathcal{G} = \{g_\theta : \theta \in \Theta\}$ be a parametric family of factor densities. The cross-entropy parameter is defined by

$$\theta_x^* \in \arg \min_{\theta \in \Theta} D_{\text{KL}}(\pi_{U,x}^* \| g_\theta). \quad (3.4)$$

Terms that do not depend on θ can be removed from the objective, so (3.4) is equivalent to

$$\theta_x^* \in \arg \max_{\theta \in \Theta} \mathbb{E}[q_x(U) \log g_\theta(U)]. \quad (3.5)$$

The factor κ_x^{-1} has been omitted because it does not affect the maximiser. In other words, population cross-entropy calibration is simply a weighted likelihood fit: factor states are weighted by their conditional probability of producing a tail loss.

For an exponential family

$$g_\theta(u) = c(u) \exp\{\theta^\top T(u) - A(\theta)\},$$

the first-order condition for an interior optimum is

$$\mathbb{E}_{g_{\theta_x^*}}[T(U)] = \mathbb{E}[T(U) | \mathcal{A}_x]. \quad (3.6)$$

Hence the cross-entropy projection matches the tail-conditional sufficient statistics. The Gaussian and inverse-Gamma proposals below are both examples of this rule; see [7, 1] for the general cross-entropy method.

3.3 Gaussian proposal for the systematic factors

Under both copula specifications, the original systematic factor is $Z \sim \mathcal{N}_d(0, I_d)$. We approximate its tail-conditioned marginal by $\mathcal{N}_d(\mu, \Sigma)$. Since the Gaussian sufficient statistics are Z and $ZZ^\top$, moment matching gives

$$\mu_x^* = \mathbb{E}[Z \mid \mathcal{A}_x] = \frac{\mathbb{E}[Z q_x(U)]}{\mathbb{E}[q_x(U)]}, \quad (3.7)$$

$$\Sigma_x^* = \text{Cov}(Z \mid \mathcal{A}_x) = \frac{\mathbb{E}[(Z - \mu_x^*)(Z - \mu_x^*)^\top q_x(U)]}{\mathbb{E}[q_x(U)]}. \quad (3.8)$$

Thus the Gaussian proposal is fitted to the global location and spread of the factor states associated with a large loss. For the Gaussian copula, $U = Z$; for the Student t -copula, the expectations are over $U = (Z, W)$.

If only a mean shift is allowed, the optimal mean is still μ_x^* and the covariance is fixed at I_d . Within a diagonal Gaussian family, the optimal variances are the diagonal entries of Σ_x^* . These restricted proposals can be useful when the pilot sample is too small to fit a stable full covariance. Finite-sample regularisation is discussed in Section 4.3.

3.4 Student t -copula product proposal

For the Student t -copula, the common state is (Z, W) . Although Z and W are independent under the original model, conditioning on a large loss generally makes them dependent: a tail loss may be caused by an adverse factor direction, a large common scale, or both. For tractability, we use the product family

$$g_{Z,W}(z, w) = \phi_d(z; \mu, \Sigma) f_W(w; \alpha, \beta), \quad (3.9)$$

where f_W is the inverse-Gamma density in (2.3). Because the log-density in (3.9) is additive, the cross-entropy objective separates into a Gaussian fit for Z and an inverse-Gamma fit for W . The Gaussian parameters are therefore still given by (3.7)–(3.8).

For the scale component, define the two tail-conditional sufficient-statistic moments

$$m_{\log, x} = \mathbb{E}[\log W \mid \mathcal{A}_x] = \frac{\mathbb{E}[(\log W) q_x(Z, W)]}{\mathbb{E}[q_x(Z, W)]}, \quad (3.10)$$

$$m_{-1, x} = \mathbb{E}[W^{-1} \mid \mathcal{A}_x] = \frac{\mathbb{E}[W^{-1} q_x(Z, W)]}{\mathbb{E}[q_x(Z, W)]}. \quad (3.11)$$

Theorem 3.1 (Inverse-Gamma cross-entropy projection). *Assume that $\mathbb{E}[|\log W| + W^{-1} \mid \mathcal{A}_x] < \infty$ and that $W \mid \mathcal{A}_x$ is non-degenerate. The inverse-Gamma component of the product proposal is uniquely determined by*

$$\beta_x^* = \frac{\alpha_x^*}{m_{-1, x}}, \quad \psi(\alpha_x^*) - \log \alpha_x^* + m_{\log, x} + \log m_{-1, x} = 0, \quad (3.12)$$

where ψ is the digamma function.

Proof. For $\alpha, \beta > 0$, the inverse-Gamma log-density is

$$\log g_{\alpha, \beta}(w) = \alpha \log \beta - \log \Gamma(\alpha) - (\alpha + 1) \log w - \frac{\beta}{w}.$$

Hence the tail-conditional cross-entropy objective is

$$J_x(\alpha, \beta) = \alpha \log \beta - \log \Gamma(\alpha) - (\alpha + 1)m_{\log, x} - \beta m_{-1, x}.$$

The assumed integrability ensures that this quantity is finite, and $m_{-1, x} > 0$.

For fixed $\alpha > 0$,

$$\frac{\partial J_x}{\partial \beta} = \frac{\alpha}{\beta} - m_{-1, x}, \quad \frac{\partial^2 J_x}{\partial \beta^2} = -\frac{\alpha}{\beta^2} < 0.$$

Therefore the unique maximiser in β is

$$\beta(\alpha) = \frac{\alpha}{m_{-1, x}}.$$

Substitution gives the profiled objective

$$\bar{J}_x(\alpha) = \alpha \log \alpha - \alpha \log m_{-1, x} - \log \Gamma(\alpha) - (\alpha + 1)m_{\log, x} - \alpha.$$

Let

$$c_x = m_{\log, x} + \log m_{-1, x}.$$

Then

$$\bar{J}'_x(\alpha) = \log \alpha - \psi(\alpha) - c_x,$$

and

$$\bar{J}''_x(\alpha) = \frac{1}{\alpha} - \psi_1(\alpha) < 0,$$

where ψ_1 is the trigamma function. Indeed,

$$\psi_1(\alpha) = \sum_{k=0}^{\infty} \frac{1}{(\alpha + k)^2} > \int_0^{\infty} \frac{dt}{(\alpha + t)^2} = \frac{1}{\alpha}.$$

Thus $\bar{J}_x$ is strictly concave.

Moreover, strict Jensen inequality gives

$$\begin{aligned} c_x &= \log \mathbb{E}[W^{-1} \mid \mathcal{A}_x] - \mathbb{E}[\log(W^{-1}) \mid \mathcal{A}_x] \\ &> 0, \end{aligned}$$

because $W \mid \mathcal{A}_x$ is non-degenerate. Define

$$h(\alpha) = \log \alpha - \psi(\alpha).$$

The preceding trigamma inequality shows that h is strictly decreasing, while the standard digamma asymptotics give

$$\lim_{\alpha \downarrow 0} h(\alpha) = +\infty, \quad \lim_{\alpha \uparrow \infty} h(\alpha) = 0.$$

Since $c_x > 0$, there is therefore a unique $\alpha_x^* > 0$ satisfying

$$h(\alpha_x^*) = c_x,$$

or equivalently,

$$\psi(\alpha_x^*) - \log \alpha_x^* + m_{\log, x} + \log m_{-1, x} = 0.$$

Because $\bar{J}_x$ is strictly concave, this root is its unique global maximiser. The corresponding scale parameter is uniquely given by

$$\beta_x^* = \frac{\alpha_x^*}{m_{-1, x}}.$$

□

The inverse-Gamma fit therefore matches $\log W$ and W^{-1} , not the ordinary mean of W . The product form in (3.9) is a deliberate approximation: it does not represent the dependence between Z and W induced by the tail event. A richer conditional proposal, such as a Gaussian law for $Z \mid W$, is possible but is outside the first implementation considered here.

3.5 Pilot reweighting and likelihood ratios

The identities above are written under the original common-state density f_U , but the pilot sample may come from another density h . Assume that $h(u) > 0$ wherever $f_U(u)q_x(u) > 0$, and define

$$R_h(u) = \frac{f_U(u)}{h(u)}. \quad (3.13)$$

Then

$$\mathbb{E}[T(U) \mid \mathcal{A}_x] = \frac{\mathbb{E}_h[T(U)R_h(U)q_x(U)]}{\mathbb{E}_h[R_h(U)q_x(U)]}. \quad (3.14)$$

When $h = f_U$, the weight is simply $q_x(U)$. Formula (3.14) also permits iterative cross-entropy schemes in which the pilot distribution is updated between stages.

Once a proposal has been fitted, production estimators must be corrected back to the original measure. If the original Gaussian law is $\mathcal{N}_d(0, I_d)$ and the proposal is $\mathcal{N}_d(\mu, \Sigma)$, the factor likelihood ratio is

$$R_Z(z; \mu, \Sigma) = |\Sigma|^{1/2} \exp \left\{ \frac{1}{2}(z - \mu)^\top \Sigma^{-1}(z - \mu) - \frac{1}{2}z^\top z \right\}. \quad (3.15)$$

For a mean shift, $\Sigma = I_d$, this reduces to

$$R_Z(z; \mu, I_d) = \exp \left(-\mu^\top z + \frac{1}{2}\mu^\top \mu \right).$$

If the original scale law is $\text{InvGamma}(\alpha_0, \beta_0)$ and the proposal uses $\text{InvGamma}(\alpha, \beta)$, then

$$R_W(w; \alpha, \beta) = \frac{\beta_0^{\alpha_0} \Gamma(\alpha)}{\Gamma(\alpha_0) \beta^\alpha} w^{\alpha - \alpha_0} \exp \left(\frac{\beta - \beta_0}{w} \right). \quad (3.16)$$

Under the product proposal, $R_U(z, w) = R_Z(z)R_W(w)$. Any change to the conditional Bernoulli default probabilities contributes a separate likelihood ratio in the production stage. All ratios are evaluated on the logarithmic scale in the implementation.

4 Estimating the Cross-Entropy Proposal

Section 3 shows that the cross-entropy proposal is obtained from moments of the common state under the tail event $\mathcal{A}_x = \{L \geq x\}$. In practice, proposal calibration therefore comes down to one question: how much weight should each pilot factor draw receive? CEIS uses the outcome of one simulated portfolio loss, whereas ISCOS uses an approximation of the conditional probability of a tail loss. Once the weights have been computed, both methods fit the proposal in exactly the same way.

Throughout this section, the threshold x is fixed. In the numerical experiments it is an attainable estimate of VaR_α obtained from a preliminary Monte Carlo run. The proposal is calibrated to the non-strict event $\{L \geq x\}$. The same threshold is then used for the exact-level and tail estimators in the production stage.

Let h be the pilot density of the common state and write $R_h(u) = f_U(u)/h(u)$. The first implementation takes $h = f_U$, so that $R_h \equiv 1$; the more general notation also covers an updated pilot proposal.

4.1 One pilot sample, two weights

Draw $U^{(1)}, \dots, U^{(M_0)}$ independently from h . For CEIS, also generate a conditional default vector and its loss $L^{(m)} = \sum_{n=1}^N l_n Y_n^{(m)}$. ISCOS does not need this additional default draw for calibration. The two weights are

$$\omega_m = R_h(U^{(m)}) \begin{cases} \mathbf{1}_{\{L^{(m)} \geq x\}}, & \text{CEIS,} \\ q_{x,K}(U^{(m)}), & \text{ISCOS.} \end{cases} \quad (4.1)$$

For any nonnegative weights $\omega_1, \dots, \omega_{M_0}$, let $S_\omega = \sum_m \omega_m$. Provided that $S_\omega > 0$, the Gaussian proposal is fitted by the weighted mean and covariance

$$\hat{\mu}_\omega = \frac{1}{S_\omega} \sum_{m=1}^{M_0} \omega_m Z^{(m)}, \quad (4.2)$$

$$\hat{\Sigma}_{\omega, \text{raw}} = \frac{1}{S_\omega} \sum_{m=1}^{M_0} \omega_m (Z^{(m)} - \hat{\mu}_\omega)(Z^{(m)} - \hat{\mu}_\omega)^\top. \quad (4.3)$$

Substituting the first or second line of (4.1) gives the CEIS or ISCOS estimate, respectively. The denominator is the total weight, rather than an unbiased-sample correction, because the target is the population covariance in Section 3.3.

For the Student t -copula, the same weights are also used to estimate

$$\widehat{m}_{\log, \omega} = \frac{\sum_m \omega_m \log W^{(m)}}{S_\omega}, \quad \widehat{m}_{-1, \omega} = \frac{\sum_m \omega_m (W^{(m)})^{-1}}{S_\omega}.$$

Inserting these two moments into (3.12) gives the fitted inverse-Gamma parameters.

When $h = f_U$, CEIS assigns weight one only to simulated tail hits. At an extreme threshold, there may be too few such observations to estimate a full covariance matrix reliably. ISCOS instead assigns a probability between zero and one to each factor state. States that are stressful but happen not to produce a tail loss in one conditional simulation

can therefore still contribute to the fit. Section 5 makes this conditional-Monte-Carlo comparison precise.

4.2 Computing the smooth COS weights

ISCOS requires $q_x(u) = \mathbb{P}(L \geq x \mid U = u)$. Conditional on $U = u$, the loss characteristic function is available from (2.7). Choose an interval $[a, b]$ that contains the loss support in its interior, and let K be the number of COS modes, including the zero mode. With $\omega_k = k\pi/(b-a)$ and filter coefficients $\sigma_{k,K} = \sigma(k/K)$, the filtered conditional CDF is approximated by

$$F_{L|u,K}^\sigma(y) = \frac{y-a}{b-a} + \frac{2}{\pi} \sum_{k=1}^{K-1} \frac{\sigma_{k,K}}{k} \operatorname{Re} \left\{ \varphi_{L|u}(\omega_k) e^{-i\omega_k a} \right\} \sin \left(k\pi \frac{y-a}{b-a} \right). \quad (4.4)$$

Because the event is $L \geq x$, the relevant CDF value is the left limit $F_{L|u}(x^-)$. We therefore evaluate the expansion at a point $y_x < x$ and set

$$\tilde{q}_{x,K}(u) = 1 - F_{L|u,K}^\sigma(y_x), \quad (4.5)$$

$$q_{x,K}(u) = \Pi_{[0,1]}(\tilde{q}_{x,K}(u)). \quad (4.6)$$

The left-limit convention matters for a discrete loss. If all losses lie on a lattice with step Δ and x is attainable, we use

$$a = -\frac{\Delta}{2}, \quad b = L_{\max} + \frac{\Delta}{2}, \quad y_x = x - \frac{\Delta}{2}. \quad (4.7)$$

Then $F_{L|u}(y_x) = \mathbb{P}(L < x \mid U = u)$, so the COS expansion is not evaluated at a jump of the CDF. For a non-lattice portfolio, y_x can be chosen just below x , with the convention reported alongside the numerical results.

Clipping removes small Fourier over- and undershoots. Since the true probability lies in $[0, 1]$, it cannot make the pointwise error larger:

$$|q_{x,K}(u) - q_x(u)| \leq |\tilde{q}_{x,K}(u) - q_x(u)|. \quad (4.8)$$

The number of modes controls the trade-off between numerical cost and the accuracy of the smooth weight. Section 5 studies how the finite-COS error propagates to the fitted proposal parameters.

4.3 Practical calibration

A preliminary simulation is first used to choose an attainable threshold $\hat{x}_\alpha = \inf\{x : \hat{F}_{M_0}(x) \geq \alpha\}$. The pilot factor draws can then be reused for CEIS and ISCOS, but the final importance-sampling run should be independent of this calibration sample.

A weighted covariance can be nearly singular when only a small effective sample contributes. We first symmetrise the raw estimate and, in the baseline implementation, add a small ridge:

$$\hat{\Sigma}_{x,\text{sym}} = \frac{1}{2} \left(\hat{\Sigma}_{x,\text{raw}} + \hat{\Sigma}_{x,\text{raw}}^\top \right), \quad (4.9)$$

$$\hat{\Sigma}_{x,\rho} = \hat{\Sigma}_{x,\text{sym}} + \rho I_d, \quad \rho > 0. \quad (4.10)$$

If more regularisation is needed, one may shrink towards a spherical covariance,

$$\hat{\Sigma}_{x,\lambda} = (1 - \lambda) \hat{\Sigma}_{x,\text{sym}} + \lambda \tau I_d, \quad \tau = \frac{\text{tr}(\hat{\Sigma}_{x,\text{sym}})}{d}, \quad (4.11)$$

or floor very small eigenvalues. These are finite-sample safeguards rather than part of the population cross-entropy projection.

A useful summary of the calibration weights is their self-normalised effective sample size,

$$M_{\text{eff}} = \frac{\left(\sum_{m=1}^{M_0} \omega_m \right)^2}{\sum_{m=1}^{M_0} \omega_m^2}. \quad (4.12)$$

If M_{eff} is comparable to, or smaller than, the factor dimension, a full covariance fit should be treated cautiously. For ISCOS we also record the range of the raw COS probabilities, the amount of clipping, and the average smooth weight. Conditional characteristic functions are evaluated in batches; identical obligors can be grouped, reducing the direct $\mathcal{O}(M_0 NK)$ cost when the portfolio has repeated exposures and loadings.

The calibration procedure can be summarised as follows.

1. Estimate an attainable tail threshold and retain the pilot common states.
2. Compute either the CEIS indicators or the ISCOS conditional-tail probabilities.
3. Fit the weighted Gaussian moments and, for the Student t -copula, the two inverse-Gamma moments.
4. Regularise the covariance when needed and check the weight diagnostics.
5. Use the fitted proposal in an independent production importance-sampling run.

The two methods therefore differ at only one point: CEIS uses one binary loss outcome per pilot state, whereas ISCOS uses a smooth estimate of the chance that the same state produces a tail loss. The next section studies the statistical gain from this replacement and the approximation error introduced by the finite COS expansion.

5 Error Analysis for ISCOS Calibration

The cross-entropy proposal is determined by moments of the common state under $\mathcal{A}_x = \{L \geq x\}$. In practice, these moments are not observed directly. CEIS estimates them with the binary indicator $I_x = \mathbf{1}_{\mathcal{A}_x}$, whereas ISCOS replaces this indicator by a finite COS approximation $q_{x,K}(U)$ of the conditional tail probability

$$q_x(U) = \mathbb{P}(\mathcal{A}_x \mid U).$$

These two substitutions affect the calibration in different ways. Replacing I_x by the exact conditional probability $q_x(U)$ removes the idiosyncratic default noise that remains after

the common state has been drawn. Replacing q_x by $q_{x,K}$, on the other hand, introduces a deterministic numerical error.

It is useful to keep these effects separate. Let ϑ_x^* denote the exact population cross-entropy parameters, let $\vartheta_{x,K}^*$ denote the population parameters obtained with the finite COS weight $q_{x,K}$, and let $\hat{\vartheta}_{x,K,M_0}$ be the corresponding estimator from a pilot sample of size M_0 . Then

$$\hat{\vartheta}_{x,K,M_0} - \vartheta_x^* = \underbrace{(\hat{\vartheta}_{x,K,M_0} - \vartheta_{x,K}^*)}_{\text{pilot Monte Carlo error}} + \underbrace{(\vartheta_{x,K}^* - \vartheta_x^*)}_{\text{finite-COS error}}. \quad (5.1)$$

The first term is statistical and decreases with M_0 ; the second is deterministic and decreases with the number of COS modes K . Before studying this decomposition, we compare CEIS with the ideal estimator that uses the exact conditional probability.

Throughout this section, the loss threshold x is fixed. Thus, all results refer to the calibration of a proposal for a prescribed event $\mathcal{A}_x$. The additional effect of replacing x by a preliminary empirical quantile is discussed in Section 5.5.

5.1 Exact conditional weighting and Rao–Blackwellisation

The proposal parameters are ratios of unnormalised tail moments. It is therefore convenient to compare the underlying raw moments first, before the normalisation is applied. Let $T(U) \in \mathbb{R}^r$ collect the common-state statistics required by the chosen proposal family. For the Gaussian component, one may take

$$T_G(U) = \left(1, Z^\top, \text{vech}(ZZ^\top)^\top\right)^\top. \quad (5.2)$$

The first coordinate estimates the tail probability, the next d coordinates estimate the numerator of the tail-conditional mean, and the remaining coordinates estimate the numerator of the second moment. For the Student t -copula proposal, the inverse-Gamma sufficient statistics are appended:

$$T_t(U) = \left(T_G(U)^\top, \log W, W^{-1}\right)^\top. \quad (5.3)$$

Let h be the pilot density and $R_h(u) = f_U(u)/h(u)$ the corresponding likelihood ratio. Define the one-observation raw-moment vectors

$$G_x^{\text{CEIS}} := R_h(U)I_x T(U), \quad G_x^{\text{RB}} := R_h(U)q_x(U)T(U). \quad (5.4)$$

Here RB denotes the ideal Rao–Blackwellised estimator. Conditional on U , the factor $R_h(U)T(U)$ is fixed, and therefore

$$\mathbb{E}[G_x^{\text{CEIS}} \mid U] = R_h(U)T(U)\mathbb{E}[I_x \mid U] = G_x^{\text{RB}}. \quad (5.5)$$

Consequently, the two vectors estimate exactly the same population moments:

$$\mathbb{E}_h[G_x^{\text{CEIS}}] = \mathbb{E}_h[G_x^{\text{RB}}] = \mathbb{E}_{f_U}[q_x(U)T(U)]. \quad (5.6)$$

The difference lies only in their sampling covariance.

Theorem 5.1 (Exact Rao–Blackwell covariance reduction). *Assume that*

$$\mathbb{E}_h[R_h(U)^2 \|T(U)\|_2^2] < \infty.$$

Then

$$\begin{aligned} \text{Cov}_h(G_x^{\text{CEIS}}) - \text{Cov}_h(G_x^{\text{RB}}) \\ = \mathbb{E}_h[R_h(U)^2 q_x(U)(1 - q_x(U))T(U)T(U)^\top] \succeq 0. \end{aligned} \quad (5.7)$$

Equivalently, for every $a \in \mathbb{R}^r$,

$$\text{Var}_h(a^\top G_x^{\text{RB}}) \leq \text{Var}_h(a^\top G_x^{\text{CEIS}}). \quad (5.8)$$

Proof. Conditional on U , the only remaining randomness in G_x^{CEIS} is the Bernoulli variable I_x . Since

$$\mathbb{E}[I_x | U] = q_x(U), \quad \text{Var}(I_x | U) = q_x(U)(1 - q_x(U)),$$

we have (5.5) and

$$\text{Cov}(G_x^{\text{CEIS}} | U) = R_h(U)^2 q_x(U)(1 - q_x(U))T(U)T(U)^\top.$$

The law of total covariance gives

$$\begin{aligned} \text{Cov}_h(G_x^{\text{CEIS}}) &= \text{Cov}_h(\mathbb{E}[G_x^{\text{CEIS}} | U]) + \mathbb{E}_h[\text{Cov}(G_x^{\text{CEIS}} | U)] \\ &= \text{Cov}_h(G_x^{\text{RB}}) + \mathbb{E}_h[R_h(U)^2 q_x(U)(1 - q_x(U))T(U)T(U)^\top], \end{aligned}$$

which proves (5.7). The last matrix is positive semidefinite because it is an expectation of positive-semidefinite rank-one matrices. Equation (5.8) follows by pre- and post-multiplying by $a^\top$ and a . $\square$

The result has a direct componentwise interpretation. When the original pilot law is used, so that $R_h \equiv 1$, every scalar statistic $T_\ell(U)$ satisfies

$$\text{Var}(T_\ell(U)I_x) - \text{Var}(T_\ell(U)q_x(U)) = \mathbb{E}[T_\ell(U)^2 q_x(U)(1 - q_x(U))]. \quad (5.9)$$

Thus, the denominator, every first moment, and every second moment used in the Gaussian fit all benefit from the same conditional-variance removal. The gain is strict in a direction a whenever $a^\top R_h(U)T(U) \neq 0$ with positive probability on a region where $0 < q_x(U) < 1$. If $q_x(U) \in \{0, 1\}$ almost surely, the common state already determines whether the event occurs, so there is no conditional default noise to remove.

The exact order in Theorem 5.1 concerns the unnormalised moments. The fitted CE parameters are nonlinear functions of these moments. To make this connection explicit, write the Gaussian raw moment vector as

$$m_x = (\kappa_x, b_x, \text{vech}(C_x)), \quad b_x = \mathbb{E}_h[R_h q_x Z], \quad C_x = \mathbb{E}_h[R_h q_x Z Z^\top], \quad (5.10)$$

where $\kappa_x = \mathbb{E}_h[R_h q_x] > 0$. The parameter map $\mathcal{G}$ is

$$\mu_x^* = \frac{b_x}{\kappa_x}, \quad \Sigma_x^* = \frac{C_x}{\kappa_x} - \mu_x^* (\mu_x^*)^\top. \quad (5.11)$$

For perturbations $(\dot{\kappa}, \dot{b}, \dot{C})$, its differential is

$$D\mu_x^* = \frac{\dot{b} - \mu_x^* \dot{\kappa}}{\kappa_x}, \quad (5.12)$$

$$D\Sigma_x^* = \frac{\dot{C}}{\kappa_x} - \frac{C_x \dot{\kappa}}{\kappa_x^2} - (D\mu_x^*)(\mu_x^*)^\top - \mu_x^*(D\mu_x^*)^\top. \quad (5.13)$$

Hence the map is continuously differentiable whenever $\kappa_x > 0$.

Corollary 5.2 (Asymptotic covariance of the fitted proposal). *Assume the conditions of Theorem 5.1, let $\kappa_x > 0$, and suppose that the parameter map $\mathcal{G}$ is continuously differentiable at the common raw-moment vector m_x . Define*

$$\widehat{m}_x^j = \frac{1}{M_0} \sum_{m=1}^{M_0} G_{x,m}^j, \quad j \in \{\text{CEIS}, \text{RB}\}. \quad (5.14)$$

Then

$$\sqrt{M_0} (\mathcal{G}(\widehat{m}_x^j) - \mathcal{G}(m_x)) \xrightarrow{d} \mathcal{N}(0, \mathcal{V}_j), \quad (5.15)$$

where

$$\mathcal{V}_j = D\mathcal{G}(m_x) \text{Cov}_h(G_x^j) D\mathcal{G}(m_x)^\top. \quad (5.16)$$

Moreover,

$$\mathcal{V}_{\text{RB}} \preceq \mathcal{V}_{\text{CEIS}}. \quad (5.17)$$

Proof. The multivariate central limit theorem applies to the raw-moment averages in (5.14). The delta method then yields (5.15). Finally,

$$\mathcal{V}_{\text{CEIS}} - \mathcal{V}_{\text{RB}} = D\mathcal{G}(m_x) [\text{Cov}_h(G_x^{\text{CEIS}}) - \text{Cov}_h(G_x^{\text{RB}})] D\mathcal{G}(m_x)^\top,$$

which is positive semidefinite by Theorem 5.1. $\square$

For the Student t -copula, the same conclusion applies to the Gaussian parameters and to the inverse-Gamma parameters. The latter are smooth functions of the two sufficient-statistic moments $m_{\log, x}$ and $m_{-1, x}$ because the derivative of the inverse-Gamma shape equation is non-zero at the population solution; this point is made explicit in Corollary 5.6 below.

Corollary 5.2 is a first-order asymptotic statement for the normalised proposal parameters. It does not give an exact finite-sample variance order for the ratio estimators $\widehat{\mu}_x$ and $\widehat{\Sigma}_x$. It also does not yet describe practical ISCOS, because practical ISCOS uses $q_{x,K}$, not the exact conditional probability q_x .

5.2 A uniform COS error for the conditional tail probability

We next study the deterministic difference between $q_{x,K}$ and q_x . The required bound must hold uniformly over the common state: the subsequent CE moments integrate over all possible values of U , so a pointwise-in- u convergence statement would not by itself control the moment ratios. The finite support of the conditional portfolio loss makes the desired uniformity available.

Let

$$\mathcal{S}_L = \{s_1, \dots, s_J\} \subset (a, b) \quad (5.18)$$

be the distinct attainable losses. This set depends only on the deterministic losses-at-default and not on the common state. Conditional on $U = u$, write

$$\pi_j(u) = \mathbb{P}(L = s_j \mid U = u), \quad \pi_j(u) \geq 0, \quad \sum_{j=1}^J \pi_j(u) = 1. \quad (5.19)$$

The conditional CDF is therefore the convex combination

$$F_{L|u}(y) = \sum_{j=1}^J \pi_j(u) \mathbf{1}_{\{s_j \leq y\}}. \quad (5.20)$$

Let $y_x < x$ be the continuity point used in Section 4 to represent the left limit, so that $q_x(u) = 1 - F_{L|u}(y_x)$. For a unit point mass at s_j , denote by $H_{j,K}^\sigma(y_x)$ the filtered COS approximation to its CDF at y_x . The discrete-COS results in [8] give, for the chosen filter, interval, and evaluation point, an atomwise estimate of the form

$$\left| H_{j,K}^\sigma(y_x) - \mathbf{1}_{\{s_j \leq y_x\}} \right| \leq C_{j,x,\sigma} K^{-p}, \quad j = 1, \dots, J, \quad (5.21)$$

where $p > 0$ is the proved convergence order of the filtered CDF approximation.

Theorem 5.3 (Uniform conditional COS error). *Suppose that $\mathcal{S}_L$ is finite, lies strictly inside (a, b) , and $y_x \notin \mathcal{S}_L$. If (5.21) holds, then there is a finite constant $C_{x,\sigma}$, independent of u , such that*

$$\sup_u |\tilde{q}_{x,K}(u) - q_x(u)| \leq C_{x,\sigma} K^{-p}. \quad (5.22)$$

The clipped COS probability in (4.6) satisfies the same order:

$$\epsilon_{x,K} := \sup_u |q_{x,K}(u) - q_x(u)| = O(K^{-p}). \quad (5.23)$$

Proof. The filtered COS reconstruction is linear in the characteristic function and therefore linear in the atom probabilities. Hence

$$F_{L|u,K}^\sigma(y_x) = \sum_{j=1}^J \pi_j(u) H_{j,K}^\sigma(y_x). \quad (5.24)$$

Combining (5.20) and (5.24) gives

$$\begin{aligned} \left| F_{L|u,K}^\sigma(y_x) - F_{L|u}(y_x) \right| &\leq \sum_{j=1}^J \pi_j(u) \left| H_{j,K}^\sigma(y_x) - \mathbf{1}_{\{s_j \leq y_x\}} \right| \\ &\leq \left(\max_{1 \leq j \leq J} C_{j,x,\sigma} \right) K^{-p} \sum_{j=1}^J \pi_j(u) \\ &= C_{x,\sigma} K^{-p}. \end{aligned}$$

The constant is independent of u because the conditional atom probabilities form a convex combination. Since $\tilde{q}_{x,K} = 1 - F_{L|u,K}^\sigma(y_x)$, the same bound holds for the raw tail approximation. Finally, projection onto $[0, 1]$ is non-expansive relative to any point in $[0, 1]$, so (4.8) gives the same bound for $q_{x,K}$. $\square$

The assumption $y_x \notin \mathcal{S}_L$ is essential for approximating the non-strict event $\{L \geq x\}$. At a jump of a discrete CDF, a symmetric Fourier reconstruction naturally approaches a midpoint value rather than the left limit required by $1 - F_{L|u}(x^-)$. For a lattice portfolio, the half-step convention in (4.7) places y_x strictly between two attainable losses and therefore avoids this ambiguity.

Two features of Theorem 5.3 are worth emphasising. First, the common state changes the probabilities $\pi_j(u)$, but not the support points s_j ; this is why the bound is uniform in both the Gaussian and Student t -copula models. Second, the symbol p denotes the convergence order actually established for the selected filtered CDF approximation. It should not be identified mechanically with a parameter in the name of a filter. For example, if the available theorem for a formal order- r filter gives only $O(K^{1-r})$, then the subsequent results should be read with $p = r - 1$. A sharper value of p should be used only when it is proved for the selected filter and parameterisation. Filter-specific endpoint terms and the position of y_x relative to the support and the COS interval affect the constant and, in some cases, the available order.

5.3 Propagation of the COS error to the CE parameters

The finite-COS proposal is defined through normalised weighted moments, so the next step is to propagate the uniform error in (5.23) through these ratios. We first give a simple identity that will be used repeatedly.

Lemma 5.4 (Perturbation identity for a weighted moment). *Let q and q_K be integrable weights with $\kappa = \mathbb{E}[q(U)] > 0$ and $\kappa_K = \mathbb{E}[q_K(U)] > 0$. For an integrable vector- or matrix-valued function g , define*

$$\bar{g} = \frac{\mathbb{E}[g(U)q(U)]}{\kappa}, \quad \bar{g}_K = \frac{\mathbb{E}[g(U)q_K(U)]}{\kappa_K}.$$

With $e_K = q_K - q$,

$$\bar{g}_K - \bar{g} = \frac{\mathbb{E}[(g(U) - \bar{g})e_K(U)]}{\kappa_K}. \quad (5.25)$$

Consequently, if $\|e_K\|_\infty \leq \epsilon_K$, then

$$\|\bar{g}_K - \bar{g}\| \leq \frac{\epsilon_K}{\kappa_K} \mathbb{E}[\|g(U) - \bar{g}\|]. \quad (5.26)$$

Proof. Using $q_K = q + e_K$ and $\mathbb{E}[gq] = \kappa\bar{g}$,

$$\begin{aligned} \kappa_K(\bar{g}_K - \bar{g}) &= \mathbb{E}[gq_K] - \kappa_K\bar{g} \\ &= \mathbb{E}[gq] + \mathbb{E}[ge_K] - (\kappa + \mathbb{E}[e_K])\bar{g} \\ &= \mathbb{E}[(g - \bar{g})e_K]. \end{aligned}$$

The norm bound follows immediately. □

For later use, define the finite- K normaliser and Gaussian cross-entropy parameters by

$$\kappa_{x,K} := \mathbb{E}_h[R_h(U)q_{x,K}(U)] = \mathbb{E}_{f_U}[q_{x,K}(U)], \quad (5.27)$$

$$\mu_{x,K}^* := \frac{\mathbb{E}_h[Z R_h(U)q_{x,K}(U)]}{\kappa_{x,K}}, \quad (5.28)$$

$$\Sigma_{x,K}^* := \frac{\mathbb{E}_h\left[\left(Z - \mu_{x,K}^*\right)\left(Z - \mu_{x,K}^*\right)^\top R_h(U)q_{x,K}(U)\right]}{\kappa_{x,K}}. \quad (5.29)$$

Their exact counterparts are obtained by replacing $q_{x,K}$ with q_x .

Theorem 5.5 (Propagation to the Gaussian CE parameters). *Assume that $\kappa_x = \mathbb{P}(L \geq x) > 0$, $\mathbb{E}[\|Z\|_2^2] < \infty$, and (5.23) holds. Then*

$$|\kappa_{x,K} - \kappa_x| \leq \epsilon_{x,K}. \quad (5.30)$$

For all sufficiently large K , so that $\epsilon_{x,K} \leq \kappa_x/2$,

$$\|\mu_{x,K}^* - \mu_x^*\|_2 \leq \frac{2\epsilon_{x,K}}{\kappa_x} \mathbb{E}[\|Z - \mu_x^*\|_2], \quad (5.31)$$

and

$$\begin{aligned} \|\Sigma_{x,K}^* - \Sigma_x^*\|_F &\leq \frac{2\epsilon_{x,K}}{\kappa_x} \mathbb{E}\left[\|(Z - \mu_x^*)(Z - \mu_x^*)^\top - \Sigma_x^*\|_F\right] \\ &\quad + \left(\frac{2\epsilon_{x,K}}{\kappa_x} \mathbb{E}[\|Z - \mu_x^*\|_2]\right)^2. \end{aligned} \quad (5.32)$$

In particular,

$$|\kappa_{x,K} - \kappa_x| = O(K^{-p}), \quad \|\mu_{x,K}^* - \mu_x^*\|_2 = O(K^{-p}), \quad \|\Sigma_{x,K}^* - \Sigma_x^*\|_F = O(K^{-p}). \quad (5.33)$$

Proof. Let $e_{x,K} = q_{x,K} - q_x$. The normalising constants satisfy

$$|\kappa_{x,K} - \kappa_x| = |\mathbb{E}[e_{x,K}(U)]| \leq \epsilon_{x,K},$$

which proves (5.30). Hence $\kappa_{x,K} \geq \kappa_x/2$ for sufficiently large K .

Apply Lemma 5.4 with $g(U) = Z$, $q = q_x$, and $q_K = q_{x,K}$. Since the exact weighted mean is μ_x^* ,

$$\mu_{x,K}^* - \mu_x^* = \frac{\mathbb{E}[(Z - \mu_x^*)e_{x,K}(U)]}{\kappa_{x,K}}. \quad (5.34)$$

Taking norms, using the uniform error bound, and replacing $1/\kappa_{x,K}$ by $2/\kappa_x$ gives (5.31).

For the covariance, set

$$A_x(Z) = (Z - \mu_x^*)(Z - \mu_x^*)^\top, \quad \delta_{x,K} = \mu_{x,K}^* - \mu_x^*.$$

Because the $q_{x,K}$ -weighted mean of $Z - \mu_x^*$ is $\delta_{x,K}$, expanding around μ_x^* yields

$$\Sigma_{x,K}^* = \frac{\mathbb{E}[A_x(Z)q_{x,K}(U)]}{\kappa_{x,K}} - \delta_{x,K}\delta_{x,K}^\top. \quad (5.35)$$

The exact q_x -weighted mean of $A_x(Z)$ is Σ_x^* . Applying Lemma 5.4 to A_x and using (5.35) gives

$$\Sigma_{x,K}^* - \Sigma_x^* = \frac{\mathbb{E}[(A_x(Z) - \Sigma_x^*)e_{x,K}(U)]}{\kappa_{x,K}} - \delta_{x,K}\delta_{x,K}^\top. \quad (5.36)$$

Taking Frobenius norms and using $\|\delta\delta^\top\|_F = \|\delta\|_2^2$ proves (5.32). The moment assumption makes the constants finite, and $\epsilon_{x,K} = O(K^{-p})$ gives (5.33). $\square$

The explicit bounds show what is hidden by the order notation. The algebraic order is inherited without loss, but the constants contain $1/\kappa_x$. Therefore, the theorem is a fixed-threshold result; it is not a uniform statement as x moves indefinitely deeper into the tail. For a rarer event, a given absolute error in $q_{x,K}$ can produce a larger error in the normalised CE moments.

The same perturbation identity controls the scale component of the Student t -copula proposal. Define

$$m_{\log,x,K} = \frac{\mathbb{E}[(\log W)q_{x,K}(Z, W)]}{\kappa_{x,K}}, \quad m_{-1,x,K} = \frac{\mathbb{E}[W^{-1}q_{x,K}(Z, W)]}{\kappa_{x,K}}. \quad (5.37)$$

Corollary 5.6 (Student t -copula scale parameters). *Assume the conditions of Theorem 3.1,*

$$\mathbb{E}[|\log W| + W^{-1}] < \infty,$$

and (5.23). Let $(\alpha_{x,K}^, \beta_{x,K}^*)$ be the inverse-Gamma CE parameters obtained from the finite-COS moments in (5.37). Then*

$$|m_{\log,x,K} - m_{\log,x}| + |m_{-1,x,K} - m_{-1,x}| = O(K^{-p}), \quad (5.38)$$

and

$$|\alpha_{x,K}^* - \alpha_x^*| + |\beta_{x,K}^* - \beta_x^*| = O(K^{-p}). \quad (5.39)$$

Proof. Applying Lemma 5.4 to $g(W) = \log W$ and $g(W) = W^{-1}$ gives (5.38). The inverse-Gamma shape parameter is defined by

$$H(\alpha, m_{\log}, m_{-1}) := \psi(\alpha) - \log \alpha + m_{\log} + \log m_{-1} = 0. \quad (5.40)$$

At the exact solution,

$$\frac{\partial H}{\partial \alpha} = \psi_1(\alpha_x^*) - \frac{1}{\alpha_x^*} > 0, \quad (5.41)$$

where ψ_1 is the trigamma function. The implicit function theorem therefore makes α locally continuously differentiable, and hence locally Lipschitz, in $(m_{\log}, m_{-1})$. Finally, $\beta = \alpha/m_{-1}$ is smooth because $m_{-1,x} > 0$. This proves (5.39). $\square$

5.4 Finite-COS covariance and the combined calibration error

The exact covariance order in Theorem 5.1 does not hold literally for a finite COS expansion, because $q_{x,K}(U)$ is not exactly $\mathbb{E}[I_x | U]$. Nevertheless, the finite- K raw-moment covariance approaches the Rao–Blackwell covariance at the same rate as the conditional-probability approximation.

Define

$$G_{x,K}^{\text{ISCOS}} := R_h(U)q_{x,K}(U)T(U), \quad V_{x,K} := \text{Cov}_h(G_{x,K}^{\text{ISCOS}}), \quad V_x^{\text{RB}} := \text{Cov}_h(G_x^{\text{RB}}). \quad (5.42)$$

Proposition 5.7 (Finite- K perturbation of the Rao–Blackwell covariance). *Assume (5.23) and*

$$\mathbb{E}_h[R_h(U)^2\|T(U)\|_2^2] < \infty.$$

Then

$$\|V_{x,K} - V_x^{\text{RB}}\|_2 = O(K^{-p}). \quad (5.43)$$

If the parameter map $\mathcal{G}$ is twice continuously differentiable in a neighbourhood of the target raw moments, the corresponding finite- K asymptotic covariance of the fitted parameters is also an $O(K^{-p})$ perturbation of the exact Rao–Blackwell asymptotic covariance.

Proof. Write

$$D_{x,K} := G_{x,K}^{\text{ISCOS}} - G_x^{\text{RB}} = R_h(U)(q_{x,K}(U) - q_x(U))T(U). \quad (5.44)$$

The uniform COS bound implies

$$\|D_{x,K}\|_{L^2(h)} \leq \epsilon_{x,K} \left(\mathbb{E}_h[R_h(U)^2\|T(U)\|_2^2] \right)^{1/2} = O(K^{-p}). \quad (5.45)$$

Now expand

$$\text{Cov}(G_x^{\text{RB}} + D_{x,K}) - \text{Cov}(G_x^{\text{RB}}) = \text{Cov}(G_x^{\text{RB}}, D_{x,K}) + \text{Cov}(D_{x,K}, G_x^{\text{RB}}) + \text{Cov}(D_{x,K}).$$

Cauchy–Schwarz bounds the two cross-covariance terms by a constant times $\|D_{x,K}\|_{L^2(h)}$, while the final term is of order $\|D_{x,K}\|_{L^2(h)}^2$. This proves (5.43). The parameter statement follows by combining this bound with the $O(K^{-p})$ perturbation of the population raw moments and the continuity of the Jacobian of $\mathcal{G}$. $\square$

Proposition 5.7 gives a norm convergence result, not a finite- K Loewner order. At a particular resolution, the deterministic approximation error can move the covariance in either direction. What is guaranteed is that, as K increases, the finite-COS covariance approaches the lower Rao–Blackwell covariance.

We now add the pilot sampling error. For clarity, define the finite- K raw moment average

$$\widehat{m}_{x,K} = \frac{1}{M_0} \sum_{m=1}^{M_0} R_h(U^{(m)})q_{x,K}(U^{(m)})T(U^{(m)}), \quad (5.46)$$

with population mean $m_{x,K}$. When the ISCOS weights in (4.1) are used, denote the estimators in (4.2)–(4.3) by

$$\widehat{\mu}_{x,K}^{\text{ISCOS}} := \widehat{\mu}_\omega, \quad \widehat{\Sigma}_{x,K,\text{raw}}^{\text{ISCOS}} := \widehat{\Sigma}_{\omega,\text{raw}}.$$

These estimators are components of $\mathcal{G}(\widehat{m}_{x,K})$, while $(\mu_{x,K}^*, \Sigma_{x,K}^*)$ are components of $\mathcal{G}(m_{x,K})$.

Theorem 5.8 (Combined ISCOS calibration error). *Suppose that $\kappa_x > 0$, (5.23) holds, and*

$$\mathbb{E}_h \left[R_h(U)^2 (1 + \|Z\|_2^4) \right] < \infty. \quad (5.47)$$

Then, for the fixed threshold x ,

$$\left\| \hat{\mu}_{x,K}^{\text{ISCOS}} - \mu_x^\star \right\|_2 = O_{\mathbb{P}}(M_0^{-1/2}) + O(K^{-p}), \quad (5.48)$$

and

$$\left\| \hat{\Sigma}_{x,K,\text{raw}}^{\text{ISCOS}} - \Sigma_x^\star \right\|_{\text{F}} = O_{\mathbb{P}}(M_0^{-1/2}) + O(K^{-p}). \quad (5.49)$$

The same decomposition holds for the Student t -copula Gaussian and inverse-Gamma parameters under the additional moment conditions used in Corollary 5.6.

Proof. Because the clipped COS weights satisfy $0 \leq q_{x,K} \leq 1$, the moment condition (5.47) gives a multivariate central limit theorem for the sample normaliser, first moment, and second moment. By Theorem 5.3, $\kappa_{x,K} \geq \kappa_x/2$ for all sufficiently large K , so the ratio map is smooth in a neighbourhood of the finite- K population moments. The delta method therefore gives

$$\hat{\mu}_{x,K}^{\text{ISCOS}} - \mu_{x,K}^\star = O_{\mathbb{P}}(M_0^{-1/2}), \quad \hat{\Sigma}_{x,K,\text{raw}}^{\text{ISCOS}} - \Sigma_{x,K}^\star = O_{\mathbb{P}}(M_0^{-1/2}). \quad (5.50)$$

Using the decomposition

$$\begin{aligned} \hat{\mu}_{x,K}^{\text{ISCOS}} - \mu_x^\star &= (\hat{\mu}_{x,K}^{\text{ISCOS}} - \mu_{x,K}^\star) + (\mu_{x,K}^\star - \mu_x^\star), \\ \hat{\Sigma}_{x,K,\text{raw}}^{\text{ISCOS}} - \Sigma_x^\star &= (\hat{\Sigma}_{x,K,\text{raw}}^{\text{ISCOS}} - \Sigma_{x,K}^\star) + (\Sigma_{x,K}^\star - \Sigma_x^\star), \end{aligned}$$

and applying Theorem 5.5 proves (5.48)–(5.49). The Student t -copula conclusion follows in the same way from the smooth parameter maps and Corollary 5.6. $\square$

Theorem 5.8 separates the two tuning parameters clearly. For a fixed K , increasing M_0 removes the pilot sampling error but leaves convergence to the finite-COS target rather than to the exact CE target. To make the COS error negligible on the root- M_0 scale, one needs

$$K^{-p} = o(M_0^{-1/2}). \quad (5.51)$$

Balancing the two displayed orders gives the heuristic choice

$$K \asymp M_0^{1/(2p)}. \quad (5.52)$$

This balances deterministic and statistical error only. It is not necessarily work-optimal, because the direct calibration cost grows approximately linearly in both M_0 and K .

Under the usual additional uniform-integrability conditions for the ratio estimators, the componentwise mean-squared error corresponding to (5.48)–(5.49) has the form

$$\text{MSE} = O(M_0^{-1}) + O(K^{-2p}). \quad (5.53)$$

The qualification is relevant because a ratio estimator can have unusually large finite-sample moments if its random denominator is allowed to approach zero. In implementation, the total calibration weight and the effective sample size should therefore be monitored alongside the asymptotic rate.

Covariance regularisation contributes an additional deterministic term. For the ridge estimator in (4.10), and fixed dimension d ,

$$\left\| \widehat{\Sigma}_{x,K,\rho}^{\text{ISCOS}} - \Sigma_x^* \right\|_{\text{F}} = O_{\mathbb{P}}(M_0^{-1/2}) + O(K^{-p}) + O(\rho). \quad (5.54)$$

For the shrinkage estimator in (4.11), the analogous additional term is $O(\lambda)$. Consistency for the unregularised population covariance requires $\rho \rightarrow 0$ and $\lambda \rightarrow 0$. If the regularisation parameters are held fixed, the estimator instead converges to the corresponding regularised target, which may nevertheless be preferable for finite-sample importance-sampling stability.

5.5 Threshold estimation and scope of the result

The preceding results condition on a fixed threshold x . In the numerical implementation, x is obtained from a preliminary empirical VaR. For a discrete loss distribution, this additional step is qualitatively different from the smooth moment perturbations considered above.

Let $x_\alpha = \inf\{y : F_L(y) \geq \alpha\}$. If the confidence level lies strictly inside the jump at x_α , namely

$$F_L(x_\alpha^-) < \alpha < F_L(x_\alpha), \quad (5.55)$$

then the empirical quantile is locally stable. More precisely, for an independent preliminary sample of size M_{pre} , define

$$\delta_- = \alpha - F_L(x_\alpha^-) > 0, \quad \delta_+ = F_L(x_\alpha) - \alpha > 0.$$

Hoeffding's inequality gives the elementary bound

$$\mathbb{P}(\widehat{x}_\alpha \neq x_\alpha) \leq \exp(-2M_{\text{pre}}\delta_-^2) + \exp(-2M_{\text{pre}}\delta_+^2). \quad (5.56)$$

Indeed, selecting a loss below x_α requires the empirical CDF just below x_α to cross α , while selecting a larger loss requires the empirical CDF at x_α to fall below α . Under (5.55), both events are large deviations.

If α coincides with one side of a CDF jump, one of these gaps vanishes and the empirical quantile can move to an adjacent atom under a small CDF perturbation. In that case, the threshold error cannot in general be absorbed into the smooth $O_{\mathbb{P}}(M_0^{-1/2}) + O(K^{-p})$ expansion. It should be reported separately, controlled by a larger independent preliminary sample, or removed from the calibration comparison by fixing a common threshold across methods.

Finally, the covariance comparison in this section concerns proposal calibration, not every downstream importance-sampling estimator. A more accurate or lower-variance estimate of the CE moments need not produce a smaller variance for every final VaR, expected-shortfall, or obligor-level ratio estimator. The production variance also depends on the nonlinear likelihood ratio, the conditional Bernoulli twisting, the selected regularisation, and the particular final integrand. Moreover, the CE proposal is the Kullback–Leibler projection of the tail-conditioned factor law within the chosen family; it is not the unrestricted variance-minimising proposal for every possible downstream quantity.

The main conclusions can therefore be summarised as

$$\boxed{\text{Cov}_h(G_x^{\text{RB}}) \preceq \text{Cov}_h(G_x^{\text{CEIS}})} \quad (5.57)$$

for exact conditional weighting,

$$\boxed{\epsilon_{x,K} = O(K^{-p}) \implies \|\mu_{x,K}^* - \mu_x^*\|_2 + \|\Sigma_{x,K}^* - \Sigma_x^*\|_F = O(K^{-p})} \quad (5.58)$$

for the deterministic COS approximation, and

$$\boxed{\hat{\vartheta}_{x,K,M_0} - \vartheta_x^* = O_{\mathbb{P}}(M_0^{-1/2}) + O(K^{-p})} \quad (5.59)$$

for the combined calibration error, componentwise for the Gaussian proposal parameters and, under the stated moment conditions, for the Student t scale parameters.

6 Numerical Experiments: Gaussian Copula

Section 5 separates the sampling error of proposal calibration from the deterministic error of the finite COS approximation. This section studies both effects on an eleven-factor Gaussian-copula benchmark. The first experiment fixes the pilot factors and compares filtered COS weights with exact conditional tail probabilities obtained by block convolution. The second uses a common pilot, common production budgets, and common random-number streams to compare three practical factor proposals: the paper-aligned Glasserman mean shift $\mathcal{N}(\mu_x^{\text{GL}}, I_{11})$, indicator-weighted CEIS, and filtered ISCOS with exactly 32 modes. The Hessian covariance extension of the Glasserman construction is not used in this comparison.

We retain the historical terminology used throughout the manuscript:

$$\text{CVaR}_{k,\alpha} = \mathbb{E}[l_k Y_k \mid L = \text{VaR}_\alpha], \quad \text{CES}_{k,\alpha} = \mathbb{E}[l_k Y_k \mid L \geq \text{VaR}_\alpha]. \quad (6.1)$$

For a discrete loss, the threshold-tail mean $\mathbb{E}[L \mid L \geq \text{VaR}_\alpha]$ need not equal the quantile-integral definition of expected shortfall because the VaR atom may require partial inclusion.

All intervals reported below are conditional, nominal, marginal pointwise 95% intervals from the production ratio estimators. They do not propagate the uncertainty of the preliminary VaR, proposal calibration, or the finite COS approximation. They are not intervals for pairwise method differences, and interval overlap is not used as a significance test.

6.1 Benchmark portfolio and matched experimental design

We use Example 4 of Glasserman [4]. The portfolio contains 100 obligors divided into ten homogeneous blocks of ten. Every obligor has unconditional default probability 0.01. The first of eleven independent standard Gaussian factors is market-wide, while factor $g + 1$ is specific to block g . The loss-at-default in block g is c_g , with

$$(c_1, \dots, c_{10}) = (1, 1, 4, 4, 9, 9, 16, 16, 25, 25). \quad (6.2)$$

Thus each exposure level 1, 4, 9, 16, 25 occurs for twenty obligors.

The implementation uses the convention that positive factor realisations are adverse. For an obligor in block g ,

$$p_g(z) = \Phi\left(\frac{\Phi^{-1}(0.01) + 0.3z_1 + 0.8z_{g+1}}{\sqrt{0.27}}\right). \quad (6.3)$$

This is equivalent to the lower-tail convention of Section 2 after reversing the signs of the non-zero loadings. Conditional on $Z = z$, the block default counts are independent and

$$D_g \mid Z = z \sim \text{Binomial}(10, p_g(z)), \quad L \mid Z = z = \sum_{g=1}^{10} c_g D_g. \quad (6.4)$$

The loss lies on the integer lattice and satisfies $0 \leq L \leq 1100$.

A preliminary Monte Carlo sample gives the attainable 99.9% threshold $x = 250$, consistent with the benchmark in [4]. The main settings are collected in Table 1. The mode count includes the zero mode throughout.

Table 1 – Experimental design for the Gaussian-copula experiments.

Quantity	Specification
Confidence level and threshold	$\alpha = 0.999$, $x = \widehat{\text{VaR}}_\alpha = 250$
Shared pilot sample	$M_0 = 250\,000$, seed 42
Production budget per method	$M = 250\,000$ for $L = 250$ and $M = 250\,000$ for $L \geq 250$
Production seeds	exact-level event: 420; threshold-tail event: 421
Random-number coupling	the same base Gaussian draws and Bernoulli uniforms are used across methods within each event; the transformed factors and defaults still differ by proposal
Exact-convolution validation	$K \in \{16, 32, 64, 128, 256, 512, 1024\}$, EXP-4 filter
Matched-budget ISCOS	exactly $K = 32$ modes, $k = 0, \dots, 31$, with $\sigma_{k,32} = \exp[-8(k/32)^4]$
COS interval and lattice cut-off	$[a, b] = [-0.5, 1100.5]$, $y_x = 249.5$, loss step $\Delta = 1$
Implementation details	batches of 5000; ten homogeneous blocks; ridge $\rho = 10^{-8}$; raw COS probabilities projected onto $[0, 1]$

The three methods differ only in the Gaussian factor proposal used in the production stage. The benchmark proposals are

$$g_{\text{GL}}(z) = \phi_{11}(z; \mu_x^{\text{GL}}, I_{11}), \quad (6.5)$$

$$g_{\text{CEIS}}(z) = \phi_{11}(z; \hat{\mu}_x^{\text{CEIS}}, \hat{\Sigma}_x^{\text{CEIS}}), \quad \omega_m^{\text{CEIS}} = \mathbf{1}_{\{L^{(m)} \geq 250\}}, \quad (6.6)$$

$$g_{\text{ISCOS}}(z) = \phi_{11}(z; \hat{\mu}_{x,32}^{\text{ISCOS}}, \hat{\Sigma}_{x,32}^{\text{ISCOS}}), \quad \omega_m^{\text{ISCOS}} = q_{x,32}(Z^{(m)}). \quad (6.7)$$

The Glasserman mean is obtained by the saddle-point optimisation used in the paper's Algorithms 6.1 and 6.2, while its covariance remains the identity. CEIS and ISCOS use the same weighted Gaussian fit and the same ridge; only their calibration weights differ.

All three proposals are followed by the same conditional Bernoulli twist,

$$p_n^{(\theta)}(z) = \frac{p_n(z)e^{\theta l_n}}{1 + p_n(z)(e^{\theta l_n} - 1)}, \quad \psi(\theta, z) = \sum_{n=1}^N \log[1 + p_n(z)(e^{\theta l_n} - 1)]. \quad (6.8)$$

For $\{L = 250\}$, θ solves $\sum_n l_n p_n^{(\theta)}(z) = 250$. For $\{L \geq 250\}$, the implementation uses $\theta^+(z) = \max\{\theta(z), 0\}$, following [5, 4]. If R_Z denotes the Gaussian factor likelihood ratio, the complete production likelihood ratio is

$$R_Z(z) \exp\{-\theta L + \psi(\theta, z)\}.$$

Thus the credit model, conditional twist, event definitions, and final ratio estimators are identical across methods.

6.2 Accuracy of the filtered COS calibration

The block structure permits an exact conditional benchmark. For each pilot factor draw $Z^{(m)}$, finite-state convolution of the ten binomial block counts gives

$$q_x^{\text{ex}}(Z^{(m)}) = \mathbb{P}\left(\sum_{g=1}^{10} c_g D_g \geq 250 \mid Z^{(m)}\right). \quad (6.9)$$

The dynamic program retains states below the threshold and accumulates mass as soon as the threshold is crossed. This avoids subtracting two nearly equal probabilities when the conditional tail is very small.

On the common finite pilot, the exact-weight Gaussian fit is

$$\hat{\mu}_x^{\text{ex}} = \frac{\sum_m q_x^{\text{ex}}(Z^{(m)}) Z^{(m)}}{\sum_m q_x^{\text{ex}}(Z^{(m)})}, \quad (6.10)$$

$$\hat{\Sigma}_x^{\text{ex}} = \frac{\sum_m q_x^{\text{ex}}(Z^{(m)}) (Z^{(m)} - \hat{\mu}_x^{\text{ex}})(Z^{(m)} - \hat{\mu}_x^{\text{ex}})^\top}{\sum_m q_x^{\text{ex}}(Z^{(m)})} + 10^{-8} I_{11}. \quad (6.11)$$

These quantities are exact only with respect to the conditional loss calculation; they remain finite-pilot estimates of the population CE moments. For the filtered COS fit, define

$$e_\mu(K) = \frac{\|\hat{\mu}_{x,K}^{\text{ISCOS}} - \hat{\mu}_x^{\text{ex}}\|_2}{\|\hat{\mu}_x^{\text{ex}}\|_2}, \quad e_\Sigma(K) = \frac{\|\hat{\Sigma}_{x,K}^{\text{ISCOS}} - \hat{\Sigma}_x^{\text{ex}}\|_F}{\|\hat{\Sigma}_x^{\text{ex}}\|_F}. \quad (6.12)$$

The exact conditional weights have sample mean 1.19926×10^{-3} and calibration ESS 587.60. In Table 2, “spurious mass” is the share of total clipped COS weight carried by pilot states with $q_x^{\text{ex}}(Z^{(m)}) \leq 10^{-12}$.

The 16-mode approximation is inadequate: most of its fitted weight is assigned to factor states whose exact conditional tail probability is negligible. The largest improvement occurs between $K = 16$ and $K = 32$. Thereafter the mean absolute weight error and the proposal-moment errors decrease overall, although not monotonically at every adjacent mode count. At $K = 1024$, the relative mean and covariance errors are 2.53% and 3.07%, respectively. The finite-pilot, clipping, and near-zero-weight effects visible in

Table 2 – EXP-4 COS calibration against exact conditional block convolution on the same 250 000 pilot factors.

K	$M_0^{-1} \sum_m q_{x,K}^{(m)}$	mean $ q_{x,K} - q_x^{\text{ex}} $	spurious mass	$e_\mu(K)$	$e_\Sigma(K)$
16	1.1699×10^{-2}	1.071×10^{-2}	86.7408%	94.59%	58.67%
32	1.1070×10^{-3}	2.082×10^{-4}	2.9546%	5.67%	8.95%
64	1.0369×10^{-3}	1.801×10^{-4}	0.2448%	6.44%	7.82%
128	1.0542×10^{-3}	1.568×10^{-4}	0.0012%	5.72%	6.72%
256	1.0764×10^{-3}	1.354×10^{-4}	0.0762%	4.78%	5.81%
512	1.0985×10^{-3}	1.073×10^{-4}	0.0276%	3.88%	4.71%
1024	1.1301×10^{-3}	7.125×10^{-5}	0.0008%	2.53%	3.07%




Figure 1 – Relative errors of the EXP-4 ISCOS proposal moments against exact conditional-convolution moments on the same pilot sample.

the table make this a convergence diagnostic rather than an estimate of the asymptotic exponent in Section 5.

The matched-budget ISCOS run uses exactly the filtered $K = 32$ specification in the second row of Table 2. Before clipping, its COS probabilities have sample mean 1.06296×10^{-3} , minimum -4.29299×10^{-2} , and maximum 1.004806. The fractions below zero and above one are 8.7728% and 0.0020%, respectively. After projection onto $[0, 1]$, the weight mean is 1.10700×10^{-3} and the calibration ESS is 601.8. These clipping fractions are useful numerical diagnostics but are not pointwise error measures; the exact conditional comparison is the relevant accuracy check.

6.3 Proposal-calibration diagnostics

For a fitted Gaussian proposal $(\hat{\mu}, \hat{\Sigma})$, define its relative distance from the exact-weight finite-pilot fit by

$$d_{\mu} = \frac{\|\hat{\mu} - \hat{\mu}_x^{\text{ex}}\|_2}{\|\hat{\mu}_x^{\text{ex}}\|_2}, \quad d_{\Sigma} = \frac{\|\hat{\Sigma} - \hat{\Sigma}_x^{\text{ex}}\|_{\text{F}}}{\|\hat{\Sigma}_x^{\text{ex}}\|_{\text{F}}}. \quad (6.13)$$

For the Glasserman proposal these are descriptive distances, because its mean-only mode approximation is not designed to match the CE covariance.

We also audit the Gaussian factor likelihood ratio. If $f = \mathcal{N}(0, I_d)$, $g = \mathcal{N}(\mu, \Sigma)$, and $R_Z = f/g$, then

$$\mathbb{E}_g[R_Z^2] < \infty \iff 2I_d - \Sigma^{-1} \succ 0 \iff \lambda_{\min}(\Sigma) > \frac{1}{2}. \quad (6.14)$$

We report the margin

$$m_{\text{LR}}(\Sigma) = \lambda_{\min}(2I_d - \Sigma^{-1}), \quad (6.15)$$

so a positive value means that the untruncated Gaussian factor ratio passes this second-moment check.

Table 3 – Proposal-calibration diagnostics under the common pilot budget. The exact conditional CE row uses block-convolution weights on the same finite pilot sample.

Method	mean weight	calibration ESS	d_{μ}	d_{Σ}	$\lambda_{\min}(\hat{\Sigma})$	$m_{\text{LR}}(\hat{\Sigma})$
Exact conditional CE	1.1993×10^{-3}	587.6	0.00%	0.00%	0.5050	0.0197
Glasserman $\mathcal{N}(\mu^{\text{GL}}, I)$	–	–	36.26%	59.66%	1.0000	1.0000
CEIS	1.1760×10^{-3}	294.0	7.04%	10.99%	0.4726	-0.1158
ISCOS-32 (EXP-4)	1.1070×10^{-3}	601.8	5.67%	8.95%	0.7296	0.6294

The CEIS pilot contains 294 realised tail observations, so its binary-weight ESS is exactly 294. The smooth ISCOS weights raise the calibration ESS to 601.8, an increase of 104.7% under the same pilot budget. In this run, filtered ISCOS-32 is also closer to the exact-weight fit than CEIS in both reported norms: d_{μ} falls from 7.04% to 5.67%, and d_{Σ} from 10.99% to 8.95%. The exact-weight covariance is itself close to the likelihood-ratio boundary, with $m_{\text{LR}} = 0.0197$. Sampling noise moves the CEIS covariance just across that boundary, whereas the filtered ISCOS covariance remains more diffuse and has a positive margin of 0.6294.

The Glasserman mean-only proposal automatically satisfies the Gaussian second-moment condition because its covariance is I_{11} . Its fitted mean shifts the market factor and the two industry factors attached to the largest-loss blocks most strongly, in line with the portfolio’s exposure profile. Its distances from the exact CE moments should not be interpreted as failure of its own objective: it is a local mode-based approximation with a fixed covariance, whereas CEIS and ISCOS are global moment fits.

6.4 Risk contributions and nominal pointwise intervals

Let A denote either $\{L = 250\}$ or $\{L \geq 250\}$, and let $\Lambda^{(m)}$ be the complete factor-and-default likelihood ratio. The obligor-level ratio estimator and the standard error used for the displayed bands are

$$\hat{r}_k(A) = \frac{\sum_{m=1}^M \Lambda^{(m)} l_k Y_k^{(m)} \mathbf{1}_{A^{(m)}}}{\sum_{m=1}^M \Lambda^{(m)} \mathbf{1}_{A^{(m)}}}, \quad (6.16)$$

$$\widehat{\text{se}}(\hat{r}_k(A)) = \frac{\left[\sum_{m=1}^M \left\{ \Lambda^{(m)} \mathbf{1}_{A^{(m)}} \left(l_k Y_k^{(m)} - \hat{r}_k(A) \right) \right\}^2 \right]^{1/2}}{\sum_{m=1}^M \Lambda^{(m)} \mathbf{1}_{A^{(m)}}}. \quad (6.17)$$

The reported interval is $\hat{r}_k \pm 1.96 \widehat{\text{se}}(\hat{r}_k)$. The event-weight ESS is

$$M_{\text{eff},A} = \frac{\left(\sum_m \Lambda^{(m)} \mathbf{1}_{A^{(m)}} \right)^2}{\sum_m \left(\Lambda^{(m)} \mathbf{1}_{A^{(m)}} \right)^2}.$$

Table 4 – Downstream importance-sampling diagnostics. Half-lengths are averages over the 100 obligors. All intervals are nominal and subject to the second-moment qualifications discussed in the text.

Method	CVaR hit rate	CVaR ESS	mean CVaR half-length	CES hit rate	CES ESS	mean CES half-length	tail-loss mean
Glasserman $\mathcal{N}(\mu^{\text{GL}}, I)$	12.6040%	402.5	0.2667	75.2372%	1291.6	0.2560	280.003
CEIS	13.8720%	5337.8	0.07623	70.3120%	29 354.6	0.05068	280.470
ISCOS-32 (EXP-4)	13.1556%	9404.8	0.05993	70.2516%	45 413.0	0.03916	280.508

The estimated exact-event probabilities are 2.77015×10^{-4} , 2.72541×10^{-4} , and 2.74853×10^{-4} for Glasserman, CEIS, and ISCOS, respectively. The corresponding tail-event estimates are 1.10660×10^{-3} , 1.15660×10^{-3} , and 1.15360×10^{-3} . All three CVaR allocation vectors add to 250 up to numerical rounding, while each CES vector adds to its estimated threshold-tail mean. For CEIS and ISCOS, these means are 280.470 and 280.508, a difference below 0.02%. Their contribution vectors are also close: the ISCOS–CEIS relative ℓ_2 differences are 1.52% for CVaR and 0.39% for CES. In the $l_k = 25$ group, an obligor contributes about 12.35 to CVaR and 11.47 to CES under both methods.

Raw proposal hit rates do not rank the methods by efficiency. Glasserman produces many event hits, but its event-weight ESS is only 402.5 for CVaR and 1291.6 for CES because a small number of likelihood weights dominate. Relative to CEIS, ISCOS raises the exact-event ESS by 76.2% and the tail-event ESS by 54.7%, despite slightly lower

hit rates. Its average nominal half-length is 21.4% smaller for CVaR and 22.7% smaller for CES. The componentwise CVaR ordering is mixed—ISCOS is narrower for 50 of 100 obligors—whereas it is narrower for 99 of 100 CES components. The reported average reductions therefore describe this matched run and should not be read as a uniform componentwise ordering.

The intervals require a separate integrability qualification. The Gaussian factor-ratio check in (6.14) passes for Glasserman and ISCOS but fails for CEIS. More strongly, a direction-wise calculation finds a CEIS covariance eigenvalue below $1/2$ whose signed eigenvector has strictly positive loadings for every obligor. This proves that the complete CEIS CES importance weight has an infinite second moment in the reported configuration. The CEIS CES bands are therefore finite-run dispersion summaries rather than CLT-valid confidence intervals. The same sufficient divergence diagnostic does not produce a witness for Glasserman or ISCOS, but failure to find a witness is not a proof that every complete event-weight second moment is finite. For the exact-level CVaR estimator, the event $\{L = 250\}$ may supply additional decay, and a complete integrability analysis is not undertaken. For these reasons all bands are described as nominal.

6.5 Computational cost and limitations

The homogeneous block structure reduces the cost of ISCOS calibration. Its conditional characteristic function is


$$\varphi_{L|z}(\omega) = \prod_{g=1}^{10} \left[1 - p_g(z) + p_g(z) e^{i\omega c_g} \right]^{10}, \quad (6.18)$$

so the implementation processes ten distinct Bernoulli groups rather than 100 individual obligors. The shared preliminary Monte Carlo stage takes approximately 1.45 seconds. In the saved fixed-order timing run, the equivalent stand-alone pipelines for all three methods are roughly 96–97 seconds; the production CVaR and CES simulations account for almost all of each bar. The additional filtered-COS calibration therefore has little effect on total cost in this example.

The timing comparison is descriptive. It consists of one sequential run in a fixed method order, without warm-up repetitions or randomisation of that order, and the recorded metadata do not identify the processor, memory, Python version, or numerical-library versions. The absolute times are therefore within-run measurements, not portable performance benchmarks.


The Gaussian experiments support three conclusions. First, the exact conditional-convolution study confirms that the filtered COS proposal moments approach their exact-weight counterparts as the resolution increases. Second, under the matched pilot, filtered ISCOS-32 uses substantially more calibration information than CEIS, is slightly closer to the exact finite-pilot CE moments, and remains on the finite-Gaussian-ratio side of the second-moment boundary. Third, its production weights are more balanced in this run, yielding higher event-weight ESS and shorter average nominal intervals, with the clearer componentwise improvement occurring for CES.

These conclusions remain limited to one pilot and one production run at a fixed threshold. The experiment does not estimate repeated-run bias, variance, coverage, or mean squared error; the displayed intervals exclude pilot and calibration uncertainty; common



(a) CVaR contributions conditional on $L = 250$. (b) CES contributions conditional on $L \geq 250$.

Figure 2 – Obligor-level contribution estimates with nominal marginal pointwise 95% intervals. The methods share the threshold, budgets, and base random-number streams within each event. The bands are not intervals for pairwise method differences.



(a) Nominal CVaR interval half-lengths.

(b) Nominal CES interval half-lengths.

Figure 3 – Pointwise interval half-lengths by obligor. The sharp increase for obligors 81–100 reflects the two blocks with loss-at-default 25.




Figure 4 – Equivalent full-pipeline wall-clock time by method. The same shared preliminary Monte Carlo cost is included in every bar so that each bar can be read as a stand-alone pipeline. Plotting and output writing are excluded.

random numbers do not convert marginal bands into paired-difference intervals; and closeness to the CE moment target does not imply variance optimality for every downstream ratio estimator. The second-moment warning for CEIS CES is especially important. The results thus support smooth conditional-probability weighting as a proposal-calibration strategy in this benchmark, not a universal downstream variance ordering.

7 Numerical Experiments: Student t -Copula

This section examines the Student t -copula implementation of the Gaussian–inverse-Gamma product proposal derived in Section 3. The experiment is deliberately matched to the preceding Gaussian study: it uses the same eleven-factor portfolio, the same pilot and production sample sizes, and the same definitions of the exact-level VaR contribution and the tail conditional expected-shortfall contribution. The comparison is between CEIS- t and ISCOS- t . A direct Student t analogue of the Glasserman mean-shift benchmark is not included in the supplied implementation.

The purpose of the experiment is therefore narrower than that of the Gaussian section. We ask whether replacing binary pilot indicators by COS conditional tail probabilities produces a more stable product-proposal fit and more balanced production likelihood weights when the common state contains both a Gaussian factor vector and a heavy-tailed scale variable. The experiment is a single matched-budget comparison rather than a COS-resolution or repeated-run convergence study.

Throughout this section, “CVaR contribution” retains the convention

$$\mathbb{E}[l_k Y_k \mid L = \text{VaR}_\alpha],$$

whereas CES denotes

$$\mathbb{E}[l_k Y_k \mid L \geq \text{VaR}_\alpha].$$

All confidence intervals reported below are conditional, nominal, marginal pointwise intervals from the production ratio estimators. They do not include uncertainty from the pilot VaR, proposal calibration, or the finite COS expansion, and they are not intervals for a CEIS-t minus ISCOS-t difference.

7.1 Portfolio, Product Proposal, and Reproducible Setup

We reuse the 100-obligor, eleven-factor portfolio described in the Gaussian experiment. The first factor is market-wide and the remaining ten factors are industry-specific. Each obligor has unconditional default probability 0.01, and the block losses-at-default are

$$(c_1, \dots, c_{10}) = (1, 1, 4, 4, 9, 9, 16, 16, 25, 25),$$

with ten obligors in every block. Hence $0 \leq L \leq 1100$ and the loss lies on the integer lattice.

For the Student t -copula, the common state is

$$U = (Z, W), \quad Z \sim \mathcal{N}_{11}(0, I_{11}), \quad W = \frac{\nu}{V}, \quad V \sim \chi_\nu^2,$$

with Z and W independent under the original measure. We set $\nu = 4$, so that

$$W \sim \text{InvGamma}(2, 2)$$

under the shape-scale parameterisation used in this paper. Conditional on $(Z, W) = (z, w)$, the number of defaults in block g is

$$D_g \mid Z = z, W = w \sim \text{Binomial}(10, p_g(z, w)), \quad g = 1, \dots, 10,$$

independently across blocks, where

$$p_g(z, w) = \Phi \left(\frac{T_4^{-1}(0.01)/\sqrt{w} - 0.3z_1 - 0.8z_{g+1}}{\sqrt{0.27}} \right). \quad (7.1)$$

This is the lower-tail sign convention used in the theoretical sections: a more negative factor realisation increases conditional default probabilities. Consequently, adverse proposal mean shifts appear with negative signs in the reported parameter estimates.

Both methods fit the independent product family

$$g(z, w) = \phi_{11}(z; \mu, \Sigma) f_{\text{IG}}(w; a_W, b_W), \quad (7.2)$$

where the Gaussian component matches weighted first and second moments of Z , while the inverse-Gamma component matches the weighted sufficient statistics $\log W$ and W^{-1} . CEIS-t uses the calibration weight $\mathbf{1}_{\{L \geq x\}}$. ISCOS-t replaces it by the clipped EXP-4 COS approximation of $\mathbb{P}(L \geq x \mid Z, W)$, using 64 modes including the zero mode.

Table 5 records the common experimental design. The two methods share the same pilot sample and empirical VaR. In the production stage, the implementation also shares

Table 5 – Parameters used in the Student t -copula comparison. The mode count includes the zero mode.

Quantity	Value
Degrees of freedom	$\nu = 4$
Confidence level and threshold	$\alpha = 0.999$, $x = \widehat{\text{VaR}}_\alpha = 504$
Shared pilot sample	$M_0 = 250,000$, seed 42
Production samples per method	250,000 for $L = 504$ and 250,000 for $L \geq 504$
Production seeds	exact-level: 420; tail: 421
Random-number coupling	proposal-independent inverse-CDF common random numbers
COS specification	$K = 64$, EXP-4 filter $\exp[-8(k/K)^4]$
COS interval and cutoff	$[a, b] = [-0.5, 1100.5]$, $y_x = 503.5$
Batch size and homogeneous groups	5,000; ten groups of ten obligors
Covariance regularisation	ridge $\rho = 10^{-8}$
Probability safeguard	raw COS tail probabilities projected onto $[0, 1]$

proposal-independent base random numbers: standard normals for the Gaussian component, uniforms transformed by the inverse-Gamma inverse CDF, and uniforms for the conditional Bernoulli defaults. This common-random-number construction makes the method comparison more directly aligned, but it does not turn the marginal intervals into confidence intervals for method differences.

The production estimator changes both components of the common state and also exponentially twists the conditional Bernoulli probabilities. Its full likelihood ratio is therefore the product of the Gaussian factor ratio, the inverse-Gamma scale ratio, and the conditional Bernoulli ratio. The same production estimator and the same twisting rule are used after CEIS- t and ISCOS- t calibration; only the fitted product proposal differs.

7.2 Proposal Calibration

Table 6 summarises the fitted proposals. The CEIS- t pilot contains 255 realised tail hits, so its binary calibration-weight ESS is exactly 255. The smooth ISCOS- t weights have an ESS of 353.5, an increase of 38.6% under the same pilot budget. The two fitted Gaussian means are similar: their Euclidean distance is 7.66% of the CEIS- t mean norm. The relative Frobenius distance between the two covariance matrices is 12.61%. These are between-method distances, not errors, because no exact conditional-moment benchmark is available for this Student t run.

Both methods shift the market factor most strongly, with fitted means -1.298 for CEIS- t and -1.333 for ISCOS- t . The largest industry shifts occur in the two blocks with loss-at-default 25, where the fitted means are approximately $(-0.757, -0.838)$ and $(-0.797, -0.841)$, respectively. Thus, the calibrated stress direction is economically consistent with the portfolio’s lumpy exposure profile.

The COS diagnostics require some care. Before clipping, the ISCOS- t weights have sample mean 1.06625×10^{-3} ; after clipping, the mean is 1.06887×10^{-3} , so clipping changes the mean by only 2.62×10^{-6} . Nevertheless, 94.14% of the raw values are negative and 94.15% of the clipped values are exactly zero. This large fraction is compatible with

Table 6 – Student t -copula proposal-calibration diagnostics. Here a_W and b_W are the inverse-Gamma shape and scale, and $\kappa_2(\hat{\Sigma})$ is the spectral condition number.

Method	mean weight	calibration ESS	$\hat{a}_W$	$\hat{b}_W$	$\lambda_{\min}(\hat{\Sigma})$	$\kappa_2(\hat{\Sigma})$
CEIS-t	1.0200×10^{-3}	255.0	1.9500	36.1980	0.5508	2.658
ISCOS-t	1.0689×10^{-3}	353.5	1.8928	33.8338	0.5933	2.305

Table 7 – Downstream Student t -copula importance-sampling diagnostics. CI half-lengths are averages over the 100 obligors and are nominal; the second-moment qualification in the text is essential.

Method	CVaR hit rate	CVaR ESS	mean CVaR half-length	CES hit rate	CES ESS	mean CES half-length
CEIS-t	0.8152%	257.4	0.6404	77.7988%	22,688.8	0.06891
ISCOS-t	0.8060%	326.7	0.5775	77.5692%	43,964.5	0.05031

the fact that most original-measure pilot states have an extremely small conditional probability of producing a loss above 504, and the raw negative values can therefore be small oscillatory approximations around zero. It is not, however, a substitute for a pointwise accuracy check. In contrast to the Gaussian experiment, the supplied Student t results contain neither an exact conditional-convolution benchmark nor a sweep over K . The 64-mode specification should therefore be interpreted as the implemented numerical choice rather than as an empirically validated asymptotic regime.

A second qualification concerns likelihood-ratio moments. For the product proposal in (7.2), the unconditional common-state likelihood ratio has a finite second moment only if

$$\lambda_{\min}(\Sigma) > \frac{1}{2}, \quad a_W < \nu, \quad b_W < \nu. \quad (7.3)$$

The Gaussian condition is satisfied by both fitted covariances, and both inverse-Gamma shapes are below $\nu = 4$. The fitted scales, however, are 36.198 and 33.834, so the scale condition fails for both methods. Consequently, the common-state second-moment diagnostic does not support a central limit theorem for either set of displayed intervals. This diagnostic is a warning rather than a complete analysis of every event-weighted estimator: the event indicator and the conditional Bernoulli twist may provide additional tail decay. We therefore describe all intervals below as nominal finite-run dispersion measures.

7.3 Risk Contributions and Nominal Pointwise Intervals

Table 7 reports the main production diagnostics. The raw proposal hit rates are nearly the same for the two methods and are slightly lower under ISCOS-t. The event-weight ESS nevertheless increases from 257.4 to 326.7 for the exact-level event and from 22,688.8 to 43,964.5 for the tail event. Thus, relative to CEIS-t, ISCOS-t raises the exact-event ESS by 26.9% and the tail-event ESS by 93.8%. The gain comes from more balanced likelihood weights rather than from a higher event frequency.

The two proposals give close estimates of the underlying risk quantities. The estimated exact-level probabilities are 8.4364×10^{-6} under CEIS-t and 8.5581×10^{-6} under ISCOS-t. The corresponding CVaR contribution vectors satisfy the allocation identity

$$\sum_{k=1}^{100} \widehat{\text{CVaR}}_{k,\alpha} = 504$$

up to numerical rounding. For the tail event, the estimated probabilities are 1.0298×10^{-3} and 1.0223×10^{-3} , and the threshold-based tail means are 601.026 and 601.465. The two tail means differ by less than 0.1%.

The contribution profile follows the five exposure levels. Under CEIS-t and ISCOS-t, the average CES contributions per obligor in the exposure groups 1, 4, 9, 16, 25 are approximately

$$(0.342, 1.460, 3.827, 8.148, 16.274)$$


and

$$(0.341, 1.472, 3.821, 8.189, 16.250),$$

respectively. The CVaR contribution estimates are noisier because the exact-level event has an event-weight ESS of only a few hundred. Even so, both methods reproduce the same blockwise allocation pattern and place the largest contributions in the two highest-exposure blocks.


Figure 5 shows the point estimates and nominal pointwise intervals. The methods are almost indistinguishable for CES, whereas the exact-level CVaR curves display greater within-block variation. ISCOS-t has a smaller interval half-length for 92 of the 100 CVaR components and for all 100 CES components. Averaged over obligors, its nominal half-length is 9.8% smaller for CVaR and 27.0% smaller for CES.

CEIS-t vs ISCOS-t, Student-t copula
 $\nu = 4$, $\alpha = 0.999$, VaR = 504



(a) CVaR contributions conditional on $L = 504$.

CEIS-t vs ISCOS-t, Student-t copula
 $\nu = 4$, $\alpha = 0.999$, VaR = 504



(b) CES contributions conditional on $L \geq 504$.

Figure 5 – CEIS-t and ISCOS-t obligor-level contribution estimates with nominal pointwise 95% intervals. The methods share the pilot sample, VaR threshold, production budgets, and proposal-independent base random numbers. The intervals are descriptive because the common-state likelihood-ratio second-moment condition fails for both proposals.

Figure 6 plots the componentwise ratio

$$\frac{\text{CEIS-t CI half-length}}{\text{ISCOS-t CI half-length}}.$$

Values above one favour ISCOS-t. The median ratios are 1.114 for CVaR and 1.396 for CES. The systematic separation of the CES curve from one is consistent with the much larger tail-event ESS under ISCOS-t. These ratios are comparisons of marginal nominal intervals; they are not test statistics for a method difference.




Figure 6 – Componentwise nominal CI half-length ratio, CEIS-t divided by ISCOS-t. Values above one indicate a shorter interval under ISCOS-t.

7.4 Computational Cost

The shared initial simulation takes 2.015 seconds. CEIS-t calibration is almost cost-less relative to production, taking 0.101 seconds, whereas ISCOS-t calibration takes 2.656 seconds because it evaluates the grouped conditional characteristic function at 64 frequencies. The grouping reduces the conditional characteristic-function product from 100 obligors to ten distinct blocks.

The production stages dominate the total cost. The exact-level and tail runs take 87.469 and 96.652 seconds after CEIS-t calibration, compared with 90.527 and 99.029 seconds after ISCOS-t calibration. Counting the shared initial simulation once for each stand-alone pipeline gives equivalent total times of 186.237 seconds for CEIS-t and 194.226 seconds for ISCOS-t. ISCOS-t therefore increases total runtime by 7.99 seconds, or 4.3%, while approximately doubling the tail-event ESS and reducing the average nominal CES interval half-length by 27.0%.




Figure 7 – Matched-budget wall-clock times for the Student t -copula comparison. The same initial Monte Carlo cost is displayed for both methods so that each set of bars can be read as a stand-alone pipeline.

As in the Gaussian experiment, the recorded metadata do not identify the processor, memory, Python version, or numerical-library versions. The absolute times should therefore be read as within-run comparisons rather than as portable performance benchmarks.

7.5 Summary and Limitations of the Student t Experiment

The matched Student t -copula run gives the same broad empirical message as the Gaussian experiment. Smooth conditional-probability weights increase the amount of information used in proposal calibration and lead to more balanced production likelihood weights. Under the supplied configuration, ISCOS-t raises calibration ESS by 38.6%, exact-event ESS by 26.9%, and tail-event ESS by 93.8%. Its average nominal pointwise intervals are shorter for both risk contributions, with the larger reduction occurring for CES, while total runtime remains within 4.3% of CEIS-t.

These findings remain descriptive for four reasons. First, the experiment contains one pilot and one production run, so it does not estimate repeated-run bias, variance, coverage, or mean squared error. Second, there is no exact conditional benchmark or mode sweep for the Student t COS weights. Third, the independent product proposal cannot represent the dependence between Z and W induced by conditioning on a large loss. Fourth, the fitted inverse-Gamma scales violate the common-state likelihood-ratio second-moment condition in (7.3). The numerical results therefore support the practical promise of ISCOS-t, especially for tail contributions, but they do not establish a universal variance ordering or CLT-valid confidence intervals.

8 Conclusion

This paper has developed ISCOS as a cross-entropy proposal-calibration method for rare-event simulation in multi-factor credit portfolios. The method starts from the

common-state distribution conditional on a large portfolio loss and projects it onto a tractable family. This gives a Gaussian proposal for the systematic factors and, for the Student t -copula, a Gaussian–inverse-Gamma product proposal for the factor vector and common scale. The construction remains in the low-dimensional common-state space and can be combined with conditional default twisting in the production stage.

ISCOS differs from CEIS only in the information used to fit that proposal. CEIS uses the realised indicator $\mathbf{1}_{\{L \geq x\}}$; ISCOS uses an approximation of

$$q_x(U) = \mathbb{P}(L \geq x \mid U).$$

Conditional independence makes the loss characteristic function available in closed form, so a filtered COS expansion can compute this smooth weight without an additional conditional default draw. The half-step convention for lattice losses evaluates the CDF away from its jump and therefore preserves the non-strict event $\{L \geq x\}$.

The analysis separates the benefit of conditional weighting from the error of the finite expansion. With exact q_x , ISCOS is a Rao–Blackwellisation of the CEIS raw sufficient-statistic estimator and has an explicitly smaller covariance matrix in the positive-semidefinite order. The corresponding ordering for the fitted proposal parameters is first-order asymptotic. If the conditional COS approximation is uniformly $O(K^{-p})$, the population proposal parameters converge at the same order, while a pilot sample of size M_0 contributes the usual $O_{\mathbb{P}}(M_0^{-1/2})$ term.

The numerical experiments support this calibration principle while also showing its limits. In the Gaussian model, exact conditional convolution confirms that the COS-calibrated moments approach their exact-weight counterparts as the resolution increases. In the matched run, filtered ISCOS-32 more than doubles the calibration ESS relative to CEIS, is slightly closer to the finite-pilot exact-weight moments, and remains on the finite-Gaussian-ratio side of the reported second-moment boundary. It also produces higher event-weight ESS and shorter average nominal intervals, particularly for CES, at essentially the same total cost. The componentwise CVaR interval ordering remains mixed, and the CEIS CES second-moment warning prevents interpreting these finite-run bands as a universal variance result.

The Student t -copula experiment reaches the same broad conclusion for a common state containing both Gaussian factors and a heavy-tailed scale. Under the reported configuration, ISCOS-t increases calibration ESS by 38.6% and tail-event ESS by 93.8%, while reducing the average nominal CES half-length by 27.0%; the equivalent stand-alone runtime is 4.3% higher than for CEIS-t. The gain comes from more balanced event weights rather than a higher proposal hit rate.

These results do not establish that ISCOS minimises the variance of every final VaR or tail-contribution estimator. The theoretical covariance order is a calibration result, the experiments are single matched runs rather than repeated-run studies, and the displayed intervals are nominal. In the Student t experiment, there is no exact conditional benchmark or COS mode sweep, the product proposal cannot represent the dependence between Z and W induced by the tail event, and the fitted inverse-Gamma scales fail the stated common-state likelihood-ratio second-moment condition. These qualifications are essential when interpreting the interval comparisons.

Further work should therefore focus on repeated, work-normalised comparisons; a Student t resolution study and conditional benchmark; and richer proposals that allow the

Gaussian factors to depend on the common scale. Adaptive choices of the pilot size, COS resolution, and covariance regularisation are also natural. The error decomposition suggests increasing K only until the finite-COS error is comparable with pilot sampling and regularisation error. More generally, ISCOS shows how conditional tail information can stabilise proposal calibration without changing the final importance-sampling estimator.

References

- [1] Pieter-Tjerk de Boer, Dirk P. Kroese, Shie Mannor, and Reuven Y. Rubinstein. A tutorial on the cross-entropy method. *Annals of Operations Research*, 134(1):19–67, 2005.
- [2] Stefano Demarta and Alexander J. McNeil. The t copula and related copulas. *International Statistical Review*, 73(1):111–129, 2005.
- [3] Fang Fang and Cornelis W. Oosterlee. A novel pricing method for european options based on fourier-cosine series expansions. *SIAM Journal on Scientific Computing*, 31(2):826–848, 2008.
- [4] Paul Glasserman. Measuring marginal risk contributions in credit portfolios. Working Paper 2005-01, Federal Deposit Insurance Corporation, Center for Financial Research, January 2005. FDIC Center for Financial Research Working Paper No. 2005-01.
- [5] Paul Glasserman and Jingyi Li. Importance sampling for portfolio credit risk. *Management Science*, 51(11):1643–1656, 2005.
- [6] Michael Kalkbrener. An axiomatic approach to capital allocation. *Mathematical Finance*, 15(3):425–437, 2005.
- [7] Reuven Y. Rubinstein and Dirk P. Kroese. *The Cross-Entropy Method: A Unified Approach to Combinatorial Optimization, Monte-Carlo Simulation and Machine Learning*. Information Science and Statistics. Springer, New York, NY, 2004.
- [8] Xiaoyu Shen, Fang Fang, and Chengguang Liu. The fourier cosine method for discrete probability distributions, 2024. arXiv:2410.04487v2.