

Adaptive singular-point method for pricing and hedging surrenderable equity-linked contracts

Andrea Molent*

Marcellino Gaudenzi†

Abstract

We propose a deterministic numerical method for pricing and hedging surrenderable equity-linked life-insurance contracts with periodic premiums and fund contributions, maturity and death guarantees, and Bermudan surrender under correlated stochastic volatility and stochastic interest rates. The main computational challenge is the non-recombining accumulated fund, which couples with multiple stochastic factors and early exercise. Our key idea is to avoid a full multidimensional fund lattice: variance and interest-rate factors are discretized on recombining lattices, while at each factor node the contract value is represented as an adaptive one-dimensional function of the fund. Periodic contributions then act as translations of the fund argument, whereas surrender is handled directly through a backward obstacle condition. Piecewise-cubic representations propagate payoff and exercise singularities and are compressed by continuous pruning criteria that control the representation error. We establish weak convergence of the financial chains, convergence of the adaptive valuation under vanishing representation error, and Delta consistency on regular fund regions. For the strict binomial scheme, additional regularity yields first-order weak accuracy and a Talay–Tubaro expansion supporting Richardson extrapolation. Numerical experiments show compact representations, favorable cost–accuracy, and close agreement with independent Monte Carlo and cross-fitted least-squares Monte Carlo benchmarks. Hedging results further show that contracts with similar values can generate materially different exposures to equity, volatility, and interest-rate risk.

Keywords: Adaptive approximation; Dynamic programming; Optimal stopping; Equity-linked insurance; Stochastic volatility.

1 Introduction

Equity-linked life-insurance contracts combine market participation with long-dated guarantees and, frequently, a right to surrender before maturity. Related unit-linked products remain economically significant in Europe: EIOPA reports about EUR 42 billion of gross written premium for these products in its 2024 sample [13]. These figures highlight the continuing practical relevance of reliable methods for the valuation, surrender analysis, and hedging of market-linked insurance contracts.

*Dipartimento di Scienze Economiche e Statistiche, Università degli Studi di Udine, Udine, Italy. Email: andrea.molent@uniud.it. ORCID: 0000-0002-0887-826X.

†Dipartimento di Scienze Economiche e Statistiche, Università degli Studi di Udine, Udine, Italy. Email: marcellino.gaudenzi@uniud.it. ORCID: 0000-0003-0366-3435.

The early literature established the valuation of asset-linked guarantees [4] and later incorporated stochastic interest rates [25], surrender and bonus mechanisms [18, 28], and periodic-premium contracts [1]. These features are economically important but create a challenging sequential decision problem: discounting, guarantee value, market volatility, mortality, future contributions, and the exercise decision evolve jointly over horizons of one or two decades.

From an operational-research perspective, the valuation is a finite-horizon stochastic dynamic program with a continuous, non-recombining resource state. Recombining trees can represent the financial factors, but annual additions to the accumulated fund prevent the fund itself from recombining. A direct grid in all state variables is therefore expensive. Approximate dynamic programming usually addresses such state-space growth by replacing the exact value function with a structured approximation [26]. Regression-based simulation is one important implementation, but in the present stopping problem it combines sampling, approximation, and stopping-rule errors [2, 3, 24]. Related work develops exact or error-controlled simulation schemes for stochastic-volatility models and path-dependent claims [6, 23]. Singular-point methods offer a deterministic, structure-exploiting alternative: they retain the value function only where contractual kinks or the backward recursion generate relevant nonlinear structure [8, 9, 14].

We develop this idea for a fully correlated Heston–CIR++ model. Heston stochastic variance captures leverage and time-varying equity risk [20]; the CIR++ rate specification preserves a non-negative square-root factor while fitting the observed initial term structure [7, 10]. Hybrid equity–rate models show that volatility, rates, and dependence can materially affect long-dated prices and risk exposures, including in settings with optimal surrender [15, 16, 17, 19, 21].

Within the proposed scheme, the variance and CIR rate factors are discretized jointly on recombining lattices so as to preserve their dependence structure, while at each joint factor node the contract value is represented as an adaptive one-dimensional function of the accumulated fund. Consequently, the contract value depends on the past only through the current fund, variance, and interest-rate states; no additional variable is needed to record earlier contributions. This representation naturally accommodates periodic contributions and surrender decisions at prescribed dates while concentrating computational effort where the value function exhibits relevant nonlinear features.

The contributions of the paper are threefold. First, we represent the non-recombining fund dimension through an adaptive value function rather than a fixed global grid. Periodic contributions then act simply by shifting the fund argument, while annual surrender decisions can be incorporated directly within the same deterministic backward recursion. This preserves the relevant nonlinear structure of the contract value while avoiding the state-space growth associated with a full discretization of the fund variable.

Second, we extend singular-point dynamic programming to a fully correlated Heston–CIR++ setting by combining recombining discretizations of stochastic variance and interest rates with the adaptive representation of the fund dimension. The resulting construction retains a deterministic backward recursion while accommodating stochastic volatility, stochastic rates, their dependence, mortality, and Bermudan surrender.

Third, we develop a convergence and error-control framework that distinguishes the financial time-discretization error from the error introduced by the adaptive representation and the

numerical boundaries. We establish convergence of the contract values and, under explicit regularity and representation conditions, consistency of the fund Delta. For the binomial discretization, stronger regularity assumptions also yield a first-order weak-error expansion and a theoretically supported Richardson benchmark.

The numerical study assesses accuracy and computational efficiency against independent Monte Carlo and least-squares Monte Carlo benchmarks, and uses the resulting framework to investigate surrender behavior, model effects, and multifactor hedging.

The remainder of the paper is organized as follows. Section 2 defines the model and contract. Section 3 presents the numerical method, Section 4 states the convergence and error-control results, and Section 5 reports the numerical study.

2 Model and contract

We consider an equity-linked life-insurance contract in which the policyholder pays premiums over time and deterministic amounts are credited to an investment fund linked to an equity index. The contract provides guarantees at death and maturity and may also allow the policyholder to surrender at predetermined dates. If the contract is continued at such a date, the scheduled premium and fund contribution are paid and the fund subsequently evolves with the financial market until the next contractual event.

For the specification considered below, the contract has maturity $T \in \mathbb{N}$ years, with policy anniversaries indexed by $i = 0, \dots, T$, and a_0 denotes the issue age. We assume, for simplicity, that premiums and fund contributions are paid annually at policy anniversaries and remain constant over time: P denotes the annual premium and D the corresponding amount credited to the fund. Other deterministic payment frequencies can be accommodated by including the corresponding dates in the time grid.

2.1 Financial dynamics and curve fit

Under the risk-neutral measure $\mathbb{Q}$, the equity index S , instantaneous variance factor V , and non-negative rate factor X satisfy

$$dS_t = (r_t - q) S_t dt + \sqrt{V_t} S_t dW_t^S, \quad (1)$$

$$dV_t = \kappa_V(\theta_V - V_t) dt + \sigma_V \sqrt{V_t} dW_t^V, \quad (2)$$

$$dX_t = \kappa_r(\theta_r - X_t) dt + \sigma_r \sqrt{X_t} dW_t^r, \quad (3)$$

$$r_t = X_t + \varphi(t). \quad (4)$$

The Brownian correlation matrix is

$$\mathbf{R} = \begin{pmatrix} 1 & \rho_{SV} & \rho_{Sr} \\ \rho_{SV} & 1 & \rho_{Vr} \\ \rho_{Sr} & \rho_{Vr} & 1 \end{pmatrix}, \quad 1 + 2\rho_{SV}\rho_{Sr}\rho_{Vr} - \rho_{SV}^2 - \rho_{Sr}^2 - \rho_{Vr}^2 \geq 0. \quad (5)$$

Between contract dates, the accumulated fund F has the same proportional return as S ,

$$dF_t = (r_t - q) F_t dt + \sqrt{V_t} F_t dW_t^S, \quad F_i^+ = F_i^- + D, \quad i = 0, \dots, T-1. \quad (6)$$

Here F_i^- and F_i^+ denote the fund values immediately before and after the contribution at anniversary i , respectively. The jump relation applies whenever the contract is in force and, at a surrender anniversary, is continued. The deterministic shift φ in (4) is chosen so that the model reproduces the initial market discount curve exactly. Let $P^{\text{CIR}}(0, u)$ denote the zero-coupon bond price generated by the unshifted CIR factor X . Since $r_t = X_t + \varphi(t)$, the model zero-coupon price with maturity u is

$$P(0, u) = P^{\text{CIR}}(0, u) \exp\left(-\int_0^u \varphi(s) ds\right).$$

Therefore, fitting the initial market discount curve exactly requires

$$P^M(0, u) = P^{\text{CIR}}(0, u) \exp\left(-\int_0^u \varphi(s) ds\right), \quad 0 \leq u \leq T. \quad (7)$$

The empirical curve is represented by a Svensson specification; its formula and parameters are reported in the Supplementary Material.

2.2 Annual premiums, guarantees, mortality, and surrender

At inception and, subsequently, at each anniversary $i = 1, \dots, T-1$ at which the contract is continued, the policyholder pays the annual premium P and the corresponding amount D is credited to the investment fund; no premium or contribution is due at maturity. The contract may terminate upon death, through surrender at a prescribed anniversary, or at maturity. No surrender decision is available at inception.

Let g_m denote the accumulation rate used for the maturity and death guarantees, and let g_s denote the rate used for the guarantee component of the surrender payoff. For $\gamma \in \{g_m, g_s\}$ and $i = 1, \dots, T$, define

$$G_i^{(\gamma)} = \sum_{j=0}^{i-1} D \exp\{\gamma(i-j)\}.$$

For a contract still in force at anniversary i , $G_i^{(\gamma)}$ is the accumulated value of the contributions credited at anniversaries $0, \dots, i-1$. If the policyholder is alive and the contract remains in force at maturity, the maturity benefit, as a function of the fund value immediately before settlement, is $\max\{F, G_T^{(g_m)}\}$.

Mortality is assumed to be independent of the financial factors. Let q_x denote the one-year probability of death between ages x and $x+1$, conditional on survival to age x . Hence, for $i = 0, \dots, T-1$, q_{a_0+i} is the probability of death during policy year $(i, i+1]$, conditional on being alive at anniversary i . If the contract is in force immediately after anniversary i and death occurs during that policy year, the death benefit paid at anniversary $i+1$ is $B_{i+1}(F) = \max\{F, G_{i+1}^{(g_m)}\}$, where F denotes the fund value immediately before settlement. Since mortality is independent of the financial factors, the mortality probabilities enter the backward recursion as deterministic

actuarial weights.

At each surrender anniversary $i = 1, \dots, T - 1$, conditional on the policyholder being alive and the contract being in force, the surrender decision is made before the new premium and fund contribution. Let F_i^- denote the fund value immediately before this decision and let $\alpha_i^s \in (0, 1]$ be the fraction of the fund returned under the fund-based component. We consider the fund-based, guarantee-based, and mixed surrender payoffs

$$R_{\text{FB}}(i, F) = \alpha_i^s F, \quad R_{\text{GB}}(i, F) = G_i^{(gs)}, \quad R_{\text{MX}}(i, F) = \max\{\alpha_i^s F, G_i^{(gs)}\}. \quad (8)$$

If surrender is exercised under contract J , the amount $R_J(i, F_i^-)$ is paid at anniversary i . No premium is paid and no fund contribution is credited at that anniversary, and the contract terminates. Because the decision precedes the contribution due at anniversary i , $G_i^{(gs)}$ contains only the contributions previously credited at anniversaries $0, \dots, i - 1$.

If the contract is continued, the policyholder pays the premium P and the amount D is credited to the fund, so that $F_i^+ = F_i^- + D$. The financial state then evolves from (F_i^+, V_i, X_i) over the following policy year $(i, i + 1]$. If death occurs during that year, $B_{i+1}(F_{i+1}^-)$ is paid at anniversary $i + 1$. If the policyholder survives, the contract proceeds to the next surrender anniversary, or to the maturity benefit when $i + 1 = T$.

For a fixed annual premium P , let $U_i(F, v, x)$ denote the contract value immediately before the surrender decision at anniversary i , and let $C_i(F, v, x)$ denote the value obtained by choosing continuation from the same pre-decision state. Thus, for $i = 1, \dots, T - 1$,

$$U_i(F, v, x) = \max\{C_i(F, v, x), R_J(i, F)\}, \quad J \in \{\text{FB}, \text{GB}, \text{MX}\}. \quad (9)$$

Let P_{NS} denote the fair annual premium of the no-surrender contract, and let P_J , $J \in \{\text{FB}, \text{GB}, \text{MX}\}$, denote the corresponding fair annual premiums of the surrenderable contracts.

Conditional on the contract being in force, the valuation state, apart from time, is (F, V, X) . The current fund value incorporates all previously credited contributions and their subsequent investment performance, while the guarantee levels are deterministic functions of the anniversary index. Therefore, no additional state variable is required to record the contribution history.

3 Adaptive singular-point dynamic programming

Let N_{yr} be the number of numerical time steps per year, set $N = TN_{\text{yr}}$ and $h = 1/N_{\text{yr}} = T/N$, and define $t_n = nh$, $n = 0, \dots, N$. Thus anniversary i corresponds to the grid time $t_{iN_{\text{yr}}} = i$. At time t_n , (j, k) indexes a joint variance-rate node. For a generic square-root factor $Z \in \{V, X\}$, $(\kappa_Z, \theta_Z, \sigma_Z)$ denotes its mean-reversion speed, long-run mean, and volatility coefficient. For convenience, $D_n = D$ when t_n is a payment anniversary $i = 0, \dots, T - 1$, and $D_n = 0$ otherwise; the premium outflow P likewise enters the continuation value only at payment anniversaries.

3.1 Recombining square-root lattices

Both CIR-type factors $Z \in \{V, X\}$ are represented on non-negative recombining square-root lattices. For parameters $(\kappa_Z, \theta_Z, \sigma_Z)$, the exact one-step conditional mean and variance are

$$m_Z(z) = \theta_Z + (z - \theta_Z)e^{-\kappa_Z h}, \quad v_Z(z) = \frac{\sigma_Z^2 z}{\kappa_Z} e^{-\kappa_Z h} (1 - e^{-\kappa_Z h}) + \frac{\theta_Z \sigma_Z^2}{2\kappa_Z} (1 - e^{-\kappa_Z h})^2. \quad (10)$$

The two lattice constructions use these moments differently. The adaptive-trinomial marginals match both $m_Z(z)$ and $v_Z(z)$ exactly. The binomial comparison matches $m_Z(z)$ exactly, while its conditional variance agrees with $v_Z(z)$ up to an $O(h^2)$ error on interior compact sets.

3.1.1 Binomial factor lattices

The fully binomial construction is retained as an independent deterministic comparison. For each factor, the grid spacing in the square-root coordinate is $\Delta_Z = \frac{\sigma_Z}{2} \sqrt{h}$, and consecutive time layers use opposite grid parities. From a parent state z , let $z_d < z_u$ be the two neighboring nodes on the next time layer that bracket the exact conditional mean, $z_d \leq m_Z(z) \leq z_u$. The upper transition probability is determined by

$$p_{Z,h}(z) = \frac{m_Z(z) - z_d}{z_u - z_d}, \quad \mathbb{P}_h(Z_{n+1} = z_u \mid Z_n = z) = p_{Z,h}(z), \quad (11)$$

with lower probability $1 - p_{Z,h}(z)$. Consequently, $\mathbb{E}_h[Z_{n+1} \mid Z_n = z] = m_Z(z)$ exactly, while the bracketing condition guarantees non-negative marginal probabilities.

At an interior state and for sufficiently small h , the selected successors are the adjacent opposite-parity nodes

$$z_{\pm} = \left(\sqrt{z} \pm \frac{\sigma_Z}{2} \sqrt{h} \right)^2. \quad (12)$$

The corresponding discrete conditional variance is

$$\hat{v}_{Z,h}(z) = p_{Z,h}(z)(1 - p_{Z,h}(z))(z_u - z_d)^2 = v_Z(z) + O(h^2) \quad (13)$$

uniformly on compact subsets of $(0, \infty)$. Thus the binomial marginal matches the exact CIR mean but only the leading-order CIR variance. Near the lower boundary or on coarse grids, the bracketing rule, rather than the fixed adjacent pair in (12), determines the two successors.

Let $p_V = p_{V,h}(v)$, $p_X = p_{X,h}(x)$ be the marginal upper probabilities at the current variance and rate-factor nodes. The joint 2×2 table is

$$\begin{aligned} p_{uu} &= p_V p_X + \delta, & p_{ud} &= p_V(1 - p_X) - \delta, \\ p_{du} &= (1 - p_V)p_X - \delta, & p_{dd} &= (1 - p_V)(1 - p_X) + \delta, \end{aligned} \quad (14)$$

where

$$\delta = \rho_{Vr} \sqrt{p_V(1 - p_V)p_X(1 - p_X)}. \quad (15)$$

This construction preserves both marginal transition laws and imposes the correlation ρ_{Vr} between their standardized binomial innovations. Equivalently, the discrete factor covariance is $\rho_{Vr} \sqrt{\hat{v}_{V,h}(v)\hat{v}_{X,h}(x)}$, and is therefore locally consistent with the target CIR covariance. Non-

negativity of all four joint probabilities is equivalent to the local Fréchet bounds

$$\max\{-p_V p_X, -(1-p_V)(1-p_X)\} \leq \delta \leq \min\{p_V(1-p_X), (1-p_V)p_X\}. \quad (16)$$

At a given time and parent state, let

$$\mathcal{I}_h = \{\rho \in [-1, 1] : p_{ab}(\rho) \geq 0 \text{ for all joint binomial branches } (a, b)\}$$

denote the local admissible correlation interval. Whenever the requested ρ_{Vr} lies outside $\mathcal{I}_h$, the binomial comparison replaces it by its projection onto this interval:

$$\rho_{Vr,h}^{\text{eff}} = \text{proj}_{\mathcal{I}_h}(\rho_{Vr}) = \arg \min_{\rho \in \mathcal{I}_h} |\rho - \rho_{Vr}|. \quad (17)$$

The dependence of $\mathcal{I}_h$ and $\rho_{Vr,h}^{\text{eff}}$ on time and the parent state is suppressed in the notation. The projection changes only the dependence between the two factors and leaves both marginal transition laws unchanged. Its asymptotic irrelevance is analyzed in Section 4.3.

3.1.2 Adaptive-trinomial factor lattices

For the trinomial construction, the grid is uniform in the Lamperti coordinate $\chi_Z = 2\sqrt{Z}/\sigma_Z$:

$$\chi_{Z,j} = \chi_{Z,0} + j\lambda_Z\sqrt{h}, \quad Z_j = \left(\frac{\sigma_Z}{2} \max\{\chi_{Z,j}, 0\}\right)^2. \quad (18)$$

The grid itself is fixed; adaptivity enters through the nodewise choice of the transition stencil. At each parent state z , the algorithm selects the narrowest ordered triplet of grid nodes, not necessarily consecutive, for which non-negative probabilities p_i^Z , $i = 1, 2, 3$, satisfy

$$\sum_{i=1}^3 p_i^Z = 1, \quad \sum_{i=1}^3 p_i^Z Z'_i = m_Z(z), \quad \sum_{i=1}^3 p_i^Z (Z'_i - m_Z(z))^2 = v_Z(z), \quad (19)$$

where Z'_i are the selected successor values. Thus both exact marginal CIR moments are matched at every active node.

The scale restriction $0 < \lambda_Z < 2$ has a direct positivity interpretation. If the conditional mean m lies between adjacent grid values $z_\ell < m < z_u$, the smallest variance attainable by a grid-supported distribution with mean m is $(m - z_\ell)(z_u - m)$. Away from the lower boundary, the worst-case ratio of this minimum to the leading CIR conditional variance is asymptotically $\lambda_Z^2/4$. Values of λ_Z close to two therefore leave little probability-positivity margin, whereas smaller values increase the number of factor nodes.

Because the Heston variance factor V and the CIR rate factor X underlying the CIR++ specification have markedly different mean-reversion dynamics, their grid scales are allowed to differ. A numerical screening of candidate pairs selected $(\lambda_V, \lambda_X) = (1, 1.4)$ for the central parameterization. Among the screened configurations, this pair provided the best end-to-end pricing accuracy while maintaining exact non-negative moment matching, strict local coupling, and a moderate joint state-space size.

For a parent state z and a selected successor Z'_i , define the centered variance-one marginal

innovation by

$$z_i^Z = \frac{Z'_i - m_Z(z)}{\sqrt{v_Z(z)}}.$$

For trinomial marginal probabilities p_i^V, p_j^X , the strict implementation uses the one-parameter covariance-preserving joint table

$$\pi_{ij} = p_i^V p_j^X (1 + \rho_{Vr} z_i^V z_j^X). \quad (20)$$

Because the marginal innovations are centered and have unit variance, this construction preserves both marginal transition laws and reproduces the target innovation correlation ρ_{Vr} . Since $p_i^V, p_j^X \geq 0$, the coupling is strictly admissible whenever $1 + \rho_{Vr} z_i^V z_j^X > 0$ for every active pair (i, j) . For $(\lambda_V, \lambda_X) = (1, 1.4)$, a sufficient uniform interior range is $|\rho_{Vr}| < 5/14$. This range is broad relative to commonly used variance-rate correlations, which are often set to zero in hybrid stochastic-volatility/stochastic-rate models; see, for example, [21]. This is a sufficient range for the one-parameter covariance-preserving construction, not necessarily the widest covariance interval attainable among all 3×3 tables.

The stated interior range follows from the limiting marginal stencils. The next lemma records the calculation because strict local admissibility is part of the convergence argument, not merely an implementation diagnostic.

Lemma 3.1. *Consider a CIR factor Z on the square-root grid $Z_q = (\sqrt{Z_0} + q\lambda\sigma\sqrt{h}/2)^2$ and let K_Z be a compact subset of $(0, \infty)$. For the symmetric stencil $(q-1, q, q+1)$, denote by $p_-(z, h)$, $p_0(z, h)$, and $p_+(z, h)$ the transition probabilities to Z_{q-1} , Z_q , and Z_{q+1} , respectively. Uniformly for $z \in K_Z$:*

(i) *if $\lambda > 1$, the symmetric stencil $(q-1, q, q+1)$ is admissible for all sufficiently small h , with*

$$p_0(z, h) = 1 - \frac{1}{\lambda^2} + O(h); \quad (21)$$

$$p_{\pm}(z, h) = \frac{1}{2\lambda^2} \pm \frac{\kappa(\theta - z) - \sigma^2/4}{2\lambda\sigma\sqrt{z}}\sqrt{h} + O(h), \quad (22)$$

(ii) *if $\lambda = 1$, define*

$$A(z) = \kappa - \frac{\kappa^2(\theta - z)^2}{\sigma^2 z} - \frac{\sigma^2}{16z}. \quad (23)$$

When $A(z) > 0$, the symmetric stencil is admissible and

$$p_0(z, h) = A(z)h + O(h^2), \quad (24)$$

$$p_{\pm}(z, h) = \frac{1}{2} \pm \frac{\kappa(\theta - z) - \sigma^2/4}{2\sigma\sqrt{z}}\sqrt{h} - \frac{1}{2}A(z)h + O(h^{3/2}). \quad (25)$$

When $A(z) < 0$, the minimal-span search selects a one-sided width-three stencil whose limiting probabilities are, up to reversal, $(1/6, 1/2, 1/3)$.

Proof. See Appendix A.

Proposition 3.2. *Let K be a compact subset of $(0, \infty) \times (0, \infty)$. For the adaptive-trinomial chain with $(\lambda_V, \lambda_X) = (1, 1.4)$, every correlation satisfying $|\rho_{Vr}| < \frac{5}{14}$ is strictly admissible on K*

for all sufficiently small h .

Proof. See Appendix A.

The bound $|\rho_{Vr}| < 5/14$ corresponds to the worst-case enlarged variance stencil, where $\max_{i,j} |z_i^V z_j^X|$ tends to $2 \times 1.4 = 2.8$. If the variance stencil remains symmetric, the same quantity tends to $1 \times 1.4 = 1.4$, and the local admissible range widens to $|\rho_{Vr}| \leq 5/7$.

Both joint factor constructions are combined with the conditional fund transition described in the next subsection.

3.2 Correlated fund transition

The following construction is common to the binomial and adaptive-trinomial factor lattices. At each time discretization, the deterministic CIR++ shift is calibrated on the discrete rate lattice so that the initial market discount curve is reproduced exactly at all grid maturities; details are given in the Supplementary Material.

At time t_n and parent node (j, k) , let b index a positive-probability joint factor branch. Its probability is $p_{n,b}$ and its child indices are (j_b, k_b) . The branchwise integrated short rate is approximated by

$$R_{n,b} = \frac{h}{2}(x_k + x_{k_b}) + \int_{t_n}^{t_{n+1}} \varphi(u) du.$$

The one-step fund transition uses the parent variance v_j .

In what follows, $\rho_{Vr,h}^{\text{eff}}$ denotes the variance–rate correlation actually used by the local joint coupling. For a strictly admissible coupling, $\rho_{Vr,h}^{\text{eff}} = \rho_{Vr}$; for the projected binomial fallback, it is the projected value defined in (17). We then set

$$\alpha_V = \frac{\rho_{SV} - \rho_{Sr}\rho_{Vr,h}^{\text{eff}}}{1 - (\rho_{Vr,h}^{\text{eff}})^2}, \quad \alpha_r = \frac{\rho_{Sr} - \rho_{SV}\rho_{Vr,h}^{\text{eff}}}{1 - (\rho_{Vr,h}^{\text{eff}})^2}, \quad \alpha_{\perp} = \left[1 - \frac{\rho_{SV}^2 - 2\rho_{SV}\rho_{Sr}\rho_{Vr,h}^{\text{eff}} + \rho_{Sr}^2}{1 - (\rho_{Vr,h}^{\text{eff}})^2} \right]^{1/2} \geq 0. \quad (26)$$

If η is independent with mean zero and variance one, then $\xi_S = \alpha_V \xi_V + \alpha_r \xi_r + \alpha_{\perp} \eta$ has unit variance and the required equity correlations. For branch b and quadrature point ℓ ,

$$F_{n+1} = (F_n + D_n)M_{n,b,\ell},$$

$$M_{n,b,\ell} = \exp\left\{R_{n,b} - qh - \frac{1}{2}v_j h + \sqrt{v_j h}(\alpha_V \xi_{V,b} + \alpha_r \xi_{r,b} + \alpha_{\perp} \eta_{\ell})\right\}. \quad (27)$$

The binomial comparison uses the two-point residual rule $\eta \in \{-1, 1\}$ with weights $(1/2, 1/2)$, yielding at most 8 hybrid branches per factor node. The production adaptive-trinomial scheme uses $\eta \in \{-\sqrt{3}, 0, \sqrt{3}\}$ with weights $(1/6, 2/3, 1/6)$, yielding at most 27 hybrid branches.

A local multiplicative martingale correction is applied to the fund multipliers. If $M_{n,b,\ell}^{(0)}$ denotes the uncorrected multiplier, set

$$\Lambda_{n,j,k} = \frac{e^{-qh}}{\sum_{b,\ell} p_{n,b,\ell} e^{-R_{n,b}} M_{n,b,\ell}^{(0)}}, \quad M_{n,b,\ell} = \Lambda_{n,j,k} M_{n,b,\ell}^{(0)}.$$

Then

$$\sum_{b,\ell} p_{n,b,\ell} e^{-R_{n,b}} M_{n,b,\ell} = e^{-qh}, \quad (28)$$

so that $\mathbb{E}_n^h[e^{-R_{n,n+1}^h} F_{n+1}] = (F_n + D_n)e^{-qh}$ exactly.

3.3 Backward recursion and adaptive fund representation

Let $\omega = (b, \ell)$ index a positive-probability hybrid branch, and set $p_{n,\omega} = p_{n,b,\ell}$, $R_{n,\omega} = R_{n,b}$, $M_{n,\omega} = M_{n,b,\ell}$, and $(j_\omega, k_\omega) = (j_b, k_b)$. The set of such branches from node (j, k) is denoted by $\Omega_n(j, k)$.

At each joint factor node (v_j, x_k) , the value function is represented by a sorted set of fund knots and a piecewise polynomial. Candidate knots are generated by inverse images of child knots under the affine branch maps $F \mapsto (F + D_n)M_{n,\omega}$, together with payoff kinks, continuation–surrender intersections, and domain endpoints.

On the last numerical step of policy year i , characterized by $t_{n+1} = i + 1$, define the mortality-adjusted child value by

$$U_{n+1}^{\text{mort}}(F, v, x; P) = (1 - q_{a_0+i})U_{n+1}(F, v, x; P) + q_{a_0+i}B_{i+1}(F). \quad (29)$$

On all other numerical steps, set $U_{n+1}^{\text{mort}} = U_{n+1}$.

The continuation operator is

$$\begin{aligned} C_n(F, v_j, x_k; P) = & -P \mathbf{1}_{\{t_n \in \{0, \dots, T-1\}\}} \\ & + \sum_{\omega \in \Omega_n(j,k)} p_{n,\omega} e^{-R_{n,\omega}} U_{n+1}^{\text{mort}}((F + D_n)M_{n,\omega}, v_{j_\omega}, x_{k_\omega}; P). \end{aligned} \quad (30)$$

On each retained fund interval $[a, b]$, define the four local interpolation points

$$f_0 = a, \quad f_1 = a + \left(1 - \frac{1}{\sqrt{2}}\right)(b - a), \quad f_2 = a + \frac{1}{\sqrt{2}}(b - a), \quad f_3 = b. \quad (31)$$

The production scheme represents the value function on $[a, b]$ by the unique cubic interpolant through f_0, f_1, f_2, f_3 . These symmetric locations minimize $\max_{0 \leq t \leq 1} |t(t - \alpha)(t - \beta)(t - 1)|$ within the symmetric two-point family, and hence minimize the standard fourth-order interpolation error bound on a smooth interval of given width.

At surrender dates, continuation–payoff intersections are inserted as knots before applying (9). Exercise intervals are then compressed exactly to their endpoints. The terminal payoff kink and affine right tail are retained, so the fund cap limits only the representation domain and does not clip values.

Remark 3.3. *The resulting interpolating cubic is not, in general, the best cubic approximation to the value function under the local error criterion used for pruning. Its coefficients are, however, obtained directly from four function values, without solving a local optimization problem. Since this construction is repeated across fund intervals, factor states, and time steps, its simplicity provides a favorable accuracy–cost trade-off for the overall algorithm.*

3.4 Certified pruning, Delta control, and complexity

Backward propagation maps each retained child knot through every hybrid branch, so the number of candidate singular points may grow exponentially with the number of time steps in the worst case. The production scheme controls this growth by applying a certified Douglas–Peucker-type pruning [12] to the piecewise-cubic value functions.

On each candidate fund interval $[a, b]$, let $U(F)$ denote the current piecewise-cubic value function and let $\tilde{U}(F)$ denote its pruned cubic approximation. Since $U - \tilde{U}$ is cubic on each common subinterval, its maximum absolute deviation is obtained by checking the endpoints and all interior stationary points, which are found by solving a quadratic equation. If the prescribed tolerance is violated, the interval is split at a point where this maximum is attained and the test is repeated recursively. The retained approximation therefore satisfies, for every $F \in [a, b]$,

$$|U(F) - \tilde{U}(F)| \leq \varepsilon_{\text{abs}} + \varepsilon_{\text{rel}} F, \quad (32)$$

where ε_{abs} and ε_{rel} are the prescribed absolute and relative pruning tolerances, respectively. A conservative interval envelope based on the minimum fund level over each common subinterval guarantees this mixed tolerance continuously. Optional probability-weighted, preimage, and emergency safeguards are not used in the production results.

Proposition 3.4. *On a common-refinement cell $I = [a, b]$ containing no preserved singularity,*

$$\|\partial_F U - \partial_F \tilde{U}\|_{\infty, I} \leq \frac{18}{b-a} \|U - \tilde{U}\|_{\infty, I}. \quad (33)$$

Proof. The difference $U - \tilde{U}$ is a polynomial of degree at most three on I . Rescaling I to $[-1, 1]$ and applying Markov’s inequality (see [27]) yields the factor $2 \times 3^2 = 18$. $\square$

Thus, away from preserved singularities, value-function pruning also controls the local Delta error, with the bound scaling inversely with the interval width. End-to-end Delta accuracy is tested numerically in Section 5.4.

At time level n , each marginal factor lattice has $O(n)$ nodes and the joint grid has $O(n^2)$ node pairs. If $N_{n,j,k}^{\text{knot}}$ knots are retained at node (j, k) , permanent memory at that level is proportional to $\sum_{j,k} N_{n,j,k}^{\text{knot}}$. The main temporary cost arises from merging the branch preimages. The implementation uses a streaming k -way merge and OpenMP parallelism across joint factor nodes.

4 Convergence, Delta consistency, and Richardson extrapolation

The results are organized in six steps: (i) local consistency and identification of the limiting generator; (ii) weak convergence of the strict binomial and adaptive-trinomial chains, with convergence of European and finite-date Bermudan values; (iii) convergence of the projected binomial fallback and stability of the contractual recursion; (iv) convergence of the adaptive singular-point representation, including pruning and fair-premium determination; (v) under stronger smoothness assumptions, first-order weak accuracy for the strict, unpruned binomial scheme and regular-region Delta consistency; and (vi) the stronger asymptotic expansion required

for Richardson extrapolation. No global first-order rate is claimed for the production adaptive-trinomial scheme. The surrender feature studied here leads to a finite-date Bermudan stopping problem, whereas continuous-exercise American stopping is outside the present theoretical scope.

Between contractual dates set $Y_t = \log F_t$ and $\mathbf{Z}_t = (Y_t, V_t, X_t)$. The unpruned strict-chain value at time t_n is U_n^h , and its adaptive representation is $\tilde{U}_n^h$. For a scalar valuation problem, $\mathcal{V}$ denotes the exact value and $\mathcal{V}_h$ the unpruned time-discrete value. On a regular fund interval, Delta denotes the fund derivative $\partial_F U$; at a preserved kink, left and right derivatives are considered separately.

Unless stated otherwise, all local $O(\cdot)$ and $o(\cdot)$ estimates are uniform on compact subsets of $\mathbb{R} \times (0, \infty) \times (0, \infty)$. For the rate statements, the market curve is assumed sufficiently smooth and the calibrated shift increments satisfy

$$s_{n,h} = \int_{t_n}^{t_{n+1}} \varphi(u) du + O(h^2) \quad (34)$$

uniformly on the time grid.

When the martingale correction is applied, its parent-node rescaling factor $\Lambda_{n,j,k}$ satisfies $\Lambda_{n,j,k} = 1 + o(h)$ under the local-consistency assumptions, so it does not change the limiting drift or covariance. For the binomial weak-order and Richardson results, the stronger expansion $\Lambda_{n,j,k} = 1 + O(h^2)$ is assumed and follows from the corresponding integer-power one-step expansion. The stronger rate statements impose the additional smoothness and operator expansions given below.

All results concern a sequence of grids with $h = T/N \downarrow 0$ whose contractual dates belong to every grid. Unless stated otherwise, the financial chain is strict, unprojected, and unpruned; projection and adaptive representation are introduced only after convergence of the strict chain has been established. Selected technical proofs are deferred to Appendix A.

Throughout this section, $O(h^\alpha)$ denotes a quantity whose absolute value is bounded by Ch^α for all sufficiently small h , with a constant C independent of h . When the estimate is stated on a compact set, the bound is understood to hold uniformly on that set.

4.1 Local consistency and the limiting generator

Let (Y_n^h, V_n^h, X_n^h) be the chain generated by the two factor lattices and the conditional fund quadrature. Uniformly on compact subsets of $\mathbb{R} \times (0, \infty) \times (0, \infty)$, the marginal moment properties of the two lattice constructions, together with strict covariance matching, give

$$\mathbb{E}[\Delta V_n^h \mid v, x] = \kappa_V(\theta_V - v)h + o(h), \quad \mathbb{E}[(\Delta V_n^h)^2 \mid v, x] = \sigma_V^2 v h + o(h), \quad (35)$$

$$\mathbb{E}[\Delta X_n^h \mid v, x] = \kappa_r(\theta_r - x)h + o(h), \quad \mathbb{E}[(\Delta X_n^h)^2 \mid v, x] = \sigma_r^2 x h + o(h), \quad (36)$$

$$\mathbb{E}[\Delta V_n^h \Delta X_n^h \mid v, x] = \rho_{Vr} \sigma_V \sigma_r \sqrt{vx} h + o(h). \quad (37)$$

The equity decomposition in (26), the symmetric residual quadrature, and the local martingale correction yield

$$\mathbb{E}[\Delta Y_n^h \mid y, v, x] = \left(x + \varphi(t_n) - q - \frac{1}{2}v \right) h + o(h), \quad (38)$$

$$\mathbb{E}[(\Delta Y_n^h)^2 \mid y, v, x] = vh + o(h), \quad (39)$$

$$\mathbb{E}[\Delta Y_n^h \Delta V_n^h \mid y, v, x] = \rho_{SV} \sigma_V v h + o(h), \quad (40)$$

$$\mathbb{E}[\Delta Y_n^h \Delta X_n^h \mid y, v, x] = \rho_{Sr} \sigma_r \sqrt{vx} h + o(h). \quad (41)$$

Consequently, the discrete generator converges to

$$\begin{aligned} \mathcal{L}_t \psi = & \left(x + \varphi(t) - q - \frac{1}{2}v \right) \partial_y \psi + \kappa_V (\theta_V - v) \partial_v \psi + \kappa_r (\theta_r - x) \partial_x \psi \\ & + \frac{1}{2}v \partial_{yy} \psi + \frac{1}{2}\sigma_V^2 v \partial_{vv} \psi + \frac{1}{2}\sigma_r^2 x \partial_{xx} \psi \\ & + \rho_{SV} \sigma_V v \partial_{yv} \psi + \rho_{Sr} \sigma_r \sqrt{vx} \partial_{yx} \psi + \rho_{Vr} \sigma_V \sigma_r \sqrt{vx} \partial_{vx} \psi. \end{aligned} \quad (42)$$

The discounted pricing generator is $\mathcal{A}_t \psi = \mathcal{L}_t \psi - [x + \varphi(t)]\psi$. The vanishing-jump property follows because every active successor lies $O(\sqrt{h})$ from its parent on an interior compact set.

4.2 Weak convergence of the strict chains and contract values

The local consistency relations above, together with tightness and vanishing jumps, identify the diffusion limit. The same argument also gives convergence of discounted contract values when the payoff family is uniformly integrable.

Theorem 4.1. *Assume local consistency in (35)–(41), vanishing jumps, compact containment, uniform polynomial moment bounds, and uniqueness of the martingale problem for the Heston–CIR++ diffusion. Then both the strict binomial and the strict adaptive-trinomial chains converge weakly to the diffusion on every finite horizon. If discounted payoffs have linear growth and are uniformly integrable, the corresponding European and finite-date Bermudan values converge.*

Proof. The first two conditional moments converge locally to the drift and covariance of (42), and all jumps vanish uniformly on interior compact sets. Compact containment prevents loss of mass at the localization boundary. The Markov-chain approximation theorem of Kushner and Dupuis [22] therefore identifies every weak subsequential limit with a solution of the Heston–CIR++ martingale problem; uniqueness yields convergence of the full sequence. Uniform integrability allows discounted expectations of linear-growth payoffs to pass to the limit. A finite collection of contractual and exercise dates is handled by backward induction through the continuous event maps. $\square$

The European case corresponds to valuation without early exercise. With annual surrender dates, the contract instead gives a finite-date Bermudan stopping problem, which is covered by the final statement of Theorem 4.1.

4.3 Projected binomial chain and stability of the contractual recursion

For the binomial marginal defined in Section 3.1.1, the exact-mean interpolation probability satisfies, uniformly on interior compact sets,

$$p_{Z,h}(z) = \frac{1}{2} + \beta_Z(z) \sqrt{h} + O(h^{3/2}), \quad \beta_Z(z) = \frac{\kappa_Z(\theta_Z - z) - \sigma_Z^2/4}{2\sigma_Z \sqrt{z}}. \quad (43)$$

For marginal up probabilities p_V and p_X , the attainable correlation interval of the 2×2 joint table has endpoints

$$\begin{aligned}\rho_+(p_V, p_X) &= \exp\{-\tfrac{1}{2}|\ell(p_V) - \ell(p_X)|\}, \\ \rho_-(p_V, p_X) &= -\exp\{-\tfrac{1}{2}|\ell(p_V) + \ell(p_X)|\}, \quad \ell(u) = \log \frac{u}{1-u}.\end{aligned}\tag{44}$$

Consequently, on every compact subset of $(0, \infty) \times (0, \infty)$, the corresponding local endpoints satisfy $\rho_-(v, x; h) \rightarrow -1$ and $\rho_+(v, x; h) \rightarrow 1$ uniformly as $h \downarrow 0$.

Proposition 4.2. *Let K be a compact subset of $(0, \infty) \times (0, \infty)$. For every fixed $\rho_{Vr} \in (-1, 1)$, the unprojected 2×2 joint table is strictly admissible on K for all sufficiently small h .*

Proof. See Appendix A.

Let A_h be the set of factor states at which the requested ρ_{Vr} lies outside the local binomial admissibility interval, and define the first projected time by

$$\tau_h = \inf\{t_n \leq T : (V_n^h, X_n^h) \in A_h\}.\tag{45}$$

Proposition 4.3. *Assume the strict Feller conditions $2\kappa_V\theta_V > \sigma_V^2$ and $2\kappa_r\theta_r > \sigma_r^2$. Suppose the two marginal CIR lattices are tight on $[0, T]$ and converge weakly to their CIR limits. Since projection preserves both marginals, the same compact-containment property holds for the projected joint chain. Then*

$$\mathbb{P}(\tau_h \leq T) \longrightarrow 0.\tag{46}$$

Proof. See Appendix A.

Define the time-weighted occupation error

$$\mathcal{M}_h = h \sum_{n:t_n < T} \mathbb{E}\left[|\rho_{Vr,h}^{\text{eff}} - \rho_{Vr}| \mathbf{1}_{A_h}(V_n^h, X_n^h)\right].\tag{47}$$

Since correlations are bounded by one, $\mathcal{M}_h \leq 2T\mathbb{P}(\tau_h \leq T) \rightarrow 0$.

Theorem 4.4. *Assume Proposition 4.3, local consistency of the unprojected chain on interior compact sets, vanishing jumps, uniform moment bounds, and uniqueness of the martingale problem for the Heston–CIR++ diffusion. Assume also that every projected local value $\rho_{Vr,h}^{\text{eff}}$ leaves the residual equity variance in (26) non-negative, so the three-factor transition is well defined. Then the projected binomial chain converges weakly to the same limit as the strict localized chain. If discounted contract payoffs are uniformly integrable, European and finite-date Bermudan values converge as well.*

Proof. See Appendix A.

Because the CIR++ shift can be negative, the discounted continuation operator need not be a contraction in the ordinary supremum norm. Define $\varphi_- := \max_{0 \leq t \leq T} (-\varphi(t))_+$. The branchwise rate integral is the sum of a non-negative CIR contribution and a calibrated shift increment; by (34), $e^{-R_{n,b}} \leq e^{\varphi_- h + O(h^2)}$ uniformly on the grid. Hence one numerical step is stable with factor $1 + O(h)$ and the product of all stability factors is bounded uniformly over the finite horizon.

Contribution translations are isometries, premium updates are affine, the mortality map is a convex combination, and the obstacle is non-expansive:

$$|\max\{f, R\} - \max\{g, R\}| \leq |f - g|. \quad (48)$$

The complete finite-horizon recursion is therefore stable.

4.4 Adaptive representation and fair-premium convergence

For $J \in \{\text{NS}, \text{FB}, \text{GB}, \text{MX}\}$, let $\Phi_J(P)$ and $\Phi_J^h(P)$ denote the exact and numerical initial net values at annual premium P , and let P_J and P_J^h be their respective roots.

Let U_n^h be the unpruned value of the discrete chain and $\tilde{U}_n^h$ its adaptive approximation. Suppose ordinary interpolation and pruning errors at step n are bounded by $\iota_{n,h}$ and $\varepsilon_{n,h}$ on the represented fund interval. Stability gives

$$\|\tilde{U}_0^h - U_0^h\|_\infty \leq C_T \sum_{n=0}^{N-1} (\iota_{n,h} + \varepsilon_{n,h}) + \mathcal{E}_h^{\text{bdry}}, \quad (49)$$

where C_T is independent of h and $\mathcal{E}_h^{\text{bdry}}$ is the remaining factor- and fund-boundary error. For the implemented payoffs, the right tail is exactly affine and stored through its slope; once the cap lies beyond all non-affine singularities it creates no value truncation. On a quasi-uniform time grid, uniform per-step representation error $o(h)$ is a sufficient, though not necessary, condition for the sum in (49) to vanish.

Theorem 4.5. *Assume the strict financial chain is locally consistent, tight, and uniformly integrable; the factor and fund boundary errors vanish; and the right-hand side of (49) tends to zero. Then the adaptive singular-point value converges to the Heston-CIR++ contract value with the prescribed finite collection of premium, mortality, and surrender dates.*

Proof. Local consistency, tightness, and vanishing jumps imply weak convergence of the unpruned chain. Uniform integrability allows discounted expectations of the linear-growth payoffs to pass to the limit. Backward application of the finite collection of stable translations, premium updates, mortality maps, and non-expansive obstacle operators preserves convergence. The remaining difference between the unpruned discrete value and its adaptive representation is bounded by (49), which vanishes by assumption. $\square$

Corollary 4.6. *Under the hypotheses of Theorem 4.4, if the boundary and representation errors in (49) vanish, the projected adaptive binomial scheme converges to the same contract value.*

Define

$${}_i p_{a_0} = \prod_{u=0}^{i-1} (1 - q_{a_0+u}), \quad {}_0 p_{a_0} = 1, \quad (50)$$

as the probability of surviving to anniversary i . Because premiums are paid only while the insured is alive, the no-surrender discrete net value is exactly affine:

$$\Phi_{\text{NS}}^h(P) = \mathcal{B}_h - P A^M, \quad A^M = \sum_{i=0}^{T-1} {}_i p_{a_0} P^M(0, i), \quad P_{\text{NS}}^h = \mathcal{B}_h / A^M. \quad (51)$$

The deterministic survival probabilities and the discrete CIR++ calibration make A^M grid independent. Surrender premiums are obtained by safeguarded Brent–Dekker root finding [5].

Proposition 4.7. *Suppose $\Phi_J^h \rightarrow \Phi_J$ uniformly on a compact premium interval, Φ_J has a unique interior root P_J , and $|\Phi_J'(P)| \geq m_J > 0$ near P_J . Then $P_J^h \rightarrow P_J$ and*

$$|P_J^h - P_J| \leq \frac{\sup_P |\Phi_J^h(P) - \Phi_J(P)|}{m_J} + o\left(\sup_P |\Phi_J^h(P) - \Phi_J(P)|\right). \quad (52)$$

Proof. Uniform convergence implies that, for sufficiently small h , a numerical root lies in the neighborhood where the derivative is bounded away from zero. Since $\Phi_J(P_J) = 0 = \Phi_J^h(P_J^h)$,

$$\begin{aligned} |\Phi_J(P_J^h) - \Phi_J(P_J)| &= |\Phi_J(P_J^h) - \Phi_J^h(P_J^h)| \\ &\leq \sup_P |\Phi_J^h(P) - \Phi_J(P)|. \end{aligned}$$

The mean-value theorem and $|\Phi_J'| \geq m_J$ give the leading bound. The stated remainder follows from the local expansion of the inverse map at the simple root. $\square$

4.5 Weak-error rates

The preceding results establish convergence of the valuation method. A first-order weak-error bound requires additional moment structure and smoothness and is therefore stated separately.

The binomial scheme has the stronger moment structure needed for a first-order rate. Let ΔZ_h^{bin} denote one strict binomial increment.

Lemma 4.8. *Uniformly on interior compact sets,*

$$\mathbb{E}[\Delta Z_{h,i}^{\text{bin}}] = b_i(t, Z)h + O(h^2), \quad (53)$$

$$\mathbb{E}[\Delta Z_{h,i}^{\text{bin}} \Delta Z_{h,j}^{\text{bin}}] = a_{ij}(t, Z)h + O(h^2), \quad (54)$$

$$\mathbb{E}[\Delta Z_{h,i}^{\text{bin}} \Delta Z_{h,j}^{\text{bin}} \Delta Z_{h,k}^{\text{bin}}] = O(h^2), \quad (55)$$

$$\mathbb{E}[|\Delta Z_{h,i}^{\text{bin}} \Delta Z_{h,j}^{\text{bin}} \Delta Z_{h,k}^{\text{bin}} \Delta Z_{h,\ell}^{\text{bin}}|] = O(h^2), \quad (56)$$

and $\mathbb{E}|\Delta Z_h^{\text{bin}}|^5 = O(h^{5/2})$.

Proof. See Appendix A.

Theorem 4.9. *Suppose that the strict binomial chain satisfies the local moment estimates in Lemma 4.8. Assume also uniform polynomial moment bounds, stability of the discounted operators, and a backward value with five state derivatives of polynomial growth between contractual dates. If the finitely many deterministic event maps preserve this regularity, then*

$$|\mathcal{V}_h^{\text{bin}} - \mathcal{V}| \leq C_T h = \frac{C_T T}{N}. \quad (57)$$

Proof. See Appendix A.

The proof is based on the one-step expansion, valid for smooth test functions,

$$Q_{t,h}^{\text{bin}} f = f + h \mathcal{A}_t f + O(h^2). \quad (58)$$

Its derivation from Lemma 4.8 is given in Appendix A.

For an adaptive trinomial with $\lambda > 1$, the limiting symmetric marginal probabilities in Lemma 3.1 imply the same $O(\sqrt{h})$ third standardized moments, so the preceding first-order argument applies on localized regions where all selected stencils are eventually symmetric. The production variance grid uses $\lambda_V = 1$ and requires a separate qualification.

Proposition 4.10. *At an interior variance state where $A(v) < 0$ in (23), the limiting standardized support of the selected minimal-span stencil is, up to reflection, $(-2, 0, 1)$ with probabilities $(1/6, 1/2, 1/3)$. Its third standardized moment is -1 (or $+1$ after reflection). Consequently, the corresponding raw third moment of the variance increment is generically $O(h^{3/2})$, not $O(h^2)$.*

Proof. The limiting support and probabilities follow from Lemma 3.1. Direct calculation gives

$$\frac{1}{6}(-2)^3 + \frac{1}{2}0^3 + \frac{1}{3}1^3 = -1.$$

Multiplication by the cube of the $O(\sqrt{h})$ local scale gives an $O(h^{3/2})$ raw third moment. $\square$

Corollary 4.11. *The first-order weak bound in Theorem 4.9 extends to an adaptive-trinomial refinement only under an eventual symmetric-stencil condition, for example on a localized region where $A_V(v) \geq c_A > 0$ for some constant c_A , where A_V denotes the function in (23) evaluated with $(\kappa_V, \theta_V, \sigma_V)$, or when both grid multipliers exceed one. For the production choice $\lambda_V = 1$, strict weak convergence follows from Theorem 4.1, but the moment argument above does not establish a global first-order rate.*

Remark 4.12. *The one-sided-stencil issue is not a negligible tail event for the baseline variance process; it is a structural feature of the $\lambda_V = 1$ grid. We therefore report direct trinomial refinements without fitting or claiming a formal convergence order. The numerical sequence is used as an accurate strict approximation, while the binomial chain supplies the proved first-order/Richardson cross-check.*

4.6 Delta consistency on regular fund regions

On a compact regular fund region define $\Delta_n^h = \partial_F U_n^h$ and $\tilde{\Delta}_n^h = \partial_F \tilde{U}_n^h$. A regular region does not cross a preserved payoff kink or exercise boundary; at such singularities the statements apply to one-sided derivatives.

Proposition 3.4 controls the derivative error created by one cubic pruning operation. The following statements make explicit what this implies for a refinement sequence.

Corollary 4.13. *At a fixed pruning operation, suppose the source partition is fixed on a compact regular fund interval K and has minimum cell length $m_K > 0$. If $\|U - \tilde{U}\|_{\infty, K} \leq \varepsilon$, then*

$$\|\partial_F U - \partial_F \tilde{U}\|_{\infty, K} \leq \frac{18\varepsilon}{m_K}. \quad (59)$$

Hence the pruned Delta converges uniformly to the source-curve Delta as $\varepsilon \downarrow 0$.

Proof. Apply Proposition 3.4 on every cell of the common refinement and use $h_I \geq m_K$. $\square$

Theorem 4.14. *Let K be a compact interval of initial fund values. Assume the hypotheses of Theorem 4.9 and, in addition, that the exact backward solution and the strict binomial backward values are six-times continuously differentiable on an open tube containing every branch image generated from K . At each contractual date, assume that this tube stays a positive distance from payoff and exercise singularities, so the active branch of every event map is fixed there. If the one-step weak defect in (58) and the operator-stability estimate hold in the C^1 norm on that tube, then*

$$\|\partial_F U_0^{h,\text{bin}} - \partial_F U_0\|_{\infty,K} \leq C_K h. \quad (60)$$

Proof. See Appendix A.

Theorem 4.15. *Let K be a compact fund interval at the initial factor state that stays a positive distance from every preserved payoff, guarantee, or exercise singularity. Let U_0^h and $\tilde{U}_0^h$ be the unpruned and adaptive initial-node curves. Assume*

- (D1) $U_0^h \rightarrow U_0$ in $C^1(K)$; for the strict binomial chain this follows from Theorem 4.14 under its regular-tube hypotheses;
- (D2) on every cell I of the common refinement of U_0^h and $\tilde{U}_0^h$, with length h_I , the final representation error $E_h(I) = \|U_0^h - \tilde{U}_0^h\|_{\infty,I}$ satisfies

$$\max_{I \subset K} \frac{E_h(I)}{h_I} \rightarrow 0. \quad (61)$$

Then

$$\|\partial_F \tilde{U}_0^h - \partial_F U_0\|_{\infty,K} \rightarrow 0. \quad (62)$$

At a preserved singularity, the same conclusion applies separately to the left and right derivatives on adjacent regular intervals.

Proof. By the triangle inequality,

$$\|\partial_F \tilde{U}_0^h - \partial_F U_0\|_{\infty,K} \leq \|\partial_F \tilde{U}_0^h - \partial_F U_0^h\|_{\infty,K} + \|\partial_F U_0^h - \partial_F U_0\|_{\infty,K}.$$

The second term tends to zero by (D1). On each common-refinement cell, Proposition 3.4 bounds the first term by $18E_h(I)/h_I$; taking the maximum and using (D2) proves (62). $\square$

Remark 4.16. *Condition (D1) is not implied by value convergence alone. Theorem 4.14 supplies it for the strict binomial chain on regular tubes; for the production adaptive-trinomial sequence it remains an explicit regularity condition rather than a claimed rate theorem. Condition (D2) is a joint time-space requirement: a fixed value tolerance is not sufficient if the smallest relevant fund cell also shrinks. The stability bound (49) supplies a computable sufficient upper bound for the numerator in (D2). The numerical tests in Section 5.4 assess the complete recursion rather than only one pruning operation.*

4.7 Richardson extrapolation for the binomial scheme

Weak order one is not enough to justify Richardson extrapolation; a stable asymptotic error expansion is required. We state the assumptions explicitly because they delimit the theoretical

claim.

Let $Q_{t,h}$ be one discounted step of the strict, unpruned binomial chain and $P_{t,t+h}$ the exact discounted semigroup. Assume:

- (R1) the correlation matrix is uniformly positive definite and the localized binomial joint probabilities are strictly admissible for all sufficiently small h ;
- (R2) φ is three-times continuously differentiable and bounded below, and the backward value and its images under the deterministic event maps have state derivatives up to order eight and time derivatives up to order three, all of polynomial growth;
- (R3) the numerical chain and diffusion have uniform polynomial moment bounds, and localization errors are $o(h^2)$ in the initial discounted value;
- (R4) for every test function in this regularity class,

$$Q_{t,h}f = f + h\mathcal{A}_t f + h^2\mathcal{B}_t f + h^3\mathcal{C}_t f + O(h^4), \quad (63)$$

$$P_{t,t+h}f = f + h\mathcal{A}_t f + h^2\mathcal{E}_t f + h^3\mathcal{F}_t f + O(h^4), \quad (64)$$

with polynomially weighted, uniform remainders;

- (R5) the finite set of contractual dates is common to every refined grid and the associated interface maps preserve the stated regularity.

For the binomial construction, the integer-power moment structure required by (R4) is obtained on interior compact sets by expanding $p_{Z,h}$ around $1/2$, inserting the explicit 2×2 coupling, and Taylor expanding the one-step operator. Odd standardized moments are $O(\sqrt{h})$ and even moments have expansions in integer powers of h ; therefore raw third and fifth moments first enter at orders h^2 and h^3 . This is the discrete structure required by the Talay–Tubaro expansion [29].

Lemma 4.17. *Under (R2)–(R4), there are differential operators Ψ_t and Ξ_t such that*

$$(Q_{t,h} - P_{t,t+h})f = h^2\Psi_t f + h^3\Xi_t f + O(h^4) \quad (65)$$

uniformly on the localized state domain.

Proof. Subtract (64) from (63) and set $\Psi_t = \mathcal{B}_t - \mathcal{E}_t$ and $\Xi_t = \mathcal{C}_t - \mathcal{F}_t$. □

Theorem 4.18. *Under (R1)–(R5), the strict, unpruned binomial price of a sufficiently smooth contract satisfies*

$$\mathcal{V}_h = \mathcal{V} + c_1 h + c_2 h^2 + o(h^2), \quad (66)$$

where c_1 and c_2 are independent of the refinement level.

Proof. See Appendix A.

Corollary 4.19. *Under Theorem 4.18,*

$$\mathcal{V}_h^{\text{Rich}} = 2\mathcal{V}_{h/2} - \mathcal{V}_h = \mathcal{V} + O(h^2). \quad (67)$$

More generally, for a refinement ratio $r > 1$,

$$\mathcal{V}_{h,r}^{\text{Rich}} = \frac{r\mathcal{V}_{h/r} - \mathcal{V}_h}{r - 1} = \mathcal{V} + O(h^2). \quad (68)$$

Proof. Insert (66) into either linear combination; the coefficient of $c_1 h$ cancels identically. $\square$

If the net-value function has a uniform expansion $\Phi_h(P) = \Phi(P) + a_1(P)h + a_2(P)h^2 + o(h^2)$ near a simple fair-premium root, the implicit-function theorem gives $P_J^h = P_J + d_1 h + d_2 h^2 + o(h^2)$, so the same extrapolation applies to the premium. For P_{NS} , this follows directly from (51), because A^M is grid independent.

Remark 4.20. *Theorem 4.18 is a theorem for strict, unpruned transitions and sufficiently smooth values. Projection and representation errors must be $o(h)$ to preserve the leading coefficient and $O(h^2)$ to retain the second-order extrapolation. The payoff $\max\{F, G\}$ and a Bermudan obstacle are non-smooth and can generate lattice-alignment oscillations [11]. A full second-order theorem for the exact insurance payoff therefore requires a smoothing or interface argument in addition to (R1)–(R5). The binomial Richardson values reported below should be read as theoretically motivated and numerically validated accelerations, not as an unconditional second-order theorem for the kinked contract. The adaptive-trinomial sequence is not extrapolated because its non-nested triplet selection does not display a stable leading coefficient on the available grids.*

5 Numerical study

5.1 Setup and independent benchmarks

We use SP, MC, and LSMC for singular points, Monte Carlo, and least-squares Monte Carlo. In the diagnostic tables, $N_{\text{max}}^{\text{knot}}$ is the largest number of retained fund knots at a joint factor node, ε_{max} is the largest posterior continuous pruning error, and t_{12} is twelve-thread wall time. Table 1 summarizes the central specification. Unless otherwise stated, contract-level results use the production configuration: strict adaptive trinomial factor lattices, 24 steps per year ($N = 480$), cubic fund functions, and $(\varepsilon_{\text{abs}}, \varepsilon_{\text{rel}}) = (10^{-4}, 10^{-6})$. Mortality uses the 2025 ISTAT resident-population table at issue age 50.

Block	Parameters	Values
Contract	T, D, g_m, g_s, q	20, 100, 0.02, 0.02, 0
Heston variance	$V_0, \kappa_V, \theta_V, \sigma_V$	0.04, 2.00, 0.04, 0.30
CIR rate factor	$X_0, \kappa_r, \theta_r, \sigma_r$	0.018, 0.45, 0.025, 0.10
Correlations	$\rho_{SV}, \rho_{Sr}, \rho_{Vr}$	−0.70, −0.20, 0.02
Factor grid	type; (λ_V, λ_X)	adaptive trinomial; (1.0, 1.4)
Fund representation	anchors; tolerances	cubic; $10^{-4}, 10^{-6}$
Production grid	N_{yr}, N	24, 480
Surrender	dates; α_i^s	annual; 1

Table 1. Central contract, model, and numerical parameters. The initial curve is the ECB euro-area AAA government-bond Svensson curve dated 15 July 2026.

The financial engine reproduces all grid-maturity discount factors and the corrected discounted-fund martingale condition to roundoff. The production trinomial transitions match the two CIR moments and all three correlations to roundoff and remain strictly admissible over the complete refinement range. Detailed factor-grid, admissibility, curve-fit, martingale, and refinement diagnostics are reported in the Supplementary Material.

The independent no-surrender benchmark is antithetic full-truncation Euler Monte Carlo with exact CIR++ shift increments and analytic control variates. One hundred replications of one million paths at 256 time steps per year give $P_{\text{MC}} = 111.6586729$ with standard error 0.0012648. All deterministic timings use 12 OpenMP threads on an Intel Core Ultra 5 125H notebook with 32 GB RAM; absolute times are machine-specific.

For validation, we use independent Monte Carlo for the no-surrender benchmark and cross-fitted least-squares Monte Carlo for the surrender products. These approaches provide external reference values based on numerical procedures that are structurally different from the proposed deterministic method, and therefore offer a useful check on its accuracy without relying on the same discretization or approximation mechanisms.

5.2 Convergence and cost–accuracy

Table 2 reports representative levels of the common refinement sequence. It shows the slow monotone approach of the direct binomial values, the substantial cancellation obtained by Richardson, and the higher coarse-grid accuracy of the adaptive trinomial scheme.

N_{yr}	N	Binomial direct	Binomial Richardson	Trinomial direct
4	80	110.748278	111.157080	111.594769
8	160	111.140506	111.532734	111.584134
16	320	111.374914	111.609323	111.640138
24	480	111.465744	111.636463	111.644323
32	640	111.512365	111.649816	111.653842

Table 2. Representative temporal refinements for the mortality-extended no-surrender premium. The independent Monte Carlo benchmark is 111.6586729. Richardson uses the corresponding half-resolution binomial run.

Figure 1 compares the direct binomial, binomial Richardson, and direct adaptive-trinomial sequences. Direct binomial values approach the Monte Carlo benchmark from below. At $N = 640$, Richardson reduces the absolute discrepancy to 0.008857 in 116.2 seconds, while the direct trinomial value has error 0.004830 in 191.3 seconds. The trinomial representation stays between 176 and 187 knots at every displayed refinement. Thus Richardson is competitive at intermediate cost, whereas the adaptive trinomial method gives the most accurate direct value. Because the $\lambda_V = 1$ minimal-span rule can select one-sided stencils, these direct trinomial refinements are used as a convergent accuracy sequence without fitting or claiming a formal weak order.




Figure 1. Cost-accuracy comparison for the mortality-extended no-surrender premium. Richardson cost includes both required grids; dotted step segments are best-so-far error envelopes.

5.3 Pruning sensitivity and selected tolerance

Table 3 varies both coefficients of the mixed certificate at $N_{\text{yr}} = 24$ while keeping every other setting fixed. The comparison uses the exact interval-wise posterior certificate and disables all probability-dependent compression policies. Within each block, the tightest row is used only as an internal representation reference.

Panel A: binomial scheme					Panel B: trinomial scheme				
ε	P	$ P - P_{\text{tight}} $	$N_{\text{max}}^{\text{knot}}$		ε	P	$ P - P_{\text{tight}} $	$N_{\text{max}}^{\text{knot}}$	
$\varepsilon_{\text{abs}}; \varepsilon_{\text{rel}} = 10^{-6}$					$\varepsilon_{\text{abs}}; \varepsilon_{\text{rel}} = 10^{-6}$				
$3 \cdot 10^{-4}$	111.465926	$2.155 \cdot 10^{-4}$	206		$3 \cdot 10^{-4}$	111.644490	$2.296 \cdot 10^{-4}$	181	
10^{-4}	111.465754	$4.395 \cdot 10^{-5}$	215		10^{-4}	111.644323	$6.213 \cdot 10^{-5}$	187	
$3 \cdot 10^{-5}$	111.465728	$1.786 \cdot 10^{-5}$	218		$3 \cdot 10^{-5}$	111.644267	$6.592 \cdot 10^{-6}$	188	
10^{-5}	111.465710	0	220		10^{-5}	111.644261	0	190	
$\varepsilon_{\text{rel}}; \varepsilon_{\text{abs}} = 10^{-4}$					$\varepsilon_{\text{rel}}; \varepsilon_{\text{abs}} = 10^{-4}$				
$3 \cdot 10^{-6}$	111.467447	$2.215 \cdot 10^{-3}$	132		$3 \cdot 10^{-6}$	111.645861	$2.063 \cdot 10^{-3}$	121	
10^{-6}	111.465754	$5.222 \cdot 10^{-4}$	215		10^{-6}	111.644323	$5.249 \cdot 10^{-4}$	187	
$3 \cdot 10^{-7}$	111.465345	$1.126 \cdot 10^{-4}$	373		$3 \cdot 10^{-7}$	111.643913	$1.152 \cdot 10^{-4}$	320	
10^{-7}	111.465232	0	616		10^{-7}	111.643798	0	489	

Table 3. Pruning sensitivity at $N_{\text{yr}} = 24$ ($N = 480$). The upper block in each panel varies ε_{abs} and the lower block varies ε_{rel} . P_{tight} is the tightest result within the corresponding block.

Tightening ε_{abs} from 10^{-4} to 10^{-5} changes the premium by 4.40×10^{-5} in the binomial scheme and 6.21×10^{-5} in the trinomial scheme. The relative coefficient has a larger effect on complexity because its contribution grows with $|F|$. Relative to 10^{-7} , the production value $\varepsilon_{\text{rel}} = 10^{-6}$ changes the premium by about 5.2×10^{-4} in either scheme while keeping $N_{\text{max}}^{\text{knot}}$ at 215 and 187; the 10^{-7} calculations require 616 and 489 knots. A separate fund-cap check over $10^8 D$ – $10^{12} D$ has spread 4.21×10^{-6} . These comparisons motivate the production setting $(\varepsilon_{\text{abs}}, \varepsilon_{\text{rel}}) = (10^{-4}, 10^{-6})$.

5.4 Delta consistency and independent no-surrender benchmark

Table 4 combines three tests. Panel A isolates one local pruning operation on the conditional policy-year-10 value. Panel B adds temporal refinement and compares the analytic cubic Delta with an independent pathwise Monte Carlo benchmark on $\mathcal{F} = \{500, 1000, 1500, 2000\}$. These four fund values span a lower-fund region, the representative policy-year-10 state used in the hedging experiment, and progressively more fund-dominated regions. Panel C reports the inception Delta immediately after the first contribution. For Panel A, $E_U = \max_{F \in \mathcal{F}} |U^\varepsilon(F) - U^0(F)|$ and $E_\Delta = \max_{F \in \mathcal{F}} |\Delta^\varepsilon(F) - \Delta^0(F)|$. The Monte Carlo benchmarks use 1.6 million antithetic paths and 256 full-truncation Euler steps per year; for Panel C, $\Delta_{0,\text{MC}} = 0.810044207$ with standard error 1.70×10^{-4} .

<i>Panel A: local pruning at policy year 10</i>				<i>Panel B: policy-year-10 temporal convergence</i>			
ε_{abs}	E_U	E_Δ	knots	N_{yr}	Binomial	Richardson	Trinomial
10^{-3}	$1.295 \cdot 10^{-4}$	$1.508 \cdot 10^{-5}$	18	4	$1.682 \cdot 10^{-2}$	$1.005 \cdot 10^{-2}$	$2.076 \cdot 10^{-3}$
10^{-4}	$3.536 \cdot 10^{-5}$	$1.162 \cdot 10^{-6}$	26	8	$9.367 \cdot 10^{-3}$	$1.918 \cdot 10^{-3}$	$1.165 \cdot 10^{-3}$
10^{-5}	$2.780 \cdot 10^{-6}$	$1.675 \cdot 10^{-6}$	52	16	$4.879 \cdot 10^{-3}$	$3.900 \cdot 10^{-4}$	$5.674 \cdot 10^{-4}$
				24	$3.135 \cdot 10^{-3}$	$1.377 \cdot 10^{-4}$	$4.788 \cdot 10^{-4}$
				32	$2.230 \cdot 10^{-3}$	$4.187 \cdot 10^{-4}$	$6.682 \cdot 10^{-4}$

<i>Panel C: inception Delta after the first contribution</i>						
N_{yr}	Binomial		Richardson		Trinomial	
	Δ_0	e_0	Δ_0	e_0	Δ_0	e_0
4	0.808856	$1.188 \cdot 10^{-3}$	0.807713	$2.332 \cdot 10^{-3}$	0.811659	$1.615 \cdot 10^{-3}$
8	0.809175	$8.688 \cdot 10^{-4}$	0.809495	$5.493 \cdot 10^{-4}$	0.810985	$9.412 \cdot 10^{-4}$
16	0.809589	$4.552 \cdot 10^{-4}$	0.810003	$4.166 \cdot 10^{-5}$	0.810557	$5.132 \cdot 10^{-4}$
24	0.809775	$2.693 \cdot 10^{-4}$	0.810120	$7.615 \cdot 10^{-5}$	0.810507	$4.630 \cdot 10^{-4}$
32	0.809888	$1.566 \cdot 10^{-4}$	0.810186	$1.420 \cdot 10^{-4}$	0.810460	$4.162 \cdot 10^{-4}$

Table 4. Delta consistency of the certified cubic no-surrender engine. Panel A reports the local value and Delta errors relative to the unpruned source curve. Panel B reports $E_\Delta = \max_{F \in \mathcal{F}} |\Delta(F) - \Delta_{\text{MC}}(F)|$. Panel C reports $e_0 = |\Delta_0 - \Delta_{0,\text{MC}}|$; the 95% Monte Carlo half-width at inception is 0.000333.

Panel A confirms that a value-only certificate preserves the local derivative: all displayed Delta errors are below 1.6×10^{-5} . In Panel B, the direct binomial error decreases steadily and Richardson reduces the maximum discrepancy to 1.38×10^{-4} at $N_{\text{yr}} = 24$. The mild non-monotonicity on the finest Richardson and trinomial levels is comparable to the Monte Carlo sampling uncertainty and does not include Monte Carlo time bias. Panel C gives the same conclusion at the initial hedging point: Richardson lies within one Monte Carlo standard error from $N_{\text{yr}} = 16$ onward, while the trinomial discrepancy decreases monotonically to 4.16×10^{-4} .

5.5 Correlation sensitivity and local admissibility

The fully correlated specification is numerically useful only if the local coupling remains strict over economically relevant parameter changes. Figure 2 varies one correlation at a time around the baseline $(\rho_{SV}, \rho_{Sr}, \rho_{Vr}) = (-0.70, -0.20, 0.02)$ under the production configuration. The ordinate is the premium change relative to $P_{\text{NS}}^{\text{base}} = 111.644322965$.

The dependence is close to linear over the displayed ranges. Least-squares slopes are approxi-




Figure 2. Sensitivity of the no-surrender fair premium to the three pairwise correlations. In each panel, the displayed correlation varies while the other two remain at their baseline values.

mately 0.348, 2.166, and -0.175 with respect to ρ_{SV} , ρ_{Sr} , and ρ_{Vr} , respectively. Thus a change of 0.1 in the equity-rate correlation changes the annual premium by about 0.217, compared with 0.035 for the equity-variance correlation and -0.017 for the variance-rate correlation. The variance-rate sweep is restricted to $[-0.13, 0.20]$, well inside the common strict range in Proposition 3.2; no point in the sweep uses local projection. Direct binomial and binomial Richardson sweeps give the same qualitative ordering.

5.6 Surrender contracts and independent LSMC validation

For $J \in \{\text{FB}, \text{GB}, \text{MX}\}$, define $H_J = P_J - P_{\text{NS}}$ as the surrender-option component of the fair annual premium. Table 5 summarizes the contract-level comparison. The surrender-option components are considerably more stable than the premium levels: across the five finest trinomial grids they lie in $H_{\text{FB}} \in [0.2201, 0.2209]$, $H_{\text{GB}} \in [5.7745, 5.7792]$, and $H_{\text{MX}} \in [6.4243, 6.4287]$.

<i>Panel A: selected deterministic valuations</i>				
Method	P_{NS}	P_{FB}	P_{GB}	P_{MX}
Binomial Richardson, $N_{\text{yr}} = 32$	111.6499	111.8719	117.4202	118.0735
Trinomial direct, $N_{\text{yr}} = 24$	111.6443	111.8644	117.4207	118.0693
Trinomial direct, $N_{\text{yr}} = 32$	111.6538	111.8744	117.4331	118.0826
<i>Surrender-option components</i>				
Binomial Richardson, $N_{\text{yr}} = 32$	–	0.2220	5.7703	6.4236
Trinomial direct, $N_{\text{yr}} = 24$	–	0.2201	5.7763	6.4250
Trinomial direct, $N_{\text{yr}} = 32$	–	0.2206	5.7792	6.4287
<i>Panel B: independent LSMC validation of surrender premiums</i>				
Product	LSMC-256	Held-out 95% CI	SP production	SP-LSMC
P_{FB}	111.7090	[111.6279, 111.7902]	111.8644	0.1554
P_{GB}	117.3181	[117.1971, 117.4390]	117.4207	0.1026
P_{MX}	117.7707	[117.6766, 117.8647]	118.0693	0.2986

Table 5. Fair annual premiums and surrender-option components. Panel B is an independent two-fold cross-fitted validation using the guarantee-kink quadratic basis, one million paths, and 256 time steps per policy year. Its intervals quantify held-out sampling uncertainty only; regression and time-discretization bias are not included.

Every deterministic refinement gives $P_{MX} > P_{GB} > P_{FB} > P_{NS}$. The $N_{yr} = 32$ binomial Richardson components closely match the direct trinomial values, providing an internal deterministic cross-check. The cross-fitted LSMC premiums are lower by 0.1554, 0.1026, and 0.2986 for P_{FB}, P_{GB}, P_{MX} , respectively; the largest relative discrepancy is 0.253%. This is consistent with a fitted, generally suboptimal stopping policy. In the associated simulation control, the no-surrender estimate is 111.6789 with held-out interval $[111.5842, 111.7736]$, containing both the production value and the high-precision Monte Carlo benchmark.

Figure 3 shows policy-year-10 sections of the obstacle problem. Fund-based surrender occurs only in the high-fund tail, guarantee-based surrender in the low-fund region, and the mixed payoff on both sides of a central continuation band. Higher variance generally expands continuation, while higher rates favor exercise.




Figure 3. Continuation and surrendered regions at policy year 10 under the production trinomial scheme. Pale blue denotes continuation, dotted orange immediate surrender, and black curves the interpolated boundaries.

5.7 Issue-age and mortality robustness

The production configuration is also stable across materially different mortality levels. Table 6 compares issue ages 40, 50, and 60 with a no-mortality benchmark while holding the financial model and numerical discretization fixed.

The base no-surrender premium is only mildly affected, whereas mortality has a more visible effect on the guarantee-based surrender components. From the no-mortality case to issue age 60, H_{GB} and H_{MX} decrease by about 5.6% and 6.2%, respectively, because death shortens the horizon

Panel A: fair annual premiums

Scenario	20-year survival	P_{NS}	P_{FB}	P_{GB}	P_{MX}
Issue age 40	96.0843%	111.6528	111.8774	117.5034	118.1701
Issue age 50	90.4215%	111.6443	111.8644	117.4207	118.0693
Issue age 60	75.6815%	111.6206	111.8287	117.1937	117.7952
No mortality	100.0000%	111.6588	111.8864	117.5613	118.2410

Panel B: surrender-option components

Scenario	H_{FB}	H_{GB}	H_{MX}
Issue age 40	0.2246	5.8506	6.5173
Issue age 50	0.2201	5.7763	6.4250
Issue age 60	0.2081	5.5730	6.1745
No mortality	0.2276	5.9025	6.5822

Table 6. Issue-age and mortality sensitivity under the production adaptive-trinomial discretization. The no-mortality row disables both mortality termination and the death benefit.

over which the surrender guarantees can be exercised. The monotone age pattern confirms that the central results are not specific to the issue-age-50 calibration.

5.8 Risk-neutral timing of optimal surrender

The adaptive representation also gives direct access to the deterministic exercise policy. To summarize its dynamic implications, we simulate the production policies under the risk-neutral measure with two million paths and annual exercise only. Figure 4 reports the unconditional probability of first surrender in each policy year; the paths that do not surrender terminate through death or maturity.

Contract	Surrender	Death	Maturity	Mean surrender year
P_{FB}	29.9308%	8.5539%	61.5153%	15.825
P_{GB}	66.7444%	4.7233%	28.5322%	7.871
P_{MX}	89.7464%	3.5006%	6.7530%	9.001

Table 7. Risk-neutral terminal outcomes under the fixed production surrender policies. The largest annual 95% Monte Carlo half-width is 0.0392 percentage points.

Fund-based surrender is comparatively infrequent and late: more than 61% of P_{FB} paths reach maturity, and conditional surrender occurs on average near policy year 16. Guarantee-based surrender is much earlier and more frequent. The mixed payoff is exercised on almost 90% of paths, but its mean surrender year is later than for P_{GB} because the fund branch creates an additional continuation region at intermediate fund levels. No simulated observation falls outside the stored policy domain. These results are descriptive risk-neutral policy diagnostics, not behavioral lapse predictions.




Figure 4. Unconditional risk-neutral distribution of the first surrender year under the production policies. The remaining probability mass terminates through death or maturity as reported in Table 7.

5.9 Model relevance and factor hedging

In this subsection, L is the liability value, C the one-year call value, and B the zero-coupon bond value. Subscripts denote partial derivatives with respect to the indicated state variable, and n_C, n_B, n_S are the call, bond, and stock holdings. For strategy h , $\Pi_h(t)$ is the hedge-portfolio value and $\text{HE}_h(t) = \Pi_h(t) - L(t)$ is its mark-to-model hedging error. Table 8 compares four nested models while holding the contract, mortality, initial discount curve, and numerical engine fixed. Deterministic-rate models follow the same market forward curve, so the comparison is not contaminated by a different initial term structure.

Model	P_{NS}	P_{FB}	P_{GB}	P_{MX}
Black–Scholes, deterministic rates	112.1970	112.6570	116.8521	117.8997
Heston, deterministic rates	111.8767	112.1013	117.3195	117.9572
Black–Scholes–CIR++	111.9231	112.3773	117.0078	118.0790
Heston–CIR++	111.6443	111.8644	117.4207	118.0693

Table 8. Model-component comparison under the production numerical configuration and a common initial discount curve.

Relative to Black–Scholes with deterministic rates, the full model changes $(P_{\text{NS}}, P_{\text{FB}}, P_{\text{GB}}, P_{\text{MX}})$ by approximately $(-0.553, -0.793, +0.569, +0.170)$. Hence model enrichment is payoff dependent rather than a uniform price correction. For the mixed contract, the volatility and rate contributions partly offset.

To assess the risk-management relevance, we form a one-year static hedge for the no-surrender liability at policy year 10. Physical-measure paths are simulated under a stated drift scenario, while all liability values and hedge ratios are computed under $\mathbb{Q}$. We compare a Black–Scholes Delta hedge, a Heston–CIR++ Delta hedge, and a stock–call–bond hedge matching first-order fund, variance, and rate exposures. The positions are held fixed to policy year 11; mortality is treated as diversified.

Replacing the Black–Scholes Delta alone reduces RMSE by about 2%. Matching variance and rate exposures reduces RMSE by 80.4% relative to the model-consistent Delta-only hedge and reduces the magnitude of the left-tail mean by 75.1%. This is a static, frictionless factor-exposure

Strategy	RMSE	Left-tail mean (1%)
Black–Scholes Delta	36.549	-92.222
Heston–CIR++ Delta	35.825	-95.107
Delta + variance + rate	7.019	-23.687

Table 9. One-period mark-to-model hedging errors. The error is hedge value minus gross liability; the last column is the conditional mean below the empirical 1% quantile.

diagnostic, not a complete asset–liability-management study.

6 Conclusions

We proposed an adaptive dynamic-programming method for long-dated annual-premium contracts with stochastic volatility, stochastic interest rates, mortality, and Bermudan surrender. The central state-space reduction is to discretize only the recombining financial factors and retain the accumulated fund as an adaptive value function. Contributions become translations, and the exercise decision is applied directly to the same function. This avoids a global fund grid while preserving a fully deterministic backward recursion.

The numerical construction combines locally covariance-consistent square-root lattices, residual equity quadrature, discrete curve fitting, and a martingale correction. Piecewise-cubic singular-point propagation retains payoff and exercise kinks, exact affine exercise compression removes redundant regions, and continuous pruning certificates control the represented value between knots. Both strict financial chains converge weakly. The strict binomial chain is weakly first-order accurate and admits the asymptotic expansion required for Richardson under the stated smoothness assumptions; the production adaptive trinomial is retained as a direct convergent approximation without a claimed global rate. The adaptive valuation converges when representation and boundary errors vanish, and the cubic certificate yields a regular-region Delta consistency result under an explicit value-error-to-cell-length condition.

The experiments support the method from complementary directions. The adaptive-trinomial scheme gives the most accurate direct no-surrender value on the finest grid, with fewer than 200 retained fund knots at every displayed node. Binomial Richardson provides a useful independent deterministic cross-check. Pathwise Monte Carlo supports the reported Deltas, while cross-fitted LSMC validates the ordering and level of the surrender premiums. The model-decomposition and hedge experiments show why small premium changes should not be interpreted as negligible model risk: the sign of the price effect depends on the contractual payoff, and unhedged variance and rate exposures can dominate a Delta-only hedge.

The method is designed for a finite set of contractual and exercise dates. Two extensions are particularly natural: reusing adaptive representations across neighboring premium evaluations, and extending the error expansion to a broader class of non-smooth stopping problems. The same functional-state principle may also be useful in other stochastic dynamic programs in which deterministic cash additions prevent one state variable from recombining.

Declarations

Funding. This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Competing interests. The authors declare that they have no competing interests.

Data availability. The numerical inputs and aggregate results supporting the findings are reported in the article and the Supplementary Material.

AI-assisted editing. During the preparation of this work, the authors used ChatGPT 5.6 Sol (OpenAI) solely to assist with language, clarity, and textual organization. The authors reviewed and edited the resulting text and take full responsibility for the content of the publication.

References

- [1] A. R. Bacinello, Endogenous model of surrender conditions in equity-linked life insurance, *Insurance: Mathematics and Economics* 37(2) (2005) 270–296. doi:10.1016/j.insmatheco.2005.02.002.
- [2] A. R. Bacinello, E. Biffis, and P. Millossovich, Pricing life insurance contracts with early exercise features, *Journal of Computational and Applied Mathematics* 233(1) (2009) 27–35. doi:10.1016/j.cam.2008.05.036.
- [3] A. R. Bacinello, E. Biffis, and P. Millossovich, Regression-based algorithms for life insurance contracts with surrender guarantees, *Quantitative Finance* 10(9) (2010) 1077–1090. doi:10.1080/14697680902960242.
- [4] M. J. Brennan and E. S. Schwartz, The pricing of equity-linked life insurance policies with an asset value guarantee, *Journal of Financial Economics* 3(3) (1976) 195–213. doi:10.1016/0304-405X(76)90003-9.
- [5] R. P. Brent, *Algorithms for Minimization without Derivatives*, Prentice–Hall, Englewood Cliffs, NJ, 1973.
- [6] R. Brignone and G. Junike, Exact simulation of stochastic volatility models based on the conditional Fourier–cosine method, *European Journal of Operational Research* 328(3) (2026) 1036–1053. doi:10.1016/j.ejor.2025.08.061.
- [7] D. Brigo and F. Mercurio, A deterministic-shift extension of analytically tractable and time-homogeneous short-rate models, *Finance and Stochastics* 5(3) (2001) 369–387. doi:10.1007/PL00013541.
- [8] M. Costabile, M. Gaudenzi, I. Massabò, and A. Zanette, Evaluating fair premiums of equity-linked policies with surrender option in a bivariate model, *Insurance: Mathematics and Economics* 45(2) (2009) 286–295. doi:10.1016/j.insmatheco.2009.07.008.
- [9] M. Costabile, I. Massabò, and E. Russo, A binomial model for valuing equity-linked policies embedding surrender options, *Insurance: Mathematics and Economics* 42(3) (2008) 873–886. doi:10.1016/j.insmatheco.2007.10.003.
- [10] J. C. Cox, J. E. Ingersoll, Jr., and S. A. Ross, A theory of the term structure of interest rates, *Econometrica* 53(2) (1985) 385–407. doi:10.2307/1911242.

- [11] F. Diener and M. Diener, Asymptotics of the price oscillations of a European call option in a tree model, *Mathematical Finance* 14(2) (2004) 271–293. doi:10.1111/j.0960-1627.2004.00192.x.
- [12] D. H. Douglas and T. K. Peucker, Algorithms for the reduction of the number of points required to represent a digitized line or its caricature, *The Canadian Cartographer* 10(2) (1973) 112–122. doi:10.3138/FM57-6770-U75U-7727.
- [13] European Insurance and Occupational Pensions Authority, *Costs and Past Performance Report 2026*, EIOPA-BoS-26/122, Frankfurt am Main, 1 April 2026. Available at: https://www.eiopa.europa.eu/publications/costs-and-past-performance-report-2026_en.
- [14] M. Gaudenzi, A. Zanette, and M. A. Lepellere, The singular points binomial method for pricing American path-dependent options, *The Journal of Computational Finance* 14(1) (2010) 29–56. doi:10.21314/JCF.2010.214.
- [15] L. Goudenège, A. Molent, X. Wei, and A. Zanette, Enhancing valuation of variable annuities in Lévy models with stochastic interest rate, *Scandinavian Actuarial Journal* 2025(2) (2025) 213–235. doi:10.1080/03461238.2024.2402816.
- [16] L. Goudenège, A. Molent, and A. Zanette, Pricing and hedging GLWB in the Heston and in the Black–Scholes with stochastic interest rate models, *Insurance: Mathematics and Economics* 70 (2016) 38–57. doi:10.1016/j.insmatheco.2016.05.018.
- [17] L. Goudenège, A. Molent, and A. Zanette, Robust pricing of equity-indexed annuities under uncertain volatility and stochastic interest rate, *Insurance: Mathematics and Economics* 127 (2026) 103229. doi:10.1016/j.insmatheco.2026.103229.
- [18] A. Grosen and P. L. Jørgensen, Fair valuation of life insurance liabilities: the impact of interest rate guarantees, surrender options, and bonus policies, *Insurance: Mathematics and Economics* 26(1) (2000) 37–57. doi:10.1016/S0167-6687(99)00041-4.
- [19] L. A. Grzelak and C. W. Oosterlee, On the Heston model with stochastic interest rates, *SIAM Journal on Financial Mathematics* 2 (2011) 255–286. doi:10.1137/090756119.
- [20] S. L. Heston, A closed-form solution for options with stochastic volatility with applications to bond and currency options, *The Review of Financial Studies* 6(2) (1993) 327–343. doi:10.1093/rfs/6.2.327.
- [21] B. Kang and J. Ziveyi, Optimal surrender of guaranteed minimum maturity benefits under stochastic volatility and interest rates, *Insurance: Mathematics and Economics* 79 (2018) 43–56. doi:10.1016/j.insmatheco.2017.12.012.
- [22] H. J. Kushner and P. Dupuis, *Numerical Methods for Stochastic Control Problems in Continuous Time*, 2nd ed., Springer, New York, 2001. doi:10.1007/978-1-4613-0007-6.
- [23] C. Li and L. Wu, Exact simulation of the Ornstein–Uhlenbeck driven stochastic volatility model, *European Journal of Operational Research* 275(2) (2019) 768–779. doi:10.1016/j.ejor.2018.11.057.
- [24] F. A. Longstaff and E. S. Schwartz, Valuing American options by simulation: a simple least-squares approach, *The Review of Financial Studies* 14(1) (2001) 113–147. doi:10.1093/rfs/14.1.113.
- [25] J. Aase Nielsen and K. Sandmann, Equity-linked life insurance: a model with stochastic interest rates, *Insurance: Mathematics and Economics* 16(3) (1995) 225–253. doi:10.1016/0167-6687(95)00007-F.

- [26] W. B. Powell, *Approximate Dynamic Programming: Solving the Curses of Dimensionality*, 2nd ed., Wiley, Hoboken, NJ, 2011.
- [27] T. J. Rivlin, *Chebyshev Polynomials: From Approximation Theory to Algebra and Number Theory*, 2nd ed., Wiley, New York, 1990.
- [28] W. Shen and H. Xu, The valuation of unit-linked policies with or without surrender options, *Insurance: Mathematics and Economics* 36(1) (2005) 79–92. doi:10.1016/j.insmatheco.2004.10.002.
- [29] D. Talay and L. Tubaro, Expansion of the global error for numerical schemes solving stochastic differential equations, *Stochastic Analysis and Applications* 8(4) (1990) 483–509. doi:10.1080/07362999008809220.

A Proofs of technical results

This appendix collects the technical proofs deferred from the construction and convergence sections. The shorter arguments that are useful for following the main logical steps remain in the body of the paper.

A.1 Adaptive-trinomial factor construction

Proof of Lemma 3.1. The exact CIR mean and variance satisfy, uniformly on K_Z ,

$$\begin{aligned} m_h(z) &= z + \kappa(\theta - z)h + O(h^2), \\ s_h^2(z) &= \sigma^2 zh + O(h^2), \end{aligned}$$

while adjacent grid values obey $Z_{q\pm 1} - Z_q = \pm \lambda \sigma \sqrt{zh} + \lambda^2 \sigma^2 h/4$. Write the three probabilities as the solution of the linear system matching unit mass, $m_h(z)$, and $s_h^2(z) + m_h(z)^2$. Substitution and expansion of the explicit solution give (22)–(25). For $\lambda > 1$, every leading probability is strictly positive. For $\lambda = 1$, the leading central mass vanishes and its first non-zero coefficient is $A(z)$; if it is negative, the symmetric stencil is inadmissible. On the standardized limiting supports $(-2, 0, 1)$ and $(-1, 0, 2)$, matching zero mean and unit variance gives respectively $(1/6, 1/2, 1/3)$ and its reversal. Strict positivity of these limits makes the corresponding exact CIR stencils admissible for sufficiently small h . □

Proof of Proposition 3.2. For the rate factor, Lemma 3.1 implies that the standardized innovations converge uniformly to $(-1.4, 0, 1.4)$. For the variance factor, they converge either to $(-1, 0, 1)$ or, up to reflection, to $(-2, 0, 1)$. Hence, for every $\epsilon > 0$ and all sufficiently small h ,

$$\max_{i,j} |z_i^V z_j^X| \leq (2 + \epsilon)(1.4 + \epsilon) \tag{69}$$

uniformly on K . If $|\rho_{Vr}| < 5/14$, choose ϵ so that $|\rho_{Vr}|(2 + \epsilon)(1.4 + \epsilon) < 1$. Then $1 + \rho_{Vr} z_i^V z_j^X > 0$ for every active pair, which is precisely (??). □

A.2 Projected binomial chain

Proof of Proposition 4.2. The functions β_V and β_X in (43) are bounded on K . Expanding (44) around $p_V = p_X = 1/2$ therefore gives, uniformly on K ,

$$\begin{aligned}\rho_-(v, x; h) &= -1 + 2|\beta_V(v) + \beta_X(x)|\sqrt{h} + O(h), \\ \rho_+(v, x; h) &= 1 - 2|\beta_V(v) - \beta_X(x)|\sqrt{h} + O(h).\end{aligned}$$

Every fixed interior correlation lies between these endpoints for sufficiently small h . □

Proof of Proposition 4.3. Fix $\eta > 0$. Under the strict Feller conditions, both continuous CIR factors remain positive. Compact containment of the diffusions and tightness of the marginal lattices allow a rectangle $K = [\underline{v}, \bar{v}] \times [\underline{x}, \bar{x}]$ that is a compact subset of $(0, \infty) \times (0, \infty)$ such that the probability that either the diffusion or the marginal chains leave K before T is below η for all sufficiently small h . By Proposition 4.2, $A_h \cap K = \emptyset$ for sufficiently small h . A projected node can then be reached only after the joint chain leaves K , so $\limsup_{h \downarrow 0} \mathbb{P}(\tau_h \leq T) \leq \eta$. Letting $\eta \downarrow 0$ proves the claim. □

Proof of Theorem 4.4. On each interior compact set, projected and strict local generators coincide for all sufficiently small h except for the mixed V – X covariance on A_h . For a smooth compactly supported test function, the integrated generator discrepancy is bounded by a constant times $\mathcal{M}_h$, which vanishes. Compact containment controls the complement of the localization set. The martingale-problem characterization identifies every subsequential limit with the Heston–CIR++ diffusion; uniqueness gives convergence of the whole sequence. Uniform integrability permits passage to discounted expectations. A finite collection of contractual and exercise dates is then handled by backward induction through the stable event maps. □

A.3 Weak-error rates

Proof of Lemma 4.8. On an interior compact set, the two binomial CIR successors are $O(\sqrt{h})$ from the parent and the exact-mean interpolation probability has the expansion $p_{Z,h} = 1/2 + O(\sqrt{h})$. The resulting conditional variance equals the CIR variance to leading order, with an $O(h^2)$ remainder. The centered variance-one innovation has third moment $O(\sqrt{h})$ and uniformly bounded higher moments. The 2×2 factor coupling preserves this order for mixed third moments, and the residual stock quadrature is symmetric. The equity decomposition and the local martingale correction give the stated first two moments for Y and its mixed covariances. Because each state increment is a drift term of order h plus $\sqrt{h}$ times a standardized innovation, the raw third moments are $O(h^2)$, the fourth moments are $O(h^2)$, and the fifth absolute moment is $O(h^{5/2})$. □

Proof of Theorem 4.9. For a smooth test function, a fourth-order multivariate Taylor expansion

and Lemma 4.8 give (58). The exact discounted semigroup has the same expansion, so the one-step defect is $O(h^2)$. Insert the defect into the telescoping identity for the numerical and exact operator products. Stability transports each local error with a uniform factor, and summing T/h defects yields $O(h)$. A fixed finite number of regular event interfaces changes only the stability constant.

□

A.4 Delta consistency

Proof of Theorem 4.14. On a regular tube bounded away from $F = 0$, the coordinate change $Y = \log F$ is smooth, while the fund branch map itself is affine and has the branch multiplier as derivative. Polynomial moment bounds therefore make the exact and numerical one-step operators stable in the C^1 norm. Differentiating the fourth-order Taylor expansion used in Theorem 4.9 gives a one-step C^1 defect of order h^2 . Apply the derivative to the telescoping identity for the numerical and exact operator products. Each differentiated defect is transported with a uniform C^1 stability factor, and summing T/h defects yields $C_K h$. Because the active branch of each contractual map is fixed on the tube, the finite set of event interfaces preserves differentiability and changes only the constant.

□

A.5 Richardson extrapolation

Proof of Theorem 4.18. First consider an interval with no contractual event and write $Q_n = Q_{t_n, h}$ and $P_n = P_{t_n, t_{n+1}}$. The telescoping identity is

$$Q_0 \cdots Q_{N-1} - P_0 \cdots P_{N-1} = \sum_{n=0}^{N-1} Q_0 \cdots Q_{n-1} (Q_n - P_n) P_{n+1} \cdots P_{N-1}. \quad (70)$$

Apply it to the terminal value and insert Lemma 4.17. Stability and the polynomial moment bounds allow the local remainder to be summed. The $h^2 \Psi_{t_n}$ terms produce

$$h^2 \sum_{n=0}^{N-1} \mathbb{E}[\Psi_{t_n} u(t_{n+1}, Y_{t_{n+1}}, V_{t_{n+1}}, X_{t_{n+1}})] = hc_1 + O(h^2), \quad (71)$$

because the bracketed quantity is sufficiently regular in time and the sum is a Riemann sum. The $h^3 \Xi_{t_n}$ terms and the first-order replacement of numerical transport by exact semigroups contribute $c_2 h^2 + o(h^2)$; the accumulated $O(h^4)$ local remainder is $O(h^3)$. A contractual date is a fixed interface between two such semigroup products. Assumption (R5) preserves the regularity needed to repeat the argument over the finitely many annual intervals and changes only the coefficients c_1 and c_2 .

□