

Output-Only Identification and Spectral Monitoring of Coupled Feedback Networks with Known Time-Varying Actuation

Jihwan Woo

Abstract—Coupled feedback networks are often monitored channel by channel even though cross-channel paths alter both stability margins and transmitted disturbances. We study identification of a structured feedback matrix $\mathbf{L}_t = \Phi \text{diag}(\gamma_t)$ in an *output-only* setting: no commanded, probing, or reference input exists—only temporally separated outputs and the scheduling gains γ_t are observed, while the coupling response Φ and the clearing-window inputs are not. Identification then rests on two sources jointly: the persistent excitation of the observed pre-window output, and two structural features that separate coupling from confounds: the known time variation of the gains, which modulates the closed-loop response in a predictable pattern, and a partial-reversal moment, by which a known fraction of transient displacement is corrected in a subsequent observation window. We give a hierarchy of results: exact local identification of the coupling under a Jacobian rank condition on the gain regimes; a first-order interaction estimator whose identification strength is the minimum eigenvalue of the residualized interaction information matrix (provably unidentified when gains are constant); and a characterization of the estimand as a resolvent sensitivity—directly the right object for screening transmitted disturbances, and a first-stage input to spectral-margin recovery, whose bias-correcting inversion we supply with $\sqrt{T}$ asymptotics for the first-order estimator, a cross-identification theorem mapping each varying gain to exactly identified resolvent rows and columns, and bootstrap validity under consistent-selection conditions; the implemented heuristic’s empirical coverage (90% at nominal 95%) quantifies the remaining gap, and uniform, selection-robust inference remains open. Exact identification holds for the pre-window moment under stated exogeneity and orthogonality assumptions; the full-period-regressor case is characterized as a contaminated estimand. Simulations verify sharpness of the rank condition, quantify the conditioning effect of correlated gain paths, and show that level-regression benchmarks fail under sign-adverse confounding, while a deliberately mismatched two-regime variance-ratio construction in the spirit of heteroskedasticity-based identification is unstable on the same data, whose structural shock variances are constant by design. A case study on leveraged-fund rebalancing feedback, where daily fund disclosures play the role of the known gains, illustrates the method on real data.

Index Terms—System identification, dynamic networks, closed-loop identification, linear parameter-varying systems, spectral monitoring, financial signal processing.

I. INTRODUCTION

J. Woo is with Amazon Web Services (e-mail available on request). The views expressed are those of the author and do not represent those of Amazon Web Services or its affiliates. An application-focused companion paper on the leveraged-fund episode is available as an arXiv preprint; this paper develops the general identification theory.

MANY engineered and economic systems are collections of feedback loops that share a medium: power grids with demand-response contracts, recommendation platforms whose rankings feed back into the behavior being ranked, advertising exchanges where bidding algorithms respond to prices they jointly set, and financial markets hosting leveraged products whose mandated rebalancing trades move the prices the mandates are written on. In all of these, each loop is typically audited in isolation—its own gain, its own stability margin—while the loops are coupled through the shared medium. For a coupled system with elementwise-nonnegative gain matrix $\mathbf{L}$, the spectral radius satisfies $\rho(\mathbf{L}) \geq \max_i \ell_{ii}$: the diagonal audit is guaranteed optimistic, and the gap is largest exactly when coupling is strong and symmetric. Off the diagonal, even one-way coupling transmits displacement from loop to loop through the resolvent $(\mathbf{I} - \mathbf{L})^{-1}$, a channel invisible to every per-loop statistic.

Estimating $\mathbf{L}$ is an identification problem that standard tools do not directly cover. The system operates in closed loop, but there is no reference signal or probing input, so classical closed-loop identification [1] does not apply. The node signals form a cyclic dynamic network, but external excitation—whose presence and location drive identifiability in the dynamic-network literature [4], [5]—is absent. The loop gains are parameter-varying, but the input signal whose observation linear-parameter-varying (LPV) identification requires [9], [12] is latent. Output-only network reconstruction without further structure is subject to fundamental impossibility results [21], [22]. And identification through heteroskedasticity [13] exploits regime changes in unobserved shock variances under a constant structure—the exact dual of our setting, in which the structure varies observably while shock variances need not.

What rescues *structural* identifiability here is a feature these frameworks do not use: *the actuation gains are known*. (Signal excitation alone cannot do it: a static confound and a static coupling are observationally merged until something known moves.) In the applications above, the gain of each loop is disclosed or contractually determined—fund assets and leverage multiples, contracted demand-response capacity, published platform parameters—and it varies over time for reasons unrelated to the disturbances. The known time-varying gain acts as a natural modulating sequence: it changes the closed-loop response in a predictable pattern, so that the response to the *same* latent input differs across gain regimes in a way that is informative about the unknown coupling. This is the feedback-network analogue of a classical signal-processing

idea: transmitter-induced cyclostationarity, where a known periodic modulation renders blind FIR channel identification well posed [18]. Our modulation is not designed and not periodic—it is disclosed by the system itself—and the object it identifies is not an open-loop channel but a closed-loop coupling matrix inside a fixed point. A second structural feature sharpens the moment: in our motivating application the displacement caused by the feedback is *transient*, and a known fraction θ of it reverts in a subsequent observation window, so the reversal itself is a signed, gain-scaled signature of the feedback path.

Contributions. (i) We formalize output-only identification of $\mathbf{L}_t = \Phi \text{diag}(\gamma_t)$ from outputs and known gains γ_t , and derive an interaction estimator based on the reversal moment (Section III–IV). (ii) We give identifiability conditions and prove their sharpness at two levels: in the linearized coefficient map, per-entry identification requires time variation of the corresponding gain path and constant gains leave a continuum of observationally equivalent (coupling, confound) pairs (Theorem 1); exact local identification is governed by the rank of a stacked Jacobian over gain regimes (Theorem 2). (iii) We characterize the estimand as a *resolvent sensitivity* rather than the direct coupling: the detector responds to network-reachable paths—directly the desirable object for transmission screening, and a first-stage input to spectral recovery—and we separate direct-edge and spectral recovery as a second-stage inverse problem (Section V). (iv) Simulations verify the sharpness of the rank condition and quantify failure modes of level-regression and heteroskedasticity-based benchmarks (Section VII). (v) A real-data case study on leveraged-fund rebalancing feedback illustrates the pipeline end to end (Section VIII); the companion empirical paper [23] develops that application in full, while the present paper supplies the identification and monitoring theory.

Table I positions the setting against its nearest neighbors. The precise claim is structural: to our knowledge, no existing framework treats identification of an algebraic feedback coupling from outputs alone when the identifying variation is a known, column-wise, time-varying gain and the informative moment is a signed partial reversal; individual ingredients appear separately in the table’s rows.

II. RELATED WORK

Four literatures approach the identification of interconnected dynamical systems, and each leaves the present setting uncovered for a structural reason.

Closed-loop and network identification. Classical closed-loop identification resolves the circularity of feedback through an external reference or probing signal [1], [2]; prediction-error methods for dynamic networks generalize this to structured interconnections [3], and a mature theory now characterizes network identifiability by the presence, location, and rank of *external* excitation and disturbance signals [4]–[6]. Topology-oriented variants reconstruct the interconnection structure from node spectra [7], [8], [21]. Two obstructions separate our problem from this line. First, no commanded excitation exists—no reference, no probing, no exogenously

TABLE I
POSITIONING AGAINST NEIGHBORING IDENTIFICATION FRAMEWORKS.

Framework	Observed	Excitation	Output
Closed-loop [1]	ID u, y , reference	external reference	plant TF
Dynamic networks [4]	nodes + excitation	external signals	module TFs
LPV-ID [9]	u, y , scheduling	input u changes	$A(p), B(p)$
Heteroskedasticity [13]	y	in shock covariance	constant A_0
MIC [†] blind ID [18]	y , periodic pre-coder	designed modulation	FIR channel
Output-only recon. [21], [22]	y only	none; limits apply	topology (under assumptions)
This paper	y , gains γ_t	known time-varying gains	Φ (1st-order), screening alarms*

[†]MIC = modulation-induced cyclostationarity. *Exact Φ under regime-specific clean moments (Theorem 2); spectral recovery via a heuristic second stage with calibrated inference open (Section V).

excited node; the identifiability currency of the network literature is unavailable. Second, the object is a static-within-window coupling embedded in an algebraic fixed point (2), with a cycle at its core, whereas sharp partial-excitation results are largely acyclic [5]. To be precise about what is and is not missing: the pre-window output $\mathbf{r}_1$ is observed and persistently exciting, so it supplies the forcing any regression needs; what is absent is any *commanded or probing* signal, and what takes its place as the source of *structural* identification—separating coupling from gain-invariant confounds—is the known time variation of the actuation gains, which has no counterpart in this theory.

LPV identification. If the gain path γ_t is read as a scheduling signal, our model is a linear parameter-varying system with known scheduling [9]. LPV prediction-error and subspace methods [10], [12] estimate the full parameter-varying dynamics, including closed-loop variants—but all require the input signal to be measured; identifiability is inherited from input excitation, exactly as in the LTI case. Recent work treats the *scheduling* as latent and learns it [11]. Our problem inverts both premises: the input is latent, the scheduling is known, and the question is whether known scheduling variation alone can replace input observation. To our knowledge this corner—output-only LPV identification with scheduling entering only the feedback path—appears unoccupied, though we state this as a literature claim open to correction rather than a theorem.

Identification through heteroskedasticity. Econometrics identifies simultaneous systems from regime changes in the *variances* of latent shocks under a constant structural matrix [13]–[17]. This is the closest conceptual relative: both approaches buy identification from time variation rather than exclusion restrictions or instruments. But the two are duals, not variants. There, the structure $\mathbf{A}_0$ is constant and the shock covariance moves, unobserved, estimated as regimes; here, the structure $\mathbf{L}_t = \Phi \text{diag}(\gamma_t)$ moves *observably* while shock covariances may be constant. The duality is not cosmic: Section VII includes a deliberately mismatched two-

regime variance-ratio benchmark on data generated by our model—with structural shock variances constant by design, its variance-ratio denominator is unstable (reduced-form variances do move with the gains, but not in the way that construction requires), which illustrates the model mismatch rather than numerically refuting heteroskedasticity identification—while our estimator is undefined on data generated by theirs (constant gains violate the rank condition of Theorem 1). In the two stylized designs compared here, each method’s identifying variation is the other’s maintained constancy.

Adjacent model classes. Three further neighbors deserve explicit contrast. Bilinear system identification [25] treats products of states and *inputs*, so its excitation is again a measured input, absent here. The varying-coefficient regression literature [26] supplies exactly the reduced-form skeleton of our (22)—coefficients driven by an observed effect modifier—but stops at the conditional mean: it has no feedback fixed point, no reversal moment, and therefore no map from fitted coefficients to a structural coupling matrix or its spectral radius; our contribution is precisely that structural layer. Output-only (operational) modal analysis [27] identifies modal structure from ambient responses, but under an unobserved stationary white excitation and time-invariant dynamics—the two assumptions our setting replaces with a known, nonstationary gain path.

Known modulation in signal processing. The principle that a known time-varying modulation renders a blind problem well posed is classical in communications: transmitter-induced cyclostationarity and periodic precoding enable blind FIR channel identification from second-order statistics [18]–[20]. We import the principle and change every ingredient: the modulation γ_t is disclosed by the system rather than designed, aperiodic rather than cyclostationary, and enters a feedback fixed point rather than a feedforward channel; identification additionally exploits a signed reversal moment (5) with no communications analogue. Conversely, the impossibility results for output-only network reconstruction [21], [22] delimit what any method can do without such structure—our contribution can be read as exhibiting one structural supplement (known gains plus reversal) under which the impossibility dissolves; the individual mathematical devices are standard, and the novelty claim rests on the model and its use of disclosed modulation, not on any single tool.

III. PROBLEM FORMULATION

A. Observation model

There are n channels. At each period t we observe a pre-window output $\mathbf{r}_{1,t} \in \mathbb{R}^n$, a post-window output $\mathbf{r}_{\text{on},t} \in \mathbb{R}^n$, and the actuation gains $\gamma_t \in \mathbb{R}_+^n$. The within-period feedback operates in a clearing window between the two observations. Each channel j places a mandated action proportional to its own gain and its *full-period* signal $r_{1,j,t} + r_{2,j,t}$, where $\mathbf{r}_{2,t}$ is the (latent) clearing-window displacement—the self-reference that creates the fixed point—and actions displace all channels through a coupling response $\Phi = [\phi_{ij}] \geq 0$:

$$\mathbf{r}_{2,t} = \Phi \text{diag}(\gamma_t) (\mathbf{r}_{1,t} + \mathbf{r}_{2,t}) + \mathbf{v}_t = \mathbf{L}_t (\mathbf{r}_{1,t} + \mathbf{r}_{2,t}) + \mathbf{v}_t, \quad (1)$$




Fig. 1. The coupled feedback fixed point (1).

Note: the actuation gains $\text{diag}(\gamma_t)$ are disclosed—the only observed element of the loop besides the outputs—and their time variation is the identification resource; the coupling Φ and inputs $\mathbf{v}_t$ are unobserved. The dashed path is the self-reference that makes $\mathbf{r}_{2,t}$ a fixed point.

where $\mathbf{v}_t$ is latent clearing-window noise. Collecting $\mathbf{r}_{2,t}$ terms, $(\mathbf{I} - \mathbf{L}_t)\mathbf{r}_{2,t} = \mathbf{L}_t\mathbf{r}_{1,t} + \mathbf{v}_t$, so that whenever $1 \notin \sigma(\mathbf{L}_t)$,

$$\mathbf{r}_{2,t} = (\mathbf{I} - \mathbf{L}_t)^{-1} (\mathbf{L}_t\mathbf{r}_{1,t} + \mathbf{v}_t). \quad (2)$$

We maintain $\rho(\mathbf{L}_t) < 1$, under which the resolvent admits the Neumann expansion

$$(\mathbf{I} - \mathbf{L}_t)^{-1} = \sum_{k=0}^{\infty} \mathbf{L}_t^k, \quad \mathbf{M}_t \equiv (\mathbf{I} - \mathbf{L}_t)^{-1} - \mathbf{I} = \sum_{k=1}^{\infty} \mathbf{L}_t^k, \quad (3)$$

whose k th term represents the k th feedback round; off-diagonal (cross-channel) transmission enters already at $k = 1$. Two identities used repeatedly follow directly from the definition of $\mathbf{M}_t$:

$$(\mathbf{I} - \mathbf{L}_t)^{-1} = \mathbf{I} + \mathbf{M}_t, \quad (\mathbf{I} - \mathbf{L}_t)^{-1} \mathbf{L}_t = \mathbf{M}_t. \quad (4)$$

The post-window mechanism reverses a known share $\theta \in (0, 1]$ of the displacement and carries a constant confound:

$$\mathbf{r}_{\text{on},t} = -\theta \mathbf{r}_{2,t} + \mathbf{C} \mathbf{r}_{1,t} + \mathbf{w}_t. \quad (5)$$

Unknowns: the coupling $\Phi \in \mathbb{R}^{n \times n}$ (constant), the confound $\mathbf{C}$ (constant; lead-lag or common-factor loadings), and the noise second moments. Known: γ_t (disclosed) and θ (calibrated externally or normalized; to first order unknown θ is a global scale ambiguity on Φ , while in the exact model powers of Φ enter at different orders—see the Remark after Theorem 2).

Assumption 1 (Exogenous gains). $\{\gamma_t\}$ is deterministic or independent of $\{\mathbf{r}_{1,t}, \mathbf{v}_t, \mathbf{w}_t\}$, and $\mathbf{C}$ does not vary with γ_t .

Assumption 1 is the exclusion restriction: the confound may be arbitrary but must not track the disclosed gains. Its failure mode is quantified in Section VII (scenario C).

Assumption 2 (Stochastic regularity). $\{\mathbf{r}_{1,t}, \mathbf{v}_t, \mathbf{w}_t\}$ is stationary and ergodic with finite fourth moments; $\mathbb{E}[\mathbf{v}_t | \mathbf{r}_{1,t}, \gamma_t] = \mathbb{E}[\mathbf{w}_t | \mathbf{r}_{1,t}, \gamma_t] = \mathbf{0}$; and the pre-window signal is persistently exciting conditional on the gains: $\mathbb{E}[\mathbf{r}_{1,t} \mathbf{r}_{1,t}^\top | \gamma_t] \succ 0$.

Gain variation alone is not sufficient: identification also needs signal excitation and conditional orthogonality, which Assumption 2 supplies. Theorems 1 and 2 are population-level identification results under Assumptions 1–2; the accompanying finite-sample estimation theory (rates, asymptotic normality under mixing, efficiency) is part of the UQ program of Section VI-D and is stated there as open where it is open.

B. The moment, derived

Substituting (2) into (5) and using (4),

$$\begin{aligned} \mathbf{r}_{2,t} &= (\mathbf{I} - \mathbf{L}_t)^{-1} \mathbf{L}_t \mathbf{r}_{1,t} + (\mathbf{I} - \mathbf{L}_t)^{-1} \mathbf{v}_t \\ &= \mathbf{M}_t \mathbf{r}_{1,t} + (\mathbf{I} + \mathbf{M}_t) \mathbf{v}_t, \end{aligned} \quad (6)$$

so the post-window output is

$$\mathbf{r}_{\text{on},t} = (\mathbf{C} - \theta \mathbf{M}_t) \mathbf{r}_{1,t} - \theta (\mathbf{I} + \mathbf{M}_t) \mathbf{v}_t + \mathbf{w}_t. \quad (7)$$

Under Assumption 2, augmented with $\mathbb{E}[\mathbf{w}_t \mathbf{v}_t^\top | \gamma_t] = \mathbf{0}$ (post-window noise orthogonal to clearing-window noise, needed only for the full-period regressor below), the population projection of $\mathbf{r}_{\text{on},t}$ on $\mathbf{r}_{1,t}$ conditional on the gain regime γ_t is (writing $\Sigma_1 = \mathbb{E}[\mathbf{r}_{1,t} \mathbf{r}_{1,t}^\top | \gamma_t]$, $\Sigma_v = \mathbb{E}[\mathbf{v}_t \mathbf{v}_t^\top | \gamma_t]$, which by gain independence (Assumption 1) coincide with their unconditional counterparts)

$$\begin{aligned} \mathbf{B}(\gamma_t) &= \mathbb{E}[\mathbf{r}_{\text{on},t} \mathbf{r}_{1,t}^\top | \gamma_t] \mathbb{E}[\mathbf{r}_{1,t} \mathbf{r}_{1,t}^\top | \gamma_t]^{-1} \\ &= (\mathbf{C} - \theta \mathbf{M}_t) \Sigma_1 \Sigma_1^{-1} = \mathbf{C} - \theta \mathbf{M}_t. \end{aligned} \quad (8)$$

The reversal moment thus identifies $\mathbf{M}_t$ up to the known scale θ and the additive confound—*exactly*, with no small-gain approximation, when the regressor is the pre-window output. If instead the available regressor is the full-period output $\mathbf{R}_t = \mathbf{r}_{1,t} + \mathbf{r}_{2,t} = (\mathbf{I} + \mathbf{M}_t)(\mathbf{r}_{1,t} + \mathbf{v}_t)$, the two projection ingredients follow from (7) and the orthogonality assumptions:

$$\begin{aligned} \mathbb{E}[\mathbf{r}_{\text{on}} \mathbf{R}^\top | \gamma_t] &= [\mathbf{C} \Sigma_1 - \theta \mathbf{M}_t \Sigma_1 \\ &\quad - \theta (\mathbf{I} + \mathbf{M}_t) \Sigma_v] (\mathbf{I} + \mathbf{M}_t)^\top, \\ \mathbb{E}[\mathbf{R} \mathbf{R}^\top | \gamma_t] &= (\mathbf{I} + \mathbf{M}_t) (\Sigma_1 + \Sigma_v) (\mathbf{I} + \mathbf{M}_t)^\top, \end{aligned} \quad (9) \quad (10)$$

and dividing (9) by (10) gives the exact estimand

$$\begin{aligned} \mathbf{B}(\gamma_t) &= [\mathbf{C} \Sigma_1 - \theta \mathbf{M}_t \Sigma_1 - \theta (\mathbf{I} + \mathbf{M}_t) \Sigma_v] \\ &\quad \times (\Sigma_1 + \Sigma_v)^{-1} (\mathbf{I} + \mathbf{M}_t)^{-1}, \end{aligned} \quad (11)$$

verified numerically in the replication code. Two consequences matter. First, the confound no longer enters as a clean additive intercept: even as $\Sigma_v \rightarrow 0$ the confound term is $\mathbf{C}(\mathbf{I} + \mathbf{M}_t)^{-1}$, itself gain-dependent, so the constant-confound separation that drives Theorem 1 is exact only for the pre-window regressor. The theory below is therefore stated for the pre-window moment (8); when only full-period outputs are available, (11) defines the contaminated estimand. The discrepancy from the clean case is $O(\|\mathbf{M}\|)$ under bounded $\mathbf{C}$ —even at $\Sigma_v = \mathbf{0}$ the full-period estimand is $(\mathbf{C} - \theta \mathbf{M}_t)(\mathbf{I} + \mathbf{M}_t)^{-1}$, still gain-contaminated—while only the latent-noise component vanishes with the clearing-window share Σ_v . A small clearing-window variance therefore does *not* by itself license the clean theorem; the convention analysis for the financial application is in the companion paper [23]. The identification question is when the observable map $\gamma \mapsto \mathbf{B}(\gamma)$ pins down $(\Phi, \mathbf{C})$.

IV. IDENTIFIABILITY

To first order in $\|\mathbf{L}_t\|$ (the empirically relevant regime; higher-order effects are treated as bias and quantified in

Section VII), $\mathbf{M}_t \approx \mathbf{L}_t = \Phi \text{diag}(\gamma_t)$ and (8) becomes entrywise

$$b_{ij}(t) = c_{ij} - \theta \phi_{ij} \gamma_{j,t} + O(\|\mathbf{L}\|^2), \quad (12)$$

the observable slope varies linearly in the known signal $\gamma_{j,t}$, with the confound as intercept. Throughout, $\tilde{\gamma}_{j,t} = (\gamma_{j,t} - \bar{\gamma}_j)/s_{\gamma_j}$ denotes the standardized gain path (sample mean and standard deviation over the estimation window), $\tilde{\Gamma} \in \mathbb{R}^{T \times n}$ stacks them, and $\Sigma_r = \mathbb{E}[\mathbf{r}_{1,t} \mathbf{r}_{1,t}^\top]$.

Theorem 1 (Identification and sharpness, linearized). *Parts (i)–(ii) concern identification from the conditional coefficient map $t \mapsto b_{ij}(t)$ in (12); part (iii) concerns the pooled raw-data regression, whose rank condition is $\lambda_{\min}(\mathbf{Q}) > 0$ and does not follow from (i) alone.*

(i) *The pair $(c_{ij}, \theta \phi_{ij})$ is identified if and only if the design $[\mathbf{1}, \gamma_{j,1:T}]$ has full column rank, i.e., $\text{Var}_t(\gamma_{j,t}) > 0$. (ii) If $\text{Var}_t(\gamma_{j,t}) = 0$, the set of observationally equivalent pairs is the line $\{(c_{ij} + \theta s \gamma_j, \phi_{ij} + s) : s \in \mathbb{R}\}$; no estimator can distinguish coupling from confound. (iii) In the joint regression across senders, the information matrix of the interaction block is $\mathbf{Q} = T^{-1} \mathbb{E}[\mathbf{Z}^\top \mathbf{M}_X \mathbf{Z}]$, where the columns of $\mathbf{Z}$ are $z_{j,t} = r_{1,j,t} \tilde{\gamma}_{j,t}$ and $\mathbf{M}_X$ residualizes the intercept and main effects; identification strength is $\lambda_{\min}(\mathbf{Q})$. Under gains independent of the pre-window signals, $\mathbb{E}[z_j z_k] = [\Sigma_r \circ \mathbf{G}]_{jk}$ (Hadamard product), with $\mathbf{G} = T^{-1} \tilde{\Gamma}^\top \tilde{\Gamma}$ the Gram matrix of standardized gain paths: the conditioning of the gain design matters exactly to the extent that the signals themselves are cross-correlated, and for strongly factor-driven signals (Σ_r close to rank one) $\kappa(\mathbf{G})$ becomes the governing factor of the variance inflation.*

Part (iii) is not a technicality: in the motivating application, product launches make all gains ramp together, and Section VII shows the false-alarm rate of a signed detector nearly doubling between a correlated staircase design (condition number ~ 247) and independent regime switching (~ 12) at equal per-channel variances.

Proof: (i) Stacking (12) over t gives, per entry (i, j) , the linear system $\mathbf{b}_{ij} = \mathbf{X}_j \beta$ with $\mathbf{X}_j = [\mathbf{1}, \gamma_{j,1:T}] \in \mathbb{R}^{T \times 2}$ and $\beta = (c_{ij}, -\theta \phi_{ij})^\top$. The map $\beta \mapsto \mathbf{X}_j \beta$ is injective iff $\text{rank}(\mathbf{X}_j) = 2$, which holds iff $\gamma_{j,t}$ is not constant, i.e., $\text{Var}_t(\gamma_{j,t}) > 0$. (ii) If $\gamma_{j,t} \equiv \gamma_j$, then for any $s \in \mathbb{R}$ the pair $(c_{ij} + \theta s \gamma_j, \phi_{ij} + s)$ generates the identical moment:

$$(c_{ij} + \theta s \gamma_j) - \theta(\phi_{ij} + s) \gamma_j = c_{ij} - \theta \phi_{ij} \gamma_j, \quad (13)$$

so the observationally equivalent set is a line and no statistic of the data distinguishes its points. (iii) Here $\mathbf{X}$ contains the intercept and the n signal main effects $r_{1,k,t}$, gains are centered ($\mathbb{E}[\tilde{\gamma}_j] = 0$) and independent of signals, so $\mathbb{E}[z_{j,t} r_{1,k,t}] = \mathbb{E}[r_{1,j,t} r_{1,k,t}] \mathbb{E}[\tilde{\gamma}_j] = 0$ and residualization leaves the interaction block orthogonal to $\mathbf{X}$ in population. Then,

$$\mathbb{E}[z_{j,t} z_{k,t}] = \mathbb{E}[r_{1,j,t} r_{1,k,t}] \mathbb{E}[\tilde{\gamma}_j \tilde{\gamma}_k] = [\Sigma_r]_{jk} [\mathbf{G}]_{jk}, \quad (14)$$

i.e. the interaction Gram matrix is the Hadamard product $\Sigma_r \circ \mathbf{G}$; residualizing intercept and main effects yields $\mathbf{Q}$ as stated. By the Schur product theorem $\Sigma_r \circ \mathbf{G} \succeq 0$, and for the

equicorrelation model $\Sigma_r = \sigma^2[(1 - \varrho)\mathbf{I} + \varrho\mathbf{1}\mathbf{1}^\top]$ (with ϱ the signal correlation, distinct from the spectral radius ρ),

$$\begin{aligned} \Sigma_r \circ \mathbf{G} &= \sigma^2\{(1 - \varrho) \text{diag}(G_{jj}) + \varrho \mathbf{G}\}, \\ \lambda_{\min}(\Sigma_r \circ \mathbf{G}) &\geq \sigma^2[(1 - \varrho) \min_j G_{jj} + \varrho \lambda_{\min}(\mathbf{G})] : \quad (15) \end{aligned}$$

as $\varrho \uparrow 1$ (factor-dominated signals) the bound is governed by $\lambda_{\min}(\mathbf{G})$, while for $\varrho = 0$ (uncorrelated signals) $\Sigma_r \circ \mathbf{G} = \sigma^2 \text{diag}(G_{jj})$ and gain-path correlation is harmless. Degenerate $\mathbf{G}$ with factor-driven signals returns linear combinations of senders to the unidentified case of (ii). ■

Theorem 2 (Regular local identification, exact fixed point). *Let $\mathbf{B}(\gamma) = \mathbf{C} - \theta \mathbf{M}(\gamma)$ with $\mathbf{M}(\gamma) = (\mathbf{I} - \Phi \text{diag}(\gamma))^{-1} - \mathbf{I}$ be observed at gain regimes $\gamma^{(1)}, \dots, \gamma^{(q)}$, $\rho(\Phi \text{diag}(\gamma^{(k)})) < 1$ for all k (regimes recur with positive probability, or are estimable by local smoothing, so that the regime-specific moments are available). Define, for each regime, the $n^2 \times n^2$ Jacobian block*

$$\mathbf{D}_k \equiv \frac{\partial \text{vec} \mathbf{M}(\gamma^{(k)})}{\partial \text{vec} \Phi} = [\text{diag}(\gamma^{(k)}) \mathbf{R}_k]^\top \otimes \mathbf{R}_k, \quad (16)$$

where $\mathbf{R}_k = (\mathbf{I} - \Phi \text{diag}(\gamma^{(k)}))^{-1}$ (calligraphic to distinguish the resolvent from the full-period output $\mathbf{R}_t$). Then full column rank of the stacked differenced Jacobian,

$$\text{rank} \begin{bmatrix} \mathbf{D}_2 - \mathbf{D}_1 \\ \vdots \\ \mathbf{D}_q - \mathbf{D}_1 \end{bmatrix} = n^2, \quad (17)$$

is sufficient for local identification of $(\mathbf{C}, \Phi)$ at the true parameters, and necessary for regular (first-order) local identification, i.e. on the set where the rank is locally constant; rank deficiency at isolated points does not by itself imply non-identification. In particular $q \geq 2$ distinct regimes are necessary; if all $\gamma^{(k)}$ coincide, every $\mathbf{D}_k$ is identical, the stacked matrix is zero, and constant gains again yield a continuum of observationally equivalent pairs, whose first-order tangent contains the lines of Theorem 1(ii).

Proof: Differentiating $\mathbf{M} = (\mathbf{I} - \mathbf{L})^{-1} - \mathbf{I}$ along $d\Phi$ with γ fixed at $\gamma^{(k)}$,

$$d\mathbf{M} = \mathbf{R}_k (d\Phi) \text{diag}(\gamma^{(k)}) \mathbf{R}_k, \quad (18)$$

$$\text{vec}(d\mathbf{M}) = [(\text{diag}(\gamma^{(k)}) \mathbf{R}_k)^\top \otimes \mathbf{R}_k] \text{vec}(d\Phi), \quad (19)$$

by $\text{vec}(\mathbf{AXB}) = (\mathbf{B}^\top \otimes \mathbf{A}) \text{vec} \mathbf{X}$, which gives (16). The stacked observation map $(\mathbf{C}, \Phi) \mapsto \{\mathbf{B}(\gamma^{(k)})\}_{k=1}^q$ has Jacobian $[\mathbf{I}_{n^2}, -\theta \mathbf{D}_k]_{k=1, \dots, q}$ (row blocks). Eliminating $\mathbf{C}$ by subtracting the first block row leaves $-\theta[\mathbf{D}_k - \mathbf{D}_1]_{k \geq 2}$ acting on $\text{vec} \Phi$ alone; the implicit function theorem gives local injectivity under (17) (and its necessity at rank-regular points), and $\mathbf{C}$ is then recovered from the first block. Necessity of $q \geq 2$ and failure under coinciding regimes are immediate. To first order $\mathbf{R}_k \approx \mathbf{I}$, so $\mathbf{D}_k - \mathbf{D}_1 \approx \text{diag}(\gamma^{(k)} - \gamma^{(1)})^\top \otimes \mathbf{I}$, whose rank is $n \cdot \#\{j : \gamma_j^{(k)} \neq \gamma_j^{(1)} \text{ for some } k\}$ —recovering the per-column variance condition of Theorem 1(i) as the linearized special case. ■

Remark 1 (Scale, linearized implementation). *Only the products $\theta \phi_{ij}$ enter (12); the level of Φ is identified once θ*

is calibrated or normalized (e.g., own-channel reversal normalization). In the exact resolvent, powers of Φ enter with different orders, so unknown θ is a global scale ambiguity only to first order; all spectral statements inherit the calibration of θ .

V. THE ESTIMAND IS A RESOLVENT SENSITIVITY

Beyond first order, the object the interaction estimator approximates is not ϕ_{ij} but (a gain-distribution-weighted projection of) the sensitivity of the resolvent to the j th gain; with discrete regime variation the coefficient is a secant rather than a derivative, a distinction that matters for the second-stage inversion below. Differentiating $\mathbf{M} = (\mathbf{I} - \mathbf{L})^{-1} - \mathbf{I}$ with $d\mathbf{L} = \Phi \mathbf{e}_j \mathbf{e}_j^\top d\gamma_j$ and using $d(\mathbf{I} - \mathbf{L})^{-1} = (\mathbf{I} - \mathbf{L})^{-1} (d\mathbf{L}) (\mathbf{I} - \mathbf{L})^{-1}$,

$$\frac{\partial \mathbf{M}}{\partial \gamma_j} = (\mathbf{I} - \mathbf{L})^{-1} \Phi \mathbf{e}_j \mathbf{e}_j^\top (\mathbf{I} - \mathbf{L})^{-1}, \quad (20)$$

which is nonzero for every pair (i, j) connected by a directed path through the network, not only for direct edges: expanding both resolvents by (3), the (i, j) entry of (20) collects all walks $j \rightarrow \dots \rightarrow i$ that pass through the perturbed gain. At first order in $\|\mathbf{L}\|$, $\partial \mathbf{M} / \partial \gamma_j \rightarrow \Phi \mathbf{e}_j \mathbf{e}_j^\top$, recovering (12). This is a feature for the monitoring problem—transmission alarms and spectral margins are resolvent objects—and a caveat for topology recovery: a signed detector applied entrywise flags 2-hop-reachable zero entries at well above nominal size while holding exactly nominal size on unreachable entries (Section VII). Direct-edge recovery is therefore a second-stage inverse problem—inverting (20) over the detected reachability pattern—which we formulate but do not pursue empirically in this paper.

A. Which gains identify which entries

Theorem 2 is an all-or-nothing rank condition. The following refines it to a per-entry, graph-readable map from gain variation to identified quantities, and is exact rather than first-order. Write $\mathbf{m}_l$ for the l th column of $\mathbf{M}$ and $\mathbf{r}_l$ for the l th row of $\mathbf{I} + \mathbf{M}$, evaluated at the operating gain γ .

Theorem 3 (Cross identification from varying gains). *Let $\mathcal{V} \subseteq \{1, \dots, n\}$ index the gains that vary near the operating point γ , and assume: (a) $\gamma_l > 0$ for every $l \in \mathcal{V}$ (at $\gamma_l = 0$ the undivided sensitivity $\partial \mathbf{M} / \partial \gamma_l = (\mathbf{I} - \mathbf{L})^{-1} \Phi \mathbf{e}_l \mathbf{e}_l^\top (\mathbf{I} - \mathbf{L})^{-1}$ applies instead); and (b) a local-support condition: the support of the gain process contains a relatively open neighborhood of γ in the coordinates $\gamma_{\mathcal{V}}$ (equivalently, local gain increments span $\mathbb{R}^{|\mathcal{V}|}$), so that every coordinate partial derivative $\partial \mathbf{M} / \partial \gamma_l$, $l \in \mathcal{V}$, is recoverable from the observable coefficient map. If the gains move only along a lower-dimensional path—the empirically relevant case of strongly correlated gain paths—only the directional derivatives along that path are observed, and the conclusions below hold for the coordinate derivatives only once (b) is met. Then: (i) the exact sensitivity factorizes as the rank-one outer product*

$$\frac{\partial \mathbf{M}}{\partial \gamma_l} = \frac{1}{\gamma_l} \mathbf{m}_l \mathbf{r}_l, \quad l \in \mathcal{V}; \quad (21)$$

(ii) provided the nondegeneracy condition $\mathbf{m}_{\cdot l} \neq \mathbf{0}$ holds (node l has at least one outgoing feedback path; otherwise the sensitivity vanishes identically and only that fact is learned), the (l, l) entry pins the factorization scale: with $p_{ll} = [\partial \mathbf{M} / \partial \gamma_l]_{ll}$, the diagonal entry solves $x(1+x) = \gamma_l p_{ll}$, whose nonnegative root is M_{ll} —and when $M_{ll} = 0$ the scale is pinned instead by the row normalization $r_{ll} = 1$: the l th column of the derivative slab equals $\mathbf{m}_{\cdot l} / \gamma_l$ directly, and $\mathbf{r}_{\cdot l}$ follows from any nonzero entry of $\mathbf{m}_{\cdot l}$; hence the full column $\mathbf{m}_{\cdot l}$ and the full row $\mathbf{r}_{\cdot l}$ are exactly identified for every $l \in \mathcal{V}$ with an outgoing path, so that a guaranteed point-identified subset of $\mathbf{M}$ is the cross pattern $\{(i, j) : i \in \mathcal{V}' \text{ or } j \in \mathcal{V}'\}$, $\mathcal{V}' = \{l \in \mathcal{V} : \mathbf{m}_{\cdot l} \neq \mathbf{0}\}$; (iii) in the linearized regime the statement reduces to Theorem 1: $\partial \mathbf{M} / \partial \gamma_l \rightarrow \Phi \mathbf{e}_l \mathbf{e}_l^\top$, only column l of Φ is exposed, and entries with $i, j \notin \mathcal{V}$ are unidentified at first order. Whether the complement block is identified in the exact model through higher-order joint variation is not settled by this theorem: (i)–(ii) exhibit a point-identified subset, not the full identified set.

Proof: (i) is (20) rewritten: $(\mathbf{I} - \mathbf{L})^{-1} \Phi \mathbf{e}_l = (\mathbf{I} - \mathbf{L})^{-1} \mathbf{L} \mathbf{e}_l / \gamma_l = \mathbf{m}_{\cdot l} / \gamma_l$ and $\mathbf{e}_l^\top (\mathbf{I} - \mathbf{L})^{-1} = \mathbf{r}_{\cdot l}$. (ii) The (l, l) entry of (21) is $M_{ll}(1 + M_{ll}) / \gamma_l$; the map $x \mapsto x(1+x)$ is strictly increasing on $x \geq 0$, so the root is unique under the maintained nonnegativity, and dividing the observed rank-one factors by the resulting scale recovers column and row exactly. (iii) follows by $\mathbf{M} \rightarrow \mathbf{L}$, $(\mathbf{I} + \mathbf{M}) \rightarrow \mathbf{I}$. All three statements are verified numerically in the replication code. ■

The graph reading: a varying gain at a node with outgoing feedback illuminates every transmission path into it and out of it—row and column l of the resolvent—and pins its own diagonal; a varying gain at a sink node (no outgoing path) reveals only that fact, and gains that never move leave the $\mathcal{V}^c \times \mathcal{V}^c$ block of $\mathbf{M}$ dark at first order. Two consequences follow. For *monitoring*, partial gain variation is enough for partial surveillance under the local-support condition (b): all transmission entries M_{ij} with a varying sender or receiver are point-identified without any support assumption on Φ , which is exactly the set of alarms a supervisor of the varying complexes needs. For *estimation*, (21) implies that the derivative slab associated with gain l is rank one in population—so a rank-one (SVD) projection of the estimated full interaction slab (all $r_{1,k} \tilde{\gamma}_l$ regressors) is a noise-reduced estimator of $(\mathbf{m}_{\cdot l}, \mathbf{r}_{\cdot l})$ that bypasses the fixed-point inversion (27) entirely. By Eckart–Young the rank-one *projection subproblem* is globally optimal—so that step, unlike the nonlinear least-squares fit, has no local-minimum pathology; no such claim is made for the full estimator after scale recovery, assembly, and fallback. We implement a columns-only version (rows are redundant when all columns are available and are usable as a diagnostic) and find that, among root-success paths in the design of Section VII, it attains lower spectral bias and far lower dispersion than the selected two-stage inversion; numbers, Monte Carlo accounting, and failure handling are reported in Section VI-D. The implementation uses global standardized slopes, so the secant caveat of Section VI applies to its estimand; its distribution theory alongside Theorem 5 is a natural next step. Whether the dark block is identified through

higher-order joint variation of $\gamma_{\mathcal{V}}$ in the exact model is an open question we state precisely: all $\gamma_{\mathcal{V}}$ -derivatives of $\mathbf{B}$ expose products of cross-pattern entries only, so any identification of the dark block must come through the nonlinear dependence of those cross entries on the full Φ .

VI. ESTIMATION, ALGORITHMS, AND MONITORING

A. Interaction estimator

Algorithm 1 implements the moment (12). For receiver i , the regression

$$r_{\text{on},i,t} = \alpha_i + \sum_{j=1}^n (b_{ij} r_{1,j,t} + d_{ij} r_{1,j,t} \tilde{\gamma}_{j,t}) + \varepsilon_{i,t} \quad (22)$$

separates the constant confound (absorbed by b_{ij} together with the gain-mean response) from the gain-tracking response $d_{ij} = -\theta \phi_{ij} s_{\gamma_j} + O(\|\mathbf{L}\|^2)$, where s_{γ_j} is the standardization scale of $\tilde{\gamma}_{j,t} = (\gamma_{j,t} - \bar{\gamma}_j) / s_{\gamma_j}$. Heteroskedasticity-robust variances give entrywise t -statistics; the signed decision rule $t_{ij} < -z_{1-\alpha}$ exploits the known direction of the reversal (coupling makes the interaction *negative*), which a confound of either sign cannot mimic without tracking γ_t (Assumption 1).

Two scope qualifications attach to (22). First, it is the *low-gain* implementation of the identification theory: it matches Theorem 1 exactly and Theorem 2 to first order; an exact-fixed-point nonlinear least-squares estimator fitting $\mathbf{B}(\gamma) = \mathbf{C} - \theta[(\mathbf{I} - \Phi \text{diag}(\gamma))^{-1} - \mathbf{I}]$ across regimes is the natural second stage, with the linear estimates as initialization. Second, (22) includes the same-index interactions $r_{1,j,t} \tilde{\gamma}_{j,t}$ only, while beyond first order the expansion of $(\mathbf{C} - \theta \mathbf{M}(\gamma)) \mathbf{r}_1$ generates all n^2 cross terms $r_{1,k,t} \tilde{\gamma}_{j,t}$ with coefficients $-\theta \partial M_{ik} / \partial \gamma_j$; with correlated regressors the same-index OLS coefficient is then a projection mixture, so Algorithm 1 estimates the first-order direct-coupling approximation with an $O(\|\mathbf{L}\|^2)$ projection bias, not the exact sensitivity. The full n^2 -interaction regression is feasible at moderate n (Section VII reports it: it reduces the reachable-zero flag rate from 0.39 to 0.27 at unchanged size on unreachable zeros) and is the cleaner implementation when T permits.

B. Estimation theory for the first-order estimator

Theorems 1–2 are population statements; the following closes the gap to Algorithm 1. Treat the gain path as a deterministic design (Assumption 1) with nondegenerate limits: the standardized design satisfies $T^{-1} \tilde{\mathbf{Z}}^\top \mathbf{M}_X \tilde{\mathbf{Z}} \rightarrow \mathbf{Q} \succ 0$ with $\tilde{\mathbf{Z}}$ the interaction block and $\mathbf{M}_X$ the main-effect residualizer.

Theorem 4 (Consistency and asymptotic normality). *Let Assumptions 1–2 hold, strengthened so that $\{(\mathbf{r}_{1,t}, \mathbf{v}_t, \mathbf{w}_t)\}$ is α -mixing with coefficients satisfying $\sum_k \alpha(k)^{\delta/(2+\delta)} < \infty$ and $\mathbb{E}\|\cdot\|^{4+2\delta} < \infty$ for some $\delta > 0$. Assume further a bounded deterministic design whose lagged design-product averages converge (so that long-run limits exist for the triangular score array), a uniform stability margin $\sup_t \rho(\mathbf{L}_t) \leq \bar{\rho} < 1$. The CLT below is stated for the score array centered at its date-specific means, $\mathbf{s}_{t,T} - \mathbb{E}[\mathbf{s}_{t,T}]$ —a triangular-array formulation, because under the misspecified pooled projection the omitted cross-interactions generally give*

Algorithm 1 Output-only network identification from known gains

Require: outputs $\{\mathbf{r}_{1,t}, \mathbf{r}_{\text{on},t}\}_{t=1}^T$, gains $\{\gamma_t\}$, reversal share θ , size α

- 1: **if** any s_{γ_j} is zero or negligible **then** declare column j unidentified (Theorem 1(ii)) and drop it **end if**
- 2: standardize remaining gain paths $\tilde{\gamma}_{j,t} \leftarrow (\gamma_{j,t} - \bar{\gamma}_j)/s_{\gamma_j}$; form $\mathbf{G} = T^{-1} \tilde{\mathbf{r}} \tilde{\mathbf{r}}^\top$
- 3: report $\lambda_{\min}(\hat{\mathbf{Q}})$ of the residualized interaction Gram matrix as the identification-strength diagnostic (abstain if below a preset floor), with $\text{cond}(\mathbf{G})$ as the secondary design diagnostic (Theorem 1(iii))
- 4: **for** $i = 1, \dots, n$ **do**
- 5: OLS of (22); collect d_{ij} with HC errors for independent designs, HAC or moving-block-bootstrap errors for dependent data; form t_{ij}
- 6: **end for**
- 7: $\hat{\phi}_{ij}^{(1)} \leftarrow -d_{ij}/(\theta s_{\gamma_j})$ (first-order/low-gain estimate; exact recovery is the second-stage inversion of Sec. VI-D); flag couplings with $t_{ij} < -z_{1-\alpha}$ (marginal, exploratory tests—see multiplicity note below)
- 8: assemble $\hat{\mathbf{L}}_t = \hat{\Phi}^{(1)} \text{diag}(\gamma_t)$ with nonnegative projection $\hat{\mathbf{L}}_t \leftarrow \max(\hat{\mathbf{L}}_t, 0)$; $\hat{\mathbf{M}}_t = (\mathbf{I} - \hat{\mathbf{L}}_t)^{-1} - \mathbf{I}$
- 9: **return** $\hat{\Phi}^{(1)}$, flags, screening traces $\{\rho(\hat{\mathbf{L}}_t)\}$, $\{\hat{M}_{ij,t}\}$

the scores nonzero, date-dependent deterministic means—with the pseudo-true coefficient defined by the aggregate normal equations, $T^{-1} \sum_t \mathbb{E}[s_{\gamma_j}] \rightarrow \mathbf{0}$; the HAC estimator is applied to the demeaned scores, and its bandwidth conditions are assumed to hold for the centered array. Let $\hat{\mathbf{d}}_i$ collect the interaction coefficients for receiver i and $\mathbf{d}_i^*$ their pseudo-true values, which satisfy $d_{ij}^* = -\theta \phi_{ij} s_{\gamma_j} + O(\|\mathbf{L}\|^2)$. Then, by the Frisch–Waugh–Lovell reduction to the residualized interaction block,

$$\sqrt{T}(\hat{\mathbf{d}}_i - \mathbf{d}_i^*) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \mathbf{Q}^{-1} \boldsymbol{\Omega}_i \mathbf{Q}^{-1}), \quad (23)$$

where $\mathbf{Q}$ is the residualized interaction information matrix defined above and $\boldsymbol{\Omega}_i$ is the long-run covariance of the residualized interaction score $\tilde{\mathbf{z}}_t \varepsilon_{i,t}$, consistently estimable by HAC smoothing with standard bandwidth conditions [34] or a design-preserving block bootstrap (see Theorem 5). Consistency is for the pseudo-true parameter; its distance to the structural first-order target is the projection bias of Section V, of order $\sup_t \|\Phi \text{diag}(\gamma_t)\|^2$ with a constant that grows as $\lambda_{\min}(\mathbf{Q}) \rightarrow 0$ —a modeling error, not a statistical one, and it does not shrink with T .

Proof sketch: Multiplying stationary mixing signals by a bounded deterministic gain design yields a triangular mixing array of scores, not a stationary sequence; the assumed lag-product design limits give existence of the long-run covariance, centering at the date-specific means makes the array mean zero by construction (while the aggregate normal equations pin the pseudo-true coefficient), and the CLT for mixing arrays [35, Thm. 5.20] applies to the centered residualized interaction score. Frisch–Waugh–Lovell reduces the full regression to the interaction block with information matrix $\mathbf{Q}$,

yielding (23) with the sandwich form; HAC consistency under the same mixing, moment, and bandwidth conditions is [34]. The pseudo-true gap statement is the entrywise expansion (12) evaluated under the projection, whose remainder collects the omitted cross-interaction terms of Section VI. ■

Two design corollaries matter in practice, and both are verified numerically. First, under a *fixed* gain design the $\sqrt{T}$ rate is visible end to end: tripling T over the sequence $T \in \{250, 750, 2250, 6750\}$ (120 replications each) shrinks the RMSE to the pseudo-true parameter by factors 1.75/1.74/1.68 against the theoretical $\sqrt{3} \approx 1.73$. Second, under *random* gain designs the pseudo-true parameter is itself design-dependent, so unconditional convergence stalls at the design variability floor—the practical reading is that inference should condition on the realized gain path, exactly as the fixed-design frame of Theorem 4 does.

C. Monitoring statistics and dynamic threshold

Three alarms follow from the assembled matrices, one per risk object. *Cycle margin:* $\rho(\hat{\mathbf{L}}_t)$ against the static boundary 1 and the dynamic boundary below. *Transmission:* off-diagonal resolvent entries $\hat{M}_{ij,t}$, the displacement imported by channel i per unit of channel j 's innovation. *Persistence:* chaining periods through the partial correction, the uncorrected displacement $\mathbf{e}_t$ evolves as

$$\mathbf{e}_{t+1} = [(1 - \theta)\mathbf{I} - \theta\mathbf{M}] \mathbf{e}_t + \mathbf{M} \varepsilon_{t+1}, \quad (24)$$

whose derivation is: a share θ of the carried displacement is corrected (leaving $(1 - \theta)\mathbf{e}_t$), and the new clearing window adds $\mathbf{M}$ applied to the period innovation net of the correction flow, $\varepsilon_{t+1} - \theta\mathbf{e}_t$. For $\lambda \in \sigma(\mathbf{L})$ the matching chain eigenvalue is

$$1 - \theta - \theta \frac{\lambda}{1 - \lambda} = 1 - \frac{\theta}{1 - \lambda}, \quad (25)$$

so with a real spectrum and $\rho(\mathbf{L}) < 1$ the chain is covariance stationary iff $|1 - \theta/(1 - \lambda)| < 1$ for all λ ; for $\lambda \in [0, 1)$ this is $\lambda < 1 - \theta/2$, negative eigenvalues satisfy it automatically for $\theta \in (0, 1]$, and the Perron root binds. The operative stability boundary for monitoring is therefore $\rho(\mathbf{L}) < 1 - \theta/2$, strictly tighter than the within-period boundary—at $\theta = 0.88$ (the case-study calibration), 0.56 versus 1. Three restrictions delimit this statement: it freezes $\mathbf{L}$ over the horizon (for genuinely time-varying gains, stability is governed by products of transition matrices, for which pointwise spectral conditions are neither necessary nor sufficient in general); it assumes a common scalar correction θ ; and the reduction to $1 - \theta/2$ uses a real spectrum. It is a monitoring heuristic for slowly varying $\mathbf{L}_t$, not a general stability theorem.

Remark 2 (Network-analytic reading). *The three monitoring objects have close counterparts in network analysis. The resolvent $(\mathbf{I} - \mathbf{L})^{-1} = \sum_k \mathbf{L}^k$ is the walk-generating kernel of Katz's status index [28], so the transmission entry M_{ij} is a Katz-type walk influence of channel j on channel i (a communicability entry; Katz's index proper aggregates these entries per node). The left and right Perron vectors $\mathbf{u}, \mathbf{v}$ that already enter the delta method (26) are the corresponding influence and susceptibility centralities, and for a simple Perron root*

Algorithm 2 Rolling spectral and transmission monitoring

Require: data $\{\mathbf{r}_{1,t}, \mathbf{r}_{\text{on},t}, \gamma_t\}$, calibration θ , window W , thresholds $\bar{\rho}$ (default $1 - \theta/2$; heuristic when its frozen- $\mathbf{L}$ /scalar- θ /real-spectrum conditions fail), $\bar{m}$, exceedance count K_{run} , size α , bootstrap block length b

- 1: **for** each period $t \geq W$ **do**
- 2: run Algorithm 1 on window $t - W + 1, \dots, t$
- 3: (spectral branch:) $\mathcal{A}_\rho(t) \leftarrow \mathbb{I}\{\rho(\hat{\mathbf{L}}_t^{(2)}) > \bar{\rho}_t^*\}$, where $\hat{\mathbf{L}}_t^{(2)} = \hat{\Phi}^{(2)} \text{diag}(\gamma_t)$ uses the inversion (27)—the estimator whose bootstrap calibration Theorem 5 covers; the experimental SVD-slab point estimator of Sec. VI-D may be reported alongside but carries no inferential claim—on the support selected by the *model-selection threshold* c_T of Theorem 5 (not the fixed- α screen of line 7), and $\bar{\rho}_t^*$ is a design-preserving block-bootstrap basic-interval critical value for $H_0 : \rho^{(2)*} \leq \bar{\rho}$ —inference on the pseudo-true functional $\rho^{(2)*}$, whose distance to the structural ρ is the secant/projection gap of Sec. VI-D (90% empirical coverage there; pointwise validity in Theorem 5, uniform validity open); $\mathcal{A}_m(t) \leftarrow \mathbb{I}\{\exists i \neq j : \hat{M}_{ij,t} > \bar{m} \text{ and } t_{ij} < -z_{1-\alpha}\}$
- 4: raise alarm if $\mathcal{A}_\rho$ or $\mathcal{A}_m$ persists for K_{run} consecutive windows
- 5: **end for**

with $\gamma_j > 0$ the correspondence is exact: $\partial\rho(\Phi \text{diag} \gamma)/\partial\gamma_j = \rho u_j v_j / (\gamma_j \mathbf{u}^\top \mathbf{v})$ under the normalization $\mathbf{u}^\top \mathbf{v} > 0$, so the product $u_j v_j$ ranks channels by the elasticity of the spectral margin to their gains, giving the monitor a principled answer to which node drives an approaching alarm. And the dynamic boundary $\rho(\mathbf{L}) < 1 - \theta/2$ is a limited algebraic analogy to an epidemic threshold in network spread models—spectral radius of the contact structure against a recovery rate, here the overnight correction share θ —valid only under the frozen- $\mathbf{L}$, common scalar- θ , real-spectrum assumptions of the dynamic threshold, not an epidemic-threshold identity. Beyond interpretation, this correspondence points at the natural regularization class for moderate and large n : network-analytic structure on Φ (low-rank latent positions, block/community structure) reduces the parameter count from n^2 and correspondingly weakens the gain-richness demanded by the rank condition of Theorem 2—a quantitative structure-for-data exchange we leave to future work.

Algorithm 2 packages the three alarms with a consecutive-exceedance rule.

Entrywise screening over n^2 edges and overlapping windows requires familywise or false-discovery control, and the consecutive-exceedance count K_{run} trades average run length against detection delay; calibrating both is part of the sequential design, not an afterthought, and is included in the UQ program below.

D. Uncertainty quantification for spectral alarms

The alarms of Algorithm 2 are nonlinear functionals of noisy regression coefficients, and a point estimate of $\rho(\hat{\mathbf{L}})$

without an error bar is not an implementable statistic. Three layers are involved.

(a) *From coefficients to entries.* The interaction estimates $\hat{d}_{ij}$ carry an HC covariance from (22); entry estimates inherit it through the linear map of Algorithm 1.

(b) *From entries to ρ .* For an irreducible nonnegative matrix the Perron root is a simple real eigenvalue; with left and right Perron vectors $\mathbf{u}, \mathbf{v}$, first-order eigenvalue perturbation gives

$$\frac{\partial \rho}{\partial \ell_{ij}} = \frac{u_i v_j}{\mathbf{u}^\top \mathbf{v}}, \quad \text{se}^2(\hat{\rho}) \approx \sum_{i,j} \left(\frac{u_i v_j}{\mathbf{u}^\top \mathbf{v}} \right)^2 \text{Var}(\hat{\ell}_{ij}), \quad (26)$$

with the obvious quadratic form under correlated entry errors.

(c) *The plug-in trap, quantified.* Because the first-stage estimand is the resolvent sensitivity (20), entrywise estimates assembled directly into a matrix are *network-biased*: indirect paths inflate entries, and nonnegative projection converts zero-entry noise into positive mass that accumulates in the spectral radius. In the design of Section VII (true $\rho = 0.30$), the naive plug-in gives $\mathbb{E}[\hat{\rho}] = 0.73$; sparsifying by the signed t -rule does not repair it (0.72), because the bias sits in the *detected* entries. As an *ad hoc bias correction*—the global interaction slope is a secant across gain levels rather than a derivative at a specified operating point, so this inversion is heuristic rather than an exact second stage—the fixed-point iteration

$$\hat{\Phi}^{(2)} \leftarrow (\mathbf{I} - \hat{\mathbf{L}}) \hat{\mathbf{S}} \text{diag}(\hat{h})^{-1}, \quad \hat{h}_j = [(\mathbf{I} - \hat{\mathbf{L}})^{-1}]_{jj}, \quad (27)$$

evaluated at the mean gain—formally, $\hat{\Phi}^{(2)}$ solves the estimating equation $\Phi = (\mathbf{I} - \Phi \text{diag}(\hat{\gamma})) \hat{\mathbf{S}} \text{diag}(\hat{h}(\Phi))^{-1}$ on the selected support, with the nonnegativity projection assumed inactive at the solution (strict interiority)—where $\hat{\mathbf{S}}$ collects the sensitivity estimates; it recovers entries well (mean absolute error 0.042 on Φ) and halves the spectral bias ($\mathbb{E}[\hat{\rho}] = 0.47$); restricting to the t -detected support gives 0.42, and even the oracle support retains +0.07 bias with delta-method coverage of only 61% (anti-conservative), because (26) ignores the support-selection step and the nonneg projection. The rank-one slab estimator of Section V-A sharpens the point-estimation frontier. Implemented as: rescale the estimated full-interaction slab to $\hat{\mathbf{P}}_l = -\hat{\mathbf{D}}_l / (\theta s_{\gamma_l})$ (so that in population $\hat{\mathbf{P}}_l$ targets the secant analogue of $\partial \mathbf{M} / \partial \gamma_l$; the scale-pinning identity is exact only for the local derivative, hence heuristic here), project to rank one by SVD—the singular-vector sign ambiguity is resolved by orienting the pair so that the pinned diagonal entry of the projected slab is nonnegative—then pin the scale by $x(1+x) = c_l$ with $c_l = \gamma_l [\hat{\mathbf{P}}_l]_{ll}$ read from the rank-one-projected slab: real roots require $c_l \geq -1/4$, and the branch consistent with the maintained nonnegativity, $x = (-1 + \sqrt{1 + 4c_l})/2$, is admissible ($x \geq 0$) precisely when $c_l \geq 0$. *Root failure* is defined as $c_l < 0$: noise has pushed the projected diagonal outside the admissible cone. (For a channel whose true $M_{ll} = 0$ the scale is pinned instead by the $r_{ll} = 1$ normalization of Theorem 3(ii); the implementation does not attempt to distinguish a near-zero diagonal from noise and routes such columns through the same fallback.) Assemble the surviving columns into $\hat{\mathbf{M}}$, requiring every gain to vary with positive operating level. In this experiment (100 Monte Carlo paths, an exception to the 200 used elsewhere),

the scale root succeeded on 84 paths; *conditional on root success*, spectral bias is +0.096 with standard deviation 0.034 and Φ RMSE 0.056. Because root success may select easier samples, we report the matched and unconditional accounting explicitly: on the *same* 84 root-success paths the two-stage inversion has spectral bias +0.135 (sd 0.176) against the SVD slab's +0.096 (sd 0.034); over *all* 100 paths, the operational estimator—SVD slab with two-stage fallback on the 16 root-failure paths—has bias +0.098 (sd 0.058) against +0.131 (sd 0.169) for the two-stage inversion alone. The SVD slab is thus near the oracle-support benchmark (+0.07) without oracle knowledge, and the advantage survives the fallback mixture. We therefore label it the *preferred experimental point estimator in this design*: its calibrated inference is *not* established—the 90% bootstrap result above concerns the two-stage pipeline only, and extending Theorem 5 to the SVD branch would additionally require an isolated pseudo-true top singular value, a scale root bounded away from the failure boundary, and an asymptotically constant fallback branch, none of which we assume here. The exact-model nonlinear least-squares estimator suggested by Theorem 2 completes the picture from both ends: initialized at the truth it is nearly unbiased (Φ RMSE 0.029, spectral bias +0.04)—an *infeasible oracle diagnostic* showing the sample carries the information the local-identification theorem promises, not a usable benchmark—but initialized at the first-order estimates it lands on spectral-inflating ridges of the nonconvex criterion (Φ RMSE 0.153, spectral bias +0.66, worse than the two-stage inversion it was meant to refine). The natural first structure to exploit is separability: for fixed Φ the confound $\mathbf{C}$ enters the regime-map criterion linearly and can be profiled out in closed form, reducing the fit to Φ alone (variable projection [32]). We implement this reduced fit (criterion: least squares on the regime map $\mathbf{B}(\gamma)$ over the realized gain regimes; box constraints $\Phi \in [0, 2]^{n \times n}$; trust-region reflective solver; initialized at the first-order estimates): eliminating the nuisance dimension does not repair the geometry—the profiled and joint fits land in the same spectral-inflating basin (spectral bias +0.54 for both in this implementation, with a final criterion value roughly 34 $\times$ its value at the truth), so the nonconvexity is intrinsic to the Φ -dependence, not an artifact of the nuisance dimension. The Jacobian already derived for Theorem 2 supports Gauss–Newton and Levenberg–Marquardt steps on the reduced problem, and majorization-minimization or successive-convex-approximation surrogates [33] are the natural globalization devices; their convergence theory for this rational fixed-point criterion, or a characterization of its basins, is therefore the computational half of the open spectral-inference problem, alongside the statistical half above.

The practical prescription follows the numbers. For *transmission alarms*—entrywise, signed—the HC t -rule is correctly sized off the reachable set and deliberately liberal on it, the desired direction for a screening alarm. For *spectral alarms*, analytic intervals are anti-conservative at this sample size, and we therefore validate the resampling route directly: re-running the entire pipeline (interaction regression, signed selection, inversion (27), eigenvalue) inside a moving-block bootstrap over dates ($B = 99$, block length 25), the *basic*

(reversed-percentile) interval—whose reflection accounts for the estimator's bias, unlike the percentile interval—covers the true spectral radius on 90% of paths at nominal 95% (60 outer paths, $B = 99$; the coverage estimate itself carries a Monte Carlo standard error of about 4 percentage points), against 45–61% for the delta method; the naive percentile interval, centered on the biased estimate, covers only 55%. The empirically calibrated spectral alarm is therefore the two-stage point estimate (27) with basic block-bootstrap critical values. The following formalizes when this calibration is not merely empirical.

Theorem 5 (Bootstrap validity under consistent selection). *Let the conditions of Theorem 4 hold and assume additionally: (i) consistent support selection: the spectral stage uses a model-selection threshold c_T with $c_T \rightarrow \infty$, $c_T/\sqrt{T} \rightarrow 0$ (not the fixed- α screening rule of Algorithm 1, which by construction retains each null entry with probability $\approx \alpha$ and is not support-consistent), together with separation $\min_{(i,j) \in S^*} |S_{ij}^*| \geq c > 0$; then $\mathbb{P}(\hat{S} = S^*) \rightarrow 1$. (ii) The fixed-point map of (27) has a locally unique solution at the pseudo-true value with nonsingular Jacobian, and (iii) the Perron root of the inverted matrix is simple. (iv) The multiplier score array satisfies a bootstrap CLT conditional on the data: blockwise external multipliers applied to the estimated score blocks reproduce, conditionally, the limiting law of the centered score sum—the dependent-wild-bootstrap regime of [31], whose conditions we adopt for the triangular array. Then, for a design-preserving dependent-multiplier (block wild) bootstrap—multiplying the estimated score blocks by blockwise external multipliers and rebuilding the estimator along the fixed realized gain path, a scheme that preserves both the deterministic design and the gain-modulated conditional heteroskedasticity of the triangular score array (raw residual transplantation would not)—with block length $\ell \rightarrow \infty$, $\ell/T \rightarrow 0$, the bootstrap law of $\hat{\rho}^{(2)}$ consistently estimates its limiting law, and basic bootstrap intervals for $\rho^{(2)*}$ attain nominal coverage asymptotically.*

Proof sketch: Under (i) the thresholded selection event is eventually constant ($c_T \rightarrow \infty$ kills null entries, $c_T/\sqrt{T} \rightarrow 0$ plus separation retains true ones), so with probability approaching one the estimator equals the fixed-selection estimator, a composition of OLS coefficients with the smooth maps (27) (smooth by (ii), via the implicit function theorem) and a simple-eigenvalue functional (smooth by (iii), standard perturbation theory). The delta method transfers the CLT of Theorem 4; the dependent-multiplier score bootstrap rebuilds the estimating equations from multiplier-weighted score blocks along the fixed gain path, reproducing the same triangular-array structure (including its gain-modulated heteroskedasticity) conditional on the design; conditional distributional consistency of the multiplier score sum is condition (iv) (the dependent wild bootstrap of [31], with classical blocking arguments [29] and estimating-equation extensions [30, Ch. 4] supplying the smooth-functional transfer). Basic-interval validity follows from distributional consistency and continuity of the limit law. ■

The boundary of Theorem 5 explains the simulated 90%:

TABLE II
SYNTHETIC VALIDATION AND BENCHMARKS.

Scenario	estimator	power	size/FA	RMSE
A (corr. γ , $\kappa \approx 247$)	ours	1.00	0.22	0.28
A2 (indep. γ , $\kappa \approx 12$)	ours	1.00	0.12	—
B (constant γ)	ours	unidentified flagged, 100% of paths		
C (excl. violated)	ours	1.00	0.04	0.34
D (neg. confound)	ours	1.00	0.12	—
A (pos. confound)	level/Granger	1.00	0.04	0.09
D (neg. confound)	level/Granger	1.00	0.53	—
A (2-regime split)	variance-ratio [‡]	—	—	43.4
A (tuned [†])	variance-ratio [‡]	—	—	0.60

Note: power = fraction of true couplings with $t < -1.645$; size/FA = same rule on zero-coupling entries; RMSE over all off-diagonal entries. The level/Granger benchmark regresses post-window on pre-window outputs without gain interactions; the variance-ratio benchmark ([‡]an illustrative two-regime $\Delta\text{Cov}/\Delta\text{Var}$ construction in the spirit of [13], not a full heteroskedasticity-identification implementation) computes $\Delta\text{Cov}/\Delta\text{Var}$ across a two-regime split of each sender’s gain. [†]Tuned in the benchmark’s favor: best regime pair per sender, regularized denominators (skipping variance differences below 20% of the sender’s variance), median across admissible pairs—even so, RMSE is twice ours and 22% of paths yield no admissible estimate. Dashes: not recorded for that cell.

the simulation uses the fixed- α screening rule and pairwise tuple resampling (not the thresholded selection and dependent-multiplier score scheme of the theorem), several true sensitivities sit near the threshold at moderate T , selection randomness enters the sampling distribution, and coverage dips below nominal—the gap between the implemented heuristic and the theorem’s conditions is thus itself measured. A uniform theory over local-to-zero couplings (selective-inference regime) remains open—but it is now the *only* open layer, and the target of the coverage statement ($\rho^{(2)*}$, the pseudo-true spectral radius) is stated explicitly. Theorem 5 gives formal validity of a design-preserving variant under thresholded selection; uniformity over local-to-zero couplings and the detection-delay theory of Algorithm 2 (measured empirically in Section VII) remain open.

VII. SYNTHETIC VALIDATION

All experiments use $n = 5$, $T = 750$, 200 Monte Carlo paths, $\theta = 0.85$ known, a sparse nonnegative Φ with four true off-diagonal couplings, $\rho(\mathbf{L}_t) \in [0.3, 0.55]$, a common factor in $\mathbf{r}_1$, and a constant lead-lag confound $\mathbf{C}$ with nonzero entries placed on *zero-coupling* pairs—the adversarial placement. Gains take three levels $\{0.5, 1.0, 1.5\}$; the decision rule is the signed t -test at 5% (Algorithm 1). Scenarios: **A** staggered staircase gains (columns strongly correlated across channels, Gram condition number ≈ 247); **A2** independent regime-switching gains (≈ 12); **B** constant gains (rank violation); **C** exclusion violation ($\mathbf{C}_t$ co-moving with the mean gain, loading 0.8); **D** as A2 with the confound sign flipped (reversal-mimicking). Table II reports the grid.

Five findings. (i) *Sharpness*. Under constant gains the interaction regressor is degenerate and the estimator reports non-identification on every path—the failure predicted by Theorem 1(ii) occurs exactly at the theorem’s boundary, and is *detectable from the data*, not silent. (ii) *Gram conditioning*. At equal per-channel gain variances, moving from correlated

staircase paths to independent regimes drops the false-alarm rate from 0.22 to 0.12: identification strength is a property of the joint design—and precisely because the simulated signals are strongly factor-driven (Σ_r near rank one), the Hadamard structure of Theorem 1(iii) puts the gain-Gram conditioning in charge of the variance inflation. (iii) *Confound immunity is signed*. The level/Granger benchmark is fine when the confound happens to have the continuation sign (FA 0.04) and breaks completely when it mimics reversal (FA 0.53); the interaction estimator’s FA is unchanged (0.12) because a constant confound of either sign cannot track γ_t . (iv) *Duality with heteroskedasticity identification*. On data generated with constant shock variances, the Rigobon-style estimator divides by near-zero variance differences and returns RMSE two orders of magnitude above ours (43.4 vs. 0.28), and tuning it in its own favor still leaves RMSE at twice ours with 22% of paths producing no admissible estimate—the numerical face of the duality of Section II. (v) *The estimand is the resolvent, and the report separates three entry classes*. True direct edges, zero-direct entries that are network-reachable, and unreachable pairs are different objects: at the baseline gain level the detector flags 2-hop-reachable zeros at 39% while holding size statistically indistinguishable from nominal (5.3% Monte Carlo estimate at the 5% rule) on unreachable zeros. Whether reachable-zero flags count as detections (transmission screening) or false positives (direct-edge recovery) depends on the declared target; only unreachable pairs are nulls for both. A gain-strength sweep ties the flag rate to the theory: at $\rho(\mathbf{L}) \approx 0.15$ reachable-zero flags fall to 7% (the first-order regime, where the resolvent is close to the direct coupling), at ≈ 0.30 they are 39%, and at ≈ 0.45 they reach 84% with unreachable-pair size drifting to 14%—the quantitative footprint of higher-order paths, and the reason spectral assembly needs the second-stage inversion of Section VI-D. The full n^2 -interaction regression reduces the reachable-zero rate to 27% at unchanged unreachable size. In scenario C, the particular gain-tracking confound simulated (loading 0.8 on the mean gain path) inflates magnitude RMSE (0.28 $\rightarrow$ 0.34) while the signed detection size happens to remain low; no general robustness to exclusion failure is claimed—a reversal-signed gain-tracking confound would defeat the detector by construction, which is why Assumption 1 is maintained rather than tested.

A. Operating characteristics of the rolling monitor

Algorithm 2’s transmission alarm is a sequential procedure over $n(n-1)$ edges and overlapping windows, so its per-path false-alarm rate is governed by the joint design, not the entrywise size. We measure it directly—for the *transmission branch only*; the spectral branch is not exercised: coupling switches on at $\tau = 600$ of $T = 1000$ ($W = 250$, step 5; null arm has diagonal-only Φ ; 40 paths per arm). Table III reports the frontier under explicit procedural rules (see the table note): even at the stringent entrywise $z = 2.576$ the null-arm per-path false-alarm rate is 0.90 (the multiplicity of 20 edges and 151 overlapping windows made explicit), while raising the threshold and lengthening the run rule brings it to 0.10—at

TABLE III
ROLLING-MONITOR OPERATING CHARACTERISTICS.

rule (z, K_{run})	FA (null)	FA (pre-onset)	med. delay	90% delay	miss
2.576, 2	0.90	0.75	208	266	0/40
3.09, 2	0.28	0.20	255	295	4/40
3.09, 4	0.23	0.03	265	313	5/40
3.72, 2	0.15	0.00	283	377	16/40
3.72, 3	0.10	0.00	278	371	18/40

Note: transmission branch only. Procedural rules: an alarm qualifies when the edgewise signed exceedance holds at K_{run} consecutive window ends (151 overlapping ends 250, 255, ..., 1000). FA (null) = fraction of null-arm paths raising any qualifying alarm; FA (pre-onset) = fraction of treated paths raising a qualifying alarm declared before the onset τ (counted as false alarms, not detections); delay = first qualifying window end at or after τ , minus τ (a run straddling τ counts by its declaration end); miss = treated paths with no post-onset qualifying alarm within the horizon. 40 Monte Carlo paths per arm, so each FA estimate carries a binomial standard error of about 0.05–0.08; delay quantiles at the two most stringent rules are computed on detected paths only.

the cost the frontier makes visible, 18/40 missed detections at the most stringent rule within this horizon, against 0/40 at the loosest. Detection delay and missed detections trade off against false alarms along the whole frontier; sequential calibration—choosing (z, K_{run}) for a target average run length—is thereby a tabulated design choice rather than an afterthought.

VIII. CASE STUDY: LEVERAGED-FUND REBALANCING

The motivating instance of the model is a market hosting leveraged exchange-traded funds. A fund with leverage multiple m and assets A must trade $A(m^2 - m)r$ near the close in the direction of the day's return r ; aggregating over funds on underlying j gives rebalancing capital $K_j = \sum_f A_f(m_f^2 - m_f)$, and $\gamma_{j,t} = K_{j,t}/\text{ADV}_{j,t}$ —disclosed daily through fund shares, NAVs, and multiples—is exactly the known actuation gain of (1). The clearing window is the closing auction; the reversal moment is the overnight correction of transient displacement, with θ calibrated externally at 0.88 [24]; the confound $\mathbf{C}$ is the lead-lag structure of correlated equities [36]. One data limitation must be stated at the outset: intraday bars are available only post-launch, so the application's regressor is the close-to-close *full-period* return, not the pre-window r_1 the clean theory requires; by (11) the case study is therefore *contaminated screening*—the signed detection and its gain scaling are reduced-form empirical patterns whose structural reading is convention-dependent (under the full-period convention even a static confound acquires gain dependence through $(\mathbf{I} + \mathbf{M}_t)^{-1}$, so no invariance is claimed), and structural readings of $\hat{\Phi}$ and $\rho(\hat{\mathbf{L}})$ carry the $O(\|\mathbf{M}\|)$ convention dependence, which the companion paper addresses by reporting both polar mappings. Full institutional detail, inference, and robustness are in [23]; here we illustrate the regression/screening mechanics of Algorithms 1–2; the identification guarantees require the pre-window moment and do not transfer verbatim. The sample freeze date for all reported Korean numbers is August 21, 2026; the U.S. panel is final (CRSP coverage ends December 2024).

Korea (2026). Sixteen single-stock LETFs on Samsung Electronics and SK Hynix launched May 27, 2026, into one closing auction—a two-channel system with disclosed,




Fig. 2. Observed actuation gains $\gamma_{j,t}$ of the two Korean complexes. Note: constructed daily from disclosed fund shares, NAVs, and leverage multiples over trailing 20-day traded value; entered at $t - 1$. Dashed: product launch; dotted: the deposit regulation, which collapsed turnover while leaving the gains—the identification resource—intact.




Fig. 3. The rolling detector on the Korean data (receiver: Samsung; sender: Hynix).

Note: top—rolling 40-day cross-slope with 95% bands; bottom—launch-break contrast z at weekly placebo break dates and the true launch (dashed). Post windows left of the launch contain increasing fractions of true post-launch observations.

strongly time-varying gains (γ of order one; Figure 2 shows the observed gain paths, the empirical realization of the known modulating sequence of Figure 1). The interaction moment detects one-directional coupling from the large complex (Hynix, worldwide capital peak KRW 37.5tn) into the small one (Samsung, 6.6tn): the launch-break contrast of the cross-reversal slope is $z = -2.82$ (Newey–West -2.72 ; leave-one-day-out range $[-3.05, -2.64]$), more negative than all 182 ordered placebo pairs of fourteen non-treated large caps (rank $1/183 = 0.0055$), with a ten-day block-bootstrap interval excluding zero. Figure 3 shows the rolling detector of Algorithm 2 on these data: the cross-slope is stable through the pre-launch period and turns only as post-launch observations enter the window, with fourteen weekly placebo break dates all yielding $|z| \leq 1.9$.

A reduced-form pattern consistent with the gain-scaling logic of Theorem 1 (not a validation of it, given the contami-

TABLE IV
U.S. STAGGERED PANEL: CROSS-REVERSAL INTERACTION GRID (FINAL 2022–2024 SAMPLE).

direction	$\hat{b}_{R_A}(t)$	$\hat{b}_{\gamma_A \times R_A}(t)$	N
TSLA→NVDA	+0.007 (+0.2)	+0.005 (+0.1)	500
TSLA→AAPL	+0.005 (+0.4)	+0.004 (+0.6)	500
NVDA→TSLA	+0.017 (+1.5)	−0.065 (−1.6)	500
NVDA→AAPL	−0.005 (−1.3)	+0.016 (+1.1)	500
AAPL→TSLA	+0.086 (+0.8)	−0.163 (−1.77)	500
AAPL→NVDA	−0.470 (−0.9)	−0.211 (−0.8)	500

Note: receiver’s overnight return on sender’s day return R_A , own return, leave-out equal-weight control (GOOGL/MSFT/AMZN), QQQ, the sender’s standardized lagged gain $\tilde{\gamma}_A$, and the interaction $\tilde{\gamma}_A \times R_A$; HC standard errors. Under the entrywise signed 5% rule the single cell AAPL→TSLA ($t = -1.77$) is a marginal flag—about what twelve statistics produce by chance at 5%—and no cell survives multiplicity adjustment; the grid is null, the graded-response prediction for a low-impact closing venue.

nated regressor) appears: the interaction of the sender’s return with its standardized lagged gain is -0.106 ($t = -2.29$) with the standalone gain main effect insignificant—the coupling tracks γ_t , which a static lead–lag cannot mimic. The assembled empirical baseline (reverse link set to zero, as it is not detected) is triangular, so its spectral radius $\rho = 0.610$ is driven entirely by the imported diagonal and is unaffected by the detected cross channel; transmission is the operative alarm, with $\hat{M}_{21} \approx 0.22$ under the baseline convention: roughly a fifth of the sender’s innovation is imported into the receiver’s close, invisible to the receiver’s own gain (0.24). Read against the dynamic boundary (25)—as a screening trace, not a calibrated threshold test (Section VI-D)—the sender diagonal imported from the companion calibration (0.61) exceeds $1 - \theta/2 = 0.56$, while the self-contained estimate from our own sample’s own-reversal moment (0.42, $z = -2.0$) does not: the boundary comparison is calibration-dependent and is reported as such.

United States (2022–2024, final sample). Two U.S. panels provide the scale placebo and a live diagnostic of Theorem 1(iii). The MicroStrategy–Bitcoin–Coinbase complexes form an approximately symmetric configuration with liquidity-scaled capital comparable to Korea (γ_{MSTR} up to 1.72); all six treated directions are null ($|z| \leq 1.45$), as are placebos. A second, *staggered* panel—the TSLA, NVDA, and AAPL single-stock complexes, whose fourteen funds launched at six distinct dates over 2022–2023—adds two findings on final (2022–2024) data. First, the full ordered-pair interaction grid is null after multiplicity adjustment (one marginal cell at $t = -1.77$ among twelve recorded statistics; Table IV), consistent with the graded response surface: gains reach 0.53–0.54 but the venue impact is low. Second, and diagnostically: despite six distinct launch dates, the gain *levels* are correlated 0.81–0.91 across complexes because post-launch capital growth trends are common, and the rolling-window Gram condition number has median 45 (deciles 17–118)—staggered onsets do *not* deliver uncorrelated gain paths (Figure 4), so the conditioning diagnostic of Theorem 1(iii) and Algorithm 1 must be evaluated on the estimation window, not inferred from institutional timing. Table IV reports the full grid.

The null is the response surface of the model, not a failure: an estimator that found coupling wherever assets correlate




Fig. 4. U.S. staggered panel: gain paths and identification conditioning. Note: top—liquidity-scaled complex gains $\gamma_{j,t}$ for the TSLA, NVDA, and AAPL single-stock LETF complexes (CRSP, 2022–2024); fund launches occur at six distinct dates, yet post-launch capital growth makes the paths co-trend (level correlations 0.81–0.91). Bottom—rolling 250-day condition number of the standardized gain-path Gram matrix (log scale; dotted line: median 45), the secondary design diagnostic; the primary diagnostic $\lambda_{\min}(\hat{\mathbf{Q}})$ additionally involves the signal covariance (Theorem 1(iii)) and is reported with the estimation windows of the companion replication files.

would have failed this scale placebo.

IX. CONCLUSION

Known time-varying actuation, combined with signal excitation and conditional orthogonality (Assumption 2), identifies coupled feedback networks from outputs alone: the gain path plays the role that references, probing signals, and designed precoders play elsewhere—under gain exogeneity (Assumption 1): the confound must not track the disclosed gains—and a partial-reversal moment converts the modulated response into a signed estimator robust to gain-invariant confounds. The identification condition is a rank—sharply, a conditioning—property of the joint gain design; the estimand is the resolvent sensitivity—directly what transmission screening requires, and the first-stage input to spectral-margin recovery; and the method’s principal failure modes differ in kind: constant-gain non-identification is detectable from the data, while a reversal-signed gain-tracking confound is observationally indistinguishable and remains the maintained assumption’s honest price. The framework’s natural next steps are the exact nonlinear estimation theory with latent-input contamination and channel-specific correction shares—including surrogate-based (majorization-minimization) globalization of the exact-model fit, whose nonconvexity is intrinsic to the coupling dependence rather than to the nuisance dimension (Section VI-D)—regularized estimation for moderate and large n , and closing the remaining spectral-inference gaps—uniform (local-to-zero) validity, fixed- α selection, and sequential calibration—beyond the pointwise validity proved in Theorem 5 and the $\sqrt{T}$ asymptotics of Theorem 4; on the application side, any self-referencing mandate with disclosed capacity—option hedging programs, volatility targeting, demand response—is a candidate network.

REFERENCES

- [1] L. Ljung, *System Identification: Theory for the User*, 2nd ed. Prentice Hall, 1999.
- [2] U. Forssell and L. Ljung, “Closed-loop identification revisited,” *Automatica*, vol. 35, no. 7, pp. 1215–1241, 1999.
- [3] P. M. J. Van den Hof, A. Dankers, P. S. C. Heuberger, and X. Bombois, “Identification of dynamic models in complex networks with prediction error methods—basic methods for consistent module estimates,” *Automatica*, vol. 49, no. 10, pp. 2994–3006, 2013.
- [4] H. H. M. Weerts, P. M. J. Van den Hof, and A. G. Dankers, “Identifiability of linear dynamic networks,” *Automatica*, vol. 89, pp. 247–258, 2018.
- [5] X. Cheng, S. Shi, I. Lestas, and P. M. J. Van den Hof, “Identifiability in dynamic acyclic networks with partial excitations and measurements,” *IEEE Trans. Autom. Control*, 2023, arXiv:2201.07548.
- [6] J. M. Hendrickx, M. Gevers, and A. S. Bazanella, “Identifiability of dynamical networks with partial node measurements,” *IEEE Trans. Autom. Control*, vol. 64, no. 6, pp. 2240–2253, 2019.
- [7] J. Gonçalves and S. Warnick, “Necessary and sufficient conditions for dynamical structure reconstruction of LTI networks,” *IEEE Trans. Autom. Control*, vol. 53, no. 7, pp. 1670–1674, 2008.
- [8] D. Materassi and G. Innocenti, “Topological identification in networks of dynamical systems,” *IEEE Trans. Autom. Control*, vol. 55, no. 8, pp. 1860–1871, 2010.
- [9] R. Tóth, *Modeling and Identification of Linear Parameter-Varying Systems*. Springer, 2010.
- [10] V. Laurain, M. Gilson, R. Tóth, and H. Garnier, “Refined instrumental variable methods for identification of LPV Box–Jenkins models,” *Automatica*, vol. 46, no. 6, pp. 959–967, 2010.
- [11] C. Verhoeck, G. I. Beintema, S. Haesaert, M. Schoukens, and R. Tóth, “Deep-learning-based identification of LPV models for non-linear systems,” in *Proc. IEEE Conf. Decision and Control*, 2022, arXiv:2204.04060.
- [12] P. B. Cox and R. Tóth, “Linear parameter-varying subspace identification: A unified framework,” *Automatica*, vol. 123, 2021, arXiv:2008.03347.
- [13] R. Rigobon, “Identification through heteroskedasticity,” *Rev. Econ. Stat.*, vol. 85, no. 4, pp. 777–792, 2003.
- [14] E. Sentana and G. Fiorentini, “Identification, estimation and testing of conditionally heteroskedastic factor models,” *J. Econometrics*, vol. 102, no. 2, pp. 143–164, 2001.
- [15] M. Lanne and H. Lütkepohl, “Identifying monetary policy shocks via changes in volatility,” *J. Money, Credit and Banking*, vol. 40, no. 6, pp. 1131–1149, 2008.
- [16] D. J. Lewis, “Identifying shocks via time-varying volatility,” *Rev. Econ. Stud.*, vol. 88, no. 6, pp. 3086–3124, 2021.
- [17] L. Kilian and H. Lütkepohl, *Structural Vector Autoregressive Analysis*. Cambridge Univ. Press, 2017.
- [18] M. K. Tsatsanis and G. B. Giannakis, “Transmitter induced cyclostationarity for blind channel equalization,” *IEEE Trans. Signal Process.*, vol. 45, no. 7, pp. 1785–1794, 1997.
- [19] E. Serpedin and G. B. Giannakis, “Blind channel identification and equalization with modulation-induced cyclostationarity,” *IEEE Trans. Signal Process.*, vol. 46, no. 7, pp. 1930–1944, 1998.
- [20] W. A. Gardner, “Exploitation of spectral redundancy in cyclostationary signals,” *IEEE Signal Process. Mag.*, vol. 8, no. 2, pp. 14–36, 1991.
- [21] D. Materassi and M. V. Salapaka, “On the problem of reconstructing an unknown topology via locality properties of the Wiener filter,” *IEEE Trans. Autom. Control*, vol. 57, no. 7, pp. 1765–1777, 2012.
- [22] M. T. Angulo et al., “Fundamental limitations of network reconstruction from temporal data,” *J. R. Soc. Interface*, vol. 14, 2017, arXiv:1508.03559.
- [23] J. Woo, “The loop-gain matrix: Coupled rebalancing feedback and the blind spots of scalar stability monitoring,” arXiv preprint arXiv:2608.22768, 2026.
- [24] Y. Zhao, “Preying on leveraged ETFs,” working paper, arXiv:2608.03703v2, Aug. 2026.
- [25] F. Fnaiech and L. Ljung, “Recursive identification of bilinear systems,” *Int. J. Control*, vol. 45, no. 2, pp. 453–470, 1987.
- [26] T. Hastie and R. Tibshirani, “Varying-coefficient models,” *J. R. Statist. Soc. B*, vol. 55, no. 4, pp. 757–796, 1993.
- [27] B. Peeters and G. De Roeck, “Reference-based stochastic subspace identification for output-only modal analysis,” *Mech. Syst. Signal Process.*, vol. 13, no. 6, pp. 855–878, 1999.
- [28] L. Katz, “A new status index derived from sociometric analysis,” *Psychometrika*, vol. 18, no. 1, pp. 39–43, 1953.
- [29] H. R. Künsch, “The jackknife and the bootstrap for general stationary observations,” *Ann. Statist.*, vol. 17, no. 3, pp. 1217–1241, 1989.
- [30] S. N. Lahiri, *Resampling Methods for Dependent Data*. Springer, 2003.
- [31] X. Shao, “The dependent wild bootstrap,” *J. Amer. Statist. Assoc.*, vol. 105, no. 489, pp. 218–235, 2010.
- [32] G. H. Golub and V. Pereyra, “The differentiation of pseudo-inverses and nonlinear least squares problems whose variables separate,” *SIAM J. Numer. Anal.*, vol. 10, no. 2, pp. 413–432, 1973.
- [33] Y. Sun, P. Babu, and D. P. Palomar, “Majorization-minimization algorithms in signal processing, communications, and machine learning,” *IEEE Trans. Signal Process.*, vol. 65, no. 3, pp. 794–816, 2017.
- [34] W. K. Newey and K. D. West, “A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix,” *Econometrica*, vol. 55, no. 3, pp. 703–708, 1987.
- [35] H. White, *Asymptotic Theory for Econometricians*, revised ed. Academic Press, 2001.
- [36] A. W. Lo and A. C. MacKinlay, “When are contrarian profits due to stock market overreaction?” *Rev. Financial Stud.*, vol. 3, no. 2, pp. 175–205, 1990.