

Gatheral's Conjecture Revisited

Vladimir Lucic
Imperial College London
vlucic@ic.ac.uk

September 7, 2026

Abstract

We compare the Heston model with $\rho = -1$ to the one-dimensional local-volatility model calibrated to the same European option prices. We show that, for each fixed expiry $T > 0$, their integrated variances satisfy

$$I_T^H \prec_{\text{cx}} I_T^{\text{LV}}.$$

This strict ordering gives a Heston-model counterexample to the convex-order inequality conjectured by J. Gatheral.

1 Introduction

In 2005, Gatheral [12] formulated the conjecture that, among models calibrated to the same European option prices, local volatility minimizes the value of options on realized variance; see also Gatheral [13, p. 155]. This conjecture and its variants attracted considerable attention over the years and were supported by numerical evidence and partial results; see, for example, Brockhaus and Long [5], Gatheral [13], and Austing [2, Section 10.3.4]. However, Beiglböck, Friz, and Sturm [3] subsequently disproved the conjecture by providing two carefully constructed counterexamples. While their result settled the original claim, it still left open the question of whether commonly used stochastic volatility models for which the variance itself follows a Markov process can exhibit such behavior. We show that they can: in the Heston model with perfect negative price–variance correlation, the integrated variance is smaller in convex order than the integrated variance generated by its one-dimensional local-volatility projection. Thus, in this example, local volatility does not minimize variance-option prices; it increases them.

In the remainder of this section, we informally outline the main steps and ideas of the proof, deferring the full rigor to subsequent sections. First, we fix some notation. Throughout, $\mathbb{R}_+ := [0, \infty)$. For sets A and B , the notation $A \Subset B$ means that $\bar{A}$ is a compact subset of B . For integrable random variables X and Y , the notation $X \preceq_{\text{cx}} Y$ means

$$\mathbb{E}[\varphi(X)] \leq \mathbb{E}[\varphi(Y)]$$

for every convex function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ for which both expectations are finite, and $X \prec_{\text{cx}} Y$ holds when, in addition, $\text{Law}(X) \neq \text{Law}(Y)$.

Under zero interest rates and dividends, the stochastic-volatility model of Heston [16] is given by

$$\frac{dS_t}{S_t} = \sqrt{v_t} dW_t, \tag{1}$$

$$dv_t = \kappa(\theta - v_t) dt + \xi\sqrt{v_t} dZ_t, \tag{2}$$

$$d\langle W, Z \rangle_t = \rho dt.$$

Here W and Z are Brownian motions, $S_0, v_0, \kappa, \theta, \xi > 0$, and $\rho \in [-1, 1]$. We denote the integrated variance by

$$I_t^H := \int_0^t v_u \, du.$$

Its projected local variance is

$$\lambda(t, s) := \mathbb{E}[v_t \mid S_t = s],$$

and the associated local-volatility model and its accumulated variance are

$$\frac{d\bar{S}_t}{\bar{S}_t} = \sqrt{\lambda(t, \bar{S}_t)} \, dB_t, \quad I_t^{LV} := \int_0^t \lambda(u, \bar{S}_u) \, du.$$

This is called the “Markovian projection”. Brunick and Shreve [6, Corollary 3.7] provide at least one weak mimicking diffusion $\bar{S}$ satisfying this equation and having the same one-time laws as S ; their corollary alone asserts neither uniqueness nor the Markov property. Thus the two models generate the same European option prices. Moreover, I_T^H and I_T^{LV} have the same mean.

Henceforth, assume that $\rho = -1$, so that we may take $Z = -W$. This is the perfectly negatively correlated Heston model, also known as the Heston–Nandi model; see Heston and Nandi [17] and Cox and Wang [7]. The main result of this note is the following convex ordering for the Heston–Nandi model:

$$I_T^H \prec_{cx} I_T^{LV}, \quad T > 0. \quad (3)$$

Our argument rests on two key ingredients: a backward Duhamel comparison argument and a conditional form of Chebyshev’s covariance inequality. Perfect negative correlation is what makes the two fit together. Indeed, eliminating the common stochastic integral from (1)–(2) gives the pathwise identity

$$v_t = v_0 + \kappa\theta t - \xi \log(S_t/S_0) - \beta I_t^H, \quad \beta := \kappa + \frac{\xi}{2} > 0. \quad (4)$$

After conditioning on the current spot, this becomes

$$v_t - \lambda(t, S_t) = -\beta \left(I_t^H - \mathbb{E}[I_t^H \mid S_t] \right). \quad (5)$$

To see how this identity enters the option comparison, let Φ be nonnegative and convex and let

$$V(t, s, i) := \mathbb{E}_{t,s,i}^{LV}[\Phi(I_T^{LV})]$$

be the local-volatility value at time t of a contingent claim paying $\Phi(I_T^{LV})$ at time $T > t$. The calculation here is formal for the canonical local-variance surface; it is made rigorous later for the smooth approximants, using Lemmas 14 and 15. Formally, V solves the Black–Scholes PDE

$$V_t(t, s, i) + \lambda(t, s) \left(\frac{1}{2} s^2 V_{ss}(t, s, i) + V_i(t, s, i) \right) = 0, \quad 0 \leq t < T, \quad (6)$$

with the terminal condition $V(T, s, i) = \Phi(i)$. Itô’s formula and (6) give the Duhamel identity

$$\mathbb{E}[\Phi(I_T^H)] - \mathbb{E}[\Phi(I_T^{LV})] = \int_0^T \mathbb{E}[(v_t - \lambda(t, S_t)) H(t, S_t, I_t^H)] \, dt, \quad (7)$$

where

$$H(t, s, i) := \left(\frac{1}{2} s^2 \partial_{ss} + \partial_i \right) V(t, s, i).$$

This is the typical “hedging with the wrong volatility” construction of El Karoui, Jeanblanc-Picqué, and Shreve [10]. Substitution of (5), followed by conditioning on S_t , yields

$$\mathbb{E}[\Phi(I_T^{LV})] - \mathbb{E}[\Phi(I_T^H)] = \beta \int_0^T \mathbb{E}[\text{Cov}(I_t^H, H(t, S_t, I_t^H) \mid S_t)] \, dt. \quad (8)$$

The remaining analytical ingredient is a monotonicity property of H : because Φ is convex, the map

$$i \mapsto H(t, s, i)$$

is nondecreasing for fixed (t, s) . Once this monotonicity is established, the sign in (8) is immediate from Chebyshev's identity: if Y' is an independent copy of Y and h is nondecreasing, then

$$2 \operatorname{Cov}(Y, h(Y)) = \mathbb{E}[(Y - Y')(h(Y) - h(Y'))] \geq 0.$$

Applying this conditionally in (8) with $Y = I_t^H$ and $h(i) = H(t, S_t, i)$ gives the convex-payoff inequality

$$\mathbb{E}[\Phi(I_T^H)] \leq \mathbb{E}[\Phi(I_T^{LV})].$$

Together with equality of the means, this establishes the non-strict convex ordering $I_T^H \preceq_{\text{cx}} I_T^{LV}$. Strictness is addressed next.

Strictness is extracted from the same representation after smoothing the payoff and the local-variance coefficient. For a call strike K , the proof chooses conditional-variance intervals with $j < K < i$. Strict convexity of the smooth call approximants gives a positive integrated increment between j and i , and the localized density lower bound then forces a positive amount of conditional covariance on a time-state box. Passing to the approximation limits makes the call inequality strict, so the two integrated-variance laws cannot coincide. All well-posedness, regularity, approximation, localization, and support statements needed to justify this informal calculation are rather involved and are established in the subsequent sections.

Our main result establishes strict convex order at every positive expiry. In this respect, it differs from the extremality result of Cox and Wang [7] and the robust Root-barrier bounds of Cox and Wang [8]. It is also closely analogous to the convex-order comparison between instantaneous variance and squared VIX discussed by Guyon [14] and Acciaio and Guyon [1]. Perfect correlation is structural rather than merely computational: it is what converts the generator discrepancy into a conditionally centered scalar variable. The present note treats only the endpoint $\rho = -1$; the cases $\rho = +1$ and $-1 < \rho < 1$, including whether strict inequalities persist away from the endpoint, are left open.

2 The two models

We work under a risk-neutral measure, with zero interest rates and zero dividends. The stochastic-volatility model of Heston [16] is given by

$$\frac{dS_t}{S_t} = \sqrt{v_t} dW_t, \tag{9}$$

$$\begin{aligned} dv_t &= \kappa(\theta - v_t) dt + \xi \sqrt{v_t} dZ_t, \\ d\langle W, Z \rangle_t &= \rho dt, \end{aligned} \tag{10}$$

where $S_0 > 0$, $v_0 > 0$, $\kappa, \theta, \xi > 0$, and $\rho \in [-1, 1]$. For the main results,

$$\rho = -1, \tag{11}$$

so going forward we take $Z = -W$. Define the Heston integrated variance by

$$I_t^H := \int_0^t v_u du.$$

Standard localization, the Burkholder–Davis–Gundy inequality, and Gronwall's lemma for the CIR equation give, for every $p \geq 1$ and finite T ,

$$\mathbb{E} \left[\sup_{t \leq T} v_t^p \right] + \mathbb{E} \left[(I_T^H)^p \right] < \infty.$$

The associated ordinary local variance is the Dupire Markovian projection,

$$\lambda(t, s) := \mathbb{E}[v_t \mid S_t = s], \quad (12)$$

with the corresponding local-volatility model

$$\frac{d\bar{S}_t}{\bar{S}_t} = \sqrt{\lambda(t, \bar{S}_t)} dB_t, \quad \bar{S}_0 = S_0. \quad (13)$$

We now justify weak existence together with the required mimicking property. This follows, for example, from Brunick–Shreve [6, Corollary 3.7] or Kurtz–Stockbridge [19, Corollary 4.3, p. 629], recalled here as Corollary 5 in Appendix A. It is convenient to apply either result to $L_t = \log S_t$, with $\hat{\sigma}(t) = \sqrt{v_t}$ and $\hat{b}(t) = -v_t/2$. Since $\mathbb{E}[v_t] = \theta + (v_0 - \theta)e^{-\kappa t}$, all the required finite-horizon integrability conditions hold. Equation (12) gives

$$\mathbb{E}[\hat{\sigma}(t)^2 \mid L_t] = \lambda(t, e^{L_t}), \quad \mathbb{E}[\hat{b}(t) \mid L_t] = -\frac{1}{2}\lambda(t, e^{L_t}).$$

Choosing the nonnegative square root, the corollary supplies at least one weak mimicking diffusion $\bar{L}$ with the same one-time marginal laws as L and dynamics

$$d\bar{L}_t = -\frac{1}{2}\lambda(t, e^{\bar{L}_t}) dt + \sqrt{\lambda(t, e^{\bar{L}_t})} dB_t.$$

Itô's formula then shows that $\bar{S}_t := e^{\bar{L}_t}$ solves (13) and has the required mimicking property

$$\text{Law}(\bar{S}_t) = \text{Law}(S_t), \quad t \geq 0. \quad (14)$$

Thus the two models reproduce the same European option prices at every expiry. We then put

$$I_t^{\text{LV}} := \int_0^t \lambda(u, \bar{S}_u) du. \quad (15)$$

Conditional expectations determine $\lambda(t, \cdot)$ only up to $\text{Law}(S_t)$ -null sets. From here on, λ denotes the canonical version of Proposition 11 in Appendix B: it is jointly continuous at positive times, vanishes on and above the moving barrier produced by the $\rho = -1$ identity of Section 3, and is strictly positive and smooth in spot below it. By the mimicking property (14), replacing the version supplied by the mimicking theorem with this one changes neither integral in (13)–(15). Uniqueness in law for (13) from every state attained at a *positive* time is proved in Theorem 12. No well-posedness from time zero is used in this note; the inception quantity I_T^{LV} always refers to a fixed mimicking solution $\bar{S}$.

3 The pathwise identity for the Heston–Nandi model

This identity is the continuous-time Heston–Nandi relation; see Cox–Wang [7, Eq. (6.2)]. We include its short derivation because it also fixes the sign that drives the comparison.

Set

$$X_t := \log(S_t/S_0).$$

Itô's formula applied to (9) yields

$$X_t = -\frac{1}{2}I_t^{\text{H}} + \int_0^t \sqrt{v_u} dW_u. \quad (16)$$

On the other hand, because $Z = -W$, integration of (10) gives

$$v_t - v_0 = \kappa\theta t - \kappa I_t^{\text{H}} - \xi \int_0^t \sqrt{v_u} dW_u. \quad (17)$$

Eliminating the stochastic integral between (16) and (17), we obtain the central identity

$$v_t = v_0 + \kappa\theta t - \xi \log(S_t/S_0) - \left(\kappa + \frac{\xi}{2}\right) I_t^H. \quad (18)$$

Its general-correlation counterpart is the identity underlying the Monte Carlo simulation scheme of Broadie and Kaya [4, Sec. 3.3]. For later use, write

$$\beta := \kappa + \frac{\xi}{2} > 0, \quad A(t, s) := v_0 + \kappa\theta t - \xi \log(s/S_0). \quad (19)$$

Then $v_t = A(t, S_t) - \beta I_t^H$. Since $v_t \geq 0$ and $I_t^H > 0$ for $t > 0$, the Heston spot satisfies $A(t, S_t) = v_t + \beta I_t^H > 0$ almost surely: at every positive time it lies strictly below the moving barrier

$$s^*(t) := S_0 e^{(v_0 + \kappa\theta t)/\xi}, \quad \{A(t, \cdot) > 0\} = (0, s^*(t)). \quad (20)$$

Taking the conditional expectation of (18) given S_t gives

$$\lambda(t, S_t) = A(t, S_t) - \beta \mathbb{E}[I_t^H \mid S_t]. \quad (21)$$

It will be useful to fix the Borel version

$$\iota(t, s) := \frac{A(t, s) - \lambda(t, s)}{\beta}, \quad t > 0, \quad s > 0. \quad (22)$$

Thus $\iota(t, S_t) = \mathbb{E}[I_t^H \mid S_t]$ almost surely for every $t > 0$. Consequently,

$$v_t - \lambda(t, S_t) = -\beta(I_t^H - \iota(t, S_t)). \quad (23)$$

4 The comparison identity and convex ordering

For later use, fix a maturity $T > 0$ and a convex payoff $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}$. For $t > 0$, the projected SDE is well posed in law from every spot (Theorem 12), so we may define, whenever the expectation is finite,

$$V(t, s, i) := \mathbb{E}_{t,s,i}^{\text{LV}}[\Phi(I_T^{\text{LV}})], \quad (24)$$

where the local-volatility state at time t is $(\bar{S}_t, I_t^{\text{LV}}) = (s, i)$. Equivalently,

$$V(t, s, i) = \mathbb{E}_{t,s}^{\text{LV}} \left[\Phi \left(i + \int_t^T \lambda(u, \bar{S}_u) \, du \right) \right], \quad (25)$$

which also fixes the notation when Φ is specialized below.

Theorem 1. *For the Heston–Nandi model of Section 2 and every $T > 0$,*

$$\mathbb{E}[(I_T^H - K)^+] \leq \mathbb{E}[(I_T^{\text{LV}} - K)^+], \quad K \in \mathbb{R}, \quad (26)$$

and

$$I_T^H \preceq_{\text{cx}} I_T^{\text{LV}}. \quad (27)$$

Equivalently, for every convex payoff $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}$ for which the two expectations are finite,

$$\mathbb{E}[\Phi(I_T^H)] \leq \mathbb{E}[\Phi(I_T^{\text{LV}})]. \quad (28)$$

Proof. Fix $K \in \mathbb{R}$, $m \geq 1$, and put

$$\Phi(i) := \Phi_{m,K}(i) := \frac{1}{m} \log(1 + e^{m(i-K)}).$$

This payoff is smooth, convex, nondecreasing, and 1-Lipschitz, has at most linear growth, and has bounded derivatives of every positive order. Fix also $\varepsilon \in (0, T)$ and write

$$\mu(t, x) := \lambda(t, e^x), \quad \mathcal{D}_{\log} := \frac{1}{2}(\partial_{xx} - \partial_x) + \partial_i.$$

On $[\varepsilon, T]$, Proposition 11 gives the continuity and linear-growth bound required by Lemma 15, while Theorem 12 gives the required well-posedness from every starting point. Let $(\nu_k)_{k \geq 1}$ be the smooth majorants constructed in Lemma 15; their construction is independent of the starting point, and

$$\begin{aligned} \mu(t, x) + \frac{1}{k} &\leq \nu_k(t, x) \leq C_0(1 + |x|), \quad (t, x) \in [\varepsilon, T] \times \mathbb{R}, \\ \lim_{k \rightarrow \infty} \sup_{\substack{t \in [\varepsilon, T] \\ |x| \leq R}} |\nu_k(t, x) - \mu(t, x)| &= 0, \quad \text{for every } R > 0, \end{aligned} \quad (29)$$

with C_0 independent of k . For the diffusion

$$dX_u^k = -\frac{1}{2}\nu_k(u, X_u^k) du + \sqrt{\nu_k(u, X_u^k)} dB_u,$$

define

$$U_k(t, x, i) := \mathbb{E}_{t,x} \left[\Phi \left(i + \int_t^T \nu_k(u, X_u^k) du \right) \right], \quad h_k := \mathcal{D}_{\log} U_k. \quad (30)$$

Lemma 14 shows that $U_{k,x}$ and h_k are bounded, $h_k \geq 0$, and $i \mapsto h_k(t, x, i)$ is nondecreasing.

Let $L_t := \log S_t$ and, for $n \geq 1$, set

$$\tau_n := T \wedge \inf\{t \in [\varepsilon, T] : |L_t| + v_t + I_t^H \geq n\},$$

with $\inf \emptyset = \infty$. Continuity of the Heston paths gives $\tau_n \uparrow T$ a.s. Since

$$dL_t = -\frac{1}{2}v_t dt + \sqrt{v_t} dW_t, \quad dI_t^H = v_t dt,$$

For $\delta \in (0, T - \varepsilon)$, Itô's formula and the backward equation in Lemma 14, applied up to $\tau_n \wedge (T - \delta)$, give the corresponding stopped identity. Letting $\delta \downarrow 0$, using the continuity and bounds from Lemma 14, yields

$$\begin{aligned} U_k(\tau_n, L_{\tau_n}, I_{\tau_n}^H) - U_k(\varepsilon, L_\varepsilon, I_\varepsilon^H) \\ = \int_\varepsilon^{\tau_n} (v_t - \nu_k(t, L_t)) h_k(t, L_t, I_t^H) dt + M_{\tau_n}^k, \end{aligned}$$

where

$$M_t^k := \int_\varepsilon^t \sqrt{v_u} U_{k,x}(u, L_u, I_u^H) dW_u.$$

For fixed k , the bounds from Lemma 14, (29), and the Heston moment bounds make M^k square-integrable and the drift absolutely integrable. Indeed, (18) gives $|L_t| \leq C_T(1 + v_t + I_t^H)$, and hence $\mathbb{E} \int_\varepsilon^T (v_t + \nu_k(t, L_t)) dt < \infty$. Explicitly,

$$\mathbb{E} \left[\int_\varepsilon^T \left(v_t |U_{k,x}(t, L_t, I_t^H)|^2 + (v_t + \nu_k(t, L_t)) |h_k(t, L_t, I_t^H)| \right) dt \right] < \infty.$$

Consequently $M_{\tau_n}^k \rightarrow M_T^k$ in L^2 , and the stopped drift converges in L^1 to the integral over $[\varepsilon, T]$. The linear-growth bound for U_k and the Heston moment bounds similarly give

$$U_k(\tau_n, L_{\tau_n}, I_{\tau_n}^H) \rightarrow U_k(T, L_T, I_T^H) = \Phi(I_T^H) \quad \text{in } L^1.$$

Letting $n \rightarrow \infty$ and then taking expectations therefore yields

$$\begin{aligned} \mathbb{E}[\Phi(I_T^H)] - \mathbb{E}[U_k(\varepsilon, L_\varepsilon, I_\varepsilon^H)] \\ = \int_\varepsilon^T \mathbb{E}[(v_t - \nu_k(t, L_t))h_k(t, L_t, I_t^H)] \, dt. \end{aligned} \quad (31)$$

Set

$$C_k(t) := \mathbb{E}[(I_t^H - \iota(t, S_t))h_k(t, L_t, I_t^H)].$$

This is a measurable function of t : all the terms in the expectation have jointly measurable versions, and the preceding integrability estimates apply. We use the following classical scalar covariance identity, valid for similarly ordered functions; see, for example, Stroock [23, Exercise 1.1.9(i)]. If $\eta_{t,s}$ is any regular conditional law of I_t^H given $S_t = s$, then, for $\text{Law}(S_t)$ -almost every s ,

$$\begin{aligned} 2 \text{Cov}(I_t^H, h_k(t, \log s, I_t^H) \mid S_t = s) \\ = \int_{\mathbb{R}_+^2} (y - y')(h_k(t, \log s, y) - h_k(t, \log s, y')) \eta_{t,s}(dy) \eta_{t,s}(dy') \geq 0. \end{aligned} \quad (32)$$

Integrating (32) with respect to $\text{Law}(S_t)(ds)$ and using (22) gives $C_k(t) \geq 0$ for every $t > 0$. Consequently, (23), (29), and $h_k \geq 0$ give

$$\begin{aligned} \mathbb{E}[(v_t - \nu_k(t, L_t))h_k(t, L_t, I_t^H)] \\ = -\beta C_k(t) + \mathbb{E}[(\lambda(t, S_t) - \nu_k(t, L_t))h_k(t, L_t, I_t^H)] \leq 0. \end{aligned} \quad (33)$$

Thus (31) yields

$$\mathbb{E}[\Phi(I_T^H)] \leq \mathbb{E}[U_k(\varepsilon, L_\varepsilon, I_\varepsilon^H)]. \quad (34)$$

Lemma 15 and its fourth-moment estimate imply, for each (x, i) ,

$$U_k(\varepsilon, x, i) \rightarrow V(\varepsilon, e^x, i).$$

The same estimates and the 1-Lipschitz property give, with a constant independent of k ,

$$|U_k(\varepsilon, x, i)| + |V(\varepsilon, e^x, i)| \leq C(1 + |x| + i), \quad i \geq 0. \quad (35)$$

The pathwise identity (18) and the CIR moment bounds imply $\mathbb{E}[|L_\varepsilon| + I_\varepsilon^H] < \infty$. Dominated convergence in (34) therefore gives

$$\mathbb{E}[\Phi(I_T^H)] \leq B_\varepsilon := \mathbb{E}[V(\varepsilon, S_\varepsilon, I_\varepsilon^H)]. \quad (36)$$

It remains to identify the time-zero boundary term (without assuming well-posedness from time zero). The pointwise convergence above and the continuity of each $U_k(\varepsilon, \cdot, \cdot)$ show that

$$V(\varepsilon, s, i) = \lim_{k \rightarrow \infty} U_k(\varepsilon, \log s, i), \quad (s, i) \in (0, \infty) \times \mathbb{R}_+,$$

and hence that $(s, i) \mapsto V(\varepsilon, s, i)$ is Borel. Apply Lemma 13 to $\bar{X} := \log \bar{S}$ and put $\mathbf{m}_\varepsilon := \text{Law}(\bar{X}_\varepsilon)$. By Lemma 13(a), outside one $\mathbf{m}_\varepsilon$ -null set the kernel $P_x^{(\varepsilon, \bar{X})}$ is the unique path law from (ε, x) . Consequently, on that same full set, for every $i \geq 0$,

$$\int_{C_\varepsilon} \Phi\left(i + \int_\varepsilon^T \mu(u, \omega(u)) \, du\right) P_x^{(\varepsilon, \bar{X})}(d\omega) = V(\varepsilon, e^x, i).$$

Since $I_\varepsilon^{\text{LV}}$ is $\mathcal{F}_\varepsilon$ -measurable, it can be frozen in Lemma 13(b). The required parameterized form follows first for products of a bounded Borel function of $I_\varepsilon^{\text{LV}}$ and a bounded path functional, and then for all bounded jointly Borel functionals by the monotone-class theorem. Applying this first to bounded truncations and then using monotone convergence gives

$$\mathbb{E}[\Phi(I_T^{\text{LV}})] = \mathbb{E}[V(\varepsilon, \bar{S}_\varepsilon, I_\varepsilon^{\text{LV}})]. \quad (37)$$

Because Φ is nondecreasing and 1-Lipschitz, (25) implies

$$0 \leq V(\varepsilon, s, i) - V(\varepsilon, s, 0) \leq i, \quad s > 0, \quad i \geq 0. \quad (38)$$

Since $s \mapsto V(\varepsilon, s, 0)$ is Borel, (14) yields

$$\mathbb{E}[V(\varepsilon, S_\varepsilon, 0)] = \mathbb{E}[V(\varepsilon, \bar{S}_\varepsilon, 0)].$$

Putting $L_\Phi := \mathbb{E}[\Phi(I_T^{\text{LV}})]$, combining this equality with (37), and applying (38) to the two accumulated variances at time ε gives

$$-\mathbb{E}[I_\varepsilon^{\text{LV}}] \leq B_\varepsilon - L_\Phi \leq \mathbb{E}[I_\varepsilon^{\text{H}}]. \quad (39)$$

Both error terms satisfy

$$\mathbb{E}[I_\varepsilon^{\text{H}}] = \mathbb{E}[I_\varepsilon^{\text{LV}}] = \int_0^\varepsilon \mathbb{E}[v_t] \, dt \longrightarrow 0 \quad (\varepsilon \downarrow 0).$$

Hence $B_\varepsilon \rightarrow L_\Phi$, and (36) gives

$$\mathbb{E}[\Phi_{m,K}(I_T^{\text{H}})] \leq \mathbb{E}[\Phi_{m,K}(I_T^{\text{LV}})]. \quad (40)$$

Since

$$0 \leq \Phi_{m,K}(i) - (i - K)^+ \leq \frac{\log 2}{m}, \quad i \geq 0,$$

letting $m \rightarrow \infty$ proves (26).

Finally, by (12), Tonelli's theorem, and the mimicking property,

$$\begin{aligned} \mathbb{E}[I_T^{\text{H}}] &= \int_0^T \mathbb{E}[v_t] \, dt = \int_0^T \mathbb{E}[\lambda(t, S_t)] \, dt \\ &= \int_0^T \mathbb{E}[\lambda(t, \bar{S}_t)] \, dt = \mathbb{E}[I_T^{\text{LV}}]. \end{aligned} \quad (41)$$

The call inequalities together with the equal means are the standard characterization of convex order for integrable random variables, proving (27); the equivalent payoff formulation follows. $\square$

4.1 Strictness

The strict comparison uses the following nondegeneracy property of the Heston accumulated variance. Its proof is deferred to Appendix C.

Lemma 2 (Localized lower density for Heston accumulated variance). *Set*

$$\mathcal{O} := \{(t, s, i) : t > 0, \quad 0 < s < s^*(t), \quad 0 < i < A(t, s)/\beta\}.$$

For every compact set $\mathcal{K}$ contained entirely in the open set $\mathcal{O}$ (and hence separated by a positive distance from its boundary), there is a nonnegative, jointly Borel function $q_{\mathcal{K}}$ on $(0, \infty)^3$ such that, for every $t > 0$, writing $\Gamma_t := \text{Law}(S_t, I_t^{\text{H}})$,

$$\Gamma_t(ds, di) \geq q_{\mathcal{K}}(t, s, i) \, ds \, di, \quad \inf_{(t,s,i) \in \mathcal{K}} q_{\mathcal{K}}(t, s, i) > 0. \quad (42)$$

The function q_K may be taken to vanish outside the time-projection of K . The marginal spot law has a density $f_t^S(s)$ admitting a jointly continuous, strictly positive version on $\{(t, s) : 0 < t \leq T, 0 < s < s^*(t)\}$ and hence locally bounded there. If $\eta_{t,s}$ is a regular conditional law of I_t^H given $S_t = s$, then, for every compact set K contained entirely in $\mathcal{O}$ and every $t \in (0, T]$,

$$f_t^S(s) \eta_{t,s}(di) \geq q_K(t, s, i) \, di \quad \text{for almost every } s. \quad (43)$$

Moreover, on each compact interval $J = [t_0, T] \subset (0, T]$, there exists a jointly Borel probability kernel $(t, s) \mapsto \eta_{t,s}$ such that, for dt -almost every $t \in J$, the map $s \mapsto \eta_{t,s}$ is a regular conditional law of I_t^H given $S_t = s$; for this choice, (43) holds for $dt \, ds$ -almost every $(t, s) \in J \times (0, \infty)$.

Theorem 3 (Strict integrated-variance comparison). *For the Heston–Nandi model of Section 2 and every $T > 0$,*

$$\mathbb{E}[(I_T^H - K)^+] < \mathbb{E}[(I_T^{LV} - K)^+], \quad K > 0. \quad (44)$$

Consequently,

$$I_T^H \prec_{\text{cx}} I_T^{LV}. \quad (45)$$

Moreover, for every strictly convex $\Psi : \mathbb{R}_+ \rightarrow \mathbb{R}$ such that $\Psi(I_T^H), \Psi(I_T^{LV}) \in L^1$,

$$\mathbb{E}[\Psi(I_T^H)] < \mathbb{E}[\Psi(I_T^{LV})]. \quad (46)$$

Proof. Fix $K > 0$ and $t_0 \in (0, T)$. Since $A(t_0, s) \rightarrow \infty$ as $s \downarrow 0$, we may choose numbers

$$0 < s_1 < s_2 < s^*(t_0), \quad 0 < i_1 < i_2 < K < i_3 < i_4 < \frac{A(t_0, s_2)}{\beta}. \quad (47)$$

The barrier $s^*(t)$ and $A(t, s)$ are increasing in t , while $A(t, s)$ is decreasing in s . Hence this entire box remains in the interior region $\mathcal{O}$ of Lemma 2 for $t \in [t_0, T]$.

For $m \geq 1$, let $\Phi_m := \Phi_{m,K}$ be the soft call from the proof of Theorem 1. Fix for the moment m and $\varepsilon \in (0, t_0)$. Apply the smooth-majorant construction on $[\varepsilon, T]$ with payoff Φ_m . Write $U_{k,m}$ for the value in (30) and $h_{k,m} := \mathcal{D}_{\log} U_{k,m}$. Lemma 14 shows that $h_{k,m} \geq 0$ and that it is nondecreasing in its last argument. Put

$$C_{k,m}(t) := \mathbb{E}[(I_t^H - \iota(t, S_t)) h_{k,m}(t, L_t, I_t^H)] \geq 0.$$

This unconditional formula makes $t \mapsto C_{k,m}(t)$ measurable; for each fixed t it equals the expected conditional covariance by (22). Rearranging (31) by means of (23) gives the quantitative gap identity

$$\begin{aligned} & \mathbb{E}[U_{k,m}(\varepsilon, L_\varepsilon, I_\varepsilon^H)] - \mathbb{E}[\Phi_m(I_T^H)] \\ &= \beta \int_\varepsilon^T C_{k,m}(t) \, dt + \int_\varepsilon^T \mathbb{E}[(\nu_k(t, L_t) - \lambda(t, S_t)) h_{k,m}(t, L_t, I_t^H)] \, dt \geq \beta \int_\varepsilon^T C_{k,m}(t) \, dt. \end{aligned} \quad (48)$$

Thus it is enough to retain a positive amount of covariance as $k \rightarrow \infty$; no derivative of the canonical value function is involved.

For $(s, j, i) \in (s_1, s_2) \times (i_1, i_2) \times (i_3, i_4)$ define

$$G_{k,m}(r, s; j, i) := h_{k,m}(r, \log s, i) - h_{k,m}(r, \log s, j) \geq 0, \quad \Psi_{j,i}^m(z) := \Phi_m(i + z) - \Phi_m(j + z).$$

Let $J_{t_0,s}^k$ be the accumulated variance from t_0 to T for the diffusion with coefficient ν_k , started from $\log s$ at t_0 . The integral form (85), subtracted at i and j , gives

$$\begin{aligned} R_{k,m}(s, j, i) &:= \int_{t_0}^T \nu_k(r, \log s) G_{k,m}(r, s; j, i) \, dr \\ &= \mathbb{E}[\Psi_{j,i}^m(J_{t_0,s}^k)] - \Psi_{j,i}^m(0). \end{aligned} \quad (49)$$

The restrictions of ν_k to $[t_0, T]$ still converge locally uniformly to μ and have a common linear-growth bound. The stability argument of Lemma 15 therefore gives $J_{t_0,s}^k \Rightarrow J_{t_0,s}$, where

$$J_{t_0,s} := \int_{t_0}^T \lambda(r, \bar{S}_r^{t_0,s}) \, dr.$$

Here $\bar{S}^{t_0,s}$ denotes the solution of (13) started from s at time t_0 . By Proposition 11(iii), $\lambda(t_0, s) > 0$. Joint continuity of λ and continuity of $\bar{S}^{t_0,s}$ therefore make $u \mapsto \lambda(u, \bar{S}_u^{t_0,s})$ positive on a nontrivial interval immediately after t_0 , almost surely. Hence $J_{t_0,s} > 0$ almost surely. Since Φ_m is strictly convex, $\Psi_{j,i}^m$ is strictly increasing; it is also bounded by $i - j$. Hence

$$R_{k,m}(s, j, i) \longrightarrow R_m(s, j, i) := \mathbb{E}[\Psi_{j,i}^m(J_{t_0,s})] - \Psi_{j,i}^m(0) > 0.$$

The common growth bound in (29) supplies a finite C_{box} , independent of k and ε , such that $\nu_k(r, \log s) \leq C_{\text{box}}$ on $[t_0, T] \times [s_1, s_2]$. Dominated convergence in (s, j, i) and (49) therefore show that

$$\liminf_{k \rightarrow \infty} \int_{t_0}^T \int_{s_1}^{s_2} \int_{i_1}^{i_2} \int_{i_3}^{i_4} G_{k,m}(r, s; j, i) \, di \, dj \, ds \, dr \geq \frac{\Delta_m}{C_{\text{box}}} > 0, \quad (50)$$

where

$$\Delta_m := \int_{s_1}^{s_2} \int_{i_1}^{i_2} \int_{i_3}^{i_4} R_m(s, j, i) \, di \, dj \, ds > 0.$$

Apply Lemma 2 to

$$\mathcal{K}_0 := [t_0, T] \times [s_1, s_2] \times ([i_1, i_2] \cup [i_3, i_4]).$$

This set is compact and contained entirely in $\mathcal{O}$. It supplies a subdensity $q := q_{\mathcal{K}_0}$. Compact positivity of q and local boundedness of the chosen continuous version of f^S give constants $q_0, F_0 > 0$ such that

$$q(t, s, z) \geq q_0 \quad ((t, s, z) \in \mathcal{K}_0), \quad f_t^S(s) \leq F_0, \quad (t, s) \in [t_0, T] \times [s_1, s_2].$$

Choose on $[t_0, T]$ the jointly Borel kernel supplied by Lemma 2. For dt -almost every t ,

$$2C_{k,m}(t) = \int_0^\infty f_t^S(s) \int_{\mathbb{R}_+^2} (y - y') (h_{k,m}(t, \log s, y) - h_{k,m}(t, \log s, y')) \eta_{t,s}(dy) \eta_{t,s}(dy') \, ds. \quad (51)$$

Moreover, the joint form of (43) gives

$$f_t^S(s) \eta_{t,s}(dz) \geq q(t, s, z) \, dz \quad \text{for } dt \, ds\text{-almost every } (t, s).$$

On the fixed box, $f_t^S(s) > 0$, and tensoring the preceding measure inequality with itself gives, for almost every (t, s) ,

$$f_t^S(s) \eta_{t,s}(dj) \eta_{t,s}(di) = \frac{[f_t^S(s) \eta_{t,s}(dj)][f_t^S(s) \eta_{t,s}(di)]}{f_t^S(s)} \geq \frac{q_0^2}{F_0} \, dj \, di$$

on $[i_1, i_2] \times [i_3, i_4]$. Tonelli's theorem, (51), and restriction to this box therefore yield

$$2 \int_{t_0}^T C_{k,m}(t) \, dt \geq \frac{(i_3 - i_2) q_0^2}{F_0} \int_{t_0}^T \int_{s_1}^{s_2} \int_{i_1}^{i_2} \int_{i_3}^{i_4} G_{k,m}(t, s; j, i) \, di \, dj \, ds \, dt. \quad (52)$$

Combining (50) and (52) gives

$$\liminf_{k \rightarrow \infty} \int_{t_0}^T C_{k,m}(t) \, dt \geq \frac{(i_3 - i_2) q_0^2 \Delta_m}{2F_0 C_{\text{box}}} > 0.$$

The constants C_{box}, q_0, F_0 are independent of both m and ε : the constant C_λ in Proposition 11(ii) is uniform on $(0, T]$, while the localized lower-density bounds depend only on the fixed interior box.

Let

$$B_{\varepsilon, m} := \mathbb{E}[V_m(\varepsilon, S_\varepsilon, I_\varepsilon^H)],$$

where V_m is defined probabilistically by (25) with payoff Φ_m . The same stability and dominated-convergence step used in (36) gives

$$\mathbb{E}[U_{k, m}(\varepsilon, L_\varepsilon, I_\varepsilon^H)] \longrightarrow B_{\varepsilon, m}.$$

Letting $k \rightarrow \infty$ in (48) therefore gives, for every $\varepsilon < t_0$,

$$B_{\varepsilon, m} - \mathbb{E}[\Phi_m(I_T^H)] \geq \frac{\beta(i_3 - i_2)q_0^2}{2F_0C_{\text{box}}} \Delta_m.$$

Finally, the boundary sandwich (39), applied to Φ_m , identifies the limit of $B_{\varepsilon, m}$ as $\varepsilon \downarrow 0$. We have proved

$$\mathbb{E}[\Phi_m(I_T^{LV})] - \mathbb{E}[\Phi_m(I_T^H)] \geq \frac{\beta(i_3 - i_2)q_0^2}{2F_0C_{\text{box}}} \Delta_m. \quad (53)$$

It remains to pass from the soft call to the call itself. Let $\Phi_K(i) := (i - K)^+$ and $\Psi_{j, i}^K(z) := \Phi_K(i + z) - \Phi_K(j + z)$. Since $j < K < i$ throughout the box (47), for $z \geq 0$,

$$\Psi_{j, i}^K(z) - \Psi_{j, i}^K(0) = z \wedge (K - j).$$

The bound $0 \leq \Phi_m(i) - \Phi_K(i) \leq (\log 2)/m$ and dominated convergence therefore give

$$\Delta_m \longrightarrow \Delta_K := \int_{s_1}^{s_2} \int_{i_1}^{i_2} \int_{i_3}^{i_4} \mathbb{E}[J_{t_0, s} \wedge (K - j)] \, di \, dj \, ds > 0,$$

where positivity follows from $J_{t_0, s} > 0$ a.s. Letting $m \rightarrow \infty$ in (53) proves (44). Since $K > 0$ was arbitrary, the inequality holds at every positive strike. In particular, the two integrated-variance laws are different, and Theorem 1 gives (45).

For completeness, Strassen's characterization supplies a martingale coupling (X, Y) whose marginals are, respectively, the laws of I_T^H and I_T^{LV} , and for which $\mathbb{E}[Y \mid X] = X$. If π_x denotes a regular conditional law of Y given $X = x$, then equality in conditional Jensen for a strictly convex Ψ holds precisely when π_X is almost surely a point mass. The martingale condition then forces $Y = X$ almost surely, which would make the marginal laws equal. This proves (46). $\square$

For comparison, the instantaneous variances have the elementary ordering

$$\lambda(t, S_t) = \mathbb{E}[v_t \mid S_t] \preceq_{\text{cx}} v_t, \quad (54)$$

by conditional Jensen. Thus (27) is an ordering inversion on passing from instantaneous variance to its time integral.

Remark 4. The feedback coefficient $\beta = \kappa + \xi/2$ in (23) is positive at $\rho = -1$: at fixed spot, high accumulated variance lowers the current variance, by (18). This feedback, rather than perfect correlation by itself, is what drives the ordering.

A The Kurtz–Stockbridge mimicking corollary

For completeness, we record the parallel result mentioned in Section 2.

Corollary 5 (Kurtz–Stockbridge [19, Corollary 4.3]). *Suppose that W is an $\mathbb{R}^d$ -valued $\{\mathcal{F}_t\}$ -Brownian motion; $\hat{\sigma}$ and $\hat{b}$ are progressively measurable processes taking values in $\mathbb{R}^{d \times d}$ and $\mathbb{R}^d$, respectively; $X(0)$ is $\mathbb{R}^d$ -valued and $\mathcal{F}_0$ -measurable; and, for every $T > 0$,*

$$\mathbb{E} \int_0^T \left(\|\hat{\sigma}(s)\|_{\mathbb{F}}^2 + |\hat{b}(s)| \right) ds < \infty.$$

Let

$$X(t) = X(0) + \int_0^t \hat{\sigma}(s) dW(s) + \int_0^t \hat{b}(s) ds.$$

Then there exist measurable functions

$$\sigma : [0, \infty) \times \mathbb{R}^d \longrightarrow \mathbb{R}^{d \times d}, \quad b : [0, \infty) \times \mathbb{R}^d \longrightarrow \mathbb{R}^d,$$

an $\mathbb{R}^d$ -valued Brownian motion $\widetilde{W}$, and a process Y satisfying

$$Y(t) = Y(0) + \int_0^t \sigma(s, Y(s)) d\widetilde{W}(s) + \int_0^t b(s, Y(s)) ds,$$

such that, for each $t \geq 0$, the distributions of

$$\left(X(t), \mathbb{E}[\hat{b}(t) \mid X(t)], \mathbb{E}[\hat{\sigma}(t)\hat{\sigma}(t)^\top \mid X(t)] \right)$$

and

$$\left(Y(t), b(t, Y(t)), \sigma(t, Y(t))\sigma(t, Y(t))^\top \right)$$

are the same.

B Positive-time regularity and well-posedness of the projected SDE for Heston–Nandi

This appendix proves the properties of the projected surface λ and of the SDE (13) that are needed at positive times. For compactness of notation, write the Heston variance equation as $dv_t = (a + bv_t) dt + c\sqrt{v_t} dZ_t$, with $a > 0$, $b < 0$, $c > 0$; the present parameters correspond to

$$a = \kappa\theta > 0, \quad b = -\kappa < 0, \quad c = \xi,$$

and therefore $\beta = c/2 - b = \kappa + \xi/2$, consistently with (19). In log spot $x = \log s$, the barrier (20) reads $x^*(t) := \log S_0 + (v_0 + \kappa\theta t)/\xi$ and $A(t, s) = \xi(x^*(t) - \log s)$. Once the canonical version is constructed below, we write

$$\mu(t, x) := \lambda(t, e^x) \tag{55}$$

for the surface in log spot. The *log form* of (13) on $[r, T]$ from x is

$$d\bar{X}_u = -\frac{1}{2}\mu(u, \bar{X}_u) du + \sqrt{\mu(u, \bar{X}_u)} dB_u, \quad u \in [r, T], \quad \bar{X}_r = x. \tag{56}$$

By Itô's formula, $\bar{S}$ solves (13) on $[r, T]$ with continuous paths in $(0, \infty)$ if and only if $\bar{X} = \log \bar{S}$ solves (56) with continuous real-valued paths, so the two forms may be used interchangeably. At $r = 0$, all coefficient integrals refer to the fixed mimicking solution $\bar{X} := \log \bar{S}$. For this solution,

$$\mathbb{E} \int_0^T \mu(u, \bar{X}_u) du = \int_0^T \mathbb{E}[v_u] du < \infty.$$

Thus its nonnegative time integral is finite almost surely.

B.1 The gap transform and the canonical surface

Put

$$Q_t := v_t + \beta I_t^H = A(t, S_t) = c(x^*(t) - \log S_t) > 0. \quad (57)$$

Thus conditioning on S_t is equivalent to conditioning on Q_t . The regularity of that conditional expectation can be obtained from the affine transform of the integrated CIR process, which we now record explicitly. For $\Re z > 0$, set

$$\begin{aligned} d = d(z) &:= \sqrt{b^2 + 2\beta c^2 z}, & r_{\pm} = r_{\pm}(z) &:= \frac{b \pm d}{c^2}, & \mathfrak{h} = \mathfrak{h}(z) &:= \frac{z - r_+}{z - r_-}, \\ B_t(z) &:= \frac{r_+ - \mathfrak{h}e^{-dt}r_-}{1 - \mathfrak{h}e^{-dt}}, & \mathcal{A}_t(z) &:= r_+t + \frac{2}{c^2} \log \left(\frac{1 - \mathfrak{h}e^{-dt}}{1 - \mathfrak{h}} \right), \\ \mathcal{H}_t(z) &:= e^{-dt} \left(\frac{1 - \mathfrak{h}}{1 - \mathfrak{h}e^{-dt}} \right)^2, & \mathcal{J}_t(z) &:= \frac{2(1 - e^{-dt})}{c^2(z - r_-)(1 - \mathfrak{h}e^{-dt})}. \end{aligned} \quad (58)$$

The principal square root is used, and the logarithm is selected by continuation from the positive real axis.

Lemma 6 (Affine transforms of the gap). *For $\Re z > 0$,*

$$L_t(z) := \mathbb{E}[e^{-zQ_t}] = \exp\{-a\mathcal{A}_t(z) - v_0B_t(z)\}, \quad (59)$$

$$\mathcal{M}_t(z) := \mathbb{E}[v_te^{-zQ_t}] = L_t(z)(a\mathcal{J}_t(z) + v_0\mathcal{H}_t(z)). \quad (60)$$

Both functions are analytic on $\{\Re z > 0\}$. For every $t_0 > 0$ and $\sigma > 0$, there are constants $C, N, c_0 > 0$ such that

$$|L_t(\sigma + iy)| + |\mathcal{M}_t(\sigma + iy)| \leq C(1 + |y|)^N e^{-c_0\sqrt{|y|}}, \quad t \in [t_0, T], \quad y \in \mathbb{R}. \quad (61)$$

Proof. For positive real terminal parameter u and running parameter q , the joint CIR Laplace transform of Lamberton and Lapeyre [20, Prop. 6.2.4, p. 162], applied with $X = v$, $x = v_0$, $a_{\text{book}} = a$, $b_{\text{book}} = -b$, $\sigma = c$, $\lambda = u$, and $\mu = q$, gives

$$\mathbb{E}\left[e^{-uv_t - q \int_0^t v_s ds}\right] = \exp\{-a\mathcal{A}_t(u, q) - v_0\mathcal{B}_t(u, q)\},$$

where

$$\dot{\mathcal{B}} = q + b\mathcal{B} - \frac{c^2}{2}\mathcal{B}^2, \quad \mathcal{B}_0 = u, \quad \mathcal{A}_t(u, q) = \int_0^t \mathcal{B}_s(u, q) ds.$$

Taking $(u, q) = (z, \beta z)$ for positive real z and solving the Riccati equation gives (59) and (58). Differentiating with respect to the terminal parameter while keeping the running parameter equal to βz yields

$$\partial_u \mathcal{B}_t(u, \beta z)|_{u=z} = \mathcal{H}_t(z), \quad \partial_u \mathcal{A}_t(u, \beta z)|_{u=z} = \mathcal{J}_t(z),$$

and hence (60). The affine identity is first obtained for positive real transform parameters; after this specialization, both the probabilistic and closed-form sides are holomorphic on $\{\Re z > 0\}$, so (59)–(60) extend there by the identity theorem.

For completeness, the closed forms are analytic throughout the right half-plane. There $\Re d > 0$ and $\text{sgn}(\Im d) = \text{sgn}(\Im z)$. Moreover, $\Re(z - r_-) > 0$ and

$$|z - r_+|^2 - |z - r_-|^2 = -\frac{4}{c^2}\Re(z\bar{d}) + \frac{4b}{c^4}\Re d < 0,$$

so $|\mathfrak{h}| < 1$. Consequently $1 - \mathfrak{h}e^{-dt} \neq 0$ and $1 - \mathfrak{h} = 2d/(c^2(z - r_-)) \neq 0$. The quotient inside the logarithm is analytic and nonzero on the simply connected half-plane, which fixes

the claimed analytic branch. The zero-exclusion and branch-selection argument above is in line with Lucic [21, Thm. 2.1], who characterizes the singularities of the Heston characteristic function as purely imaginary. The use here of analytic continuation and transform estimates to establish density regularity parallels the complex-moment-generating-function analysis of del Baño Rollin, Ferreiro-Castilla, and Utzet [9]. Finally fix $t_0 > 0$ and $\sigma > 0$, put $z = \sigma + iy$, and write $\langle y \rangle = 1 + |y|$. The explicit formula for d gives constants $c_j, C_j > 0$, depending only on t_0, σ, T , such that, uniformly for $t \in [t_0, T]$,

$$\Re d(z) \geq c_1 \langle y \rangle^{1/2}, \quad |d(z)| \leq C_1 \langle y \rangle^{1/2}, \quad |e^{-d(z)t}| \leq e^{-c_1 t_0 \langle y \rangle^{1/2}}.$$

Furthermore, there are constants $0 < \gamma \leq \Gamma < \infty$ such that, for every $y \in \mathbb{R}$,

$$\begin{aligned} \gamma \langle y \rangle &\leq |z - r_-(z)| \leq \Gamma \langle y \rangle, \\ \gamma \langle y \rangle^{-1/2} &\leq |1 - \mathfrak{h}(z)| = \frac{2|d(z)|}{c^2 |z - r_-(z)|} \leq \Gamma \langle y \rangle^{-1/2}. \end{aligned}$$

Since $|\mathfrak{h}| < 1$,

$$|1 - \mathfrak{h}e^{-dt}| \geq 1 - e^{-c_1 t_0 \langle y \rangle^{1/2}},$$

and the right-hand side is bounded away from zero. Using

$$B_t = r_+ + \frac{\mathfrak{h}e^{-dt}(r_+ - r_-)}{1 - \mathfrak{h}e^{-dt}}$$

and $\Re \log w = \log |w|$ on the selected analytic branch, we obtain

$$\Re B_t(\sigma + iy) \geq c_2 \langle y \rangle^{1/2} - C_2, \quad |B_t(\sigma + iy)| \leq C_2 \langle y \rangle^{1/2},$$

and

$$\begin{aligned} \Re \mathcal{A}_t(\sigma + iy) &= t \Re r_+ + \frac{2}{c^2} (\log |1 - \mathfrak{h}e^{-dt}| - \log |1 - \mathfrak{h}|) \\ &\geq c_3 \langle y \rangle^{1/2} - C_3. \end{aligned}$$

The remaining two factors satisfy the explicit bounds

$$|\mathcal{H}_t(\sigma + iy)| \leq C_4 \langle y \rangle^{-1} e^{-c_1 t_0 \langle y \rangle^{1/2}}, \quad |\mathcal{J}_t(\sigma + iy)| \leq C_4 \langle y \rangle^{-1}.$$

Consequently,

$$|L_t(\sigma + iy)| \leq C_5 e^{-c_5 \sqrt{|y|}}, \quad |\mathcal{M}_t(\sigma + iy)| \leq C_5 (1 + |y|)^{-1} e^{-c_5 \sqrt{|y|}},$$

after enlarging C_5 on a bounded y -interval. This proves (61), uniformly for $t \in [t_0, T]$. $\square$

Proposition 7 (Smooth gap densities). *For every $t > 0$, the law of Q_t and the v_t -weighted law of Q_t have densities g_t and m_t , respectively:*

$$\text{Law}(Q_t)(dq) = g_t(q) dq, \quad \mathbb{E}[v_t; Q_t \in dq] = m_t(q) dq.$$

On $(0, T] \times (0, \infty)$ these densities, together with all their q -derivatives, are jointly continuous. If the CIR process is instead started from $y > 0$, the corresponding densities $g_t^y(q)$ and $m_t^y(q)$ are jointly continuous in $(y, q) \in (0, \infty)^2$.

Proof. For $\sigma > 0$, consider the finite measures

$$e^{-\sigma q} \text{Law}(Q_t)(dq), \quad e^{-\sigma q} \mathbb{E}[v_t; Q_t \in dq].$$

Their Fourier transforms are, respectively, $L_t(\sigma + iy)$ and $\mathcal{M}_t(\sigma + iy)$. Since the exponential factor in (61) dominates every polynomial in $|y|$, the bound remains integrable after multiplication by any power of $|y|$. Fourier inversion and arbitrary differentiation in q are therefore legitimate. Dominated convergence gives joint continuity, locally uniformly for $t > 0$. Replacing v_0 by y in (59)–(60) proves the last assertion, since the same domination is locally uniform for $y > 0$. $\square$

Lemma 8 (Positivity of the gap densities). *For every $t \in (0, T]$ and $q > 0$,*

$$g_t(q) > 0, \quad m_t(q) > 0.$$

Proof. We first note a support property of the CIR process. Let $t_1 > 0$ and let $\varphi : [0, t_1] \rightarrow (0, \infty)$ be continuous with $\varphi(0) = v_0$. The Lamperti transform $R_u := 2\sqrt{v_u}/c$, up to its first hitting time of zero, satisfies

$$dR_u = \left(\frac{2a/c^2 - 1/2}{R_u} + \frac{b}{2} R_u \right) du + dZ_u.$$

A compact interval in $(0, \infty)$ containing a small uniform neighborhood of $2\sqrt{\varphi}/c$ permits the drift to be extended to a bounded Lipschitz function on $\mathbb{R}$. The localized diffusion agrees with R until it exits that interval. Girsanov's theorem and the support of Wiener measure give positive probability to every sufficiently narrow tube about $2\sqrt{\varphi}/c$, and hence to every sufficiently narrow tube about φ for the variance process. It follows that, for every nonempty open $U \subset (0, \infty)^2$,

$$\mathbb{P}((v_{t_1}, I_{t_1}^H) \in U) > 0. \quad (62)$$

Indeed, a positive continuous path with prescribed positive endpoint and prescribed positive integral can be formed by a dip-and-recovery path, with a positive bump used to adjust the integral.

Fix $q > 0$. Choose $y_1 \in (0, q/4)$ and then $\tau \in (0, t)$ sufficiently small that the gap process started from y_1 satisfies $\mathbb{P}_{y_1}(Q_\tau < q/2) > 1/2$. Since $g_\tau^{y_1}$ integrates to one, there is $z_1 \in (0, q/2)$ with $g_\tau^{y_1}(z_1) > 0$. Proposition 7 makes this density bounded below on a small neighborhood of (y_1, z_1) . With $t_1 = t - \tau$, the Markov property gives the convolution identity below for almost every q . Choose points from that full-measure set converging to the prescribed q ; continuity of g_t and Fatou's lemma then give

$$g_t(q) \geq \mathbb{E} \left[g_\tau^{v_{t_1}}(q - \beta I_{t_1}^H); v_{t_1} > 0, \beta I_{t_1}^H < q \right].$$

Applying (62) to a sufficiently small neighborhood of $(y_1, (q - z_1)/\beta)$ makes the right-hand side strictly positive.

The argument for m_t is identical. Since

$$\mathbb{E}_{y_1}[v_\tau; Q_\tau < q/2] \rightarrow y_1 > 0 \quad (\tau \downarrow 0),$$

one has $m_\tau^{y_1}(z_1) > 0$ for some $z_1 \in (0, q/2)$, and the same support argument applies. $\square$

Define

$$\phi_t(q) := \frac{m_t(q)}{g_t(q)}, \quad q > 0. \quad (63)$$

By Proposition 7 and Lemma 8, this is jointly continuous, smooth in q , and a version of $\mathbb{E}[v_t \mid Q_t = q]$.

We next determine its behavior at the endpoint $q = 0$. The general saddle-point method is described, for example, by Olver [22, Ch. 4]; the uniform differentiated version needed here is stated and proved explicitly below.

Lemma 9 (Square-root saddle transfer). *Let $\mathcal{T} = [\vartheta_-, \vartheta_+] \subset \mathbb{R}$ be a compact interval and fix $\sigma_0, R_0 > 0$. For $\vartheta \in \mathcal{T}$, let F_ϑ, G_ϑ be analytic on $\{\Re z > 0\}$. Suppose that there are constants $K, c_0 > 0$ and $N \geq 0$, independent of ϑ , such that the following conditions hold:*

- (i) *For each ϑ there exist $A_\vartheta, C_\vartheta, D_\vartheta > 0$, $\alpha_\vartheta \in \mathbb{R}$, and remainder functions $R_{F,\vartheta}, R_{G,\vartheta}$, analytic on a neighborhood of $\{z : \Re z \geq \sigma_0, |z| \geq R_0\}$, such that, with the principal branches of $\sqrt{z}$ and $z^{-\alpha_\vartheta}$,*

$$\begin{aligned} F_\vartheta(z) &= C_\vartheta z^{-\alpha_\vartheta} e^{-A_\vartheta \sqrt{z}} (1 + R_{F,\vartheta}(z)), \\ G_\vartheta(z) &= F_\vartheta(z) \frac{D_\vartheta}{z} (1 + R_{G,\vartheta}(z)) \end{aligned} \quad (64)$$

for $\Re z \geq \sigma_0$ and $|z| \geq R_0$. The coefficients $A_\vartheta, C_\vartheta, D_\vartheta$ are uniformly bounded above and away from zero, α_ϑ is uniformly bounded, and, on the same region,

$$|R_{F,\vartheta}(z)| + |R_{G,\vartheta}(z)| \leq K|z|^{-1/2}. \quad (65)$$

(ii) On the cone where $\Re z \geq \sigma_0$, $|\Im z| \leq \frac{1}{2}\Re z$, and $|z| \geq R_0$, the differentiated remainders satisfy

$$|\partial_z R_{F,\vartheta}(z)| + |\partial_z R_{G,\vartheta}(z)| \leq K|z|^{-3/2}. \quad (66)$$

(iii) On the Bromwich line $\mathcal{C}_{\sigma_0} := \{\sigma_0 + iy : y \in \mathbb{R}\}$,

$$|F_\vartheta(\sigma_0 + iy)| + |G_\vartheta(\sigma_0 + iy)| \leq K(1 + |y|)^N e^{-c_0\sqrt{|y|}}. \quad (67)$$

Let f_ϑ, g_ϑ be their inverse Laplace transforms and suppose these are positive near zero. Then, uniformly for $\vartheta \in \mathcal{T}$,

$$\frac{g_\vartheta(q)}{f_\vartheta(q)} = \frac{4D_\vartheta}{A_\vartheta^2} q^2 (1 + O(q)), \quad \left(\frac{g_\vartheta}{f_\vartheta}\right)'(q) = \frac{8D_\vartheta}{A_\vartheta^2} q + O(q^2). \quad (68)$$

Proof. Suppress ϑ and set

$$z_* := \frac{A^2}{4q^2}.$$

The fixed-line bound (67) makes the Bromwich integrals, and the integrals obtained after multiplication by z , absolutely convergent on $\mathcal{C}_{\sigma_0}$. Hence differentiation under inversion is valid.

We next justify displacement to $\Re z = z_*$. On a horizontal side $z = x \pm iR$, $\sigma_0 \leq x \leq z_*$, the expansion (64)–(65) gives, for $j = 0, 1$,

$$|e^{qz} z^j F(z)| + |e^{qz} z^j G(z)| \leq C_q R^{N_j} e^{-c\sqrt{R}}.$$

Thus the two horizontal integrals tend to zero as $R \rightarrow \infty$. Cauchy's theorem therefore moves both the original and differentiated inversion integrals to $\Re z = z_*$.

For $x > 0$, $\Re \sqrt{x + iy} = \left(\frac{\sqrt{x^2 + y^2} + x}{2}\right)^{1/2}$, and consequently there exists a constant $c > 0$, independent of x and y , such that

$$\Re \sqrt{x + iy} - \sqrt{x} \geq c \begin{cases} y^2/x^{3/2}, & |y| \leq x, \\ \sqrt{|y|}, & |y| \geq x. \end{cases} \quad (69)$$

Writing the shifted contour as $z = z_* + iy$, define its central portion by $|y| \leq z_*^{13/16}$. On this portion put $y = z_*^{3/4}u$, equivalently $|u| \leq z_*^{1/16}$. The saddle identity $q = A/(2\sqrt{z_*})$ cancels the linear term and gives

$$\begin{aligned} iqz_*^{3/4}u - A(\sqrt{z_* + iz_*^{3/4}u} - \sqrt{z_*}) \\ = -\frac{A}{8}u^2 + \frac{iA}{16}z_*^{-1/4}u^3 + O(z_*^{-1/2}(u^4 + u^6)). \end{aligned} \quad (70)$$

Also,

$$\begin{aligned} (1 + iuz_*^{-1/4})^{-\alpha} &= 1 - i\alpha uz_*^{-1/4} + O(u^2 z_*^{-1/2}), \\ \frac{z_*}{z_* + iz_*^{3/4}u} &= 1 - iuz_*^{-1/4} + O(u^2 z_*^{-1/2}). \end{aligned} \quad (71)$$

The order- $z_*^{-1/4}$ terms in (70)–(71) are odd and integrate to zero on the symmetric contour. Equations (69)–(71), the Gaussian majorant e^{-cu^2} , and (65) therefore yield

$$\begin{aligned} f(q) &= \mathcal{P}(q)(1 + \varepsilon_F(q)), \\ g(q) &= \frac{D}{z_*} \mathcal{P}(q)(1 + \varepsilon_G(q)), \end{aligned} \quad (72)$$

where

$$\mathcal{P}(q) := \frac{C}{2\pi} \sqrt{\frac{8\pi}{A}} z_*^{3/4-\alpha} e^{-A^2/(4q)}$$

and

$$|\varepsilon_F(q)| + |\varepsilon_G(q)| \leq Cq. \quad (73)$$

For completeness, Taylor's formula gives (70)–(71) on the central portion. On $z_*^{1/16} \leq |u| \leq z_*^{1/4}$, (69) gives $O(e^{-cz_*^{1/8}})$, while the part $|y| \geq z_*$ is $O(e^{-c\sqrt{z_*}})$ relative to $\mathcal{P}(q)$. These bounds are uniform over $\mathcal{T}$.

For the derivative estimate, return first to $\mathcal{C}_{\sigma_0}$. Absolute convergence in (67) gives

$$f'(q) = \frac{1}{2\pi i} \int_{\mathcal{C}_{\sigma_0}} z e^{qz} F(z) dz, \quad g'(q) = \frac{1}{2\pi i} \int_{\mathcal{C}_{\sigma_0}} z e^{qz} G(z) dz.$$

The horizontal-side estimate above, now with $j = 1$, moves these integrals to $\Re z = z_*$. To track the moving saddle explicitly, put

$$\delta_* := z_*^{-1/4} = \sqrt{\frac{2q}{A}}, \quad \Psi(q, u) := iqz_*^{3/4}u - A(\sqrt{z_* + iz_*^{3/4}u} - \sqrt{z_*}).$$

The change of variables used above gives, for all sufficiently small q , the exact normalized identities

$$\begin{aligned} \frac{f(q)}{\mathcal{P}(q)} &= \sqrt{\frac{A}{8\pi}} \int_{\mathbb{R}} e^{\Psi(q, u)} (1 + iu\delta_*)^{-\alpha} [1 + R_F(z_* + iz_*^{3/4}u)] du, \\ \frac{g(q)}{(D/z_*)\mathcal{P}(q)} &= \sqrt{\frac{A}{8\pi}} \int_{\mathbb{R}} e^{\Psi(q, u)} (1 + iu\delta_*)^{-\alpha-1} [1 + R_F(z_* + iz_*^{3/4}u)] [1 + R_G(z_* + iz_*^{3/4}u)] du. \end{aligned}$$

Implement the split with a smooth even cutoff that equals one for $|u| \leq z_*^{1/16}$ and vanishes for $|u| \geq 2z_*^{1/16}$. The root bound, applied also to the shifted formulas for f' and g' , shows that the outer pieces and their q -derivatives are $O(q^m)$ for every m . On the inner piece, (70)–(66) give, respectively,

$$e^{-Au^2/8} \left[1 + i\delta_* P_F(u) + O\left(\delta_*^2(1 + |u|)^M\right) \right], \quad e^{-Au^2/8} \left[1 + i\delta_* P_G(u) + O\left(\delta_*^2(1 + |u|)^M\right) \right],$$

where

$$P_F(u) = \frac{A}{16} u^3 - \alpha u, \quad P_G(u) = \frac{A}{16} u^3 - (\alpha + 1)u.$$

After differentiation in q , the remainders have a common bound $C(1 + |u|)^{M'} e^{-cu^2}$, uniformly over $\mathcal{T}$. Indeed, $z'_* = -2z_*/q$ and $\delta'_* = \delta_*/(2q)$; the only apparent $q^{-1/2}$ terms are $i\delta'_* P_F$ and $i\delta'_* P_G$. They are odd in u and therefore integrate to zero against the even cutoff. Derivatives of the moving cutoff are supported in the overlap region and are exponentially small by (69). Dominated convergence and (72) consequently give

$$|\varepsilon'_F(q)| + |\varepsilon'_G(q)| \leq C. \quad (74)$$

Since $D/z_* = 4Dq^2/A^2$, (72)–(74) give

$$\frac{g}{f} = \frac{4D}{A^2}q^2(1 + O(q))$$

and, on differentiating,

$$\left(\frac{g}{f}\right)' = \frac{8D}{A^2}q + O(q^2),$$

uniformly over $\mathcal{T}$. □

Proposition 10 (Quadratic boundary pinning). *Uniformly for $t \in [t_0, T]$ with $t_0 > 0$,*

$$\phi_t(q) = \frac{4a}{\beta(v_0 + at)^2}q^2(1 + O(q)), \quad \phi_t'(q) = \frac{8a}{\beta(v_0 + at)^2}q + O(q^2), \quad q \downarrow 0. \quad (75)$$

Consequently, there are $\delta_B > 0$ and $C_B < \infty$ such that, whenever $t \in [t_0, T]$ and $0 < x^*(t) - x \leq \delta_B$,

$$|\partial_x[\phi_t(c(x^*(t) - x))]| \leq C_B(x^*(t) - x). \quad (76)$$

Proof. As $|z| \rightarrow \infty$, uniformly for $\Re z \geq \sigma_0 > 0$ and $t \in [t_0, T]$, there are analytic remainders $R_{L,t}, R_{M,t}$ such that

$$L_t(z) = C_t z^{-a/c^2} \exp\left\{-\frac{(v_0 + at)\sqrt{2\beta}}{c}\sqrt{z}\right\} (1 + R_{L,t}(z)),$$

where $C_t > 0$ is continuous, and

$$\frac{\mathcal{M}_t(z)}{L_t(z)} = a\mathcal{J}_t(z) + v_0\mathcal{H}_t(z) = \frac{2a}{c^2 z} (1 + R_{M,t}(z)).$$

Uniformly for $t \in [t_0, T]$,

$$|R_{L,t}(z)| + |R_{M,t}(z)| \leq C|z|^{-1/2}$$

on $\Re z \geq \sigma_0$. Indeed, the zeroth-order bounds follow directly from

$$d = c\sqrt{2\beta}z(1 + O(|z|^{-1})), \quad 1 - \mathfrak{h} = \frac{2\sqrt{2\beta}}{c}z^{-1/2}(1 + O(|z|^{-1/2})),$$

and the exponential smallness of e^{-dt} . The same expansions and remainder bounds hold uniformly, for large $|z|$, on a wider cone $|\Im z| \leq \kappa\Re z$ for some $\kappa > 1/2$. After increasing R_0 if necessary, choose $\eta \in (0, 1)$ so that, for every z in the cone specified in Lemma 9, the closed disk $\{\zeta : |\zeta - z| \leq \eta|z|\}$ lies in this wider large- $|z|$ region. On that disk the triangle inequality gives $(1 - \eta)|z| \leq |\zeta| \leq (1 + \eta)|z|$, and Cauchy's first-derivative estimate gives

$$\begin{aligned} & |\partial_z R_{L,t}(z)| + |\partial_z R_{M,t}(z)| \\ & \leq \frac{1}{\eta|z|} \left(\sup_{|\zeta - z| = \eta|z|} |R_{L,t}(\zeta)| + \sup_{|\zeta - z| = \eta|z|} |R_{M,t}(\zeta)| \right) \leq C|z|^{-3/2}, \end{aligned}$$

uniformly for $t \in [t_0, T]$. Together with (61), Lemma 9 applies with $\mathcal{T} = [t_0, T]$, $\vartheta = t$, $F_t = L_t$, $G_t = \mathcal{M}_t$, and

$$A_t = \frac{(v_0 + at)\sqrt{2\beta}}{c}, \quad D_t = \frac{2a}{c^2}, \quad \alpha_t = \frac{a}{c^2},$$

and proves (75). Substitution of $q = c(x^*(t) - x)$ gives (76). □

Since $0 \leq v_t \leq Q_t$, the density inequality $0 \leq m_t(q) \leq qg_t(q)$ holds almost everywhere. Continuity makes it valid for every $q > 0$, and hence

$$0 \leq \phi_t(q) \leq q. \quad (77)$$

The canonical log-local variance announced in (55) is

$$\mu(t, x) := \begin{cases} \phi_t(c(x^*(t) - x)), & x < x^*(t), \\ 0, & x \geq x^*(t). \end{cases} \quad (78)$$

Proposition 11 (The canonical projected surface). *There is a Borel function $\lambda : (0, T] \times (0, \infty) \rightarrow [0, \infty)$ with the following properties.*

- (i) *For every $t \in (0, T]$, $\lambda(t, \cdot)$ is a version of $\mathbb{E}[v_t \mid S_t = \cdot]$.*
- (ii) *$0 \leq \lambda(t, s) \leq A(t, s)^+$ everywhere; in particular $\lambda(t, s) = 0$ for $s \geq s^*(t)$, and there is a constant $C_\lambda = C_\lambda(T)$ with*

$$\mu(t, x) \leq C_\lambda(1 + |x|), \quad (t, x) \in (0, T] \times \mathbb{R}. \quad (79)$$

- (iii) *$\lambda(t, s) > 0$ for $0 < s < s^*(t)$, and $\lambda(t, \cdot)$ is C^∞ on $(0, s^*(t))$ for each $t \in (0, T]$.*
- (iv) *For every $\varepsilon \in (0, T)$, λ is jointly continuous on $[\varepsilon, T] \times (0, \infty)$, and $x \mapsto \mu(t, x)$ is Lipschitz on compact sets, uniformly in $t \in [\varepsilon, T]$.*

Proof. Define $\lambda(t, s) := \mu(t, \log s)$. Equations (57) and (63) show that this is a Borel version of $\mathbb{E}[v_t \mid S_t = s]$. The bound (77) gives

$$0 \leq \mu(t, x) \leq c(x^*(t) - x)^+ = A(t, e^x)^+,$$

and hence item (ii), because $A(t, e^x)^+ \leq v_0 + \kappa\theta T + \xi|\log S_0| + \xi|x|$. Strict positivity and interior smoothness follow from Proposition 7 and Lemma 8.

It remains to check the behavior at the barrier. The bound (77) and continuity of x^* show that extension by zero is jointly continuous there. Fix $\varepsilon \in (0, T)$ and a compact interval $J \subset \mathbb{R}$. Taking $t_0 = \varepsilon$ in Proposition 10, choose $\delta_B > 0$ and $C_B < \infty$ so that, whenever $t \in [\varepsilon, T]$ and $0 < x^*(t) - x \leq \delta_B$,

$$|\partial_x \mu(t, x)| \leq C_B(x^*(t) - x) \leq C_B \delta_B.$$

By Proposition 7 and Lemma 8, $\partial_q \phi_t(q)$ is jointly continuous on $(0, T] \times (0, \infty)$. The region given by $t \in [\varepsilon, T]$, $x \in J$, and $x \leq x^*(t) - \delta_B$ is compact, and there $q = c(x^*(t) - x) \geq c\delta_B$. Hence $\partial_x \mu(t, x) = -c\partial_q \phi_t(c(x^*(t) - x))$ is uniformly bounded on that region. For $x \geq x^*(t)$ the function is zero. If $x < x^*(t) \leq y$, then

$$|\mu(t, x) - \mu(t, y)| = \mu(t, x) \leq c(x^*(t) - x) \leq c(y - x).$$

Taking the maximum of the two derivative bounds and c gives a Lipschitz constant on J independent of $t \in [\varepsilon, T]$. Since ε and J were arbitrary, this proves the uniform local Lipschitz assertion. Joint continuity of λ follows by composition with $(t, s) \mapsto (t, \log s)$. $\square$

Theorem 12 (Well-posedness on every interval $[\varepsilon, T]$ with $\varepsilon > 0$). *Let $0 < \varepsilon \leq T$ and $x \in \mathbb{R}$. Equation (56) on $[\varepsilon, T]$ from x has a weak solution with continuous real-valued paths; pathwise uniqueness holds among such solutions, and consequently the equation has a unique strong solution and is well posed in law from every (ε, x) .*

Proof. By Proposition 11(iv), μ is continuous and locally Lipschitz in its spatial variable, locally uniformly in time. Moreover,

$$|\sqrt{\mu(t, x)} - \sqrt{\mu(t, y)}|^2 \leq |\mu(t, x) - \mu(t, y)|.$$

Thus the one-dimensional criterion of Ethier and Kurtz [11, Ch. 5, Thm. 3.8 and Rem. 3.9, pp. 298–299], applied up to successive exits from bounded intervals, gives pathwise uniqueness.

Shift the initial time to zero and extend the coefficient constantly after T : for $r \geq 0$, set

$$\mu_\varepsilon(r, x) := \mu(\varepsilon + (r \wedge (T - \varepsilon)), x).$$

Then define

$$\sigma_\varepsilon(r, x) := \sqrt{\mu_\varepsilon(r, x)}, \quad b_\varepsilon(r, x) := -\frac{1}{2}\mu_\varepsilon(r, x).$$

The growth estimate (79) implies, for a suitable C'_λ ,

$$|\sigma_\varepsilon(r, x)|^2 = \mu_\varepsilon(r, x) \leq C'_\lambda(1 + x^2), \quad xb_\varepsilon(r, x) = -\frac{1}{2}x\mu_\varepsilon(r, x) \leq C'_\lambda(1 + x^2).$$

For the second inequality, the left-hand side is nonpositive when $x \geq 0$, whereas for $x < 0$ it follows directly from $\mu_\varepsilon(r, x) \leq C_\lambda(1 + |x|)$. These bounds and continuity verify the hypotheses of the existence and martingale-problem/SDE representation theorems of Ethier and Kurtz [11, Ch. 5, Thms. 3.10 and 3.3, pp. 299 and 293–295]. Together they give a nonexplosive weak solution Y from $Y_0 = x$; then $\bar{X}_u := Y_{u-\varepsilon}$, $u \in [\varepsilon, T]$, solves (56). Yamada and Watanabe [24] then yield a unique strong solution and uniqueness in law. $\square$

For $\varepsilon \in (0, T)$ let $C_\varepsilon := C([\varepsilon, T]; \mathbb{R})$ with its Borel σ -algebra, and let Π_ε denote the restriction of a path on a larger interval to $[\varepsilon, T]$.

Lemma 13 (Restart kernels). *Fix $\varepsilon \in (0, T)$, let $\bar{X}$ be a solution of (56) on $[r, T]$, where $0 \leq r < \varepsilon$, from any initial value, and put $\mathbf{m}_\varepsilon := \text{Law}(\bar{X}_\varepsilon)$. There is a Borel probability kernel $x \mapsto P_x^{(\varepsilon, \bar{X})}$ from $\mathbb{R}$ to the probability measures on C_ε such that:*

- (a) *For $\mathbf{m}_\varepsilon$ -almost every x , $P_x^{(\varepsilon, \bar{X})}$ is the unique law of a solution of (56) on $[\varepsilon, T]$ from x .*
- (b) *For every bounded Borel functional G on C_ε ,*

$$\mathbb{E}[G(\Pi_\varepsilon \bar{X}) | \mathcal{F}_\varepsilon] = \int G \, dP_{\bar{X}_\varepsilon}^{(\varepsilon, \bar{X})} \quad a.s.$$

Proof. Fix $\varepsilon > 0$ and extend μ after T by $\hat{\mu}(\tau, x) := \mu(\tau \wedge T, x)$ on $[\varepsilon, \infty) \times \mathbb{R}$. Define the autonomous operator

$$\mathcal{A}F(\tau, x) := F_\tau(\tau, x) + \frac{1}{2}\hat{\mu}(\tau, x)(F_{xx}(\tau, x) - F_x(\tau, x)). \quad (80)$$

If X is a weak solution of (56) on $[\tau, T]$ from $X_\tau = x$, set

$$(\mathbb{T}_u, Y_u) := (\tau + u, X_{\tau+u}), \quad 0 \leq u \leq T - \tau.$$

Itô's formula shows that $(\mathbb{T}, Y)$ solves the $\mathcal{A}$ -martingale problem from (τ, x) . Conversely, testing a solution of that problem with localized versions of $(\tau, x) \mapsto \tau$ and $(\tau, x) \mapsto \tau^2$ shows that its first coordinate is $\tau + u$; the martingale-problem/SDE representation of Ethier and Kurtz [11, Ch. 5, Thm. 3.3] then realizes its second coordinate as a weak solution of (56). Theorem 12 therefore gives uniqueness for the lifted martingale problem from every (τ, x) with $\tau \in [\varepsilon, T]$. Let Q^ω be a regular conditional distribution of $\Pi_\varepsilon \bar{X}$ given $\mathcal{F}_\varepsilon$, and let $x \mapsto \tilde{Q}_x$ be a regular

conditional distribution of $\Pi_\varepsilon \bar{X}$ given $\bar{X}_\varepsilon = x$. These kernels exist because $\mathbb{R}$ and C_ε are standard Borel spaces. The defining initial-coordinate relation gives

$$\begin{aligned} Q^\omega \{\eta : \eta(\varepsilon) = \bar{X}_\varepsilon(\omega)\} &= 1 \quad \text{for almost every } \omega, \\ \tilde{Q}_x \{\eta : \eta(\varepsilon) = x\} &= 1 \quad \text{for } \mathbf{m}_\varepsilon\text{-almost every } x. \end{aligned}$$

The deterministic-time conditioning argument in the proof of Ethier and Kurtz [11, Thm. 4.4.2], applied to a countable determining class of test functions, rational time pairs, and a countable cylinder algebra, shows that, after the deterministic lift $\eta \mapsto (u \mapsto (\varepsilon + u, \eta(\varepsilon + u)))$, Q^ω solves the lifted martingale problem from $(\varepsilon, \bar{X}_\varepsilon(\omega))$ for almost every ω , and that the analogous lift of $\tilde{Q}_x$ solves it from (ε, x) for $\mathbf{m}_\varepsilon$ -almost every x . Since $\mathbf{m}_\varepsilon = \text{Law}(\bar{X}_\varepsilon)$, the random variable $\bar{X}_\varepsilon$ belongs to the latter full set almost surely. Thus the two conditional laws solve the problem from the same point for almost every ω , and uniqueness gives

$$Q^\omega = \tilde{Q}_{\bar{X}_\varepsilon(\omega)} \quad \text{for almost every } \omega.$$

Taking $P_x^{(\varepsilon, \bar{X})} := \tilde{Q}_x$ proves both assertions. $\square$

C Localized density and strict positivity

Proof of Lemma 2. The sign of the Brownian term does not affect the law of the CIR process. Apply the Lamperti transformation

$$R_t := \frac{2\sqrt{v_t}}{\xi}, \quad \gamma := \frac{\xi^2}{4}, \quad \delta_{\text{CIR}} := \frac{4\kappa\theta}{\xi^2}.$$

Up to replacing the Brownian motion by its negative, on $R_t > 0$ the pair (I_t^H, R_t) satisfies

$$dI_t^H = \gamma R_t^2 dt, \quad dR_t = \left(\frac{\delta_{\text{CIR}} - 1}{2R_t} - \frac{\kappa R_t}{2} \right) dt + dB_t. \quad (81)$$

Fix $t > 0$ and a target $(i_1, r_1) \in (0, \infty)^2$, and put $r_0 := 2\sqrt{v_0}/\xi$. First choose a positive C^1 path $r^{(0)} : [0, t] \rightarrow (0, \infty)$ such that

$$r^{(0)}(0) = r_0, \quad r^{(0)}(t) = r_1, \quad \gamma \int_0^t (r_q^{(0)})^2 dq < i_1.$$

To construct it, choose $\tau \in (0, t/2)$ and $h > 0$, use positive monotone C^1 profiles to join r_0 to h on $[0, \tau]$ and h to r_1 on $[t - \tau, t]$, with zero derivative where they meet the constant middle segment, and set $r^{(0)} = h$ on $[\tau, t - \tau]$. Taking τ and h sufficiently small gives the displayed strict inequality.

Choose a nonzero $\psi \in C_c^\infty((\tau, t - \tau))$ with $\psi \geq 0$. The map

$$\ell \mapsto \gamma \int_0^t (r_q^{(0)} + \ell \psi_q)^2 dq, \quad \ell \geq 0,$$

is continuous and strictly increasing from a value below i_1 to infinity. Hence, by the intermediate value theorem, it equals i_1 for some $\ell_* > 0$. Set $\bar{r} := r^{(0)} + \ell_* \psi$. This is a positive C^1 path with the prescribed endpoints and $\gamma \int_0^t \bar{r}_q^2 dq = i_1$. The control

$$u_q = \dot{\bar{r}}_q - \frac{\delta_{\text{CIR}} - 1}{2\bar{r}_q} + \frac{\kappa \bar{r}_q}{2}$$

is continuous, since $\bar{r}$ is bounded away from zero on $[0, t]$. The corresponding skeleton path is

$$\left(\gamma \int_0^q \bar{r}_u^2 du, \bar{r}_q \right), \quad 0 \leq q \leq t,$$

and therefore runs from $(0, r_0)$ to (i_1, r_1) .

Write

$$b_R(r) := \frac{\delta_{\text{CIR}} - 1}{2r} - \frac{\kappa r}{2}.$$

For a control v near u in $L^2(0, t)$, let r^v solve

$$\dot{r}_q^v = b_R(r_q^v) + v_q, \quad r_0^v = r_0,$$

and define its endpoint by

$$\mathcal{E}_t(v) := \left(\gamma \int_0^t (r_q^v)^2 \, dq, r_t^v \right).$$

The localized inverse-function argument used below requires more than reachability of the target: the chosen control must be *regular*, meaning that

$$D\mathcal{E}_t(u) : L^2(0, t) \longrightarrow \mathbb{R}^2$$

is onto. To verify this, fix $k \in L^2(0, t)$ and let $z := Dr^u[k]$ be the corresponding first variation. Since $r^u = \bar{r}$,

$$\dot{z}_q = b'_R(\bar{r}_q)z_q + k_q, \quad z_0 = 0,$$

and

$$D\mathcal{E}_t(u)[k] = \left(2\gamma \int_0^t \bar{r}_q z_q \, dq, z_t \right).$$

Solving the variational equation and changing the order of integration show that these two components are the $L^2(0, t)$ pairings of k with

$$q_1(s) := 2\gamma \int_s^t \bar{r}_q \exp\left\{ \int_s^q b'_R(\bar{r}_\ell) \, d\ell \right\} \, dq, \quad q_2(s) := \exp\left\{ \int_s^t b'_R(\bar{r}_\ell) \, d\ell \right\}. \quad (82)$$

The functions q_1 and q_2 are linearly independent: $q_1/q_2 > 0$ for $s < t$ and tends to zero as $s \uparrow t$. Hence the endpoint derivative is onto, so u is regular; equivalently, its deterministic Malliavin covariance is invertible.

To localize, let $\bar{i}_q := \gamma \int_0^q \bar{r}_\ell^2 \, d\ell$ and choose a bounded open set D such that

$$\{(\bar{i}_q, \bar{r}_q) : 0 \leq q \leq t\} \Subset D \Subset \mathbb{R} \times (0, \infty).$$

Choose open sets D_0, D_1 with

$$\{(\bar{i}_q, \bar{r}_q) : 0 \leq q \leq t\} \subset D_0 \Subset D_1 \Subset D.$$

On a neighborhood of $\bar{D}$, define the original vector fields by

$$X_1 := \partial_r, \quad X_0 := \gamma r^2 \partial_i + b_R(r) \partial_r;$$

then

$$[X_1, X_0] = 2\gamma r \partial_i + b'_R(r) \partial_r.$$

Choose a smooth cutoff supported in $\mathbb{R} \times (0, \infty)$ and equal to one on a neighborhood of $\bar{D}$. Multiplying X_0 by this cutoff gives a globally C_b^∞ drift, while X_1 is left unchanged. Thus the cutoff and original fields agree on D , where the parabolic Hörmander condition holds.

Let τ_D be the first exit time from D . We spell out why the regular control found above gives positivity for the killed, rather than merely the globally cut-off, diffusion. The controlled solution map and its first two endpoint derivatives are continuous at $H_q := \int_0^q u_\ell \, d\ell$. Because the controlled path is contained in D_0 , the Wiener-space neighborhood used in the inverse-function argument may therefore be chosen so that the controlled paths, together with all Cameron–Martin translates whose parameters lie in the compact ball used in that argument, remain in D_1

up to time t . On this neighborhood the cutoff diffusion and the original diffusion coincide and $t < \tau_D$. This neighborhood has positive Wiener measure because H is a Cameron–Martin path and Wiener measure has full support in the uniform topology.

Now apply the localized Girsanov and inverse-function calculation used by Herzog and Mattingly [15, proof of Lemma 4.5 in Appendix A]. Their ball-survival indicator is used only to ensure that the translated paths remain in the region where the original and cutoff coefficients coincide; replacing it here by $\mathbf{1}_{\{t < \tau_D\}}$ leaves that calculation unchanged. Since the derivative of the endpoint map is onto by (82), the calculation gives

$$p_t^D((0, 2\sqrt{v_0}/\xi), (i_1, r_1)) > 0,$$

where p^D is the density of the sub-Markov kernel

$$P_u^D(x, E) := \mathbb{P}_x\{(I_u^H, R_u) \in E, u < \tau_D\}, \quad E \subset D.$$

For the fixed initial point $x_0 = (0, 2\sqrt{v_0}/\xi)$, local parabolic Hörmander regularity makes $(u, y) \mapsto p_u^D(x_0, y)$ smooth locally on $(0, \infty) \times D$; this is the stopped-kernel formulation described by Herzog and Mattingly [15, Remark 2.4], applied on nested balls compactly contained in D . Hence the displayed positivity persists on a space–time neighborhood of (t, i_1, r_1) . Finally, $P_u^D(x, \cdot) \leq \text{Law}_x(I_u^H, R_u)$ on D , so this killed kernel supplies the required positive subdensity of the full transition law.

For each $u > 0$, the pathwise identity makes

$$\Theta_u(i, r) := \left(S_0 \exp \left[\frac{v_0 + \kappa \theta u - \gamma r^2 - \beta i}{\xi} \right], i \right)$$

a smooth bijection from $(0, \infty)^2$ onto the spatial section of $\mathcal{O}$ at time u , with a strictly positive Jacobian. Transporting the killed density through Θ_u and extending it by zero therefore gives, around every point $z \in \mathcal{O}$, a jointly Borel subdensity q^z of Γ_u that is bounded below by a positive constant on some space–time neighborhood U_z of z .

Fix a compact set $\mathcal{K}$ contained entirely in $\mathcal{O}$ and choose a finite subcover $U_{z_1}, \dots, U_{z_N}$. Put

$$q_{\mathcal{K}} := \max_{1 \leq j \leq N} q^{z_j}.$$

Multiplying this function by the indicator of the time-projection of $\mathcal{K}$, if necessary, makes it vanish outside that projection. For fixed u , every $q^{z_j}(u, \cdot, \cdot)$ $ds \, di$ is dominated by Γ_u . Hence it is bounded almost everywhere by the density of the absolutely continuous part of Γ_u ; the same is true of their finite maximum. Thus $q_{\mathcal{K}}$ satisfies the measure domination in (42), while the finite cover gives its strictly positive lower bound on $\mathcal{K}$.

It remains to record the asserted regularity of the spot marginal. Put $Q_t := v_t + \beta I_t^H = A(t, S_t)$. By Proposition 7 and Lemma 8, Q_t has a density $g_t(q)$ that is jointly continuous and strictly positive on $(0, T] \times (0, \infty)$. Since $|\partial_s A(t, s)| = \xi/s$, the jointly continuous version of the spot density is

$$f_t^S(s) = \frac{\xi}{s} g_t(A(t, s)), \quad 0 < t \leq T, \quad 0 < s < s^*(t).$$

It is strictly positive and locally bounded on the displayed region; from now on, extend f_t^S by zero for $s \geq s^*(t)$.

Fix $t \in (0, T]$ and a regular conditional law $s \mapsto \eta_{t,s}$. Since the first marginal of Γ_t is $f_t^S(s) \, ds$,

$$\Gamma_t(ds, di) = f_t^S(s) \eta_{t,s}(di) \, ds.$$

For a compact set $\mathcal{K} \Subset \mathcal{O}$, subtracting the measure in (42) leaves a positive measure. Testing on rectangles whose i -component ranges over a countable determining class, and then applying the monotone-class theorem, gives (43) for this fixed t .

It remains to justify the asserted joint choice. Let $J = [t_0, T] \Subset (0, \infty)$. Continuity of the sample paths implies that $t \mapsto \Gamma_t$ is Borel (indeed, $t \mapsto \int \varphi \, d\Gamma_t$ is continuous for every bounded continuous φ). Hence

$$G_J(dt, ds, di) := \mathbf{1}_J(t) \, dt \, \Gamma_t(ds, di)$$

is a finite Borel measure whose (t, s) -marginal is $\mathbf{1}_J(t) f_t^S(s) \, dt \, ds$. Since the spaces are standard Borel, disintegration gives a jointly Borel probability kernel $(t, s) \mapsto \eta_{t,s}$ such that

$$G_J(dt, ds, di) = \mathbf{1}_J(t) f_t^S(s) \eta_{t,s}(di) \, dt \, ds.$$

Fubini's theorem, applied first to a countable determining class, shows that $s \mapsto \eta_{t,s}$ is a regular conditional law at time t for dt -almost every $t \in J$. Integrating (42) over J gives

$$G_J(dt, ds, di) \geq \mathbf{1}_J(t) q_K(t, s, i) \, dt \, ds \, di.$$

Testing the difference on rectangles, again using a countable determining class in the i -variable and the monotone-class theorem, yields

$$f_t^S(s) \eta_{t,s}(di) \geq q_K(t, s, i) \, di \quad \text{for } dt \, ds\text{-almost every } (t, s) \in J \times (0, \infty).$$

This proves the joint assertion. $\square$

D Smooth comparison and approximation

This appendix proves the smooth comparison and approximation results used in the proofs of Theorems 1 and 3.

D.1 The smooth comparison

Work on an interval $[r_0, T] \subseteq [0, T]$ with the extended state

$$\begin{aligned} dX_u &= -\frac{1}{2}\nu(u, X_u) \, du + \sqrt{\nu(u, X_u)} \, dB_u, \\ dI_u &= \nu(u, X_u) \, du, \quad (X_r, I_r) = (x, i), \end{aligned} \tag{83}$$

for a surface ν on $[r_0, T] \times \mathbb{R}$, and let

$$\mathcal{D} := \frac{1}{2}(\partial_{xx} - \partial_x) + \partial_i,$$

so that the backward generator of (83) is $\nu(r, x)\mathcal{D}$.

Lemma 14 (Smooth case). *Let $\nu : [r_0, T] \times \mathbb{R} \rightarrow (0, \infty)$ be jointly continuous and C^∞ in x , with $\inf \nu > 0$ and, for some $C_\nu < \infty$,*

$$\nu(r, x) \leq C_\nu(1 + |x|).$$

Assume that every positive-order spatial derivative of ν is bounded and jointly continuous. Let $\psi \in C^\infty(\mathbb{R})$ be nondecreasing and of at most linear growth, with every positive-order derivative bounded, and put

$$U(r, x, i) := \mathbb{E}_{r,x} \left[\psi \left(i + \int_r^T \nu(u, X_u) \, du \right) \right], \quad h := \mathcal{D}U. \tag{84}$$

Then $U \in C^{1,2,1}$, with one-sided time derivatives at the endpoints, and

$$U_r + \nu \mathcal{D}U = 0, \quad U(T, x, i) = \psi(i).$$

The functions U_x and h are bounded, $h \geq 0$ on $[r_0, T] \times \mathbb{R} \times \mathbb{R}$, and, if ψ is convex, the map $i \mapsto h(r, x, i)$ is nondecreasing. Moreover, for $r_0 \leq t \leq t' \leq T$,

$$U(t, x, i) - U(t', x, i) = \int_t^{t'} \nu(r, x) h(r, x, i) \, dr \geq 0. \tag{85}$$

Proof. Regularity. The coefficients $-\nu/2$ and $\sqrt{\nu}$ have linear growth and are globally Lipschitz in x . Their positive-order spatial derivatives are bounded; for $\sqrt{\nu}$ this also uses $\inf \nu > 0$. Consequently the equation is nonexplosive and the stochastic flow $x \mapsto X_u^{r,x}$ is C^∞ , with

$$\sup_{r \in [r_0, T]} \sup_{x \in \mathbb{R}} \mathbb{E} \left[\sup_{u \in [r, T]} |\partial_x^m X_u^{r,x}|^p \right] < \infty \quad (m \geq 1, p < \infty)$$

and local uniform continuity in (r, x) ; see Kunita [18, Ch. 4, Thm. 4.6.5]. The same estimates apply to the positive-order spatial derivatives of $\int_r^T \nu(u, X_u^{r,x}) du$. Differentiating under the expectation in (84), and using the translation structure in i , shows that U is smooth in (x, i) , that U_x and all the positive-order spatial derivatives used below are bounded, and that these derivatives are jointly continuous in (r, x, i) up to $r = T$. The function U itself has at most linear growth.

The backward equation. For $r < q \leq T$, the Markov property and Itô's formula, first localized and then delocalized using the preceding linear-growth and moment bounds, applied on $[r, q]$ to the fixed function $(x, i) \mapsto U(q, x, i)$ give

$$U(r, x, i) - U(q, x, i) = \mathbb{E}_{r,x,i} \left[\int_r^q \nu(u, X_u) \mathcal{D}U(q, X_u, I_u) du \right]. \quad (86)$$

Dividing by $q - r$ and letting $q \downarrow r$ identifies the right time derivative for $r < T$; the analogous identity from an earlier starting time identifies the left derivative for $r > r_0$, and the two agree on the interior. Hence, with one-sided derivatives at the endpoints,

$$U_r = -\nu \mathcal{D}U = -\nu h \quad \text{on } [r_0, T], \quad U(T, x, i) = \psi(i), \quad (87)$$

and U_r is jointly continuous by the continuity established above. The integral form of (87),

$$U(r, \cdot) - U(q, \cdot) = \int_r^q \nu(u, \cdot) h(u, \cdot) du,$$

may be differentiated in (x, i) under the time integral, because the integrand has locally bounded continuous spatial derivatives of the orders appearing in $\mathcal{D}$. Hence $h_r = -\mathcal{D}(\nu h)$ on $[r_0, T]$, with one-sided time derivatives at the endpoints; in particular $h \in C^{1,2,1}$ up to $r = T$ and, expanding,

$$\begin{aligned} h_r + \frac{1}{2} \nu h_{xx} + (\nu_x - \frac{1}{2} \nu) h_x + \nu h_i + \varpi h &= 0, \\ h(T, x, i) = \psi'(i) &\geq 0, \quad \varpi := \frac{1}{2} (\nu_{xx} - \nu_x). \end{aligned} \quad (88)$$

For the present smooth surface, ϖ is bounded.

Positivity of h . The auxiliary diffusion

$$\begin{aligned} d\hat{X}_u &= (\nu_x - \frac{1}{2} \nu)(u, \hat{X}_u) du + \sqrt{\nu(u, \hat{X}_u)} d\hat{B}_u, \\ d\hat{I}_u &= \nu(u, \hat{X}_u) du, \quad (\hat{X}_r, \hat{I}_r) = (x, i), \end{aligned} \quad (89)$$

has globally Lipschitz coefficients of linear growth and is nonexplosive. Put

$$\mathcal{E}_u := \exp \left\{ \int_r^u \varpi(q, \hat{X}_q) dq \right\}.$$

For $r < T$, $0 < \delta < T - r$, and $r \leq u \leq T - \delta$, Itô's formula and (88) give

$$\mathcal{E}_u h(u, \hat{X}_u, \hat{I}_u) = h(r, x, i) + \int_r^u \mathcal{E}_q \sqrt{\nu(q, \hat{X}_q)} h_x(q, \hat{X}_q, \hat{I}_q) d\hat{B}_q.$$

Because ϖ and h_x are bounded, while ν has linear growth and $\hat{X}$ has finite moments, the stochastic integral is a true martingale. Let $\hat{\mathbb{E}}_{r,x}$ denote expectation for the law of $\hat{X}$ in (89)

started from $\widehat{X}_r = x$; the initial i -coordinate is displayed explicitly below. Taking expectations at $u = T - \delta$ and then letting $\delta \downarrow 0$, using boundedness and continuity, gives

$$h(r, x, i) = \widehat{\mathbb{E}}_{r,x} \left[\exp \left\{ \int_r^T \varpi(u, \widehat{X}_u) du \right\} \psi' \left(i + \int_r^T \nu(u, \widehat{X}_u) du \right) \right] \geq 0. \quad (90)$$

For $r = T$, the same conclusion is the terminal condition $h(T, x, i) = \psi'(i)$. (In particular, $0 \leq h \leq e^{(T-r_0)\|\varpi\|_\infty} \|\psi'\|_\infty$.) If ψ is convex, then ψ' is nondecreasing; the auxiliary law in (90) does not otherwise depend on the initial value i , so the same representation shows that $i \mapsto h(r, x, i)$ is nondecreasing. (No uniqueness for the terminal-value problem (88) is invoked; the representation is read off the martingale identity.) No estimate uniform in an approximation parameter will be used.

Conclusion. At fixed (x, i) , (87) and the joint continuity of U_r give $U(t, x, i) - U(t', x, i) = \int_t^{t'} \nu(r, x) h(r, x, i) dr$, which is nonnegative by (90) and $\nu \geq 0$. This is (85). $\square$

D.2 Removal of smoothness

Lemma 15 (Smooth approximation under linear growth). *Let $r \in [0, T)$, and let $\nu : [r, T] \times \mathbb{R} \rightarrow [0, \infty)$ be continuous with, for some $C_\nu < \infty$,*

$$\nu(u, x) \leq C_\nu(1 + |x|), \quad (u, x) \in [r, T] \times \mathbb{R}. \quad (91)$$

Fix $(x, i) \in \mathbb{R} \times [0, \infty)$, and suppose that the SDE

$$dX_u = -\frac{1}{2}\nu(u, X_u) du + \sqrt{\nu(u, X_u)} dB_u, \quad u \in [r, T], \quad X_r = x, \quad (92)$$

has a weak solution with continuous real-valued paths whose law on $C([r, T]; \mathbb{R})$ is unique, and set $I_u := i + \int_r^u \nu(w, X_w) dw$. Then there exist a constant $C'_\nu < \infty$ and surfaces ν_n on $[r, T] \times \mathbb{R}$, jointly continuous, C^∞ in x with every positive-order spatial derivative bounded and jointly continuous, satisfying

$$\begin{aligned} \nu_n &\geq \nu + \frac{1}{n}, & \nu_n(u, y) &\leq C'_\nu(1 + |y|) \quad \text{for all } n, \\ \sup_{(u,y) \in [r,T] \times [-R,R]} |\nu_n(u, y) - \nu(u, y)| &\longrightarrow 0 \quad \text{for every } R > 0, \end{aligned} \quad (93)$$

whose extended systems (83) from (r, x, i) satisfy

$$(X^n, I^n) \Longrightarrow (X, I) \quad \text{in } C([r, T]; \mathbb{R} \times \mathbb{R}_+). \quad (94)$$

The construction of ν_n does not involve the starting point.

Proof. Construction. Let $\chi \in C_c^\infty(\mathbb{R})$ be nonnegative, supported in $[-1, 1]$, with $\int \chi = 1$, and set

$$c_\chi := \int_{\mathbb{R}} |z| \chi(z) dz, \quad L_n := C_\nu + n, \quad \delta_n := L_n^{-3}.$$

The least L_n -Lipschitz majorant of $\nu(u, \cdot)$ is

$$g_n(u, x) := \sup_{y \in \mathbb{R}} \{ \nu(u, y) - L_n |x - y| \}.$$

By (91),

$$\nu(u, x) \leq g_n(u, x) \leq C_\nu(1 + |x|). \quad (95)$$

Indeed,

$$\nu(u, y) - L_n |x - y| \leq C_\nu(1 + |x|) - (L_n - C_\nu) |x - y|.$$

The function $g_n(u, \cdot)$ is L_n -Lipschitz. Its supremum is attained, and, if $|x| \leq R$, every maximizer y satisfies

$$|x - y| \leq \frac{C_\nu(1 + R)}{L_n - C_\nu} = \frac{C_\nu(1 + R)}{n}.$$

Thus the maximizers remain in a common compact set locally in x . The maximum theorem makes g_n jointly continuous, and the displayed bound together with uniform continuity of ν on compact sets shows that $g_n \rightarrow \nu$ locally uniformly.

Mollify only in the spatial variable and correct the possible undershoot:

$$\nu_n(u, x) := \int_{\mathbb{R}} g_n(u, x - \delta_n z) \chi(z) \, dz + L_n \delta_n c_\chi + \frac{1}{n}. \quad (96)$$

The Lipschitz property gives

$$\left| \int_{\mathbb{R}} g_n(u, x - \delta_n z) \chi(z) \, dz - g_n(u, x) \right| \leq L_n \delta_n c_\chi,$$

so $\nu_n \geq g_n + 1/n \geq \nu + 1/n$. Equations (95)–(96) give the uniform linear-growth bound and local uniform convergence in (93), since $L_n \delta_n = L_n^{-2} \rightarrow 0$. Spatial convolution gives all the asserted smoothness. Each positive-order spatial derivative is bounded: after differentiating the mollifier, subtract the constant $g_n(u, x)$ and use the L_n -Lipschitz bound. Joint continuity of those derivatives follows from the joint continuity of g_n and dominated convergence.

Step 1: uniform moments and tightness. Constants below depend only on T , C'_ν and the starting point. For each fixed n , the coefficient is globally Lipschitz and of linear growth, so the usual localized SDE estimate gives $\mathbb{E}[\sup_{[r, T]} |X^n|^4] < \infty$. For $p \in \{2, 4\}$ and $r \leq u \leq w \leq T$, Hölder's inequality and (93) give

$$\mathbb{E}\left[\left(\int_u^w \nu_n(q, X_q^n) \, dq\right)^p\right] \leq (w - u)^{p-1} (C'_\nu)^p \int_u^w \mathbb{E}[(1 + |X_q^n|)^p] \, dq.$$

Combining this (drift and quadratic variation) with the Burkholder–Davis–Gundy inequality and Gronwall's lemma yields $\sup_n \mathbb{E}[\sup_{[r, T]} |X^n|^4] \leq C_T(1 + |x|^4)$, and then, for $r \leq u \leq w \leq T$,

$$\mathbb{E}[|X_w^n - X_u^n|^4] \leq C(1 + |x|^4)(w - u)^2, \quad \mathbb{E}[|I_w^n - I_u^n|^4] \leq C(1 + |x|^4)(w - u)^4.$$

Writing $Z^n := (X^n, I^n)$ and using equivalence of norms in $\mathbb{R}^2$, the two displayed estimates imply

$$\mathbb{E}[\|Z_w^n - Z_u^n\|^4] \leq C(1 + |x|^4)(w - u)^2.$$

Since all Z^n start from the fixed point (x, i) , Kolmogorov's tightness criterion applies directly: the displayed bound gives a modulus of continuity uniformly in probability. Hence their laws are tight in $C([r, T]; \mathbb{R} \times \mathbb{R}_+)$. Equivalently, for every $\eta > 0$,

$$\lim_{\delta \downarrow 0} \sup_n \mathbb{P}\left(\sup_{\substack{u, w \in [r, T] \\ |u - w| \leq \delta}} \|Z_w^n - Z_u^n\| > \eta\right) = 0.$$

In particular, every subsequential limit is supported on continuous paths.

Step 2: identification of limits. Let $\tilde{Z} = (\tilde{X}, \tilde{I})$ be the limit of a weakly convergent subsequence; its paths are continuous and $\tilde{Z}_r = (x, i)$. Fix $f \in C_c^\infty(\mathbb{R}^2)$, times $r \leq s_1 \leq \dots \leq s_m \leq u \leq w \leq T$ and $g_1, \dots, g_m \in C_b(\mathbb{R}^2)$. Each (X^n, I^n) satisfies

$$\mathbb{E}\left[\left(f(Z_w^n) - f(Z_u^n) - \int_u^w \nu_n(q, X_q^n) \mathcal{D}f(Z_q^n) \, dq\right) \prod_{k \leq m} g_k(Z_{s_k}^n)\right] = 0.$$

The expression inside the expectation is bounded and continuous on path space, and $\nu_n \mathcal{D}f \rightarrow \nu \mathcal{D}f$ uniformly because the difference is supported by the compact support of f . Passing to the limit, for every $f \in C_c^\infty(\mathbb{R}^2)$,

$$f(\tilde{Z}_w) - \int_r^w \nu(q, \tilde{X}_q) \mathcal{D}f(\tilde{Z}_q) \, dq$$

is a martingale for the natural filtration of $\tilde{Z}$. Because $\tilde{Z}$ has continuous paths, the stopping times

$$S_m := T \wedge \inf\{q \in [r, T] : |\tilde{X}_q| \vee \tilde{I}_q \geq m\}, \quad \inf \emptyset := \infty,$$

satisfy $S_m \uparrow T$ almost surely. Applying the martingale property, stopped at S_m , to smooth compactly supported test functions that agree on $[-m, m] \times [0, m]$ with

$$(y, z) \mapsto \varphi(y), \quad (y, z) \mapsto z, \quad (y, z) \mapsto z^2,$$

where $\varphi \in C_c^\infty(\mathbb{R})$, shows successively that

$$M_u^\varphi := \varphi(\tilde{X}_u) - \varphi(x) - \int_r^u \nu(q, \tilde{X}_q) \left[\frac{1}{2} \varphi''(\tilde{X}_q) - \frac{1}{2} \varphi'(\tilde{X}_q) \right] \, dq$$

and $N_u := \tilde{I}_u - i - \int_r^u \nu(q, \tilde{X}_q) \, dq$ are continuous local martingales, while the z^2 test gives $\langle N \rangle \equiv 0$; hence $\tilde{I} = i + \int_r^\cdot \nu(q, \tilde{X}_q) \, dq$. Moreover, the process M^φ is bounded, because ν is bounded on $[r, T] \times \text{supp } \varphi$; a bounded local martingale is a martingale, so $\tilde{X}$ solves the martingale problem associated with (92) for test functions in $C_c^\infty(\mathbb{R})$. Extend ν outside $[r, T]$ by freezing it at the endpoints and define

$$\mathcal{A}_\nu F(\tau, x) := F_\tau(\tau, x) + \frac{1}{2} \nu(\tau, x) (F_{xx}(\tau, x) - F_x(\tau, x)).$$

Put $\mathsf{T}_u := r + u$ and $Y_u := \tilde{X}_{r+u}$, $0 \leq u \leq T - r$. Integration by parts, first for $F(\tau, x) = \alpha(\tau)\varphi(x)$ and then by approximation for every $F \in C_c^\infty(\mathbb{R}^2)$, shows that

$$F(\mathsf{T}_u, Y_u) - F(r, x) - \int_0^u \mathcal{A}_\nu F(\mathsf{T}_q, Y_q) \, dq$$

is a martingale. Thus (T, Y) solves the autonomous $\mathcal{A}_\nu$ -martingale problem. Its coefficients are Borel and locally bounded, so the representation theorem of Ethier and Kurtz [11, Ch. 5, Thm. 3.3] yields, on an enlargement of the probability space, a Brownian motion B such that

$$d\mathsf{T}_u = du, \quad dY_u = -\frac{1}{2} \nu(\mathsf{T}_u, Y_u) \, du + \sqrt{\nu(\mathsf{T}_u, Y_u)} \, dB_u.$$

Since $\mathsf{T}_u = r + u$, translating back to calendar time makes $\tilde{X}$ a weak solution of (92) from (r, x) . Uniqueness in law gives $\tilde{X} \stackrel{d}{=} X$, and therefore $\tilde{Z} \stackrel{d}{=} (X, I)$, since $\tilde{I}$ and I are the same functional of $\tilde{X}$ and X . Every subsequence thus has a further subsequence converging to the one law $\text{Law}(X, I)$, which proves (94). $\square$

Declaration on the use of generative AI

During the development and preparation of this paper, the author used OpenAI's ChatGPT and Anthropic's Claude as exploratory mathematical tools. The central analytical ingredients—notably the use of the Duhamel equation and the Chebyshev monotonicity argument—were introduced by the author in his interactions with the systems. ChatGPT assisted in combining and developing these author-supplied ideas, carrying out intermediate calculations, testing the resulting implications, and identifying the construction that led to the reverse-ordering result presented in the paper. Claude was used to verify the work. Thus, although the underlying proof strategy originated with the author, the result emerged from AI-assisted exploration. The author subsequently reconstructed and independently verified every mathematical argument included in this manuscript. All mathematical judgments and conclusions are those of the author, who takes full responsibility for the accuracy and originality of the work.

References

- [1] Beatrice Acciaio and Julien Guyon. Inversion of convex ordering: local volatility does not maximize the price of VIX futures. *SIAM Journal on Financial Mathematics*, 11:SC1–SC13, 2020.
- [2] Peter Austing. *Smile Pricing Explained*. Financial Engineering Explained. Palgrave Macmillan, 2014.
- [3] Mathias Beiglböck, Peter Friz, and Stephan Sturm. Is the minimum value of an option on variance generated by local volatility? *SIAM Journal on Financial Mathematics*, 2:213–220, 2011.
- [4] Mark Broadie and Özgür Kaya. Exact simulation of stochastic volatility and other affine jump diffusion processes. *Operations Research*, 54(2):217–231, 2006.
- [5] Oliver Brockhaus and Douglas Long. Volatility swaps made simple. *Risk*, pages 92–95, January 2000. [Risk.net article](#).
- [6] Gerard Brunick and Steven Shreve. Mimicking an Itô process by a solution of a stochastic differential equation. *Annals of Applied Probability*, 23:1584–1628, 2013. [doi:10.1214/12-AAP881](#).
- [7] A. M. G. Cox and Jiajie Wang. Optimal robust bounds for variance options. arXiv:1308.4363, 2013.
- [8] A. M. G. Cox and Jiajie Wang. Root’s barrier: construction, optimality and applications to variance options. *Annals of Applied Probability*, 23:859–894, 2013.
- [9] Sebastian del Baño Rollin, Albert Ferreira-Castilla, and Frederic Utzet. On the density of log-spot in the Heston volatility model. *Stochastic Processes and their Applications*, 120(10):2037–2063, 2010.
- [10] Nicole El Karoui, Monique Jeanblanc-Picqué, and Steven E. Shreve. Robustness of the Black and Scholes formula. *Mathematical Finance*, 8:93–126, 1998. [doi:10.1111/1467-9965.00047](#).
- [11] Stewart N. Ethier and Thomas G. Kurtz. *Markov Processes: Characterization and Convergence*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, New York, 1986. [doi:10.1002/9780470316658](#).
- [12] Jim Gatheral. Valuation of volatility derivatives. Presentation at Global Derivatives & Risk Management 2005, Paris, May 2005.
- [13] Jim Gatheral. *The Volatility Surface: A Practitioner’s Guide*. Wiley, 2006.
- [14] Julien Guyon. Inversion of convex ordering in the VIX market. *Quantitative Finance*, 20:1597–1623, 2020.
- [15] David P. Herzog and Jonathan C. Mattingly. A practical criterion for positivity of transition densities. *Nonlinearity*, 28(8):2823–2845, 2015.
- [16] Steven L. Heston. A closed-form solution for options with stochastic volatility with applications to bond and currency options. *Review of Financial Studies*, 6:327–343, 1993. [doi:10.1093/rfs/6.2.327](#).
- [17] Steven L. Heston and Saikat Nandi. A closed-form GARCH option valuation model. *The Review of Financial Studies*, 13(3):585–625, 2000. [doi:10.1093/rfs/13.3.585](#).

- [18] Hiroshi Kunita. *Stochastic Flows and Stochastic Differential Equations*, volume 24 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.
- [19] Thomas G. Kurtz and Richard H. Stockbridge. Existence of Markov controls and characterization of optimal Markov controls. *SIAM Journal on Control and Optimization*, 36(2):609–653, 1998. doi:[10.1137/S0363012995295516](https://doi.org/10.1137/S0363012995295516);
- [20] Damien Lamberton and Bernard Lapeyre. *Introduction to Stochastic Calculus Applied to Finance*. Chapman & Hall/CRC, Boca Raton, 2nd edition, 2008.
- [21] Vladimir Lucic. On singularities in the Heston model. In Peter K. Friz, Jim Gatheral, Archil Gulisashvili, Antoine Jacquier, and Josef Teichmann, editors, *Large Deviations and Asymptotic Methods in Finance*, volume 110 of *Springer Proceedings in Mathematics & Statistics*, pages 439–448. Springer, Cham, 2015.
- [22] Frank W. J. Olver. *Asymptotics and Special Functions*. A K Peters, Wellesley, MA, 1997.
- [23] Daniel W. Stroock. *Probability Theory: An Analytic View*. Cambridge University Press, Cambridge, 3 edition, 2024. doi:[10.1017/9781009549035](https://doi.org/10.1017/9781009549035).
- [24] Toshio Yamada and Shinzo Watanabe. On the uniqueness of solutions of stochastic differential equations. *Journal of Mathematics of Kyoto University*, 11(1):155–167, 1971.