

Finite-Particle Convergence Rates for Conservative and Non-Conservative Drifting Models

Krishnakumar Balasubramanian
 Department of Statistics, University of California, Davis
 kbala@ucdavis.edu

May 22, 2026

Abstract

We propose and analyze a conservative drifting method for one-step generative modeling. The method replaces the original displacement-based drifting velocity by a kernel density estimator (KDE)-gradient velocity, namely the difference of the kernel-smoothed data score and the kernel-smoothed model score. This velocity is a gradient field, addressing the non-conservatism issue identified for general displacement-based drifting fields. We prove continuous-time finite-particle convergence bounds for the conservative method on $\mathbb{R}^d$: a joint-entropy identity yields bounds for the empirical Stein drift, the smoothed Fisher discrepancy of the KDE, and the squared center velocity. The main finite-particle correction is a reciprocal-KDE self-interaction term, and we give deterministic and high-probability local-occupancy conditions under which this term is controlled. We keep the quadrature constants explicit and track their possible bandwidth dependence: the root residual-velocity rate $N^{-1/(d+4)}$ holds under an additional h -uniform quadrature regularity condition, while a more general growth condition yields the optimized root rate $N^{-(2-\beta)/(2(d+4-\beta))}$, where $0 \leq \beta < 2$. We also analyze the non-conservative drifting method with Laplace kernel, corresponding to the original displacement-based velocity proposed in [Deng et al. \(2026\)](#). For this method, a sharp companion kernel decomposes the velocity into a positive scalar preconditioning of a sharp-score mismatch plus a Laplace scale-mismatch residual, producing an analogous finite-particle rate with an unavoidable residual term. Finally, we explain how the continuous-time residual-velocity bounds translate into one-step generation guarantees through the explicit drift size η .

1 Introduction

Drifting models ([Deng et al., 2026](#)) provide a direct approach to one-step generative modeling by moving the model distribution during training rather than applying a long iterative sampler at inference time. Let ν denote the data distribution on $\mathbb{R}^d$ and let μ denote the current model distribution. For a bandwidth- h kernel K_h , define the local kernel mean-shift vector of a probability measure α by

$$m_{\alpha,h}(z) := \frac{\int_{\mathbb{R}^d} (y - z) K_h(z - y) \alpha(dy)}{\int_{\mathbb{R}^d} K_h(z - y) \alpha(dy)}.$$

The displacement-based drifting velocity from μ toward ν is

$$u_{\nu,\mu,h}^{\text{disp}}(z) := m_{\nu,h}(z) - m_{\mu,h}(z).$$

The data term attracts a point z toward regions locally represented by the data, while the model term repels z from regions already represented by the model. In practical training, the drifted

target is treated as a stop-gradient target: one forms $z + \eta u_{\nu,\mu,h}^{\text{disp}}(z)$ and trains the generator to regress to this frozen target. In the idealized exact-regression limit, this produces the frozen-field Euler update $z^+ = z + \eta u_{\nu,\mu,h}^{\text{disp}}(z)$, and the continuous-time particle systems studied below are obtained by sending the step size to zero; see Section A for additional explanation.

The motivation for the first method in this paper is the fact that displacement-based drifting fields, as proposed in Deng et al. (2026), are generally not conservative. In particular, Franz et al. (2026) show that the position-dependent normalization in the original displacement field generally destroys the gradient-field structure, with the Gaussian kernel being a special case in which the field remains conservative. We therefore propose¹ the conservative drifting method, which replaces the displacement velocity by an explicit KDE-gradient or score velocity. For any probability measure α , define its kernel-smoothed density and score by

$$\rho_{\alpha,h}(z) := \int_{\mathbb{R}^d} K_h(z-y) \alpha(dy), \quad s_{\alpha,h}(z) := \nabla \log \rho_{\alpha,h}(z).$$

The conservative velocity from μ toward ν is

$$b_{\nu,\mu,h}(z) := s_{\nu,h}(z) - s_{\mu,h}(z) = \nabla \log \rho_{\nu,h}(z) - \nabla \log \rho_{\mu,h}(z).$$

It is called conservative because

$$b_{\nu,\mu,h}(z) = \nabla \{\log \rho_{\nu,h}(z) - \log \rho_{\mu,h}(z)\}$$

is a gradient field. Throughout the paper, b denotes a conservative velocity, whereas u denotes a non-conservative displacement velocity.

For Gaussian kernels, the conservative velocity and the displacement-based velocity coincide up to a deterministic scale factor. Indeed, if

$$K_h(u) = (2\pi h^2)^{-d/2} \exp\left(-\frac{\|u\|^2}{2h^2}\right),$$

then

$$\nabla_z K_h(z-y) = \frac{y-z}{h^2} K_h(z-y).$$

Therefore

$$s_{\alpha,h}(z) = \frac{\int \nabla_z K_h(z-y) \alpha(dy)}{\int K_h(z-y) \alpha(dy)} = \frac{1}{h^2} m_{\alpha,h}(z),$$

and hence

$$b_{\nu,\mu,h}(z) = \frac{1}{h^2} u_{\nu,\mu,h}^{\text{disp}}(z).$$

Thus the two Gaussian dynamics generate the same trajectories after rescaling time or step size. For non-Gaussian kernels this identity generally fails: the non-conservative displacement field averages the Euclidean vector $y-z$, while the conservative field uses the KDE-gradient direction.

We also remark that Franz et al. (2026) proposed a sharp-normalized field to restore conservativeness. While both our approach and that of Franz et al. (2026) lead to score-difference fields, they score different smoothed densities. Our conservative KDE-gradient field uses the ordinary KDE

¹**Concurrent and independent work.** During the final stages of completing this work, the author became aware of the work by Esteban-Casadevall et al. (2026) posted to arxiv on 11th May 2026. While their work also introduces the kernel-gradient drifting models, our contribution is a finite-particle entropy-rate analysis for the conservative dynamics and a parallel treatment of the non-conservative Laplace field.

$K_h * \alpha$, whereas the sharp-normalized field uses the companion KDE $K_h^\# * \alpha$, where $K_h^\#$ is chosen so that $\nabla K_h^\#(z - y) = (y - z)K_h(z - y)$. These two fields coincide up to scale for Gaussian kernels, because the Gaussian satisfies $\nabla K_h(z - y) \propto (y - z)K_h(z - y)$. For Laplace kernels they differ: the ordinary KDE-gradient field averages unit directions through ∇K_h , while the sharp-normalized field averages full displacement vectors and normalizes by the sharp KDE. Nevertheless, we leverage the idea of [Franz et al. \(2026\)](#) in our proof for analyzing the original drifting method proposed in [Deng et al. \(2026\)](#); see Section 6.

For the finite-particle conservative method, let $x = (x_1, \dots, x_N)$ and define

$$q_x(z) := \frac{1}{N} \sum_{j=1}^N K_h(z - x_j), \quad s_x(z) := \nabla \log q_x(z).$$

With the smoothed target density $\rho := \rho_{\nu, h}$ and $s_\rho := \nabla \log \rho$, the finite-particle conservative velocity is

$$b_x(z) := s_\rho(z) - s_x(z),$$

and the center-evaluation dynamics are

$$\dot{X}_i(t) = b_{X(t)}(X_i(t)), \quad i = 1, \dots, N.$$

The main residual-drift quantity is

$$V_N(x) := \frac{1}{N} \sum_{i=1}^N \|b_x(x_i)\|^2.$$

This is exactly the mean squared speed of the generated particle cloud under the conservative drifting flow. If one more explicit conservative drift step is applied,

$$\tilde{x}_i = x_i + \eta b_x(x_i), \quad \tilde{\mu}_x^N = \frac{1}{N} \sum_{i=1}^N \delta_{\tilde{x}_i},$$

then the identity coupling gives

$$W_2(\mu_x^N, \tilde{\mu}_x^N) \leq \eta \sqrt{V_N(x)}, \quad \mu_x^N := \frac{1}{N} \sum_{i=1}^N \delta_{x_i}.$$

Thus a bound on V_N says that another drift refinement would change the one-step generated distribution only slightly. With a quadrature condition, V_N also approximates the smoothed Fisher discrepancy $\int \|s_\rho - s_x\|^2 q_x$ and therefore controls a smoothed distribution-level score mismatch.

Our first contribution is a continuous-time finite-particle rate for the conservative method. By leveraging and extending the joint-entropy structure of the finite-particle Stein Variational Gradient Descent (SVGD) analysis in [Banerjee et al. \(2025\)](#), we prove that under regularity, reciprocal-KDE control, and point-evaluation quadrature smoothness,

$$\frac{1}{T} \int_0^T \mathbb{E}[V_N(X(t))] dt \leq \frac{H_N(0)}{NT} + \frac{\{-\Delta K(0)\}_+ \Lambda_T}{Nh^{d+2}} + C_T(h)h^2.$$

Here $H_N(0)$ is the initial joint entropy relative to the product smoothed target, Λ_T controls the reciprocal KDE term, and $C_T(h)$ is a quadrature constant that may depend on the bandwidth.

This bandwidth dependence is important. If $C_N(h) = O(1)$ and $T = N$, optimizing the last two terms gives the root residual-velocity rate $N^{-1/(d+4)}$. More generally, if $C_N(h) = O(h^{-\beta})$ for some $0 \leq \beta < 2$, the optimized root rate becomes

$$N^{-(2-\beta)/(2(d+4-\beta))}.$$

If $\beta \geq 2$, the present quadrature argument does not yield a vanishing residual by shrinking h .

Our second contribution is a corresponding result for the non-conservative drifting method with Laplace kernel, namely the original displacement field specialized to

$$K_h(u) = c_d h^{-d} \exp\left(-\frac{\|u\|}{h}\right).$$

This velocity is not generally a gradient field. To analyze it, we leverage the sharp companion kernel idea proposed in [Franz et al. \(2026\)](#) and define L_h satisfying

$$\nabla_z L_h(z - y) = (y - z)K_h(z - y).$$

This yields a decomposition of the full non-conservative Laplace velocity of the form

$$u_x(z) = a_{x,h}(z)b_x^\#(z) + e_x(z),$$

where $b_x^\#$ is a sharp-score mismatch, $a_{x,h}$ is a positive local scale factor, and e_x is a scale-mismatch residual measuring the difference between the data and model local Laplace-weighted radii. For the self-masked, leave-one-out Laplace dynamics, we prove a finite-particle rate

$$\frac{1}{N} \int_0^N \mathbb{E}[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{\kappa_0}{\gamma_h N} + \frac{\beta_h}{\gamma_h} \Delta_h^2 + \frac{\varepsilon_{S,h,N}}{\gamma_h} + \varepsilon_{V,h,N}.$$

The new term Δ_h is unavoidable for the non-conservative Laplace method: it vanishes only when the local Laplace scales of the data and model are aligned. The leave-one-out form removes the reciprocal self-interaction term from the entropy identity, but introduces leave-one-out approximation errors controlled by denominator and occupancy conditions.

1.1 Related Work

Drifting models were introduced by [Deng et al. \(2026\)](#) as a training-time transport paradigm for one-step generation: generated samples are repeatedly moved by a data-attractive and model-repulsive drift field, and the generator is trained to reproduce the drifted samples through a stop-gradient target. This viewpoint is also related to score-difference flows for implicit generative modeling, where the velocity is written as the difference between target and source scores ([Weber, 2023](#)). Subsequent work has connected drifting to score matching and Wasserstein gradient flows of smoothed divergences (in the infinite-particle setting), especially in the Gaussian-kernel case where the displacement field can be written as a difference of KDE log-density gradients ([Cao et al., 2026](#); [Lai et al., 2026](#); [Turan and Ovsjanikov, 2026](#); [Gretton et al., 2026](#)).

A complementary line of work emphasizes the non-conservative nature of the original displacement-based field. [Franz et al. \(2026\)](#) show that the position-dependent normalization in the original field generally destroys conservatism, with the Gaussian kernel being a special case where the field remains a gradient field; they also introduce sharp normalizations that restore conservatism for broader radial kernels. [Lee and Chun \(2026\)](#) studies identifiability and stability for companion-elliptic kernel families, including the Laplace kernel, showing that vanishing drift can identify the

target under appropriate kernel structure while also highlighting possible instability modes. These works motivate our distinction between the conservative drifting method, based on KDE-score gradients, and the non-conservative drifting method by [Deng et al. \(2026\)](#), based on the original displacement velocity. For the latter, our analysis shows that finite-particle rates require a scale-alignment residual measuring the failure of the Laplace displacement field to be an exact score-difference field.

Recent drifting-related works study complementary algorithmic and variational perspectives. [Dumont et al. \(2026\)](#) study learning Monge maps through constrained drifting models: they lift a divergence, such as relative entropy, to the space of transport maps, constrain the dynamics to the set of optimal transport maps, and prove long-time existence and convergence toward the OT map under convexity assumptions. [He et al. \(2026a\)](#) introduce Sinkhorn-Drifting Generative Models, showing that the attraction–repulsion structure of drifting can be viewed as a one-sided surrogate for a Sinkhorn-divergence gradient flow, while two-sided Sinkhorn scaling gives improved identifiability and stability. [Zhang et al. \(2026\)](#) propose Lookahead Drifting Models, which compute several sequential drift terms within each training iteration and regress toward a weighted sum of these terms, with the goal of incorporating higher-order drift information while preserving one-step inference.

More broadly, modern generative modeling is often organized around learned transport dynamics. Diffusion and score-based models learn reverse-time denoising dynamics and generate samples by numerically solving an SDE or probability-flow ODE ([Ho et al., 2020](#); [Song et al., 2021](#)), while flow matching and rectified flow learn time-dependent vector fields that transport a base distribution to the data distribution along prescribed probability paths ([Albergo and Vanden-Eijnden, 2023](#); [Lipman et al., 2023](#); [Liu et al., 2023](#)). Flow-map and distillation-based approaches instead try to learn longer-time solution operators, or compress many solver steps into one or a few neural evaluations, as in flow-map matching, consistency models, and progressive distillation ([Boffi et al., 2025](#); [Song et al., 2023](#); [Salimans and Ho, 2022](#)). From this viewpoint, drifting models can be interpreted as moving part of the inference-time transport computation into training: the long-short flow-map perspective of [Li and Zhu \(2026\)](#) decomposes a global transport into a long-horizon map followed by a short terminal correction, and recovers the drifting field in the limit of a vanishing terminal interval.

The proposed conservative velocity approach is also closely related to deterministic interacting-particle methods based on Stein discrepancies. Stein variational gradient descent (SVGD) was introduced by [Liu and Wang \(2016\)](#) as a deterministic particle method for approximate Bayesian inference, and [Liu \(2017\)](#) interpreted its population limit as a gradient flow of KL divergence under a Stein-induced geometry. Amortized SVGD ([Feng et al., 2017](#)) trains a sampler network to reproduce SVGD particle updates, which is conceptually close to the stop-gradient regression view of drifting. At the level of finite-particle analysis, [Banerjee et al. \(2025\)](#) obtain improved SVGD rates by differentiating the joint entropy of the particle law relative to the product target, improving upon earlier results by [Shi and Mackey \(2023\)](#). Extension to a regularized setting was studied by [He et al. \(2026b\)](#). We leverage and extend this high-level entropy-dissipation strategy in our proofs, but the controlled quantities and error terms are specific to drifting: for conservative drifting, the controlled quantity is the mean squared conservative residual with a reciprocal-KDE self-interaction and an h -dependent quadrature error; for the non-conservative Laplace method, the rate additionally contains the Laplace scale-mismatch residual.

2 Preliminaries

2.1 Set-up for conservative drifting

Let ρ be a positive probability density on $\mathbb{R}^d$. For the conservative drifting method proposed in this paper, ρ should be read as the kernel-smoothed data density $\rho_{\nu,h} = K_h * \nu$. Write

$$s_\rho(z) := \nabla \log \rho(z).$$

Let $K : \mathbb{R}^d \rightarrow (0, \infty)$ be a normalized kernel and, for a bandwidth $h > 0$, define

$$K_h(u) := h^{-d} K(u/h).$$

For a particle configuration

$$x = (x_1, \dots, x_N) \in (\mathbb{R}^d)^N,$$

define the kernel density estimate

$$q_x(z) := \frac{1}{N} \sum_{j=1}^N K_h(z - x_j), \quad s_x(z) := \nabla \log q_x(z),$$

and the conservative drift field

$$b_x(z) := s_\rho(z) - s_x(z).$$

The center-evaluation finite-particle ODE for the conservative drifting method is

$$\dot{X}_i(t) = b_i(X(t)), \quad b_i(x) := b_x(x_i), \quad i = 1, \dots, N. \quad (1)$$

This is a deterministic interacting particle system: each particle follows an ordinary differential equation, but its velocity depends on the full configuration through the KDE and its score. In this sense, the dynamics are structurally similar to SVGD (Liu and Wang, 2016; Liu, 2017), where particles also evolve deterministically through an empirical-measure-dependent velocity field; the key difference is that the conservative drifting velocity is the KDE-score residual $s_\rho - s_x$, rather than the kernelized Stein velocity used in SVGD.

Let p_t^N denote the joint density of $X(t) = (X_1(t), \dots, X_N(t))$. We measure the joint law against the product target density

$$\rho^{\otimes N}(x) := \prod_{i=1}^N \rho(x_i)$$

through the joint relative entropy

$$H_N(t) := \text{KL}(p_t^N \| \rho^{\otimes N}) = \int_{(\mathbb{R}^d)^N} p_t^N(x) \log \frac{p_t^N(x)}{\rho^{\otimes N}(x)} dx. \quad (2)$$

For a vector field $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$, define the Stein divergence (Barbour, 1988; Gorham and Mackey, 2015) associated with ρ by

$$\mathcal{A}_\rho f(z) := \nabla \cdot f(z) + s_\rho(z) \cdot f(z). \quad (3)$$

For a configuration x , define the empirical Stein drift

$$S_N(x) := \frac{1}{N} \sum_{i=1}^N \mathcal{A}_\rho b_x(x_i), \quad (4)$$

the KDE-smoothed Fisher discrepancy

$$\mathbf{I}_N(x) := \int_{\mathbb{R}^d} \|b_x(z)\|^2 q_x(z) \, dz, \quad (5)$$

and the squared center velocity

$$\mathbf{V}_N(x) := \frac{1}{N} \sum_{i=1}^N \|b_x(x_i)\|^2. \quad (6)$$

Finally set

$$\mathbf{R}_N(x) := \frac{1}{N} \sum_{i=1}^N \frac{1}{q_x(x_i)}. \quad (7)$$

2.2 Assumptions

The following assumptions are used in the entropy calculation and in the conversion from empirical Stein drift to Fisher-type discrepancies.

Assumption 2.1 (Kernel). *The base kernel K is positive, even, normalized, and C^3 :*

$$K(u) > 0, \quad K(u) = K(-u), \quad \int_{\mathbb{R}^d} K(u) \, du = 1.$$

Moreover

$$m_2(K) := \int_{\mathbb{R}^d} \|u\|^2 K(u) \, du < \infty,$$

and $\Delta K(0)$ is finite. Since K is even, $\nabla K(0) = 0$.

Assumption 2.2 (Target and regular flow). *The density ρ is positive and sufficiently smooth so that s_ρ is C^2 . The ODE (1) has a unique global solution for the initial law under consideration. The joint density p_t^N is differentiable in t and solves the Liouville equation*

$$\partial_t p_t^N(x) + \sum_{i=1}^N \nabla_{x_i} \cdot \{p_t^N(x) b_i(x)\} = 0.$$

All integrals appearing below are finite, and the boundary terms in the integrations by parts on $(\mathbb{R}^d)^N$ vanish. A sufficient, but not necessary, condition is rapid decay of $p_t^N(x) b_i(x)$ and of the products obtained by multiplying $p_t^N(x) b_i(x)$ by $\log(p_t^N(x)/\rho^{\otimes N}(x))$. See [Banerjee et al. \(2025\)](#) for related conditions for SVGD.

Assumption 2.3 (Reciprocal KDE control). *For the time horizon $T > 0$ under consideration, there is a finite constant Λ_T such that*

$$\sup_{0 \leq t \leq T} \mathbb{E}_t[\mathbf{R}_N(X(t))] \leq \Lambda_T, \quad (8)$$

where $\mathbb{E}_t$ denotes expectation with respect to p_t^N and $\mathbf{R}_N$ is defined in (7).

Remark 2.4 (Conditional nature of reciprocal-KDE control). *The reciprocal-KDE assumption is a stability assumption, not a consequence of the entropy identity. The local-occupancy results in Section 5 give deterministic or high-probability sufficient conditions under which*

$$q_{X(t)}(X_i(t)) \geq \lambda > 0 \quad \text{for all } i$$

at a fixed time, and under a propagated occupancy condition. However, propagation arguments that use a flow Lipschitz constant are conditional: the Lipschitz constant of the conservative vector field itself depends on reciprocal KDE quantities such as $1/q_x$, and sometimes on higher inverse powers of q_x . Thus there is a potential bootstrap structure.

A precise way to read the assumption is through the stopping time

$$\tau_\lambda := \inf \left\{ t \geq 0 : \min_{1 \leq i \leq N} q_{X(t)}(X_i(t)) < \lambda \right\}.$$

All conservative finite-particle bounds hold on the stopped interval $[0, T \wedge \tau_\lambda]$, with constants depending on λ . If one can prove separately that $\tau_\lambda > T$, for example by a deterministic occupancy argument or by a high-probability propagation argument that closes a bootstrap, then the stopped estimates become estimates on the full interval $[0, T]$. We do not claim that reciprocal-KDE control follows automatically from the dynamics; it is an independent stability condition required to prevent denominator singularities.

Assumption 2.5 (Quadrature smoothness). *For the time horizon $T > 0$, there are finite constants $B_{A,T}(h)$ and $B_{V,T}(h)$ such that, for every configuration $X(t)$ visited by the flow for $0 \leq t \leq T$,*

$$\sup_{z \in \mathbb{R}^d} \left\| D^2 \{ \mathcal{A}_\rho b_{X(t)} \} (z) \right\|_{\text{op}} \leq B_{A,T}(h), \quad (9)$$

$$\sup_{z \in \mathbb{R}^d} \left\| D^2 \{ \|b_{X(t)}\|^2 \} (z) \right\|_{\text{op}} \leq B_{V,T}(h). \quad (10)$$

Here D^2 denotes the Hessian, and $\|\cdot\|_{\text{op}}$ is the operator norm.

Remark 2.6 (Bandwidth dependence of the quadrature constants). *The constants $B_{A,T}(h)$ and $B_{V,T}(h)$ may depend on N, h, T, ρ, K , and on the region of state space visited by the dynamics. This dependence is not a lower-order issue. Writing*

$$r_x(z) := \log \rho(z) - \log q_x(z), \quad b_x(z) = \nabla r_x(z),$$

we have

$$\mathcal{A}_\rho b_x = \Delta r_x + \nabla \log \rho \cdot \nabla r_x,$$

and

$$D^2 \|b_x\|^2 = 2(Db_x)^\top (Db_x) + 2 \sum_{\ell=1}^d b_{x,\ell} D^2 b_{x,\ell}.$$

Thus the quadrature constants involve log-derivatives of ρ and q_x up to fourth order. If these log-derivatives vary on the kernel scale, for example if

$$\sup_z \left\| D^j (\log \rho - \log q_x)(z) \right\| = O(h^{-j}), \quad j = 1, 2, 3, 4,$$

then one typically has $B_{A,T}(h) + B_{V,T}(h) = O(h^{-4})$. In that regime the quadrature contribution

$$\{B_{A,T}(h) + B_{V,T}(h)\} m_2(K_h) = O(h^{-2})$$

scales like h^{-2} , not h^2 , and shrinking the bandwidth worsens this term. Consequently, the rates below are non-asymptotic inequalities with the quadrature constants displayed. Any asymptotic bandwidth optimization must explicitly account for the h -dependence of $B_{A,T}(h)$ and $B_{V,T}(h)$.

Remark 2.7 (Examples of quadrature-growth regimes). *The exponent β in*

$$B_{A,N}(h) + B_{V,N}(h) = O(h^{-\beta})$$

should be interpreted as a regularity exponent for the KDE-score field along the particle trajectory, not as a property of the kernel alone.

First, $\beta = 0$ is realized in an h -uniform log-smooth regime. For example, take a Gaussian kernel or a C^∞ compactly supported kernel, and suppose that the target density and the model KDE satisfy

$$\sup_{0 \leq t \leq N} \sup_z \max_{1 \leq j \leq 4} \left\{ \|D^j \log \rho(z)\|_{\text{op}}, \|D^j \log q_{X(t)}(z)\|_{\text{op}} \right\} \leq C_{\log},$$

with $C_{\log}$ independent of h . Then

$$B_{A,N}(h) + B_{V,N}(h) = O(1),$$

so $\beta = 0$. This situation can occur when the particles are sufficiently dense and regular so that the finite KDE behaves like the convolution of the kernel with a fixed smooth density bounded away from zero, rather than like a collection of isolated bandwidth- h spikes.

Second, exponents $0 < \beta < 2$ correspond to intermediate-scale regularity. Suppose the log-score residual

$$r_x(z) := \log \rho(z) - \log q_x(z)$$

varies on a length scale $\ell_h = h^\alpha$, where $0 < \alpha < 1/2$. More concretely, assume

$$\sup_z \|D^j r_x(z)\|_{\text{op}} = O(\ell_h^{-j}) = O(h^{-\alpha j}), \quad j = 1, 2, 3, 4.$$

Since $b_x = \nabla r_x$, the quantities $D^2\{\mathcal{A}_\rho b_x\}$ and $D^2\{\|b_x\|^2\}$ involve derivatives of r_x up to fourth order. Hence

$$B_{A,N}(h) + B_{V,N}(h) = O(h^{-4\alpha}).$$

Thus

$$\beta = 4\alpha.$$

Because $0 < \alpha < 1/2$, this gives examples with $0 < \beta < 2$. The same smooth kernels, such as Gaussian kernels or smooth compactly supported kernels, can realize either regime. What changes is the regularity of the log-density ratio $\log \rho - \log q_x$ along the particle trajectory. If this ratio has only macroscopic variation, then $\beta = 0$. If it has features at an intermediate scale $\ell_h = h^\alpha$ with $0 < \alpha < 1/2$, then $\beta = 4\alpha \in (0, 2)$. If it varies at the kernel scale $\ell_h = h$, then $\alpha = 1$, hence $\beta = 4$, and the optimized vanishing-rate statement no longer applies.

Finally, we emphasize that our conservative drifting analysis assumes that the smoothing kernel is sufficiently smooth, for example $K \in C^3$ or stronger when the quadrature constants involve higher derivatives. This assumption excludes the exact Laplace kernel $K_h(u) = c_d h^{-d} \exp(-\|u\|/h)$, which is not differentiable at the origin. This is why the exact Laplace kernel is treated separately in Section 6 through the non-conservative displacement field and the sharp-companion-kernel decomposition. A smooth regularization of the Laplace kernel could be included in the conservative analysis, but the resulting estimates would have constants depending on the regularization scale.

Another useful feature of the forthcoming results is that it imposes only the aforementioned local smoothness and positivity assumptions on the smoothed target density needed to define its score and justify the entropy calculation; in particular, we do not assume curvature conditions such as the target is log-concave, strongly log-concave, or satisfies a Log-Sobolev, Poincaré, or dissipativity condition. Our bounds that control the time-averaged residual drift, provide a global convergence in KL or Wasserstein distance under such additional curvature conditions.

3 Entropy identity for conservative drifting

The next theorem is the basic finite-particle identity. The self-interaction term is the only term not present in the corresponding population smoothed-KL calculation.

Theorem 3.1 (Joint-entropy identity). *Let Assumptions 2.1 and 2.2 hold. Then, for every t for which the quantities are finite,*

$$\frac{d}{dt}H_N(t) = -N\mathbb{E}_t[\mathbf{S}_N(X(t))] - \Delta K_h(0) \mathbb{E}_t[\mathbf{R}_N(X(t))]. \quad (11)$$

Equivalently, with

$$a_h := \{-\Delta K_h(0)\}_+,$$

one has the upper bound

$$\frac{d}{dt}H_N(t) \leq -N\mathbb{E}_t[\mathbf{S}_N(X(t))] + a_h \mathbb{E}_t[\mathbf{R}_N(X(t))]. \quad (12)$$

Proof. Let

$$r_N(x) := \rho^{\otimes N}(x).$$

By definition,

$$H_N(t) = \int p_t^N(x) \log \frac{p_t^N(x)}{r_N(x)} dx.$$

Differentiating under the integral gives

$$\frac{d}{dt}H_N(t) = \int \partial_t p_t^N(x) \left\{ \log \frac{p_t^N(x)}{r_N(x)} + 1 \right\} dx.$$

Since p_t^N is a probability density,

$$\int \partial_t p_t^N(x) dx = \frac{d}{dt} \int p_t^N(x) dx = 0.$$

Thus

$$\frac{d}{dt}H_N(t) = \int \partial_t p_t^N(x) \log \frac{p_t^N(x)}{r_N(x)} dx.$$

Using the Liouville equation,

$$\frac{d}{dt}H_N(t) = - \sum_{i=1}^N \int \nabla_{x_i} \cdot \{p_t^N(x) b_i(x)\} \log \frac{p_t^N(x)}{r_N(x)} dx.$$

Integrating by parts in x_i ,

$$\frac{d}{dt}H_N(t) = \sum_{i=1}^N \int p_t^N(x) b_i(x) \cdot \nabla_{x_i} \log \frac{p_t^N(x)}{r_N(x)} dx.$$

Now

$$\nabla_{x_i} \log \frac{p_t^N(x)}{r_N(x)} = \nabla_{x_i} \log p_t^N(x) - \nabla_{x_i} \log r_N(x).$$

Since $r_N(x) = \prod_{j=1}^N \rho(x_j)$,

$$\nabla_{x_i} \log r_N(x) = s_\rho(x_i).$$

Therefore

$$\frac{d}{dt}H_N(t) = \sum_{i=1}^N \int p_t^N b_i \cdot \nabla_{x_i} \log p_t^N dx - \sum_{i=1}^N \int p_t^N b_i \cdot s_\rho(x_i) dx.$$

For the first integral,

$$p_t^N \nabla_{x_i} \log p_t^N = \nabla_{x_i} p_t^N.$$

Hence

$$\int p_t^N b_i \cdot \nabla_{x_i} \log p_t^N dx = \int b_i \cdot \nabla_{x_i} p_t^N dx = - \int p_t^N \nabla_{x_i} \cdot b_i dx.$$

Consequently,

$$\frac{d}{dt}H_N(t) = -\mathbb{E}_t \left[\sum_{i=1}^N \{ \nabla_{x_i} \cdot b_i(X(t)) + s_\rho(X_i(t)) \cdot b_i(X(t)) \} \right]. \quad (13)$$

It remains to compute $\nabla_{x_i} \cdot b_i(x)$. Since

$$b_i(x) = s_\rho(x_i) - s_x(x_i),$$

we have

$$\nabla_{x_i} \cdot b_i(x) = \nabla \cdot s_\rho(x_i) - \nabla_{x_i} \cdot s_x(x_i).$$

The derivative of $s_x(x_i)$ with respect to x_i contains two pieces: the derivative of the evaluation point and the derivative of the i th KDE center. Write

$$s_x^m(z) = \frac{\partial_m q_x(z)}{q_x(z)}, \quad m = 1, \dots, d.$$

For fixed z ,

$$\partial_{x_i^\ell} q_x(z) = -\frac{1}{N} \partial_\ell K_h(z - x_i),$$

and

$$\partial_{x_i^\ell} \partial_m q_x(z) = -\frac{1}{N} \partial_{\ell m}^2 K_h(z - x_i).$$

Therefore

$$\partial_{x_i^\ell} s_x^m(z) = -\frac{1}{N} \frac{\partial_{\ell m}^2 K_h(z - x_i)}{q_x(z)} + \frac{1}{N} \frac{\partial_m q_x(z) \partial_\ell K_h(z - x_i)}{q_x(z)^2}.$$

Set $m = \ell$ and sum over ℓ . Since K_h is even, $\nabla K_h(0) = 0$, and hence at $z = x_i$,

$$\sum_{\ell=1}^d \partial_{x_i^\ell} s_x^\ell(z) \Big|_{z=x_i} = -\frac{1}{N} \frac{\Delta K_h(0)}{q_x(x_i)}.$$

By the chain rule,

$$\nabla_{x_i} \cdot s_x(x_i) = \nabla_z \cdot s_x(z) \Big|_{z=x_i} - \frac{1}{N} \frac{\Delta K_h(0)}{q_x(x_i)}.$$

Thus

$$\begin{aligned} \nabla_{x_i} \cdot b_i(x) &= \nabla \cdot s_\rho(x_i) - \nabla_z \cdot s_x(z) \Big|_{z=x_i} + \frac{1}{N} \frac{\Delta K_h(0)}{q_x(x_i)} \\ &= \nabla_z \cdot b_x(z) \Big|_{z=x_i} + \frac{1}{N} \frac{\Delta K_h(0)}{q_x(x_i)}. \end{aligned}$$

Since $b_i(x) = b_x(x_i)$,

$$\nabla_{x_i} \cdot b_i(x) + s_\rho(x_i) \cdot b_i(x) = \mathcal{A}_\rho b_x(x_i) + \frac{1}{N} \frac{\Delta K_h(0)}{q_x(x_i)}.$$

Substituting this identity into (13) gives

$$\begin{aligned} \frac{d}{dt} H_N(t) &= -\mathbb{E}_t \left[\sum_{i=1}^N \mathcal{A}_\rho b_{X(t)}(X_i(t)) + \frac{\Delta K_h(0)}{N} \sum_{i=1}^N \frac{1}{q_{X(t)}(X_i(t))} \right] \\ &= -N \mathbb{E}_t[\mathcal{S}_N(X(t))] - \Delta K_h(0) \mathbb{E}_t[\mathcal{R}_N(X(t))]. \end{aligned}$$

This proves (11). Since

$$-\Delta K_h(0) \leq \{-\Delta K_h(0)\}_+ = a_h,$$

(12) follows. $\square$

Remark 3.2 (Interpretation of the self-interaction term). *The first term on the right-hand side of (11) is the desired entropy dissipation. The second term is finite-particle specific: it appears because the velocity at x_i depends on the same particle through the KDE denominator. Its sign and size are controlled by $-\Delta K_h(0)$ and by the reciprocal quantity $\mathcal{R}_N$. This is why the subsequent rate theorem requires reciprocal-KDE control. If particles are locally well populated at bandwidth h , then $q_x(x_i)$ is bounded below and the self-interaction correction remains of order $1/N$ after the entropy normalization.*

For Gaussian kernels, the self-interaction correction has a definite positive sign. Indeed, for $K(u) = (2\pi)^{-d/2} \exp(-\|u\|^2/2)$, we have, for each coordinate j , $\partial_{jj} K(0) = -K(0) = -(2\pi)^{-d/2}$. Therefore

$$\Delta K(0) = \sum_{j=1}^d \partial_{jj} K(0) = -d(2\pi)^{-d/2} < 0,$$

and hence $a_1 := \{-\Delta K(0)\}_+ = d(2\pi)^{-d/2} > 0$. For the bandwidth-scaled kernel $K_h(u) = h^{-d} K(u/h)$,

$$-\Delta K_h(0) = h^{-d-2} a_1.$$

Thus the reciprocal-KDE term in the conservative finite-particle bound is a genuine positive self-interaction correction for Gaussian kernels.

4 Continuous-time convergence rates for conservative drifting

4.1 Rate for the empirical Stein drift

Theorem 4.1 (Entropy rate for the empirical Stein drift). *Let Assumptions 2.1, 2.2, and 2.3 hold. Then, for every $T > 0$,*

$$\frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{S}_N(X(t))] dt \leq \frac{H_N(0)}{NT} + \frac{a_h \Lambda_T}{N}, \quad a_h := \{-\Delta K_h(0)\}_+. \quad (14)$$

Proof. By Theorem 3.1,

$$H'_N(t) \leq -N\mathbb{E}_t[\mathbf{S}_N(X(t))] + a_h \mathbb{E}_t[\mathbf{R}_N(X(t))].$$

Using Assumption 2.3,

$$H'_N(t) \leq -N\mathbb{E}_t[\mathbf{S}_N(X(t))] + a_h \Lambda_T.$$

Rearrange:

$$N\mathbb{E}_t[\mathbf{S}_N(X(t))] \leq -H'_N(t) + a_h \Lambda_T.$$

Integrate over $t \in [0, T]$:

$$N \int_0^T \mathbb{E}_t[\mathbf{S}_N(X(t))] dt \leq H_N(0) - H_N(T) + a_h \Lambda_T T.$$

Since relative entropy is nonnegative,

$$H_N(0) - H_N(T) \leq H_N(0).$$

Therefore

$$N \int_0^T \mathbb{E}_t[\mathbf{S}_N(X(t))] dt \leq H_N(0) + a_h \Lambda_T T.$$

Dividing by NT proves (14). $\square$

Remark 4.2 (What the Stein-drift rate controls). *The quantity $\mathbf{S}_N$ is the empirical Stein drift associated with the vector field b_x . It is not written as a square, so (14) is not yet a residual-velocity bound. The role of the quadrature step below is to compare this empirical Stein drift with the KDE-averaged Fisher discrepancy, which is nonnegative and is directly related to the squared velocity of the particles.*

4.2 Quadrature conversion to Fisher and center velocity

The empirical Stein drift $\mathbf{S}_N$ is not manifestly nonnegative. The next lemma identifies the nonnegative population quantity that it approximates.

Lemma 4.3 (KDE-averaged Stein identity). *For every fixed configuration x for which the integrations by parts are justified,*

$$\int_{\mathbb{R}^d} \mathcal{A}_\rho b_x(z) q_x(z) dz = \mathbf{l}_N(x). \quad (15)$$

Proof. By definition of $\mathcal{A}_\rho$,

$$\int q_x \mathcal{A}_\rho b_x = \int q_x \nabla \cdot b_x + \int q_x s_\rho \cdot b_x.$$

Integrating the first term by parts,

$$\int q_x \nabla \cdot b_x = - \int \nabla q_x \cdot b_x.$$

Since $\nabla q_x = q_x s_x$,

$$- \int \nabla q_x \cdot b_x = - \int q_x s_x \cdot b_x.$$

Thus

$$\int q_x \mathcal{A}_\rho b_x = \int q_x (s_\rho - s_x) \cdot b_x.$$

Because $b_x = s_\rho - s_x$,

$$\int q_x \mathcal{A}_\rho b_x = \int q_x \|b_x\|^2 = \mathfrak{l}_N(x).$$

This proves the lemma. $\square$

Lemma 4.4 (Point-evaluation quadrature error). *Let K_h be even and have finite second moment*

$$m_2(K_h) := \int_{\mathbb{R}^d} \|u\|^2 K_h(u) \, du.$$

Under Assumption 2.5, for every configuration $X(t)$ visited by the flow,

$$|\mathfrak{S}_N(X(t)) - \mathfrak{l}_N(X(t))| \leq \frac{1}{2} B_{A,T}(h) m_2(K_h), \quad (16)$$

$$|\mathfrak{V}_N(X(t)) - \mathfrak{l}_N(X(t))| \leq \frac{1}{2} B_{V,T}(h) m_2(K_h). \quad (17)$$

Proof. We prove (16); the proof of (17) is identical with $\|b_x\|^2$ replacing $\mathcal{A}_\rho b_x$.

Fix x and define

$$\phi_x(z) := \mathcal{A}_\rho b_x(z).$$

By Lemma 4.3,

$$\mathfrak{l}_N(x) = \int q_x(z) \phi_x(z) \, dz.$$

Since

$$q_x(z) = \frac{1}{N} \sum_{i=1}^N K_h(z - x_i),$$

we have

$$\mathfrak{l}_N(x) = \frac{1}{N} \sum_{i=1}^N \int K_h(z - x_i) \phi_x(z) \, dz.$$

Meanwhile,

$$\mathfrak{S}_N(x) = \frac{1}{N} \sum_{i=1}^N \phi_x(x_i).$$

Therefore

$$\mathbf{S}_N(x) - \mathbf{l}_N(x) = \frac{1}{N} \sum_{i=1}^N \left\{ \phi_x(x_i) - \int K_h(z - x_i) \phi_x(z) \, dz \right\}.$$

Set $u = z - x_i$. Then

$$\int K_h(z - x_i) \phi_x(z) \, dz = \int K_h(u) \phi_x(x_i + u) \, du.$$

Taylor's formula gives

$$\phi_x(x_i + u) = \phi_x(x_i) + \nabla \phi_x(x_i) \cdot u + \int_0^1 (1-r) u^\top D^2 \phi_x(x_i + ru) u \, dr.$$

Integrating against $K_h(u) \, du$, the zeroth-order term gives $\phi_x(x_i)$ because $\int K_h = 1$. The first-order term vanishes because K_h is even, hence $\int u K_h(u) \, du = 0$. Thus

$$\begin{aligned} & \left| \int K_h(u) \phi_x(x_i + u) \, du - \phi_x(x_i) \right| \\ & \leq \int K_h(u) \int_0^1 (1-r) \|D^2 \phi_x(x_i + ru)\|_{\text{op}} \|u\|^2 \, dr \, du. \end{aligned}$$

Using Assumption 2.5,

$$\left| \int K_h(u) \phi_x(x_i + u) \, du - \phi_x(x_i) \right| \leq B_{A,T}(h) \int_0^1 (1-r) \, dr \int \|u\|^2 K_h(u) \, du.$$

Since $\int_0^1 (1-r) \, dr = 1/2$,

$$\left| \int K_h(u) \phi_x(x_i + u) \, du - \phi_x(x_i) \right| \leq \frac{1}{2} B_{A,T}(h) m_2(K_h).$$

Averaging over $i = 1, \dots, N$ proves (16). $\square$

Theorem 4.5 (Continuous-time rates for Fisher discrepancy and center velocity). *Let Assumptions 2.1, 2.2, 2.3, and 2.5 hold. Then, for every $T > 0$,*

$$\frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{l}_N(X(t))] \, dt \leq \frac{H_N(0)}{NT} + \frac{a_h \Lambda_T}{N} + \frac{1}{2} B_{A,T}(h) m_2(K_h), \quad (18)$$

$$\frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N(X(t))] \, dt \leq \frac{H_N(0)}{NT} + \frac{a_h \Lambda_T}{N} + \frac{1}{2} \{B_{A,T}(h) + B_{V,T}(h)\} m_2(K_h). \quad (19)$$

Proof. From Lemma 4.4,

$$\mathbf{l}_N(X(t)) \leq \mathbf{S}_N(X(t)) + \frac{1}{2} B_{A,T}(h) m_2(K_h).$$

Taking expectations, integrating over $[0, T]$, dividing by T , and applying Theorem 4.1 gives (18).

Similarly,

$$\mathbf{V}_N(X(t)) \leq \mathbf{l}_N(X(t)) + \frac{1}{2} B_{V,T}(h) m_2(K_h).$$

Combining this inequality with (18) gives (19). $\square$

Remark 4.6 (Implication for one-step generation). *Theorem 4.5 bounds the average squared speed of the generated particle cloud under the conservative drift. If a time τ is sampled uniformly from $[0, T]$, then*

$$\mathbb{E}[\mathbf{V}_N(X(\tau))] = \frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N(X(t))] dt.$$

Consequently, a small right-hand side in (19) implies that, at a typical training time, one more explicit drift step changes the empirical generated law by at most

$$\mathbb{E} \left[W_2^2 \left(\mu_{X(\tau)}^N, \frac{1}{N} \sum_{i=1}^N \delta_{X_i(\tau) + \eta b_{X(\tau)}(X_i(\tau))} \right) \right] \leq \eta^2 \frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N(X(t))] dt.$$

Thus the theorem is a convergence-to-small-residual-drift statement for one-step generation. With the quadrature comparison, it also controls the smoothed Fisher discrepancy between q_x and ρ .

Corollary 4.7 (Bandwidth form). *Let $K_h(u) = h^{-d}K(u/h)$ and let the assumptions of Theorem 4.5 hold. Then*

$$m_2(K_h) = h^2 m_2(K), \quad \Delta K_h(0) = h^{-d-2} \Delta K(0).$$

Consequently, with $a_1 := \{-\Delta K(0)\}_+$,

$$\frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N(X(t))] dt \leq \frac{H_N(0)}{NT} + \frac{a_1 \Lambda_T}{N h^{d+2}} + \frac{1}{2} \{B_{A,T}(h) + B_{V,T}(h)\} m_2(K) h^2. \quad (20)$$

If, in addition, $H_N(0) \leq \kappa_0 N$ and $T = N$, then

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N(X(t))] dt \leq \frac{\kappa_0}{N} + \frac{a_1 \Lambda_N}{N h^{d+2}} + \frac{1}{2} \{B_{A,N}(h) + B_{V,N}(h)\} m_2(K) h^2. \quad (21)$$

Thus the corresponding root-discrepancy bound is

$$\left(\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N(X(t))] dt \right)^{1/2} \leq \sqrt{\frac{\kappa_0}{N}} + \sqrt{\frac{a_1 \Lambda_N}{N h^{d+2}}} + h \sqrt{\frac{1}{2} \{B_{A,N}(h) + B_{V,N}(h)\} m_2(K)}. \quad (22)$$

Proof. The scaling identities follow by the change of variables $u = hv$:

$$m_2(K_h) = \int \|u\|^2 h^{-d} K(u/h) du = h^2 \int \|v\|^2 K(v) dv = h^2 m_2(K).$$

Also,

$$\Delta K_h(u) = h^{-d-2} (\Delta K)(u/h),$$

so $\Delta K_h(0) = h^{-d-2} \Delta K(0)$. Substituting these identities into (19) gives (20). If $H_N(0) \leq \kappa_0 N$ and $T = N$, then

$$\frac{H_N(0)}{NT} \leq \frac{\kappa_0 N}{N^2} = \frac{\kappa_0}{N},$$

which gives (21). Finally, (22) follows from $\sqrt{a+b+c} \leq \sqrt{a} + \sqrt{b} + \sqrt{c}$, for $a, b, c \geq 0$. $\square$

4.3 Optimizing the conservative bandwidth and the one-step size

The bandwidth form (21) separates three effects: the entropy initialization term, the finite-particle self-interaction term, and the quadrature term. Since the quadrature constants may depend on h , we write them explicitly as functions of the bandwidth. Define

$$A_N(h) := a_1 \Lambda_N(h), \quad C_N(h) := \frac{1}{2} \{B_{A,N}(h) + B_{V,N}(h)\} m_2(K).$$

Then (21) reads

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N(X(t))] dt \leq \frac{\kappa_0}{N} + \frac{A_N(h)}{N h^{d+2}} + C_N(h) h^2. \quad (23)$$

Thus the commonly stated bandwidth tradeoff is valid only after specifying how $A_N(h)$ and $C_N(h)$ scale with h .

Assume that, on the admissible bandwidth range, there are constants $A, C > 0$ and $0 \leq \beta < 2$ such that

$$A_N(h) \leq A, \quad C_N(h) \leq C h^{-\beta}. \quad (24)$$

Equivalently, the quadrature constants satisfy

$$B_{A,N}(h) + B_{V,N}(h) = O(h^{-\beta}).$$

Then (23) implies

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N(X(t))] dt \leq \frac{\kappa_0}{N} + \frac{A}{N h^{d+2}} + C h^{2-\beta}. \quad (25)$$

The last two terms are minimized at

$$h_{\star,N} := \left(\frac{(d+2)A}{(2-\beta)CN} \right)^{1/(d+4-\beta)}. \quad (26)$$

Indeed, differentiating

$$h \mapsto \frac{A}{N h^{d+2}} + C h^{2-\beta}$$

gives

$$-\frac{(d+2)A}{N h^{d+3}} + (2-\beta)C h^{1-\beta}.$$

Setting this derivative equal to zero gives (26). At this bandwidth,

$$\frac{A}{N h_{\star,N}^{d+2}} = \frac{2-\beta}{d+2} C h_{\star,N}^{2-\beta}.$$

Therefore

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N(X(t))] dt \leq \frac{\kappa_0}{N} + \frac{d+4-\beta}{d+2} C \left(\frac{(d+2)A}{(2-\beta)CN} \right)^{(2-\beta)/(d+4-\beta)}. \quad (27)$$

Thus, under (24), the squared residual velocity is

$$O\left(N^{-(2-\beta)/(d+4-\beta)}\right),$$

and the root residual velocity is

$$O\left(N^{-(2-\beta)/(2(d+4-\beta))}\right).$$

The h -uniform quadrature case is the special case $\beta = 0$, which recovers the root rate $N^{-1/(d+4)}$. If $\beta \geq 2$, the quadrature term $h^{2-\beta}$ does not vanish as $h \rightarrow 0$, and this quadrature argument does not yield a vanishing residual by bandwidth shrinkage. For example, the natural kernel-scale derivative scaling $B_{A,N}(h) + B_{V,N}(h) = O(h^{-4})$ corresponds to $\beta = 4$ and falls outside the regime of the optimized vanishing-rate statement.

The occupancy condition remains compatible with (26) whenever $Nh_{\star,N}^d \rightarrow \infty$ sufficiently quickly. For $0 \leq \beta < 2$,

$$Nh_{\star,N}^d \asymp N^{1-d/(d+4-\beta)} = N^{(4-\beta)/(d+4-\beta)},$$

which diverges faster than $\log N$ for fixed d and fixed $\beta < 2$.

The explicit one-step size η does not enter the continuous-time entropy inequality. It enters when the learned generator is interpreted through one more explicit correction $z \mapsto z + \eta b_x(z)$. For a fixed h , the theorem gives

$$\mathbb{E} [W_2^2(\mu_x^N, \tilde{\mu}_x^N)] \leq \eta^2 R_{N,h},$$

where $R_{N,h}$ denotes the right-hand side of (23). Taking $h = h_{\star,N}$ gives a residual correction of order

$$\eta N^{-(2-\beta)/(2(d+4-\beta))}$$

in root mean square, up to constants. If the explicit map is required to be a small stable Euler step and b_x has Lipschitz constant $L_b(h)$, a standard sufficient condition is $\eta L_b(h) \leq c$ for a numerical constant $c < 1$. In conservative models one often has $L_b(h) = O(h^{-2})$ under denominator and smoothness control, which suggests

$$\eta_{\star,N} \asymp h_{\star,N}^2 \asymp N^{-2/(d+4-\beta)}.$$

For Gaussian kernels, the non-conservative displacement velocity satisfies $u^{\text{disp}} = h^2 b$, so this conservative-step choice corresponds to a constant-order step size for the displacement formulation.

Remark 4.8 (Initialization). *If the initial particles are i.i.d. from a density μ_0 satisfying $\text{KL}(\mu_0 \parallel \rho) < \infty$, then*

$$p_0^N = \mu_0^{\otimes N} \implies H_N(0) = N \text{KL}(\mu_0 \parallel \rho).$$

Thus the condition $H_N(0) \leq \kappa_0 N$ holds with $\kappa_0 = \text{KL}(\mu_0 \parallel \rho)$.

5 Sufficient conditions for reciprocal KDE control on $\mathbb{R}^d$

The reciprocal-KDE condition (8) is needed because the center-evaluation divergence differentiates the i th KDE center and produces a factor $1/q_x(x_i)$. This section gives checkable sufficient conditions.

5.1 Deterministic local-occupancy condition

Assumption 5.1 (Kernel lower bound near the origin). *There exist constants $r_K > 0$ and $\kappa_K > 0$ such that*

$$K(u) \geq \kappa_K \quad \text{whenever } \|u\| \leq r_K. \quad (28)$$

Equivalently,

$$K_h(u) \geq \kappa_K h^{-d} \quad \text{whenever } \|u\| \leq r_K h.$$

For a configuration x , define the local neighbor count

$$M_i(x; h) := \#\{j \in \{1, \dots, N\} : \|x_j - x_i\| \leq r_K h\}. \quad (29)$$

The count includes $j = i$.

Proposition 5.2 (Local occupancy implies reciprocal control). *Let Assumption 5.1 hold. Suppose that a configuration x satisfies*

$$M_i(x; h) \geq \alpha N h^d \quad \text{for every } i = 1, \dots, N \quad (30)$$

for some $\alpha > 0$. Then

$$q_x(x_i) \geq \kappa_K \alpha \quad \text{for every } i = 1, \dots, N, \quad (31)$$

and hence

$$R_N(x) \leq \frac{1}{\kappa_K \alpha}. \quad (32)$$

Proof. Fix i . By definition,

$$q_x(x_i) = \frac{1}{N} \sum_{j=1}^N K_h(x_i - x_j).$$

For every j with $\|x_j - x_i\| \leq r_K h$, Assumption 5.1 gives

$$K_h(x_i - x_j) \geq \kappa_K h^{-d}.$$

Therefore

$$q_x(x_i) \geq \frac{1}{N} M_i(x; h) \kappa_K h^{-d}.$$

Using (30),

$$q_x(x_i) \geq \frac{1}{N} (\alpha N h^d) \kappa_K h^{-d} = \kappa_K \alpha.$$

Taking reciprocals and averaging over i yields (32). $\square$

5.2 A high-probability occupancy bound for i.i.d. particles

The previous proposition is deterministic. The next result shows that its hypothesis holds with high probability for i.i.d. samples from a density with uniform lower local mass.

Assumption 5.3 (Uniform lower local mass). *Let μ be a probability measure on $\mathbb{R}^d$. For the bandwidth h under consideration, there is a constant $p_0 > 0$ such that*

$$\mu(B(y, r_K h)) \geq p_0 h^d \quad \text{for } \mu\text{-a.e. } y. \quad (33)$$

Remark 5.4 (One way to verify Assumption 5.3). *Suppose μ has density f on a closed set $S \subset \mathbb{R}^d$, $f(y) \geq m_0 > 0$ for $y \in S$, and S satisfies the thickness condition*

$$\text{Leb}(S \cap B(y, r)) \geq \theta v_d r^d \quad \text{for all } y \in S \text{ and } 0 < r \leq r_0,$$

where v_d is the volume of the unit Euclidean ball. If $r_K h \leq r_0$, then

$$\mu(B(y, r_K h)) \geq m_0 \text{Leb}(S \cap B(y, r_K h)) \geq m_0 \theta v_d r_K^d h^d.$$

Thus Assumption 5.3 holds with

$$p_0 = m_0 \theta v_d r_K^d.$$

Proposition 5.5 (I.i.d. local occupancy). *Let $Y_1, \dots, Y_N$ be i.i.d. from a probability measure μ satisfying Assumption 5.3. Let*

$$Y = (Y_1, \dots, Y_N).$$

Assume $N \geq 2$. Define the event

$$\mathcal{E}_h := \left\{ M_i(Y; h) \geq \frac{p_0}{4} N h^d \text{ for every } i = 1, \dots, N \right\}.$$

Then

$$\mathbb{P}(\mathcal{E}_h^c) \leq N \exp\left(-\frac{p_0}{16} N h^d\right). \quad (34)$$

Consequently, under Assumption 5.1, on $\mathcal{E}_h$,

$$\mathbf{R}_N(Y) \leq \frac{4}{\kappa_K p_0}. \quad (35)$$

Moreover, since $K_h(0) = h^{-d} K(0)$,

$$\mathbf{R}_N(Y) \leq \frac{N h^d}{K(0)} \quad \text{always}, \quad (36)$$

and hence

$$\mathbb{E}[\mathbf{R}_N(Y)] \leq \frac{4}{\kappa_K p_0} + \frac{N h^d}{K(0)} N \exp\left(-\frac{p_0}{16} N h^d\right). \quad (37)$$

Proof. Fix i . Conditional on $Y_i = y$, define

$$Z_i := \sum_{j \neq i} \mathbf{1}\{\|Y_j - y\| \leq r_K h\}.$$

Then Z_i is binomial with parameters $N - 1$ and

$$p(y) := \mu(B(y, r_K h)).$$

By Assumption 5.3, $p(y) \geq p_0 h^d$ for μ -a.e. y . The neighbor count including the particle itself is

$$M_i(Y; h) = 1 + Z_i.$$

Let $m_i := (N - 1)p(y)$. For $N \geq 2$,

$$m_i \geq (N - 1)p_0 h^d \geq \frac{1}{2} N p_0 h^d.$$

If

$$Z_i \geq \frac{1}{2} m_i,$$

then

$$M_i(Y; h) = 1 + Z_i \geq \frac{1}{2} m_i \geq \frac{1}{4} N p_0 h^d.$$

Therefore

$$\mathbb{P}\left(M_i(Y; h) < \frac{1}{4} N p_0 h^d \mid Y_i = y\right) \leq \mathbb{P}\left(Z_i < \frac{1}{2} m_i \mid Y_i = y\right).$$

For a binomial random variable, the multiplicative Chernoff bound gives

$$\mathbb{P}\left(Z_i < \frac{1}{2}m_i \mid Y_i = y\right) \leq \exp\left(-\frac{m_i}{8}\right).$$

Using $m_i \geq (N-1)p_0h^d \geq Np_0h^d/2$,

$$\exp\left(-\frac{m_i}{8}\right) \leq \exp\left(-\frac{p_0}{16}Nh^d\right).$$

This bound is uniform in y . Integrating over Y_i and taking a union bound over $i = 1, \dots, N$ proves (34).

On $\mathcal{E}_h$, Proposition 5.2 applies with $\alpha = p_0/4$, giving (35). For the deterministic self-bound, observe that

$$q_Y(Y_i) = \frac{1}{N} \sum_{j=1}^N K_h(Y_i - Y_j) \geq \frac{1}{N} K_h(0) = \frac{K(0)}{Nh^d}.$$

Thus

$$\frac{1}{q_Y(Y_i)} \leq \frac{Nh^d}{K(0)} \quad \text{for every } i,$$

and averaging over i gives (36). Finally,

$$\begin{aligned} \mathbb{E}[\mathbf{R}_N(Y)] &= \mathbb{E}[\mathbf{R}_N(Y)\mathbf{1}_{\mathcal{E}_h}] + \mathbb{E}[\mathbf{R}_N(Y)\mathbf{1}_{\mathcal{E}_h^c}] \\ &\leq \frac{4}{\kappa_K p_0} + \frac{Nh^d}{K(0)} \mathbb{P}(\mathcal{E}_h^c). \end{aligned}$$

Substituting (34) proves (37). □

5.3 Propagation of occupancy along the continuous-time flow

The i.i.d. result controls the reciprocal KDE at a single time. To control it along the flow, one may combine initial occupancy at a slightly smaller scale with a Lipschitz-distortion bound for pairwise distances.

Assumption 5.6 (Lipschitz distortion of the drift). *For the realized trajectory on $[0, T]$, there exists an integrable function L_t such that*

$$\|b_{X(t)}(z) - b_{X(t)}(z')\| \leq L_t \|z - z'\| \quad \text{for all } z, z' \in \mathbb{R}^d, \quad (38)$$

for all $0 \leq t \leq T$. Define

$$\Gamma_T := \int_0^T L_t \, dt. \quad (39)$$

Lemma 5.7 (Pairwise distance distortion). *Under Assumption 5.6, for every pair i, j and every $0 \leq t \leq T$,*

$$\|X_i(t) - X_j(t)\| \leq e^{\Gamma_T} \|X_i(0) - X_j(0)\|. \quad (40)$$

Proof. For i, j , set $D_{ij}(t) := X_i(t) - X_j(t)$. From (1),

$$\frac{d}{dt} D_{ij}(t) = b_{X(t)}(X_i(t)) - b_{X(t)}(X_j(t)).$$

Using Assumption 5.6,

$$\frac{d}{dt} \|D_{ij}(t)\| \leq L_t \|D_{ij}(t)\|$$

whenever $D_{ij}(t) \neq 0$. The same inequality holds in the upper Dini derivative sense when $D_{ij}(t) = 0$. Gronwall's inequality gives

$$\|D_{ij}(t)\| \leq \exp\left(\int_0^t L_s ds\right) \|D_{ij}(0)\| \leq e^{\Gamma_T} \|D_{ij}(0)\|.$$

This proves the lemma. $\square$

Proposition 5.8 (Trajectory-level reciprocal control from initial occupancy). *Assume Assumptions 5.1 and 5.6. Suppose the initial configuration satisfies*

$$\#\{j : \|X_j(0) - X_i(0)\| \leq r_K h e^{-\Gamma_T}\} \geq \alpha N h^d e^{-d\Gamma_T} \quad \text{for every } i. \quad (41)$$

Then, for every $0 \leq t \leq T$,

$$R_N(X(t)) \leq \frac{e^{d\Gamma_T}}{\kappa_K \alpha}. \quad (42)$$

Proof. Fix i and $t \in [0, T]$. If

$$\|X_j(0) - X_i(0)\| \leq r_K h e^{-\Gamma_T},$$

then by Lemma 5.7,

$$\|X_j(t) - X_i(t)\| \leq e^{\Gamma_T} r_K h e^{-\Gamma_T} = r_K h.$$

Thus the number of particles within distance $r_K h$ of $X_i(t)$ is at least the number of particles within distance $r_K h e^{-\Gamma_T}$ of $X_i(0)$. By (41),

$$M_i(X(t); h) \geq \alpha N h^d e^{-d\Gamma_T}.$$

The proof of Proposition 5.2 gives

$$q_{X(t)}(X_i(t)) \geq \frac{1}{N} \{\alpha N h^d e^{-d\Gamma_T}\} \kappa_K h^{-d} = \kappa_K \alpha e^{-d\Gamma_T}.$$

Taking reciprocals and averaging over i proves (42). $\square$

Corollary 5.9 (I.i.d. initial data plus controlled distortion). *Let $X_1(0), \dots, X_N(0)$ be i.i.d. from a probability measure μ . Suppose that, with*

$$\tilde{h} := h e^{-\Gamma_T},$$

the lower local mass condition

$$\mu(B(y, r_K \tilde{h})) \geq p_0 \tilde{h}^d \quad \text{for } \mu\text{-a.e. } y \quad (43)$$

holds. Assume Assumptions 5.1 and 5.6. Then, with probability at least

$$1 - N \exp\left(-\frac{p_0}{16} N h^d e^{-d\Gamma_T}\right), \quad (44)$$

one has

$$\sup_{0 \leq t \leq T} R_N(X(t)) \leq \frac{4e^{d\Gamma_T}}{\kappa_K p_0}. \quad (45)$$

Furthermore,

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} R_N(X(t)) \right] \leq \frac{4e^{d\Gamma_T}}{\kappa_K p_0} + \frac{N h^d}{K(0)} N \exp\left(-\frac{p_0}{16} N h^d e^{-d\Gamma_T}\right). \quad (46)$$

Proof. Apply Proposition 5.5 at bandwidth $\tilde{h} = he^{-\Gamma_T}$. With probability at least (44), for every i ,

$$\#\{j : \|X_j(0) - X_i(0)\| \leq r_K h e^{-\Gamma_T}\} \geq \frac{p_0}{4} N (h e^{-\Gamma_T})^d.$$

This is (41) with $\alpha = p_0/4$. Therefore Proposition 5.8 gives (45).

For the expectation bound, use the same decomposition as in the proof of Proposition 5.5. On the high-probability event, use (45). On the complement, the deterministic self-bound holds at every time:

$$q_{X(t)}(X_i(t)) \geq \frac{1}{N} K_h(0) = \frac{K(0)}{N h^d},$$

so

$$\sup_{0 \leq t \leq T} R_N(X(t)) \leq \frac{N h^d}{K(0)}.$$

Combining with the probability estimate proves (46). $\square$

Remark 5.10 (Use in Theorem 4.5). *If the right-hand side of (46) is finite, then Assumption 2.3 holds with*

$$\Lambda_T = \frac{4e^{d\Gamma_T}}{\kappa_K p_0} + \frac{N h^d}{K(0)} N \exp\left(-\frac{p_0}{16} N h^d e^{-d\Gamma_T}\right).$$

In particular, if Γ_T is bounded independently of N and $N h^d \rightarrow \infty$, then the exponential term is negligible and $\Lambda_T = O(1)$.

6 Finite-particle rates for the non-conservative drifting method with Laplace kernel

We now prove a continuous-time finite-particle rate for the non-conservative displacement-based drifting field with Laplace kernel. The main point is that, unlike the Gaussian kernel, the Laplace mean-shift field is not exactly a score-difference field. Nevertheless, it admits a useful “sharp-score” representation, which yields a finite-particle convergence theorem with an additional scale-mismatch residual. The sharp companion kernel construction used in this section follows the sharp-normalization framework of Franz et al. (2026), specialized to the Laplace kernel. Our use of it is different: rather than replacing the original Laplace normalization by the sharp normalization, we use the sharp kernel to decompose the original non-conservative Laplace field into a preconditioned sharp-score mismatch plus a scale-mismatch residual.

Throughout this section, let

$$K_h(u) := c_d h^{-d} \exp\left(-\frac{\|u\|}{h}\right), \quad u \in \mathbb{R}^d,$$

where $c_d > 0$ is chosen so that $\int_{\mathbb{R}^d} K_h(u) du = 1$. For a probability measure α on $\mathbb{R}^d$, define its Laplace KDE

$$Q_{\alpha,h}(z) := \int_{\mathbb{R}^d} K_h(z - y) \alpha(dy),$$

and the non-conservative displacement mean-shift vector

$$M_{\alpha,h}(z) := \frac{\int_{\mathbb{R}^d} (y - z) K_h(z - y) \alpha(dy)}{Q_{\alpha,h}(z)}.$$

The non-conservative Laplace drifting field from a model distribution μ toward the data distribution ν is

$$u_{\nu,\mu,h}^{\text{Lap}}(z) := M_{\nu,h}(z) - M_{\mu,h}(z).$$

For a particle configuration $x = (x_1, \dots, x_N) \in (\mathbb{R}^d)^N$, let

$$\mu_x^N := \frac{1}{N} \sum_{j=1}^N \delta_{x_j}.$$

We analyze the self-masked, or leave-one-out, finite-particle version of the non-conservative Laplace drifting field. For each i , define

$$\mu_{x,-i}^N := \frac{1}{N-1} \sum_{j \neq i} \delta_{x_j},$$

and

$$u_{x,-i}(z) := M_{\nu,h}(z) - M_{\mu_{x,-i}^N,h}(z).$$

The continuous-time dynamics are

$$\dot{X}_i(t) = u_{X(t),-i}(X_i(t)), \quad i = 1, \dots, N. \quad (47)$$

The residual drift quantity we want to control is

$$\mathbf{V}_N^{\text{Lap}}(x) := \frac{1}{N} \sum_{i=1}^N \|u_{x,-i}(x_i)\|^2. \quad (48)$$

This is the mean squared norm of the non-conservative Laplace drift applied to the generated particles. Hence if one more explicit drift step is applied,

$$\tilde{x}_i = x_i + \eta u_{x,-i}(x_i),$$

then the identity coupling gives

$$W_2^2 \left(\frac{1}{N} \sum_{i=1}^N \delta_{x_i}, \frac{1}{N} \sum_{i=1}^N \delta_{\tilde{x}_i} \right) \leq \eta^2 \mathbf{V}_N^{\text{Lap}}(x).$$

Thus a bound on $\mathbf{V}_N^{\text{Lap}}$ controls how much an additional non-conservative Laplace drifting correction changes the generated distribution.

6.1 Sharp-score representation of the Laplace mean-shift field

The Laplace mean-shift field is not itself a KDE score, but it is a scaled score of a companion kernel. Define the sharp companion kernel

$$L_h(u) := h(\|u\| + h)K_h(u). \quad (49)$$

Let

$$R_{\alpha,h}(z) := \int_{\mathbb{R}^d} L_h(z-y) \alpha(dy),$$

and define the sharp-smoothed score

$$\sigma_{\alpha,h}(z) := \nabla \log R_{\alpha,h}(z).$$

Also define the scalar scale factor

$$a_{\alpha,h}(z) := \frac{R_{\alpha,h}(z)}{Q_{\alpha,h}(z)}.$$

Lemma 6.1 (Sharp-score representation). *For every probability measure α for which the following quantities are well-defined,*

$$M_{\alpha,h}(z) = a_{\alpha,h}(z)\sigma_{\alpha,h}(z). \quad (50)$$

Moreover,

$$a_{\alpha,h}(z) = h\{\bar{r}_{\alpha,h}(z) + h\}, \quad (51)$$

where

$$\bar{r}_{\alpha,h}(z) := \frac{\int_{\mathbb{R}^d} \|y - z\| K_h(z - y) \alpha(dy)}{\int_{\mathbb{R}^d} K_h(z - y) \alpha(dy)}$$

is the Laplace-weighted local mean radius.

Proof. Let $r = \|u\|$. Since

$$L_h(u) = h(r + h)c_d h^{-d} e^{-r/h},$$

the radial derivative is

$$\frac{d}{dr} \{h(r + h)c_d h^{-d} e^{-r/h}\} = h c_d h^{-d} e^{-r/h} - h(r + h)c_d h^{-d} \frac{1}{h} e^{-r/h}.$$

Combining the two terms,

$$\frac{d}{dr} \{h(r + h)c_d h^{-d} e^{-r/h}\} = c_d h^{1-d} e^{-r/h} - (r + h)c_d h^{-d} e^{-r/h}.$$

Since

$$c_d h^{1-d} e^{-r/h} = h c_d h^{-d} e^{-r/h},$$

we get

$$\frac{d}{dr} \{h(r + h)c_d h^{-d} e^{-r/h}\} = \{h - (r + h)\} c_d h^{-d} e^{-r/h} = -r K_h(u).$$

Therefore, for $u \neq 0$,

$$\nabla_u L_h(u) = -r K_h(u) \frac{u}{r} = -u K_h(u).$$

By continuity, the same identity holds at $u = 0$, because both sides vanish there. Taking $u = z - y$, we obtain

$$\nabla_z L_h(z - y) = -(z - y) K_h(z - y) = (y - z) K_h(z - y).$$

Hence

$$\nabla R_{\alpha,h}(z) = \int_{\mathbb{R}^d} \nabla_z L_h(z - y) \alpha(dy) = \int_{\mathbb{R}^d} (y - z) K_h(z - y) \alpha(dy).$$

Dividing by $Q_{\alpha,h}(z)$ gives

$$M_{\alpha,h}(z) = \frac{\nabla R_{\alpha,h}(z)}{Q_{\alpha,h}(z)} = \frac{R_{\alpha,h}(z)}{Q_{\alpha,h}(z)} \frac{\nabla R_{\alpha,h}(z)}{R_{\alpha,h}(z)} = a_{\alpha,h}(z) \sigma_{\alpha,h}(z).$$

This proves (50).

It remains to prove (51). By definition of L_h ,

$$R_{\alpha,h}(z) = \int_{\mathbb{R}^d} h(\|z - y\| + h) K_h(z - y) \alpha(dy).$$

Thus

$$R_{\alpha,h}(z) = h \int_{\mathbb{R}^d} \|z - y\| K_h(z - y) \alpha(dy) + h^2 \int_{\mathbb{R}^d} K_h(z - y) \alpha(dy).$$

The second integral is $Q_{\alpha,h}(z)$. Therefore

$$\frac{R_{\alpha,h}(z)}{Q_{\alpha,h}(z)} = h \frac{\int \|z - y\| K_h(z - y) \alpha(dy)}{Q_{\alpha,h}(z)} + h^2.$$

Since $\|z - y\| = \|y - z\|$, this is

$$a_{\alpha,h}(z) = h\bar{r}_{\alpha,h}(z) + h^2 = h\{\bar{r}_{\alpha,h}(z) + h\}.$$

The lemma follows. □

Let

$$Z_{\#,h} := \int_{\mathbb{R}^d} L_h(u) du.$$

The normalized sharp-smoothed data density is

$$\rho_{\nu,h}^{\#}(z) := \frac{R_{\nu,h}(z)}{Z_{\#,h}}.$$

Since the normalizing constant does not depend on z ,

$$\nabla \log \rho_{\nu,h}^{\#}(z) = \nabla \log R_{\nu,h}(z) = \sigma_{\nu,h}(z).$$

For the full empirical model, define

$$R_{x,h} := R_{\mu_x^N,h}, \quad Q_{x,h} := Q_{\mu_x^N,h}, \quad \sigma_{x,h} := \nabla \log R_{x,h}, \quad a_{x,h} := \frac{R_{x,h}}{Q_{x,h}}.$$

Also define the full non-conservative Laplace field

$$u_x(z) := M_{\nu,h}(z) - M_{\mu_x^N,h}(z).$$

By Lemma 6.1,

$$u_x(z) = a_{\nu,h}(z)\sigma_{\nu,h}(z) - a_{x,h}(z)\sigma_{x,h}(z).$$

Define the sharp score mismatch

$$b_x^{\#}(z) := \sigma_{\nu,h}(z) - \sigma_{x,h}(z),$$

and the Laplace scale-mismatch residual

$$e_x(z) := \{a_{\nu,h}(z) - a_{x,h}(z)\}\sigma_{\nu,h}(z). \tag{52}$$

Then

$$u_x(z) = a_{x,h}(z)b_x^{\#}(z) + e_x(z). \tag{53}$$

This decomposition is the key difference between the non-conservative Laplace field and the conservative field. If the local scales $a_{\nu,h}$ and $a_{x,h}$ match, then the non-conservative Laplace drift is a positive scalar preconditioning of a score mismatch. If they do not match, the residual e_x is unavoidable.

6.2 Entropy identity for the leave-one-out non-conservative field

Let p_t^N denote the joint density of $X(t)$. We measure it against the sharp-smoothed product target

$$(\rho_{\nu,h}^\#)^{\otimes N}(x) = \prod_{i=1}^N \rho_{\nu,h}^\#(x_i),$$

and define

$$H_N^\#(t) := \text{KL} \left(p_t^N \parallel (\rho_{\nu,h}^\#)^{\otimes N} \right). \quad (54)$$

Let

$$\mathcal{A}_\# f(z) := \nabla \cdot f(z) + \sigma_{\nu,h}(z) \cdot f(z)$$

be the Stein divergence associated with $\rho_{\nu,h}^\#$. Define

$$\mathbb{S}_N^{\text{Lap}}(x) := \frac{1}{N} \sum_{i=1}^N \mathcal{A}_\# u_{x,-i}(x_i). \quad (55)$$

Assumption 6.2 (Regularity for the leave-one-out Laplace flow). *The ODE (47) has a unique solution on $[0, T]$, the joint law admits a differentiable density p_t^N , and p_t^N solves the Liouville equation associated with the vector field $x \mapsto (u_{x,-1}(x_1), \dots, u_{x,-N}(x_N))$. All integrations by parts below are justified, and the trajectory avoids particle collisions and data atoms at which the exact Laplace field is not differentiable. Equivalently, one may replace K_h by a smooth regularized Laplace kernel and then pass to the exact Laplace limit under uniform bounds.*

Theorem 6.3 (Joint-entropy identity for non-conservative Laplace drifting). *Under Assumption 6.2,*

$$\frac{d}{dt} H_N^\#(t) = -N \mathbb{E}_t[\mathbb{S}_N^{\text{Lap}}(X(t))]. \quad (56)$$

Proof. Let

$$r_N^\#(x) := (\rho_{\nu,h}^\#)^{\otimes N}(x).$$

The joint density p_t^N satisfies

$$\partial_t p_t^N(x) + \sum_{i=1}^N \nabla_{x_i} \cdot \{p_t^N(x) u_{x,-i}(x_i)\} = 0.$$

By definition,

$$H_N^\#(t) = \int p_t^N(x) \log \frac{p_t^N(x)}{r_N^\#(x)} dx.$$

Differentiating under the integral gives

$$\frac{d}{dt} H_N^\#(t) = \int \partial_t p_t^N(x) \left\{ \log \frac{p_t^N(x)}{r_N^\#(x)} + 1 \right\} dx.$$

Since p_t^N is a probability density,

$$\int \partial_t p_t^N(x) dx = 0.$$

Therefore

$$\frac{d}{dt}H_N^\#(t) = \int \partial_t p_t^N(x) \log \frac{p_t^N(x)}{r_N^\#(x)} dx.$$

Using the Liouville equation,

$$\frac{d}{dt}H_N^\#(t) = - \sum_{i=1}^N \int \nabla_{x_i} \cdot \{p_t^N(x) u_{x,-i}(x_i)\} \log \frac{p_t^N(x)}{r_N^\#(x)} dx.$$

Integrating by parts in x_i ,

$$\frac{d}{dt}H_N^\#(t) = \sum_{i=1}^N \int p_t^N(x) u_{x,-i}(x_i) \cdot \nabla_{x_i} \log \frac{p_t^N(x)}{r_N^\#(x)} dx.$$

Now

$$\nabla_{x_i} \log r_N^\#(x) = \nabla \log \rho_{\nu,h}^\#(x_i) = \sigma_{\nu,h}(x_i).$$

Hence

$$\nabla_{x_i} \log \frac{p_t^N(x)}{r_N^\#(x)} = \nabla_{x_i} \log p_t^N(x) - \sigma_{\nu,h}(x_i).$$

Thus

$$\begin{aligned} \frac{d}{dt}H_N^\#(t) &= \sum_{i=1}^N \int p_t^N u_{x,-i}(x_i) \cdot \nabla_{x_i} \log p_t^N dx \\ &\quad - \sum_{i=1}^N \int p_t^N u_{x,-i}(x_i) \cdot \sigma_{\nu,h}(x_i) dx. \end{aligned}$$

Since

$$p_t^N \nabla_{x_i} \log p_t^N = \nabla_{x_i} p_t^N,$$

the first integral in the last display satisfies

$$\int p_t^N u_{x,-i}(x_i) \cdot \nabla_{x_i} \log p_t^N dx = \int u_{x,-i}(x_i) \cdot \nabla_{x_i} p_t^N dx.$$

Integrating by parts again,

$$\int u_{x,-i}(x_i) \cdot \nabla_{x_i} p_t^N dx = - \int p_t^N \nabla_{x_i} \cdot u_{x,-i}(x_i) dx.$$

Because $u_{x,-i}$ depends on x_i only through the evaluation point $z = x_i$, not through the leave-one-out model measure $\mu_{x,-i}^N$, we have

$$\nabla_{x_i} \cdot u_{x,-i}(x_i) = \nabla \cdot_z u_{x,-i}(z) \Big|_{z=x_i}.$$

Therefore

$$\frac{d}{dt}H_N^\#(t) = -\mathbb{E}_t \left[\sum_{i=1}^N \{ \nabla \cdot u_{X(t),-i}(X_i(t)) + \sigma_{\nu,h}(X_i(t)) \cdot u_{X(t),-i}(X_i(t)) \} \right].$$

The expression inside braces is exactly

$$\mathcal{A}_\# u_{X(t),-i}(X_i(t)).$$

Hence

$$\frac{d}{dt}H_N^\#(t) = -N \mathbb{E}_t[\mathbf{S}_N^{\text{Lap}}(X(t))],$$

which proves the identity. $\square$

6.3 Laplace coercivity with a scale-mismatch residual

Let

$$\rho_{x,h}^\#(z) := \frac{R_{x,h}(z)}{Z_{\#,h}}$$

be the full sharp-smoothed empirical density. Define the sharp-smoothed drift energy

$$\mathcal{V}_h^{\text{Lap}}(x) := \int_{\mathbb{R}^d} \|u_x(z)\|^2 \rho_{x,h}^\#(z) \, dz, \quad (57)$$

and the sharp-smoothed Stein drift

$$\mathcal{J}_h^{\text{Lap}}(x) := \int_{\mathbb{R}^d} \mathcal{A}_\# u_x(z) \rho_{x,h}^\#(z) \, dz. \quad (58)$$

Lemma 6.4 (Sharp-smoothed Stein identity). *For every fixed configuration x for which the integrations by parts are justified,*

$$\mathcal{J}_h^{\text{Lap}}(x) = \int_{\mathbb{R}^d} b_x^\#(z) \cdot u_x(z) \rho_{x,h}^\#(z) \, dz. \quad (59)$$

Proof. By definition,

$$\mathcal{A}_\# u_x = \nabla \cdot u_x + \sigma_{\nu,h} \cdot u_x.$$

Therefore

$$\mathcal{J}_h^{\text{Lap}}(x) = \int \rho_{x,h}^\# \nabla \cdot u_x + \int \rho_{x,h}^\# \sigma_{\nu,h} \cdot u_x.$$

For the first term, integration by parts gives

$$\int \rho_{x,h}^\# \nabla \cdot u_x = - \int \nabla \rho_{x,h}^\# \cdot u_x.$$

Since

$$\nabla \rho_{x,h}^\# = \rho_{x,h}^\# \nabla \log \rho_{x,h}^\# = \rho_{x,h}^\# \sigma_{x,h},$$

we obtain

$$\int \rho_{x,h}^\# \nabla \cdot u_x = - \int \rho_{x,h}^\# \sigma_{x,h} \cdot u_x.$$

Thus

$$\mathcal{J}_h^{\text{Lap}}(x) = \int \rho_{x,h}^\# (\sigma_{\nu,h} - \sigma_{x,h}) \cdot u_x.$$

By definition, $b_x^\# = \sigma_{\nu,h} - \sigma_{x,h}$. Hence

$$\mathcal{J}_h^{\text{Lap}}(x) = \int b_x^\# \cdot u_x \rho_{x,h}^\#,$$

which proves the claim. $\square$

Assumption 6.5 (Laplace scale alignment). *For all configurations visited by the flow on $[0, T]$, there exist constants $0 < \lambda_h \leq L_h < \infty$ and $\Delta_h \geq 0$ such that*

$$\lambda_h \leq a_{x,h}(z) \leq L_h \quad \text{for all } z \in \mathbb{R}^d, \quad (60)$$

and

$$\int_{\mathbb{R}^d} \|e_x(z)\|^2 \rho_{x,h}^\#(z) \, dz \leq \Delta_h^2. \quad (61)$$

Lemma 6.6 (Laplace coercivity). *Under Assumption 6.5,*

$$\mathcal{J}_h^{\text{Lap}}(x) \geq \gamma_h \mathcal{V}_h^{\text{Lap}}(x) - \beta_h \Delta_h^2, \quad (62)$$

where

$$\gamma_h := \frac{\lambda_h}{4L_h^2}, \quad \beta_h := \frac{\lambda_h}{2L_h^2} + \frac{1}{2\lambda_h}. \quad (63)$$

Proof. By Lemma 6.4,

$$\mathcal{J}_h^{\text{Lap}}(x) = \int b_x^\#(z) \cdot u_x(z) \rho_{x,h}^\#(z) \, dz.$$

Using the decomposition (53),

$$u_x(z) = a_{x,h}(z) b_x^\#(z) + e_x(z),$$

we get

$$b_x^\#(z) \cdot u_x(z) = a_{x,h}(z) \|b_x^\#(z)\|^2 + b_x^\#(z) \cdot e_x(z).$$

Thus

$$\mathcal{J}_h^{\text{Lap}}(x) = \int a_{x,h} \|b_x^\#\|^2 \rho_{x,h}^\# + \int b_x^\# \cdot e_x \rho_{x,h}^\#.$$

By $a_{x,h} \geq \lambda_h$,

$$\int a_{x,h} \|b_x^\#\|^2 \rho_{x,h}^\# \geq \lambda_h \int \|b_x^\#\|^2 \rho_{x,h}^\#.$$

For the second term, Young's inequality gives, pointwise,

$$-b_x^\# \cdot e_x \leq \frac{\lambda_h}{2} \|b_x^\#\|^2 + \frac{1}{2\lambda_h} \|e_x\|^2.$$

Equivalently,

$$b_x^\# \cdot e_x \geq -\frac{\lambda_h}{2} \|b_x^\#\|^2 - \frac{1}{2\lambda_h} \|e_x\|^2.$$

After integration,

$$\int b_x^\# \cdot e_x \rho_{x,h}^\# \geq -\frac{\lambda_h}{2} \int \|b_x^\#\|^2 \rho_{x,h}^\# - \frac{1}{2\lambda_h} \int \|e_x\|^2 \rho_{x,h}^\#.$$

Using (61),

$$\int b_x^\# \cdot e_x \rho_{x,h}^\# \geq -\frac{\lambda_h}{2} \int \|b_x^\#\|^2 \rho_{x,h}^\# - \frac{\Delta_h^2}{2\lambda_h}.$$

Combining the two lower bounds yields

$$\mathcal{J}_h^{\text{Lap}}(x) \geq \frac{\lambda_h}{2} \int \|b_x^\#\|^2 \rho_{x,h}^\# - \frac{\Delta_h^2}{2\lambda_h}.$$

Next, from $u_x = a_{x,h} b_x^\# + e_x$ and $a_{x,h} \leq L_h$,

$$\|u_x\|^2 \leq 2a_{x,h}^2 \|b_x^\#\|^2 + 2\|e_x\|^2 \leq 2L_h^2 \|b_x^\#\|^2 + 2\|e_x\|^2.$$

Integrating with respect to $\rho_{x,h}^\#$,

$$\mathcal{V}_h^{\text{Lap}}(x) \leq 2L_h^2 \int \|b_x^\#\|^2 \rho_{x,h}^\# + 2\Delta_h^2.$$

Rearranging,

$$\int \|b_x^\#\|^2 \rho_{x,h}^\# \geq \frac{\mathcal{V}_h^{\text{Lap}}(x) - 2\Delta_h^2}{2L_h^2}.$$

Substituting this into the lower bound for $\mathcal{J}_h^{\text{Lap}}$,

$$\mathcal{J}_h^{\text{Lap}}(x) \geq \frac{\lambda_h}{2} \frac{\mathcal{V}_h^{\text{Lap}}(x) - 2\Delta_h^2}{2L_h^2} - \frac{\Delta_h^2}{2\lambda_h}.$$

Therefore

$$\mathcal{J}_h^{\text{Lap}}(x) \geq \frac{\lambda_h}{4L_h^2} \mathcal{V}_h^{\text{Lap}}(x) - \left(\frac{\lambda_h}{2L_h^2} + \frac{1}{2\lambda_h} \right) \Delta_h^2.$$

This is exactly (62). □

6.4 From the sharp-smoothed coercivity to particle rates

The entropy identity controls the empirical leave-one-out Stein quantity $\mathbf{S}_N^{\text{Lap}}$, whereas the coercivity lemma is written for the full-field sharp-smoothed quantities $\mathcal{J}_h^{\text{Lap}}$ and $\mathcal{V}_h^{\text{Lap}}$. The following assumption records the required quadrature and leave-one-out approximation errors.

Assumption 6.7 (Quadrature and leave-one-out approximation). *For all configurations visited by the flow on $[0, T]$, there are constants $\varepsilon_{S,h,N} \geq 0$ and $\varepsilon_{V,h,N} \geq 0$ such that*

$$\left| \mathbf{S}_N^{\text{Lap}}(x) - \mathcal{J}_h^{\text{Lap}}(x) \right| \leq \varepsilon_{S,h,N}, \quad (64)$$

$$\left| \mathbf{V}_N^{\text{Lap}}(x) - \mathcal{V}_h^{\text{Lap}}(x) \right| \leq \varepsilon_{V,h,N}. \quad (65)$$

Remark 6.8. *Assumption 6.7 separates the entropy argument from the technical problem of controlling leave-one-out perturbations. The quantities $\varepsilon_{S,h,N}$ and $\varepsilon_{V,h,N}$ contain two distinct effects: the point-evaluation quadrature error and the difference between the full field U_x and the leave-one-out fields $U_{x,-i}$. Lemma 6.12 reduces these errors to Hessian bounds for the full-field integrands and the pointwise leave-one-out errors*

$$\ell_{S,h,N} := \frac{1}{N} \sum_{i=1}^N |\mathcal{A}_\# U_{x,-i}(x_i) - \mathcal{A}_\# U_x(x_i)|,$$

and

$$\ell_{V,h,N} := \frac{1}{N} \sum_{i=1}^N \left| \|U_{x,-i}(x_i)\|^2 - \|U_x(x_i)\|^2 \right|.$$

In this paper these leave-one-out errors are kept as explicit assumptions. A fully explicit bound in terms of N and h would require uniform lower bounds on the relevant Laplace KDE denominators, upper bounds on local weighted moments, and derivative stability estimates for the sharp-score and mean-shift maps after removing one particle. Under such bounds one expects leave-one-out perturbations to be of order $1/N$ times an h -dependent stability constant, but deriving the sharp dependence is a separate stability problem. The finite-particle rate in Theorem 6.9 below should therefore be interpreted as conditional on the displayed $\varepsilon_{S,h,N}$ and $\varepsilon_{V,h,N}$, rather than as an unconditional N, h -explicit theorem for the practical minibatch algorithm.

Theorem 6.9 (Continuous-time finite-particle rate for non-conservative Laplace drifting). *Let Assumptions 6.2, 6.5, and 6.7 hold on $[0, T]$. Then*

$$\frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{H_N^\#(0)}{\gamma_h NT} + \frac{\beta_h}{\gamma_h} \Delta_h^2 + \frac{\varepsilon_{S,h,N}}{\gamma_h} + \varepsilon_{V,h,N}, \quad (66)$$

where

$$\gamma_h = \frac{\lambda_h}{4L_h^2}, \quad \beta_h = \frac{\lambda_h}{2L_h^2} + \frac{1}{2\lambda_h}.$$

In particular, if $H_N^\#(0) \leq \kappa_0 N$ and $T = N$, then

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{\kappa_0}{\gamma_h N} + \frac{\beta_h}{\gamma_h} \Delta_h^2 + \frac{\varepsilon_{S,h,N}}{\gamma_h} + \varepsilon_{V,h,N}. \quad (67)$$

Equivalently,

$$\left(\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \right)^{1/2} \leq \sqrt{\frac{\kappa_0}{\gamma_h N}} + \sqrt{\frac{\beta_h}{\gamma_h}} \Delta_h + \sqrt{\frac{\varepsilon_{S,h,N}}{\gamma_h}} + \sqrt{\varepsilon_{V,h,N}}. \quad (68)$$

Proof. By Theorem 6.3,

$$\frac{d}{dt} H_N^\#(t) = -N \mathbb{E}_t[\mathbf{S}_N^{\text{Lap}}(X(t))].$$

Integrating from 0 to T ,

$$H_N^\#(T) - H_N^\#(0) = -N \int_0^T \mathbb{E}_t[\mathbf{S}_N^{\text{Lap}}(X(t))] dt.$$

Rearranging,

$$\int_0^T \mathbb{E}_t[\mathbf{S}_N^{\text{Lap}}(X(t))] dt = \frac{H_N^\#(0) - H_N^\#(T)}{N}.$$

Since relative entropy is nonnegative,

$$H_N^\#(T) \geq 0.$$

Therefore

$$\frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{S}_N^{\text{Lap}}(X(t))] dt \leq \frac{H_N^\#(0)}{NT}. \quad (69)$$

Next, fix a configuration x visited by the flow. By Assumption 6.7,

$$\mathbf{S}_N^{\text{Lap}}(x) \geq \mathcal{J}_h^{\text{Lap}}(x) - \varepsilon_{S,h,N}.$$

By Lemma 6.6,

$$\mathcal{J}_h^{\text{Lap}}(x) \geq \gamma_h \mathcal{V}_h^{\text{Lap}}(x) - \beta_h \Delta_h^2.$$

Therefore

$$\mathbf{S}_N^{\text{Lap}}(x) \geq \gamma_h \mathcal{V}_h^{\text{Lap}}(x) - \beta_h \Delta_h^2 - \varepsilon_{S,h,N}.$$

Again by Assumption 6.7,

$$\mathcal{V}_h^{\text{Lap}}(x) \geq \mathbf{V}_N^{\text{Lap}}(x) - \varepsilon_{V,h,N}.$$

Substituting this lower bound gives

$$\mathbf{S}_N^{\text{Lap}}(x) \geq \gamma_h \{\mathbf{V}_N^{\text{Lap}}(x) - \varepsilon_{V,h,N}\} - \beta_h \Delta_h^2 - \varepsilon_{S,h,N}.$$

Thus

$$\gamma_h \mathbf{V}_N^{\text{Lap}}(x) \leq \mathbf{S}_N^{\text{Lap}}(x) + \beta_h \Delta_h^2 + \varepsilon_{S,h,N} + \gamma_h \varepsilon_{V,h,N}.$$

Apply this with $x = X(t)$, take expectations, integrate over $t \in [0, T]$, and divide by T :

$$\gamma_h \frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{S}_N^{\text{Lap}}(X(t))] dt + \beta_h \Delta_h^2 + \varepsilon_{S,h,N} + \gamma_h \varepsilon_{V,h,N}.$$

Using (69),

$$\gamma_h \frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{H_N^\#(0)}{NT} + \beta_h \Delta_h^2 + \varepsilon_{S,h,N} + \gamma_h \varepsilon_{V,h,N}.$$

Dividing by γ_h proves (66). If $H_N^\#(0) \leq \kappa_0 N$ and $T = N$, then

$$\frac{H_N^\#(0)}{\gamma_h NT} \leq \frac{\kappa_0 N}{\gamma_h N^2} = \frac{\kappa_0}{\gamma_h N},$$

which proves (67). Finally, (68) follows from $\sqrt{a+b+c+d} \leq \sqrt{a} + \sqrt{b} + \sqrt{c} + \sqrt{d}$ for nonnegative a, b, c, d . $\square$

Remark 6.10 (Meaning of the Laplace rate). *The first term in (66) is the same entropy-driven optimization term as in the conservative theorem, except that it is divided by the Laplace coercivity constant γ_h . The terms $\varepsilon_{S,h,N}$ and $\varepsilon_{V,h,N}$ are finite-particle approximation errors: they compare the leave-one-out, center-evaluated quantities to the full-field sharp-smoothed quantities.*

The term

$$\frac{\beta_h}{\gamma_h} \Delta_h^2$$

in Theorem 6.9 is an irreducible residual term for the original non-conservative drifting method with Laplace kernel. It is not a proof artifact; it measures the fact that the non-conservative Laplace displacement field is not an exact score mismatch. It is also not a finite-particle fluctuation term and it does not vanish merely by taking $N \rightarrow \infty$ with h fixed. Consequently, the theorem proves convergence to a residual neighborhood whose size is controlled by Δ_h , not convergence to zero residual unless an additional alignment condition forces $\Delta_h \rightarrow 0$.

Using the identity

$$a_{\alpha,h}(z) = h\{\bar{r}_{\alpha,h}(z) + h\},$$

the residual can be written as

$$e_x(z) = h\{\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)\}\sigma_{\nu,h}(z).$$

Therefore, if

$$\|\sigma_{\nu,h}\|_\infty \leq G_h \quad \text{and} \quad \sup_z |\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)| \leq \delta_{r,h},$$

then

$$\Delta_h^2 \leq h^2 G_h^2 \delta_{r,h}^2.$$

Thus the non-conservative Laplace method has a vanishing residual only in regimes where the model and data local Laplace-weighted radii align. Without a separate argument giving $\delta_{r,h} \rightarrow 0$, the theorem should be read as a convergence-to-neighborhood result.

6.5 Checkable forms of the Laplace assumptions

The scale-alignment residual has a concrete interpretation in terms of local weighted radii. By Lemma 6.1,

$$a_{\alpha,h}(z) = h\{\bar{r}_{\alpha,h}(z) + h\}.$$

Therefore

$$a_{\nu,h}(z) - a_{x,h}(z) = h\{\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)\},$$

and the residual is

$$e_x(z) = h\{\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)\}\sigma_{\nu,h}(z).$$

Corollary 6.11 (Shell-alignment sufficient condition). *Assume that, along the trajectory on $[0, T]$, there are constants*

$$0 \leq r_{-,h} \leq r_{+,h} < \infty, \quad \delta_{r,h} \geq 0, \quad G_h < \infty,$$

such that, for all configurations visited by the flow and all $z \in \mathbb{R}^d$,

$$r_{-,h} \leq \bar{r}_{x,h}(z) \leq r_{+,h},$$

$$|\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)| \leq \delta_{r,h},$$

and

$$\|\sigma_{\nu,h}(z)\| \leq G_h.$$

Then Assumption 6.5 holds with

$$\lambda_h = h(r_{-,h} + h), \quad L_h = h(r_{+,h} + h),$$

and

$$\Delta_h^2 \leq h^2 G_h^2 \delta_{r,h}^2.$$

Consequently, if the hypotheses of Theorem 6.9 also hold, then

$$\frac{1}{T} \int_0^T \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{H_N^\#(0)}{\gamma_h N T} + \frac{\beta_h}{\gamma_h} h^2 G_h^2 \delta_{r,h}^2 + \frac{\varepsilon_{S,h,N}}{\gamma_h} + \varepsilon_{V,h,N}.$$

Proof. Since

$$a_{x,h}(z) = h\{\bar{r}_{x,h}(z) + h\},$$

the bound $r_{-,h} \leq \bar{r}_{x,h}(z) \leq r_{+,h}$ gives

$$h(r_{-,h} + h) \leq a_{x,h}(z) \leq h(r_{+,h} + h).$$

Thus (60) holds with the stated λ_h and L_h .

Next,

$$e_x(z) = h\{\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)\}\sigma_{\nu,h}(z).$$

Taking norms,

$$\|e_x(z)\| \leq h |\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)| \|\sigma_{\nu,h}(z)\|.$$

Using the assumed bounds,

$$\|e_x(z)\| \leq h \delta_{r,h} G_h.$$

Squaring,

$$\|e_x(z)\|^2 \leq h^2 \delta_{r,h}^2 G_h^2.$$

Since $\rho_{x,h}^\#$ is a probability density,

$$\int \|e_x(z)\|^2 \rho_{x,h}^\#(z) dz \leq h^2 \delta_{r,h}^2 G_h^2.$$

Thus $\Delta_h^2 \leq h^2 G_h^2 \delta_{r,h}^2$. Substitution into Theorem 6.9 proves the final display. $\square$

We next give one sufficient route to Assumption 6.7. Let

$$\bar{L}_h(u) := \frac{L_h(u)}{Z_{\#,h}}, \quad m_{2,\#,h} := \int_{\mathbb{R}^d} \|u\|^2 \bar{L}_h(u) du.$$

Because $\bar{L}_h$ is a bandwidth- h radial kernel, $m_{2,\#,h} = h^2 m_{2,\#,1}$ for a constant $m_{2,\#,1}$ depending only on dimension.

Lemma 6.12 (A sufficient quadrature bound). *Fix a configuration x . Define*

$$\phi_x(z) := \mathcal{A}_\# u_x(z), \quad \psi_x(z) := \|u_x(z)\|^2.$$

Assume that

$$\sup_{z \in \mathbb{R}^d} \|D^2 \phi_x(z)\|_{\text{op}} \leq B_{\phi,h}, \quad \sup_{z \in \mathbb{R}^d} \|D^2 \psi_x(z)\|_{\text{op}} \leq B_{\psi,h}.$$

Assume also that the leave-one-out point errors satisfy

$$\frac{1}{N} \sum_{i=1}^N |\mathcal{A}_\# u_{x,-i}(x_i) - \mathcal{A}_\# u_x(x_i)| \leq \ell_{S,h,N},$$

and

$$\frac{1}{N} \sum_{i=1}^N \left| \|u_{x,-i}(x_i)\|^2 - \|u_x(x_i)\|^2 \right| \leq \ell_{V,h,N}.$$

Then

$$\left| \mathcal{S}_N^{\text{Lap}}(x) - \mathcal{J}_h^{\text{Lap}}(x) \right| \leq \frac{1}{2} B_{\phi,h} m_{2,\#,h} + \ell_{S,h,N},$$

and

$$\left| \mathcal{V}_N^{\text{Lap}}(x) - \mathcal{V}_h^{\text{Lap}}(x) \right| \leq \frac{1}{2} B_{\psi,h} m_{2,\#,h} + \ell_{V,h,N}.$$

Proof. We prove the first inequality. The second is identical with ψ_x in place of ϕ_x .

Since

$$\rho_{x,h}^\#(z) = \frac{1}{N} \sum_{i=1}^N \bar{L}_h(z - x_i),$$

we have

$$\mathcal{J}_h^{\text{Lap}}(x) = \int \phi_x(z) \rho_{x,h}^\#(z) dz = \frac{1}{N} \sum_{i=1}^N \int \phi_x(z) \bar{L}_h(z - x_i) dz.$$

Also,

$$\frac{1}{N} \sum_{i=1}^N \phi_x(x_i) = \frac{1}{N} \sum_{i=1}^N \mathcal{A}_\# u_x(x_i).$$

Therefore

$$\left| \frac{1}{N} \sum_{i=1}^N \phi_x(x_i) - \mathcal{J}_h^{\text{Lap}}(x) \right| \leq \frac{1}{N} \sum_{i=1}^N \left| \phi_x(x_i) - \int \phi_x(z) \bar{L}_h(z - x_i) \, dz \right|.$$

Set $u = z - x_i$. Then

$$\int \phi_x(z) \bar{L}_h(z - x_i) \, dz = \int \phi_x(x_i + u) \bar{L}_h(u) \, du.$$

Taylor's formula gives

$$\phi_x(x_i + u) = \phi_x(x_i) + \nabla \phi_x(x_i) \cdot u + \int_0^1 (1-r) u^\top D^2 \phi_x(x_i + ru) u \, dr.$$

Integrating against $\bar{L}_h(u) \, du$, the constant term gives $\phi_x(x_i)$. The first-order term vanishes because $\bar{L}_h$ is even, hence

$$\int u \bar{L}_h(u) \, du = 0.$$

Thus

$$\left| \phi_x(x_i) - \int \phi_x(x_i + u) \bar{L}_h(u) \, du \right| \leq \int \bar{L}_h(u) \int_0^1 (1-r) \|D^2 \phi_x(x_i + ru)\|_{\text{op}} \|u\|^2 \, dr \, du.$$

Using the Hessian bound,

$$\left| \phi_x(x_i) - \int \phi_x(x_i + u) \bar{L}_h(u) \, du \right| \leq B_{\phi,h} \int_0^1 (1-r) \, dr \int \|u\|^2 \bar{L}_h(u) \, du.$$

Since $\int_0^1 (1-r) \, dr = 1/2$,

$$\left| \phi_x(x_i) - \int \phi_x(x_i + u) \bar{L}_h(u) \, du \right| \leq \frac{1}{2} B_{\phi,h} m_{2,\#,h}.$$

Averaging over i gives

$$\left| \frac{1}{N} \sum_{i=1}^N \mathcal{A}_{\#} u_x(x_i) - \mathcal{J}_h^{\text{Lap}}(x) \right| \leq \frac{1}{2} B_{\phi,h} m_{2,\#,h}.$$

Finally,

$$\mathbf{S}_N^{\text{Lap}}(x) = \frac{1}{N} \sum_{i=1}^N \mathcal{A}_{\#} u_{x,-i}(x_i).$$

By the assumed leave-one-out point-error bound,

$$\left| \mathbf{S}_N^{\text{Lap}}(x) - \frac{1}{N} \sum_{i=1}^N \mathcal{A}_{\#} u_x(x_i) \right| \leq \ell_{S,h,N}.$$

The triangle inequality gives

$$\left| \mathbf{S}_N^{\text{Lap}}(x) - \mathcal{J}_h^{\text{Lap}}(x) \right| \leq \frac{1}{2} B_{\phi,h} m_{2,\#,h} + \ell_{S,h,N}.$$

This proves the first bound. $\square$

Finally, the leave-one-out denominators are controlled by the same local-occupancy mechanism used for conservative drifting.

Lemma 6.13 (Leave-one-out Laplace denominator control). *Fix $r_0 > 0$. For the Laplace kernel,*

$$K_h(u) \geq c_d h^{-d} e^{-r_0} \quad \text{whenever } \|u\| \leq r_0 h.$$

For a configuration x , define

$$n_i^{(-)}(x) := \#\{j \neq i : \|x_j - x_i\| \leq r_0 h\}.$$

If

$$n_i^{(-)}(x) \geq \alpha(N-1)h^d \quad \text{for every } i,$$

then

$$Q_{\mu_{x,-i},h}^N(x_i) \geq c_d e^{-r_0} \alpha \quad \text{for every } i.$$

Proof. By definition,

$$Q_{\mu_{x,-i},h}^N(x_i) = \frac{1}{N-1} \sum_{j \neq i} K_h(x_i - x_j).$$

For every $j \neq i$ satisfying $\|x_j - x_i\| \leq r_0 h$,

$$K_h(x_i - x_j) = c_d h^{-d} \exp\left(-\frac{\|x_i - x_j\|}{h}\right) \geq c_d h^{-d} e^{-r_0}.$$

Therefore

$$Q_{\mu_{x,-i},h}^N(x_i) \geq \frac{1}{N-1} n_i^{(-)}(x) c_d h^{-d} e^{-r_0}.$$

Using $n_i^{(-)}(x) \geq \alpha(N-1)h^d$,

$$Q_{\mu_{x,-i},h}^N(x_i) \geq \frac{1}{N-1} \{\alpha(N-1)h^d\} c_d h^{-d} e^{-r_0} = c_d e^{-r_0} \alpha.$$

This proves the claim. □

6.6 Bandwidth and step-size choices for the non-conservative drifting method with Laplace kernel

The leave-one-out Laplace theorem has a different bandwidth tradeoff from the conservative theorem. Because the self-masked field removes the self-interaction term from the entropy identity, there is no analogue of the conservative self-interaction term $1/(Nh^{d+2})$ in (66). Instead, the bandwidth appears through the scale-mismatch residual and the quadrature and leave-one-out errors.

A useful simplified form is obtained from Corollary 6.11 and Lemma 6.12. Suppose that along the trajectory

$$\frac{\beta_h}{\gamma_h} \Delta_h^2 \leq C_{\Delta,h} h^2 G_h^2 \delta_{r,h}^2,$$

and

$$\frac{\varepsilon_{S,h,N}}{\gamma_h} + \varepsilon_{V,h,N} \leq C_{Q,h} h^2 + \frac{C_{\text{loo},h}}{N}.$$

Then, for $T = N$ and $H_N^\#(0) \leq \kappa_0 N$, Theorem 6.9 gives

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{\kappa_0}{\gamma_h N} + C_{\Delta,h} h^2 G_h^2 \delta_{r,h}^2 + C_{Q,h} h^2 + \frac{C_{\text{loo},h}}{N}. \quad (70)$$

If γ_h is bounded below and $C_{\Delta,h}, C_{Q,h}, C_{\text{loo},h}$ remain bounded, then the deterministic part of the rate improves as h decreases, provided $h G_h \delta_{r,h}$ also decreases or remains controlled. The bandwidth should therefore be chosen as small as permitted by denominator control and regularity. A typical local-occupancy requirement is

$$N h^d \gtrsim \log N,$$

which suggests

$$h_N \asymp \left(\frac{\log N}{N} \right)^{1/d} \quad (71)$$

when the constants in (70) are stable at that scale. In the benign case where $G_h \delta_{r,h} = O(1)$, this gives

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \lesssim \frac{1}{N} + \left(\frac{\log N}{N} \right)^{2/d},$$

up to the displayed constants and residual assumptions. The corresponding root residual velocity is

$$O(N^{-1/2}) + O\left((\log N/N)^{1/d}\right).$$

If the shell mismatch is worse, the term $h G_h \delta_{r,h}$ should be kept explicitly; if it is better, the rate improves accordingly.

The one-step drift size η again enters only when the continuous-time bound is converted into a statement about an explicit correction. For the non-conservative Laplace field,

$$\tilde{x}_i = x_i + \eta u_{x,-i}(x_i)$$

satisfies

$$W_2^2(\mu_x^N, \tilde{\mu}_x^N) \leq \eta^2 \mathbf{V}_N^{\text{Lap}}(x).$$

Thus a typical residual correction at the optimized bandwidth (71) is of order

$$\eta \left\{ N^{-1/2} + (\log N/N)^{1/d} + h_N G_{h_N} \delta_{r,h_N} \right\}$$

in root mean square. If the explicit displacement map must be a stable Euler step and the leave-one-out Laplace field has Lipschitz constant $L_u(h)$, a sufficient condition is $\eta L_u(h) \leq c < 1$. Under denominator control and bounded local-radius derivatives, the non-conservative displacement field often has $L_u(h) = O(1)$, in which case a constant-order η is admissible. If $L_u(h)$ grows with decreasing h , the admissible step size should be reduced according to $\eta \asymp 1/L_u(h)$.

6.7 Comparison between the conservative and non-conservative results

The conservative velocity field studied earlier is

$$b_x(z) = s_\rho(z) - s_x(z),$$

where $s_\rho = \nabla \log \rho$ is the score of the smoothed target and $s_x = \nabla \log q_x$ is the score of the model KDE. Its squared center velocity is

$$\mathbf{V}_N^{\text{cons}}(x) := \mathbf{V}_N(x) = \frac{1}{N} \sum_{i=1}^N \|b_x(x_i)\|^2.$$

The finite-particle rate for this conservative method has the form

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N^{\text{cons}}(X(t))] dt \leq \frac{\kappa_0}{N} + \frac{C_{\text{self}}}{N h^{d+2}} + C_{\text{quad}}(h) h^2,$$

where the second term is the reciprocal-KDE self-interaction correction and

$$C_{\text{quad}}(h) = \frac{1}{2} \{B_{A,N}(h) + B_{V,N}(h)\} m_2(K)$$

is the point-evaluation quadrature constant. Hence the conservative method has a clean score/Fisher structure, but its optimized bandwidth rate depends critically on the h -dependence of the quadrature constants. If $C_{\text{quad}}(h) = O(1)$, the root residual-velocity rate is $N^{-1/(d+4)}$. If $C_{\text{quad}}(h) = O(h^{-\beta})$ with $0 \leq \beta < 2$, the optimized root rate is instead

$$N^{-(2-\beta)/(2(d+4-\beta))}.$$

If $\beta \geq 2$, the present quadrature argument does not give a vanishing rate by shrinking h .

For the non-conservative drifting method with Laplace kernel, the corresponding rate is

$$\frac{1}{N} \int_0^N \mathbb{E}_t[\mathbf{V}_N^{\text{Lap}}(X(t))] dt \leq \frac{\kappa_0}{\gamma_h N} + \frac{\beta_h}{\gamma_h} \Delta_h^2 + \frac{\varepsilon_{S,h,N}}{\gamma_h} + \varepsilon_{V,h,N}.$$

The first term is the same entropy-driven finite-particle optimization term, up to the Laplace coercivity constant γ_h . The main new term is

$$\frac{\beta_h}{\gamma_h} \Delta_h^2,$$

which measures the failure of the non-conservative Laplace mean-shift field to be an exact score-difference field. Explicitly,

$$\Delta_h^2 \geq \int \|\{a_{\nu,h}(z) - a_{x,h}(z)\} \sigma_{\nu,h}(z)\|^2 \rho_{x,h}^\#(z) dz.$$

Equivalently, since

$$a_{\alpha,h}(z) = h\{\bar{r}_{\alpha,h}(z) + h\},$$

this residual is controlled by the mismatch between the data and model local Laplace-weighted radii:

$$a_{\nu,h}(z) - a_{x,h}(z) = h\{\bar{r}_{\nu,h}(z) - \bar{r}_{x,h}(z)\}.$$

The two theories therefore share the same entropy backbone but differ in their coercivity mechanisms. For the conservative method, the drift is already a score mismatch, so the Stein identity directly gives a Fisher-type quantity after quadrature. For the non-conservative Laplace method, the drift decomposes as

$$u_x(z) = a_{x,h}(z) b_x^\#(z) + e_x(z),$$

where $a_{x,h}$ is a positive scalar preconditioner and e_x is the Laplace scale-mismatch residual. Thus the non-conservative method has a comparable rate only when the local scale factors of the data and model are aligned well enough that Δ_h is small.

There is also a difference in the finite-particle correction. The conservative theorem above uses the center-evaluation field with the full KDE, so differentiating the i th particle’s own KDE contribution produces a reciprocal-KDE self-interaction correction. In the non-conservative Laplace result we use the self-masked, leave-one-out field $u_{x,-i}$. This removes the self-interaction term from the entropy identity. The price is the leave-one-out approximation error contained in $\varepsilon_{S,h,N}$ and $\varepsilon_{V,h,N}$, which is controlled by denominator lower bounds and local occupancy conditions such as Lemma 6.13.

Finally, the target density used by the two analyses is different. The conservative theorem uses the ordinary smoothed target $\rho = K_h * \nu$. The non-conservative Laplace theorem uses the sharp-smoothed target

$$\rho_{\nu,h}^\#(z) = \frac{1}{Z_{\#,h}} \int L_h(z-y)\nu(dy), \quad L_h(u) = h(\|u\| + h)K_h(u).$$

This difference is not an artifact of the proof. It reflects the identity

$$\nabla_z L_h(z-y) = (y-z)K_h(z-y),$$

which is the structural reason the non-conservative Laplace displacement field can be related to a score mismatch at all. In the Gaussian case, the non-conservative displacement field is simply a constant multiple of the conservative field; in the Laplace case, it is instead a variable-scale preconditioned sharp-score field plus the residual e_x .

References

- Michael Samuel Albergo and Eric Vanden-Eijnden. Building Normalizing Flows with Stochastic Interpolants. In *The Eleventh International Conference on Learning Representations*, 2023. URL <https://openreview.net/forum?id=li7qeBbCR1t>.
- Sayan Banerjee, Krishnakumar Balasubramanian, and Promit Ghosal. Improved Finite-Particle Convergence Rates for Stein Variational Gradient Descent. In *The Thirteenth International Conference on Learning Representations*, 2025. URL <https://openreview.net/forum?id=sbG8qhMjkZ>.
- Andrew D Barbour. Stein’s method and Poisson process convergence. *Journal of Applied Probability*, 25(A):175–184, 1988.
- Nicholas Matthew Boffi, Michael Samuel Albergo, and Eric Vanden-Eijnden. Flow map matching with stochastic interpolants: A mathematical framework for consistency models. *Transactions on Machine Learning Research*, 2025. ISSN 2835-8856. URL <https://openreview.net/forum?id=cqDH0e6ak2>.
- Jiarui Cao, Zixuan Wei, and Yuxin Liu. Gradient flow drifting: Generative modeling via Wasserstein gradient flows of KDE-approximated divergences. *arXiv preprint arXiv:2603.10592*, 2026.
- Mingyang Deng, He Li, Tianhong Li, Yilun Du, and Kaiming He. Generative modeling via drifting. *arXiv preprint arXiv:2602.04770*, 2026.

- Théo Dumont, Théo Lacombe, and François-Xavier Vialard. Learning Monge maps with constrained drifting models. *arXiv preprint arXiv:2603.25182*, 2026.
- Maria Esteban-Casadevall, Jorge Carrasco-Pollo, Max Welling, Jan-Willem van de Meent, Erik J. Bekkers, and Floor Eijkelboom. Kernel-gradient drifting models. *arXiv preprint arXiv:2605.10727*, 2026.
- Yihao Feng, Dilin Wang, and Qiang Liu. Learning to draw samples with amortized Stein variational gradient descent. In *The Conference on Uncertainty in Artificial Intelligence (UAI)*, 2017.
- Leonard Franz, Sebastian Hoffmann, and Georg Martius. Drifting fields are not conservative. *arXiv preprint arXiv:2604.06333*, 2026.
- Jackson Gorham and Lester Mackey. Measuring sample quality with Stein’s method. *Advances in neural information processing systems*, 28, 2015.
- Arthur Gretton, Li Kevin Wenliang, Alexandre Galashov, James Thornton, Valentin De Bortoli, and Arnaud Doucet. On the Wasserstein Gradient Flow Interpretation of Drifting Models. *arXiv preprint arXiv:2605.05118*, 2026.
- Ping He, Om Khangaonkar, Hamed Pirsiavash, Yikun Bai, and Soheil Kolouri. Sinkhorn-Drifting Generative Models. *arXiv preprint arXiv:2603.12366*, 2026a.
- Ye He, Krishnakumar Balasubramanian, Sayan Banerjee, and Promit Ghosal. Finite-Particle Rates for Regularized Stein Variational Gradient Descent. *arXiv preprint arXiv:2602.05172*, 2026b.
- Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising Diffusion Probabilistic Models. In *Advances in Neural Information Processing Systems*, volume 33, pages 6840–6851, 2020.
- Chieh-Hsin Lai, Bac Nguyen, Naoki Murata, Yuhta Takida, Toshimitsu Uesaka, Yuki Mitsufuji, Stefano Ermon, and Molei Tao. A unified view of drifting and score-based models. *arXiv preprint arXiv:2603.07514*, 2026.
- Hak Geun Lee and Hyonho Chun. Identifiability and Stability of Generative Drifting with Companion-Elliptic Kernel Families. *arXiv preprint arXiv:2604.24196*, 2026.
- Zhiqi Li and Bo Zhu. A Long-Short Flow-Map Perspective for Drifting Models. *arXiv preprint arXiv:2602.20463*, 2026.
- Yaron Lipman, Ricky T. Q. Chen, Heli Ben-Hamu, Maximilian Nickel, and Matthew Le. Flow Matching for Generative Modeling. In *The Eleventh International Conference on Learning Representations*, 2023. URL <https://openreview.net/forum?id=PqvMRDCJT9t>.
- Qiang Liu. Stein variational gradient descent as gradient flow. *Advances in neural information processing systems*, 30, 2017.
- Qiang Liu and Dilin Wang. Stein variational gradient descent: A general purpose bayesian inference algorithm. *Advances in neural information processing systems*, 29, 2016.
- Xingchao Liu, Chengyue Gong, and Qiang Liu. Flow Straight and Fast: Learning to Generate and Transfer Data with Rectified Flow. In *The Eleventh International Conference on Learning Representations*, 2023. URL <https://openreview.net/forum?id=XVjTT1nw5z>.

- Tim Salimans and Jonathan Ho. Progressive distillation for fast sampling of diffusion models. In *International Conference on Learning Representations*, 2022. URL <https://openreview.net/forum?id=TIdIXIpzhoI>.
- Jiaxin Shi and Lester Mackey. A finite-particle convergence rate for Stein variational gradient descent. *Advances in Neural Information Processing Systems*, 36:26831–26844, 2023.
- Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-Based Generative Modeling through Stochastic Differential Equations. In *International Conference on Learning Representations*, 2021. URL <https://openreview.net/forum?id=PXTIG12RRHS>.
- Yang Song, Prafulla Dhariwal, Mark Chen, and Ilya Sutskever. Consistency models. In *International Conference on Machine Learning*, 2023.
- Erkan Turan and Maks Ovsjanikov. Generative drifting is secretly score matching: A spectral and variational perspective. *arXiv preprint arXiv:2603.09936*, 2026.
- Romann M. Weber. The Score-Difference Flow for Implicit Generative Modeling. *Transactions on Machine Learning Research*, 2023. ISSN 2835-8856. URL <https://openreview.net/forum?id=dpGSNLUCzu>.
- Guoqiang Zhang, Kenta Niwa, and W. Bastiaan Kleijn. Lookahead drifting model. *arXiv preprint arXiv:2605.04060*, 2026.

A Stop-gradient training and the particle ODE

We briefly explain how the continuous-time particle dynamics arise from the practical stop-gradient training rule. Let $g_\theta : \mathcal{Z} \rightarrow \mathbb{R}^d$ be the generator and let

$$x_i^k := g_{\theta_k}(\xi_i), \quad i = 1, \dots, N,$$

be the generated batch at training step k , for fixed latent variables $\xi_1, \dots, \xi_N$. Let v_{x^k} denote the drift field computed from the current generated batch

$$x^k = (x_1^k, \dots, x_N^k)$$

and the data batch. In the original displacement-based drifting method (Deng et al., 2026), v_{x^k} is the non-conservative field $u_{\nu, \mu_{x^k}, h}^{\text{disp}}$; in the conservative method studied in this paper, it is the KDE-score field b_{x^k} . The practical target for the i -th generated sample is

$$y_i^k := \text{sg}\{x_i^k + \eta v_{x^k}(x_i^k)\}.$$

Here sg denotes the stop-gradient operator. For any quantity $a(\theta)$ appearing in the computational graph,

$$\text{sg}\{a(\theta)\} = a(\theta) \quad \text{in the forward pass,} \quad \nabla_\theta \text{sg}\{a(\theta)\} = 0 \quad \text{in the backward pass.}$$

Thus sg preserves the numerical value of its argument but removes all derivatives through that argument. In particular, the target $y_i = \text{sg}\{x_i + \eta v_x(x_i)\}$ is evaluated as $x_i + \eta v_x(x_i)$, but is treated as a constant when differentiating the regression loss with respect to the generator parameters.

The corresponding regression loss is

$$\mathcal{L}_k(\theta) := \frac{1}{2N} \sum_{i=1}^N \|g_\theta(\xi_i) - y_i^k\|^2. \quad (72)$$

The stop-gradient operation means that y_i^k is treated as a constant when differentiating $\mathcal{L}_k$ with respect to θ . Thus the derivative of the loss is

$$\nabla_\theta \mathcal{L}_k(\theta) = \frac{1}{N} \sum_{i=1}^N D_\theta g_\theta(\xi_i)^\top \{g_\theta(\xi_i) - y_i^k\},$$

with no derivative through the map $x^k \mapsto v_{x^k}$. Therefore the training step is a frozen-target regression problem: the generator is trained to move its current outputs toward the fixed points

$$x_i^k + \eta v_{x^k}(x_i^k).$$

The idealized exact-regression limit is the limit in which this regression problem is solved exactly at every step and the generator class is rich enough to interpolate the frozen targets on the sampled latent variables. In that limit, the next generator satisfies

$$g_{\theta_{k+1}}(\xi_i) = y_i^k = x_i^k + \eta v_{x^k}(x_i^k), \quad i = 1, \dots, N.$$

Consequently the generated particles obey the explicit Euler update

$$x_i^{k+1} = x_i^k + \eta v_{x^k}(x_i^k), \quad i = 1, \dots, N. \quad (73)$$

If the regression is not exact, then (73) holds with an additional approximation error

$$e_i^k := g_{\theta_{k+1}}(\xi_i) - \{x_i^k + \eta v_{x^k}(x_i^k)\}.$$

Our analysis neglects this optimization and approximation error and studies the idealized particle system (73).

Finally, define $t_k = k\eta$ and let $X_i^\eta(t)$ be the piecewise-linear interpolation of the Euler iterates. Under the usual consistency and stability conditions for explicit Euler schemes, for example local Lipschitz continuity of the interacting-particle vector field

$$F_i(x) := v_x(x_i), \quad x = (x_1, \dots, x_N),$$

the interpolation $X^\eta(t)$ converges as $\eta \rightarrow 0$ on finite time intervals to the solution of

$$\dot{X}_i(t) = F_i(X(t)) = v_{X(t)}(X_i(t)), \quad i = 1, \dots, N.$$

Here, v_x can either be the conservative KDE-score field or the non-conservative leave-one-out Laplace field. These are the continuous-time particle dynamics analyzed in the main text. Thus stop-gradient is used to justify the frozen-field Euler update; once this particle ODE is assumed, the entropy identities and convergence bounds are properties of the resulting interacting-particle flow.

We emphasize that the passage from the stop-gradient Euler update to the continuous-time ODE is an idealized modeling step. Let $F(x) := (F_1(x), \dots, F_N(x))$. The exact-regression stop-gradient update is

$$x^{k+1} = x^k + \eta F(x^k).$$

If F is Lipschitz on a forward-invariant region $\mathcal{D} \subset (\mathbb{R}^d)^N$ with Lipschitz constant $L_{h,\mathcal{D}}$, and if the Euler iterates and the ODE trajectory remain in $\mathcal{D}$, then the standard explicit-Euler estimate gives, on a fixed time interval $[0, T]$,

$$\max_{0 \leq k\eta \leq T} \|x^k - X(k\eta)\| \leq C_{T,h,\mathcal{D}}\eta,$$

where $C_{T,h,\mathcal{D}}$ depends on T , $L_{h,\mathcal{D}}$, and a bound on F on $\mathcal{D}$.

For the interacting-particle systems studied here, $L_{h,\mathcal{D}}$ can grow rapidly as $h \rightarrow 0$ and depends on denominator lower bounds such as $q_x(x_i) \geq \lambda$. Therefore the continuous-time approximation requires a step-size condition of the form

$$\eta L_{h,\mathcal{D}} \ll 1.$$

If, for instance, $L_{h,\mathcal{D}} = O(h^{-\alpha})$, then one needs

$$\eta = o(h^\alpha)$$

for the Euler approximation error to vanish. Thus our continuous-time results should not be interpreted as unconditional finite-step guarantees for the practical algorithm. They describe the idealized stop-gradient, exact-regression, small-step limit under independent stability and denominator-control assumptions.

If the regression step is not exact, the particle update has the form $x^{k+1} = x^k + \eta F(x^k) + e^k$, as mentioned above. Under the same Lipschitz assumptions, the discrete-to-continuous error contains an additional accumulated term of size

$$\sum_{k\eta \leq T} e^{L_{h,\mathcal{D}}(T-k\eta)} \|e^k\|.$$

Thus tracking the practical neural-network training dynamics would require bounds on optimization error, approximation error, minibatch noise, and the h -dependent stability constant. These effects are outside the scope of the continuous-time entropy analysis.