

LOW-RANK AND GRAPHON LIMITS FOR DYNAMIC THRESHOLD DISTRESS CONTAGION IN HETEROGENEOUS FINANCIAL NETWORKS

PENGBIN FENG

ABSTRACT. We study a deterministic contagion model for a large population of financial institutions connected by a weighted directed exposure matrix. The sign convention and loss term are motivated by a default cascade with partial recovery and interest servicing, whereas the dynamic model records distress through occupation time and therefore permits recovery. A rank- K factorization yields an exact reduction of the finite network to K macroscopic feedback coordinates. For bounded Lipschitz losses, the reduced dynamics form a nonautonomous K -dimensional ODE; we prove Wasserstein stability with respect to the type law and derive a transport representation for the joint state–factor distribution. On a fixed latent space, the associated directed-kernel equation is well posed and L^1 -stable, and a quantitative bridge theorem separates finite-population error from kernel-approximation error. For the indicator loss, we establish fixed-rank well-posedness under threshold regularity and a Vapnik–Chervonenkis-type estimate for measurable selections of sampled solutions. At the graphon level, we prove well-posedness for factorized kernels and for piecewise- C^1 kernel–profile pairs satisfying uniform transversality, together with a perturbation theorem for uniformly transverse approximation families. A sovereign-overlap illustration based on the 2025 EBA transparency exercise computes factor loadings and a priori sensitivity bounds from public disclosures; resampling errors on the empirical 117-bank population are consistent with the predicted $N^{-1/2}$ scale.

1. INTRODUCTION

A persistent difficulty in systemic-risk modeling is to retain both network heterogeneity and dynamic tractability. Static financial-network models preserve detailed bilateral structure and yield sharp resilience criteria, but time evolution is usually handled only through one-shot clearing maps or cascade iterations [18, 13, 4, 19, 26]. Dynamic models based on homogeneous or nearly homogeneous mean-field interaction are analytically much more tractable and accommodate diffusion, control, and game-theoretic extensions, but they flatten the exposure matrix precisely where real infrastructures are most heterogeneous [12, 28, 22]. Several financial infrastructures occupy this intermediate regime. Large-value payment systems, tiered interbank markets, central clearing, and bank–NBFI architectures are neither fully homogeneous nor completely arbitrary; they are organized around a small number of dominant transmission channels [11, 15, 1, 29, 2].

The homogeneous mean-field model provides the reference case. Let N institutions have cash buffers $X^{i,N}$ evolving under a common trend and a symmetric distress feedback,

$$(1) \quad X_t^{i,N} = x_i^N + \mu t - \frac{1}{N} \sum_{j=1}^N \int_0^t \ell(X_s^{j,N}) ds, \quad 1 \leq i \leq N,$$

2020 *Mathematics Subject Classification.* 60K35, 60J60, 91G40, 05C82.

Key words and phrases. systemic risk, financial networks, threshold distress contagion, occupation-time distress, directed graphons, nonlinear feedback systems, interacting particle systems, low-rank approximation.

The original idea of this work appeared in the author’s Ph.D. dissertation [21]. The present paper substantially revises and extends that material.

where ℓ is a bounded distress-cost function and $\ell(x) = \mathbf{1}_{\{x \leq 0\}}$ is the indicator case considered below. Section 2.1 motivates the signs and the loss channel in (1) from a balance-sheet cascade with bilateral exposures, partial recovery, interest servicing, and limited liability [18, 13, 4]. The passage to continuous time is a modeling choice rather than a discrete-to-continuous convergence result. The indicator acts on the current state, so the accumulated loss records occupation time and permits recovery; the model is therefore distinct from an absorbing clearing model. As $N \rightarrow \infty$, the interaction closes through the scalar quantity $\beta(t) = \int_0^t \int \ell \, d\nu_s \, ds$. This scalar closure accounts for the tractability of homogeneous mean-field models [12, 28, 22], but it suppresses the distinction between institutions that transmit stress and those that absorb it and cannot represent tiered or hub-dominated architectures.

Many financial infrastructures transmit stress through a limited number of intermediaries, clearing venues, or funding and collateral layers [11, 15, 1, 29, 2]. Such concentration can produce low effective rank even when the bilateral exposure matrix is high dimensional. In the factorized model of section 2, the homogeneous mean-field system is the constant rank-one case. General rank one distinguishes the loading of a stress sender from the sensitivity of a receiver, while rank K introduces one macroscopic feedback coordinate for each transmission channel. Increasing-rank approximations lead to the directed-kernel model of section 4. Thus the rank parameter has both a structural interpretation and a precise role in the dimensional reduction.

The paper is related to three strands of literature. Graphon mean-field theory provides law-of-large-numbers limits, uniform-in-time convergence, concentration bounds, and nonparametric estimation for dense heterogeneous particle systems [9, 7, 8, 3, 14, 10]; financial contagion models have moved beyond homogeneous mean field through heterogeneous default contagion and dynamic clearing [22, 6]; and supervisory work on bank-NBFI and CCP structures identifies architectures in which low-dimensional representations are particularly relevant [1, 29, 2]. Limit theorems for static heterogeneous default cascades appear in Amini, Cao, and Sulem [5]; Erol, Parise, and Teytelboym [20] analyze static threshold contagion on networks sampled from a graphon.

The contribution is a reduction theory for a deterministic dense-network model rather than a new clearing mechanism. Compared with graphon mean-field systems, the analysis is specialized to directed financial exposure kernels and threshold losses and yields a finite-rank-to-kernel approximation result on a common latent space. Compared with static graphon contagion, the state evolves continuously and the threshold loss is accumulated through occupation time rather than an absorbing best-response map. In the discontinuous fixed-rank regime, the main quantitative result is a VC-type law-of-large-numbers estimate for the macroscopic feedback coordinates. The low-rank formulation originates in the author’s dissertation [21]; the present paper adds directed-kernel stability, indicator-loss well-posedness, and numerical diagnostics.

We focus here on the deterministic dense-network setting. Sparse or randomized networks require different tools and are left for future work. Within that scope, the main contributions are as follows.

- (1) *Fixed-rank reduction.* If the exposure matrix has a rank- K factorization, the N -dimensional finite network admits an exact reduction to K macroscopic feedback coordinates. For bounded Lipschitz losses, the corresponding large-population equation is a nonautonomous K -dimensional ODE. Its solution is Wasserstein-stable with respect to the type law, with constants uniform in K under the $1/K$ normalization, and the joint state-factor law has an explicit transport representation. The indicator case is treated separately under threshold regularity.
- (2) *Bounded-Lipschitz graphon bridge.* We formulate the infinite-rank version as a directed graphon contagion equation, prove well-posedness and L^1 -stability for bounded Lipschitz

loss functions, and combine the finite-rank and graphon estimates into a quantitative bridge theorem separating finite-population sampling error from low-rank truncation error.

- (3) *Indicator-loss theory.* For the discontinuous loss $\ell(x) = \mathbf{1}_{\{x \leq 0\}}$, we prove fixed-rank well-posedness under threshold regularity, derive a Vapnik–Chervonenkis (VC)-type finite- N estimate for measurable selections of sampled solutions, and extend the graphon-level theory to factorized kernels and to non-factorized piecewise-smooth kernel–profile pairs. The sampled estimate is conditional on the existence of a measurable absolutely continuous solution of the atomic system, and the graphon bridge assumes uniform transversality along the entire approximation family.
- (4) *Financial interpretation and numerical diagnostics.* We interpret the factors through core–periphery payment systems, multi-CCP clearing markets, and multiplex bank–NBFI networks. The numerical experiments verify the exact finite-rank reduction, quantify bounded-Lipschitz kernel truncation, and examine directed imbalance, layered amplification, and indicator approximations within transverse families. Section 6.5 constructs factor loadings from disclosed EBA sovereign exposures and evaluates the resulting a priori sensitivity bounds. The resampling experiment on the 117-bank empirical population is consistent with the predicted $N^{-1/2}$ scale. In the indicator regime, the deterministic limit is interpreted as a large-population benchmark rather than a point prediction for a single sampled network.

The paper is organized as follows. Sections 2 and 3 introduce the finite-rank model, prove its Wasserstein stability and transport representation, and treat the indicator loss at fixed rank. Section 4 develops the directed-kernel formulation, first for bounded Lipschitz losses and then for indicator losses under factorization or transversality assumptions. Section 5 discusses financial-network interpretations. Section 6 contains the numerical experiments and the sovereign-overlap illustration. Section 7 concludes. Detailed proofs are collected in [section A](#).

Throughout the paper, the contagion term enters with a minus sign: if a counterparty is in distress and exposures are nonnegative, then the marked-to-market cash position of the receiving institution weakly decreases.

2. DYNAMIC CONTAGION MODEL AND LOW-RANK FACTORIZATION

2.1. From default cascades to a dynamic contagion benchmark. This subsection motivates the sign conventions in (6) by embedding a one-period default cascade in a continuous-time debt-service model. It is not a convergence theorem from the discrete cascade to the occupation-time dynamics. In the bilateral interpretation, $e_{ij} \geq 0$ is the raw exposure of institution i to institution j , namely the amount owed by j to i , and $e_{ii} = 0$. Bilateral positions are not netted, so e_{ij} and e_{ji} are recorded separately. The interbank assets and liabilities of i are $\sum_j e_{ij}$ and $\sum_j e_{ji}$, respectively, and x_i denotes net wealth outside the interbank book. The zero-diagonal convention is used only in this bilateral interpretation; the transmission model introduced in [section 2.2](#) does not require it.

One clearing date: the static cascade. Consider first a single clearing date. Let $\theta_0 \in [0, 1)$ be the recovery rate and $\theta := 1 - \theta_0$ the loss rate: a defaulted institution j pays its creditor i only $\theta_0 e_{ij}$, while the obligations owed by i remain due in full—limited liability protects the defaulter, not its counterparties. Writing $\mathbb{D}_0 := \{i : x_i \leq 0\}$ for the initially insolvent set, the losses caused by $\mathbb{D}_0$ can push further institutions below the threshold, and the resolution is the cascade

$$(2) \quad X_i^{(k+1)} := x_i + \sum_{j=1}^N (e_{ij} - e_{ji}) - \theta \sum_{j=1}^N e_{ij} \mathbf{1}_{\mathbb{D}_k}(j) \quad \text{for } i \notin \mathbb{D}_0, \quad \mathbb{D}_{k+1} := \{i : X_i^{(k+1)} \leq 0\},$$

with $X_i^{(k+1)} := x_i$ for $i \in \mathbb{D}_0$: initially insolvent institutions are already in resolution and their books are not updated, whereas an institution that defaults *during* the cascade continues to mark losses—only the sign of its buffer propagates. The default sets increase, the buffers decrease from the first round on, and since every round that changes the default set enlarges it by at least one institution, the cascade stabilizes after at most N rounds. Its terminal outcome is the *maximal* solution of the fixed-point system

$$(3) \quad X_i = x_i + \mathbf{1}_{\{x_i > 0\}} \left[\sum_{j=1}^N (e_{ij} - e_{ji}) - \theta \sum_{j=1}^N e_{ij} \mathbf{1}_{\{X_j \leq 0\}} \right], \quad 1 \leq i \leq N;$$

the short induction argument is recorded in [proposition A.1](#). This is the one-shot picture of the static literature [18, 13]: a single date, an instantaneous resolution, and a terminal default set. The static theory yields resilience criteria for the terminal cascade, including asymptotic default fractions on large random networks [4]. Because the formulation is static, it serves here only as motivation for the continuous-time benchmark introduced next.

From one date to a horizon. Now place the same balance sheets on a horizon $[0, T]$ and let $r > 0$ be the interest rate. Suppose for simplicity that no interbank loan matures before T , so that interbank cash flows over $[0, T]$ consist only of debt service. Over a short window $[t, t + \Delta t]$, a solvent counterparty j services the exposure e_{ij} in full, whereas a counterparty currently below the threshold services at the recovery haircut; the resulting interest income of i on that position is

$$r e_{ij} \mathbf{1}_{\{X_j > 0\}} + \theta_0 r e_{ij} \mathbf{1}_{\{X_j \leq 0\}} = r e_{ij} - \theta r e_{ij} \mathbf{1}_{\{X_j \leq 0\}},$$

per unit time. Thus, while j remains below the threshold, this position reduces the buffer of institution i at rate $\theta r e_{ij}$. The obligations of i itself, $r e_{ji}$ per unit time, remain due regardless of the state of j , and the non-interbank book drifts at an idiosyncratic rate μ_i . The window update is therefore

$$(4) \quad X_{t+\Delta t}^i = X_t^i + \Delta t \left[\mu_i + r \sum_{j=1}^N (e_{ij} - e_{ji}) \right] - \theta r \Delta t \sum_{j=1}^N e_{ij} \mathbf{1}_{\{X_{t+\Delta t}^j \leq 0\}},$$

in which the indicator is contemporaneous: the cascade is assumed to settle within each window, and (4) is evaluated at its maximal solution in the sense of (3). Equation (4) is an Euler-type accounting relation; its formal continuous-time analogue is the stopped system

$$(5) \quad X_t^i = x_i + \int_0^{\tau_i \wedge t} \left[\mu_i + r \sum_{j=1}^N (e_{ij} - e_{ji}) - \theta r \sum_{j=1}^N e_{ij} \mathbf{1}_{\{X_s^j \leq 0\}} \right] ds, \quad \tau_i := \inf\{t \geq 0 : X_t^i \leq 0\} \wedge T,$$

where the stopping time τ_i freezes a defaulted institution, as in the static model. The system (5) is well posed. Between successive default times the dynamics are affine, and an event-driven construction over at most N default times yields the unique solution; see [proposition A.2](#). The corresponding aggregate statistic is the instantaneous threshold-distress fraction $\frac{1}{N} \sum_{i=1}^N \mathbf{1}_{\{X_t^i \leq 0\}}$, whose large- N behavior is studied below. Two model extensions are immediate. The outside book may carry idiosyncratic noise—for example, multiplicative dynamics $\sigma X_t^i dW_t^i$ or a general diffusion coefficient $\sigma(t, X_t^i)$ —and the exposures may vary over time as loans mature or are rolled over. To isolate the contagion channel, the present paper studies the deterministic skeleton, with $\sigma \equiv 0$ and a static exposure matrix, and returns to stochastic and time-varying extensions in [section 7](#).

Three modeling choices. The model used in the remainder of the paper is obtained from (5) by the following three modifications. (i) *Dense scaling.* Write $e_{ij} = e_{ij}^N/N$, with e_{ij}^N uniformly bounded, so that each institution has an order-one aggregate exposure. (ii) *General loss term.*

Set $\mu_i \equiv \mu$ and absorb the scalar coefficient θr into a bounded function $\ell : \mathbb{R} \rightarrow [0, \ell_*]$. The choice $\ell = \mathbf{1}_{\{x \leq 0\}}$ corresponds to unit loss intensity; an independent intensity parameter is recovered by replacing ℓ with $\kappa \ell$. Smooth choices of ℓ describe gradual mark-downs or rating-dependent haircuts. (iii) *Occupation-time dynamics*. Remove the stopping time τ_i . An institution then transmits losses while its state is below zero and may subsequently recover. The resulting paths need not be monotone, which is why the indicator analysis in Sections 3 and 4 requires threshold regularity. Under these choices, the bilateral model is a special case of (6); for $e_{ij}^N \equiv 1$ and $r = 0$, the latter reduces to (1).

2.2. The finite network. Consider N financial institutions with states $(X_t^{i,N})_{1 \leq i \leq N}$, where $X_t^{i,N}$ denotes the cash position, reserve, or marked-to-market buffer of bank i at time t . Let $e_{ij}^N \geq 0$ be a dense-normalized transmission weight from institution j to institution i , and let $\ell : \mathbb{R} \rightarrow [0, \ell_*]$ be a bounded loss function. Under the three modeling choices above, the deterministic contagion dynamics are

$$(6) \quad X_t^{i,N} = x_i^N + \mu t + \frac{rt}{N} \sum_{j=1}^N (e_{ij}^N - e_{ji}^N) - \frac{1}{N} \sum_{j=1}^N e_{ij}^N \int_0^t \ell(X_s^{j,N}) ds,$$

for $1 \leq i \leq N$. From this point onward, e_{ij}^N is a nonnegative transmission weight, and no zero-diagonal condition is imposed. In a bilateral-liability model one may retain $e_{ii}^N = 0$; factor or common-exposure representations may have $e_{ii}^N > 0$, in which case the diagonal represents an institution's own contribution to and exposure through the common channel. Since $e_{ii}^N - e_{ii}^N = 0$, the diagonal does not enter the imbalance term, although it may contribute to the loss term. None of the reduction or stability results below uses a zero-diagonal assumption. The drift $\mu \in \mathbb{R}$ is a common exogenous trend, while the term proportional to $r \in \mathbb{R}$ records the net imbalance between outgoing and incoming exposures. The indicator case is

$$\ell(x) = \mathbf{1}_{\{x \leq 0\}}.$$

More regular choices of ℓ can represent smoothed distress costs or mark-to-market losses prior to formal default.

The indicator convention in this paper is instantaneous rather than absorbing. Thus $\mathbf{1}_{\{X_t^{j,N} \leq 0\}}$ records whether institution j is below the threshold at time t , and the accumulated quantity $\int_0^t \mathbf{1}_{\{X_s^{j,N} \leq 0\}} ds$ is an occupation-time distress exposure, not the event that the institution has ever crossed the default boundary. If the net deterministic drift is non-positive along the relevant paths, this convention is close to the usual absorbing default interpretation; if strong positive drift or recovery effects are present, a bank may leave the distress region. An absorbing-default variant could be modeled by replacing the instantaneous indicator by $\mathbf{1}_{\{\inf_{0 \leq s \leq t} X_s^{j,N} \leq 0\}}$, but the present paper studies the occupation-time model because its cumulative loss channel admits a tractable feedback representation. Throughout the numerical section, the phrase “hard default fraction” denotes the instantaneous fraction of types with $X_t \leq 0$. In the graphon indicator formulation below, the cumulative occupation-time profile is denoted by

$$H_t(u) := \int_0^t \mathbf{1}_{\{X_s(u) \leq 0\}} ds,$$

and is introduced formally in (30).

2.3. Rank- K exposure matrices. The phrase “rank K ” below refers to a finite product representation of the exposure matrix, whereas the resulting limiting dynamics are finite-dimensional because the contagion feedback closes through K macroscopic coordinates. These two meanings

agree in the exact factorized case but should be kept conceptually separate when we later discuss approximate low-rank kernels.

Fix $K \in \mathbb{N}$. We assume that the exposure matrix admits the factorization

$$(7) \quad e_{ij}^N = \frac{1}{K} \sum_{k=1}^K a_{i,k}^N b_{j,k}^N,$$

where $a_{i,k}^N$ measures the exposure sensitivity of bank i to factor k and $b_{j,k}^N$ measures the contribution of bank j to losses transmitted through factor k . In the nonnegative-exposure interpretation one usually takes the factors nonnegative, or otherwise verifies that the resulting sum in (7) is nonnegative. The analytical estimates below only require uniform boundedness, so the same formulas also cover signed factor representations when they are used as analytic approximations. Rank one gives a generalized mean-field model, while finite K represents several transmission channels. The following example records the homogeneous, rank-one, and rank-two special cases.

Example 2.1 (From mean field to rank two). *(i) Homogeneous mean field.* Take $K = 1$ and $a_{i,1} = b_{j,1} = 1$ for all i, j . Then $e_{ij}^N \equiv 1$ and (6) with $r = 0$ is exactly the introductory model (1): every institution feels the same aggregate distress $\frac{1}{N} \sum_j \ell(X^{j,N})$, and the large-population dynamics close through one scalar feedback.

(ii) Generalized mean field. Keep $K = 1$ but allow general loadings, $e_{ij}^N = a_{i,1}^N b_{j,1}^N$. There is still a single transmission channel, but it is now weighted: $b_{j,1}^N$ measures how much stress institution j sends into the channel, while $a_{i,1}^N$ measures how strongly institution i depends on it. For example, in a market intermediated by a single clearing hub, b measures the contribution of an institution to hub-level stress and a measures its sensitivity to losses allocated through the hub. The network is summarized by one scalar function of time, but senders and receivers are no longer exchangeable.

(iii) Rank two: core-periphery. Split the population into a core (mass π_c) and a periphery (mass $1 - \pi_c$) and take $K = 2$ with

$$a_{i,1} = \gamma_1 \mathbf{1}_{\{i \in \text{core}\}}, \quad b_{j,1} = \delta_1 \mathbf{1}_{\{j \in \text{periph}\}}, \quad a_{i,2} = \gamma_2 \mathbf{1}_{\{i \in \text{periph}\}}, \quad b_{j,2} = \delta_2 \mathbf{1}_{\{j \in \text{core}\}}.$$

Then $e_{ij}^N = \frac{1}{2}(a_{i,1}b_{j,1} + a_{i,2}b_{j,2})$ is nonzero only *across* the two tiers: factor 1 is the collection channel through which peripheral distress hits the core, and factor 2 is the redistribution channel through which core distress hits the periphery. The contagion state is driven by exactly two macroscopic loss coordinates, β_1 (losses collected from the periphery) and β_2 (losses redistributed by the core), matching the rank-two row of [table 2](#); [section 6.2](#) quantifies this specification numerically.

(iv) Toward the graphon. Viewed on the continuum type space of [section 4](#), cases (i)–(iii) correspond, respectively, to the constant kernel $W \equiv 1$, the product kernel $W(u, v) = a(u)b(v)$, and a 2×2 block kernel. General bounded kernels, treated in [section 4](#), arise by letting the number of channels grow, and finite-rank models return there as quantifiable approximations of such kernels.

Introduce the type vector

$$z_i^N = (x_i^N, a_{i,1}^N, \dots, a_{i,K}^N, b_{i,1}^N, \dots, b_{i,K}^N) \in \mathcal{Z} := \mathbb{R}^{2K+1},$$

and the empirical type law

$$\mu_0^N := \frac{1}{N} \sum_{i=1}^N \delta_{z_i^N}.$$

We write a generic point of $\mathcal{Z}$ as $z = (x, a, b)$ with $a = (a_1, \dots, a_K)$ and $b = (b_1, \dots, b_K)$. The norm on $\mathcal{Z}$ is the ℓ^1 norm,

$$|z| := |x| + \sum_{k=1}^K |a_k| + \sum_{k=1}^K |b_k|.$$

2.4. The rank- K feedback system: one construction, two instances. The finite network and its large-population limit are generated by one and the same construction, applied to two different type measures. To avoid stating the model twice, we define it once for a generic measure. Fix a probability measure $\varrho \in \mathcal{P}_1(\mathcal{Z})$ and define the means

$$\bar{a}_k^\varrho := \int a_k \varrho(dz), \quad \bar{b}_k^\varrho := \int b_k \varrho(dz),$$

and the imbalance coefficient

$$(8) \quad \Lambda^\varrho(z) := \frac{r}{K} \sum_{k=1}^K (a_k \bar{b}_k^\varrho - b_k \bar{a}_k^\varrho), \quad z = (x, a, b) \in \mathcal{Z}.$$

Given a deterministic vector $\beta = (\beta_1, \dots, \beta_K)$ of feedback variables, the associated state map is

$$(9) \quad X_t^\varrho(z) = x + \mu t + t \Lambda^\varrho(z) - \frac{1}{K} \sum_{k=1}^K a_k \beta_k(t),$$

and the system is closed by requiring

$$(10) \quad \beta_k(t) = \int_0^t m_k(s) ds, \quad m_k(t) = \int b_k \ell(X_t^\varrho(z)) \varrho(dz).$$

Exactly two instances of (8)–(10) are used in this paper. The *empirical instance* takes $\varrho = \mu_0^N$; we then write $\bar{a}_k^N, \bar{b}_k^N, \Lambda^N, X_t^N, \beta_k^N, m_k^N$ for the corresponding objects. The *limiting instance* takes $\varrho = \mu_0$, a fixed probability measure on $\mathcal{Z}$ that we call the large-population type distribution; we then drop the superscript and write $\bar{a}_k, \bar{b}_k, \Lambda, X_t, \beta_k, m_k$. The empirical instance is not a new model but an exact rewriting of the finite network:

Lemma 2.2 (Exact reformulation of the finite network). *Let ℓ be bounded and measurable and take $\varrho = \mu_0^N$. If β^N solves the closed system (9)–(10), then $X_t^{i,N} := X_t^N(z_i^N)$, $1 \leq i \leq N$, solves the finite-network equation (6). Conversely, if $(X^{i,N})_{1 \leq i \leq N}$ solves (6), then the vector β^N defined from these paths by (10) satisfies the closed system, and $X_t^{i,N} = X_t^N(z_i^N)$.*

Proof. Integrals against μ_0^N are averages over the atoms $z_1^N, \dots, z_N^N$. Substituting the factorization (7) into the contagion term of (6) gives

$$\frac{1}{N} \sum_{j=1}^N e_{ij}^N \int_0^t \ell(X_s^{j,N}) ds = \frac{1}{K} \sum_{k=1}^K a_{i,k}^N \left(\frac{1}{N} \sum_{j=1}^N b_{j,k}^N \int_0^t \ell(X_s^{j,N}) ds \right) = \frac{1}{K} \sum_{k=1}^K a_{i,k}^N \beta_k^N(t),$$

where the last equality holds once $X_s^{j,N} = X_s^N(z_j^N)$ and β^N is given by (10). Similarly, $\frac{1}{N} \sum_j e_{ij}^N = \frac{1}{K} \sum_k a_{i,k}^N \bar{b}_k^N$ and $\frac{1}{N} \sum_j e_{ji}^N = \frac{1}{K} \sum_k b_{i,k}^N \bar{a}_k^N$, so the imbalance term of (6) equals $t \Lambda^N(z_i^N)$. Hence (6) for the family $(X^{i,N})_i$ is identical, atom by atom, to (9)–(10) evaluated at $z = z_i^N$; reading this identity in the two directions proves both claims. $\square$

By lemma 2.2, equations (8)–(10) are an exact representation of the finite network. They replace the N coupled state paths by the K -dimensional feedback path β ; each institution's state is then recovered from (9). The finite system and the limiting system have the same form and differ only through the driving type measures μ_0^N and μ_0 . Consequently, the dynamical dimension is reduced from N to K , convergence can be studied through continuity with respect

to the type law, and K is simultaneously the number of factors in (7) and the dimension of the macroscopic feedback.

The limiting instance defines the candidate large-population model studied in section 3. In contrast with classical homogeneous mean field, the law of X_t is not closed by a single scalar statistic; instead, the number of macroscopic coordinates equals the effective rank of the exposure structure.

3. THE FINITE-RANK GENERALIZED MEAN-FIELD LIMIT

Throughout this section we fix $K \in \mathbb{N}$ and assume that the static exposure factors are uniformly bounded. The initial state coordinate is allowed to be unbounded unless a stronger hypothesis is imposed explicitly later.

Assumption 3.1. There exists $M > 0$ such that for every N and every $1 \leq k \leq K$,

$$|a_k| \leq M, \quad |b_k| \leq M$$

for μ_0^N -almost every $z = (x, a, b)$ and for μ_0 -almost every $z = (x, a, b)$.

Remark 3.2. Assumption 3.1 constrains only the static factor loadings (A, B) . It does not require the initial-capital coordinate X_0 to be compactly supported unless such a restriction is stated separately, as in the transport-PDE theorem below. When Wasserstein stability is invoked, the relevant first-moment requirement is supplied by the assumption $\mu_0^N, \mu_0 \in \mathcal{P}_1(\mathcal{Z})$ implicit in $\mathcal{W}_1$ convergence.

3.1. Reduction to a K -dimensional feedback system. The finite-rank model is deterministic once the feedback vector β is known.

Proposition 3.3 (Finite-dimensional feedback equation). *Let $\ell : \mathbb{R} \rightarrow [0, \ell_*]$ be bounded and measurable. The limit system (9)–(10) is equivalent to*

$$(11) \quad \beta_k(t) = \int_0^t F_k(s, \beta(s)) \, ds, \quad 1 \leq k \leq K,$$

where

$$(12) \quad F_k(t, \beta) := \int b_k \ell \left(x + \mu t + t\Lambda(z) - \frac{1}{K} \sum_{j=1}^K a_j \beta_j \right) \mu_0(dz).$$

Every solution is absolutely continuous and satisfies the nonautonomous ODE $\dot{\beta}_k(t) = F_k(t, \beta(t))$ for almost every t , with $\beta_k(0) = 0$. The same representation holds for the finite- N system after replacing μ_0 and Λ by μ_0^N and Λ^N . If ℓ is Lipschitz, then F is globally Lipschitz in β , uniformly on compact time intervals, and the rank- K system is globally well posed.

Proof. The equivalence is immediate from (9)–(10): inserting (9) into (10) yields (11)–(12), and conversely any solution of (11) defines X by (9). If ℓ is Lipschitz with constant L_ℓ , then for any $\beta, \tilde{\beta} \in \mathbb{R}^K$,

$$\begin{aligned} |F_k(t, \beta) - F_k(t, \tilde{\beta})| &\leq \frac{L_\ell}{K} \int |b_k| \sum_{\ell=1}^K |a_\ell| |\beta_\ell - \tilde{\beta}_\ell| \mu_0(dz) \\ &\leq \frac{L_\ell}{K} \cdot M \cdot \sum_{\ell=1}^K M |\beta_\ell - \tilde{\beta}_\ell| \\ &\leq \frac{L_\ell}{K} \cdot M \cdot KM \|\beta - \tilde{\beta}\|_\infty = M^2 L_\ell \|\beta - \tilde{\beta}\|_\infty. \end{aligned}$$

The Carathéodory existence and uniqueness theorem therefore applies. $\square$

3.2. Wasserstein stability for bounded Lipschitz losses. The following theorem establishes continuity of the limiting dynamics with respect to the type distribution.

Theorem 3.4. *Suppose [assumption 3.1](#) holds and $\ell : \mathbb{R} \rightarrow [0, \ell_*]$ is Lipschitz with constant L_ℓ . If $\mu_0^N \rightarrow \mu_0$ in $\mathcal{W}_1$, then for every $T > 0$ there exists $C_T < \infty$, depending only on T, M, r, ℓ_* , and L_ℓ , such that*

$$(13) \quad \sup_{0 \leq t \leq T} \mathcal{W}_1(\nu_t^N, \nu_t) \leq C_T \mathcal{W}_1(\mu_0^N, \mu_0),$$

where

$$\nu_t^N := (\Xi_t^N)_\# \mu_0^N, \quad \nu_t := (\Xi_t)_\# \mu_0,$$

with

$$\Xi_t^N(z) := (X_t^N(z), a, b), \quad \Xi_t(z) := (X_t(z), a, b).$$

In particular, the law of a representative bank together with its static exposure coefficients converges in $\mathcal{W}_1$ uniformly on compact time intervals.

Proof sketch. Fix an optimal coupling π^N of μ_0^N and μ_0 and compare the two instances of the state map (9) pointwise along the coupling. The imbalance and mean terms contribute $O(\mathcal{W}_1(\mu_0^N, \mu_0))$, the feedback difference $|m_k^N - m_k|$ is bounded along the same coupling by $\ell_* \mathcal{W}_1(\mu_0^N, \mu_0) + ML_\ell D_s^N$, and Gronwall's lemma closes the estimate; the pushforward step adds one further $\mathcal{W}_1(\mu_0^N, \mu_0)$ term. The complete proof is given on page 42 in [section A](#). $\square$

Remark 3.5. Under the $1/K$ normalization and the uniform factor bounds, the constant C_T in [theorem 3.4](#) is independent of K . This uniformity is used in the finite-rank-to-kernel approximation theorem.

Remark 3.6. If the type vectors $z_1^N, \dots, z_N^N$ are sampled i.i.d. from μ_0 and have a finite first moment, then $\mathcal{W}_1(\mu_0^N, \mu_0) \rightarrow 0$ almost surely; see, for example, Fournier and Guillin [23]. Hence [theorem 3.4](#) yields almost sure convergence of the generalized mean-field approximation along the sampled sequence.

3.3. Transport PDE for the limiting density. The next result identifies the density of the joint law of the limiting state and the static factors.

Theorem 3.7. *Assume [assumption 3.1](#). Let $T > 0$ and suppose the limiting rank- K system (9)–(10) admits a solution $\beta \in C([0, T]; \mathbb{R}^K)$ on $[0, T]$. This is guaranteed by [proposition 3.3](#) for bounded Lipschitz ℓ and later by [theorem 3.11](#) for $\ell = \mathbf{1}_{\{x \leq 0\}}$ under [assumption 3.9](#). Let ℓ be bounded and measurable, and suppose that μ_0 admits a compactly supported C^1 density Φ on $\mathcal{Z}$.*

Then β is absolutely continuous, with $\beta_k(t) = m_k(t)$ for a.e. t , and for every $t \in [0, T]$ the measure $\nu_t = (\Xi_t)_\# \mu_0$ admits a density $f(t, \cdot)$ on $\mathcal{Z}$, given by

$$(14) \quad f(t, x, a, b) = \Phi(x - \Theta_t(a, b), a, b),$$

where

$$\Theta_t(a, b) := \mu t + \frac{rt}{K} \sum_{k=1}^K (a_k \bar{b}_k - b_k \bar{a}_k) - \frac{1}{K} \sum_{k=1}^K a_k \beta_k(t).$$

Moreover, f solves the transport equation

$$(15) \quad \partial_t f + \partial_x(v_f f) = 0, \quad f(0, x, a, b) = \Phi(x, a, b),$$

in the distributional sense on $(0, T) \times \mathcal{Z}$, with velocity field

$$(16) \quad v_f(t, a, b) = \mu + \frac{r}{K} \sum_{k=1}^K (a_k \bar{b}_k - b_k \bar{a}_k) - \frac{1}{K} \sum_{k=1}^K a_k m_k(t),$$

where

$$(17) \quad m_k(t) = \int_{\mathcal{Z}} \tilde{b}_k \ell(y) f(t, y, \tilde{a}, \tilde{b}) \, dy \, d\tilde{a} \, d\tilde{b}.$$

Equivalently, for every $\varphi \in C_c^1([0, T] \times \mathcal{Z})$,

$$(18) \quad \int_0^T \int_{\mathcal{Z}} f(t, x, a, b) \left(\partial_t \varphi(t, x, a, b) + v_f(t, a, b) \partial_x \varphi(t, x, a, b) \right) dx \, da \, db \, dt \\ + \int_{\mathcal{Z}} \Phi(x, a, b) \varphi(0, x, a, b) \, dx \, da \, db = 0.$$

If, in addition, $t \mapsto m_k(t)$ is continuous for every k , then f is a classical solution of (15).

Proof sketch. For each fixed (a, b) the map $z \mapsto X_t(z)$ is a translation in the x -variable, so the pushforward of Φ has the explicit form (14). Testing against $\varphi \in C_c^1$ and differentiating along the flow yields the weak formulation (18); when $t \mapsto m_k(t)$ is continuous, Θ_t is C^1 in time, (14) can be differentiated pointwise, and the classical equation follows. The complete proof is given on page 43 in section A. $\square$

Remark 3.8. The compact-support and C^1 assumptions in theorem 3.7 are imposed on the density Φ of μ_0 . They justify the explicit density formula (14) and, when the feedback velocities are continuous in time, the classical transport calculation. For merely bounded measurable losses, the feedback $m_k(t)$ need not be continuous, and the transport equation should be read in the weak sense (18). These assumptions are separate from the threshold-regularity assumptions used later in remark 3.10 and theorem 3.11; in particular, the heavy-tailed laws discussed in remarks 3.10 and 3.13 remain admissible for the indicator-loss well-posedness theory even though they need not satisfy the stronger PDE hypothesis of theorem 3.7.

3.4. Discontinuous default losses. The default indicator is the economically natural loss function but destroys global Lipschitz continuity. At the level of the deterministic rank- K limit, the threshold event can still be controlled through a one-dimensional projection argument.

Assumption 3.9. Let $T > 0$ and $C > 0$. There exists $M_\rho(T, C) < \infty$ such that for every $t \in [0, T]$ and every $\beta \in [-C, C]^K$, the scalar random variable

$$\Psi_t(z, \beta) := x + \mu t + t\Lambda(z) - \frac{1}{K} \sum_{k=1}^K a_k \beta_k$$

under μ_0 admits a density bounded by $M_\rho(T, C)$.

Remark 3.10 (Sufficient conditions for assumption 3.9). Assumption 3.9 is a regularity requirement on one-dimensional threshold projections; it is not itself part of the model, and in applications it is verified through the following conditional-density criterion. Suppose that under μ_0 the initial state X_0 admits a regular conditional density $f_{X_0|A,B}(\cdot | a, b)$ given $(A, B) = (a, b)$ such that

$$(19) \quad \operatorname{ess\,sup}_{(a,b) \in [-M, M]^{2K}} \left\| f_{X_0|A,B}(\cdot | a, b) \right\|_{L^\infty(\mathbb{R})} \leq \bar{M} < \infty.$$

Then assumption 3.9 holds with $M_\rho(T, C) = \bar{M}$ for every $T, C > 0$. For fixed t and β , write

$$\Psi_t = X_0 + G_t(A, B; \beta), \quad G_t(a, b; \beta) := \mu t + t\Lambda(a, b) - \frac{1}{K} \sum_{k=1}^K a_k \beta_k.$$

Conditional on $(A, B) = (a, b)$, the variable Ψ_t is a translation of X_0 and therefore has density $y \mapsto f_{X_0|A,B}(y - G_t(a, b; \beta) | a, b)$. Averaging over (A, B) preserves the bound $\bar{M}$. The same

argument remains uniform when the population means entering Λ are replaced by deterministic vectors $(\alpha, \gamma) \in [-M, M]^{2K}$, because these parameters only translate the conditional law. This uniform form is used in [theorem 3.17](#). Condition (19) holds, in particular, in each of the following cases:

- (i) X_0 is independent of (A, B) and has bounded density;
- (ii) X_0 is lognormal, independently of (A, B) , or conditionally lognormal with uniformly bounded conditional densities;
- (iii) X_0 is Pareto-type with lower cutoff $x_m > 0$ and uniformly controlled lower cutoff; for a $\text{Pareto}(x_m, \alpha)$ input the density bound is α/x_m .

If one additionally wants to invoke the $\mathcal{W}_1$ -stability estimate of [theorem 3.4](#), one imposes $\mathbb{E}|X_0| < \infty$; for Pareto tails this corresponds to $\alpha > 1$. Deterministic translations coming from liabilities, reserves, or accounting conventions do not change the density bound.

Theorem 3.11. *Assume [assumptions 3.1](#) and [3.9](#) and let $\ell(x) = \mathbf{1}_{\{x \leq 0\}}$. Then the limiting rank- K system (9)–(10) is well posed.*

Proof sketch. Any solution is confined a priori to the cube $\mathcal{C}_T = [-MT, MT]^K$. On this cube the indicator feedback map is Lipschitz: if two feedback vectors differ by δ in the sup norm, the corresponding indicators can disagree only on a threshold tube of width $M\delta$, whose μ_0 -mass is at most $2M_\rho(T, MT)M\delta$ by [assumption 3.9](#). Carathéodory theory for the feedback ODE then yields global existence and uniqueness. The complete proof is given on page 44 in [section A](#). $\square$

Corollary 3.12 (Indicator-loss density and transport under additional regularity). *Assume the hypotheses of [theorem 3.11](#). If, in addition, μ_0 admits a compactly supported C^1 density Φ on $\mathcal{Z}$, then the density formula (14) holds for the unique solution from [theorem 3.11](#). The corresponding transport equation (15), with velocity field (16) and $m_k(t)$ given by (17) with $\ell(y) = \mathbf{1}_{\{y \leq 0\}}$, holds in the distributional sense. Moreover, in this compactly supported C^1 setting, $t \mapsto m_k(t)$ is continuous for every k , so the transport equation holds classically.*

Proof. The existence and uniqueness of the feedback path follow from [theorem 3.11](#). The density formula and the weak transport equation then follow from [theorem 3.7](#). It remains only to note that m_k is continuous in the present indicator case. By (14),

$$m_k(t) = \int \tilde{b}_k \int_{-\infty}^0 \Phi(y - \Theta_t(\tilde{a}, \tilde{b}), \tilde{a}, \tilde{b}) \, dy \, d\tilde{a} \, d\tilde{b}.$$

The integrand is pointwise continuous in t , bounded by an integrable compact-support envelope, and $\tilde{b}_k$ is bounded by [assumption 3.1](#). Dominated convergence gives continuity of m_k . The classical conclusion in [theorem 3.7](#) therefore applies. $\square$

Remark 3.13. The bounded-density condition is a regularity requirement on one-dimensional threshold projections, not a thin-tail requirement. Heavy-tailed balance-sheet inputs such as lognormal or Pareto-type initial-capital laws are compatible with [assumption 3.9](#) as long as they remain continuously distributed; this is consistent with [Assumption 3.1](#), which constrains the factor loadings but not the state coordinate X_0 . The obstruction is the presence of atoms, near-atoms, or extended flat regions that place macroscopic mass on a threshold layer, rather than tail asymmetry itself.

Remark 3.14 (Regulatory bunching and atoms). A realistic way in which [assumption 3.9](#) can fail is regulatory or accounting bunching. If a substantial set of institutions is reported with capital buffers pinned to the same supervisory minimum, then after the deterministic shift in Ψ_t the projected law may develop an atom, or at least a very sharp near-atom, near the default threshold. Exact atoms at the threshold fall outside the present bounded-density argument

and can destroy the local Lipschitz property of the indicator feedback map used in the proof of [theorem 3.11](#). Near-atoms do not invalidate the theorem, but they enlarge the constant $M_\rho(T, C)$ and therefore worsen the regularization and stability bounds. A full treatment of genuinely atomic threshold mass would require a different formulation, for example with an explicit selection rule or a set-valued feedback at the discontinuity.

Remark 3.15. The projection-density assumption above is a statement about the limiting type law μ_0 , so it is not automatic for arbitrary empirical measures μ_0^N , which are atomic. Still, if $Z_1, \dots, Z_N$ are i.i.d. samples from μ_0 and $Y_i := \Psi_t(Z_i, \beta)$ for fixed (t, β) , then the empirical threshold mass inherits a high-probability small-ball bound from the density of Y_i . Indeed, if [assumption 3.9](#) holds and $\varepsilon, \eta > 0$, the Dvoretzky–Kiefer–Wolfowitz inequality yields

$$\mathbb{P}\left(\sup_{x \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \mathbf{1}_{\{|Y_i - x| \leq \varepsilon\}} > 2M_\rho(T, C)\varepsilon + 2\eta\right) \leq 2e^{-2N\eta^2}.$$

Thus the projected empirical mass in an ε -neighborhood of the threshold is typically of order $\varepsilon + N^{-1/2}$. The quantile-matched constructions in [section 6](#) suppress the sampling term. For the sampled feedback equation, the analysis requires a uniform bound over the entire feedback cube, which is provided by the next lemma.

Lemma 3.16 (Uniform empirical process bound for the fixed-rank indicator classes). *Assume [assumption 3.1](#). Let $Z_1, \dots, Z_N$ be i.i.d. with law μ_0 , write $\mathbb{P}_N := N^{-1} \sum_{i=1}^N \delta_{Z_i}$ and $\mathbb{P}f := \int f d\mu_0$, and fix $T > 0$. Set $\mathcal{C}_T := [-MT, MT]^K$. For $\alpha, \gamma \in [-M, M]^K$, $t \in [0, T]$, $\beta \in \mathcal{C}_T$, and $1 \leq k \leq K$, define*

$$g_{t, \beta, \alpha, \gamma, k}(z) := b_k \mathbf{1}\left\{x + \mu t + \frac{rt}{K} \sum_{j=1}^K (a_j \gamma_j - b_j \alpha_j) - \frac{1}{K} \sum_{j=1}^K a_j \beta_j \leq 0\right\}, \quad z = (x, a, b) \in \mathcal{Z}.$$

Let $\mathcal{G}_{T, K}$ be the class of all such functions. Then $\mathcal{G}_{T, K}$ is a bounded VC-subgraph class with envelope M . Consequently there exists $C = C(M, r, T)$ such that, for every $N \geq 2$,

$$\mathbb{E}\left[\sup_{g \in \mathcal{G}_{T, K}} |(\mathbb{P}_N - \mathbb{P})g|\right] \leq C \sqrt{\frac{K \log N}{N}}.$$

Proof sketch. The threshold maps are affine in $z \in \mathbb{R}^{2K+1}$, so the underlying indicators form a half-space class of VC dimension at most $2K + 2$; multiplying by the bounded coordinate b_k preserves the VC-subgraph property, and the union over k costs only a logarithmic factor in the covering numbers. A standard uniform-entropy maximal inequality then gives the stated rate. The complete proof is given on page 45 in [section A](#). $\square$

Theorem 3.17 (Fixed-rank finite- N convergence estimate for selected indicator solutions). *Assume [assumption 3.1](#) and the conditional-density condition (19) of [remark 3.10](#). Let $Z_1, \dots, Z_N$ be i.i.d. with law μ_0 , define*

$$\mu_0^N := \frac{1}{N} \sum_{i=1}^N \delta_{Z_i}, \quad \bar{a}_k^N := \int a_k d\mu_0^N, \quad \bar{b}_k^N := \int b_k d\mu_0^N,$$

and, for $\beta \in \mathcal{C}_T = [-MT, MT]^K$, set

$$\begin{aligned} \Lambda^N(z) &:= \frac{r}{K} \sum_{k=1}^K (a_k \bar{b}_k^N - b_k \bar{a}_k^N), & \Psi_t^N(z, \beta) &:= x + \mu t + t \Lambda^N(z) - \frac{1}{K} \sum_{k=1}^K a_k \beta_k, \\ F_k^N(t, \beta) &:= \int b_k \mathbf{1}_{\{\Psi_t^N(z, \beta) \leq 0\}} \mu_0^N(dz), & 1 \leq k \leq K. \end{aligned}$$

Let β be the unique solution of the deterministic limit equation from [theorem 3.11](#). Let $S_N : \mathcal{Z}^N \rightarrow C([0, T]; \mathcal{C}_T)$ be Borel measurable and set $\beta^N = S_N(Z_1, \dots, Z_N)$. Assume that, for $\mu_0^{\otimes N}$ -almost every sample, β^N is absolutely continuous and

$$\beta_k^N(t) = \int_0^t F_k^N(s, \beta^N(s)) ds, \quad 1 \leq k \leq K.$$

Then there exists $C_T < \infty$, depending only on T , M , $|r|$, and $\bar{M}$, such that

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} \left\| \beta^N(t) - \beta(t) \right\|_\infty \right] \leq C_T \sqrt{\frac{K \log N}{N}}.$$

If, in addition, $\mathbb{E}|X_0| < \infty$, then

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} \mathcal{W}_1(\nu_t^N, \nu_t) \right] \leq C_T \sqrt{\frac{K \log N}{N}} + C_T \mathbb{E} \mathcal{W}_1(\mu_0^N, \mu_0),$$

where $\nu_t^N := (\Xi_t^N)_{\#} \mu_0^N$, $\nu_t := (\Xi_t)_{\#} \mu_0$, and

$$\begin{aligned} \Xi_t^N(z) &:= \left(x + \mu t + t \Lambda^N(z) - \frac{1}{K} \sum_{k=1}^K a_k \beta_k^N(t), a, b \right), \\ \Xi_t(z) &:= \left(x + \mu t + t \Lambda(z) - \frac{1}{K} \sum_{k=1}^K a_k \beta_k(t), a, b \right). \end{aligned}$$

Proof sketch. The difference between the sampled and limiting feedback fields is decomposed into a uniform empirical-process term and a perturbation of the empirical factor means. The first term is controlled by [lemma 3.16](#); the second follows from the conditional-density bound and Hoeffding's inequality. The Lipschitz estimate for the limiting feedback field then gives the pathwise bound by Gronwall's lemma, uniformly over all solutions satisfying the sampled equation. Borel measurability of S_N ensures that the random path and the error functional are measurable, so the pathwise estimate can be integrated. The state-law estimate follows by coupling the two pushforwards on the sampled atoms. The complete proof is given on page 45 in [section A](#). $\square$

Remark 3.18 (Canonical selections via smoothing). One canonical source of the selected sampled paths in [theorem 3.17](#) is to replace $\mathbf{1}_{\{x \leq 0\}}$ by the Lipschitz regularizations

$$(20) \quad \ell_\varepsilon(x) := \begin{cases} 1, & x \leq 0, \\ 1 - x/\varepsilon, & 0 < x < \varepsilon, \\ 0, & x \geq \varepsilon, \end{cases} \quad \varepsilon > 0,$$

solve the regularized sampled ODE for each $\varepsilon > 0$, and consider the limit as $\varepsilon \downarrow 0$. The positive-side regularization (20) is used in [proposition 4.6](#) and in the stylized experiments of [section 6](#); [section 6.5](#) uses the corresponding negative-side ramp. If, for a deterministic sequence $\varepsilon_n \downarrow 0$, the regularized solutions converge almost surely in $C([0, T]; \mathbb{R}^K)$ to a path satisfying the sampled indicator equation, then the limit is a measurable selection to which [theorem 3.17](#) applies. [Proposition A.3](#) gives pathwise subsequential compactness and identifies indicator solutions under a zero-contact condition; it does not by itself provide a measurable subsequence.

Remark 3.19 (Rate separation and the low-rank advantage). For fixed K , the feedback error in [theorem 3.17](#) is of order $\sqrt{K \log N / N}$. The state-law bound also contains $\mathbb{E} \mathcal{W}_1(\mu_0^N, \mu_0)$, an empirical-measure term on $\mathbb{R}^{2K+1}$ whose rate may deteriorate with dimension for absolutely

continuous laws; see Fournier and Guillin [23]. The two terms quantify different statistical tasks: estimating the K aggregate feedback coordinates and estimating the full type distribution.

Remark 3.20. The estimate in [theorem 3.17](#) is conditional on the existence of a measurable selection and does not establish well-posedness for arbitrary atomic empirical measures. [Proposition A.3](#) identifies regularization limits as indicator solutions when the contact set has zero Lebesgue measure. If the contact set has positive measure, a regularization limit may solve only the associated Filippov inclusion.

4. THE GRAPHON, OR INFINITE-RANK, FORMULATION

Exact finite-rank reduction requires a factorized exposure matrix. General dense heterogeneity is represented below by a bounded directed kernel, with finite-rank kernels used as approximants. The bounded-Lipschitz theory is developed first. The indicator analysis is then divided into factorized kernels, a non-factorized transverse class, and a restricted-family perturbation theorem.

Throughout this paper, a graphon is a bounded directed kernel on the fixed probability space (I, λ) with $I = [0, 1]$. All convergence statements compare representatives on this common latent space in the stated L^1 or L^∞ topology. We do not consider convergence modulo measure-preserving relabelings or dynamics on W -random graphs.

[Table 1](#) summarizes the regimes, the corresponding results, and their standing assumptions.

TABLE 1. Map of the main results and their standing assumptions.

Regime	Results	Key assumptions
Bounded-Lipschitz, fixed rank	theorems 3.4 and 3.7	bounded factors; $\mathcal{W}_1$ convergence of type laws
Bounded-Lipschitz, graphon	theorems 4.2 and 4.4	bounded kernels; L^1 kernel/profile convergence; admissible bounded-factor approximants
Indicator, fixed rank	theorems 3.11 and 3.17	threshold-density condition; selected sampled paths
Indicator, factorized graphon	theorem 4.16	density/transversality of scalar threshold projections
Indicator, non-factorized graphon	theorem 4.19 and proposition 4.23	graphon-level small-ball or piecewise-smooth transversality
Indicator truncation bridge	theorems 4.28 and 4.30 and corollary 4.32	uniformly transverse approximation family; L^∞ kernel/profile control; selected sampled paths

Within the last row of the table, [corollary 4.32](#) adds sampled finite- N convergence to the restricted-family bridge, while [corollary 4.34](#) exhibits a verifiable structure-preserving approximation family to which [theorem 4.30](#) applies.

4.1. Directed graphon contagion equation. Let $I = [0, 1]$ with Lebesgue measure λ . A bounded measurable kernel $W \in L^\infty(I^2)$ represents the continuum exposure from type u to type v ; the first argument is the exposed institution and the second is the counterparty whose distress it absorbs, matching the orientation of e_{ij}^N in (6). In the direct financial-exposure interpretation one takes $W \geq 0$ a.e.; the estimates below are written for bounded signed kernels as well, because signed kernels naturally arise as analytic low-rank or spectral approximations.

The unadorned letter W denotes a bounded measurable directed kernel on I^2 . We write W_1, W_2 for two kernels in stability estimates, $W^{(K)}$ for rank- K approximants, $\Pi_K W$ for block averages, and $W^{\text{blk}}, W^{\text{sm}}$ for the block and smooth components used in the numerical examples. The graphon data are the deterministic pair (W, x_0) with $x_0 \in L^\infty(I)$. The Wasserstein distance is denoted by $\mathcal{W}_1$. Define

$$(21) \quad R_W(u) := \int_I (W(u, v) - W(v, u)) dv,$$

and the nonlinear operator

$$(22) \quad (\Gamma_W \varphi)(u) := \int_I W(u, v) \ell(\varphi(v)) dv.$$

The graphon contagion equation reads

$$(23) \quad X_t(u) = x_0(u) + \mu t + rtR_W(u) - \int_0^t (\Gamma_W X_s)(u) \, ds.$$

This is the directed weighted continuum analogue of (6). As in the finite system, $\lambda(\{u : X_t(u) \leq 0\})$ is an instantaneous hard-threshold fraction, whereas the cumulative profile H_t introduced below is an occupation-time variable.

Remark 4.1. Nothing in (23) requires symmetry. This is essential in finance, where asset and liability channels are directional and the imbalance term R_W records net exposure asymmetry.

Remark (Deterministic data and sampling). Equation (23) is deterministic: W and x_0 are fixed measurable functions, and $u \mapsto X_t(u)$ is a deterministic profile. If $U \sim \text{Unif}(I)$, then $\nu_t = \mathcal{L}(X_t(U))$ is the pushforward of Lebesgue measure and provides the continuum state law. Randomness enters only through sampled finite-network types, as in [remark 3.6](#), [theorem 3.17](#), and [corollary 4.32](#). Random initial capital can be represented by augmenting the latent type so that the initial profile remains a deterministic function of the enlarged label. This convention is used in assumption (ii) of [corollary 4.32](#) and in (19). Dynamics on W -random graphs are not considered here [20].

4.2. Well-posedness and stability.

Theorem 4.2. *Let $\ell : \mathbb{R} \rightarrow [0, \ell_*]$ be bounded and Lipschitz with constant L_ℓ , and let $W \in L^\infty(I^2)$ and $x_0 \in L^\infty(I)$. Then for every $T > 0$ the graphon equation (23) has a unique solution*

$$X \in C([0, T]; L^\infty(I)).$$

Moreover, if (W_1, x_0^1) and (W_2, x_0^2) generate solutions X^1 and X^2 , and if

$$\|W_1\|_{L^\infty} \leq M_W, \quad \|W_2\|_{L^\infty} \leq M_W,$$

then

$$(24) \quad \sup_{0 \leq t \leq T} \|X_t^1 - X_t^2\|_{L^1} \leq C_T \left(\|x_0^1 - x_0^2\|_{L^1} + \|W_1 - W_2\|_{L^1(I^2)} \right),$$

for a constant C_T depending only on $T, |r|, \ell_*, L_\ell$, and M_W .

Proof sketch. Well-posedness follows from a Picard iteration in $C([0, T]; L^\infty(I))$, because Γ_W is Lipschitz from L^1 , hence from L^∞ , into L^∞ with constant $\|W\|_{L^\infty} L_\ell$. For the stability estimate, subtract the two equations, use $\|R_{W_1} - R_{W_2}\|_{L^1} \leq 2\|W_1 - W_2\|_{L^1(I^2)}$ and $\|\Gamma_{W_1}\varphi - \Gamma_{W_2}\psi\|_{L^1} \leq \ell_* \|W_1 - W_2\|_{L^1(I^2)} + M_W L_\ell \|\varphi - \psi\|_{L^1}$, and apply Gronwall's lemma in L^1 . The complete proof is given on page 47 in [section A](#). $\square$

Corollary 4.3. *Let $W_K \in L^\infty(I^2)$ and $x_0^K \in L^\infty(I)$ satisfy*

$$\sup_{K \geq 1} \|W_K\|_{L^\infty} < \infty, \quad W_K \rightarrow W \text{ in } L^1(I^2), \quad x_0^K \rightarrow x_0 \text{ in } L^1(I).$$

If X^K and X solve (23) with data (W_K, x_0^K) and (W, x_0) , respectively, then

$$\sup_{0 \leq t \leq T} \|X_t^K - X_t\|_{L^1(I)} \rightarrow 0 \quad \text{for every } T > 0.$$

Proof. Immediate from [theorem 4.2](#). $\square$

Theorem 4.4 (Low-rank reduction principle for admissible bounded-factor families). *Let $\ell : \mathbb{R} \rightarrow [0, \ell_*]$ be bounded and Lipschitz. Let $W \in L^\infty(I^2)$ and $x_0 \in L^\infty(I)$. For each $K \in \mathbb{N}$, suppose that the following assumptions hold.*

(A1) Bounded-factor approximation. *There are coefficient maps $a_k^{(K)}, b_k^{(K)} \in L^\infty(I)$ such that*

$$W^{(K)}(u, v) = \frac{1}{K} \sum_{k=1}^K a_k^{(K)}(u) b_k^{(K)}(v).$$

(A2) Uniform boundedness. *For some $M < \infty$,*

$$\sup_{K \geq 1} \left(\|x_0^{(K)}\|_{L^\infty} + \max_{1 \leq k \leq K} \|a_k^{(K)}\|_{L^\infty} + \max_{1 \leq k \leq K} \|b_k^{(K)}\|_{L^\infty} + \|W^{(K)}\|_{L^\infty} \right) \leq M.$$

Here M is chosen to dominate both the factor bounds and the kernel bounds; the theorem does not infer the latter from the former.

(A3) Continuum approximation.

$$W^{(K)} \rightarrow W \quad \text{in } L^1(I^2), \quad x_0^{(K)} \rightarrow x_0 \quad \text{in } L^1(I).$$

Let $U \sim \text{Unif}(I)$ and define the rank- K type law

$$\mu_0^{(K)} := (x_0^{(K)}(U), a^{(K)}(U), b^{(K)}(U))_{\#} \lambda,$$

where $a^{(K)} = (a_1^{(K)}, \dots, a_K^{(K)})$ and $b^{(K)} = (b_1^{(K)}, \dots, b_K^{(K)})$. Let $\mu_0^{N,K}$ be empirical measures on $\mathbb{R}^{2K+1}$ satisfying

$$\text{supp}(\mu_0^{N,K}) \subset [-M, M]^{2K+1}.$$

Denote by $\nu_t^{N,K}$ the law of the state component in the finite- N rank- K system associated with $\mu_0^{N,K}$, by

$$\nu_t^K := \mathcal{L}(X_t^K(U))$$

the state law of the rank- K graphon model with kernel $W^{(K)}$ and profile $x_0^{(K)}$, and by

$$\nu_t := \mathcal{L}(X_t(U))$$

the state law of the target graphon model with data (W, x_0) . Then for every $T > 0$ there exists $C_T < \infty$, depending only on T, r, ℓ_, L_ℓ , and M , such that*

$$(25) \quad \sup_{0 \leq t \leq T} \mathcal{W}_1(\nu_t^{N,K}, \nu_t) \leq C_T \mathcal{W}_1(\mu_0^{N,K}, \mu_0^{(K)}) + C_T \left(\|W^{(K)} - W\|_{L^1(I^2)} + \|x_0^{(K)} - x_0\|_{L^1(I)} \right).$$

Proof sketch. Apply [theorem 3.4](#) to the pair $(\mu_0^{N,K}, \mu_0^{(K)})$, which controls the finite-population sampling error; couple the rank- K and target graphon state laws through the common uniform variable U and invoke the stability estimate of [theorem 4.2](#), which controls the truncation error; conclude by the triangle inequality. The complete proof is given on page 48 in [section A](#). $\square$

The hypotheses of the theorem are not free: each approximating kernel must come with an admissible bounded-factor representation with uniform bounds, and generic SVD or block-average truncations do not satisfy this automatically; [remark 4.11](#) spells out what must be checked in practice. In particular, an SVD truncation controls the continuum L^1 error but does not by itself supply an admissible bounded-factor finite- N bridge.

Remark 4.5 (Indicator loss along graphon truncations). [Theorem 4.4](#) is a bounded-Lipschitz bridge theorem; it does *not* by itself cover the discontinuous default indicator. For $\ell(x) = \mathbf{1}_{\{x \leq 0\}}$, an analogous truncation strategy can be analyzed only after one verifies a *uniform* threshold-regularity hypothesis along the chosen rank- K approximation family. This extra hypothesis is not implied by [theorem 4.4](#) and is not automatic for generic low-rank approximations of discontinuous kernels. The next proposition therefore records only a *conditional* uniform $O(\varepsilon)$ smoothing estimate along a prescribed truncation family.

Proposition 4.6 (Conditional uniform smoothing error along rank- K truncations). *For $\varepsilon > 0$ let ℓ_ε be the piecewise-linear regularization of the default indicator defined in (20). In the setting of theorem 4.4, assume in addition that the rank- K type laws $\mu_0^{(K)}$ satisfy assumption 3.9 on $[0, T]$ with a constant $M_\rho(T, MT)$ independent of K . This is an additional hypothesis on the specific approximation family $\{W^{(K)}\}_{K \geq 1}$, not a consequence of theorem 4.4. Let $X^{K, \varepsilon}$ and $X^{K, \text{ind}}$ denote the rank- K graphon solutions associated with $(W^{(K)}, x_0^{(K)})$ and losses ℓ_ε and $\mathbf{1}_{\{x \leq 0\}}$, respectively. Then there exists $C_T < \infty$, independent of K and ε , such that*

$$\sup_{0 \leq t \leq T} \|X_t^{K, \varepsilon} - X_t^{K, \text{ind}}\|_{L^1(I)} \leq C_T \varepsilon, \quad \sup_{0 \leq t \leq T} \mathcal{W}_1(\mathcal{L}(X_t^{K, \varepsilon}(U)), \mathcal{L}(X_t^{K, \text{ind}}(U))) \leq C_T \varepsilon.$$

Proof sketch. Since $|\ell_\varepsilon(y) - \mathbf{1}_{\{y \leq 0\}}| \leq \mathbf{1}_{\{0 < y < \varepsilon\}}$, the uniform density hypothesis makes the smoothed and indicator feedback fields differ by at most $MM_\rho(T, MT)\varepsilon$ on the feedback cube, uniformly in K ; the indicator field is Lipschitz there with a K -uniform constant, so Gronwall's inequality yields an $O(\varepsilon)$ bound on the feedback paths, which the state map transfers to L^1 and to $\mathcal{W}_1$. The complete proof is given on page 48 in section A. $\square$

The proof applies verbatim to the negative-side ramp used in section 6.5, with the tube $\{-\varepsilon < y \leq 0\}$ replacing $\{0 < y < \varepsilon\}$ in the bound above.

Remark 4.7. The assumption in proposition 4.6 is substantive. It requires that the one-dimensional threshold projections of the *approximating* rank- K systems admit density bounds that remain uniform in K . This need not follow from an arbitrary spectral or Fourier truncation scheme. In particular, for discontinuous or block-structured kernels, generic low-rank approximants can develop oscillatory artifacts near the discontinuity interfaces, and such oscillations can flatten the projected threshold map and destroy the uniform small-ball constant. Accordingly, the present paper does *not* claim a general indicator-loss bridge theorem along arbitrary rank- K truncations. The truncation experiment reported in section C.3 (figure 13) stays in the bounded-Lipschitz regime, section 6.4.1 treats the indicator-loss graphon dynamics directly in a verified non-factorized piecewise-smooth/transverse class, and section 6.4.2 illustrates the restricted-family indicator bridge on a structure-preserving approximation path.

Remark 4.8 (SVD truncations and rank selection). If W is Hilbert-Schmidt, a singular-value expansion

$$W(u, v) = \sum_{m=1}^{\infty} \sigma_m \phi_m(u) \psi_m(v)$$

provides the canonical finite-rank truncations

$$W^{(K)}(u, v) = \sum_{m=1}^K \sigma_m \phi_m(u) \psi_m(v).$$

These approximants connect the finite-rank theory of theorem 3.4 appears as the natural truncation of the graphon model in the bounded-Lipschitz regime. In the discontinuous indicator regime they remain only heuristic until the additional uniform threshold regularity from proposition 4.6 and remark 4.7 has been verified.

Theorem 4.4 treats the truncation level K as exogenous. In applications one may choose K from the singular-value decay of an estimated kernel W , for example by a scree plot or held-out cross-validation. If W is Hilbert-Schmidt and $W^{(K)}$ is the singular-value truncation from remark 4.8, then

$$\|W - W^{(K)}\|_{L^1(I^2)} \leq \|W - W^{(K)}\|_{L^2(I^2)} = \left(\sum_{m>K} \sigma_m^2 \right)^{1/2}.$$

Thus singular-value truncation controls the continuum L^1 truncation error that enters [theorem 4.2](#) and [corollary 4.3](#), but not the additional bounded-factor admissibility required by [theorem 4.4](#).

Proposition 4.9 (Constructive block-constant L^1 approximation). *Let $W \in L^\infty(I^2)$. For each $K \in \mathbb{N}$, partition I into the equal subintervals*

$$I_j^{(K)} := \left[\frac{j-1}{K}, \frac{j}{K} \right), \quad 1 \leq j \leq K,$$

and define the block-constant conditional expectation

$$\Pi_K W(u, v) := \sum_{i,j=1}^K \mathbf{1}_{I_i^{(K)}}(u) \mathbf{1}_{I_j^{(K)}}(v) \frac{1}{|I_i^{(K)}| |I_j^{(K)}|} \int_{I_i^{(K)} \times I_j^{(K)}} W(s, t) \, ds \, dt.$$

Then $\|\Pi_K W\|_{L^\infty} \leq \|W\|_{L^\infty}$, the integral operator associated with $\Pi_K W$ has rank at most K , and

$$\|\Pi_K W - W\|_{L^1(I^2)} \rightarrow 0 \quad \text{as } K \rightarrow \infty.$$

If in addition W is piecewise C^1 on finitely many rectangles and the partial derivatives are bounded on each smooth piece, then

$$\|\Pi_K W - W\|_{L^1(I^2)} = O(K^{-1}).$$

Proof sketch. $\Pi_K W$ is the conditional expectation of W with respect to the product partition σ -algebras, so the L^∞ bound is Jensen's inequality and the L^1 convergence is martingale convergence; the range of the associated operator consists of functions constant on the u -partition, whence the rank bound. In the piecewise- C^1 case, cells interior to a smooth rectangle contribute $O(K^{-1})$ by the mean-value theorem, while the cells meeting the finitely many rectangle boundaries occupy total area $O(K^{-1})$. The complete proof is given on page 49 in [section A](#). $\square$

Remark 4.10. [Proposition 4.9](#) supplies a universal constructive continuum approximation family for [theorem 4.2](#) and [corollary 4.3](#). It does *not* by itself provide the uniformly bounded-factor representation required in [theorem 4.4](#), nor the uniform threshold regularity required for indicator truncations. Those checks are family specific. In particular, when W is merely bounded measurable and the singular-value expansion from [remark 4.8](#) is unavailable or inconvenient, [proposition 4.9](#) still yields rank- K kernels of operator rank at most K with vanishing L^1 error; to use such a family inside the finite- N bridge theorem, one must separately provide an admissible bounded-factor representation.

Remark 4.11 (Block and spectral approximations versus the bridge theorem). The finite- N bridge theorem, [theorem 4.4](#), assumes that each approximant already comes with a representation

$$W^{(K)}(u, v) = \frac{1}{K} \sum_{k=1}^K a_k^{(K)}(u) b_k^{(K)}(v)$$

whose factor maps are uniformly bounded in K . Generic block averages and spectral truncations do not automatically satisfy this, so admissibility for [theorem 4.4](#) must still be checked separately. It does hold automatically when W already has a finite bounded-factor representation

$$W(u, v) = \frac{1}{R} \sum_{r=1}^R \alpha_r(u) \gamma_r(v),$$

with fixed R and uniformly bounded factors. Then, along truncation levels K that are multiples of R , one may duplicate each factor pair K/R times to obtain an admissible K -factor family with the same kernel and uniformly bounded coefficients. This is the exact low-rank regime underlying the deterministic benchmarks in [section 6.2](#).

4.3. Factorized kernels and the infinite decomposition model. An infinite decomposition representation can be written as a factorized graphon.

Proposition 4.12. *Let (Θ, ν) be a finite measure space and assume that*

$$W(u, v) = \int_{\Theta} a(u, \theta) b(v, \theta) \nu(d\theta)$$

with bounded measurable coefficients a and b . Then the graphon equation (23) is equivalent to

$$(26) \quad X_t(u) = x_0(u) + \mu t + rt R_W(u) - \int_{\Theta} a(u, \theta) C_t(\theta) \nu(d\theta),$$

where

$$(27) \quad C_t(\theta) = \int_0^t \int_I b(v, \theta) \ell(X_s(v)) dv ds.$$

Proof. Insert the factorization of W into (23) and apply Fubini's theorem:

$$\begin{aligned} \int_0^t \int_I W(u, v) \ell(X_s(v)) dv ds &= \int_0^t \int_I \int_{\Theta} a(u, \theta) b(v, \theta) \ell(X_s(v)) \nu(d\theta) dv ds \\ &= \int_{\Theta} a(u, \theta) \left(\int_0^t \int_I b(v, \theta) \ell(X_s(v)) dv ds \right) \nu(d\theta). \end{aligned}$$

Substitution yields (26)–(27). $\square$

Remark 4.13. The algebraic reformulation in [proposition 4.12](#) extends to σ -finite measure spaces (Θ, ν) whenever Fubini's theorem is justified, for example under

$$\int_{\Theta} \|a(\cdot, \theta)\|_{L^\infty(I)} \|b(\cdot, \theta)\|_{L^\infty(I)} \nu(d\theta) < \infty.$$

For the indicator well-posedness argument, stronger uniform integrability conditions are needed to replace the finite-measure estimates, for instance

$$\operatorname{ess\,sup}_{u \in I} \int_{\Theta} |a(u, \theta)| \nu(d\theta) < \infty, \quad \operatorname{ess\,sup}_{\theta \in \Theta} \int_I |b(v, \theta)| dv < \infty,$$

together with the corresponding integrability of $|a(u, \theta)b(v, \theta)|$ and the threshold regularity needed for the indicator argument. We keep the finite-measure formulation in [theorem 4.16](#) because it gives cleaner constants and avoids additional notation.

Assumption 4.14. Let $T > 0$ and $C > 0$. There exists $M_\rho^\Theta(T, C) < \infty$ such that for every $t \in [0, T]$ and every measurable $c : \Theta \rightarrow \mathbb{R}$ with $\|c\|_{L^\infty(\Theta, \nu)} \leq C$, the scalar random variable

$$(28) \quad \Psi_t(U, c) := x_0(U) + \mu t + rt R_W(U) - \int_{\Theta} a(U, \theta) c(\theta) \nu(d\theta), \quad U \sim \lambda,$$

admits a density bounded by $M_\rho^\Theta(T, C)$.

Proposition 4.15 (Uniform transversality implies [assumption 4.14](#)). *Assume that for every $t \in [0, T]$ and every measurable $c : \Theta \rightarrow \mathbb{R}$ with $\|c\|_{L^\infty(\Theta, \nu)} \leq C$, the profile $u \mapsto \Psi_t(u, c)$ is piecewise C^1 , admits at most J monotone branches, and satisfies $|\partial_u \Psi_t(u, c)| \geq m > 0$ on each branch. Then [assumption 4.14](#) holds with $M_\rho^\Theta(T, C) = J/m$.*

Proof. Fix $t \in [0, T]$ and c with $\|c\|_{L^\infty(\Theta, \nu)} \leq C$. Let $I_1, \dots, I_n$ be a partition of I into monotone C^1 branches of $u \mapsto \Psi_t(u, c)$, with $n \leq J$. On each branch the restriction $g_j := \Psi_t(\cdot, c)|_{I_j}$ is

strictly monotone, so by the one-dimensional change-of-variables formula its pushforward of Lebesgue measure has density

$$\rho_j(y) = \mathbf{1}_{g_j(I_j)}(y) \frac{1}{|g_j'(g_j^{-1}(y))|} \leq \frac{1}{m}.$$

The law of $\Psi_t(U, c)$ for $U \sim \lambda$ is the sum of these branchwise pushforwards, hence it has density

$$\rho(y) = \sum_{j=1}^n \rho_j(y) \leq \frac{n}{m} \leq \frac{J}{m}.$$

Therefore [assumption 4.14](#) holds with $M_\rho^\Theta(T, C) = J/m$. In particular, smooth block, core-periphery, and trigonometric specifications fall into this regime as long as they stay uniformly transverse to the threshold layer. $\square$

Theorem 4.16. *Assume the factorized representation of [proposition 4.12](#), let $\ell(x) = \mathbf{1}_{\{x \leq 0\}}$, set*

$$B_* := \|b\|_{L^\infty(I \times \Theta)},$$

and suppose [assumption 4.14](#) holds with the radius $C = TB_$. Then for every $T > 0$ there exists a unique*

$$C \in C([0, T]; L^1(\Theta, \nu)) \cap L^\infty([0, T] \times \Theta)$$

among paths satisfying $|C_t(\theta)| \leq tB_$ for every $t \in [0, T]$ and for ν -a.e. θ , solving the integral feedback equation*

$$(29) \quad C_t(\theta) = \int_0^t F_s(C_s)(\theta) \, ds, \quad F_t(c)(\theta) := \int_I b(v, \theta) \mathbf{1}_{\{\Psi_t(v, c) \leq 0\}} \, dv,$$

for a.e. θ . Consequently the profile X defined by (26) belongs to $C([0, T]; L^\infty(I))$ and is the unique graphon solution of (23) with indicator loss generated by the factorized representation.

Proof sketch. Work on the a priori ball $\{|C_t(\theta)| \leq tB_*\}$ equipped with the $L^1(\Theta, \nu)$ distance. The threshold-differencing argument together with [assumption 4.14](#) makes the feedback vector field Lipschitz there, and a Picard iteration in $C([0, T]; L^1(\Theta, \nu))$, with measurability supplied by the separability of L^1 , produces the unique fixed point; the representation (26) then recovers the graphon profile and transfers the uniqueness. The complete proof is given on page 49 in [section A](#). $\square$

Remark 4.17. [Theorem 4.16](#) already shows that the bounded-density threshold argument is not confined to fixed rank. In the next subsection we push this mechanism further, from factorized kernels to a non-factorized graphon regime. What remains open after that extension is the indicator-loss theory for completely arbitrary bounded measurable kernels without graphon-level threshold regularity. This does not make factorization or low effective rank an artificial corner case. In many financial systems the dominant exposure channels are mediated by a small number of settlement tiers, CCPs, dealer groups, or funding layers, so factorized and approximately low-rank kernels are often the economically natural regimes in which discontinuous default contagion is most interpretable.

4.4. Indicator-loss graphon equation beyond factorization. The factorized theorem above still leaves open whether one can treat indicator losses for non-factorized kernels. The key observation is that the graphon equation can be rewritten in terms of a cumulative default profile, which restores a closed integral feedback structure directly in function space.

For $t \in [0, T]$ and $u \in I$, define the cumulative default profile

$$(30) \quad H_t(u) := \int_0^t \mathbf{1}_{\{X_s(u) \leq 0\}} \, ds.$$

Then the indicator-loss graphon equation (23) is equivalent to

$$(31) \quad X_t(u) = x_0(u) + \mu t + rtR_W(u) - \int_I W(u, v)H_t(v) dv.$$

Introducing the threshold functional

$$(32) \quad \Psi_t(u, h) := x_0(u) + \mu t + rtR_W(u) - \int_I W(u, v)h(v) dv,$$

we see that any solution must satisfy the Banach-space integral feedback equation

$$(33) \quad H_t(u) = \int_0^t \mathbf{1}_{\{\Psi_s(u, H_s) \leq 0\}} ds.$$

This is the non-factorized graphon analogue of the finite-rank threshold-feedback equations from Section 3.4.

Assumption 4.18. Let $T > 0$ and $C > 0$. There exists $M_\rho^W(T, C) < \infty$ such that for every $t \in [0, T]$ and every $h \in L^\infty(I)$ with $\|h\|_{L^\infty} \leq C$, the scalar random variable

$$(34) \quad \Psi_t(U, h) = x_0(U) + \mu t + rtR_W(U) - \int_I W(U, v)h(v) dv, \quad U \sim \lambda,$$

admits a density bounded by $M_\rho^W(T, C)$.

Theorem 4.19. Let $W \in L^\infty(I^2)$ and $x_0 \in L^\infty(I)$, and let $\ell(x) = \mathbf{1}_{\{x \leq 0\}}$. Assume [assumption 4.18](#) holds with $C = T$. Then there exists a unique

$$H \in C([0, T]; L^1(I)) \cap L^\infty([0, T] \times I)$$

such that

$$\begin{aligned} 0 \leq H_t(u) \leq t & \quad \text{for every } t \in [0, T] \text{ and for a.e. } u \in I, \\ \|H_t - H_s\|_{L^1(I)} & \leq |t - s| \quad \text{for all } s, t \in [0, T], \end{aligned}$$

and (33) holds for every $t \in [0, T]$ and almost every $u \in I$. Consequently, the profile

$$(35) \quad X_t(u) = x_0(u) + \mu t + rtR_W(u) - \int_I W(u, v)H_t(v) dv$$

belongs to $C([0, T]; L^\infty(I))$ and is the unique solution of the graphon contagion equation (23) with indicator loss.

Proof sketch. By [assumption 4.18](#), if two admissible cumulative profiles are L^1 -close, the corresponding threshold indicators can differ only on thin level tubes, so $h \mapsto \mathbf{1}_{\{\Psi_t(\cdot, h) \leq 0\}}$ is Lipschitz from the L^∞ -ball of radius T , with the L^1 metric, into $L^1(I)$. A Picard iteration for the integral feedback fixed point (33) on the closed set $\mathcal{B}_T$, with joint measurability supplied by the separability of $L^1(I)$, gives the unique cumulative profile, and (35) recovers the state profile together with its uniqueness. The complete proof is given on page 50 in [section A](#). $\square$

Corollary 4.20 (Fixed-graphon stability with respect to the initial profile). Let $W \in L^\infty(I^2)$ be fixed and let $x_0^1, x_0^2 \in L^\infty(I)$. Suppose that the two data pairs (W, x_0^1) and (W, x_0^2) jointly satisfy [assumption 4.18](#) on $[0, T]$ with a common bound $M_\rho^W(T, T)$; that is, the density bound holds uniformly for both initial profiles and all admissible h . Let H^i and X^i be the corresponding solutions from [theorem 4.19](#). Then there exists $C_T < \infty$, depending only on T , $|r|$, $\|W\|_{L^\infty}$, and $M_\rho^W(T, T)$, such that

$$\sup_{0 \leq t \leq T} \|H_t^1 - H_t^2\|_{L^1(I)} \leq C_T \|x_0^1 - x_0^2\|_{L^\infty(I)},$$

and consequently

$$\sup_{0 \leq t \leq T} \|X_t^1 - X_t^2\|_{L^1(I)} \leq (1 + \|W\|_{L^\infty} C_T) \|x_0^1 - x_0^2\|_{L^\infty(I)}.$$

Proof sketch. The threshold profiles for the two initial conditions differ pointwise by at most $\eta + \|W\|_{L^\infty} \|H_t^1 - H_t^2\|_{L^1}$ with $\eta = \|x_0^1 - x_0^2\|_{L^\infty(I)}$; the joint density bound converts this displacement into an L^1 bound on the indicator difference, and Gronwall's lemma applied to the cumulative profiles yields both estimates. The complete proof is given on page 51 in [section A](#). $\square$

Remark 4.21 (On stability). [Corollary 4.20](#) gives a one-sided perturbative estimate for fixed W and varying initial profiles. The use of L^∞ on the input side (rather than L^1 as in [theorem 4.2](#)) reflects the pointwise nature of threshold projections: controlling $|\Psi_t^1(u, h) - \Psi_t^2(u, h)|$ uniformly in u —which is what the indicator-differencing argument requires—needs pointwise closeness of x_0^1 and x_0^2 , not merely L^1 closeness. What remains unavailable is a full analogue of [\(24\)](#) with simultaneously varying (W, x_0) . Such a theorem would require a two-dataset version of [assumption 4.18](#) that remains uniform under kernel perturbations, and the present proof does not provide that stronger statement.

Remark 4.22 (Degenerate threshold layers). [Assumption 4.18](#) is a sufficient threshold-regularity hypothesis, not an automatic property of general indicator-loss graphon systems. For example, if $W \equiv 0$, $x_0 \equiv 0$, $\mu = 0$, and $r = 0$, then

$$\Psi_t(U, h) \equiv 0$$

for every t and every h , so the law of $\Psi_t(U, h)$ is a Dirac mass rather than a bounded-density law. The corresponding graphon dynamics are nevertheless uniquely defined ($X_t \equiv 0$, $H_t \equiv t$), although the density-based Lipschitz estimate used in [theorem 4.19](#) is unavailable. Thus [assumption 4.18](#) is sufficient but not necessary; a general well-posedness theory without threshold regularity would require a different argument.

Proposition 4.23 (Piecewise-smooth transversality implies [assumption 4.18](#)). *Fix $T > 0$ and $C > 0$. Let $0 = u_0 < u_1 < \dots < u_J = 1$ and set $I_j := (u_{j-1}, u_j)$ for $1 \leq j \leq J$. Assume that x_0 is piecewise C^1 on this partition and that W is piecewise C^1 on each rectangle $I_j \times I_k$. Suppose also that*

$$A_1 := \sup_{u \in \cup_j I_j} \int_I |\partial_1 W(u, v)| dv < \infty, \quad A_2 := \sup_{u \in \cup_j I_j} \int_I |\partial_2 W(v, u)| dv < \infty.$$

Here $\partial_1 W$ and $\partial_2 W$ are taken on the interiors of the rectangles $I_j \times I_k$; the rectangle boundaries have measure zero and do not affect the integral bounds. If, on each branch I_j ,

$$|x'_0(u)| \geq m_0 > CA_1 + |r|T(A_1 + A_2) \quad \text{for all } u \in I_j,$$

then for every $t \in [0, T]$ and every $h \in L^\infty(I)$ with $\|h\|_{L^\infty} \leq C$, the map $u \mapsto \Psi_t(u, h)$ is piecewise C^1 , has at most J monotone branches, and satisfies

$$|\partial_u \Psi_t(u, h)| \geq m := m_0 - CA_1 - |r|T(A_1 + A_2) > 0 \quad \text{for all } u \in \cup_j I_j.$$

Consequently [assumption 4.18](#) holds with $M_\rho^W(T, C) = J/m$.

Proof sketch. On each branch one may differentiate $\Psi_t(\cdot, h)$ under the integral sign; the kernel-sensitivity terms are bounded by $CA_1 + |r|T(A_1 + A_2)$, so the branchwise derivative stays above $m > 0$ and the profile is piecewise monotone with at most J branches. The branchwise change-of-variables argument of [proposition 4.15](#) then gives the density constant J/m . The complete proof is given on page 52 in [section A](#). $\square$

Remark 4.24 (Economic meaning of A_1 and A_2). The constants A_1 and A_2 summarize how strongly the kernel reacts to changes in the lender type u . In financial terms, A_1 measures the aggregate sensitivity of outgoing exposures $W(u, \cdot)$ to the institution's own type, while A_2 measures the corresponding sensitivity of incoming exposures $W(\cdot, u)$. The transversality condition in [proposition 4.23](#) therefore asks that branchwise initial heterogeneity dominate the amount of type mixing created by these kernel sensitivities over the horizon $[0, T]$. Financially, a small margin m means that dense network mixing nearly offsets the separation in initial capital buffers, so many institutions can bunch near the default threshold. As $m \downarrow 0$, the bound J/m deteriorates, reflecting the fragility of hard-threshold dynamics near such critical threshold layers.

Remark 4.25. [Theorem 4.19](#) and [proposition 4.23](#) show that the indicator-loss graphon problem extends beyond factorization to a non-factorized regime. What remains open is the fully general bounded measurable case without graphon-level small-ball or transversality control. In particular, the present obstruction is no longer the absence of factorization itself, but the absence of a mechanism preventing macroscopic threshold mass from accumulating under the nonlinear graphon feedback.

4.5. A restricted-family indicator bridge via level-set transversality. For indicator losses, finite-rank approximation alone does not control the threshold layer uniformly; see [remarks 4.5](#) and [4.7](#). This subsection imposes uniform level-set transversality along the approximation family and derives a deterministic threshold-layer estimate. The resulting stability theorem applies to the corresponding restricted class of graphon approximations.

Definition 4.26 (Uniformly transverse family). Fix $T > 0$. A family $\mathfrak{F} = \{(W^\alpha, x_0^\alpha)\}_{\alpha \in \mathcal{A}} \subset L^\infty(I^2) \times L^\infty(I)$ is called (J, m) -uniformly transverse on $[0, T]$ if there exists a partition $0 = u_0 < u_1 < \dots < u_J = 1$ such that for every $\alpha \in \mathcal{A}$, every $t \in [0, T]$, and every $h \in L^\infty(I)$ with $\|h\|_{L^\infty} \leq T$, the threshold profile

$$\Psi_t^\alpha(u, h) := x_0^\alpha(u) + \mu t + rtR_{W^\alpha}(u) - \int_I W^\alpha(u, v)h(v) dv$$

is piecewise C^1 on the branch partition $\{I_j\}_{j=1}^J$, has at most J monotone branches, and satisfies

$$|\partial_u \Psi_t^\alpha(u, h)| \geq m \quad \text{for all } u \in \bigcup_{j=1}^J I_j.$$

Proposition 4.27 (Geometric level-set control under transversality). *Let $f : I \rightarrow \mathbb{R}$ be piecewise C^1 on a partition with at most J monotone branches, and assume that $|f'(u)| \geq m > 0$ on each branch. Then, for every $\eta \geq 0$,*

$$\lambda(\{u \in I : |f(u)| \leq \eta\}) \leq \frac{2J}{m} \eta.$$

Consequently, if $g : I \rightarrow \mathbb{R}$ is measurable and $\|f - g\|_{L^\infty(I)} \leq \eta$, then

$$\left\| \mathbf{1}_{\{f \leq 0\}} - \mathbf{1}_{\{g \leq 0\}} \right\|_{L^1(I)} \leq \frac{2J}{m} \eta.$$

Proof. On each monotone branch I_j , the set $\{u \in I_j : |f(u)| \leq \eta\}$ is the preimage of the interval $[-\eta, \eta]$ under a C^1 monotone map whose derivative has magnitude at least m . The one-dimensional mean-value theorem therefore gives

$$\lambda(\{u \in I_j : |f(u)| \leq \eta\}) \leq \frac{2\eta}{m}.$$

Summing over at most J branches yields the first claim. For the second, if the two indicators differ at u , then $f(u)$ and $g(u)$ have opposite signs, so $|f(u)| \leq |f(u) - g(u)| \leq \eta$. Hence

$$\{u : \mathbf{1}_{\{f \leq 0\}} \neq \mathbf{1}_{\{g \leq 0\}}\} \subset \{u : |f(u)| \leq \eta\},$$

and the first estimate finishes the proof. $\square$

Theorem 4.28 (Indicator graphon systems on uniformly transverse families). *Let*

$$\mathfrak{F} = \{(W^\alpha, x_0^\alpha)\}_{\alpha \in \mathcal{A}}$$

be a (J, m) -uniformly transverse family on $[0, T]$. Set

$$M_W := \sup_{\alpha \in \mathcal{A}} \|W^\alpha\|_{L^\infty(I^2)} < \infty.$$

Then, for every $\alpha \in \mathcal{A}$, the indicator-loss graphon equation (23) with data (W^α, x_0^α) has a unique solution (H^α, X^α) with

$$H^\alpha \in C([0, T]; L^1(I)) \cap L^\infty([0, T] \times I), \quad X^\alpha \in C([0, T]; L^\infty(I)).$$

Moreover, for any $\alpha_1, \alpha_2 \in \mathcal{A}$ there exists $C_T < \infty$, depending only on T, J, m, M_W , and $|r|$, such that

$$(36) \quad \sup_{0 \leq t \leq T} \|H_t^{\alpha_1} - H_t^{\alpha_2}\|_{L^1(I)} \leq C_T \left(\|x_0^{\alpha_1} - x_0^{\alpha_2}\|_{L^\infty(I)} + \|W^{\alpha_1} - W^{\alpha_2}\|_{L^\infty(I^2)} \right),$$

and consequently

$$(37) \quad \sup_{0 \leq t \leq T} \|X_t^{\alpha_1} - X_t^{\alpha_2}\|_{L^1(I)} \leq C_T \left(\|x_0^{\alpha_1} - x_0^{\alpha_2}\|_{L^\infty(I)} + \|W^{\alpha_1} - W^{\alpha_2}\|_{L^\infty(I^2)} \right).$$

Proof sketch. Uniform transversality combined with the level-set estimate of [proposition 4.27](#) makes each indicator map Lipschitz in L^1 with the family-uniform constant $2JM_W/m$, so the Picard argument of [theorem 4.19](#) applies verbatim and yields well-posedness for every $\alpha \in \mathcal{A}$. For two members of the family, the pointwise threshold displacement is at most $\eta_{12} + M_W \|H_t^{\alpha_1} - H_t^{\alpha_2}\|_{L^1}$; [proposition 4.27](#) converts displacement into volume, and Gronwall's lemma gives (36) and then (37). The complete proof is given on page 52 in [section A](#). $\square$

Remark 4.29 (Why the indicator bridge uses L^∞ kernel control). The topology in [theorem 4.28](#) is stronger than the L^1 topology used for bounded Lipschitz losses in [theorem 4.2](#). For a discontinuous indicator, an L^1 -small perturbation of W can be concentrated on a small set of types that lies precisely on the threshold layer, producing an order-one change in the indicator feedback. The L^∞ control in [theorem 4.28](#), together with uniform transversality, ensures that perturbing the kernel and initial profile moves the threshold boundary by a uniformly small amount in the pointwise threshold coordinate. The level-set estimate in [proposition 4.27](#) then converts that boundary displacement into a controlled change of volume.

Theorem 4.30 (Restricted-family indicator bridge theorem). *Let $\{(W^{(K)}, x_0^{(K)})\}_{K \geq 1} \cup \{(W, x_0)\}$ be a (J, m) -uniformly transverse family on $[0, T]$, and let $(H^{K, \text{ind}}, X^{K, \text{ind}})$ and $(H^{\text{ind}}, X^{\text{ind}})$ denote the corresponding indicator-loss graphon solutions. Then there exists $C_T < \infty$, depending only on $T, J, m, \sup_K \|W^{(K)}\|_{L^\infty}, \|W\|_{L^\infty}$, and $|r|$, such that*

$$(38) \quad \sup_{0 \leq t \leq T} \|H_t^{K, \text{ind}} - H_t^{\text{ind}}\|_{L^1(I)} \leq C_T \left(\|x_0^{(K)} - x_0\|_{L^\infty(I)} + \|W^{(K)} - W\|_{L^\infty(I^2)} \right),$$

and

$$(39) \quad \sup_{0 \leq t \leq T} \|X_t^{K, \text{ind}} - X_t^{\text{ind}}\|_{L^1(I)} \leq C_T \left(\|x_0^{(K)} - x_0\|_{L^\infty(I)} + \|W^{(K)} - W\|_{L^\infty(I^2)} \right).$$

In particular, if $U \sim \text{Unif}(I)$, then

$$\sup_{0 \leq t \leq T} \mathcal{W}_1(\mathcal{L}(X_t^{K,\text{ind}}(U)), \mathcal{L}(X_t^{\text{ind}}(U))) \leq C_T \left(\|x_0^{(K)} - x_0\|_{L^\infty(I)} + \|W^{(K)} - W\|_{L^\infty(I^2)} \right).$$

Proof. Apply [theorem 4.28](#) with $\alpha_1 = K$ and $\alpha_2 = \infty$. The Wasserstein estimate follows by coupling both laws through the same $U \sim \text{Unif}(I)$. $\square$

Remark 4.31 (Scope of the restricted-family bridge). [Theorem 4.30](#) is not a general indicator-loss truncation theorem. Uniform transversality of the entire approximation family, including the limiting kernel, is part of the hypothesis. Generic spectral or block approximations of a discontinuous kernel may fail this requirement even when they converge in L^1 or L^2 .

Corollary 4.32 (Sampled finite- N to graphon bridge on admissible uniformly transverse families). Assume the hypotheses of [theorem 4.30](#). In addition, suppose that the following sampling assumptions hold.

(i) Each $W^{(K)}$ admits a bounded-factor representation

$$W^{(K)}(u, v) = \frac{1}{K} \sum_{k=1}^K a_k^{(K)}(u) b_k^{(K)}(v), \quad \sup_{K \geq 1} \max_{1 \leq k \leq K} \left(\|a_k^{(K)}\|_{L^\infty} + \|b_k^{(K)}\|_{L^\infty} \right) < \infty.$$

(ii) If $U \sim \text{Unif}(I)$, then

$$\mu_0^{(K)} = \mathcal{L}(x_0^{(K)}(U), a_1^{(K)}(U), \dots, a_K^{(K)}(U), b_1^{(K)}(U), \dots, b_K^{(K)}(U)).$$

With this parametrization, the deterministic rank- K system driven by $\mu_0^{(K)}$ coincides with the graphon indicator solution associated with $(W^{(K)}, x_0^{(K)})$. Any additional initial-capital coordinate must therefore be included in the latent variable U .

(iii) The fixed-rank sampled indicator estimate of [theorem 3.17](#) holds for the laws $\mu_0^{(K)}$ with constants independent of K . For instance, this is ensured by the conditional-density condition (19) of [remark 3.10](#) with constants uniform in K , together with the uniform factor bounds above.

(iv) Let $\mu_0^{N,K}$ be the empirical measure of N i.i.d. samples from $\mu_0^{(K)}$, and let $\nu_t^{N,K,\text{ind}}$ be the state law associated with a measurable selection satisfying the hypotheses of [theorem 3.17](#). Let ν_t^{ind} denote the state law of the target graphon indicator solution.

Then there exists $C_T < \infty$, independent of N and K , such that

$$\begin{aligned} \mathbb{E} \left[\sup_{0 \leq t \leq T} \mathcal{W}_1 \left(\nu_t^{N,K,\text{ind}}, \nu_t^{\text{ind}} \right) \right] &\leq C_T \sqrt{\frac{K \log N}{N}} + C_T \mathbb{E} \mathcal{W}_1(\mu_0^{N,K}, \mu_0^{(K)}) \\ &\quad + C_T \left(\|x_0^{(K)} - x_0\|_{L^\infty(I)} + \|W^{(K)} - W\|_{L^\infty(I^2)} \right). \end{aligned}$$

Proof. Let $\nu_t^{K,\text{ind}} := \mathcal{L}(X_t^{K,\text{ind}}(U))$ be the deterministic rank- K graphon indicator state law. By assumption, this is the same deterministic rank- K limit law associated with the type distribution $\mu_0^{(K)}$. Projecting the joint-law estimate from [theorem 3.17](#) onto the state coordinate gives

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} \mathcal{W}_1 \left(\nu_t^{N,K,\text{ind}}, \nu_t^{K,\text{ind}} \right) \right] \leq C_T \sqrt{\frac{K \log N}{N}} + C_T \mathbb{E} \mathcal{W}_1(\mu_0^{N,K}, \mu_0^{(K)}),$$

with C_T independent of K by the uniform hypotheses. On the other hand, [theorem 4.30](#) gives

$$\sup_{0 \leq t \leq T} \mathcal{W}_1 \left(\nu_t^{K,\text{ind}}, \nu_t^{\text{ind}} \right) \leq C_T \left(\|x_0^{(K)} - x_0\|_{L^\infty(I)} + \|W^{(K)} - W\|_{L^\infty(I^2)} \right).$$

The claim follows from the triangle inequality. $\square$

Remark 4.33 (Sampling schemes and growth of K). The conditional-density condition is not automatic when the initial profile and the factor loadings are functions of the same latent label. It may fail, for example, when the factor vector identifies the label up to finitely many points or when the initial profile is constant on factor level sets. The following construction shows that the assumptions of [corollary 4.32](#) are compatible.

Fix a partition $I = \bigcup_{j=1}^J I_j$ and set $\delta := \min_j \lambda(I_j) > 0$. Let $A_j, B_j : [0, 1] \rightarrow \mathbb{R}$ be continuous and uniformly bounded. For $u \in I_j$, define $a(u, \theta) = A_j(\theta)$ and $b(u, \theta) = B_j(\theta)$, and let $\theta_k^{(K)} = (k - \frac{1}{2})/K$. Set

$$a_k^{(K)}(u) = a(u, \theta_k^{(K)}), \quad b_k^{(K)}(u) = b(u, \theta_k^{(K)}), \quad W^{(K)}(u, v) = \frac{1}{K} \sum_{k=1}^K a_k^{(K)}(u) b_k^{(K)}(v).$$

Because there are finitely many block pairs, midpoint Riemann sums give uniform convergence to

$$W(u, v) = \int_0^1 a(u, \theta) b(v, \theta) d\theta.$$

Let $x_0^{(K)} = x_0$, where x_0 is piecewise C^1 , strictly monotone on each I_j , and $|x'_0| \geq m_0 > 0$ on every branch. The kernel-dependent terms in the threshold profile are constant on each I_j ; hence the family is (J, m_0) -uniformly transverse. If $U \sim \text{Unif}(I)$ and $\mu_0^{(K)}$ is the pushforward law specified in assumption (ii), then the factor vector is constant on each partition interval. Conditional on a factor value, the law of U is normalized Lebesgue measure on the union of intervals carrying that value. Write $A^{(K)} = (a_1^{(K)}(U), \dots, a_K^{(K)}(U))$ and $B^{(K)} = (b_1^{(K)}(U), \dots, b_K^{(K)}(U))$, and let $\rho_K = \mathcal{L}(A^{(K)}, B^{(K)})$. A branchwise change of variables gives

$$\|(\alpha, \gamma) \mapsto \|f_{X_0|A^{(K)}, B^{(K)}}(\cdot | \alpha, \gamma)\|_{L^\infty(\mathbb{R})}\|_{L^\infty(\rho_K)} \leq \frac{J}{\delta m_0},$$

uniformly in K . Thus the bounded-factor, conditional-density, compatibility, and transversality assumptions hold with constants independent of K .

If $K = K(N)$ grows with N , the bound in [corollary 4.32](#) yields convergence provided

$$\sqrt{\frac{K(N) \log N}{N}} \rightarrow 0, \quad \mathbb{E} \mathcal{W}_1(\mu_0^{N, K(N)}, \mu_0^{(K(N))}) \rightarrow 0,$$

and

$$\|x_0^{(K(N))} - x_0\|_{L^\infty(I)} + \|W^{(K(N))} - W\|_{L^\infty(I^2)} \rightarrow 0.$$

The last condition is a restricted-family truncation requirement and is stronger than the L^1 kernel convergence used in the bounded-Lipschitz bridge.

Corollary 4.34 (A verifiable block-plus-smooth trigonometric family). *Let $0 = u_0 < u_1 < \dots < u_J = 1$ and write*

$$W(u, v) = W^{\text{blk}}(u, v) + W^{\text{sm}}(u, v), \quad W^{\text{blk}}(u, v) = \sum_{i,j=1}^J c_{ij} \mathbf{1}_{I_i}(u) \mathbf{1}_{I_j}(v),$$

where W^{sm} is C^1 on each rectangle $I_i \times I_j$ and admits a factorized trigonometric expansion

$$W^{\text{sm}}(u, v) = \sum_{m=1}^{\infty} \sigma_m \phi_m(u) \psi_m(v)$$

with $\phi_m, \psi_m \in C^1(I)$ and

$$\sum_{m=1}^{\infty} |\sigma_m| \left(\|\phi_m\|_{L^\infty} \|\psi_m\|_{L^\infty} + \|\phi'_m\|_{L^\infty} \|\psi_m\|_{L^\infty} + \|\phi_m\|_{L^\infty} \|\psi'_m\|_{L^\infty} \right) < \infty.$$

Define the partial sums

$$W^{(K)}(u, v) := W^{\text{blk}}(u, v) + \sum_{m=1}^K \sigma_m \phi_m(u) \psi_m(v).$$

If x_0 is piecewise C^1 on the same branch partition and satisfies

$$\inf_{u \in \cup_j I_j} |x'_0(u)| > T A_1^* + |r| T (A_1^* + A_2^*),$$

where

$$A_1^* := \sup_{K \geq 1} \sup_{u \in \cup_j I_j} \int_I |\partial_1 W^{(K)}(u, v)| \, dv, \quad A_2^* := \sup_{K \geq 1} \sup_{u \in \cup_j I_j} \int_I |\partial_2 W^{(K)}(v, u)| \, dv,$$

then $\{(W^{(K)}, x_0)\}_{K \geq 1} \cup \{(W, x_0)\}$ is a (J, m) -uniformly transverse family for some $m > 0$. In particular, [theorem 4.30](#) applies. The factor T in the condition comes from the a priori bound $0 \leq H_t \leq t \leq T$ for the cumulative default profile. The periodic Gaussian example from [sections 6.4.1 and 6.4.2](#) is of this form.

Proof. The series assumptions imply uniform convergence of $W^{(K)}$ to W in $L^\infty(I^2)$ and uniform convergence of the branchwise first derivatives on each smooth rectangle. In particular, the quantities A_1^* and A_2^* are finite and dominate the corresponding derivative-integral bounds for both the limit kernel and all its partial sums. Applying [proposition 4.23](#) with the common partition and these uniform bounds yields the claim. $\square$

5. FINANCIAL NETWORK EXAMPLES AND MODELING INTERPRETATION

[Table 2](#) records several stylized correspondences between financial-network architectures and low-rank kernels. In each case, the effective rank is interpreted as the number of distinct transmission channels represented by the model. The sovereign-overlap construction in [section 6.5](#) applies this interpretation to disclosed EBA holdings, first for six banks and then for the full 117-bank sample.

5.1. Rank one: generalized mean field. When

$$W(u, v) = a(u)b(v),$$

there is only one feedback coordinate,

$$\beta(t) = \int_0^t \int_I b(v) \ell(X_s(v)) \, dv \, ds,$$

and the continuum system reduces to

$$X_t(u) = x_0(u) + \mu t + rt(a(u)\bar{b} - b(u)\bar{a}) - a(u)\beta(t),$$

where $\bar{a} = \int_I a(v) \, dv$ and $\bar{b} = \int_I b(v) \, dv$. The feedback satisfies the scalar nonautonomous ODE

$$\beta'(t) = \int_I b(v) \ell(x_0(v) + \mu t + rt(a(v)\bar{b} - b(v)\bar{a}) - a(v)\beta(t)) \, dv.$$

If $a \equiv b \equiv 1$, then we recover the classical homogeneous mean-field contagion model. If the loadings are concentrated on a small subset of institutions, the kernel provides a rank-one approximation of a hub-dominated system.

TABLE 2. Stylized financial architectures and their low-rank kernels.

Architecture	Kernel or factorization	Effective rank	Interpretation
Homogeneous mean field	$W(u, v) \equiv 1$ or $W(u, v) = a(u)b(v)$ with $a \equiv b \equiv 1$	1	Classical fully connected benchmark; every institution feels the same aggregate distress.
Core-periphery or tiered payment network	$W(u, v) = a_1(u)b_1(v) + a_2(u)b_2(v)$	2	A core of money-center banks intermediates flows between many peripheral institutions [11, 15].
Multi-CCP market	$W(u, v) = \sum_{m=1}^M c_m(u)d_m(v)$	M	Each CCP creates one transmission channel; c_m measures dependence on CCP m , while d_m measures stress reallocation through shared membership [29].
Multiplex bank–NBFI system	$W = \sum_{\ell=1}^L \omega_\ell W_\ell$	layer dependent	Separate layers for repo, derivatives, secured funding, payments, or common-asset channels; each layer may itself be low rank [1, 2].

5.2. Core-periphery and tiered interbank structures. Payment systems and unsecured money markets often exhibit a core–periphery or tiered structure in which a small set of money-center banks intermediates a larger peripheral population [11, 15]. A two-factor specification can represent losses transmitted from the periphery to the core and losses redistributed from the core to peripheral institutions. The resulting kernel has rank at most two, and the dynamics close through two macroscopic feedback coordinates.

5.3. Multiple CCPs. Consider a market with multiple central counterparties (CCPs). Suppose that $c_m(u)$ measures the sensitivity of institution u to losses allocated through CCP m , while $d_m(v)$ measures the contribution of institution v to stress transmitted through that CCP. Then

$$W(u, v) = \sum_{m=1}^M c_m(u)d_m(v)$$

has rank at most M , where M is the number of clearing venues. Thus each CCP supplies one transmission factor. The multiple-CCP framework of Veraart and Aldasoro [29] is of this form. State- or time-dependent factor coefficients could incorporate margin, liquidity, or loss-allocation mechanisms, but such extensions are outside the present deterministic model. Supervisory evidence that NBFIs access central clearing through bank clearers provides an additional motivation for overlap and liquidity channels [2].

5.4. Multiplex exposures and bank–NBFI networks. Financial exposures are often multi-layered. A bank can be connected to the same counterparty through unsecured lending, repo, derivatives, collateral reuse, payment obligations, and common asset holdings. A bank–NBFI

network adds yet another layer through funds, insurers, and other nonbank intermediaries. Recent supervisory work emphasizes that bank–NBFI linkages arise precisely through such multiple channels, including leverage provision, liquidity management, clearing services, market-making, derivatives, and risk transfer [2]. A corresponding specification is a sum of layer-specific kernels,

$$W = \sum_{\ell=1}^L \omega_{\ell} W_{\ell}.$$

If layer ℓ has rank r_{ℓ} , the aggregate rank is at most $\sum_{\ell} r_{\ell}$, with possible reductions when factors are shared across layers. This representation is consistent with the multiplex empirical literature [1]; the rank parameter then records the number of distinct transmission channels represented by the model.

The same factor construction applies beyond banking; insurance, supply-chain, and energy-clearing readings of the loadings are collected in [section C.1](#).

6. NUMERICAL EXPERIMENTS

The numerical experiments address three questions. First, exact factorized networks are used to verify the finite- N reduction without truncation error. Second, a directed kernel of nonfinite rank with a smoothed loss is used to examine the L^1 kernel-stability estimate, directed imbalance through rR_W , and amplification across layers. Third, indicator-loss experiments are restricted to the non-factorized transversality class and to the approximation families covered by [theorem 4.30](#) and [corollary 4.34](#). [Section 6.5](#) contains the sovereign-overlap illustration. The accompanying archive contains the scripts for all figures and tables; Monte Carlo and discretization diagnostics are reported in [section B](#).

For indicator losses, the deterministic limit is a large-population benchmark rather than a point forecast for a sampled network. Under i.i.d. sampling, the terminal threshold fraction has substantial finite- N dispersion; in the rank-one example the reported 95% Monte Carlo interval is $[0.075, 0.532]$; see [section B](#).

6.1. Simulation design. For the exact low-rank examples we use the default loss $\ell(x) = \mathbf{1}_{\{x \leq 0\}}$, solve both the finite- N system (6) and the limiting feedback equation (11) by explicit Euler with step size $\Delta t = 5 \times 10^{-4}$, and specify finitely many bank groups $g = 1, \dots, G$. Conditional on the group, the initial capital is Gaussian, $X_0 \mid g \sim N(m_g, \sigma_g^2)$, while the factor loadings $a_{g,k}$ and $b_{g,k}$ are deterministic. The exact low-rank step size was cross-checked against $\Delta t = 2.5 \times 10^{-4}$; the resulting pathwise differences are reported in [table 9](#). The graphon truncation and non-factorized graphon experiments below continue to use $\Delta t = 2 \times 10^{-3}$, with the corresponding refinements also reported in [table 9](#). Since the first four examples set $r = 0$, the limiting feedback satisfies

$$(40) \quad \dot{\beta}_k(t) = \sum_{g=1}^G p_g b_{g,k} \Phi \left(\frac{-m_g - \mu t + \frac{1}{K} \sum_{\ell=1}^K a_{g,\ell} \beta_{\ell}(t)}{\sigma_g} \right), \quad \beta_k(0) = 0,$$

where Φ denotes the standard normal distribution function.

To isolate approximation error from Monte Carlo noise, the empirical measure μ_0^N is chosen deterministically: group masses are rounded to N and, within each group, the initial conditions are placed at Gaussian quantiles. The resulting sequence satisfies $\mathcal{W}_1(\mu_0^N, \mu_0) \rightarrow 0$ and isolates the deterministic large-population approximation error from sampling dispersion. Auxiliary Monte Carlo experiments with true i.i.d. sampling are reported in [section B.1](#). Because the exposure matrices are factorized, the finite- N dynamics can be computed without forming the full $N \times N$ matrix; the grouped threshold implementation is algebraically equivalent to the original N -bank system. At the discrete level, the hard-indicator Euler update therefore imposes one

TABLE 3. Uniform quantile-matched deterministic benchmark errors for the exact low-rank experiments. The error columns report $\sup_{0 \leq t \leq T} |h_t^N - h_t|$, where h_t^N is the full finite-network default fraction and h_t is the deterministic low-rank prediction. Because these four examples are exactly low rank, the table isolates finite- N deterministic approximation effects under quantile matching in a setting with no truncation error. At $N = 1600$ the hard-default fraction (the instantaneous threshold count $\frac{1}{N} \# \{i : X_t^{i,N} \leq 0\}$) has grid size $1/N = 6.25 \times 10^{-4}$, so errors near this scale should be read against the finite-sample staircase floor. In the Lipschitz reduction theorem, this is the analogue of the first term on the right-hand side of (25).

Example	K	T	$h(T)$ limit	$N = 400$ error	$N = 1600$ error
Rank-one generalized mean field	1	2.0	0.222	0.0047	0.0016
Core-periphery	2	2.5	0.283	0.0087	0.0023
Multiple CCPs with overlap	2	2.3	0.473	0.0156	0.0014
Multiplex bank-NBFI network	2	2.2	0.393	0.0040	0.0006

deterministic threshold convention on each sampled system. This is the numerical counterpart of the selected-solution viewpoint in [theorem 3.17](#) and [remark 3.18](#), not a separate existence theorem for arbitrary atomic empirical data. All scripts used to generate the figures and tables are included with the supplementary numerical files accompanying this manuscript.

The nonfinite-rank kernel experiment isolates truncation from threshold regularization and therefore uses the piecewise-linear loss ℓ_ε from (20) with $\varepsilon = 0.02$, together with the directed kernel

$$W(u, v) = \sum_{m \geq 1} \sigma_m a_m(u) b_m(v), \quad \sigma_m = 0.35 m^{-1.8},$$

where

$$\begin{aligned} a_m(u) &= \max\{0.05, 1 + 0.45 \sin(2\pi m u) + 0.15 \cos(2\pi(m+1)u)\}, \\ b_m(v) &= \max\{0.05, 1 + 0.40 \cos(2\pi m v + 0.6) + 0.12 \sin(2\pi(m+2)v)\}. \end{aligned}$$

The initial profile is

$$x_0(u) = 0.13 + 0.08 \sin(2\pi u) - 0.04 \cos(6\pi u), \quad \mu = -0.006,$$

and the numerical reference uses 30 modes on a uniform 2000-point grid. Rank- K truncations keep only the first K modes, with $K \in \{1, 2, 4, 6, 8, 12, 16, 24\}$. The spatial discretization was further checked on 1200-, 2000-, and 2400-point grids; see [table 9](#).

To visualize directed imbalance, we use a two-group rank-one model with equal group masses, factor loadings

$$a = (2.0, 0.6), \quad b = (0.6, 2.0),$$

group means $(0.50, 0.45)$, common volatility parameter 0.18, and compare $r = 0.05$ against the symmetric benchmark $r = 0$. Finally, for the multiplex heatmap we vary the multiplier ω_2 applied to the second-layer factor $a_2 b_2$ relative to the baseline two-layer specification $\omega_2 = 1$, together with a downward shift Δ_{NBFI} in the NBFI group mean initial buffer. The shift is measured in the same units as the state variable X ; in the baseline parametrization the NBFI mean is 0.28, so $\Delta_{\text{NBFI}} = 0.08$ lowers it to 0.20.




FIGURE 1. Validation of the rank-one and core-periphery examples. Left: the rank-one default fraction converges rapidly to the deterministic limit as N increases. Right: in the core-periphery specification, the group-resolved finite- N trajectories nearly coincide with the limiting core and periphery predictions.

6.2. Exact low-rank validation. In table 3, the uniform error at $N = 1600$ ranges from 6.2×10^{-4} to 2.3×10^{-3} across the four exact low-rank examples. The multiple-CCP case has uniform threshold-fraction error of approximately one tenth of a percentage point. These quantile-matched calculations exclude sampling variability; the i.i.d. results in section B.1 show substantially wider finite-sample dispersion for indicator losses.

In the rank-one example, the finite-network threshold fraction is a staircase function, and its uniform distance from the deterministic limit decreases with N . The terminal limiting fraction is 22.16%; the $N = 1600$ curve differs only at the finite-sample step scale.

The core-periphery panel reports group-specific paths. The periphery crosses the threshold earlier and has terminal limiting fraction 30.0%; the corresponding core fraction is 18.5%. The distinct group paths reflect the two directional transmission factors.

The multiple-CCP and multiplex examples are validated in the same way, with cross-venue transmission through overlapping clearing membership and layer-dependent amplification; the figures and discussion are collected in section C.2.

6.3. Truncation, directedness, and amplification. The graphon truncation experiment—the direct test of the bounded-Lipschitz continuum truncation mechanism of theorem 4.4 on a directed thirty-mode reference kernel—is reported in section C.3; its empirical-scale counterpart is the factor-aligned comparison of section 6.5.3.

The left panel of figure 2 exhibits the directed imbalance mechanism carried by the rR_W term. In the two-group lender/borrower specification, setting $r = 0.05$ lowers the terminal default rate of net lenders from 0.52% to 0.17% while raising the rate of net borrowers from 0.79% to 11.37%. This is a purely directed-network effect: when $r = 0$, the two groups remain close, whereas at $r \neq 0$ the asymmetry in outgoing and incoming exposures materially shifts both timing and incidence of distress.

The right panel of figure 2 gives a phase diagram for the multiplex example. The horizontal axis is the multiplier ω_2 on the second-layer factor relative to the baseline value $\omega_2 = 1$, and the vertical axis is the downward shift Δ_{NBFI} in the NBFI group mean initial buffer, measured in the same units as X . The baseline NBFI mean is 0.28, so the largest plotted shift lowers it to




FIGURE 2. Directedness and structural amplification. Left: a nonzero imbalance parameter r sharply separates net lenders from net borrowers even when the contagion channel is otherwise rank one. Right: a phase diagram for the multiplex example. The horizontal axis is the multiplier ω_2 applied to the baseline second-layer factor ($\omega_2 = 1$ is the calibrated two-layer specification), and the vertical axis is the downward shift Δ_{NBFI} in the NBFI group mean initial buffer, measured in the same units as X .

0.20. Terminal default fractions remain moderate when the second layer is weak but rise sharply when a stronger second-layer interaction is combined with a deterioration in NBFI buffers. The comparison shows that a one-layer approximation can understate amplification generated by the additional funding channel.

The deterministic quantile-matched experiments isolate approximation error from sampling variation. In the indicator case, the i.i.d. Monte Carlo results in [section B.1](#) show substantial finite- N dispersion, so the deterministic limit is interpreted as a large-population benchmark rather than as a concentration result for a single realization. For smoothed losses, the observed kernel-truncation errors are consistent with the L^1 stability estimate.

6.4. Indicator-loss graphon experiments. Two indicator-loss experiments are reported. The first uses a non-factorized piecewise-smooth kernel satisfying [theorem 4.19](#) and [proposition 4.23](#). The second studies a structure-preserving truncation family covered by [theorem 4.30](#) and [corollary 4.34](#).

6.4.1. A non-factorized piecewise-smooth indicator illustration. Consider the following non-factorized example, which satisfies the hypotheses of [theorem 4.19](#) and [proposition 4.23](#). Let $I_1 = [0, \frac{1}{2})$ and $I_2 = [\frac{1}{2}, 1]$, and define the directed kernel

$$W(u, v) = B_{ij} + 0.16 \exp(-10 \sin^2(\pi(u - v))), \quad (u, v) \in I_i \times I_j,$$

with block matrix

$$B = \begin{pmatrix} 0.38 & 0.24 \\ 0.31 & 0.36 \end{pmatrix}.$$

The block part is directed, while the periodic Gaussian term is smooth and non-factorized: its Fourier expansion contains infinitely many nonzero modes, so it cannot be reduced to a finite

sum of products $a_k(u)b_k(v)$. We take $r = 0$, $\mu = -0.06$, $T = 2$, and the piecewise linear initial profile

$$x_0(u) = \begin{cases} -0.18 + 1.95u, & 0 \leq u < \frac{1}{2}, \\ 0.82 + 1.75(u - \frac{1}{2}), & \frac{1}{2} \leq u \leq 1. \end{cases}$$

The jump $x_0(\frac{1}{2}+) - x_0(\frac{1}{2}-) = 0.025$ represents a mild block-specific shift in baseline buffers, while the positive slopes on both branches enforce transverse threshold crossing. More precisely,

$$m_0 = \min\{1.95, 1.75\} = 1.75.$$

Because the block part of W is constant on each rectangle, the constants A_1 and A_2 from [proposition 4.23](#) come entirely from the smooth periodic term. Writing $x = u - v$ and differentiating the smooth term gives

$$\partial_1 W_{\text{sm}}(u, v) = -3.2\pi \sin(\pi x) \cos(\pi x) \exp(-10 \sin^2(\pi x)), \quad x = u - v,$$

so by periodicity

$$A_1 = A_2 = \int_0^1 3.2\pi |\sin(\pi x) \cos(\pi x)| \exp(-10 \sin^2(\pi x)) dx \approx 0.320,$$

where the final value is obtained by direct quadrature of the explicit periodic integral above. In the present experiment $r = 0$, $T = 2$, and the theorem-level ball radius is $C = T = 2$, so the sufficient condition from [proposition 4.23](#) reduces to

$$m_0 > CA_1, \quad \text{i.e.} \quad 1.75 > 2 \times 0.320 \approx 0.640.$$

Thus the numerical example lies strictly inside the verified transversality regime of [proposition 4.23](#). We solve the indicator-loss graphon equation in cumulative-default form [\(33\)](#)–[\(31\)](#) on a uniform 2000-point grid with the same explicit-Euler step size $\Delta t = 2 \times 10^{-3}$ used elsewhere in [Section 6](#), and we compare it with the corresponding smoothed graphon dynamics obtained by replacing ℓ by ℓ_ε for $\varepsilon \in \{0.04, 0.02, 0.01\}$.

TABLE 4. Indicator versus smoothed dynamics for the non-factorized piecewise-smooth graphon illustration. The indicator row is the baseline solution of [theorem 4.19](#); the smoothed rows solve the same graphon equation with ℓ_ε . Errors are computed relative to the indicator solution.

Regime	ε	terminal hard default fraction	$\sup_{0 \leq t \leq T} \ X_t^\varepsilon - \bar{X}_t^{\text{ind}}\ _{L^1}$	$\sup_{0 \leq t \leq T} h_t^\varepsilon - \bar{h}_t^{\text{ind}} $
Indicator	–	0.2195	0	0
Smoothed	0.04	0.2255	9.40×10^{-3}	6.00×10^{-3}
Smoothed	0.02	0.2225	4.70×10^{-3}	3.00×10^{-3}
Smoothed	0.01	0.2210	2.35×10^{-3}	1.50×10^{-3}




FIGURE 3. A non-factorized piecewise-smooth indicator-loss graphon illustration. Left: heatmap of the directed kernel $W(u, v)$ used in Section 6.4. Right: hard default fraction trajectories for the indicator-loss dynamics and for the regularized dynamics with ℓ_ϵ . The smoothed trajectories approach the indicator trajectory as $\epsilon \downarrow 0$.

Figure 3 gives the qualitative picture, and table 4 quantifies it. The indicator and smoothed hard-default trajectories are nearly indistinguishable at plotting scale. Quantitatively, the terminal hard default fraction is 21.95% for the indicator dynamics, while the smoothed runs give 22.55%, 22.25%, and 22.10% for $\epsilon = 0.04, 0.02, 0.01$, respectively. The pathwise state discrepancy decreases from 9.40×10^{-3} to 4.70×10^{-3} and then to 2.35×10^{-3} as ϵ is halved, and the corresponding hard-default-fraction discrepancy decreases from 6.00×10^{-3} to 3.00×10^{-3} and then to 1.50×10^{-3} . Re-running the same experiment on 1200- and 2000-point grids changes the displayed hard-default trajectories by at most 5.0×10^{-4} , so the visual agreement is not a discretization artifact.

This experiment is not a calibration and does not address arbitrary indicator-loss truncations. It illustrates the piecewise-smooth non-factorized regime covered by proposition 4.23 and compares the indicator solution with the regularized dynamics generated by ℓ_ϵ .

The sensitivity of this construction to the transversality margin m is examined in section C.4: shrinking the margin degrades the indicator-smoothing agreement as the theory predicts, while all tested regimes remain inside the verified transversality class.

6.4.2. Indicator bridge along a structure-preserving trigonometric family. The restricted-family indicator bridge theorem is tested on the same kernel. Write the kernel as $W = W^{\text{blk}} + W^{\text{sm}}$, where the directed block part W^{blk} is kept exact and the smooth periodic component $W^{\text{sm}}(u, v) = 0.16 \exp(-10 \sin^2(\pi(u - v)))$ is approximated by its first M Fourier modes. This block-plus-smooth trigonometric family satisfies the hypotheses of corollary 4.34: the factors are uniformly bounded, the branch partition is fixed, and the derivatives of the partial sums are uniformly controlled. We therefore compare the indicator-loss graphon solution of the full kernel with the corresponding solutions for $M \in \{1, 2, 4, 8, 16\}$ on the same 2000-point grid and with the same explicit-Euler step size $\Delta t = 2 \times 10^{-3}$.

TABLE 5. Indicator bridge along a structure-preserving trigonometric approximation family. The block part of the kernel is kept exact; only the smooth periodic component is truncated. Errors are computed relative to the full indicator-loss graphon solution.

Retained modes M	$\ W^{(M)} - W\ _{L^\infty}$	$\sup_{0 \leq t \leq T} \ X_t^{(M)} - \bar{X}_t\ _{L^1}$	$\sup_{0 \leq t \leq T} h_t^{(M)} - h_t $	terminal hard default fraction
1	7.82×10^{-2}	6.26×10^{-3}	3.50×10^{-3}	0.2230
2	4.04×10^{-2}	2.74×10^{-3}	3.00×10^{-3}	0.2225
4	7.13×10^{-3}	2.11×10^{-4}	5.00×10^{-4}	0.2195
8	5.42×10^{-5}	8.67×10^{-7}	5.00×10^{-4}	0.2195
16	5.76×10^{-11}	6.15×10^{-13}	0	0.2195




FIGURE 4. Indicator bridge along a structure-preserving trigonometric family. Left: indicator default-fraction trajectories for the full kernel and for the Fourier-truncated families. Right: kernel L^∞ and state L^1 errors decay rapidly as the retained mode count increases, while the hard-default trajectory error quickly reaches the grid-level thresholding floor.

Figure 4 and table 5 show that the indicator trajectories converge rapidly along this structure-preserving family. The kernel L^∞ error falls from 7.82×10^{-2} at $M = 1$ to 7.13×10^{-3} at $M = 4$ and to 5.42×10^{-5} at $M = 8$, while the corresponding pathwise state error drops from 6.26×10^{-3} to 2.11×10^{-4} and then to 8.67×10^{-7} . The hard-default error is already at the 5×10^{-4} grid-level thresholding floor by $M = 4$ – 8 , and by $M = 16$ the approximation is numerically indistinguishable from the full-kernel solution at the resolution of the experiment. Unlike the bounded-Lipschitz truncation test reported in section C.3 (figure 13), this experiment operates in the discontinuous indicator regime covered by the theory: it stays inside a verifiable transversality-preserving approximation family and therefore illustrates the restricted-family bridge mechanism from theorem 4.30 and corollary 4.34 rather than a purely smoothed surrogate.

6.5. A real-data-driven sovereign-overlap illustration. Our data source is the Autumn-2025 EU-wide transparency exercise of the European Banking Authority [30], which discloses, for 120 banking groups at the highest level of consolidation, sovereign exposures by counterparty

country at reference date 30 June 2025. We extract the direct on-balance-sheet gross carrying amount of non-derivative sovereign assets, summed over all maturities (template item 2520810); the aggregate is EUR 4.01tn. We first work with a six-bank, four-country subsample used to display the construction explicitly, and [section 6.5.3](#) then scales every step of the construction to the full exercise. The six banks are BNP Paribas, Deutsche Bank, UniCredit, Banco Santander, ING Groep, and Intesa Sanpaolo; the four sovereign factors are France, Germany, Italy, and the Netherlands. Table 6 reports the holdings (million EUR) used in the six-bank experiment. The construction below is an empirical sovereign-overlap proxy, not a reconstruction of bilateral interbank liabilities.

Let $s_{i,k}$ denote the holding of bank i in sovereign factor k . We convert these observations into paper-style low-rank coefficients by

$$(41) \quad a_{i,k}^{\text{emp}} = \frac{s_{i,k}}{\sum_{\ell=1}^K s_{i,\ell}}, \quad b_{i,k}^{\text{emp}} = \frac{s_{i,k}}{\frac{1}{N} \sum_{m=1}^N s_{m,k}}.$$

The coefficient $a_{i,k}^{\text{emp}}$ is the within-bank sovereign share, while $b_{i,k}^{\text{emp}}$ measures the contribution of bank i to country- k overlap relative to the cross-bank mean. Figure 5 visualizes the resulting row shares.

We then define a real-data-driven overlap network by

$$(42) \quad e_{ij}^{N,\text{emp}} = \frac{c_E}{K} \sum_{k=1}^K a_{i,k}^{\text{emp}} b_{j,k}^{\text{emp}}, \quad c_E = 0.30.$$

This object is *not* a directly observed bilateral exposure matrix. It is an empirical factor proxy: institutions with larger loadings on the same sovereign factors exert stronger mutual effects through the common balance-sheet channel.

TABLE 6. Sovereign holdings used in the experiment (million EUR).

Bank	France	Germany	Italy	Netherlands
BNP Paribas	54,638	17,862	23,455	608
Deutsche Bank	17,962	14,776	22,414	677
UniCredit	7,121	17,405	45,760	16
Banco Santander	7,898	1,642	17,284	221
ING Groep	7,288	9,514	2,339	4,630
Intesa Sanpaolo	12,810	2,175	42,616	1,108




FIGURE 5. Bank-by-country sovereign shares $a_{i,k}^{\text{emp}}$ from (41). Italian concentration is dominant for UniCredit, Banco Santander, and Intesa Sanpaolo, and Italy is also Deutsche Bank's largest single bucket; France is dominant for BNP Paribas; ING Groep is the most diversified, with Germany, France, and the Netherlands as its main buckets.

6.5.1. *Dynamic contagion experiment.* To isolate the low-rank mechanism, we specialize the finite-network dynamics to

$$(43) \quad X_t^{i,N} = x_i^N - \frac{1}{N} \sum_{j=1}^N e_{ij}^{N,\text{emp}} \int_0^t \ell_\varepsilon(X_s^{j,N}) \, ds, \quad \ell_\varepsilon(x) = \max\{0, \min\{1, -x/\varepsilon\}\},$$

with $\varepsilon = 0.05$, $\mu = r = 0$, and horizon $T = 3$. The ramp of this ℓ_ε sits on the negative side of the threshold, so $\ell_\varepsilon(0) = 0$ and $\ell_\varepsilon \leq \mathbf{1}_{\{x \leq 0\}}$ pointwise; it differs from the positive-side regularization (20) of remark 3.18 only in the side of the threshold that carries the ramp. Both versions are bounded by one and $1/\varepsilon$ -Lipschitz, and only these two properties enter the estimates used in this section. The initial buffers are set by

$$(44) \quad x_i^N = x_* - \sum_{k=1}^K \eta_k a_{i,k}^{\text{emp}}, \quad x_* = 0.40, \quad \eta = (0.35, 0.15, 0.45, 0.05),$$

so that Italian-sovereign concentration is the main source of vulnerability in this calibration.

Because (42) has the exact four-factor representation (7), lemma 2.2 applies without approximation:

$$X_t^{i,N} = x_i^N - \frac{1}{K} \sum_{k=1}^K a_{i,k}^{\text{emp}} \beta_k^N(t), \quad \beta_k^N(t) = c_E \int_0^t \frac{1}{N} \sum_{j=1}^N b_{j,k}^{\text{emp}} \ell_\varepsilon(X_s^{j,N}) \, ds.$$

The direct finite-network solver and the rank-4 reduced solver agree to a sup-norm discrepancy of order 10^{-16} . Under this parameter choice, the Italian mark-down gives Intesa Sanpaolo an initial buffer of -0.010 ; Banco Santander starts with buffer $+0.001$ and subsequently crosses the threshold, while the other four institutions remain solvent. The parameters x_* , c_E , and η define a stress scenario and are not estimated from prices or supervisory models. The accompanying script reports the sensitivity of the classification to x_* and to the interaction intensity. The exercise is therefore a calculation on disclosed holdings, not a calibrated default forecast.

To study truncation, we next replace $e_{ij}^{N,\text{emp}}$ by its rank- r singular-value approximation for $r = 1, 2, 3$ and compare the resulting distress paths against the exact rank-4 network. The leading singular values of the empirical overlap matrix are approximately

$$0.4890, \quad 0.0949, \quad 0.0487, \quad 0.0167,$$

so the first singular direction already captures about 95.4% of the Frobenius energy, rank 2 captures about 98.9%, and rank 3 about 99.9%. All three SVD surrogates are entrywise nonnegative in this small sample; signed entries appear only at the full-sample scale, see [section 6.5.3](#).

TABLE 7. Truncation diagnostics for the empirical overlap network. The last column reports the maximal gap, over $t \in [0, T]$, between the mean smoothed-distress paths of the rank- r and exact networks, i.e., the maximal vertical distance between the curves of Figure 6.

Rank	Relative Frobenius error	Terminal mean distress	Terminal hard-default fraction	Max path gap
1	0.2155	0.1203	0.3333	0.0491
2	0.1027	0.1565	0.3333	0.0129
3	0.0334	0.1565	0.3333	0.0129
4 (exact)	0.0000	0.1694	0.3333	0.0000




FIGURE 6. Low-rank truncation of the empirical six-bank network. (a) Mean smoothed distress $\frac{1}{N} \sum_i \ell_\varepsilon(X_t^{i,N})$ for the exact rank-4 kernel and its rank-1, rank-2, and rank-3 truncations. The rank-2 and rank-3 paths are nearly identical, whereas rank 1 understates the amplification. (b) Terminal buffers for the exact kernel and the rank-2 surrogate. The threshold classification is unchanged; UniCredit is the solvent institution closest to the threshold.

6.5.2. *A priori sensitivity envelopes for a sovereign re-rating scenario.* The stability estimate also yields an a priori sensitivity envelope for a family of sovereign re-rating scenarios. The envelope is computed from the baseline holdings and does not require simulation of each scenario. For $\delta \in [-0.20, 0.20]$, consider the scenario in which the Italian sovereign book of every institution is uniformly re-rated,

$$(45) \quad s_{i,\text{IT}} \mapsto (1 + \delta) s_{i,\text{IT}}, \quad i = 1, \dots, N,$$

and propagated through (41), (42), and (44), so that the kernel and the initial buffers move together: this is a composite $(\Delta x, \Delta e)$ perturbation of the type covered by the stability estimates.

Let $(\tilde{x}, \tilde{e}, \tilde{X})$ denote the perturbed data and solution, and set $\Delta x := \tilde{x} - x$, $\Delta e := \tilde{e} - e$, and $\Delta X_t^i := \tilde{X}_t^i - X_t^i$. Splitting $\tilde{e}_{ij\ell_\varepsilon}(\tilde{X}_s^j) - e_{ij\ell_\varepsilon}(X_s^j)$ into a kernel perturbation and a state perturbation, and applying the Gronwall argument from [theorem 4.2](#) to (43), gives

$$(46) \quad \max_{1 \leq i \leq N} |\Delta X_t^i| \leq \left(\|\Delta x\|_\infty + t \bar{d} \ell_* \right) e^{L_\ell \bar{e} t}, \quad \bar{d} := \max_i \frac{1}{N} \sum_j |\Delta e_{ij}|, \quad \bar{e} := \max_i \frac{1}{N} \sum_j |e_{ij}|,$$

with $\ell_* = 1$ and $L_\ell = 1/\varepsilon$. For the nonnegative baseline kernel, the normalizations in (41) imply that every row mean equals c_E/K . Hence $\bar{e} = c_E/K = 0.075$, and the Gronwall factor at $T = 3$ is $\exp(L_\ell c_E T/K) = e^{4.5} \approx 90$. Across the re-rating family, only $\|\Delta x\|_\infty$ and $\bar{d}$ vary with δ ; both are explicit functions of the holdings.




FIGURE 7. A priori envelope versus observed response to the sovereign re-rating (45). (a) Observed sup-norm state response $\sup_{t \leq T} \max_i |\Delta X_t^i|$ and envelope (46) at $T = 3$, as functions of δ (log scale). (b) Time profile of the observed response and of the a priori envelope at $\delta = 0.10$. Thin curves in panel (a) show the $T = 1$ observed response and envelope, for which the bound is conservative only by a factor of about four.

Figure 7 compares the a priori envelope with the simulated response. For $|\delta| \leq 0.05$, the observed sup-norm response has fitted slope approximately 0.15; deviations from this linear approximation remain below six percent on $|\delta| \leq 0.20$. The envelope is conservative by a factor of approximately 55–74 at $T = 3$ and by a factor of approximately 4 at $T = 1$. Small re-ratings change the terminal mean smoothed distress approximately linearly, but the hard-threshold classification changes discontinuously when a terminal buffer crosses zero. In particular, UniCredit crosses the threshold at $\delta = 0.15$, while Banco Santander returns above the threshold only at $\delta = -0.20$. This distinction is consistent with the theory: Lipschitz functionals satisfy (46), whereas raw threshold counts need not vary continuously with the scenario parameter.

6.5.3. The full transparency-exercise sample: three notions of rank and the sampling rate. For the full Autumn-2025 exercise, three groups with no reported sovereign holdings are removed, leaving $N = 117$ institutions. The $K = 37$ counterparty buckets with aggregate share at least 0.1% cover 99.6% of the EUR 4.01tn total. We apply (41)–(44) with $\lambda_E := c_E/K = 0.075$ and extend the four-sovereign stress vector by zero to $\eta \in \mathbb{R}^{37}$. The row means of the resulting kernel equal c_E/K . The direct system and the 37-dimensional reduction agree to sup-norm error 3×10^{-15} . At the baseline, four institutions are below the threshold at time zero, seven enter the smoothing band, no additional hard threshold crossing occurs, and the terminal mean smoothed distress is 0.024.

The full-sample kernel has algebraic rank at most 37. Its spectral decay is moderate: 11 singular modes account for 90% of the Frobenius energy and 19 account for 99%. Under the Italian-centred stress, however, the factor-aligned rank-2 surrogate retaining the France and Italy channels has sup-norm path error 7×10^{-4} , while the generic rank-12 SVD surrogate has error of order 10^{-2} ; see Figure 8. The SVD surrogates are signed from rank 2 onward (2.9% of entries at rank 2 and 16.8% at rank 12, accounting for at most 4.2% of the L^1 mass), so they are analytic approximations rather than nonnegative exposure kernels. The factor-aligned surrogates remain nonnegative. These calculations distinguish algebraic rank, spectral approximation rank, and the scenario-dependent effective rank of the dynamics. For signed kernels, (46) is applied with the absolute row norm $\bar{e} = \max_i N^{-1} \sum_j |e_{ij}|$.

The full sample also provides a resampling diagnostic for theorem 3.4. Let μ_0 be the uniform measure on the 117 observed types $(x_i, a_{i,\cdot}, b_{i,\cdot})$. For each N , we draw N types i.i.d. from μ_0 and solve the reduced system driven by the sampled empirical measure. This experiment tests the theorem on a finitely supported type law; it is not an external validation of a sampling mechanism for real banking systems.

Because μ_0 is finitely supported, $\mathbb{E} \mathcal{W}_1(\mu_0^N, \mu_0) \asymp N^{-1/2}$. The upper bound follows by comparison with the total-variation distance between the empirical and population atom weights, whose expectation is $O(\sqrt{m/N})$ for $m = 117$ atoms; a lower bound follows from the central limit theorem for the mass of any fixed atom. For the bounded-Lipschitz loss, 200 replications at each $N \in \{8, \dots, 512\}$ give fitted log-log slopes -0.48 for the terminal state-law error $\mathbb{E} \mathcal{W}_1(\nu_T^N, \nu_T)$ and -0.42 for the feedback-path error $\mathbb{E} \sup_{t \leq T} \max_k |\beta_k^N(t) - \beta_k(t)|$; see figure 9. The maximum over the 37 feedback coordinates is influenced by the smallest buckets, whose b -loadings are heavy-tailed, and convergence is correspondingly flatter at the smallest sample sizes.

For the indicator loss, the discrete population does not satisfy the density condition of remark 3.10, so the theorem gives no rate. The fitted slopes are nevertheless -0.47 and -0.49 in this dataset because the sampled thresholds remain separated from atoms of the terminal law. This observation is empirical and does not replace the density assumption, which controls worst-case threshold configurations; see also section B.




FIGURE 8. The full-sample overlap kernel ($N = 117$, $K = 37$). (a) Singular values and cumulative Frobenius energy; 11 modes carry 90% of the energy. (b) Sup-norm contagion-path error of two rank- r surrogates: the factor-aligned truncation, which keeps the r largest counterparty buckets in the factor sum, reaches 7×10^{-4} at $r = 2$, while generic SVD truncation stalls near 2×10^{-2} through $r = 11$.




FIGURE 9. Sampling convergence on the full sample, as a resampling diagnostic from the observed 117-type empirical population: i.i.d. draws of size N , 200 replications per size, mean ± 1.96 standard errors, against the $N^{-1/2}$ reference (dotted). (a) Feedback-path error. (b) Terminal state-law Wasserstein error. The smoothed loss is the setting covered by [theorem 3.4](#); the indicator loss is shown for contrast.

6.5.4. *Interpretation.* The empirical calculations illustrate three aspects of the theory. First, disclosed sovereign holdings provide a directly observable factor representation, although the resulting kernel is a common-exposure proxy rather than a bilateral interbank network. Second, the exact reduction reproduces the finite-system trajectories to numerical precision, while kernel approximation error and path error need not be proportional: factor-aligned truncations can outperform spectral truncations of higher rank. Third, the stability envelope is scenario-uniform but conservative, and the resampling experiment on the empirical 117-type population is consistent with the $N^{-1/2}$ scale in [theorem 3.4](#).

Accordingly, this section is a data-based overlap stress calculation, not a calibrated estimate of default probabilities. A bilateral-network application would require counterparty-level exposures or an explicit network-reconstruction procedure.

7. CONCLUDING REMARKS

The paper establishes a reduction framework for structured dense-network contagion. Exact factorization reduces the finite network to a finite-dimensional feedback system, while strong kernel approximation connects the finite-rank systems to a directed graphon equation. For bounded Lipschitz losses, the error separates into a type-sampling term and a kernel-approximation term. For the indicator loss, the fixed-rank sampled estimate is conditional on a measurable selection, and the graphon perturbation result is restricted to uniformly transverse approximation families. The numerical results verify the algebraic reduction and illustrate the scope of the stability estimates; the appendices report sampling and discretization diagnostics.

The occupation-time convention is not an absorbing default model; an absorbing formulation would require a path-dependent threshold functional. Other model extensions include idiosyncratic or common noise in the outside book and time-dependent exposures arising from loan maturity or rollover.

For the indicator loss, the principal analytical gap is the case of arbitrary bounded measurable kernels without a graphon-level small-ball or transversality condition. The results obtained here cover fixed rank, measurable selected finite- N solutions, factorized kernels, a non-factorized

piecewise-smooth transverse class, and uniformly transverse approximation families. A family-independent perturbation theory, together with sampled finite- N convergence at the graphon level, remains open. Generic spectral truncations are not covered because threshold regularity need not be preserved by the approximation. This restriction does not arise for bounded Lipschitz losses, for which the kernel theory applies to arbitrary bounded kernels.

Another direction is to combine the present low-rank reduction with endogenous clearing, recovery, fire-sale, and margin/liquidity feedback, in the spirit of dynamic clearing models and integrated fire-sale contagion models [6, 16]. One concrete route is to replace (23) by an augmented system of the form

$$X_t(u) = x_0(u) + \mu t + rtR_W(u) - \int_0^t (\Gamma_W X_s)(u) ds - \int_0^t \kappa(u) \mathcal{F}_s ds - \int_0^t \eta(u) \mathcal{M}_s(u) ds,$$

where $\mathcal{F}_t$ is a price-mediated fire-sale term generated by an endogenous common-asset price. A more market-microfounded specification is to introduce a cumulative liquidation process $L_t(v)$ and its liquidation rate $\dot{L}_t(v)$, and to set

$$\dot{P}_t = \kappa_P(\bar{P} - P_t) - \alpha \int_I q(v) \dot{L}_t(v) dv, \quad \dot{L}_t(v) = \chi(X_t(v), P_t, (\Gamma_W X_t)(v), \mathcal{M}_t(v)),$$

so that prices respond to contemporaneous liquidation pressure while reverting toward a normal-liquidity benchmark $\bar{P}$ at rate κ_P . Equivalently,

$$P_t = \bar{P} - \alpha \int_0^t e^{-\kappa_P(t-s)} \int_I q(v) \dot{L}_s(v) dv ds.$$

Then $\mathcal{F}_t := \bar{P} - P_t$ measures the current fire-sale discount, while $\mathcal{M}_t(u)$ captures state-dependent margin calls, liquidity hoarding, or funding stress, potentially as a function of $X_t(u)$, P_t , and $(\Gamma_W X_t)(u)$. If the additional feedback channels remain low rank or approximately low rank, the present reduction strategy should still provide a tractable surrogate.

A further direction is to extend the dense deterministic framework to sparse or random networks, where recent sparse-graphon work suggests that the relevant tools differ substantially from those used here [17]. Finally, stochastic solvency/liquidity dynamics, common noise, and nonparametric kernel estimation are natural directions if one wants to move from structural reduction toward inference and control [3, 10].

APPENDIX A. PROOFS OF THE MAIN RESULTS

This appendix collects the proofs deferred from Sections 3 and 4. Statements are not repeated; each proof opens with a pointer to the corresponding result, and the notation is that of the section in which the result is stated.

A.1. Proofs for Section 3.

Proof of Theorem 3.4. Fix $T > 0$ and let $\pi^N \in \Gamma(\mu_0^N, \mu_0)$ be an optimal coupling for $\mathcal{W}_1$. Define

$$D_t^N := \int_{\mathcal{Z} \times \mathcal{Z}} |X_t^N(z) - X_t(\tilde{z})| \pi^N(dz, d\tilde{z}).$$

From the two instances of the state map (9), driven by μ_0^N and μ_0 respectively,

$$\begin{aligned} |X_t^N(z) - X_t(\tilde{z})| &\leq |x - \tilde{x}| + t |\Lambda^N(z) - \Lambda(\tilde{z})| + \frac{1}{K} \sum_{k=1}^K |a_k - \tilde{a}_k| |\beta_k(t)| \\ &\quad + \frac{1}{K} \sum_{k=1}^K |a_k| |\beta_k^N(t) - \beta_k(t)|. \end{aligned}$$

Because $|a_k|, |b_k| \leq M$ and $|m_k(t)|, |m_k^N(t)| \leq M\ell_*$, we have $|\beta_k(t)|, |\beta_k^N(t)| \leq TM\ell_*$. Moreover, by definition of the ℓ^1 metric on $\mathcal{Z}$,

$$\left| \bar{a}_k^N - \bar{a}_k \right| \leq \int |a_k - \tilde{a}_k| \pi^N(dz, d\tilde{z}) \leq \mathcal{W}_1(\mu_0^N, \mu_0),$$

and similarly for $\bar{b}_k^N - \bar{b}_k$. Expanding the imbalance difference,

$$\Lambda^N(z) - \Lambda(\tilde{z}) = \frac{r}{K} \sum_{k=1}^K \left[a_k(\bar{b}_k^N - \bar{b}_k) + \bar{b}_k(a_k - \tilde{a}_k) - b_k(\bar{a}_k^N - \bar{a}_k) - \bar{a}_k(b_k - \tilde{b}_k) \right].$$

Using $|a_k|, |b_k|, |\bar{a}_k|, |\bar{b}_k| \leq M$ together with the bounds above, we obtain

$$\begin{aligned} \int \left| \Lambda^N(z) - \Lambda(\tilde{z}) \right| \pi^N(dz, d\tilde{z}) &\leq \frac{|r|}{K} \sum_{k=1}^K \left[M \left| \bar{b}_k^N - \bar{b}_k \right| + M \int |a_k - \tilde{a}_k| \pi^N(dz, d\tilde{z}) \right. \\ &\quad \left. + M \left| \bar{a}_k^N - \bar{a}_k \right| + M \int |b_k - \tilde{b}_k| \pi^N(dz, d\tilde{z}) \right] \\ &\leq 4|r|M \mathcal{W}_1(\mu_0^N, \mu_0). \end{aligned}$$

Integrating the previous pointwise estimate against π^N , we obtain

$$(47) \quad D_t^N \leq C_T \mathcal{W}_1(\mu_0^N, \mu_0) + \frac{M}{K} \sum_{k=1}^K \int_0^t |m_k^N(s) - m_k(s)| ds.$$

Next,

$$\begin{aligned} |m_k^N(s) - m_k(s)| &= \left| \int b_k \ell(X_s^N(z)) \mu_0^N(dz) - \int \tilde{b}_k \ell(X_s(\tilde{z})) \mu_0(d\tilde{z}) \right| \\ &\leq \int \left| b_k \ell(X_s^N(z)) - \tilde{b}_k \ell(X_s(\tilde{z})) \right| \pi^N(dz, d\tilde{z}) \\ &\leq \ell_* \mathcal{W}_1(\mu_0^N, \mu_0) + ML_\ell D_s^N. \end{aligned}$$

Substituting into (47) and absorbing constants yields

$$D_t^N \leq C_T \mathcal{W}_1(\mu_0^N, \mu_0) + C \int_0^t D_s^N ds.$$

Gronwall's lemma gives

$$\sup_{0 \leq t \leq T} D_t^N \leq C_T \mathcal{W}_1(\mu_0^N, \mu_0).$$

Finally, by the definition of pushforward and the coupling $(\Xi_t^N, \Xi_t)_{\#} \pi^N$,

$$\begin{aligned} \mathcal{W}_1(\nu_t^N, \nu_t) &\leq \int \left| \Xi_t^N(z) - \Xi_t(\tilde{z}) \right| \pi^N(dz, d\tilde{z}) \\ &\leq D_t^N + \int (|a - \tilde{a}|_1 + |b - \tilde{b}|_1) \pi^N(dz, d\tilde{z}) \\ &\leq D_t^N + \mathcal{W}_1(\mu_0^N, \mu_0), \end{aligned}$$

and (13) follows. $\square$

Proof of Theorem 3.7. Since $\beta_k(t) = \int_0^t m_k(s) ds$ and $|m_k(s)| \leq M\ell_*$, each β_k is absolutely continuous and $\dot{\beta}_k(t) = m_k(t)$ for a.e. t . By construction,

$$X_t(z) = x + \Theta_t(a, b),$$

so $\Xi_t(x, a, b) = (x + \Theta_t(a, b), a, b)$. This map is a translation in the x -variable, hence the pushforward density is exactly (14). The formula (17) follows from the same change of variables.

It remains to justify the transport equation. Let $\varphi \in C_c^1([0, T] \times \mathcal{Z})$. For fixed (x, a, b) , the map

$$t \mapsto \varphi(t, x + \Theta_t(a, b), a, b)$$

is absolutely continuous, and for a.e. t ,

$$\frac{d}{dt} \varphi(t, x + \Theta_t(a, b), a, b) = \partial_t \varphi(t, x + \Theta_t(a, b), a, b) + \dot{\Theta}_t(a, b) \partial_x \varphi(t, x + \Theta_t(a, b), a, b).$$

Since $\dot{\Theta}_t(a, b) = v_f(t, a, b)$ for a.e. t , integrating in time and then integrating against μ_0 gives (18); the terminal term vanishes because φ is compactly supported in $[0, T]$. Thus (15) holds in the distributional sense.

If $t \mapsto m_k(t)$ is continuous for every k , then $\Theta_t(a, b)$ is C^1 in t . Since $\Phi \in C_c^1(\mathcal{Z})$, the formula (14) may be differentiated pointwise:

$$\partial_t f(t, x, a, b) = -\dot{\Theta}_t(a, b) \partial_x f(t, x, a, b).$$

Because $v_f = \dot{\Theta}_t$ and v_f is independent of x , this is equivalent to the classical equation (15). $\square$

Proof of Theorem 3.11. For $1 \leq k \leq K$ define

$$F_k(t, \beta) := \int b_k \mathbf{1}_{\{\Psi_t(z, \beta) \leq 0\}} \mu_0(dz).$$

Since $|F_k(t, \beta)| \leq M$, any solution on $[0, T]$ must remain in the cube

$$\mathcal{C}_T := [-MT, MT]^K.$$

We therefore fix $\beta, \tilde{\beta} \in \mathcal{C}_T$ and apply assumption 3.9 with $C = MT$. Set

$$\Delta(\beta, \tilde{\beta}; z) := \frac{1}{K} \sum_{j=1}^K a_j (\beta_j - \tilde{\beta}_j), \quad \delta(\beta, \tilde{\beta}) := M \|\beta - \tilde{\beta}\|_\infty.$$

Then

$$|\Delta(\beta, \tilde{\beta}; z)| \leq \frac{1}{K} \sum_{j=1}^K |a_j| |\beta_j - \tilde{\beta}_j| \leq \frac{1}{K} (KM \|\beta - \tilde{\beta}\|_\infty) = \delta(\beta, \tilde{\beta})$$

for every z . If the indicators $\mathbf{1}_{\{\Psi_t(z, \beta) \leq 0\}}$ and $\mathbf{1}_{\{\Psi_t(z, \tilde{\beta}) \leq 0\}}$ differ, then either $|\Psi_t(z, \beta)| \leq \delta(\beta, \tilde{\beta})$ or $|\Psi_t(z, \tilde{\beta})| \leq \delta(\beta, \tilde{\beta})$. Let $\rho_{t, \beta}$ and $\rho_{t, \tilde{\beta}}$ denote the densities of $\Psi_t(\cdot, \beta)$ and $\Psi_t(\cdot, \tilde{\beta})$ under μ_0 . Then

$$\mu_0(|\Psi_t(\cdot, \beta)| \leq \delta(\beta, \tilde{\beta})) = \int_{-\delta(\beta, \tilde{\beta})}^{\delta(\beta, \tilde{\beta})} \rho_{t, \beta}(y) dy \leq 2M_\rho(T, MT) \delta(\beta, \tilde{\beta}),$$

and similarly for $\tilde{\beta}$. Hence

$$\begin{aligned} |F_k(t, \beta) - F_k(t, \tilde{\beta})| &\leq M \mu_0(|\Psi_t(\cdot, \beta)| \leq \delta(\beta, \tilde{\beta})) + M \mu_0(|\Psi_t(\cdot, \tilde{\beta})| \leq \delta(\beta, \tilde{\beta})) \\ &\leq 4MM_\rho(T, MT) \delta(\beta, \tilde{\beta}) \\ &\leq 4M^2 M_\rho(T, MT) \|\beta - \tilde{\beta}\|_\infty. \end{aligned}$$

Thus the feedback map is locally Lipschitz in β on $\mathcal{C}_T$, with Lipschitz constant depending on T through $M_\rho(T, MT)$, uniformly in $t \in [0, T]$. The map $(t, \beta) \mapsto F(t, \beta)$ is measurable in t , continuous in β , and bounded on $[0, T] \times \mathcal{C}_T$. The integral form of the feedback ODE

$$\beta_k(t) = \int_0^t F_k(s, \beta(s)) ds, \quad 1 \leq k \leq K,$$

cannot leave the cube $\mathcal{C}_T$ on $[0, T]$. The standard Carathéodory–Picard argument for feedback ODEs with a measurable-in-time and Lipschitz-in-state vector field therefore yields existence and uniqueness on $[0, T]$. Because T was arbitrary, the solution is global. $\square$

Proof of Lemma 3.16. For fixed $(t, \beta, \alpha, \gamma)$, the threshold map

$$L_{t,\beta,\alpha,\gamma}(z) := x + \mu t + \frac{rt}{K} \sum_{j=1}^K (a_j \gamma_j - b_j \alpha_j) - \frac{1}{K} \sum_{j=1}^K a_j \beta_j$$

is affine in $z = (x, a, b) \in \mathbb{R}^{2K+1}$. Hence the corresponding indicators form a half-space class

$$\mathcal{H}_{T,K} := \left\{ \mathbf{1}_{\{L_{t,\beta,\alpha,\gamma}(\cdot) \leq 0\}} : (t, \beta, \alpha, \gamma) \in [0, T] \times \mathcal{C}_T \times [-M, M]^{2K} \right\},$$

whose VC dimension is at most $2K + 2$. For each fixed k , multiplication by the coordinate map $z \mapsto b_k$ preserves VC-subgraph structure in the following concrete sense. If $H = \{z : L_{t,\beta,\alpha,\gamma}(z) \leq 0\}$, then for $g(z) = b_k \mathbf{1}_H(z)$,

$$\text{subgraph}(g) = \{(z, y) : z \in H, y < b_k\} \cup \{(z, y) : z \notin H, y < 0\}.$$

This is a finite Boolean combination of halfspaces in the (z, y) -space $\mathbb{R}^{2K+2}$. Hence

$$\mathcal{G}_{T,K,k} := \{b_k h : h \in \mathcal{H}_{T,K}\}$$

is a bounded VC-subgraph class with envelope M and VC-subgraph index of order K . The full class is the finite union

$$\mathcal{G}_{T,K} = \bigcup_{k=1}^K \mathcal{G}_{T,K,k}.$$

Finite unions of VC-subgraph classes remain VC-subgraph, and the corresponding $L^2(Q)$ covering numbers pick up at most a logarithmic factor in K . Thus, after the usual countable-dense reduction that makes the class pointwise measurable, there exist constants $C_1, C_2 < \infty$, depending only on the bounded parameter ranges induced by M, r , and T , such that

$$\sup_Q \log N(\varepsilon M, \mathcal{G}_{T,K}, L^2(Q)) \leq C_1 K \log(C_2/\varepsilon), \quad 0 < \varepsilon \leq 1;$$

see, for example, van der Vaart and Wellner [24, Sec. 2.6]. Applying the standard uniform-entropy maximal inequality for bounded empirical processes [24, Sec. 2.14] gives

$$\mathbb{E} \left[\sup_{g \in \mathcal{G}_{T,K}} |(\mathbb{P}_N - \mathbb{P})g| \right] \leq C \sqrt{\frac{K}{N}},$$

with $C = C(M, r, T)$. Since $\log N \geq \log 2 > 0$ for $N \geq 2$, enlarging C by the harmless factor $(\log 2)^{-1/2}$ yields the displayed bound. $\square$

Proof of Theorem 3.17. The sampled vector field is discontinuous, and the theorem does not assert existence or uniqueness for every atomic empirical measure. Let β^N be a measurable selection satisfying the hypotheses of the theorem. The estimates below are pathwise and uniform over all such selections.

Let $\alpha = (\bar{a}_1, \dots, \bar{a}_K)$ and $\gamma = (\bar{b}_1, \dots, \bar{b}_K)$ denote the population means, and similarly $\alpha^N = (\bar{a}_1^N, \dots, \bar{a}_K^N)$ and $\gamma^N = (\bar{b}_1^N, \dots, \bar{b}_K^N)$ for the sample. For any $\alpha', \gamma' \in [-M, M]^K$ define

$$F_k^{N,\alpha',\gamma'}(t, \beta) := \int g_{t,\beta,\alpha',\gamma',k}(z) \mathbb{P}_N(dz), \quad F_k^{\alpha',\gamma'}(t, \beta) := \int g_{t,\beta,\alpha',\gamma',k}(z) \mathbb{P}(dz).$$

Then $F_k^N = F_k^{N,\alpha^N,\gamma^N}$ and the deterministic feedback is $F_k = F_k^{\alpha,\gamma}$. Therefore

$$(48) \quad \sup_{t,\beta,k} |F_k^N(t, \beta) - F_k(t, \beta)| \leq \sup_{t,\beta,k,\alpha',\gamma'} |F_k^{N,\alpha',\gamma'}(t, \beta) - F_k^{\alpha',\gamma'}(t, \beta)|$$

$$(49) \quad + \sup_{t, \beta, k} |F_k^{\alpha^N, \gamma^N}(t, \beta) - F_k^{\alpha, \gamma}(t, \beta)|.$$

The first term is controlled directly by [lemma 3.16](#), uniformly over both population and sample mean vectors:

$$(50) \quad \mathbb{E} \left[\sup_{t, \beta, k, \alpha', \gamma'} |F_k^{N, \alpha', \gamma'}(t, \beta) - F_k^{\alpha', \gamma'}(t, \beta)| \right] \leq C \sqrt{\frac{K \log N}{N}}.$$

It remains to control the second term in (48), which is now a population-measure comparison and therefore legitimately uses the bounded-density hypothesis. Set

$$\eta_N := \max_{1 \leq j \leq K} \max\{|\bar{a}_j^N - \bar{a}_j|, |\bar{b}_j^N - \bar{b}_j|\}.$$

For every $z = (x, a, b)$,

$$\begin{aligned} \left| \Psi_t^{\alpha^N, \gamma^N}(z, \beta) - \Psi_t^{\alpha, \gamma}(z, \beta) \right| &= \frac{|r|t}{K} \left| \sum_{j=1}^K (a_j(\bar{b}_j^N - \bar{b}_j) - b_j(\bar{a}_j^N - \bar{a}_j)) \right| \\ &\leq 2|r|TM \eta_N =: \delta_N. \end{aligned}$$

By [remark 3.10](#), the scalar threshold projection associated with any fixed $\alpha', \gamma' \in [-M, M]^K$ and $\beta \in \mathcal{C}_T$ has density bounded by $\bar{M}$. Hence, under $\mathbb{P} = \mu_0$, the set on which the two indicators in $F_k^{\alpha^N, \gamma^N}$ and $F_k^{\alpha, \gamma}$ differ is contained in the union of two δ_N -tubes around the corresponding thresholds. Thus

$$\sup_{t, \beta, k} |F_k^{\alpha^N, \gamma^N}(t, \beta) - F_k^{\alpha, \gamma}(t, \beta)| \leq 4M\bar{M} \delta_N \leq 8|r|TM^2\bar{M} \eta_N.$$

Since $|a_j|, |b_j| \leq M$, Hoeffding's inequality and a union bound yield

$$(51) \quad \mathbb{E} \eta_N \leq C \sqrt{\frac{\log(2K)}{N}} \leq C \sqrt{\frac{K \log N}{N}},$$

for a constant $C = C(M)$ and all $N \geq 2$. Combining (48)–(51), we obtain

$$(52) \quad \mathbb{E} \left[\sup_{t, \beta, k} |F_k^N(t, \beta) - F_k(t, \beta)| \right] \leq C_T \sqrt{\frac{K \log N}{N}}.$$

Now let $\Delta_T^N := \sup_{0 \leq t \leq T} \|\beta^N(t) - \beta(t)\|_\infty$. For each $t \in [0, T]$,

$$\begin{aligned} \|\beta^N(t) - \beta(t)\|_\infty &\leq \int_0^t \|F^N(s, \beta^N(s)) - F(s, \beta(s))\|_\infty \, ds \\ &\leq \int_0^t \sup_{\beta \in \mathcal{C}_T} \|F^N(s, \beta) - F(s, \beta)\|_\infty \, ds + \int_0^t \|F(s, \beta^N(s)) - F(s, \beta(s))\|_\infty \, ds. \end{aligned}$$

By the proof of [theorem 3.11](#), the deterministic indicator feedback map is Lipschitz on $\mathcal{C}_T$ with constant $L_T := 4M^2\bar{M}$. Hence

$$\Delta_T^N \leq T \sup_{s \in [0, T], \beta \in \mathcal{C}_T} \|F^N(s, \beta) - F(s, \beta)\|_\infty + L_T \int_0^T \Delta_s^N \, ds.$$

Taking expectations, using (52), and applying Gronwall's lemma yields

$$\mathbb{E} \Delta_T^N \leq C_T \sqrt{\frac{K \log N}{N}}.$$

This proves the feedback-path estimate.

For the state-law bound, first observe that the map Ξ_t is Lipschitz in z with respect to the ℓ^1 metric on $\mathcal{Z}$, with a constant depending only on T , M , and $|r|$; indeed,

$$|\Xi_t(z) - \Xi_t(\tilde{z})| \leq C_T |z - \tilde{z}|_1.$$

Therefore

$$(53) \quad \sup_{0 \leq t \leq T} \mathcal{W}_1((\Xi_t)_\# \mu_0^N, (\Xi_t)_\# \mu_0) \leq C_T \mathcal{W}_1(\mu_0^N, \mu_0).$$

Next, since Ξ_t^N and Ξ_t are evaluated on the same sampled atoms,

$$\begin{aligned} \mathcal{W}_1((\Xi_t^N)_\# \mu_0^N, (\Xi_t)_\# \mu_0^N) &\leq \frac{1}{N} \sum_{i=1}^N |X_t^N(Z_i) - X_t(Z_i)| \\ &\leq t \sup_{1 \leq i \leq N} |\Lambda^N(Z_i) - \Lambda(Z_i)| + M \|\beta^N(t) - \beta(t)\|_\infty \\ &\leq 2|r|TM\eta_N + M \|\beta^N(t) - \beta(t)\|_\infty. \end{aligned}$$

The expectation of the additional imbalance term is controlled by (51) and is absorbed by the same $\sqrt{K \log N/N}$ rate. Taking the supremum over $t \in [0, T]$, then expectations, and combining with (53) proves the final bound. $\square$

A.2. Proofs for Section 4.

Proof of Theorem 4.2. Fix $T > 0$. Set

$$X_t^{(0)}(u) := x_0(u) + \mu t + rtR_W(u)$$

and define inductively

$$X_t^{(n+1)}(u) := x_0(u) + \mu t + rtR_W(u) - \int_0^t (\Gamma_W X_s^{(n)})(u) \, ds.$$

Since

$$|R_W(u)| \leq 2 \|W\|_{L^\infty}, \quad |(\Gamma_W \varphi)(u)| \leq \int_I |W(u, v)| |\ell(\varphi(v))| \, dv \leq \|W\|_{L^\infty} \ell_*,$$

each iterate belongs to $C([0, T]; L^\infty(I))$ and is uniformly bounded there. Moreover,

$$\begin{aligned} \|\Gamma_W \varphi - \Gamma_W \psi\|_{L^\infty} &\leq \sup_{u \in I} \int_I |W(u, v)| |\ell(\varphi(v)) - \ell(\psi(v))| \, dv \\ &\leq \|W\|_{L^\infty} L_\ell \|\varphi - \psi\|_{L^1} \leq \|W\|_{L^\infty} L_\ell \|\varphi - \psi\|_{L^\infty}. \end{aligned}$$

Hence, with

$$\Delta_t^{(n)} := \sup_{0 \leq s \leq t} \|X_s^{(n+1)} - X_s^{(n)}\|_{L^\infty},$$

we obtain

$$\Delta_t^{(n)} \leq \|W\|_{L^\infty} L_\ell \int_0^t \Delta_s^{(n-1)} \, ds.$$

A standard factorial estimate shows that $(X^{(n)})_n$ is Cauchy in $C([0, T]; L^\infty(I))$. The limit therefore belongs to $C([0, T]; L^\infty(I))$ and solves (23). Uniqueness in $C([0, T]; L^\infty(I))$ follows from the same estimate applied to the difference of two solutions.

For stability, write

$$\begin{aligned} X_t^1(u) - X_t^2(u) &= x_0^1(u) - x_0^2(u) + rt(R_{W_1}(u) - R_{W_2}(u)) \\ &\quad - \int_0^t ((\Gamma_{W_1} X_s^1)(u) - (\Gamma_{W_2} X_s^2)(u)) \, ds. \end{aligned}$$

Using

$$\|R_{W_1} - R_{W_2}\|_{L^1} \leq 2 \|W_1 - W_2\|_{L^1(I^2)}$$

and

$$\|\Gamma_{W_1}\varphi - \Gamma_{W_2}\psi\|_{L^1} \leq \ell_* \|W_1 - W_2\|_{L^1(I^2)} + M_W L_\ell \|\varphi - \psi\|_{L^1},$$

we find

$$\|X_t^1 - X_t^2\|_{L^1} \leq \|x_0^1 - x_0^2\|_{L^1} + (2|r|T + T\ell_*) \|W_1 - W_2\|_{L^1(I^2)} + M_W L_\ell \int_0^t \|X_s^1 - X_s^2\|_{L^1} ds.$$

Gronwall's lemma yields (24). $\square$

Proof of Theorem 4.4. Because the rank- K graphon equation with kernel $W^{(K)}$ is exactly the continuum version of the rank- K factor model, theorem 3.4 applied to the pair $(\mu_0^{N,K}, \mu_0^{(K)})$ gives

$$\sup_{0 \leq t \leq T} \mathcal{W}_1(\nu_t^{N,K}, \nu_t^K) \leq C_T \mathcal{W}_1(\mu_0^{N,K}, \mu_0^{(K)}),$$

after projecting the joint law of (X_t, a, b) onto the state coordinate. Next, couple ν_t^K and ν_t by the common random variable U . Then

$$\mathcal{W}_1(\nu_t^K, \nu_t) \leq \mathbb{E} |X_t^K(U) - X_t(U)| = \|X_t^K - X_t\|_{L^1(I)},$$

and theorem 4.2 yields

$$\sup_{0 \leq t \leq T} \mathcal{W}_1(\nu_t^K, \nu_t) \leq C_T \left(\|W^{(K)} - W\|_{L^1(I^2)} + \|x_0^{(K)} - x_0\|_{L^1(I)} \right).$$

The claim follows from the triangle inequality. $\square$

Proof of Proposition 4.6. For each K , the graphon model with kernel $W^{(K)}$ is exactly the continuum version of the rank- K factor model generated by $\mu_0^{(K)}$. Let $\beta^{K,\varepsilon}$ and $\beta^{K,\text{ind}}$ denote the corresponding feedback coordinates for the smoothed and indicator losses. Writing $F_k^{K,\varepsilon}$ and $F_k^{K,\text{ind}}$ for the associated feedback right-hand sides, observe that for every $y \in \mathbb{R}$,

$$|\ell_\varepsilon(y) - \mathbf{1}_{\{y \leq 0\}}| \leq \mathbf{1}_{\{0 < y < \varepsilon\}}.$$

Together with the uniform density hypothesis, this implies

$$\sup_{\beta \in [-MT, MT]^K} |F_k^{K,\varepsilon}(t, \beta) - F_k^{K,\text{ind}}(t, \beta)| \leq MM_\rho(T, MT) \varepsilon, \quad 1 \leq k \leq K.$$

Moreover, by the proof of theorem 3.11, the indicator feedback map is Lipschitz on $[-MT, MT]^K$ with constant $4M^2M_\rho(T, MT)$, uniformly in K . Therefore

$$\|\beta^{K,\varepsilon}(t) - \beta^{K,\text{ind}}(t)\|_\infty \leq MM_\rho(T, MT) t \varepsilon + 4M^2M_\rho(T, MT) \int_0^t \|\beta^{K,\varepsilon}(s) - \beta^{K,\text{ind}}(s)\|_\infty ds,$$

so Gronwall's lemma yields

$$\sup_{0 \leq t \leq T} \|\beta^{K,\varepsilon}(t) - \beta^{K,\text{ind}}(t)\|_\infty \leq C_T \varepsilon$$

with C_T independent of K . Returning to the state map and using $\max_k \|a_k^{(K)}\|_{L^\infty} \leq M$, we obtain

$$\begin{aligned} \sup_{0 \leq t \leq T} \|X_t^{K,\varepsilon} - X_t^{K,\text{ind}}\|_{L^1(I)} &\leq \sup_{0 \leq t \leq T} \|X_t^{K,\varepsilon} - X_t^{K,\text{ind}}\|_{L^\infty(I)} \\ &\leq M \sup_{0 \leq t \leq T} \|\beta^{K,\varepsilon}(t) - \beta^{K,\text{ind}}(t)\|_\infty \\ &\leq C_T \varepsilon. \end{aligned}$$

The Wasserstein bound follows by coupling both laws through the same $U \sim \text{Unif}(I)$. $\square$

Proof of Proposition 4.9. The bound $\|\Pi_K W\|_{L^\infty} \leq \|W\|_{L^\infty}$ is immediate from Jensen's inequality on each block average. Since $\Pi_K W$ is constant on each rectangle $I_i^{(K)} \times I_j^{(K)}$, the associated integral operator maps $L^2(I)$ into the K -dimensional space of functions that are constant on the partition $\{I_j^{(K)}\}_{j=1}^K$; hence its rank is at most K . The range dimension is determined by the u -partition, so the operator rank is K rather than K^2 even though the kernel has K^2 rectangular cells.

For the L^1 convergence, note that $\Pi_K W$ is exactly the conditional expectation of W with respect to the product σ -algebra generated by the partition $\{I_j^{(K)}\}$ in each coordinate. These product σ -algebras increase to the Borel σ -algebra on I^2 modulo null sets, and $W \in L^\infty(I^2) \subset L^1(I^2)$. Therefore the martingale convergence theorem gives $\Pi_K W \rightarrow W$ in $L^1(I^2)$.

If W is piecewise C^1 on finitely many rectangles with bounded first derivatives, then the cells entirely contained in one smooth rectangle contribute $O(K^{-1})$ by the mean-value theorem, while the cells intersecting the finitely many rectangle boundaries occupy total area $O(K^{-1})$ and contribute at most $2\|W\|_{L^\infty}$ there. Summing these two contributions yields $\|\Pi_K W - W\|_{L^1(I^2)} = O(K^{-1})$. $\square$

Proof of Theorem 4.16. Let $A_* := \|a\|_{L^\infty(I \times \Theta)}$ and $\nu_* := \nu(\Theta)$. We use the $L^1(\Theta)$ topology for the fixed point and retain the L^∞ bound as an a priori constraint; this avoids separability issues in $L^\infty(\Theta)$. For $c, \tilde{c} \in L^\infty(\Theta, \nu)$ with $\|c\|_{L^\infty}, \|\tilde{c}\|_{L^\infty} \leq TB_*$,

$$|\Psi_t(u, c) - \Psi_t(u, \tilde{c})| \leq A_* \|c - \tilde{c}\|_{L^1(\Theta, \nu)} =: \Delta(c, \tilde{c}).$$

If the indicators $\mathbf{1}_{\{\Psi_t(u, c) \leq 0\}}$ and $\mathbf{1}_{\{\Psi_t(u, \tilde{c}) \leq 0\}}$ differ, then either $|\Psi_t(u, c)| \leq \Delta(c, \tilde{c})$ or $|\Psi_t(u, \tilde{c})| \leq \Delta(c, \tilde{c})$. The density bound from assumption 4.14 therefore gives

$$\lambda(\{u : |\Psi_t(u, c)| \leq \Delta(c, \tilde{c})\}) \leq 2M_\rho^\Theta(T, TB_*) \Delta(c, \tilde{c}),$$

and similarly for $\tilde{c}$. Hence, for a.e. θ ,

$$|F_t(c)(\theta) - F_t(\tilde{c})(\theta)| \leq 4B_* M_\rho^\Theta(T, TB_*) A_* \|c - \tilde{c}\|_{L^1(\Theta, \nu)}.$$

After integrating in θ , F_t is Lipschitz from the L^∞ ball of radius TB_* , equipped with the L^1 distance, into $L^1(\Theta)$, with constant $4\nu_* B_* M_\rho^\Theta(T, TB_*) A_*$.

For each $t \in [0, T]$, define the closed subset

$$K_t^\Theta := \{c \in L^1(\Theta, \nu) : |c(\theta)| \leq tB_* \text{ for } \nu\text{-a.e. } \theta\}.$$

Define the integral feedback map on

$$\mathcal{B}_T^\Theta := \{C \in C([0, T]; L^1(\Theta, \nu)) : C_t \in K_t^\Theta \text{ for every } t \in [0, T]\}.$$

This set is closed in $C([0, T]; L^1(\Theta, \nu))$: if $C^n \rightarrow C$ uniformly in L^1 , then $C_t^n \rightarrow C_t$ in L^1 for each fixed t , and the closedness of K_t^Θ gives $C_t \in K_t^\Theta$.

Given $C \in \mathcal{B}_T^\Theta$, choose a jointly measurable representative of the L^1 -valued path $t \mapsto C_t$, which exists because $L^1(\Theta, \nu)$ is separable and the path is continuous, hence strongly measurable. Then $(s, \theta) \mapsto F_s(C_s)(\theta)$ is jointly measurable by Fubini and bounded by B_* , so the Bochner integral in $L^1(\Theta)$ is well defined. The map sends $\mathcal{B}_T^\Theta$ into itself, and the preceding Lipschitz estimate gives the usual factorial Picard bound in $C([0, T]; L^1(\Theta))$. Hence Picard iteration yields a unique fixed point in $\mathcal{B}_T^\Theta$.

Finally, $t \mapsto \int_\Theta a(\cdot, \theta) C_t(\theta) \nu(d\theta)$ is continuous as an $L^\infty(I)$ -valued map because its increments are bounded by $A_* \|C_t - C_s\|_{L^1(\Theta)}$. The representation (26) therefore yields $X \in C([0, T]; L^\infty(I))$, and inserting the fixed-point identity for C gives the graphon equation with indicator loss.

Conversely, any graphon indicator solution X defines

$$C_t(\theta) = \int_0^t \int_I b(v, \theta) \mathbf{1}_{\{X_s(v) \leq 0\}} dv ds.$$

This C satisfies $|C_t(\theta)| \leq tB_*$ and solves the integral feedback equation (29). The uniqueness of C therefore implies uniqueness of the corresponding graphon profile X . $\square$

Proof of Theorem 4.19. For $t \in [0, T]$ and $h \in L^\infty(I)$ define

$$G_t(h)(u) := \mathbf{1}_{\{\Psi_t(u, h) \leq 0\}}.$$

If $h, \tilde{h} \in L^\infty(I)$ satisfy $\|h\|_{L^\infty}, \|\tilde{h}\|_{L^\infty} \leq T$, then

$$\left| \Psi_t(u, h) - \Psi_t(u, \tilde{h}) \right| \leq \int_I |W(u, v)| |h(v) - \tilde{h}(v)| dv \leq \|W\|_{L^\infty} \|h - \tilde{h}\|_{L^1} =: \Delta(h, \tilde{h}).$$

If the indicators $G_t(h)(u)$ and $G_t(\tilde{h})(u)$ differ, then either $|\Psi_t(u, h)| \leq \Delta(h, \tilde{h})$ or $|\Psi_t(u, \tilde{h})| \leq \Delta(h, \tilde{h})$. Let $\rho_{t,h}$ and $\rho_{t,\tilde{h}}$ denote the densities of $\Psi_t(U, h)$ and $\Psi_t(U, \tilde{h})$ from assumption 4.18. Then

$$\lambda\left(\left\{u : |\Psi_t(u, h)| \leq \Delta(h, \tilde{h})\right\}\right) = \int_{-\Delta(h, \tilde{h})}^{\Delta(h, \tilde{h})} \rho_{t,h}(y) dy \leq 2M_\rho^W(T, T) \Delta(h, \tilde{h}),$$

and similarly for $\tilde{h}$. Therefore

$$\begin{aligned} \|G_t(h) - G_t(\tilde{h})\|_{L^1} &\leq \lambda\left(\left\{u : |\Psi_t(u, h)| \leq \Delta(h, \tilde{h})\right\}\right) + \lambda\left(\left\{u : |\Psi_t(u, \tilde{h})| \leq \Delta(h, \tilde{h})\right\}\right) \\ &\leq 4M_\rho^W(T, T) \|W\|_{L^\infty} \|h - \tilde{h}\|_{L^1}. \end{aligned}$$

Thus G_t is Lipschitz on the L^∞ -ball of radius T , when measured in the L^1 norm.

Now define the integral feedback map

$$(\mathcal{TH})_t := \int_0^t G_s(H_s) ds,$$

on the set

$$\mathcal{B}_T := \{H \in C([0, T]; L^1(I)) : H_t \in K_t \text{ for every } t \in [0, T]\},$$

where

$$K_t := \{h \in L^1(I) : 0 \leq h(u) \leq t \text{ for a.e. } u \in I\}.$$

For each $t \in [0, T]$, the set K_t is closed in $L^1(I)$. Hence $\mathcal{B}_T$ is closed in $C([0, T]; L^1(I))$: if $H^n \rightarrow H$ uniformly in L^1 , then $H_t^n \rightarrow H_t$ in L^1 for each fixed t , and the closedness of K_t gives $H_t \in K_t$. Given $H \in \mathcal{B}_T$, the path $s \mapsto H_s$ is continuous from $[0, T]$ into the separable Banach space $L^1(I)$, hence strongly measurable. By the standard measurable-representative theorem for strongly measurable L^1 -valued maps (equivalently, by Pettis–Bochner measurability plus separability of $L^1(I)$), there is a jointly measurable representative $\hat{H} : [0, T] \times I \rightarrow \mathbb{R}$ of this path. A different jointly measurable representative agrees with $\hat{H}(s, \cdot)$ as an element of $L^1(I)$ for a.e. s , hence gives the same kernel integral and indicator in $L^1(I)$ for a.e. s . Then

$$(s, u) \mapsto \Psi_s(u, H_s) = x_0(u) + \mu s + rsR_W(u) - \int_I W(u, v) \hat{H}(s, v) dv$$

is jointly measurable by Fubini's theorem, and therefore so is $(s, u) \mapsto G_s(H_s)(u)$. In particular, $s \mapsto G_s(H_s)$ is strongly measurable as an $L^1(I)$ -valued map and bounded by 1, so $(\mathcal{TH})_t$ is a well-defined Bochner integral. Since $0 \leq G_s(H_s)(u) \leq 1$, we have

$$0 \leq (\mathcal{TH})_t(u) \leq t \quad \text{for every } t \in [0, T] \text{ and for a.e. } u \in I,$$

and for $0 \leq t \leq t' \leq T$,

$$\|(\mathcal{T}H)_{t'} - (\mathcal{T}H)_t\|_{L^1} \leq \int_t^{t'} \|G_s(H_s)\|_{L^1} ds \leq |t' - t|.$$

Thus $\mathcal{T}$ sends $\mathcal{B}_T$ into itself. In particular, by induction each Picard iterate admits a jointly measurable representative of the same form.

Starting from $H^{(0)} \equiv 0$ and iterating

$$H^{(n+1)} := \mathcal{T}H^{(n)},$$

we obtain for

$$\Delta_n(t) := \sup_{0 \leq s \leq t} \|H_s^{(n+1)} - H_s^{(n)}\|_{L^1}$$

the bound

$$\Delta_n(t) \leq 4M_\rho^W(T, T) \|W\|_{L^\infty} \int_0^t \Delta_{n-1}(s) ds.$$

A standard factorial estimate therefore shows that $(H^{(n)})_{n \geq 0}$ is Cauchy in $C([0, T]; L^1(I))$, so it converges to some $H \in \mathcal{B}_T$ satisfying (33); here we use that $\mathcal{B}_T$ is closed in $C([0, T]; L^1(I))$. If $H, \tilde{H} \in \mathcal{B}_T$ are two fixed points of $\mathcal{T}$, then

$$\|H_t - \tilde{H}_t\|_{L^1} \leq \int_0^t \|G_s(H_s) - G_s(\tilde{H}_s)\|_{L^1} ds \leq 4M_\rho^W(T, T) \|W\|_{L^\infty} \int_0^t \|H_s - \tilde{H}_s\|_{L^1} ds,$$

so Gronwall's lemma yields $H = \tilde{H}$. Hence the fixed point is unique in $\mathcal{B}_T$.

Given this unique H , define X by (35). Since $H \in C([0, T]; L^1(I))$ and $W \in L^\infty(I^2)$, the map $t \mapsto X_t$ belongs to $C([0, T]; L^\infty(I))$. Moreover, from (33) and (35),

$$H_t(v) = \int_0^t \mathbf{1}_{\{X_s(v) \leq 0\}} ds.$$

Hence by Fubini,

$$\int_I W(u, v) H_t(v) dv = \int_0^t \int_I W(u, v) \mathbf{1}_{\{X_s(v) \leq 0\}} dv ds,$$

which is exactly (23) with indicator loss. Conversely, any solution X of (23) generates through (30) a cumulative profile H satisfying (33), so uniqueness of H implies uniqueness of X . $\square$

Proof of Corollary 4.20. For $i = 1, 2$ define

$$\Psi_t^i(u, h) := x_0^i(u) + \mu t + rtR_W(u) - \int_I W(u, v) h(v) dv, \quad G_t^i(h)(u) := \mathbf{1}_{\{\Psi_t^i(u, h) \leq 0\}}.$$

Set

$$\eta := \|x_0^1 - x_0^2\|_{L^\infty(I)}, \quad \Delta(h, \tilde{h}) := \|W\|_{L^\infty} \|h - \tilde{h}\|_{L^1(I)}.$$

If $\|h\|_{L^\infty}, \|\tilde{h}\|_{L^\infty} \leq T$, then

$$|\Psi_t^1(u, h) - \Psi_t^2(u, \tilde{h})| \leq \eta + \Delta(h, \tilde{h}) \quad \text{for every } u \in I.$$

Hence, if $G_t^1(h)(u) \neq G_t^2(\tilde{h})(u)$, then either $|\Psi_t^1(u, h)| \leq \eta + \Delta(h, \tilde{h})$ or $|\Psi_t^2(u, \tilde{h})| \leq \eta + \Delta(h, \tilde{h})$.

Using the density bounds for the laws of $\Psi_t^1(U, h)$ and $\Psi_t^2(U, \tilde{h})$ with $U \sim \lambda$, we obtain

$$\begin{aligned} \|G_t^1(h) - G_t^2(\tilde{h})\|_{L^1} &\leq 4M_\rho^W(T, T)(\eta + \Delta(h, \tilde{h})) \\ &\leq 4M_\rho^W(T, T)(\eta + \|W\|_{L^\infty} \|h - \tilde{h}\|_{L^1}). \end{aligned}$$

Now apply this estimate with $h = H_t^1$ and $\tilde{h} = H_t^2$. Since each H^i belongs to the ball $\{0 \leq h \leq T\}$ from [theorem 4.19](#), we get

$$\left\| H_t^1 - H_t^2 \right\|_{L^1} \leq 4M_\rho^W(T, T)\eta t + 4M_\rho^W(T, T) \|W\|_{L^\infty} \int_0^t \left\| H_s^1 - H_s^2 \right\|_{L^1} ds.$$

Gronwall's lemma yields the first bound. Finally,

$$\left| X_t^1(u) - X_t^2(u) \right| \leq \eta + \int_I |W(u, v)| \left| H_t^1(v) - H_t^2(v) \right| dv \leq \eta + \|W\|_{L^\infty} \left\| H_t^1 - H_t^2 \right\|_{L^1},$$

so integrating over $u \in I$ gives the stated estimate for $X^1 - X^2$. $\square$

Proof of [Proposition 4.23](#). Fix $t \in [0, T]$ and $h \in L^\infty(I)$ with $\|h\|_{L^\infty} \leq C$. On each interval I_j , the map

$$u \mapsto \Psi_t(u, h) = x_0(u) + \mu t + rtR_W(u) - \int_I W(u, v)h(v) dv$$

is C^1 by dominated differentiation under the integral sign. Moreover,

$$\partial_u R_W(u) = \int_I \partial_1 W(u, v) dv - \int_I \partial_2 W(v, u) dv,$$

so on I_j we have

$$\partial_u \Psi_t(u, h) = x'_0(u) + rt\partial_u R_W(u) - \int_I \partial_1 W(u, v)h(v) dv.$$

Therefore

$$\begin{aligned} |\partial_u \Psi_t(u, h)| &\geq |x'_0(u)| - |r|t |\partial_u R_W(u)| - \int_I |\partial_1 W(u, v)| |h(v)| dv \\ &\geq m_0 - |r|T(A_1 + A_2) - CA_1 = m > 0. \end{aligned}$$

Since $\partial_u \Psi_t(\cdot, h)$ is continuous and bounded away from zero on each branch, its sign is constant on each I_j , and hence $u \mapsto \Psi_t(u, h)$ is monotone there. Thus the profile has at most J monotone branches. Applying the one-dimensional change-of-variables argument from [proposition 4.15](#) branchwise gives a density bound J/m for $\Psi_t(U, h)$ with $U \sim \lambda$. $\square$

Proof of [Theorem 4.28](#). Fix $\alpha \in \mathcal{A}$ and define

$$G_t^\alpha(h)(u) := \mathbf{1}_{\{\Psi_t^\alpha(u, h) \leq 0\}}.$$

If $h, \tilde{h} \in L^\infty(I)$ satisfy $\|h\|_{L^\infty}, \|\tilde{h}\|_{L^\infty} \leq T$, then

$$\left| \Psi_t^\alpha(u, h) - \Psi_t^\alpha(u, \tilde{h}) \right| \leq \int_I |W^\alpha(u, v)| |h(v) - \tilde{h}(v)| dv \leq M_W \|h - \tilde{h}\|_{L^1(I)}.$$

Applying [proposition 4.27](#) to $f = \Psi_t^\alpha(\cdot, h)$ and $g = \Psi_t^\alpha(\cdot, \tilde{h})$ yields

$$\left\| G_t^\alpha(h) - G_t^\alpha(\tilde{h}) \right\|_{L^1(I)} \leq \frac{2JM_W}{m} \|h - \tilde{h}\|_{L^1(I)}.$$

The Picard iteration argument from the proof of [theorem 4.19](#) therefore applies verbatim with this Lipschitz constant, giving existence and uniqueness of (H^α, X^α) for every $\alpha \in \mathcal{A}$.

Now fix $\alpha_1, \alpha_2 \in \mathcal{A}$ and write

$$\eta_{12} := \|x_0^{\alpha_1} - x_0^{\alpha_2}\|_{L^\infty(I)} + T(1 + 2|r|) \|W^{\alpha_1} - W^{\alpha_2}\|_{L^\infty(I^2)}.$$

Because

$$\|R_{W^{\alpha_1}} - R_{W^{\alpha_2}}\|_{L^\infty(I)} \leq 2 \|W^{\alpha_1} - W^{\alpha_2}\|_{L^\infty(I^2)},$$

and $0 \leq H_t^{\alpha_2} \leq t \leq T$ a.e. (hence $\|H_t^{\alpha_2}\|_{L^1(I)} \leq T$), we obtain for every $u \in I$,

$$\begin{aligned} |\Psi_t^{\alpha_1}(u, H_t^{\alpha_1}) - \Psi_t^{\alpha_2}(u, H_t^{\alpha_2})| &\leq \|x_0^{\alpha_1} - x_0^{\alpha_2}\|_{L^\infty(I)} + |r|t \|R_{W^{\alpha_1}} - R_{W^{\alpha_2}}\|_{L^\infty(I)} \\ &\quad + \int_I |W^{\alpha_1}(u, v)| |H_t^{\alpha_1}(v) - H_t^{\alpha_2}(v)| \, dv \\ &\quad + \int_I |W^{\alpha_1}(u, v) - W^{\alpha_2}(u, v)| |H_t^{\alpha_2}(v)| \, dv \\ &\leq \eta_{12} + M_W \|H_t^{\alpha_1} - H_t^{\alpha_2}\|_{L^1(I)}. \end{aligned}$$

Applying [proposition 4.27](#) to the function $f = \Psi_t^{\alpha_1}(\cdot, H_t^{\alpha_1})$ and the perturbation $g = \Psi_t^{\alpha_2}(\cdot, H_t^{\alpha_2})$ gives

$$\|G_t^{\alpha_1}(H_t^{\alpha_1}) - G_t^{\alpha_2}(H_t^{\alpha_2})\|_{L^1(I)} \leq \frac{2J}{m} \left(\eta_{12} + M_W \|H_t^{\alpha_1} - H_t^{\alpha_2}\|_{L^1(I)} \right).$$

Since

$$H_t^{\alpha_i}(u) = \int_0^t G_s^{\alpha_i}(H_s^{\alpha_i})(u) \, ds,$$

we obtain

$$\|H_t^{\alpha_1} - H_t^{\alpha_2}\|_{L^1(I)} \leq \frac{2JT}{m} \eta_{12} + \frac{2JM_W}{m} \int_0^t \|H_s^{\alpha_1} - H_s^{\alpha_2}\|_{L^1(I)} \, ds.$$

Gronwall's lemma proves [\(36\)](#). Finally,

$$\begin{aligned} \|X_t^{\alpha_1} - X_t^{\alpha_2}\|_{L^1(I)} &\leq \|x_0^{\alpha_1} - x_0^{\alpha_2}\|_{L^1(I)} + |r|t \|R_{W^{\alpha_1}} - R_{W^{\alpha_2}}\|_{L^1(I)} \\ &\quad + \|W^{\alpha_1}\|_{L^\infty(I^2)} \|H_t^{\alpha_1} - H_t^{\alpha_2}\|_{L^1(I)} + T \|W^{\alpha_1} - W^{\alpha_2}\|_{L^\infty(I^2)} \\ &\leq \eta_{12} + M_W \|H_t^{\alpha_1} - H_t^{\alpha_2}\|_{L^1(I)}, \end{aligned}$$

which gives [\(37\)](#). $\square$

A.3. Elementary facts for the model derivation. This subsection records the two elementary facts used in [section 2.1](#). Here $e_{ij} \geq 0$ are bilateral exposures with $e_{ii} = 0$, $\theta \in (0, 1]$ is the loss rate, $\mathbb{D}_0 = \{i : x_i \leq 0\}$, and $X^{(k)}, \mathbb{D}_k$ are defined by [\(2\)](#). The arguments use only nonnegativity of the weights; the zero-diagonal convention is not needed for the conclusions.

Proposition A.1. *The default sets are nondecreasing, $\mathbb{D}_k \subseteq \mathbb{D}_{k+1}$ for all $k \geq 0$; the buffers are nonincreasing, $X^{(k+1)} \leq X^{(k)}$ coordinatewise for all $k \geq 1$; and both sequences are constant from some round $k_0 \leq N$ on. The terminal vector $X^{(N+1)}$ is the unique maximal solution of the fixed-point system [\(3\)](#) with respect to the coordinatewise partial order on $\mathbb{R}^N$.*

Proof. Step 1: monotonicity. For $i \in \mathbb{D}_0$ we have $X_i^{(k)} = x_i \leq 0$ for every k , so $\mathbb{D}_0 \subseteq \mathbb{D}_k$ for all k ; in particular $\mathbb{D}_0 \subseteq \mathbb{D}_1$. Assume $\mathbb{D}_{k-1} \subseteq \mathbb{D}_k$ for some $k \geq 1$. For $i \notin \mathbb{D}_0$, subtracting the two updates in [\(2\)](#) gives

$$X_i^{(k+1)} - X_i^{(k)} = -\theta \sum_{j=1}^N e_{ij} (\mathbf{1}_{\mathbb{D}_k}(j) - \mathbf{1}_{\mathbb{D}_{k-1}}(j)) \leq 0,$$

since $e_{ij} \geq 0$ and $\mathbf{1}_{\mathbb{D}_k} \geq \mathbf{1}_{\mathbb{D}_{k-1}}$ pointwise; for $i \in \mathbb{D}_0$ both iterates equal x_i . Hence $X^{(k+1)} \leq X^{(k)}$, and therefore $\mathbb{D}_{k+1} = \{i : X_i^{(k+1)} \leq 0\} \supseteq \{i : X_i^{(k)} \leq 0\} = \mathbb{D}_k$, closing the induction.

Step 2: stabilization. If $\mathbb{D}_{k+1} = \mathbb{D}_k$ for some k , then [\(2\)](#) returns the same vector at every later round, so both sequences are constant from round $k+1$ on. Since $\mathbb{D}_0 \subseteq \mathbb{D}_1 \subseteq \dots \subseteq \{1, \dots, N\}$, at most N of these inclusions can be strict, so $\mathbb{D}_{k_0+1} = \mathbb{D}_{k_0}$ for some $k_0 \leq N$.

Step 3: $X^{(N+1)}$ solves (3). By the definition of (2), the update is applied exactly to the coordinates with $x_i > 0$, so for every $k \geq 0$,

$$X_i^{(k+1)} = x_i + \mathbf{1}_{\{x_i > 0\}} \left[\sum_{j=1}^N (e_{ij} - e_{ji}) - \theta \sum_{j=1}^N e_{ij} \mathbf{1}_{\mathbb{D}_k}(j) \right], \quad 1 \leq i \leq N.$$

By Step 2, $\mathbb{D}_N = \mathbb{D}_{N+1} = \{j : X_j^{(N+1)} \leq 0\}$, so taking $k = N$ above shows that $X^{(N+1)}$ satisfies (3).

Step 4: maximality and uniqueness. Let X be any solution of (3). For $j \in \mathbb{D}_0$, the fixed point forces $X_j = x_j \leq 0$, so $\mathbf{1}_{\{X_j \leq 0\}} \geq \mathbf{1}_{\mathbb{D}_0}(j)$ for all j . For $x_i > 0$ this yields

$$X_i = x_i + \left[\sum_j (e_{ij} - e_{ji}) - \theta \sum_j e_{ij} \mathbf{1}_{\{X_j \leq 0\}} \right] \leq x_i + \left[\sum_j (e_{ij} - e_{ji}) - \theta \sum_j e_{ij} \mathbf{1}_{\mathbb{D}_0}(j) \right] = X_i^{(1)},$$

while $X_i = x_i = X_i^{(1)}$ for $x_i \leq 0$; hence $X \leq X^{(1)}$ coordinatewise and therefore $\mathbf{1}_{\{X_j \leq 0\}} \geq \mathbf{1}_{\mathbb{D}_1}(j)$. Iterating the same comparison gives $X \leq X^{(k+1)}$ and $\mathbf{1}_{\{X \leq 0\}} \geq \mathbf{1}_{\mathbb{D}_{k+1}}$ for every k , so in particular $X \leq X^{(N+1)}$. Thus $X^{(N+1)}$ dominates every solution; a maximal solution is unique because two maximal solutions dominate each other. $\square$

Proposition A.2. *The stopped system (5) admits a unique solution.*

Proof. Existence. Set $\tau_{(0)} := 0$, $\overline{\mathbb{D}}_0 := \mathbb{D}_0$, and $X_{\tau_{(0)}}^{(0),i} := x_i$. For $i \in \mathbb{D}_0$ the equation forces $\tau_i = 0$ and $X_t^i \equiv x_i$. Given $\tau_{(m)} < T$ and the cumulative default set $\overline{\mathbb{D}}_m$, terminate the construction if $\overline{\mathbb{D}}_m = \{1, \dots, N\}$. Otherwise, define, for $i \notin \overline{\mathbb{D}}_m$ and $t \geq \tau_{(m)}$, the affine functions

$$X_t^{(m+1),i} := X_{\tau_{(m)}}^{(m),i} + \int_{\tau_{(m)}}^t \left[\mu_i + r \sum_{j=1}^N (e_{ij} - e_{ji}) - \theta r \sum_{j=1}^N e_{ij} \mathbf{1}_{\overline{\mathbb{D}}_m}(j) \right] ds,$$

and set

$$\tau_{(m+1)} := \inf \left\{ t \geq \tau_{(m)} : \min_{i \notin \overline{\mathbb{D}}_m} X_t^{(m+1),i} = 0 \right\} \wedge T, \quad \Delta \mathbb{D}_{m+1} := \{i \notin \overline{\mathbb{D}}_m : X_{\tau_{(m+1)}}^{(m+1),i} = 0\},$$

and $\overline{\mathbb{D}}_{m+1} := \overline{\mathbb{D}}_m \cup \Delta \mathbb{D}_{m+1}$. Since every $i \notin \overline{\mathbb{D}}_m$ satisfies $X_{\tau_{(m)}}^{(m),i} > 0$ and the drift is bounded, $\tau_{(m+1)} > \tau_{(m)}$. Paste the pieces: for $i \in \mathbb{D}_0$, take $X^i \equiv x_i$ and $\tau_i = 0$; for $i \in \Delta \mathbb{D}_m$ with $m \geq 1$, take $X_t^i := \sum_{l \leq m} X_t^{(l),i} \mathbf{1}_{[\tau_{(l-1)}, \tau_{(l)})}(t)$, frozen at 0 from $\tau_{(m)}$ on, and $\tau_i := \tau_{(m)}$; for institutions that never default, take the full concatenation up to T and $\tau_i := T$. On each interval $[\tau_{(m)}, \tau_{(m+1)})$ the indicator processes satisfy $\mathbf{1}_{\{X_s^j \leq 0\}} = \mathbf{1}_{\overline{\mathbb{D}}_m}(j)$ —solvent institutions are strictly positive there and defaulted ones are frozen at a nonpositive value—so the pasted process satisfies (5). Whenever $\tau_{(m+1)} < T$ the set $\Delta \mathbb{D}_{m+1}$ is nonempty, so $\overline{\mathbb{D}}_m$ grows strictly, and the construction reaches T after at most N steps.

Uniqueness. Let (X, τ) be any solution of (5). For $i \in \mathbb{D}_0$ the equation forces $X^i \equiv x_i$ and $\tau_i = 0$, as noted. Let $\tau^{(1)} := \min_{i \notin \overline{\mathbb{D}}_0} \tau_i$. On $[0, \tau^{(1)})$ every $i \notin \overline{\mathbb{D}}_0$ satisfies $X_s^i > 0$, so the indicators in (5) equal $\mathbf{1}_{\overline{\mathbb{D}}_0}$, and each X^i coincides with the affine function $X^{(1),i}$ above. Consequently $\tau^{(1)} = \tau_{(1)}$, and the set of institutions reaching zero at that time is exactly $\Delta \mathbb{D}_1$. If $\tau^{(1)} = T$, the solution coincides with the constructed one. Otherwise, repeat the argument on $[\tau_{(1)}, \tau^{(2)})$ with $\overline{\mathbb{D}}_1$ in place of $\overline{\mathbb{D}}_0$; after at most N iterations the solution is identified with the constructed one on all of $[0, T]$. $\square$

A.4. Existence of selected sampled indicator solutions. The following proposition is a fixed-sample compactness statement for the regularized systems. It gives a pathwise indicator solution under a zero-contact condition but does not construct a measurable selection as the sample varies.

Proposition A.3. *Fix the sampled types and let ℓ_ε be the regularization (20). (i) For every $\varepsilon > 0$ the sampled system with loss ℓ_ε has a unique absolutely continuous solution X^ε on $[0, T]$, and the family $(X^\varepsilon)_{\varepsilon>0}$ is uniformly bounded and uniformly Lipschitz in time. (ii) Every sequence $\varepsilon_n \downarrow 0$ has a subsequence along which X^{ε_n} converges uniformly on $[0, T]$ to an absolutely continuous limit X . (iii) If the contact set $\{t \in [0, T] : X_t^j = 0 \text{ for some } j\}$ has Lebesgue measure zero, then X solves the sampled indicator system, for either threshold convention $\mathbf{1}_{\{x \leq 0\}}$ or $\mathbf{1}_{\{x < 0\}}$, and is therefore a valid pathwise selected solution.*

Proof. (i) The loss ℓ_ε is bounded by one and $(1/\varepsilon)$ -Lipschitz, so the sampled vector field is globally Lipschitz and Picard iteration applies. The field is bounded by $C := |\mu| + |r| \max_i \frac{1}{N} \sum_j |e_{ij}^N - e_{ji}^N| + \max_i \frac{1}{N} \sum_j e_{ij}^N$, uniformly in ε , which gives the uniform bound and the uniform time-Lipschitz constant C . (ii) Arzelà–Ascoli. (iii) Fix t outside the contact set. For each j , $X_t^{\varepsilon_n, j} \rightarrow X_t^j \neq 0$; if $X_t^j > 0$ then $\ell_{\varepsilon_n}(X_t^{\varepsilon_n, j}) = 0$ once $\varepsilon_n < X_t^j/2$ and $|X_t^{\varepsilon_n, j} - X_t^j| < X_t^j/2$, while if $X_t^j < 0$ then $X_t^{\varepsilon_n, j} < 0$ eventually and $\ell_{\varepsilon_n}(X_t^{\varepsilon_n, j}) = 1$. Hence the integrands converge almost everywhere on $[0, T]$ to $\mathbf{1}_{\{X_t^j \leq 0\}}$, which agrees almost everywhere with $\mathbf{1}_{\{X_t^j < 0\}}$; dominated convergence in the integral equation identifies the limit as a solution of the sampled indicator system for either convention. $\square$

If a coordinate remains at the threshold on a set of positive measure, a regularization limit need only solve the corresponding Filippov inclusion. Thus [proposition A.3](#) is a pathwise statement. A measurable selection requires a specified approximation or tie-breaking rule, as discussed in [remark 3.18](#).

APPENDIX B. ADDITIONAL NUMERICAL ROBUSTNESS CHECKS

This appendix supplements [section 6](#) with diagnostics that are conceptually separate from the main deterministic reduction tests. The main text intentionally uses quantile-matched exact low-rank experiments to isolate deterministic approximation error. Here we add three further checks: first, true i.i.d. sampling benchmarks for the discontinuous indicator-loss examples; second, time-step and spatial-grid refinement diagnostics for the numerical solvers; third, an empirical convergence-rate check for the new fixed-rank finite- N indicator theorem.

B.1. True i.i.d. sampling versus quantile matching. For each exact low-rank architecture we sample complete bank types i.i.d. from the corresponding group mixture, simulate the finite- N indicator-loss system at $N = 1600$, and repeat the experiment 200 times. [Figure 10](#) shows one representative trajectory diagnostic together with a summary of the terminal distributions across all four examples. [Table 8](#) reports the corresponding terminal statistics.

TABLE 8. Terminal default fractions under true i.i.d. sampling (200 replications, $N = 1600$) for the exact low-rank indicator-loss examples. The deterministic benchmark is the large-population low-rank limit used in the main text.

Example	Deterministic limit	MC mean	MC median	MC 95% interval
Rank-one generalized mean field	0.221	0.232	0.213	[0.075, 0.532]
Core-periphery	0.283	0.286	0.275	[0.163, 0.467]
Multiple CCPs with overlap	0.470	0.466	0.457	[0.149, 0.744]
Multiplex bank-NBFI network	0.391	0.413	0.390	[0.146, 0.788]




FIGURE 10. Auxiliary Monte Carlo diagnostics for true i.i.d. sampling in the exact low-rank indicator-loss experiments. Left: in the rank-one example, the deterministic limit is close to the Monte Carlo mean trajectory, while the 95% band records substantial finite-sample dispersion. Right: boxplots of the terminal default fraction across the four exact low-rank examples, with the deterministic benchmark marked by diamonds.

The deterministic surrogate represents the large-population benchmark, whereas the i.i.d. experiments quantify finite-sample fluctuations around it. For discontinuous indicator losses, threshold-layer variation remains substantial at the network sizes considered here. The quantile-matched experiments in the main text therefore isolate deterministic approximation error from sampling variability.

When the indicator loss is used directly, the deterministic benchmark should be accompanied by uncertainty quantification. Monte Carlo bands quantify finite-sample dispersion, while a Lipschitz regularization provides a stable point statistic when smoothing is acceptable. The quantile-matched experiments isolate deterministic approximation error, whereas the smoothed graphon experiment illustrates the regularized alternative. In supervisory applications, the deterministic limit should therefore be interpreted as a structural scenario benchmark or as the center of an uncertainty band, not as a point forecast for one sampled network.

B.2. Time-step and spatial-grid refinement. Table 9 collects the basic discretization diagnostics used in the revised numerical section. For the exact low-rank examples, halving the

explicit-Euler step from $\Delta t = 5 \times 10^{-4}$ to $\Delta t = 2.5 \times 10^{-4}$ changes the deterministic default trajectories by at most 4.63×10^{-4} and the quantile-matched finite- N trajectories by at most 1.25×10^{-3} . For the graphon truncation experiment, the corresponding change in the 30-mode reference state is only 3.66×10^{-4} in $\sup_t \|\cdot\|_{L^1}$. We therefore keep $\Delta t = 2 \times 10^{-3}$ in the reported figures.

For the graphon reference, the relevant quantity is the state error, not the visually displayed hard default fraction. The smooth periodic mode construction makes midpoint quadrature substantially more accurate than a naive worst-case $O(U^{-1})$ bound would suggest. The 30-mode reference state differs by only 5.59×10^{-7} in $\sup_t \|\cdot\|_{L^1}$ between the 1200- and 2000-point grids, and by 2.06×10^{-7} between the 2000- and 2400-point grids. These values are orders of magnitude smaller than the smallest reported truncation state error 2.80×10^{-3} at $K = 24$. The larger 10^{-3} -level differences in the displayed hard default fraction are a post-processing effect of thresholding on different grids and do not affect the state metric that the theorems control. For the non-factorized indicator illustration of Section 6.4, the analogous hard-default trajectory changes by at most 5.0×10^{-4} between the 1200- and 2000-point grids, so that figure is likewise not driven by a discretization artifact.

TABLE 9. Discretization diagnostics for the numerical experiments. The exact low-rank rows compare $\Delta t = 5 \times 10^{-4}$ against $\Delta t = 2.5 \times 10^{-4}$. The graphon rows compare both time-step and spatial-grid refinements for the 30-mode reference used in figure 13, and the last row reports the additional grid check for the non-factorized indicator illustration of Section 6.4.

Check	Metric	Value
Exact low-rank limit	$\max \sup_t h_{5 \times 10^{-4}}(t) - h_{2.5 \times 10^{-4}}(t) $ across the four examples	4.63×10^{-4}
Exact low-rank finite $N = 1600$	$\max \sup_t h_{5 \times 10^{-4}}^N(t) - h_{2.5 \times 10^{-4}}^N(t) $ across the four examples	1.25×10^{-3}
Graphon 30-mode reference	$\sup_t \ X_{0.002}(t) - X_{0.001}(t)\ _{L^1}$	3.66×10^{-4}
Graphon 30-mode reference	$\sup_t \bar{\ell}_{0.002}(t) - \bar{\ell}_{0.001}(t) $ for the smoothed distress fraction	2.06×10^{-3}
Grid refinement	$\sup_t \ X_{1200}(t) - X_{2000}(t)\ _{L^1}$	5.59×10^{-7}
Grid refinement	$\sup_t \ X_{2000}(t) - X_{2400}(t)\ _{L^1}$	2.06×10^{-7}
Grid refinement (visual default only)	$\sup_t h_{1200}(t) - h_{2000}(t) $	1.33×10^{-3}
Grid refinement (visual default only)	$\sup_t h_{2000}(t) - h_{2400}(t) $	1.17×10^{-3}
Section 6.4 piecewise-smooth indicator illustration	$\sup_t h_{1200}^{\text{ind}}(t) - h_{2000}^{\text{ind}}(t) $	5.0×10^{-4}

TABLE 10. Per-example time-step sensitivity for the exact low-rank $N = 1600$ benchmarks. The first data column repeats the $N = 1600$ structural benchmark error from [table 3](#); the second and third columns show, respectively, the sup-norm and terminal discrepancies induced by halving the exact low-rank time step from 5×10^{-4} to 2.5×10^{-4} .

Example	benchmark error at $N = 1600$	$\sup_t h_{5 \times 10^{-4}}^N(t) - h_{2.5 \times 10^{-4}}^N(t) $	$ h_{5 \times 10^{-4}}^N(T) - h_{2.5 \times 10^{-4}}^N(T) $
Rank-one generalized mean field	1.64×10^{-3}	1.25×10^{-3}	6.25×10^{-4}
Core-periphery	2.30×10^{-3}	1.25×10^{-3}	6.25×10^{-4}
Multiple CCPs with overlap	1.36×10^{-3}	1.25×10^{-3}	6.25×10^{-4}
Multiplex bank-NBFI network	6.22×10^{-4}	1.25×10^{-3}	0

The per-example comparison in [table 10](#) clarifies why the overall finite- N time-step sensitivity remains visible even after refining the exact low-rank solver. The exact finite- N indicator trajectories are staircase paths with jumps of size $1/N = 6.25 \times 10^{-4}$, so a one- or two-jump shift in threshold-crossing time under step refinement naturally produces a sup-norm discrepancy of order N^{-1} even when the terminal default fraction barely changes. Accordingly, the smallest value in [table 3](#) should not be interpreted as being resolved below the finite- N jump floor; the exact low-rank experiments should instead be read as structural benchmark validation in the variance-reduced deterministic setting.

B.3. Empirical fixed-rank indicator convergence rate. To complement [theorem 3.17](#), we estimate the sampled finite- N feedback error empirically in the rank-one benchmark. For $N \in \{100, 200, 400, 800, 1600, 3200\}$ we run 200 independent i.i.d. replications, compute

$$\hat{e}(N) := \frac{1}{200} \sum_{m=1}^{200} \sup_{0 \leq t \leq T} |\beta_t^{N,(m)} - \beta_t|,$$

and compare the resulting log-log slope against the nearly parametric $N^{-1/2}$ scale predicted by [theorem 3.17](#) for fixed rank. The theorem is a continuous-time statement; the numerical check uses a sufficiently small Euler step, and the time-step diagnostics in [tables 9](#) and [10](#) indicate that the temporal discretization error is dominated by the empirical sampling error over the displayed range.




FIGURE 11. Empirical convergence-rate check for the fixed-rank indicator theorem in the rank-one benchmark. The vertical axis reports the Monte Carlo mean of $\sup_{0 \leq t \leq T} |\beta_t^N - \beta_t|$ over 200 i.i.d. replications at each N ; the shaded band shows the empirical 2.5%–97.5% interval. The fitted log–log slope is close to $-1/2$, consistent with the $\sqrt{\log N/N}$ prediction of [theorem 3.17](#) when K is fixed.

The fitted slope is approximately -0.49 . For fixed $K = 1$, [theorem 3.17](#) gives the scale $N^{-1/2} \sqrt{\log N}$, so the observed slope is consistent with the stated upper bound. This calculation is a numerical diagnostic and is not used in the proof.

APPENDIX C. ADDITIONAL EXAMPLES AND NUMERICAL MATERIAL

This appendix contains additional applications and numerical diagnostics: further interpretations of the factor construction, the multiple-CCP and multiplex experiments, the graphon truncation experiment, and a sensitivity analysis for the transversality margin.

C.1. Beyond banking: other applications. The reduction arguments require only a bounded directed kernel, a bounded loss function, and dense $1/N$ scaling. Formal analogues therefore arise in several other settings. In insurance and reinsurance networks, X may represent a solvency buffer and the kernel may encode ceded-risk or retrocession exposures. In trade-credit supply chains, X may represent working capital and the factors may correspond to dominant tiers or platform intermediaries. In centrally cleared commodity and energy markets, a small number of clearing venues or margin channels yields a factor structure analogous to the multi-CCP model. Domain-specific features such as inventory dynamics, claims seasonality, and physical delivery constraints are not included here and would require separate model extensions.

C.2. Multiple-CCP and multiplex validation. The left panel of [figure 12](#) isolates overlap-induced cross-venue transmission in the multi-CCP example. Under the overlap specification, the terminal default rate of CCP2 specialists is 6.6%, whereas the no-overlap no-dual-members




FIGURE 12. Multiple-CCP and multiplex examples. Left: threshold-fraction paths with overlapping CCP membership and in the no-overlap benchmark. Right: paths for the two-layer bank-NBFI model and the payment-only rank-one benchmark.




FIGURE 13. Graphon truncation beyond the exact low-rank setting. Left: default trajectories for a directed 30-mode reference kernel and its rank- K truncations, with the reference integrated on a dense 2000-point grid. Right: the pathwise state error decays together with the kernel L^1 approximation error, validating the L^1 -stability component of [theorem 4.2](#) and therefore the truncation component of [theorem 4.4](#) under bounded Lipschitz losses; no finite- N sampling step is present in this figure.

benchmark gives only 2.9%. The overlap factors therefore encode the transmission of stress through shared clearing membership.

In the multiplex example, the terminal threshold fraction is 6.5% for the payment-only rank-one benchmark and 39.3% for the two-layer model. The additional funding factor therefore changes the nonlinear feedback materially while preserving a finite-rank representation.

C.3. The graphon truncation experiment. The experiment isolates the L^1 -stability component of [theorem 4.2](#); it contains no finite- N sampling term. The dynamics use the smoothed loss ℓ_ε , and the hard-threshold fraction in the left panel of [figure 13](#) is reported only as a diagnostic. As K increases, both the state trajectories and the kernel converge to the 30-mode reference. The kernel error decreases from 1.03×10^{-1} at $K = 4$ to 2.96×10^{-2} at $K = 12$ and 5.44×10^{-3} at $K = 24$; the corresponding pathwise state errors are 3.22×10^{-2} , 1.35×10^{-2} , and 2.80×10^{-3} . The grid-refinement errors are 5.6×10^{-7} between the 1200- and 2000-point grids and 2.1×10^{-7} between the 2000- and 2400-point grids, well below the reported truncation errors. Thus [table 3](#) isolates finite- N approximation error for exact low-rank structures, whereas [figure 13](#) isolates kernel truncation for a nonfinite-rank reference.

C.4. Sensitivity to the transversality margin. The sufficient condition in [proposition 4.23](#) becomes more demanding as the transversality margin

$$m = m_0 - CA_1 - |r|T(A_1 + A_2)$$

shrinks. We evaluate this dependence numerically by rerunning the non-factorized experiment with the kernel fixed, $r = 0$, $C = T = 2$, and three branchwise slope configurations. In each case the left branch slope exceeds the right branch slope by 0.20, the jump $x_0(\frac{1}{2}+) - x_0(\frac{1}{2}-) = 0.025$ is kept fixed, and the smoothing error is measured at $\varepsilon = 0.01$. Since $A_1 \approx 0.320$, the theorem-level threshold is $CA_1 \approx 0.640$.

TABLE 11. Sensitivity of the non-factorized indicator experiment to the transversality margin. All three regimes remain inside the verified class from [proposition 4.23](#), but the margin $m_0 - CA_1$ becomes progressively smaller.

Regime	m_0	$m_0 - CA_1$	terminal hard default fraction	$\sup_{0 \leq t \leq T} \ X_t^\varepsilon - X_t^{\text{ind}}\ _{L^1}$ at $\varepsilon = 0.01$
Baseline	1.75	1.11	0.2195	2.35×10^{-3}
Moderate	0.90	0.26	0.4915	4.86×10^{-3}
Near-critical	0.70	0.06	0.6035	5.34×10^{-3}

As the margin decreases, the discrepancy between the indicator and regularized dynamics increases at fixed ε ; see [table 11](#). This agrees with the dependence of the sufficient density bound on J/m . The observed increase is smaller than the worst-case J/m scaling, which need not be sharp and does not account for saturation on $\{X_s \leq 0\}$. Cases with $m_0 \leq CA_1$ are omitted because they fall outside the hypotheses of [proposition 4.23](#).

REFERENCES

- [1] I. Aldasoro and I. Alves, *Multiplex interbank networks and systemic importance: An application to European data*, BIS Working Paper No. 603, 2017.
- [2] Basel Committee on Banking Supervision, *Banks' interconnections with non-bank financial intermediaries*, Bank for International Settlements, 10 July 2025.
- [3] S. Allmeier and N. Gast, *Accuracy of the graphon mean field approximation for interacting particle systems*, Stochastic Systems **15** (2025), no. 4, 273–290.
- [4] H. Amini, R. Cont, and A. Minca, *Resilience to contagion in financial networks*, Mathematical Finance **26** (2016), no. 2, 329–365.
- [5] H. Amini, Z. Cao, and A. Sulem, *Limit theorems for default contagion and systemic risk*, Mathematics of Operations Research **49** (2024), no. 4, 2652–2683.
- [6] T. Banerjee, A. Bernstein, and Z. Feinstein, *Dynamic clearing and contagion in financial networks*, European Journal of Operational Research **321** (2025), no. 2, 664–675.

- [7] E. Bayraktar and R. Wu, *Stationarity and uniform in time convergence for the graphon particle system*, Stochastic Processes and their Applications **150** (2022), 532–568.
- [8] E. Bayraktar and R. Wu, *Graphon particle system: Uniform-in-time concentration bounds*, Stochastic Processes and their Applications **156** (2023), 196–225.
- [9] E. Bayraktar, S. Chakraborty, and R. Wu, *Graphon mean field systems*, Annals of Applied Probability **33** (2023), no. 5, 3587–3619.
- [10] E. Bayraktar and H. Zhou, *Non-parametric estimates for graphon mean-field particle systems*, Bernoulli **31** (2025), no. 4, 2940–2961.
- [11] M. Boss, H. Elsinger, M. Summer, and S. Thurner, *Network topology of the interbank market*, Quantitative Finance **4** (2004), no. 6, 677–684.
- [12] R. Carmona, J.-P. Fouque, and L.-H. Sun, *Mean field games and systemic risk*, Communications in Mathematical Sciences **13** (2015), no. 4, 911–933.
- [13] R. Cont, A. Moussa, and E. B. Santos, *Network structure and systemic risk in banking systems*, in *Handbook on Systemic Risk*, Cambridge University Press, Cambridge, 2013, pp. 327–368.
- [14] F. Coppini, A. De Crescenzo, and H. Pham, *Nonlinear graphon mean-field systems*, Stochastic Processes and their Applications **190** (2025), 104728.
- [15] B. Craig and G. von Peter, *Interbank tiering and money center banks*, Journal of Financial Intermediation **23** (2014), no. 3, 322–347.
- [16] N. Detering, T. Meyer-Brandis, K. Panagiotou, and D. Ritter, *An integrated model for fire sales and default contagion*, Mathematics and Financial Economics **15** (2021), 59–101.
- [17] C. Crucianelli and L. Tangpi, *Interacting particle systems on sparse W -random graphs*, arXiv:2410.11240, 2024.
- [18] L. Eisenberg and T. H. Noe, *Systemic risk in financial systems*, Management Science **47** (2001), no. 2, 236–249.
- [19] M. Elliott, B. Golub, and M. O. Jackson, *Financial networks and contagion*, American Economic Review **104** (2014), no. 10, 3115–3153.
- [20] S. Erol, F. Parise, and A. Teytelboym, *Contagion in graphons*, Journal of Economic Theory **211** (2023), 105673.
- [21] P. Feng, *Dynamic Network Model for Systemic Risk*, Ph.D. dissertation, University of Southern California, Applied Mathematics, August 2021. Available at <https://digitallibrary.usc.edu/C.aspx?VP3=pdfviewer&rid=2A3BF1SA6QN8H>.
- [22] Z. Feinstein and A. Sojmark, *Short communication: Dynamic default contagion in heterogeneous interbank systems*, SIAM Journal on Financial Mathematics **12** (2021), no. 4, SC83–SC97.
- [23] N. Fournier and A. Guillin, *On the rate of convergence in Wasserstein distance of the empirical measure*, Probability Theory and Related Fields **162** (2015), no. 3–4, 707–738.
- [24] A. W. van der Vaart and J. A. Wellner, *Weak Convergence and Empirical Processes*, Springer, New York, 1996.
- [25] P. Gai and S. Kapadia, *Contagion in financial networks*, Proceedings of the Royal Society A **466** (2010), no. 2120, 2401–2423.
- [26] P. Glasserman and H. P. Young, *How likely is contagion in financial networks?*, Journal of Banking & Finance **50** (2015), 383–399.
- [27] L. Lovasz, *Large Networks and Graph Limits*, American Mathematical Society, Providence, RI, 2012.
- [28] S. Nadtochiy and M. Shkolnikov, *Mean field systems on networks, with singular interaction through hitting times*, Annals of Probability **48** (2020), no. 3, 1520–1556.
- [29] L. A. M. Veraart and I. Aldasoro, *Systemic risk in markets with multiple central counterparties*, Mathematical Finance **35** (2025), no. 1, 214–262.
- [30] European Banking Authority, *2025 EU-wide transparency exercise* (Autumn 2025), risk assessment report and bank-level disclosure files, reference dates September 2024–June 2025. Sovereign template `tr_sov.csv` and dictionary `TR_Metadata.xlsx`; item 2520810 (direct on-balance-sheet total gross carrying amount of non-derivative financial assets). Available at <https://www.eba.europa.eu/risk-and-data-analysis/risk-analysis/eu-wide-transparency-exercise>; accessed 7 July 2026.