
Article

Faster Monotone Implied Volatility Solver

Fabien Le Floc’h 

Independent researcher; fabien@2ipi.com

Abstract

We present ThiopheneIV, a Black–Scholes implied-volatility solver with a monotone core and explicit production guards. Prices are first reduced to Jäckel’s out-of-the-money normalisation and inverted through a tail-stable logarithmic price equation. The solver starts from the non-iterative Choi–Huh–Su L3 lower-bound seed and applies three Euler–Chebyshev corrections. In exact arithmetic, the seed is below the admissible root and the Euler–Chebyshev map increases monotonically without overshooting; the proof is included. The implementation then adds the floating-point machinery needed in practice: parity normalisation, microscopic Bachelier-limit handling, saturated-price treatment, finite-update checks, fallback seeds, and an optional Jäckel–Newton polish. Against the highly accurate expanded Jäckel reference price, ThiopheneIV is faster than a normalised Java port of Jäckel’s *Let’s Be Rational* while keeping regular-grid errors close. ThiopheneIV+ adds one final Jäckel–Newton correction for systems that need closer agreement with that expanded reference price. The broader lesson is that a convergence proof gives a clean core, but robust production inversion still depends on boundary handling and on the pricing objective one chooses to match.

Keywords: implied volatility; Black–Scholes; Chebyshev method; Choi–Huh–Su bounds; Jäckel; erfcx; root-finding; computational finance

1. Introduction

Implied volatility inversion recovers the Black–Scholes volatility σ from an option price. It is a small scalar root-finding problem, but it is called so often in calibration and risk systems that both latency and tail robustness matter. The numerical difficulty is not merely local convergence of Newton-like iterations. Deep out-of-the-money prices can underflow, nearly in-the-money prices can lose time value before the solver is called, and high-volatility prices close to the upper bound are poorly conditioned in volatility.

Jäckel’s *Let’s Be Rational* [3] remains the natural reference implementation for this domain. It combines normalised prices, region-dependent asymptotics, complementary objectives, and carefully selected iterations. Choi, Huh, and Su [5] derive implied-volatility bounds from option-delta inequalities and prove monotone convergence of Newton’s method from their lower bounds.

This paper studies a production solver built around the Choi–Huh–Su L3 lower bound. We call it ThiopheneIV. The solver uses the L3 formula as a global non-Bachelier seed and then applies three Euler–Chebyshev corrections to the log-price objective. The update is the Euler–Chebyshev method of order three ($\alpha = 0$ in Traub’s notation [9, p. 81]):

$$v_{n+1} = v_n - \frac{g(v_n)}{g'(v_n)} - \frac{g''(v_n)[g(v_n)]^2}{2[g'(v_n)]^3}.$$

Starting from the Choi L3 lower-bound seed, this map is monotone increasing and bounded above by the true root in exact arithmetic (Appendix D).

The contribution is therefore twofold. First, the paper records an explicit normalised implementation of the L3 seed and proves monotone convergence of the Euler–Chebyshev refinement from that seed. Second, it documents the guards needed to turn that mathematically clean iteration into a robust double-precision solver: converting every quote to an equivalent out-of-the-money price, using a tail-safe logarithmic Black-price

formula, switching to a local Bachelier approximation for microscopic near-at-the-money prices, guarding prices close to the no-arbitrage upper bound, rejecting non-finite updates, and adding an optional Jäckel–Newton polish when the target is ulp-level agreement with the expanded Jäckel reference price. The focus is ThiopheneIV and its comparison with Jäckel’s reference solver.

2. Problem Formulation

2.1. Normalization

All prices are first reduced to the same normalised OTM representation. Let C and P denote undiscounted call and put prices with forward F , strike K , and time to expiry T . We call an option at the money (ATM) when $F = K$, and near ATM when $|\ln(F/K)|$ is small. A call is out of the money (OTM) when $F < K$, while a put is OTM when $F > K$; the opposite cases are in the money (ITM). Rather than carrying separate call, put, ITM, and OTM formulae through the solver, we order the forward–strike pair as

$$F_* = \min(F, K), \quad K_* = \max(F, K), \quad \frac{F_*}{K_*} = \min(F/K, K/F) \leq 1.$$

The corresponding OTM value is obtained by put–call parity, and by exchanging the forward and strike for OTM puts:

$$C_{\text{OTM}} = \begin{cases} C, & \text{call and } F \leq K, \\ C - (F - K), & \text{call and } F > K, \\ P, & \text{put and } F > K, \\ P - (K - F), & \text{put and } F < K. \end{cases}$$

Thus every admissible input is represented as an undiscounted OTM call on (F_*, K_*) . The variables passed to the scalar inversion are

$$x = \ln\left(\frac{F_*}{K_*}\right) \leq 0, \quad e^x = \frac{F_*}{K_*}, \quad c = \frac{C_{\text{OTM}}}{F_*}, \quad v = \sigma\sqrt{T}. \quad (1)$$

This normalisation removes intrinsic value before inversion. In particular, originally ITM options are priced through their OTM parity legs, avoiding the cancellation that would occur if one subtracted a large intrinsic component from the original option price.

Here and throughout, Φ denotes the standard normal cumulative distribution function.

In these coordinates the Black formula for the normalised OTM call price is

$$c(x, v) = \Phi\left(\frac{x}{v} + \frac{v}{2}\right) - e^{-x}\Phi\left(\frac{x}{v} - \frac{v}{2}\right). \quad (2)$$

2.2. Log-Price Objective and erfcx Decomposition

Following Jäckel [3] and Choi et al. [5], the iteration is performed in log-price space rather than on the raw price residual. For a target normalised price c_{target} , the objective is

$$f(v) = \ln(c(x, v)) - \ln(c_{\text{target}}). \quad (3)$$

Proposition 1. For $h = x/v$ and $t = v/2$, the log-price admits the representation

$$\ln(c) = -\frac{1}{2}(h^2 + t^2) - \ln(2) - \frac{x}{2} + \ln(N^+ - N^-), \quad (4)$$

where

$$N^+ = \text{erfcx}\left(-(h + t)/\sqrt{2}\right), \quad N^- = \text{erfcx}\left(-(h - t)/\sqrt{2}\right). \quad (5)$$

Proof. Using (2) and $\Phi(z) = \frac{1}{2} \operatorname{erfc}(-z/\sqrt{2})$ gives

$$c = \frac{1}{2} [\operatorname{erfc}(-(h+t)/\sqrt{2}) - e^{-x} \operatorname{erfc}(-(h-t)/\sqrt{2})].$$

Now write $\operatorname{erfc}(z) = e^{-z^2} \operatorname{erfcx}(z)$. Since $ht = x/2$,

$$e^{-(h+t)^2/2} = e^{-(h^2+t^2)/2-x/2}, \quad e^{-x} e^{-(h-t)^2/2} = e^{-(h^2+t^2)/2-x/2}.$$

The two terms therefore share the common factor $e^{-(h^2+t^2)/2-x/2}/2$. Factoring it out and taking logarithms gives (4). $\square$

Remark 1. The scaled complementary error function $\operatorname{erfcx}(z) = e^{z^2} \operatorname{erfc}(z)$ is bounded and positive for all real z . Consequently, the difference $N^+ - N^-$ remains computable even when the price itself would underflow to zero in double precision. This is the main reason for using the log-price formulation: the objective $f(v)$ remains well defined and smoothly differentiable for arbitrarily deep OTM options.

This improvement should not be confused with a guarantee of strict monotonicity in floating-point arithmetic. The real Black price is monotone in total volatility, but a particular double-precision evaluation need not be monotone when the input is advanced by one ulp (unit in the last place, the spacing between adjacent double-precision numbers at the value under discussion); this distinction is highlighted in [4]. Figure 1 repeats this diagnostic for $F_*/K_* = 0.955$ and an initial total volatility $v = 0.05$, using 512 successive calls to `nextUp`. The price variable is the same forward-normalised OTM price $c = C_{\text{OTM}}/F_*$ used by the solver API; at the start of the sweep $c \simeq 4.94 \times 10^{-3}$, so this is a small but not deep-tail option price. The plot reports the price increment divided by the local double-precision spacing of the previous price. Both the `erfcx`/log formula and the expanded Jäckel reference price from Appendix B have downward one-ulp steps in this case. The expansion is therefore valuable for reducing cancellation, but it should not be read as a floating-point monotonicity guarantee. This diagnostic is also a reminder that once the pricing formula itself reaches its double-precision evaluation limit, further root iterations cannot recover information that is no longer present in the objective value.




Figure 1. One-ulp price monotonicity diagnostic for $F_*/K_* = 0.955$ and $v = 0.05$. Each curve shows the local OTM-price step produced by advancing the total volatility by one double-precision value. Negative points mark local downward steps despite the monotonicity of the real-valued Black price.

3. Solver Design

3.1. Log-Price Derivatives

ThiopheneIV refines total volatility with the same log-price objective as Section 2.2. Let

$$g(v) = \ln(c(x, v)) - \ln(c_{\text{target}}), \quad h = x/v, \quad t = v/2.$$

The derivatives needed by the Euler–Chebyshev correction follow from the same erfcx decomposition used to evaluate the objective.

Proposition 2. *In the log-price formulation, the first three derivatives of $g_0(v) = \ln(c(x, v))$ admit closed-form expressions:*

$$g'_0(v) = \frac{2/\sqrt{2\pi}}{N^+ - N^-}, \quad (6)$$

$$\frac{g''_0(v)}{g'_0(v)} = \frac{(h+t)(h-t)}{v} - g'_0(v), \quad (7)$$

$$\frac{g^{(3)}_0(v)}{g'_0(v)} = \frac{-3h^2 - t^2 + (h^2 - t^2)^2}{v^2} - 3g'_0(v) \frac{g''_0(v)}{g'_0(v)} - [g'_0(v)]^2. \quad (8)$$

Proof. The vega of the normalised Black formula is $\partial c / \partial v = \phi(d_1)$, where ϕ is the standard normal density. Thus $g'_0(v) = c^{-1} \phi(d_1)$. Expressing $\phi(d_1) = (2\pi)^{-1/2} e^{-d_1^2/2}$ and using Equation (5) gives (6); Equations (7) and (8) follow by differentiating with respect to v and simplifying. $\square$

Once N^+ and N^- are available from the objective evaluation, the derivative ratios involve only elementary arithmetic. The Chebyshev step therefore costs little more than a Newton step with the same erfcx calls, but has cubic local convergence at the simple root.

3.2. Choi–Huh–Su L3 Seed

For $x \leq 0$, set $k = -x \geq 0$ and $E = e^k$. The Choi–Huh–Su L3 seed maps the normalised OTM price to

$$p_3 = \frac{c(c+E)}{2c+E-1}, \quad z_3 = \Phi^{-1}(p_3), \quad (9)$$

and then solves $z_3 = d_1(v) = -k/v + v/2$ for a positive total volatility. The root is evaluated in rationalised form when $z_3 < 0$ to avoid cancellation; Appendix C gives the exact transcription, including the ATM limit.

The seed is not selected because it is the most accurate raw approximation. Its advantage is structural: Choi et al. [5] prove that this value is a lower bound for the admissible implied volatility. ThiopheneIV uses that lower-side property as the entry point for a monotone refinement rather than as a merely empirical basin guess. The inverse normal CDF is evaluated by a piecewise rational approximation in the Java implementation; subsequent Chebyshev steps and the optional Jäckel–Newton polish absorb the last-bit differences introduced by that approximation.

3.3. Euler–Chebyshev Refinement

Given a finite positive seed v_n below the root, define

$$\eta_n = -\frac{g(v_n)}{g'(v_n)}, \quad q_n = \frac{g''(v_n)}{g'(v_n)}.$$

ThiopheneIV applies the Euler–Chebyshev method of order three ($\alpha = 0$ in Traub’s notation [9, p. 81]) in the compact form

$$v_{n+1} = v_n + \eta_n \left(1 - \frac{1}{2} q_n \eta_n \right). \quad (10)$$

The production path uses three full-precision steps. Appendix D proves that, in real arithmetic, the map in (10) satisfies $0 < T(v) - v \leq v_* - v$ for every $v \in (0, v_*)$ when $x \leq 0$. Therefore the sequence from the Choi L3 lower-bound seed is monotone increasing, bounded by the true root, and convergent.

3.4. Optional Jäckel–Newton Polish

The default ThiopheneIV objective is the erfcx/log-price formula. When the goal is instead to match the expanded Jäckel reference price as closely as possible, one final Newton correction is applied in beta space. The correction uses the reference price from Appendix B and the analytic normalised vega. Equivalently, it asks how much the current total volatility must move so that repricing with that reference gives the same sqrt-forward-normalised price as the input. The update is accepted only when it is finite and positive. It is deliberately optional: it improves ulp-level agreement with the expanded Jäckel reference price, but it adds roughly 50–65 ns per call in the present Java benchmark. This last correction is useful only when the surrounding system also treats that reference price as the target. Many production systems instead use a straightforward Black–Scholes formula built from a library normal CDF; in such systems the pricing formula itself is typically less accurate than the expanded Jäckel reference price, so the polish may provide no practical benefit.

3.5. Guards and Hot Path

Algorithm 1 summarises the production path. The same guards are also useful for any Choi-seeded or Jäckel-like solver that is expected to handle adversarial inputs rather than only smooth benchmark grids.

Algorithm 1 ThiopheneIV hot path.

Require: $x \leq 0, e^x$, normalised OTM price $0 < c < 1, \ln(c)$, maturity T

```

1: if  $c \leq 10^{-6}$  and  $|x| \leq 10^{-8}$  then
2:   return Bachelier-limit volatility (scaled Mills-ratio tail, Bachelier rational seed plus expansion polish,
   or defensive zero)
3: end if
4: Compute the Choi–Huh–Su L3 seed; clamp probabilities to the open double-precision interval
5: if the L3 seed is not finite and positive then
6:   Replace it with a guarded asymptotic or ATM fallback seed
7: end if
8:  $v_0 \leftarrow \max(v_0, 10^{-10})$ 
9: if  $c \geq 0.9990$  then
10:   $v_0 \leftarrow \max(v_0, \text{upper-price asymptotic seed})$ 
11:  return complementary upper-price polish initialized at  $v_0$ 
12: end if
13: for  $i = 0, 1, 2$  do
14:   Compute one full-precision Euler–Chebyshev step on the erfcx/log objective
15:   Accept the update only if it is finite and positive
16: end for
17: if expanded-Jäckel-reference matching is requested then
18:   Apply one Jäckel–Newton polish in beta space
19: end if
20: return  $v_3/\sqrt{T}$ 

```

4. Experimental Setup

4.1. Reference Price Computation

Benchmarking a solver to machine precision requires reference prices accurate to the last bit. The textbook Black–Scholes formula $c = \Phi(d_1) - e^{-x}\Phi(d_2)$ suffers from catastrophic cancellation for deep OTM options, and even the erfcx/log objective used by the unpolished solver can lose relative bits in nearly-ATM, very-low-volatility cases because it subtracts two close erfcx values. Therefore the accuracy tables and accuracy figures use Jäckel’s cancellation-avoiding expanded normalised-Black evaluation of Appendix B as the double-precision reference. We call this the *expanded Jäckel reference price*. For each reference total volatility, this produces a

reference OTM-normalised price, which is passed to the solver in the same $c = C_{\text{OTM}}/F_*$ convention as any other input.

The $\text{erfcx}/\log$ expression remains the central ThiopheneIV iteration objective:

$$c(x, v) = \frac{1}{2} e^{-(h^2+t^2)/2-x/2} [\text{erfcx}(-(h+t)/\sqrt{2}) - \text{erfcx}(-(h-t)/\sqrt{2})], \quad (11)$$

and, for extreme OTM cases where the price itself would underflow, the iteration works entirely in log space via (4). ITM prices are recovered via put–call parity (a well-conditioned addition of intrinsic value), never by direct evaluation.

4.2. Test Datasets

To ensure robustness, we benchmark on eight datasets spanning the full range of practically relevant option parameters (Table 1).

Table 1. Test datasets.

Dataset	Cases	Description
CLY-3D	51,321	Cui–Liu–Yao three-dimensional grid: $K \in [105, 800]$, $T \in [0.01, 2]$, $\sigma \in [0.01, 0.99]$, filtered at price 10^{-20}
CLY-20	1,600	Cui–Liu–Yao fixed-volatility surface: $\sigma = 20\%$, $K \in [105, 180]$, $T \in [0.1, 2]$
CLY-80	1,600	Cui–Liu–Yao fixed-volatility surface: $\sigma = 80\%$, $K \in [105, 800]$, $T \in [0.1, 2]$
Jäckel	5,182	Wide moneyness: $K/F \in [0.5, 8]$, σ up to 4.0
Market	7,151	Realistic: $K/F \in [0.7, 1.5]$, T from 1/252 to 5 yr
Corners	278	Edge cases: low-vol/short-mat, high-vol/deep-OTM, near-ATM small-price, and saturated upper-price cases
Stress	1,270	Extremes: K/F up to 100 $\times$, $T \in [0.001, 10]$
HighVol	149	Fallback stress zone: $ x \geq 3$, $c \in (0.05, 0.95)$, σ up to 2.5

The first three datasets reproduce the comparison grids from Cui–Liu–Yao [2]: the main three-dimensional grid and the two fixed-volatility surface grids used in their numerical comparison with previous literature. The HighVol dataset specifically targets the region $|x| \geq 3$ with large option prices, where Li’s rational approximation is outside its fitted domain and the asymptotic OTM guard only accepts part of the set; cases are filtered to $c < 0.95$ to reflect the practical use of put–call parity for deep ITM options. Appendix A gives the exact construction rules used by the benchmark generator.

4.3. Solvers Under Comparison

The main comparison is between ThiopheneIV and a normalised Java port of Jäckel’s *Let’s Be Rational*. The benchmark harness is the same as for the companion solvers: minimum of 500 sweeps, 3 independent runs on OpenJDK 21.

- **Jäckel’s *Let’s Be Rational* [3].** Region-dependent asymptotic expansions, log-space iteration, and a complementary objective. The Java port uses the normalised API with $\beta = c\sqrt{F/K}$ and the original two-iteration default. It runs at 238 ns on CLY-3D in the same scalar benchmark harness.
- **ThiopheneIV.** Choi–Huh–Su L3 seed, three full-precision Euler–Chebyshev steps, production guards, and optional Jäckel–Newton polish. We denote the polished configuration by **ThiopheneIV+**: ThiopheneIV with one final Jäckel–Newton polish. The unpolished path runs at 165 ns on CLY-3D; ThiopheneIV+ runs at 225 ns.

5. Results

5.1. Accuracy and Timing

Table 2 reports accuracy and latency on the eight benchmark datasets. Input prices for the accuracy block are generated from the expanded Jäckel reference price at the known total volatility v_{ref} , then passed to the

solver as OTM-normalised prices. Errors are measured in ulps of the reference total volatility. For each test case, if the reference total volatility is v_{ref} and the solver returns $\hat{v} = \hat{\sigma}\sqrt{T}$, the reported per-case error is

$$\frac{|\hat{v} - v_{\text{ref}}|}{\text{nextUp}(v_{\text{ref}}) - v_{\text{ref}}}.$$

Thus the unit is the local double-precision spacing at the reference volatility, not a fixed decimal tolerance. For example, at $v_{\text{ref}} = 0.20$, one ulp is $2.7755575615628914 \times 10^{-17}$; an 8-ulp error corresponds to an absolute total-volatility error of about 2.22×10^{-16} , or a relative error of about 1.11×10^{-15} . Table 2 reports the maximum of this quantity over each dataset. This makes the accuracy table a comparison against the expanded Jäckel reference price rather than against the simpler erfcx/log formula.

Table 2. Accuracy against the expanded Jäckel reference price and latency by dataset.

	CLY-3D	CLY-20	CLY-80	Jäckel	Market	Corners	Stress	HighVol
Accuracy – max error (ulp of reference total volatility)								
Jäckel	7	4	5	18	9	584	5	2
ThiopheneIV	119	48	6	148	212	607	180	7
ThiopheneIV+	8	3	4	20	10	607	5	3
ThiopheneIV+ accuracy – max absolute total-volatility error								
ThiopheneIV+	6.7×10^{-16}	8.3×10^{-17}	4.4×10^{-16}	1.8×10^{-14}	1.8×10^{-15}	5.4×10^{-13}	8.9×10^{-16}	2.7×10^{-15}
Latency (ns/call)								
Jäckel	238	229	229	224	204	268	238	208
ThiopheneIV	165	170	164	172	177	181	175	167
ThiopheneIV+	225	220	219	234	232	238	238	226

Unpolished ThiopheneIV is faster than the Java port of Jäckel’s solver on all eight datasets in this run: 165 ns versus 238 ns on CLY-3D, and 167 ns versus 208 ns on HighVol. The optional Newton polish is still competitive with Jäckel’s two-iteration reference path on CLY-3D and the regular CLY surfaces, but it is no longer a pure low-latency configuration; its purpose is reference-price alignment. It makes the output closer to the expanded Jäckel reference price, especially on the regular datasets. The Corners ulp count remains large because the dataset contains saturated and near-zero-vega cases where a tiny price difference corresponds to many ulps of total volatility; the absolute-error row shows that the ThiopheneIV+ maximum there is 5.4×10^{-13} in total volatility.

Figure 2 gives fixed-log-moneyness slices against Jäckel’s normalised solver. It checks that ThiopheneIV remains in the same accuracy band as the Jäckel reference on the diagnostic slices, and shows how the optional ThiopheneIV+ polish collapses most of the remaining reference-price residual.

The separation between ThiopheneIV and ThiopheneIV+ in the first panel ($x = -10^{-8}$, $v \lesssim 10^{-3}$) is the same kind of pricing-formula limit highlighted by Figure 1, not a sign that the Choi seed or the three Euler–Chebyshev steps have failed to converge. The input prices in the figure are generated by the expanded Jäckel reference price. Unpolished ThiopheneIV then solves the erfcx/log objective, whose two erfcx terms are nearly equal in this near-ATM, very-small-total-variance corner. At this scale the erfcx/log formula has already lost enough relative bits that its root is slightly different from the expanded-Jäckel-reference root. The resulting relative price mismatch is typically 10^{-10} – 10^{-12} in this panel, which appears as about 10^{-12} relative volatility error. Jäckel and ThiopheneIV+ both finish by matching the expanded Jäckel reference price, so their curves lie near the double-precision floor.

Figure 3 resolves the same style of fixed- x slices by iteration stage. The Choi L3 seed starts on the lower side of the root, and the three Euler–Chebyshev steps compress the error to the double-precision floor across the diagnostic grid.

5.2. Jäckel–Newton Polish Diagnostic

Figure 4 shows locally the same effect reported in Table 2: the polish reduces the residual of the expanded Jäckel reference price, with the incremental cost visible in the latency block of the table.

The small non-zero mean of the ThiopheneIV+ curve in the left panel is a floating-point effect, not evidence that the polished method remains a lower bound. After the final Newton polish, the sign is dominated




Figure 2. Fixed- x comparison of ThiopheneIV and ThiopheneIV+ against Jäckel’s implementation. Each panel varies the reference total volatility at fixed log-moneyness and reports relative total-volatility error.

by rounding, objective evaluation mismatch, and ulp quantisation; vega maps the remaining few-volatility-ulp bias into sub-ulp price changes, so the repriced residual can look mean-zero.

5.3. Operation Costs

Table 3. Micro-benchmarked operation costs relevant to ThiopheneIV.

Operation	Cost (ns)
Thiophene L3 guess	20.0
1 exact erfcx (Boost/Commons)	5.5
Natural logarithm	4.8
Exponential	3.7
Square root	0.7
Optional Jäckel–Newton polish (incremental)	about 50–65

The Choi L3 seed is compact, but the three exact Euler–Chebyshev steps dominate the hot path through exact erfcx and logarithm evaluations. This explains why ThiopheneIV is faster than the Jäckel Java port while still spending most of its latency budget on the proof-backed refinement.




Figure 3. Fixed- x convergence slices for the Choi L3 seed and the Euler-Chebyshev chain. Each panel fixes log-moneyness and varies the reference total volatility. Curves show relative total-volatility error after the seed and after one, two, and three full-precision Euler-Chebyshev steps; values are floored at 10^{-18} only for log-scale plotting.




Figure 4. Local round-trip diagnostics for the optional Jäckel-Newton polish. The left panel measures implied-volatility error when the input price is generated by the expanded Jäckel reference price; the right panel measures the relative price residual after inversion and repricing with the same reference formula.

6. Discussion

The proof-backed part of ThiopheneIV is deliberately narrow and explicit. In real arithmetic the Choi L3 seed starts on the lower side of the root, the log-price objective is increasing, and the Euler–Chebyshev map does not overshoot. This gives a clean monotone-convergence story that is absent from a purely empirical H3 chain.

The engineering story is less clean, and that is the important practical point. A double-precision solver has to decide what to do when an inverse-normal probability is rounded to the endpoint, when a candidate update is non-finite, when the price is so close to the no-arbitrage upper bound that volatility is poorly conditioned, or when a microscopic near-ATM price is better represented by a Bachelier-limit expansion than by the Black tails themselves. These branches do not contradict the convergence theorem; they protect the preconditions under which the theorem and the floating-point evaluation are useful.

Jäckel’s solver embodies a similar lesson from another direction. Its robustness comes from normalisation, asymptotic regions, complementary objectives, and careful iteration choices. ThiopheneIV replaces the central seed-and-refine mechanism with Choi L3 plus monotone Chebyshev steps, but it still needs comparable boundary awareness. The optional Jäckel–Newton polish is another example: convergence to the $\text{erfcx}/\log$ objective is already obtained, yet matching a different high-accuracy pricing objective to a few ulps requires one more targeted correction.

The benchmark timings are scalar Java timings, not universal constants. A faster erfcx implementation, a vectorised batch path, or a different inverse normal CDF would shift the absolute numbers. The qualitative trade-off should be more portable: a monotone seed-refine pair buys mathematical transparency, while production robustness and final-reference matching add guard and polish costs that must be budgeted explicitly.

7. Conclusions

ThiopheneIV combines the Choi–Huh–Su L3 lower-bound seed with three Euler–Chebyshev corrections on a normalised $\text{erfcx}/\log$ -price objective. In exact arithmetic, the Choi seed and the Euler–Chebyshev map give monotone convergence from below to the admissible Black–Scholes implied volatility. In the Java 21 scalar benchmark, the unpolished implementation has low-double-digit to few-hundred-ulp maximum errors against the expanded Jäckel reference price and runs at 165 ns on CLY-3D, versus 238 ns for the normalised Java port of Jäckel’s *Let’s Be Rational*. With the optional Jäckel–Newton polish, the CLY-3D timing is 225 ns and the regular-grid errors are reduced to low ulp counts against the expanded Jäckel reference price. This polish is therefore an alignment option, not a universal improvement: if a production stack prices with the textbook normal-CDF Black–Scholes formula rather than the expanded Jäckel reference price, the extra polish can be below the accuracy floor of the surrounding pricing code.

The main conclusion is not that a convergence proof makes implied-volatility solving simple. The proof guarantees the mathematical iteration once its preconditions are met. A robust production solver still needs many additional steps: normalising quotes to equivalent out-of-the-money prices, evaluating tail prices safely, guarding probabilities and updates, handling microscopic near-at-the-money prices, treating prices close to the no-arbitrage upper bound, and sometimes applying a final polish to machine-epsilon agreement with a chosen reference price. Writing such a solver is therefore less straightforward than “start from a guaranteed seed and iterate”, even when the real-arithmetic convergence is settled. The value of ThiopheneIV is that it separates these two issues: a transparent monotone core, surrounded by the necessary engineering safeguards.

Appendix A. Benchmark Construction and Worked Examples

Appendix A.1. Normalisation and dataset construction

All generated benchmark cases are converted to the same normalised OTM representation before being passed to the solvers. Given spot S , strike K , maturity T , volatility σ , and rate r , the Java helper computes

$$F = Se^{rT}, \quad F_* = \min(F, K), \quad K_* = \max(F, K), \quad e^x = F_*/K_*,$$

and then

$$x = \ln(e^x), \quad v_{\text{ref}} = \sigma\sqrt{T}.$$

For accuracy reporting and fixed-slice figures, the stored price is the OTM-normalised price obtained from the expanded Jäckel reference price of Appendix B. For the raw dataset filters, the corresponding $\ln(c)$ is required to be finite, no larger than -10^{-15} (effectively at the upper price boundary), no smaller than -708 (double underflow), and $c \in (0, 1)$. The benchmark array stores (c, e^x, T, σ) ; all six generated datasets use calls, with ITM calls represented through the OTM parity leg.

Appendix A.2. Concrete worked examples

For example, the near-ATM small-price Corners case $S = 100, K = 100.5, T = 0.01, \sigma = 0.05, r = 0$ gives

$$\begin{aligned} F &= 100, & e^x &= 100/100.5 = 0.9950248756218906, \\ x &= -0.004987541511039051, & v_{\text{ref}} &= 0.005000000000000001, \\ \ln(c) &= -7.776203975967922, & c &= 4.196019744216237 \times 10^{-4}. \end{aligned}$$

It is therefore stored as a valid Corners case and enters the near-ATM small-price seed branch ($c \leq 5 \times 10^{-4}$ and $|x| < 0.01$), not the microscopic Bachelier branch ($c \leq 10^{-6}$ and $|x| \leq 10^{-8}$).

A complementary saturated-price corner illustrates the conditioning at high volatility close to ATM. With $S = 100, K = 100.01, T = 10, \sigma = 3$, and $r = 0$, one obtains

$$\begin{aligned} e^x &= 0.9999000099990001, & x &= -9.999500033332494 \times 10^{-5}, \\ v_{\text{ref}} &= 9.486832980505138, & \ln(c) &= -2.1015432345450336 \times 10^{-6}, \\ c &= 0.9999978984589737, & 1 - c &= 2.1015410263114376 \times 10^{-6}. \end{aligned}$$

This case is near ATM but enters the upper-price branch because $c \geq 0.9990$. In this regime, a guarded production solver can still reprice to machine-precision price accuracy even when the returned volatility differs from the generating value by more than a few ulps. The larger error in volatility is unavoidable conditioning: the normalised vega is only 5.19×10^{-6} , so an absolute price perturbation at double precision is amplified when inverted for v .

Appendix A.3. Microscopic Bachelier-limit branch

The microscopic branch is deliberately narrower than the ordinary near-ATM small-price seed. It is entered only for normalised OTM prices $c \leq 10^{-6}$, $|x| \leq 10^{-8}$, and $x \leq 0$ (with a one-part-in 10^{12} tolerance on the price cut-off). In this box the two Black normal terms are almost equal and the volatility is of the same order as the log-moneyness. Iterating directly on the Black formula can therefore spend its effort resolving a difference of nearly equal tail quantities, while a central ATM approximation can be wrong by orders of magnitude as soon as $|x|/v$ is no longer small.

The branch first removes the first-order lognormal skew by using the sqrt-forward variables

$$\beta = ce^{x/2}, \quad m = -x \geq 0, \quad a = m/v.$$

The leading price is the Bachelier integral

$$I_0(m, v) = v\phi(a) - m\Phi(-a),$$

and the local Black correction is evaluated through the expansion

$$\beta_{\text{Bl}}(m, v) \approx I_0 - \frac{I_2}{8} + \frac{I_4}{128},$$

where

$$I_2 = \frac{v^3 \phi(a) - m^2 I_0}{3}, \quad I_4 = \frac{v^5 \phi(a) - m^2 I_2}{5}.$$

The Newton denominator uses the exact sqrt-forward Black vega identity

$$\frac{\partial \beta}{\partial v} = \phi(a) e^{-v^2/8},$$

so the correction remains well-scaled even when the price itself is microscopic.

There are three exits. If $m > 0$ and $\ln(m/\beta) > 20$, a normal deep-tail equation is solved for $a = m/v$:

$$\ln\left(\frac{\beta}{m}\right) = -\frac{a^2}{2} - \frac{1}{2} \ln(2\pi) + \ln\left(\frac{1}{a} - \sqrt{\frac{\pi}{2}} \operatorname{erfcx}\left(\frac{a}{\sqrt{2}}\right)\right).$$

The scaled Mills-ratio form avoids subtracting two underflow-level normal tails. This candidate is accepted only when the resulting ratio m/v is larger than 4, where the omitted Black correction is numerically negligible. Otherwise, the rational Bachelier approximation of Le Floch'h [12] supplies the normal-model seed, and two Newton corrections are applied to the expansion above. The only remaining terminal path is defensive: if neither guarded path returns a finite positive value, the branch returns the zero-volatility limit. For finite admissible inputs this path is not expected, since the rational seed covers the ATM and transition regimes while the deep-tail solve intercepts ratios below the rational approximation's finite domain.

A deep-tail example is obtained directly in normalised coordinates by taking

$$x = -10^{-14}, \quad e^x = 0.99999999999999, \quad T = 1, \quad \ln(c) = -100.$$

Equivalently, one may take $K_* = 1$ and $F_* = e^x = 0.99999999999999$, giving $C_{\text{OTM}} = F_* c = 3.7200759760207988 \times 10^{-44}$. Thus $c = e^{-100} = 3.720075976020836 \times 10^{-44}$ and

$$m = -x = 10^{-14}.$$

The Bachelier-limit branch is active because $c \leq 10^{-6}$, $|x| \leq 10^{-8}$, and $x \leq 0$. The deep-tail sub-branch is selected first (before the rational Bachelier seed) because

$$\ln(m) - \ln(c) = 67.76380869808337 > 20.$$

Solving the scaled Mills-ratio equation gives

$$\hat{v} = 9.155604419747184 \times 10^{-16}, \quad m/\hat{v} = 10.922271803739784 > 4.$$

A 120-digit check of the Black formula gives the consistent root $v_{\text{ref}} = 9.155604419747139 \times 10^{-16}$ for $\ln(c) = -100$. The relative difference between the branch value and this high-precision Black root is about 5.0×10^{-15} . By contrast, $v = 1.25 \times 10^{-15}$ gives $\ln(c) = -71.43799710528910$, so it is not the reference volatility for the $\ln(c) = -100$ input. In ordinary double precision the direct erfcx Black log-price difference collapses for this point, which is why the solver uses the Bachelier-limit tail equation instead of entering the Black H3 path.

A second example shows why the rational Bachelier seed is still needed outside the accepted deep tail. Take

$$x = -10^{-8}, \quad e^x = 0.9999999900000001, \quad T = 1, \quad c = 10^{-16}.$$

Then

$$m = 10^{-8}, \quad \beta = ce^{x/2} = 9.999999950000000 \times 10^{-17}, \quad \ln(m/\beta) = 18.420680748952365.$$

The price and moneyness are microscopic, but the conservative deep-tail test is not satisfied because $\ln(m/\beta) < 20$. This is the transition region: the central estimate $\sqrt{2\pi}\beta$ is too small by almost a factor of eight, while the asymptotic tail equation is not used. The rational Bachelier seed gives the normal root

$$v = 1.9952018436169516 \times 10^{-9}, \quad m/v = 5.012024238044884,$$

and the local Black correction leaves the value unchanged at the shown precision. A high-precision evaluation of $\Phi(x/v + v/2) - e^{-x}\Phi(x/v - v/2)$ at this value gives $1.0000000000000004 \times 10^{-16}$, i.e., the target price to double precision. The example is therefore neither an ATM case nor a safely asymptotic tail case; it is exactly the narrow zone for which the Bachelier rational seed keeps the expansion polish in its convergence basin.

Appendix A.4. Defensive fallback and upper-price guards

The Choi L3 seed is the intended non-Bachelier entry point. The production code nevertheless treats the inverse-normal probability calculation as a floating-point object: the probability is clamped to the open double-precision interval, and any non-finite or non-positive seed is replaced before the Euler–Chebyshev chain begins. The replacement is deliberately conservative. For a very small near-ATM price ($c < 10^{-4}$ and $|x| < 0.01$), it uses

$$v_0 = \sqrt{x^2 + 2\pi c^2},$$

the first-order ATM relation with a moneyness correction. Otherwise it tries the guarded OTM asymptotic seed

$$D = \sqrt{\max\{-2\ln(c) - \ln(2\pi), 0\}}, \quad v_0 = \frac{-2x}{D + \sqrt{D^2 - 2x}},$$

accepted only when the discriminant and denominator are finite and positive. If even this defensive path fails, the terminal fallback is $v_0 = \sqrt{2|x|}$, followed by the same positive floor 10^{-10} used by the hot path. These branches are not part of the real-arithmetic convergence proof; they ensure that rounded inputs still reach a finite positive starting point.

Saturated upper prices are handled as a separate terminal guard. When $c \geq \text{nextDown}(0.9990)$, the current seed is compared with the upper-price asymptotic value

$$R = -2\ln(1 - c) - \ln(2\pi), \quad D = \sqrt{R} \left(1 - \frac{\ln R}{2(R + 1)}\right), \quad v_u = D + \sqrt{\max\{D^2 - 2x, 0\}},$$

and the complementary $\ln(1 - c)$ polish is initialized at $\max(v_0, v_u)$. The complementary gap is evaluated as

$$G(v) = \frac{1}{2}e^{-b_1^2} \operatorname{erfcx}(b_1) + \frac{1}{2}e^{-x-b_2^2} \operatorname{erfcx}(b_2), \quad b_1 = \frac{d_1}{\sqrt{2}}, \quad b_2 = -\frac{d_2}{\sqrt{2}}.$$

With $f = \ln G(v) - \ln(1 - c)$, vega $v = \exp(-d_1^2/2)/\sqrt{2\pi}$, and $d'_1 = 1/2 - x/v^2$, the implementation applies two safeguarded Newton–Halley updates of the form

$$\Delta = \frac{f G(v)}{v}, \quad \Delta \leftarrow \frac{\Delta}{1 - \frac{1}{2}f(d'_1 d_1 G(v)/v - 1)}, \quad v \leftarrow \max(v + \Delta, v/2),$$

whenever the quantities are finite and the Halley denominator is safely positive. This branch is shared production guard logic rather than a Thiophene-specific theorem: robust implied-volatility implementations need the same Bachelier-limit, fallback-seed, and upper-price protections even when their central seeds and refinement chains differ.

Appendix A.5. Dataset grids

The dataset builders use the following grids, with the above filtering applied after each raw case is generated:

- **CLY-3D.** $S = 100$, $r = 0.03$, $K = \text{linspace}(105, 800, 40)$, $T = \text{linspace}(0.01, 2, 40)$, and $\sigma = \text{linspace}(0.01, 0.99, 40)$, retaining prices above 10^{-20} as in the Cui-Liu-Yao comparison script.
- **CLY-20.** $S = 100$, $r = 0.03$, $\sigma = 0.20$, $K = \text{linspace}(105, 180, 40)$, and $T = \text{linspace}(0.1, 2, 40)$.
- **CLY-80.** $S = 100$, $r = 0.03$, $\sigma = 0.80$, $K = \text{linspace}(105, 800, 40)$, and $T = \text{linspace}(0.1, 2, 40)$.
- **Jäckel.** $S = 100$, $r = 0$, $K = 100 \text{ linspace}(0.5, 8, 30)$, $T \in \{0.01, 0.1, 0.25, 0.5, 1, 2\}$, and $\sigma = \text{linspace}(0.02, 4, 30)$.
- **Market.** $S = 100$, $r = 0.03$, $K = 100 \text{ linspace}(0.7, 1.5, 30)$, $T \in \{1/252, 5/252, 21/252, 63/252, 0.5, 1, 2, 5\}$, and $\sigma = \text{linspace}(0.05, 1.5, 30)$.
- **Corners.** The union of low-volatility short-maturity ITM calls; deep OTM calls with $K \in \{200, 300, 500, 1000, 2000\}$; near-OTM short-maturity calls with $K = \text{linspace}(101, 150, 10)$; high-volatility cases with $K \in \{100, 150, 200, 500\}$; and the near-ATM small-price grid $K \in \{100.5, 101, 102, 105, 110\}$, $T \in \{0.001, 0.005, 0.01\}$, $\sigma \in \{0.005, 0.01, 0.02, 0.05\}$.
- **Stress.** $S = 100$, $r = 0.03$, $K \in \{101, 102, 103, 110, 150, 200, 500, 1000, 2000, 5000, 10000\}$ and $K \in \{10, 20, 50, 80, 90, 95, 98, 99\}$, $T \in \{0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1, 2, 5, 10\}$, and $\sigma \in \{0.01, 0.02, 0.05, 0.10, 0.20, 0.30, 0.50, 0.80, 0.99\}$.
- **HighVol.** $S = 100$, $r = 0$, $K \in \{1, 2, 3, 4\}$, $T \in \{1, 2, 3, 5, 7, 10\}$, and $\sigma \in \{0.5, 0.8, 1.0, 1.2, 1.5, 2.0, 2.5\}$, retaining only cases with finite $\ln(c)$ and $\ln(c) \leq -0.05$ before the common normalisation filter.

Appendix B. Expanded Normalised-Black Evaluation

The cancellation-avoiding computation used for the expanded Jäckel reference price in Table 2, Figure 1, and Figure 4 evaluates Jäckel's normalised-Black value in the sqrt-forward normalisation. For $x \leq 0$ and total volatility s , define

$$h = \frac{x}{s}, \quad t = \frac{s}{2}, \quad v(x, s) = \frac{1}{\sqrt{2\pi}} \exp\left[-\frac{1}{2}(h^2 + t^2)\right].$$

In the small-volatility branch exercised by the monotonicity diagnostic, the normalised price $\beta = c\sqrt{e^x}$ is evaluated as

$$\beta_{\text{exp}}(x, s) = v(x, s) t \sum_{j=0}^6 b_j(h) t^{2j}.$$

Let

$$a(h) = 1 + h\sqrt{\frac{\pi}{2}} \operatorname{erfcx}\left(-\frac{h}{\sqrt{2}}\right).$$

The coefficients are

$$\begin{aligned} b_0 &= 2a, \\ b_1 &= \frac{-1 + a(3 + h^2)}{3}, \\ b_2 &= \frac{-7 - h^2 + a(15 + 10h^2 + h^4)}{60}, \\ b_3 &= \frac{-57 - 18h^2 - h^4 + a(105 + 105h^2 + 21h^4 + h^6)}{2520}, \\ b_4 &= \frac{1}{181440} \left[-561 - 285h^2 - 33h^4 - h^6 \right. \\ &\quad \left. + a(945 + 1260h^2 + 378h^4 + 36h^6 + h^8) \right], \\ b_5 &= \frac{1}{19958400} \left[-6555 - 4680h^2 - 840h^4 - 52h^6 - h^8 \right. \\ &\quad \left. + a(10395 + 17325h^2 + 6930h^4 + 990h^6 + 55h^8 + h^{10}) \right], \\ b_6 &= \frac{1}{3113510400} \left[-89055 - 82845h^2 - 20370h^4 - 1926h^6 - 75h^8 - h^{10} \right. \\ &\quad \left. + a(135135 + 270270h^2 + 135135h^4 + 25740h^6 + 2145h^8 + 78h^{10} + h^{12}) \right]. \end{aligned}$$

The implementation evaluates $a(h)$ with rational approximations in the regions where the direct erfcx expression would lose relative accuracy. Outside the small-volatility branch it falls back to the corresponding asymptotic or Cody- erfcx branch from Jäckel’s reference implementation. This appendix records the normalisation and the local expansion used by the diagnostics; it is not a separate pricing model.

Appendix C. Choi–Huh–Su L3 Seed in ThiopheneIV

ThiopheneIV uses the Choi–Huh–Su L3 lower-bound formula as an initial value for the Euler–Chebyshev refinement. This appendix records the formula in the normalised variables used throughout. The solver receives $x = \ln(F_*/K_*) \leq 0$, $e^x = F_*/K_*$, and $c = C_{\text{OTM}}/F_*$. Define

$$k = -x \geq 0, \quad E = e^k = \frac{1}{e^x} = \frac{K_*}{F_*}. \quad (\text{A1})$$

For $k > 0$, the L3 seed first maps the observed price to

$$p_3 = \frac{c(c + E)}{2c + E - 1}, \quad z_3 = \Phi^{-1}(p_3). \quad (\text{A2})$$

The denominator is positive on the admissible OTM price interval. In floating-point arithmetic the implementation still rejects non-positive or non-finite denominators and clamps p_3 to the open interval representable by doubles before evaluating the inverse normal CDF.

The final step is purely algebraic. Since

$$d_1(v) = \frac{x}{v} + \frac{v}{2} = -\frac{k}{v} + \frac{v}{2}, \quad (\text{A3})$$

the seed solves $d_1(v) = z_3$, equivalently $v^2/2 - z_3v - k = 0$. The positive solution is evaluated as

$$v_{\text{L3}} = \begin{cases} z_3 + \sqrt{z_3^2 + 2k}, & z_3 \geq 0, \\ \frac{2k}{\sqrt{z_3^2 + 2k} - z_3}, & z_3 < 0, \end{cases} \quad (\text{A4})$$

where the second branch is the rationalised form of the same root and avoids cancellation when z_3 is negative.

At the ATM limit $k = 0$, Equation (A2) reduces to $p_3 = (1 + c)/2$ and the Black price is $c = 2\Phi(v/2) - 1$. Thus we use

$$v_{\text{ATM}} = 2\Phi^{-1}\left(\frac{1 + c}{2}\right), \quad (\text{A5})$$

with the small-price series

$$v_{\text{ATM}} \approx \sqrt{2\pi} c \left(1 + \frac{\pi c^2}{12}\right), \quad c < 10^{-4}, \quad (\text{A6})$$

to avoid spending an inverse-normal evaluation on an almost linear case. If any of these steps produces a non-finite or non-positive seed, ThiopheneIV falls back to the same asymptotic safeguard used by the standalone reference implementation. For saturated prices ($c \geq 0.9990$), the seed is also compared with the upper-price asymptotic value before the complementary $\ln(1 - c)$ polish is applied.

Appendix D. Monotone Convergence of Chebyshev’s Method from the Choi Seed

This appendix provides a self-contained proof that Chebyshev’s method (Chebyshev–Halley family, $\alpha = 0$) converges monotonically from below to the root of the log-price objective when started from the Choi–Huh–Su L3 lower-bound seed, for all admissible OTM prices in real arithmetic. Throughout, we use the notation of Section 2.2.

Setup

Let $x \leq 0$, $k = -x \geq 0$, and let the log-price objective be

$$g(v) = \ln b(x, v) - \ln c_{\text{target}}, \quad (\text{A7})$$

where $b(x, v)$ is the normalised Black call price and $c_{\text{target}} \in (0, 1)$ is fixed. The unique positive root v_* satisfies $g(v_*) = 0$. The log-vega is $y(v) = g'(v) = f(v)/b(x, v)$, where $f(v) = \phi(d_1(v)) > 0$ is the Black vega and $d_1(v) = x/v + v/2$.

For $v < v_*$ we have $g(v) < 0$. Define the Newton displacement $\eta(v) = -g(v)/g'(v) > 0$, and the normalised curvature $q(v) = g''(v)/g'(v)$. The *Euler–Chebyshev step* is

$$T(v) = v + \eta(v)\left(1 - \frac{1}{2} q(v) \eta(v)\right). \quad (\text{A8})$$

Let v_0 be the Choi–Huh–Su L3 seed (Appendix C). Choi et al. [5] prove the lower-bound property: $0 < v_0 \leq v_*$ for all admissible (x, c) .

Theorem A1. *For all $x \leq 0$, $v \in (0, v_*)$, and admissible $c_{\text{target}} \in (0, 1)$, the Chebyshev displacement satisfies*

$$0 < T(v) - v \leq v_* - v. \quad (\text{A9})$$

Consequently, the iterates $v_{n+1} = T(v_n)$ starting from v_0 are monotone increasing, bounded above by v_* , and converge to v_* .

The proof requires two lemmas.

Lemma A: log-concavity of b in v

Lemma A1. *The normalised Black price $b(x, v)$ is log-concave in v : $g''(v) \leq 0$ for all $x \leq 0$, $v > 0$.*

Proof. Let $a(v) = f'(v)/f(v)$ denote the log-derivative of the vega. Differentiating $d_1 = x/v + v/2$ gives $d_1' = -x/v^2 + 1/2$, hence

$$a(v) = \frac{k^2}{v^3} - \frac{v}{4}. \quad (\text{A10})$$

The second log-price derivative is $g'' = g'(a - y)$. Subtracting the two log-derivative relations,

$$q(v) = \frac{g''(v)}{g'(v)} = a(v) - y(v). \quad (\text{A11})$$

Setting $H(v) = f(v) - a(v)b(x, v)$, we compute

$$H'(v) = f'(v) - a'(v)b(x, v) - a(v)f(v) = \beta(v)b(x, v), \quad (\text{A12})$$

where $\beta(v) = -a'(v) = 3k^2/v^4 + 1/4 > 0$, using $f'(v) = a(v)f(v)$ and $b'(x, v) = f(v)$. As $v \downarrow 0$, $H(v)$ has a non-negative limit: for $k > 0$, the standard Mills expansion of the two cancelling normal tails gives $b(x, v) = f(v)(v^3/k^2 + O(v^5))$, so $H(v) \rightarrow 0$; for $k = 0$, $H(v) \rightarrow \phi(0)$. Since $H' > 0$, we conclude $H(v) \geq 0$, i.e. $f(v) \geq a(v)b(x, v)$, equivalently $y(v) \geq a(v)$, equivalently $q(v) = a(v) - y(v) \leq 0$. Therefore $g''(v) = g'(v)q(v) \leq 0$. $\square$

From Lemma A1, $q(v) \leq 0$ and $\eta(v) > 0$, so $1 - \frac{1}{2}q\eta \geq 1 > 0$, hence $T(v) > v$. This proves the left inequality in (A9).

Lemma B: scalar curvature inequality

Lemma A2. For all $x \leq 0$, $v > 0$, with $y(v) = g'(v)$, $a(v) = k^2/v^3 - v/4$, $\beta(v) = 3k^2/v^4 + 1/4$,

$$y^2 - 3ay + 2a^2 + \beta \geq 0. \quad (\text{A13})$$

Proof. Set $r = y - a \geq 0$ (from Lemma A1). Substituting $y = a + r$ into (A13) gives

$$y^2 - 3ay + 2a^2 + \beta = r^2 - ar + \beta. \quad (\text{A14})$$

If $a \leq 0$, then

$$r^2 - ar + \beta \geq \beta > 0, \quad (\text{A15})$$

so (A13) follows immediately. It remains to consider $a > 0$. Define $G(v) = af(v)/(a^2 + \beta)$. Using $f' = af$ and $a' = -\beta$, a direct calculation gives

$$G'(v) - f(v) = -f(v) \frac{a\beta' + 2\beta^2}{(a^2 + \beta)^2}. \quad (\text{A16})$$

Substituting $a = v(z - 1/4)$, $\beta = 3z + 1/4$, $\beta' = -12z/v$ with $z = k^2/v^4 \geq 0$ gives

$$a\beta' + 2\beta^2 = -12z^2 + 3z + 18z^2 + 3z + \frac{1}{8} = 6z^2 + 6z + \frac{1}{8} > 0. \quad (\text{A17})$$

Hence $G' \leq f$. Because $a > 0$ implies $k > 0$, we also have $G(0+) = 0$. Integrating from 0 to v yields $G(v) \leq b(x, v)$, i.e. $af/(a^2 + \beta) \leq b$, i.e. $y = f/b \leq (a^2 + \beta)/a = a + \beta/a$, hence $r \leq \beta/a$.

Case 1: $\beta \geq a^2/4$. Then $r^2 - ar + \beta = (r - a/2)^2 + \beta - a^2/4 \geq 0$.

Case 2: $\beta < a^2/4$. On $r \in [0, \beta/a]$, the parabola $D(r) = r^2 - ar + \beta$ has its minimum at $r = a/2 > \beta/a$, so D is decreasing on $[0, \beta/a]$. Therefore

$$D(r) \geq D\left(\frac{\beta}{a}\right) = \frac{\beta^2}{a^2} - \beta + \beta = \frac{\beta^2}{a^2} > 0. \quad (\text{A18})$$

□

Proof of Theorem A1

Proof. Let ψ be the inverse branch of g that maps $(-\infty, 0)$ to $(0, v_*)$. Setting $u = g(v) < 0$ and $s = -u > 0$, we have $v = \psi(u)$ and $v_* = \psi(0)$. The inverse derivatives are $\psi' = 1/g'$, $\psi'' = -g''/g'^3$, and

$$\psi'''(u) = \frac{3g''^2 - g'g'''}{g'^5}. \quad (\text{A19})$$

Observe that the Chebyshev displacement equals the second-order Taylor approximation of ψ from u to 0:

$$\begin{aligned} T(v) - v &= s\psi'(u) + \frac{1}{2}s^2\psi''(u) \\ &= \frac{-g}{g'} - \frac{g''g^2}{2g'^3} = \eta\left(1 - \frac{1}{2}q\eta\right). \end{aligned} \quad (\text{A20})$$

Taylor's theorem gives

$$v_* - v = \psi(0) - \psi(u) = s\psi'(u) + \frac{1}{2}s^2\psi''(u) + \frac{1}{6}s^3\psi'''(\xi) = T(v) - v + \frac{s^3}{6}\psi'''(\xi), \quad (\text{A21})$$

for some $\xi \in (u, 0)$. Thus $T(v) - v \leq v_* - v$ follows once $\psi''' \geq 0$ on $(u, 0)$. Since the formula for ψ''' is evaluated at $w = \psi(\xi) \in (v, v_*)$, it is enough to verify $3g''^2 - g'g''' \geq 0$ for all positive total volatilities. Using $\delta_2 = g''/g' = q$ and $\delta_3 = g'''/g'$ from Proposition 2,

$$3g''^2 - g'g''' = g'^2(3q^2 - \delta_3). \quad (\text{A22})$$

From (7)–(8),

$$\delta_3 = \frac{-3h^2 - t^2 + (h^2 - t^2)^2}{v^2} - 3yq - y^2. \quad (\text{A23})$$

Substituting $a = (h^2 - t^2)/v = k^2/v^3 - v/4$, $a' = -\beta$, $q = a - y$, and $\delta_3 = a' + a^2 - 3yq - y^2 = -\beta + a^2 - 3yq - y^2$ yields, after collecting terms,

$$3q^2 - \delta_3 = 3(a - y)^2 - (-\beta + a^2 - 3y(a - y) - y^2) = 2q^2 - q' = y^2 - 3ay + 2a^2 + \beta. \quad (\text{A24})$$

By Lemma A2, this is non-negative for all $x \leq 0$, $v > 0$. Hence $\psi'''(\xi) \geq 0$, and the right inequality $T(v) - v \leq v_* - v$ holds.

The left inequality $T(v) > v$ follows from $q \leq 0$ (Lemma A1) and $\eta > 0$.

Both inequalities proven, the iterates satisfy $v_0 \leq v_1 \leq v_2 \leq \dots \leq v_*$. The sequence is monotone and bounded above, so it converges to some $\ell \leq v_*$. If $\ell < v_*$ then $g(\ell) < 0$ and $T(\ell) > \ell$, contradicting $v_n \rightarrow \ell$ and $v_{n+1} = T(v_n) \rightarrow \ell$. Hence $\ell = v_*$.

Local cubic convergence follows from the standard Chebyshev error expansion, since $g'(v_*) > 0$ guarantees a simple root. $\square$

Remark A1. The proof is entirely real-analytic and holds for all $x \leq 0$, $v > 0$, $c \in (0, 1)$, with no compact-domain restriction. The only prerequisite beyond Choi's lower-bound seed (Appendix C) is the log-concavity of b in v (Lemma A1) and the scalar curvature inequality (A13) (Lemma A2). Neither lemma requires any upper-price exclusion; the saturated-price regime $c \approx 1$ is covered because both lemmas hold pointwise for all admissible (x, v) .

Author Contributions: Not applicable (single author).

Funding: This research received no external funding.

Data Availability Statement: The Java source code used to generate the benchmark tables, including the solver variants, benchmark harnesses, and dataset builders, is available in the accompanying source distribution. The exact dataset grids are reproduced in Appendix A.

Conflicts of Interest: The author declares no conflicts of interest.

References

- Black, F. The Pricing of Commodity Contracts. *J. Financ. Econ.* **1976**, *3*, 167–179. [https://doi.org/10.1016/0304-405X\(76\)90024-6](https://doi.org/10.1016/0304-405X(76)90024-6).
- Cui, Z.; Liu, Y.; Yao, Y. Tighter Bounds for Implied Volatility With the Dirac Delta Family Method. *J. Futures Markets* **2025**, *45*, e70024. <https://doi.org/10.1002/fut.70024>.
- Jäckel, P. Let's Be Rational. *Wilmott* **2017**, *2015*, 40–53. <https://doi.org/10.1002/wilm.10395>.
- Le Floc'h, F. Monotonicity of the Black–Scholes Option Prices in Practice. Available online: https://chasethedevil.github.io/post/vol_monotonicity_in_practice/ (accessed on 19 May 2026).
- Choi, J.; Huh, J.; Su, N. Tighter ‘Uniform Bounds for Black–Scholes Implied Volatility’ and the Applications to Root-Finding. *Oper. Res. Lett.* **2024**, *57*, 107189. <https://doi.org/10.1016/j.orl.2024.107189>.
- Birnbaum, Z.W. An Inequality for Mill's Ratio. *Ann. Math. Statist.* **1942**, *13*, 245–246. <https://doi.org/10.1214/aoms/1177731611>.
- Sampford, M.R. Some Inequalities on Mill's Ratio and Related Functions. *Ann. Math. Statist.* **1953**, *24*, 130–132. <https://doi.org/10.1214/aoms/1177729093>.
- Abramowitz, M.; Stegun, I.A. *Handbook of Mathematical Functions*; National Bureau of Standards, Applied Mathematics Series 55: Washington, DC, USA, 1964.

9. Traub, J.F. *Iterative Methods for the Solution of Equations*; Chelsea Publishing Company: New York, NY, USA, 1982.
10. Householder, A.S. *The Numerical Treatment of a Single Nonlinear Equation*; McGraw-Hill: New York, NY, USA, 1970.
11. Healy, J. *Applied Quantitative Finance for Equity Derivatives*, 4th ed.; self-published, 2024.
12. Le Floc'h, F. Fast and Accurate Analytic Basis Point Volatility. Available online: <https://ssrn.com/abstract=2420757> (accessed on 16 May 2026).