

Global Convergence of Wasserstein Policy Gradient for Entropy-Regularized Reinforcement Learning

Zhaoyu Zhu

Shanghai Jiao Tong University

ZZY12345@SJTU.EDU.CN

Rui Gao

The University of Texas at Austin

RUI.GAO@MCCOMBS.UTEXAS.EDU

Shuang Li

The Chinese University of Hong Kong, Shenzhen

LISHUANG@CUHK.EDU.CN

Abstract

Wasserstein policy gradient (WPG) is a policy optimization method for reinforcement learning (RL) that exploits the optimal-transport geometry of action distributions. For the entropy-regularized RL objective, WPG evolves each state-conditional policy by transporting it along the action gradient of the soft Q-function together with a Langevin-type diffusion. Despite its appeal for continuous-control problems, its global convergence properties remain poorly understood. Standard Langevin analyses do not directly apply, because the RL objective depends on the policy through the Bellman recursion rather than through a static convex functional, and the Langevin drift is determined by the soft Q-function, whose regularity must be controlled along the policy iterates.

In this paper, we develop a global convergence theory for WPG by exploiting the Bellman structure of entropy-regularized RL. We show that the role usually played by convexity can be replaced by a Bellman-based argument: the soft Bellman residual admits a statewise KL representation with respect to a Gibbs policy; Bellman contraction relates this residual to the global optimality gap; and a Bellman resolvent identity connects value improvement to relative Fisher information. Combined with a uniform log-Sobolev inequality (LSI) for the evolving Gibbs family, these ingredients yield a distributional Polyak–Łojasiewicz condition. We further establish the regularity and uniform bounds needed to control the discretization error, thereby obtaining geometric contraction up to a discretization bias. Conceptually, our analysis shows that although entropy-regularized RL is not convex in the usual flat sense, the Bellman recursion induces a favorable Polyak–Łojasiewicz-type (PL) geometry that supports global convergence of WPG.

1 Introduction

Policy gradient (PG) methods are among the most widely used algorithms in reinforcement learning (RL). Beginning with REINFORCE (Williams, 1992) and the policy gradient theorem (Sutton et al., 2000), classical PG performs Euclidean gradient ascent over a parameterized policy class. While simple and scalable, Euclidean PG is sensitive to step sizes and parameterization, which has motivated geometry-aware alternatives. Natural policy gradient replaces the Euclidean metric with the information geometry induced by the Fisher information (Amari, 1998; Kakade, 2001), and is closely related to trust-region (TR) methods. In deep RL, TRPO constrains the average KL divergence between successive poli-

cies (Schulman et al., 2015), while PPO provides a practical surrogate through clipping or adaptive KL penalties (Schulman et al., 2017). These developments underscore the central role of geometry in policy optimization.

Beyond information geometry, optimal transport (OT) provides a different geometry for policy space, one that is particularly natural for distributions over continuous action spaces. Existing Wasserstein policy optimization methods can be viewed as different implementations or approximations of Wasserstein gradient flow: through JKO schemes and particle approximations (Zhang et al., 2018), kernelized Wasserstein natural-gradient approximations (Arbel et al., 2020; Moskovitz et al., 2020), trust region formulations (Terpin et al., 2022; Song et al., 2024), and Wasserstein gradient updates projected onto explicit (Pfau et al., 2025) or implicit (Zhu et al., 2026) policy classes. These methods share the same underlying mechanism: policy improvement is driven by transport in action space.

Entropy regularization is another powerful scheme in modern continuous-control RL. Maximum-entropy RL augments reward maximization with an entropy term on the policy distribution over actions, promoting exploration and robustness and leading to soft Bellman operators and Gibbs-type policy improvements (Ziebart, 2010; Haarnoja et al., 2018; Geist et al., 2019). From the OT perspective, this entropy term naturally induces diffusion. Consequently, Wasserstein policy optimization for the entropy-regularized objective takes the form of a drift-diffusion, or Fokker–Planck, equation, whose particle representation is a Langevin-type actor update. More concretely, the Wasserstein Policy Gradient (WPG) flow reads

$$\partial_t \pi_t(a | s) = -\text{div}_a \left(\pi_t(a | s) \nabla_a (Q^{\pi_t}(s, a) - \tau \log \pi_t(a | s)) \right),$$

and its discrete-time Langevin counterpart is

$$A_{k+1}^s = A_k^s + \eta \nabla_a Q^{\pi_k}(s, A_k^s) + \sqrt{2\tau\eta} \xi_{k+1}, \quad \xi_{k+1} \sim \mathcal{N}(0, I_d), \quad (1)$$

where $\pi_k(\cdot | s) = \text{Law}(A_k^s)$. Thus, each state-conditional policy is transported along the action gradient of the current soft Q-function, while entropy regularization contributes to the Langevin diffusion.

Despite the algorithmic appeal of WPG, its global convergence behavior remains largely open. Existing global analyses of PG or natural PG typically rely on Euclidean or KL/Bregman geometry in policy space, using mirror-descent identity to relate policy updates to global suboptimality (Agarwal et al., 2021; Bhandari and Russo, 2024; Cen et al., 2022; Bhandari and Russo, 2021; Khodadadian et al., 2021, 2022; Yuan et al., 2023; Xiao, 2022; Lan, 2023; Zhan et al., 2023; Lan et al., 2023). These arguments do not transfer to Wasserstein geometry, which is non-Bregman and lacks the pointwise three-point structure underlying KL-based analyses. Existing OT-based convergence results also do not directly help for the Langevin WPG update in (1). The convergence result in Zhang et al. (2018) is asymptotic and places the entropy regularization on the policy parameters rather than on state-conditional policy distributions. The trust-region analysis of Song et al. (2024) applies to finite tabular MDPs without entropy regularization, and Zhu et al. (2026) studies a JKO-type implicit proximal scheme. None of these analyses transfers to Langevin dynamics.

Our analysis is closely related to the broader literature on mean-field Langevin dynamics, which has been used to study training dynamics of large interacting particle systems and over-parameterized neural networks (Chizat and Bach, 2018; Mei et al., 2018; Nitanda

et al., 2022; Chizat, 2022; Cai et al., 2025). However, global convergence arguments in that literature typically rely on a convex objective and convexity-analysis based argument. RL objective has a different structure: the policy enters the objective through the Bellman recursion which destroys the convexity, and the soft Q-function entangles the reward and entropy term which make the regularity of the Langevin drift not taken for granted.

Contribution. Our main contribution is a non-asymptotic global convergence analysis of WPG for entropy-regularized continuous-action RL. Rather than relying on the convex-analysis machinery commonly used in standard Mean-field Langevin dynamics, we exploit the Bellman structure of the RL objective. The analysis is built on the following ingredients.

- *Value improvement-to-Fisher information via a Bellman resolvent identity.* We show that policy-space dissipation propagates through the Bellman equation via a resolvent identity. This allows us to lower bound value improvement by Fisher-information dissipation, replacing the standard energy-dissipation identity used in convex settings.
- *Optimality gap-to-KL via Bellman residuals and contraction.* We represent the state-wise soft Bellman residual as a KL divergence between the current policy and its associated Gibbs policy. Bellman contraction then relates this residual to the global optimality gap in sup-norm, replacing the entropy-sandwich arguments used in convex settings.
- *Uniform control of the moving Gibbs family.* To exploit the commonly used uniform logarithmic Sobolev inequalities to connect the KL divergence (controlling the optimality gap) with the Fisher information (controlling value improvement), we establish uniform bound estimates along the WPG iterates.

We first illustrate these ideas in the continuous-time WPG flow, where the Bellman residual, contraction, resolvent mechanisms appear most transparently. Then we provide careful analysis for the discrete-time Langevin update (1), which requires additional regularity estimates for the Wasserstein gradient and a one-step interpolation argument (Vempala and Wibisono, 2019) to control time-discretization error. To this end, we establish refined uniform *a priori* bounds on value functions, soft Q -functions, drift Lipschitz constants, action moments, drift moments, and the relevant KL quantities. Combining these ingredients yields geometric contraction of the global optimality gap up to a discretization bias.

Organization. Section 2 introduces the entropy-regularized discounted RL model, the soft Bellman operators, and the Wasserstein/Langevin policy update. Section 3 presents the continuous-time analysis, isolating the core Bellman mechanisms underlying the proof. Section 4 proves the finite-time convergence theorem for the discrete-time update, with emphasis on the additional regularity estimates and discretization-error control. The appendices contain auxiliary results and detailed proofs.

2 Model Setup

We consider an infinite-horizon discounted Markov decision process

$$\mathcal{M} = (\mathcal{S}, \mathcal{A}, P, r, \gamma, \rho_0),$$

where $\mathcal{S}$ is the state space, $\mathcal{A}$ is the action space, $\gamma \in (0, 1)$ is the discount factor, $\rho_0 \in \mathcal{P}(\mathcal{S})$ is the initial state distribution, $P(ds' \mid s, a)$ is a Markov transition kernel on $\mathcal{S}$, and $r : \mathcal{S} \times \mathcal{A} \rightarrow \mathbb{R}$ is a measurable reward function. A (stationary Markov) policy is a conditional distribution $\pi(da \mid s)$ that admits a density, still denoted $\pi(a \mid s)$, with respect to Lebesgue measure on $\mathcal{A}$. Given π , a trajectory $(s_n, a_n)_{n \geq 0}$ is generated by $s_0 \sim \rho_0$, $a_n \sim \pi(\cdot \mid s_n)$, and $s_{n+1} \sim P(\cdot \mid s_n, a_n)$. We write $\mathbb{E}_{\rho_0, \pi}[\cdot]$ for expectations under this rollout distribution. The normalized discounted state occupancy of π is the probability measure $d^\pi \in \mathcal{P}(\mathcal{S})$ defined by

$$d^\pi(B) = (1 - \gamma) \sum_{n \geq 0} \gamma^n \mathbb{P}_{\rho_0, \pi}(s_n \in B), \quad B \in \mathcal{B}(\mathcal{S}).$$

We assume throughout that the initial distribution ρ_0 has full support on $\mathcal{S}$. Hence, for every policy π ,

$$d^\pi(B) = (1 - \gamma) \sum_{t \geq 0} \gamma^t \mathbb{P}_{\rho_0, \pi}(s_t \in B) \geq (1 - \gamma) \rho_0(B), \quad B \in \mathcal{B}(\mathcal{S}).$$

Thus d^π has full support whenever ρ_0 does. This ensures that the statewise policy-gradient updates below are specified on the same state space on which the sup-norm value bounds are proved. We use the following standing measurability convention. The model kernels and their action derivatives admit jointly measurable versions such that all displayed integrals are jointly measurable. Under this convention, Bellman fixed points are selected as bounded measurable fixed points of contraction maps, and Gibbs and WPG kernels are taken in their induced jointly measurable versions.

We work with an action space $\mathcal{A} = \mathbb{R}^d$. We consider the regularized reward

$$\tilde{r}(s, a) = r(s, a) - \frac{\beta}{2} \|a\|^2,$$

where $\beta > 0$ is the quadratic action penalty. This penalty term stabilizes the training (Lillicrap et al., 2015; Brockman et al., 2016; Tunyasuvunakool et al., 2020), and is also exploited in standard mean-field Langevin dynamics analysis (Yamamoto et al., 2024), closely related to the dissipativity condition that helps to control the KL divergence along the trajectory. Fix a temperature $\tau > 0$, we study the discounted regularized objective

$$J(\pi) = \mathbb{E}_{\rho_0, \pi} \left[\sum_{n \geq 0} \gamma^n (\tilde{r}(s_n, a_n) - \tau \log \pi(a_n \mid s_n)) \right],$$

restricting attention to policies for which the above expectation is finite. The quadratic action penalty is essential on the unbounded action space. Without this term, the entropy bonus can be made arbitrarily large by spreading the policy mass over larger regions of $\mathbb{R}^d$, so the entropy-regularized policy-improvement objective may be unbounded above. The quadratic penalty rules out this degeneracy. Equivalently, the quadratic term induces the Gaussian reference

$$\rho_\beta(a) = Z_\beta^{-1} \exp \left(-\frac{\beta}{2\tau} \|a\|^2 \right), \quad Z_\beta = \left(\frac{2\pi\tau}{\beta} \right)^{d/2}.$$

For the discrete-time result, we work on the admissible class

$$\Pi := \left\{ \pi : \pi(\cdot | s) \ll \text{da}, (s, a) \mapsto \pi(a | s) \text{ jointly measurable, } \sup_{s \in \mathcal{S}} \text{KL}(\pi(\cdot | s) \| \rho_\beta) < \infty \right\}.$$

Finite relative entropy gives finite second moments; see Lemma E.1 in Appendix E.

2.1 Soft value functions and Bellman operators

Given a policy π , define the soft value function

$$V^\pi(s) = \mathbb{E} \left[\sum_{n \geq 0} \gamma^n (\tilde{r}(s_n, a_n) - \tau \log \pi(a_n | s_n)) \mid s_0 = s \right],$$

so that $J(\pi) = \int_{\mathcal{S}} V^\pi(s) \rho_0(ds)$ whenever the integral is well-defined. Define the state-action value function

$$Q^\pi(s, a) = \tilde{r}(s, a) + \gamma \int_{\mathcal{S}} V^\pi(s') P(ds' | s, a).$$

Then V^π satisfies the soft Bellman identity

$$V^\pi(s) = \int_{\mathcal{A}} (Q^\pi(s, a) - \tau \log \pi(a | s)) \pi(a | s) da.$$

For a fixed policy π , define the policy evaluation operator acting on bounded measurable $V : \mathcal{S} \rightarrow \mathbb{R}$ by

$$(\mathcal{T}^\pi V)(s) = \int_{\mathcal{A}} \left(\tilde{r}(s, a) + \gamma \int_{\mathcal{S}} V(s') P(ds' | s, a) - \tau \log \pi(a | s) \right) \pi(a | s) da. \quad (2)$$

Under standard measurability and integrability conditions, $\mathcal{T}^\pi$ is a γ -contraction in $\|\cdot\|_\infty$ and its unique fixed point is $V^\pi = \mathcal{T}^\pi V^\pi$. For a bounded candidate value V , write

$$Q_V(s, a) := \tilde{r}(s, a) + \gamma \int_{\mathcal{S}} V(s') P(ds' | s, a).$$

Define the soft optimality operator by

$$(\mathcal{T}^* V)(s) = \sup_{\mu} \int_{\mathbb{R}^d} (Q_V(s, a) - \tau \log \mu(a)) \mu(a) da,$$

where the supremum ranges over action densities for which the expression is finite. The supremum is attained uniquely at a Gibbs density, and therefore

$$(\mathcal{T}^* V)(s) = \tau \log \int_{\mathbb{R}^d} \exp \left(\frac{Q_V(s, a)}{\tau} \right) da, \quad \mathcal{G}[V](a | s) = \frac{\exp(Q_V(s, a)/\tau)}{\int_{\mathbb{R}^d} \exp(Q_V(s, \tilde{a})/\tau) d\tilde{a}}.$$

Moreover, $\mathcal{T}^*$ is a γ -contraction in $\|\cdot\|_\infty$ and has a unique bounded fixed point $V^* = \mathcal{T}^* V^*$. We set $Q^* := Q_{V^*}$ and $\pi^* := \mathcal{G}[V^*]$.

For each (π, s) , define the Gibbs density by

$$p_s^\pi(a) := \mathcal{G}[V^\pi](a | s) = \frac{\exp(Q^\pi(s, a)/\tau)}{\int_{\mathbb{R}^d} \exp(Q^\pi(s, \tilde{a})/\tau) d\tilde{a}}, \quad a \in \mathbb{R}^d. \quad (3)$$

The following identity relates the Bellman residual to the KL divergence between the current policy and the Gibbs density.

Lemma 2.1 (Bellman residual identity) *For every $s \in \mathcal{S}$,*

$$(\mathcal{T}^* V^\pi)(s) - V^\pi(s) = \tau \text{KL}(\pi(\cdot | s) \| p_s^\pi) \geq 0,$$

where p_s^π is the Gibbs density defined in (3).

2.2 Wasserstein Policy Gradient

Fix a policy π . The first-variation calculation in Appendix A.2 shows that, for statewise transport perturbations of the action law, the ascent direction is

$$\nabla_a (Q^\pi(s, a) - \tau \log \pi(a | s)).$$

The discounted occupancy d^π enters this calculation through the usual policy-gradient weighting over states, as in the policy-gradient theorem and trust-region/natural-gradient policy methods (Sutton et al., 2000; Kakade, 2001; Schulman et al., 2015). Taking the Wasserstein gradient of this first variation state by state gives the Wasserstein policy-gradient flow

$$\partial_t \pi_t(a | s) = -\nabla_a \cdot [\pi_t(a | s) \nabla_a (Q^{\pi_t}(s, a) - \tau \log \pi_t(a | s))] . \quad (\text{WPGF})$$

Using the Gibbs density $p_s^{\pi_t}$ in (3), this can be rewritten as

$$\partial_t \pi_t(a | s) = \tau \nabla_a \cdot \left(\pi_t(a | s) \nabla_a \log \frac{\pi_t(a | s)}{p_s^{\pi_t}(a)} \right) .$$

From an equivalent particle viewpoint, for each fixed state s , the above Fokker–Planck equation can be viewed formally as the forward equation of the nonlinear Langevin diffusion

$$dA_t^s = \nabla_a Q^{\pi_t}(s, A_t^s) dt + \sqrt{2\tau} dB_t^s, \quad \pi_t(\cdot | s) := \text{Law}(A_t^s).$$

Indeed, since

$$\tau \nabla_a \log p_s^{\pi_t}(a) = \nabla_a Q^{\pi_t}(s, a),$$

Applying an explicit Euler–Maruyama step to this statewise Langevin diffusion gives the discrete-time update below.

The discrete-time counterpart of (WPGF) is the Langevin policy update

$$A_{k+1}^s = A_k^s + \eta \nabla_a Q^{\pi_k}(s, A_k^s) + \sqrt{2\tau\eta} \xi_{k+1}, \quad \xi_{k+1} \sim N(0, I_d), \quad (\text{WPGD})$$

where $A_k^s \sim \pi_k(\cdot | s)$ and

$$\pi_{k+1}(\cdot | s) := \text{Law}(A_{k+1}^s).$$

Thus WPGD is a statewise policy-space update specified for every $s \in \mathcal{S}$. Under the full-support convention above, the discounted occupancy weighting does not remove any state from the gradient calculation, and the convergence theorem below controls the resulting value gap in sup-norm over the same state space.

3 Continuous-Time Analysis: Bellman Geometry

This section presents a formal continuous-time analysis of the Wasserstein policy gradient flow ([WPGF](#)). Its purpose is to make transparent the Bellman mechanisms that replace the standard mean-field Langevin dynamics proof chain. We do not claim here a complete well-posedness theorem for the nonlinear Fokker–Planck equation. Instead, the identities below are stated under a smooth-flow regularity convention that supplies the differentiability, integrability, and boundary-decay properties needed for the calculation.

Why the standard mean-field Langevin dynamics proof chain does not directly apply. For a mean-field Langevin dynamics problem, exponential convergence is often proved through the schematic chain ([Nitanda et al., 2022](#); [Chizat, 2022](#)):

$$\frac{d}{dt}\mathcal{F}(\mu_t) \stackrel{(a)}{=} -\mathcal{I}(\mu_t\|\nu_t) \stackrel{(b)}{\lesssim} -\text{KL}(\mu_t\|\nu_t) \stackrel{(c)}{\lesssim} -(\mathcal{F}(\mu_t) - \mathcal{F}^*), \quad (4)$$

where (a) is the gradient-flow dissipation identity, (b) follows from a uniform LSI, and (c) is through a convex-analysis based comparison from local KL to global suboptimality. In entropy-regularized RL, these relationships do not hold in these forms. First, (a) is altered by the Bellman recursion: differentiating V^{π_t} also differentiates Q^{π_t} through the value function, so the value derivative is not merely a statewise Fisher-information dissipation. Lemma 3.1 replaces this step by the value-resolvent identity $(I - \gamma P^{\pi_t})\dot{V}^{\pi_t} = g_t$, with g_t proportional to a statewise Fisher-information term. Second, (b) is not directly available from a convex functional, because the RL objective is nonconvex due to the Bellman recursion of the state-dependent policy distribution. Lemma 3.2 and 3.3 will derive a uniform LSI from uniform value/ Q bounds. Third, (c) cannot be obtained from convexity due to the nonconvexity and aggregation over states in the RL objective. Instead, Lemma 2.1 identifies $\tau\text{KL}(\pi_t(\cdot | s)\|p_s^{\pi_t})$ with the statewise Bellman residual, and Bellman contraction in the proof of Theorem 3.1 will compare this residual at the worst state directly with the $\|\cdot\|_\infty$ -value gap.

Smooth-flow regularity convention. Throughout this section only, the flow ([WPGF](#)) is considered for smooth positive solutions $\pi_t(\cdot | s)$ satisfying the following properties on every finite time interval. For each state s , $t \mapsto V^{\pi_t}(s)$ is locally absolutely continuous, the density $\pi_t(\cdot | s)$ has the moments, entropy, Fisher information, and decay at infinity needed for all displayed integrations by parts, and $a \mapsto Q^{\pi_t}(s, a)$ is sufficiently smooth. Differentiation under the action/state integrals is assumed valid. These assumptions are used only for the formal PDE calculation in this section for illustration and will not be used in the discrete-time analysis in Section 4.

Assumption 3.1 (Boundedness and initialization) *We assume the following throughout:*

- (i) $|r(s, a)| \leq R_{\max}$ for all $(s, a) \in \mathcal{S} \times \mathcal{A}$.
- (ii) $\sup_{s \in \mathcal{S}} \text{KL}(\pi_0(\cdot | s)\|\rho_\beta) \leq K_0$, where $\rho_\beta(a) \propto \exp\left(-\frac{\beta}{2\tau}\|a\|^2\right)$.

Our proof proceeds in three steps. First, we establish a resolvent identity for the value derivative along the flow (Lemma 3.1), which reveals a dissipation given by Fisher in-

formation relative to the moving Gibbs family $p_s^{\pi_t}$ and yields monotone value improvement. Second, we use this monotonicity together with bounded rewards and the initialization condition to obtain uniform-in-time bounds on V^{π_t} and Q^{π_t} (Lemma 3.2); these bounds imply that each $p_s^{\pi_t}$ is a bounded perturbation of a Gaussian, enabling a uniform log-Sobolev inequality along the optimization trajectory via Holley-Stroock Perturbation Lemma (Lemma 3.3). Third, we combine the uniform LSI with the Bellman-residual identity, Lemma 2.1, and Bellman contraction to convert KL dissipation into an exponential-decay estimate for the optimality gap (Theorem 3.1).

Note that RL policy differs from the standard mean-field Langevin dynamics in that it is concerned with a family of state-dependent policy distributions $\{\pi(\cdot | s)\}_{s \in \mathcal{S}}$ rather than a single distribution. Thus, adapting the usual proof chain (4) to RL settings requires choosing how to aggregate statewise quantities. Another possible route is to aggregate the KL terms in (c) by a discounted state-visitation distribution via the performance difference lemma A.1, as in occupancy-weighted policy-gradient analyses. However, this creates a mismatch between the state distribution used to measure the optimality gap and the state aggregation naturally induced by the value derivative or resolvent identity in (a). Closing this gap typically requires density-ratio or coverage assumptions between state distributions (Xiao, 2022; Agarwal et al., 2021; Zhang et al., 2020; Ding et al., 2022). Our argument above circumvents this route by keeping the Bellman residual identity statewise and using Bellman contraction to lower bound the residual in terms of the sup-norm value gap, and thus does not require additional coverage assumptions.

For a policy π , define the induced state kernel

$$P^\pi(ds' | s) := \int_{\mathbb{R}^d} P(ds' | s, a) \pi(a | s) da.$$

Definition 3.1 (KL divergence and relative Fisher information) *Let μ and ν be probability measures on $\mathbb{R}^d$ with $\mu \ll \nu$. The Kullback–Leibler divergence of μ from ν is defined by*

$$\text{KL}(\mu \| \nu) := \int_{\mathbb{R}^d} \log \frac{d\mu}{d\nu} d\mu.$$

The relative Fisher information of μ with respect to ν is defined by

$$\mathcal{I}(\mu \| \nu) := \int_{\mathbb{R}^d} \left\| \nabla \log \frac{d\mu}{d\nu} \right\|^2 d\mu.$$

Definition 3.2 (Log-Sobolev inequality) *Let ν be a probability measure on $\mathbb{R}^d$ with positive density. We say that ν satisfies a log-Sobolev inequality with constant $\alpha > 0$, or α -LSI, if*

$$\mathcal{I}(\mu \| \nu) \geq 2\alpha \text{KL}(\mu \| \nu), \quad \forall \mu \ll \nu$$

for every probability measure μ for which the two sides are well-defined.

We first establish an identity for the time derivative of V^{π_t} along the flow (WPGF).

Lemma 3.1 (Monotonicity of V^{π_t}) Under Assumption 3.1 and the smooth-flow regularity convention, for every $t \geq 0$,

$$(I - \gamma P^{\pi_t}) \frac{d}{dt} V^{\pi_t} = g_t, \quad g_t(s) := \tau^2 \mathcal{I}(\pi_t(\cdot | s) \| p_s^{\pi_t}) \geq 0, \quad (5)$$

where $\mathcal{I}(\pi_t(\cdot | s) \| p_s^{\pi_t})$ is the relative Fisher information. Moreover,

$$\frac{d}{dt} V^{\pi_t}(s) \geq g_t(s) \geq 0 \quad \forall s. \quad (6)$$

Lemma 3.1 shows that the value derivative satisfies

$$\frac{d}{dt} V^{\pi_t} = (I - \gamma P^{\pi_t})^{-1} g_t$$

and is therefore pointwise nonnegative along the flow (WPGF). Note that this identity can be viewed as the statewise resolvent form of the entropy-regularized performance-difference lemma: integrating it against an initial distribution recovers the usual occupancy-weighted performance-difference formula, whereas keeping it pointwise is what enables the subsequent sup-norm contraction argument in (9) below without introducing state-distribution mismatch coefficients. Based on Lemma 3.1, we derive uniform-in-time bounds on the value function, which further imply uniform bounds on the action-value Q-function.

Lemma 3.2 (Uniform bounds on V^{π_t} and Q^{π_t}) Under Assumption 3.1 and the smooth-flow regularity convention above, for all $t \geq 0$,

$$\sup_{s \in \mathcal{S}} V^{\pi_t}(s) \leq \frac{R_{\max} + \tau \log Z_\beta}{1 - \gamma}.$$

Moreover,

$$\inf_{s \in \mathcal{S}} V^{\pi_t}(s) \geq \inf_{s \in \mathcal{S}} V^{\pi_0}(s) \geq \frac{-R_{\max} + \tau \log Z_\beta - \tau K_0}{1 - \gamma}.$$

Consequently, for all $t \geq 0$, $s \in \mathcal{S}$, and $a \in \mathcal{A}$,

$$-R_{\max} - \frac{\beta}{2} \|a\|^2 + \gamma \inf_{x \in \mathcal{S}} V^{\pi_0}(x) \leq Q^{\pi_t}(s, a) \leq R_{\max} - \frac{\beta}{2} \|a\|^2 + \gamma \sup_{x \in \mathcal{S}} V^{\pi_t}(x). \quad (7)$$

Thanks to the bound (7) on the Q-function, the Gibbs densities $p_s^{\pi_t}$ are bounded perturbations of Gaussian densities, and thus satisfy log-Sobolev inequalities with uniform constants along the optimization trajectory due to the Holley-Stroock perturbation lemma.

Lemma 3.3 (Trajectory-uniform LSI for Gibbs policies) Define

$$V_{\max} = \max \left\{ \frac{R_{\max} + \tau \log Z_\beta}{1 - \gamma}, \frac{R_{\max} - \tau \log Z_\beta + \tau K_0}{1 - \gamma} \right\}$$

and

$$U_{\max} = R_{\max} + \gamma V_{\max}.$$

Then for every $t \geq 0$ and $s \in \mathcal{S}$, the Gibbs density $p_s^{\pi_t}$ satisfies the log-Sobolev inequality

$$\mathcal{I}(\nu \| p_s^{\pi_t}) \geq 2\alpha \text{KL}(\nu \| p_s^{\pi_t}) \quad \forall \nu \ll p_s^{\pi_t},$$

with the uniform constant

$$\alpha = \frac{\beta}{\tau} \exp\left(-\frac{2U_{\max}}{\tau}\right).$$

We now state the exponential-decay consequence of the preceding identities. Let V^* be the unique fixed point of $\mathcal{T}^*$, and let π^* denote the optimal Gibbs policy attaining V^* . Since V^* is the optimal value function and the flow remains in the admissible policy class, $V^*(s) \geq V^{\pi_t}(s)$ for all s, t . Define the residual

$$R_t(s) := (\mathcal{T}^* V^{\pi_t})(s) - V^{\pi_t}(s) \geq 0.$$

By Lemma 2.1,

$$R_t(s) = \tau \text{KL}(\pi_t(\cdot | s) \| p_s^{\pi_t}). \quad (8)$$

Define

$$W_t(s) := V^*(s) - V^{\pi_t}(s) \geq 0, \quad E(t) := \|W_t\|_\infty.$$

Since $V^* = \mathcal{T}^* V^*$, $\mathcal{T}^*$ is monotone, and $\mathcal{T}^*$ is a γ -contraction in $\|\cdot\|_\infty$,

$$0 \leq V^*(s) - (\mathcal{T}^* V^{\pi_t})(s) \leq \gamma E(t) \quad \forall s \in \mathcal{S}, t \geq 0. \quad (9)$$

Theorem 3.1 (Exponential convergence along the smooth WPG flow) *Under Assumption 3.1 and the smooth-flow regularity convention above, for all $t \geq 0$,*

$$J(\pi^*) - J(\pi_t) \leq \|V^* - V^{\pi_t}\|_\infty \leq e^{-2\alpha\tau(1-\gamma)t} \|V^* - V^{\pi_0}\|_\infty.$$

Proof By Lemma 3.1, specifically (5) and (6), for almost every t ,

$$\partial_t W_t(s) = -\frac{d}{dt} V^{\pi_t}(s) \leq -g_t(s) = -\tau^2 \mathcal{I}(\pi_t(\cdot | s) \| p_s^{\pi_t}).$$

By Lemma 3.3 and (8),

$$\mathcal{I}(\pi_t(\cdot | s) \| p_s^{\pi_t}) \geq 2\alpha \text{KL}(\pi_t(\cdot | s) \| p_s^{\pi_t}) = \frac{2\alpha}{\tau} R_t(s).$$

Therefore,

$$\partial_t W_t(s) \leq -2\alpha\tau R_t(s) \quad (10)$$

Next, using the definition of W_t and R_t ,

$$W_t(s) = (V^*(s) - (\mathcal{T}^* V^{\pi_t})(s)) + R_t(s).$$

Hence, by (9),

$$R_t(s) = W_t(s) - (V^*(s) - (\mathcal{T}^* V^{\pi_t})(s)) \geq W_t(s) - \gamma E(t). \quad (11)$$

Combining (10) and (11) gives the pointwise differential inequality

$$\partial_t W_t(s) \leq -2\alpha\tau W_t(s) + 2\alpha\tau\gamma E(t), \quad s \in \mathcal{S}, \quad (12)$$

for almost every t .

Multiplying (12) by the integrating factor $e^{2\alpha\tau t}$ and integrating from 0 to t yields

$$e^{2\alpha\tau t} W_t(s) - W_0(s) \leq 2\alpha\tau\gamma \int_0^t e^{2\alpha\tau u} E(u) \, du.$$

Equivalently,

$$W_t(s) \leq e^{-2\alpha\tau t} W_0(s) + 2\alpha\tau\gamma \int_0^t e^{-2\alpha\tau(t-u)} E(u) \, du. \quad (13)$$

Taking the supremum over $s \in \mathcal{S}$ in (13) gives

$$E(t) \leq e^{-2\alpha\tau t} E(0) + 2\alpha\tau\gamma \int_0^t e^{-2\alpha\tau(t-u)} E(u) \, du. \quad (14)$$

Define

$$H(t) := e^{2\alpha\tau t} E(t).$$

Multiplying (14) by $e^{2\alpha\tau t}$ gives

$$H(t) \leq E(0) + 2\alpha\tau\gamma \int_0^t H(u) \, du.$$

By Gronwall's inequality,

$$H(t) \leq E(0) e^{2\alpha\tau\gamma t}.$$

Therefore,

$$E(t) \leq E(0) e^{-2\alpha\tau(1-\gamma)t} = \|V^\star - V^{\pi_0}\|_\infty \exp\{-2\alpha\tau(1-\gamma)t\}.$$

Finally, since ρ_0 is a probability measure and $W_t \geq 0$ pointwise,

$$J(\pi^\star) - J(\pi_t) = \int_{\mathcal{S}} (V^\star(s) - V^{\pi_t}(s)) \rho_0(ds) \leq \|V^\star - V^{\pi_t}\|_\infty.$$

This proves the theorem. ■

Theorem 3.1 is a formal continuous-time convergence for smooth solutions of (WPGF). It identifies the rate obtained from the value-resolvent identity, uniform value/ Q bounds, the uniform LSI, and the statewise Bellman-residual identity. Its role is explanatory: it isolates the Bellman-residual mechanism and motivates the discrete-time argument.

After the first version of this paper was submitted for review, we became aware of a closely related note by Šiška and Zhang (2026) for finite-state continuous-action problems. Their analysis and resulting guarantee are structurally different from ours. In their cost-minimization notation, the argument uses an entropy-sandwich inequality to control the scalar distributional gap $V^{\pi_t}(\rho) - V^{\pi^\star}(\rho)$ and states a bound of the form

$$V^{\pi_t}(\rho) - V^{\pi^\star}(\rho) \leq \bar{\kappa} \left(\int_{\mathcal{S}} (V^{\pi_0}(s) - V^{\pi^\star}(s)) d_{\rho}^{\pi^\star}(ds) \right) e^{-2\kappa\alpha\tau(1-\gamma)t},$$

where $\bar{\kappa} := \sup_{s \in \mathcal{S}} \frac{d\rho}{dd\pi^*}(s)$, $\kappa := \inf_{s \in \mathcal{S}} \frac{d\rho}{dd\pi^*}(s) \leq 1$. Thus their estimate is distribution-dependent and requires coverage-type density-ratio control, in particular $\kappa > 0$, to obtain a positive exponential rate. By contrast, Theorem 3.1 proves a state-uniform bound and therefore controls $J(\pi^*) - J(\pi_t)$ for any initial state distribution. The proof mechanisms are also different: rather than passing through an occupancy-weighted entropy sandwich, we use the exact statewise Bellman-residual identity and Bellman contraction to compare the residual directly with the $\|\cdot\|_\infty$ -optimality gap. This removes the density-ratio constants κ from the exponential convergence rate, although with a different initial gap. Finally, both their and our analyses involve formal calculations; the main rigorous guarantee in the present paper is the discrete-time Langevin analysis in Section 4.

4 Discrete-Time Global Convergence Analysis

This section proves the non-asymptotic convergence guarantee for the explicit Langevin update (WPGD). The continuous-time section isolates the Bellman residual/resolvent mechanism. Apart from their counterparts in discrete time, an additional task is to control the finite-step Langevin discretization and the policy-dependent drift uniformly along the iterates.

4.1 Assumptions

The assumptions below are the quantitative inputs for the finite-step analysis. To apply the one step drift interpolation argument, we need uniform control of the drift Lipschitz constants, action moments, drift second moments, finite KL quantities, and the LSI constants of the Gibbs family. Since

$$b_k(s, a) := \nabla_a Q^{\pi_k}(s, a) = \nabla_a r(s, a) - \beta a + \gamma \int_{\mathcal{S}} V^{\pi_k}(s') \nabla_a p(s' | s, a) \lambda(ds'),$$

these constants cannot be read off from the Wasserstein gradient of a static functional; they must be derived from the Bellman recursion and propagated along the WPG iterates. Assumptions 4.1-4.2, together with the a priori bounds that will be proved in Proposition 4.1, provide this uniform control.

Assumption 4.1 (Bounded rewards and initialization) *We assume:*

- (i) $|r(s, a)| \leq R_{\max}$ for all $(s, a) \in \mathcal{S} \times \mathbb{R}^d$.
- (ii) $\sup_{s \in \mathcal{S}} \text{KL}(\pi_0(\cdot | s) \| \rho_\beta) \leq K_0$.

For constants used later, set

$$M_0 := \sup_{s \in \mathcal{S}} \mathbb{E} \|A_0^s\|^2, \quad A_0^s \sim \pi_0(\cdot | s).$$

This quantity is finite by Lemma E.1 applied uniformly in s . In particular, applying that lemma with $c = \beta/(4\tau)$ and using Assumption 4.1(ii) gives

$$M_0 \leq \frac{4\tau}{\beta} \left(K_0 + \log \int_{\mathbb{R}^d} \exp \left(\frac{\beta}{4\tau} \|a\|^2 \right) \rho_\beta(a) da \right) = \frac{4\tau}{\beta} \left(K_0 + \frac{d}{2} \log 2 \right).$$

Assumption 4.2 (Action regularity) *There is a σ -finite reference measure λ on $\mathcal{S}$ such that $P(ds' | s, a) = p(s' | s, a)\lambda(ds')$. There are finite constants L_r, L_p, G_r, G_p such that:*

(i) *For every $s \in \mathcal{S}$, the map $a \mapsto r(s, a)$ is continuously differentiable and*

$$\|\nabla_a r(s, a)\| \leq G_r, \quad \|\nabla_a r(s, a) - \nabla_a r(s, \bar{a})\| \leq L_r \|a - \bar{a}\|.$$

(ii) *For every $s \in \mathcal{S}$ and λ -a.e. $s' \in \mathcal{S}$, the map $a \mapsto p(s' | s, a)$ is continuously differentiable, and*

$$\int_{\mathcal{S}} \|\nabla_a p(s' | s, a)\| \lambda(ds') \leq G_p,$$

$$\int_{\mathcal{S}} \|\nabla_a p(s' | s, a) - \nabla_a p(s' | s, \bar{a})\| \lambda(ds') \leq L_p \|a - \bar{a}\|.$$

A concrete example satisfying the transition part of Assumption 4.2 is the following additive-noise model. Let $\mathcal{S} = \mathbb{R}^m$, let λ be Lebesgue measure, and suppose

$$S' = F(s, a) + \sigma\zeta,$$

where $\sigma \in \mathbb{R}^{m \times m}$ is invertible and ζ has a smooth density q . Write

$$q_\sigma(x) := |\det \sigma|^{-1} q(\sigma^{-1}x)$$

for the density of $\sigma\zeta$, and assume

$$\|\nabla q_\sigma\|_{L^1} < \infty, \quad \|D^2 q_\sigma\|_{L^1} := \int_{\mathbb{R}^m} \|D^2 q_\sigma(x)\|_{\text{op}} dx < \infty.$$

Assume further that, uniformly in s ,

$$\|D_a F(s, a)\|_{\text{op}} \leq M_1, \quad \|D_a F(s, a) - D_a F(s, \bar{a})\|_{\text{op}} \leq M_2 \|a - \bar{a}\|.$$

Then

$$p(y | s, a) = q_\sigma(y - F(s, a))$$

satisfies Assumption 4.2(ii). Indeed,

$$\nabla_a p(y | s, a) = -D_a F(s, a)^\top \nabla q_\sigma(y - F(s, a)),$$

so one may take

$$G_p = M_1 \|\nabla q_\sigma\|_{L^1}, \quad L_p = M_2 \|\nabla q_\sigma\|_{L^1} + M_1^2 \|D^2 q_\sigma\|_{L^1}.$$

Nondegenerate Gaussian noise is a special case, since its first derivative and Hessian are integrable. Together with the reward condition in Assumption 4.2(i), this gives a concrete class of controlled transition models covered by the theorem. Deterministic or degenerate-noise dynamics are not covered by this example.

With these assumptions in place, write

$$W_k := V^* - V^{\pi_k}, \quad E_k := \|W_k\|_\infty,$$

and define the current-policy Langevin drift

$$b_k(s, a) := \nabla_a Q^{\pi_k}(s, a).$$

The current-policy Gibbs stationarity condition is

$$\tau \nabla_a \log p_s^{\pi_k}(a) = \nabla_a Q^{\pi_k}(s, a) = b_k(s, a). \quad (15)$$

4.2 Proof sketch and main theorem

4.2.1 BELLMAN RESIDUAL-TO-OPTIMALITY GAP MECHANISM AND UNIFORM LSI

We first record the discrete counterparts of the residual and resolvent mechanisms from Section 3. They show that the KL to the current-policy Gibbs law is exactly the Bellman residual, that this residual controls the worst-state value gap, and that decreasing the same KL produces value improvement after applying the Bellman resolvent.

Lemma 4.1 (Bellman contraction of the residual) *Fix $k \geq 0$. Let $R_k := (\mathcal{T}^* V^{\pi_k}) - V^{\pi_k}$. Then, for every $s \in \mathcal{S}$,*

$$R_k(s) \geq W_k(s) - \gamma E_k.$$

Lemma 4.2 (Discrete Bellman resolvent) *Fix $k \geq 0$. Suppose π_k and the WPG update π_{k+1} are admissible, with measurability covered by the convention in Section 2, and the KL terms below are finite. Then*

$$(I - \gamma P^{\pi_{k+1}})(V^{\pi_{k+1}} - V^{\pi_k}) = g_k, \quad g_k(s) := (\mathcal{T}^{\pi_{k+1}} V^{\pi_k})(s) - V^{\pi_k}(s), \quad (16)$$

and, for every $s \in \mathcal{S}$,

$$g_k(s) = \tau \left(\text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) - \text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) \right). \quad (17)$$

As a consequence of these two lemmas, once $p_s^{\pi_k}$ satisfies an LSI with constant $\bar{\alpha}$, the Bellman residual yields the following PL-type residual-to-gap inequality at an ε -maximizer of W_k :

$$\mathcal{I}(\pi_k(\cdot | s) \| p_s^{\pi_k}) \geq 2\bar{\alpha} \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) = \frac{2\bar{\alpha}}{\tau} R_k(s) \geq \frac{2\bar{\alpha}}{\tau} ((1 - \gamma)E_k - \varepsilon). \quad (18)$$

This is a PL inequality for the Bellman gap, not a consequence of flat convexity of J .

The LSI used in (18) follows from the same Gaussian perturbation mechanism as in continuous time, but it must be made uniform along the discrete trajectory. The a priori bounds prove a uniform value bound; once $\|V^{\pi_k}\|_\infty \leq V_{\max}$,

$$p_s^{\pi_k}(a) = \frac{e^{\psi_{k,s}(a)} \rho_\beta(a)}{\int e^{\psi_{k,s}(\tilde{a})} \rho_\beta(\tilde{a}) d\tilde{a}}, \quad \psi_{k,s}(a) := \frac{r(s, a) + \gamma \int V^{\pi_k}(s') P(ds' | s, a)}{\tau}.$$

The perturbation $\psi_{k,s}$ may be nonconvex, but bounded rewards and the value bound give $\|\psi_{k,s}\|_\infty \leq (R_{\max} + \gamma V_{\max})/\tau$ and hence $\text{osc}(\psi_{k,s}) \leq 2(R_{\max} + \gamma V_{\max})/\tau$. The Holley-Stroock bounded-perturbation principle (Holley and Stroock, 1987) then transfers the Gaussian LSI of ρ_β to every $p_s^{\pi_k}$ uniformly.

Lemma 4.3 (Uniform LSI) *Fix $k \geq 0$ and suppose $\|V^{\pi_k}\|_\infty \leq V_{\max}$. Then, for every $s \in \mathcal{S}$, the current-policy Gibbs density $p_s^{\pi_k}$ satisfies*

$$\mathcal{I}(\mu \| p_s^{\pi_k}) \geq 2\alpha(V_{\max}) \text{KL}(\mu \| p_s^{\pi_k}) \quad \forall \mu \ll p_s^{\pi_k},$$

where

$$\alpha(V_{\max}) = \frac{\beta}{\tau} \exp \left(-\frac{2(R_{\max} + \gamma V_{\max})}{\tau} \right).$$

4.2.2 DRIFT REGULARITY AND THE DISCRETIZATION ERROR

We now control the finite-step error introduced by the Langevin discretization. As explained in Section 4.1, the drift Langevin interpolation requires uniform action-Lipschitz and dissipativity estimates for the policy-dependent drift. The following lemma converts the action-regularity assumptions and a value bound into those estimates.

Lemma 4.4 (Bellman drift regularity) *Assume Assumption 4.2. Fix $k \geq 0$ and suppose $\|V^{\pi_k}\|_\infty \leq V_{\max} < \infty$. Then, for every $s \in \mathcal{S}$,*

$$b_k(s, a) = -\beta a + h_k(s, a),$$

with

$$\sup_{s,a} \|h_k(s, a)\| \leq G(V_{\max}) := G_r + \gamma G_p V_{\max}. \quad (19)$$

Moreover, for all $a, \bar{a} \in \mathbb{R}^d$,

$$\|b_k(s, a) - b_k(s, \bar{a})\| \leq L_b(V_{\max}) \|a - \bar{a}\|, \quad L_b(V_{\max}) := \beta + L_r + \gamma L_p V_{\max}.$$

The Lipschitz bound controls the Euler interpolation error; the dissipative decomposition controls the second moment and hence the squared drift moment. Together with the LSI from Lemma 4.3, they give the only place where discrete time loses exact monotonicity. The analytic part of the next lemma is based on a one-step fixed-drift interpolation. The full calculation is given in Appendix D.2.

Lemma 4.5 (One-step KL contraction with discretization error) *Fix $k \geq 0$ and $s \in \mathcal{S}$. Suppose $p_s^{\pi_k} > 0$, $\log p_s^{\pi_k} \in C^1(\mathbb{R}^d)$, $p_s^{\pi_k}$ satisfies an LSI with constant $\alpha > 0$, and the Gibbs-score identity*

$$\tau \nabla_a \log p_s^{\pi_k}(a) = b_k(s, a) \quad \text{for all } a \in \mathbb{R}^d$$

holds. Suppose also that $b_k(s, \cdot)$ is L_b -Lipschitz and

$$\mathbb{E}_{a \sim \pi_k(\cdot | s)} \|a\|^2 < \infty, \quad \mathbb{E}_{a \sim \pi_k(\cdot | s)} \|b_k(s, a)\|^2 \leq B^2, \quad \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) < \infty.$$

Set

$$\delta_\eta(L_b, B^2) := \frac{L_b^2 d}{2} \eta^2 + \frac{L_b^2 B^2}{6\tau} \eta^3.$$

Then

$$\text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) \leq e^{-\alpha \tau \eta} \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) + \delta_\eta(L_b, B^2). \quad (20)$$

Lemma 4.6 (From KL contraction to Bellman improvement) *Fix $k \geq 0$ and $s \in \mathcal{S}$. Suppose the admissibility and finite-KL hypotheses of Lemma 4.2 hold for (π_k, π_{k+1}) . If, for some $c \in [0, 1]$ and $\delta \geq 0$,*

$$\text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) \leq (1 - c) \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) + \delta, \quad (21)$$

then, with $R_k(s) = \tau \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k})$ and g_k as in Lemma 4.2,

$$g_k(s) \geq c R_k(s) - \tau \delta. \quad (22)$$

In particular, Lemma 4.5 supplies (21) with $c = 1 - e^{-\alpha \tau \eta}$ and $\delta = \delta_\eta(L_b, B^2)$ whenever its analytic hypotheses also hold.

With the uniform constants supplied by Proposition 4.1, the preceding lemmas yield a closed one-step recursion for the sup-norm gap. Throughout this argument, $\bar{\alpha}$ denotes the trajectory-uniform LSI constant, $\bar{L}_b$ the uniform Lipschitz constant of the drift, $\bar{B}^2$ the uniform second-moment bound for the drift, and $\bar{\delta}_\eta$ the corresponding one-step Langevin discretization error. Set

$$c_\eta := 1 - e^{-\bar{\alpha}\tau\eta}.$$

Then, for every $s \in \mathcal{S}$, the estimates compose as follows:

$$\begin{aligned} \text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) &\leq (1 - c_\eta) \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) + \bar{\delta}_\eta \\ &\quad \text{by Lemma 4.5,} \\ \implies g_k(s) &\geq c_\eta R_k(s) - \tau \bar{\delta}_\eta \\ &\quad \text{by Lemma 4.6,} \\ \implies V^{\pi_{k+1}}(s) - V^{\pi_k}(s) &\geq c_\eta R_k(s) - \frac{\tau}{1 - \gamma} \bar{\delta}_\eta \quad (23) \\ &\quad \text{by Lemma 4.2,} \\ \implies E_{k+1} &\leq (1 - (1 - \gamma)c_\eta) E_k + \frac{\tau}{1 - \gamma} \bar{\delta}_\eta \\ &\quad \text{by Lemma 4.1.} \end{aligned}$$

Here the KL terms are statewise, $R_k = \mathcal{T}^* V^{\pi_k} - V^{\pi_k}$, and $E_k = \|V^* - V^{\pi_k}\|_\infty$. The substantive RL step is the middle of (23): a local KL decrease becomes a global value-gap contraction through the Bellman residual and the value resolvent, not through flat convexity of a probability functional. The implications in (23) reduce the convergence proof to estimates that hold uniformly along the WPGD trajectory. The constants entering this recursion are policy dependent, since both $p_s^{\pi_k}$ and b_k are defined through Q^{π_k} and hence through the Bellman fixed point V^{π_k} . The next proposition supplies the required a priori estimates. Approximate value monotonicity controls the lower value envelope, the quadratic action penalty gives uniform action-moment bounds, and these estimates feed back into uniform drift regularity, finite KL controls, and a trajectory-uniform LSI for the moving Gibbs family.

Proposition 4.1 (Uniform a priori bounds for the WPGD iterates) *Under Assumptions 4.1 and 4.2, there exist explicit finite constants*

$$\bar{V}, \quad \bar{L}_b, \quad \bar{G}, \quad \bar{M}, \quad \bar{B}, \quad \bar{\alpha} > 0, \quad \eta_0 > 0,$$

depending only on the problem data and the initialization such that, for every $0 < \eta \leq \eta_0$, the WPG iterates are admissible and, for all $k \geq 0$,

$$\|V^{\pi_k}\|_\infty \leq \bar{V}, \quad \sup_{s \in \mathcal{S}} \mathbb{E} \|A_k^s\|^2 \leq \bar{M}, \quad \sup_{s \in \mathcal{S}} \mathbb{E}_{\pi_k} \|\nabla_a Q^{\pi_k}(s, A)\|^2 \leq \bar{B}^2.$$

Moreover, $a \mapsto \nabla_a Q^{\pi_k}(s, a)$ is $\bar{L}_b$ -Lipschitz uniformly in (k, s) , every current-policy Gibbs density $p_s^{\pi_k}$ satisfies LSI with constant at least $\bar{\alpha}$, and the generated policies satisfy

$$\sup_{k \geq 0} \sup_{s \in \mathcal{S}} \text{KL}(\pi_k(\cdot | s) \| \rho_\beta) \leq \bar{K}_\eta, \quad (24)$$

$$\sup_{k \geq 0} \sup_{s \in \mathcal{S}} \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) \leq \bar{H}_\eta. \quad (25)$$

All constants and the feasible step-size range are given in Appendix C.

Proposition 4.1 provides the uniform constants needed to run the one-step recursion in (23) at every iteration. The value bound yields a common Holley–Stroock perturbation bound and hence a uniform LSI constant $\bar{\alpha}$; the drift Lipschitz and second-moment estimates give a uniform discretization error $\bar{\delta}_\eta = O(\eta^2)$; and the admissibility and finite-KL bounds ensure that the resolvent identity and the one step drift interpolation estimate apply throughout the trajectory. Substituting these uniform estimates into (23) gives a closed scalar recursion for the sup-norm optimality gap, which yields the geometric contraction up to the accumulated Langevin discretization bias stated in the main theorem.

Theorem 4.1 (Global convergence of WPGD up to discretization bias) *Under Assumptions 4.1 and 4.2, there exist explicit finite constants $\bar{\alpha}, C_\delta, \eta_0 > 0$, such that for every $0 < \eta \leq \eta_0$ and every $k \geq 0$,*

$$J(\pi^*) - J(\pi_k) \leq \|V^* - V^{\pi_k}\|_\infty \leq \exp\left(-\frac{1}{2}\bar{\alpha}\tau(1-\gamma)\eta k\right)\|V^* - V^{\pi_0}\|_\infty + \frac{2C_\delta}{\bar{\alpha}(1-\gamma)^2}\eta.$$

Theorem 4.1 gives a finite-step geometric contraction for the WPGD update (WPGD), up to a fixed-step discretization bias. The drift KL estimate has an additive error $\bar{\delta}_\eta = O(\eta^2)$, and the scalar gap recursion divides this error by $1 - e^{-\bar{\alpha}\tau\eta} \asymp \bar{\alpha}\tau\eta$. The proof in Appendix D.4 keeps the sharper quantity $\bar{\delta}_\eta$ throughout and substitutes $\bar{\delta}_\eta \leq C_\delta\eta^2$ only at the end.

The theorem is closest in form to discrete-time mean-field Langevin guarantees, but the source of the contraction is different. Mean-field convex analyses convert KL decrease to objective decrease through flat convexity of a distributional objective (Nitanda et al., 2022; Chizat, 2022; Suzuki et al., 2023). Here the Gibbs law $p_s^{\pi_k}$ moves with the policy through the soft Bellman equation that depends on both the policy gradient and the KL regularization. The required PL-type relation is therefore reconstructed from the RL-specific ingredients.

The result is also complementary to entropy-regularized natural policy gradient and policy mirror descent, where KL/Bregman geometry gives a closed-form policy-improvement step and Bellman monotonicity (Geist et al., 2019; Cen et al., 2022; Bhandari and Russo, 2021; Lan, 2023; Zhan et al., 2023). WPGD instead uses Wasserstein geometry: local progress is Fisher-information dissipation in action space, and the main technical burden is to show that this transport-based dissipation remains uniformly controlled under the policy-dependent Bellman recursion and relate it to the global optimality gap.

5 Conclusion

We developed a convergence analysis framework for Wasserstein policy gradient in entropy-regularized discounted RL with continuous actions. The proof combines Bellman-residual identities, value-resolvent relation, one-step interpolation, uniform value/Q-function and moment bounds, and uniform LSI for the iterated Gibbs family. These ingredients yield exponential convergence for the continuous-time WPG flow and geometric convergence for the discrete-time WPG Langevin dynamics up to a discretization bias. An important direction for future work is to integrate these policy-space guarantees with practical approximations, such as projection onto parametric policy classes and finite-sample critic estimation.

References

- Alekh Agarwal, Sham M. Kakade, Jason D. Lee, and Gaurav Mahajan. On the theory of policy gradient methods: Optimality, approximation, and distribution shift. *Journal of Machine Learning Research*, 22(98):1–76, 2021.
- Shun-ichi Amari. Natural gradient works efficiently in learning. *Neural Computation*, 10(2):251–276, 1998.
- Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient Flows: In Metric Spaces and in the Space of Probability Measures*. Lectures in Mathematics ETH Zürich. Birkhäuser, 2 edition, 2008.
- Michael Arbel, Adil Salim, Anna Korba, and Zaid Harchaoui. Kernelized wasserstein natural gradient. In *International Conference on Learning Representations (ICLR)*, 2020.
- Jalaj Bhandari and Daniel Russo. On the linear convergence of policy gradient methods for finite MDPs. In *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 130 of *Proceedings of Machine Learning Research*, pages 2386–2394. PMLR, 2021.
- Jalaj Bhandari and Daniel Russo. Global optimality guarantees for policy gradient methods. *Operations Research*, 72(5):1906–1927, 2024. doi: 10.1287/opre.2021.0014.
- Greg Brockman, Vicki Cheung, Ludwig Pettersson, Jonas Schneider, John Schulman, Jie Tang, and Wojciech Zaremba. Openai gym. *arXiv preprint arXiv:1606.01540*, 2016.
- Yang Cai, Siddharth Mitra, Xiuyuan Wang, and Andre Wibisono. On the convergence of min-max langevin dynamics and algorithm. In *Proceedings of Thirty Eighth Conference on Learning Theory*, volume 291 of *Proceedings of Machine Learning Research*, pages 677–754. PMLR, 2025.
- Shicong Cen, Chen Cheng, Yuxin Chen, Yuting Wei, and Yuejie Chi. Fast global convergence of natural policy gradient methods with entropy regularization. *Operations Research*, 70(4):2563–2578, 2022. doi: 10.1287/opre.2021.2151.
- Lénaïc Chizat. Mean-field langevin dynamics: Exponential convergence and annealing. *arXiv preprint arXiv:2202.01009*, 2022.
- Lénaïc Chizat and Francis Bach. On the global convergence of gradient descent for over-parameterized models using optimal transport. *arXiv preprint arXiv:1805.09545*, 2018.
- Yuhao Ding, Junzi Zhang, and Javad Lavaei. On the global convergence of momentum-based policy gradient. In *Proceedings of The 25th International Conference on Artificial Intelligence and Statistics*, pages 1910–1934. PMLR, 2022.
- Matthieu Geist, Bruno Scherrer, and Olivier Pietquin. A theory of regularized markov decision processes. In *Proceedings of the 36th International Conference on Machine Learning (ICML)*, 2019.

- Tuomas Haarnoja, Aurick Zhou, Pieter Abbeel, and Sergey Levine. Soft actor-critic: Off-policy maximum entropy deep reinforcement learning with a stochastic actor. In *Proceedings of the 35th International Conference on Machine Learning (ICML)*, 2018.
- Richard Holley and Daniel Stroock. Logarithmic sobolev inequalities and stochastic ising models. *Journal of Statistical Physics*, 46(5–6):1159–1194, 1987.
- Sham M. Kakade. A natural policy gradient. In *Advances in Neural Information Processing Systems 14 (NeurIPS 2001)*, 2001.
- Sajad Khodadadian, Prakirt Raj Jhunjunwala, Sushil Mahavir Varma, and Siva Theja Maguluri. On the linear convergence of natural policy gradient algorithm. In *2021 60th IEEE Conference on Decision and Control (CDC)*, pages 3794–3799, 2021. doi: 10.1109/CDC45484.2021.9682908.
- Sajad Khodadadian, Prakirt Raj Jhunjunwala, Sushil Mahavir Varma, and Siva Theja Maguluri. On linear and super-linear convergence of natural policy gradient algorithm. *Systems & Control Letters*, 164:105214, 2022. doi: 10.1016/j.sysconle.2022.105214.
- Guanghui Lan. Policy mirror descent for reinforcement learning: Linear convergence, new sampling complexity, and generalized problem classes. *Mathematical Programming*, 198(1):1059–1106, 2023.
- Guanghui Lan, Yan Li, and Tuo Zhao. Block policy mirror descent. *SIAM Journal on Optimization*, 33(3):2341–2378, 2023. doi: 10.1137/22M1480409.
- Timothy P. Lillicrap, Jonathan J. Hunt, Alexander Pritzel, Nicolas Heess, Tom Erez, Yuval Tassa, David Silver, and Daan Wierstra. Continuous control with deep reinforcement learning. *arXiv preprint arXiv:1509.02971*, 2015.
- Song Mei, Andrea Montanari, and Phan-Minh Nguyen. A mean field view of the landscape of two-layer neural networks. *Proceedings of the National Academy of Sciences*, 115(33):E7665–E7671, 2018.
- Ted Moskovitz et al. Efficient wasserstein natural gradient for policy optimization. *arXiv preprint arXiv:2010.05380*, 2020.
- Atsushi Nitanda, Denny Wu, and Taiji Suzuki. Convex analysis of the mean field langevin dynamics. In *International Conference on Artificial Intelligence and Statistics*, pages 9741–9757. PMLR, 2022.
- David Pfau, Ian Davies, Diana L. Borsa, João Guilherme Madeira Araújo, Brendan Daniel Tracey, and Hado van Hasselt. Wasserstein policy optimization. *arXiv preprint arXiv:2505.00663*, 2025.
- John Schulman, Sergey Levine, Philipp Moritz, Michael I. Jordan, and Pieter Abbeel. Trust region policy optimization. In *Proceedings of the 32nd International Conference on Machine Learning (ICML)*, 2015.

- John Schulman, Filip Wolski, Prafulla Dhariwal, Alec Radford, and Oleg Klimov. Proximal policy optimization algorithms. *arXiv preprint arXiv:1707.06347*, 2017.
- David Šiška and Yufei Zhang. A note on convergence of wasserstein policy optimization. *arXiv preprint arXiv:2605.22622*, 2026.
- Jun Song, Niao He, Lijun Ding, and Chaoyue Zhao. Provably convergent policy optimization via metric-aware trust region methods. *Transactions on Machine Learning Research*, 2024. Accepted; initial version 2023.
- Richard S. Sutton, David A. McAllester, Satinder P. Singh, and Yishay Mansour. Policy gradient methods for reinforcement learning with function approximation. In *Advances in Neural Information Processing Systems 12 (NIPS 1999)*, pages 1057–1063, 2000.
- Taiji Suzuki, Denny Wu, and Atsushi Nitanda. Convergence of mean-field langevin dynamics: Time and space discretization, stochastic gradient, and variance reduction. In *Advances in Neural Information Processing Systems*, 2023.
- Antonio Terpin, Nicolas Lanzetti, Batuhan Yardim, Florian Dörfler, and Giorgia Ramponi. Trust region policy optimization with optimal transport discrepancies: Duality and algorithm for continuous actions. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2022.
- Saran Tunyasuvunakool, Alistair Muldal, Yotam Doron, Siqi Liu, Steven Bohez, Josh Merel, Tom Erez, Timothy Lillicrap, Nicolas Heess, and Yuval Tassa. dm_control: Software and tasks for continuous control. *Software Impacts*, 6:100022, 2020.
- Santosh Vempala and Andre Wibisono. Rapid convergence of the unadjusted langevin algorithm: Isoperimetry suffices. In *Advances in Neural Information Processing Systems*, volume 32, 2019.
- Cédric Villani. *Optimal Transport: Old and New*, volume 338 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2009.
- Ronald J. Williams. Simple statistical gradient-following algorithms for connectionist reinforcement learning. *Machine Learning*, 8(3–4):229–256, 1992.
- Lin Xiao. On the convergence rates of policy gradient methods. *Journal of Machine Learning Research*, 23:1–36, 2022.
- Takei Yamamoto, Kazusato Oko, Zhuoran Yang, and Taiji Suzuki. Mean field langevin actor-critic: Faster convergence and global optimality beyond lazy learning. In *Forty-first International Conference on Machine Learning*, 2024.
- Rui Yuan, Simon S. Du, Robert M. Gower, Alessandro Lazaric, and Lin Xiao. Linear convergence of natural policy gradient methods with log-linear policies. In *International Conference on Learning Representations (ICLR)*, 2023.

Wenhao Zhan, Shicong Cen, Baihe Huang, Yuxin Chen, Jason D. Lee, and Yuejie Chi. Policy mirror descent for regularized reinforcement learning: A generalized framework with linear convergence. *SIAM Journal on Optimization*, 33(2):1061–1091, 2023. doi: 10.1137/21M1456789.

Junyu Zhang, Alec Koppel, Amrit Singh Bedi, Csaba Szepesvari, and Mengdi Wang. Variational policy gradient method for reinforcement learning with general utilities. In *Advances in Neural Information Processing Systems*, volume 33, pages 4572–4583, 2020.

Ruiyi Zhang, Changyou Chen, Chunyuan Li, and Lawrence Carin. Policy optimization as wasserstein gradient flows. In *International Conference on machine learning*, pages 5737–5746. PMLR, 2018.

Zhaoyu Zhu, Shuhan Zhang, Rui Gao, and Shuang Li. Wasserstein proximal policy gradient. *arXiv preprint arXiv:2603.02576*, 2026.

Brian D. Ziebart. *Modeling Purposeful Adaptive Behavior with the Principle of Maximum Causal Entropy*. PhD thesis, Carnegie Mellon University, 2010.

Appendix A. Auxiliary Results

A.1 Entropy-Regularized Performance Difference Lemma

Lemma A.1 (Entropy-Regularized Performance Difference Lemma) ([Lan, 2023](#), Lemma2)

Let $\rho_0 \in \mathcal{P}(S)$ be the initial state distribution, and write $J(\pi) := J_{\rho_0}(\pi) = \int_S V^\pi(s) \rho_0(ds)$.

For any two feasible policies π and π' , we have

$$J(\pi') - J(\pi) = \frac{1}{1 - \gamma} \int_S \left[\int_{\mathbb{R}^d} Q^\pi(s, a) (\pi'(a | s) - \pi(a | s)) da - \tau H^{\pi'}(s) + \tau H^\pi(s) \right] d_{\rho_0}^{\pi'}(ds),$$

where

$$H^\pi(s) := \int_{\mathbb{R}^d} \pi(a | s) \log \pi(a | s) da$$

is the statewise negative entropy, and

$$d_{\rho_0}^{\pi'}(B) := (1 - \gamma) \sum_{t=0}^{\infty} \gamma^t \mathbb{P}_{\rho, \pi'}(s_t \in B), \quad B \in \mathcal{B}(S),$$

is the normalized discounted state occupancy measure of π' initialized from ρ .

For completeness, we provide the proof here and adapt it to our notation.

Proof Let $\Gamma^{\pi'}(\rho_0)$ denote the trajectory $(s_t, a_t, s_{t+1})_{t \geq 0}$ generated by $s_0 \sim \rho_0$, $a_t \sim \pi'(\cdot | s_t)$, and $s_{t+1} \sim P(\cdot | s_t, a_t)$. By the definition of J , we have

$$J(\pi') - J(\pi) = \mathbb{E}_{\Gamma^{\pi'}(\rho_0)} \left[\sum_{t=0}^{\infty} \gamma^t (\tilde{r}(s_t, a_t) - \tau \log \pi'(a_t | s_t)) \right] - \mathbb{E}_{s_0 \sim \rho_0} [V^\pi(s_0)].$$

Adding and subtracting $V^\pi(s_t)$ inside the summation and rearranging the telescoping value terms gives

$$\begin{aligned} J(\pi') - J(\pi) &= \mathbb{E}_{\Gamma^{\pi'}(\rho_0)} \left[\sum_{t=0}^{\infty} \gamma^t (\tilde{r}(s_t, a_t) - \tau \log \pi'(a_t | s_t) + \gamma V^\pi(s_{t+1}) - V^\pi(s_t)) \right] \\ &\quad + \mathbb{E}_{\Gamma^{\pi'}(\rho_0)} [V^\pi(s_0)] - \mathbb{E}_{s_0 \sim \rho_0} [V^\pi(s_0)]. \end{aligned}$$

The last two terms cancel. Taking conditional expectation over s_{t+1} given (s_t, a_t) and using

$$Q^\pi(s_t, a_t) = \tilde{r}(s_t, a_t) + \gamma \mathbb{E}[V^\pi(s_{t+1}) | s_t, a_t],$$

we obtain

$$J(\pi') - J(\pi) = \mathbb{E}_{\Gamma^{\pi'}(\rho_0)} \left[\sum_{t=0}^{\infty} \gamma^t (Q^\pi(s_t, a_t) - V^\pi(s_t) - \tau \log \pi'(a_t | s_t)) \right].$$

For each fixed s_t , since $a_t \sim \pi'(\cdot | s_t)$ and

$$V^\pi(s_t) = \int_{\mathbb{R}^d} (Q^\pi(s_t, a) - \tau \log \pi(a | s_t)) \pi(a | s_t) da,$$

we have

$$\begin{aligned} &\mathbb{E} [Q^\pi(s_t, a_t) - V^\pi(s_t) - \tau \log \pi'(a_t | s_t) | s_t] \\ &= \int_{\mathbb{R}^d} Q^\pi(s_t, a) (\pi'(a | s_t) - \pi(a | s_t)) da - \tau H^{\pi'}(s_t) + \tau H^\pi(s_t). \end{aligned}$$

Substituting this identity into the trajectory expectation yields

$$J(\pi') - J(\pi) = \mathbb{E}_{\Gamma^{\pi'}(\rho_0)} \left[\sum_{t=0}^{\infty} \gamma^t \left(\int_{\mathbb{R}^d} Q^\pi(s_t, a) (\pi'(a | s_t) - \pi(a | s_t)) da - \tau H^{\pi'}(s_t) + \tau H^\pi(s_t) \right) \right].$$

Finally, by the definition of $d_{\rho_0}^{\pi'}$, the last display is equivalent to

$$J(\pi') - J(\pi) = \frac{1}{1 - \gamma} \int_S \left[\int_{\mathbb{R}^d} Q^\pi(s, a) (\pi'(a | s) - \pi(a | s)) da - \tau H^{\pi'}(s) + \tau H^\pi(s) \right] d_{\rho_0}^{\pi'}(ds).$$

This proves the lemma. ■

A.2 Derivation of the first variation and Wasserstein policy-gradient direction

We justify the statewise direction used in Section 2.2. Throughout this subsection, we write $J(\pi) = J_{\rho_0}(\pi)$ for the entropy-regularized objective with initial distribution ρ_0 . Fix a policy π , and let $d^\pi = d_{\rho_0}^\pi$ be its normalized discounted state occupancy.

For each state s , consider a smooth transport perturbation $t \mapsto \pi_t(\cdot | s)$ with $\pi_0 = \pi$ and velocity field $u(s, \cdot)$, so that

$$\partial_t \pi_t(a | s) \big|_{t=0} = -\nabla_a \cdot (\pi(a | s) u(s, a)). \quad (26)$$

The following proposition identifies the first variation of J along such statewise perturbations.

Proposition A.1 (First variation under statewise transport perturbations) *Let $t \mapsto \pi_t$ be a smooth curve of feasible policies with $\pi_0 = \pi$, and write*

$$\dot{\pi}(a | s) := \partial_t \pi_t(a | s) \Big|_{t=0}.$$

Then

$$\frac{d}{dt} J(\pi_t) \Big|_{t=0} = \frac{1}{1-\gamma} \int_{\mathcal{S}} \int_{\mathbb{R}^d} (Q^\pi(s, a) - \tau \log \pi(a | s)) \dot{\pi}(a | s) da d^\pi(ds). \quad (27)$$

If, in addition, $\dot{\pi}$ is generated by the transport perturbation (26), then

$$\frac{d}{dt} J(\pi_t) \Big|_{t=0} = \frac{1}{1-\gamma} \int_{\mathcal{S}} \int_{\mathbb{R}^d} \langle \nabla_a (Q^\pi(s, a) - \tau \log \pi(a | s)), u(s, a) \rangle \pi(a | s) da d^\pi(ds). \quad (28)$$

Proof Apply the entropy-regularized performance difference lemma with $\pi' = \pi_t$. This gives

$$J(\pi_t) - J(\pi) = \frac{1}{1-\gamma} \int_{\mathcal{S}} \left[\int_{\mathbb{R}^d} Q^\pi(s, a) (\pi_t(a | s) - \pi(a | s)) da - \tau H^{\pi_t}(s) + \tau H^\pi(s) \right] d^{\pi_t}(ds).$$

Define

$$F_t(s) := \int_{\mathbb{R}^d} Q^\pi(s, a) (\pi_t(a | s) - \pi(a | s)) da - \tau H^{\pi_t}(s) + \tau H^\pi(s).$$

Then $F_0(s) = 0$ for every s . Hence, when differentiating $\int_{\mathcal{S}} F_t(s) d^{\pi_t}(ds)$ at $t = 0$, the derivative of the occupancy measure does not contribute, and we obtain

$$\frac{d}{dt} J(\pi_t) \Big|_{t=0} = \frac{1}{1-\gamma} \int_{\mathcal{S}} \dot{F}_0(s) d^\pi(ds).$$

It remains to compute $\dot{F}_0(s)$. Since

$$H^{\pi_t}(s) = \int_{\mathbb{R}^d} \pi_t(a | s) \log \pi_t(a | s) da,$$

we have

$$\frac{d}{dt} H^{\pi_t}(s) \Big|_{t=0} = \int_{\mathbb{R}^d} (1 + \log \pi(a | s)) \dot{\pi}(a | s) da.$$

Because $\int_{\mathbb{R}^d} \dot{\pi}(a | s) da = 0$, the constant term drops out, and therefore

$$\dot{F}_0(s) = \int_{\mathbb{R}^d} (Q^\pi(s, a) - \tau \log \pi(a | s)) \dot{\pi}(a | s) da.$$

Substituting this expression into the previous display proves (27).

Now suppose that $\dot{\pi}$ is given by (26). Substituting $\dot{\pi} = -\nabla_a \cdot (\pi u)$ into (27) and using integration by parts in divergence form with respect to the action variable yields

$$\int_{\mathbb{R}^d} (Q^\pi(s, a) - \tau \log \pi(a | s)) \dot{\pi}(a | s) da = \int_{\mathbb{R}^d} \langle \nabla_a (Q^\pi(s, a) - \tau \log \pi(a | s)), u(s, a) \rangle \pi(a | s) da.$$

This proves (28). ■

Thus, the action-space gradient direction associated with the first variation is

$$\nabla_a(Q^\pi(s, a) - \tau \log \pi(a | s)). \quad (29)$$

Equivalently, one may view (28) as taking the Wasserstein gradient with respect to the statewise 2-Wasserstein geometry on the conditional action laws, averaged over states using the policy-gradient weight d^π . This is the standard Otto-calculus interpretation of gradients on probability measures (Ambrosio et al., 2008; Villani, 2009); the appearance of d^π is the standard discounted state weighting from the policy-gradient theorem (Sutton et al., 2000), and analogous visitation-weighted averages appear in natural-gradient and trust-region policy optimization (Kakade, 2001; Schulman et al., 2015).

Under the full-support convention on ρ , the induced occupancy d^π has full support, so this statewise direction is determined on all states relevant to the sup-norm analysis. Substituting the direction (29) into the continuity equation gives the Wasserstein policy-gradient flow:

$$\partial_t \pi_t(a | s) = -\nabla_a \cdot (\pi_t(a | s) \nabla_a (Q^{\pi_t}(s, a) - \tau \log \pi_t(a | s))),$$

which is (WPGF).

Appendix B. Proofs for Section 3

Proof [Proof of Lemma 2.1] Fix $t \geq 0$ and $s \in \mathcal{S}$. Let

$$f(a) := Q^{\pi_t}(s, a), \quad Z := \int_{\mathcal{A}} \exp(f(a)/\tau) da,$$

and define the Gibbs density

$$p(a) := \frac{\exp(f(a)/\tau)}{Z},$$

so that $p(\cdot) = p_s^{\pi_t}(\cdot)$.

For any density μ on $\mathcal{A}$,

$$\text{KL}(\mu \| p) = \int_{\mathcal{A}} \mu(a) \log \frac{\mu(a)}{p(a)} da = \int_{\mathcal{A}} \mu(a) \log \mu(a) da - \int_{\mathcal{A}} \mu(a) \log p(a) da.$$

Since $\log p(a) = f(a)/\tau - \log Z$, we have

$$\int_{\mathcal{A}} \mu(a) \log p(a) da = \frac{1}{\tau} \int_{\mathcal{A}} f(a) \mu(a) da - \log Z.$$

Substituting gives

$$\text{KL}(\mu \| p) = \int_{\mathcal{A}} \mu(a) \log \mu(a) da - \frac{1}{\tau} \int_{\mathcal{A}} f(a) \mu(a) da + \log Z,$$

or equivalently,

$$\int_{\mathcal{A}} f(a) \mu(a) da - \tau \int_{\mathcal{A}} \mu(a) \log \mu(a) da = \tau \log Z - \tau \text{KL}(\mu \| p). \quad (30)$$

By definition of $\mathcal{T}^*$ (as the supremum over policies applied to the evaluation functional at V^{π_t}),

$$(\mathcal{T}^*V^{\pi_t})(s) = \sup_{\mu(\cdot|s)} \left\{ \int_{\mathcal{A}} Q^{\pi_t}(s, a) \mu(a|s) da - \tau \int_{\mathcal{A}} \mu(a|s) \log \mu(a|s) da \right\}.$$

Applying (30) with $\mu(\cdot|s)$ shows the supremum equals $\tau \log Z$ and is attained at $\mu = p$. Hence,

$$(\mathcal{T}^*V^{\pi_t})(s) = \tau \log Z.$$

On the other hand, V^{π_t} is the fixed point of $\mathcal{T}^{\pi_t}$, so

$$V^{\pi_t}(s) = (\mathcal{T}^{\pi_t}V^{\pi_t})(s) = \int_{\mathcal{A}} Q^{\pi_t}(s, a) \pi_t(a|s) da - \tau \int_{\mathcal{A}} \pi_t(a|s) \log \pi_t(a|s) da.$$

Applying (30) with $\mu(\cdot|s) = \pi_t(\cdot|s)$ yields

$$V^{\pi_t}(s) = \tau \log Z - \tau \text{KL}(\pi_t(\cdot|s) \| p).$$

Combining the last two displays gives

$$(\mathcal{T}^*V^{\pi_t})(s) - V^{\pi_t}(s) = \tau \text{KL}(\pi_t(\cdot|s) \| p_s^{\pi_t}),$$

which is nonnegative since KL divergence is nonnegative. ■

Proof [Proof of Lemma 3.1] Fix $t \geq 0$ and $s \in \mathcal{S}$. The soft policy-evaluation identity gives

$$V^{\pi_t}(s) = \int_{\mathbb{R}^d} Q^{\pi_t}(s, a) \pi_t(a|s) da - \tau \int_{\mathbb{R}^d} \pi_t(a|s) \log \pi_t(a|s) da. \quad (31)$$

Differentiating (31) in t gives

$$\begin{aligned} \dot{V}_t(s) &= \int_{\mathbb{R}^d} (\partial_t Q^{\pi_t})(s, a) \pi_t(a|s) da + \int_{\mathbb{R}^d} Q^{\pi_t}(s, a) \partial_t \pi_t(a|s) da \\ &\quad - \tau \int_{\mathbb{R}^d} \partial_t \pi_t(a|s) (1 + \log \pi_t(a|s)) da. \end{aligned} \quad (32)$$

We first evaluate the term containing $\partial_t Q^{\pi_t}$. Since

$$Q^{\pi_t}(s, a) = \tilde{r}(s, a) + \gamma \int_{\mathcal{S}} V^{\pi_t}(s') P(ds'|s, a),$$

and $\tilde{r}$ and P are time-independent,

$$(\partial_t Q^{\pi_t})(s, a) = \gamma \int_{\mathcal{S}} \dot{V}_t(s') P(ds'|s, a).$$

Therefore,

$$\int_{\mathbb{R}^d} (\partial_t Q^{\pi_t})(s, a) \pi_t(a|s) da = \gamma (P^{\pi_t} \dot{V}_t)(s).$$

It remains to evaluate the two terms in (32) involving $\partial_t \pi_t(\cdot | s)$. Since $\pi_t(\cdot | s)$ is a probability density for every t ,

$$\int_{\mathbb{R}^d} \partial_t \pi_t(a | s) da = 0.$$

Hence these two terms can be written as

$$\int_{\mathbb{R}^d} (Q^{\pi_t}(s, a) - \tau \log \pi_t(a | s)) \partial_t \pi_t(a | s) da.$$

By the Gibbs definition of $p_s^{\pi_t}$,

$$p_s^{\pi_t}(a) = \frac{\exp(Q^{\pi_t}(s, a)/\tau)}{\int_{\mathbb{R}^d} \exp(Q^{\pi_t}(s, \bar{a})/\tau) d\bar{a}},$$

and therefore

$$Q^{\pi_t}(s, a) = \tau \log p_s^{\pi_t}(a) + \tau \log \int_{\mathbb{R}^d} \exp(Q^{\pi_t}(s, \bar{a})/\tau) d\bar{a}.$$

The second term is independent of a , so it vanishes after integration against $\partial_t \pi_t(a | s)$. Thus

$$\begin{aligned} & \int_{\mathbb{R}^d} Q^{\pi_t}(s, a) \partial_t \pi_t(a | s) da - \tau \int_{\mathbb{R}^d} \partial_t \pi_t(a | s) (1 + \log \pi_t(a | s)) da \\ &= -\tau \int_{\mathbb{R}^d} \log \frac{\pi_t(a | s)}{p_s^{\pi_t}(a)} \partial_t \pi_t(a | s) da. \end{aligned}$$

Using the WPGF equation in Gibbs form,

$$\partial_t \pi_t(a | s) = \tau \nabla_a \cdot \left(\pi_t(a | s) \nabla_a \log \frac{\pi_t(a | s)}{p_s^{\pi_t}(a)} \right),$$

where

$$\tau \nabla_a \log p_s^{\pi_t}(a) = \nabla_a Q^{\pi_t}(s, a),$$

and applying the divergence-form integration-by-parts formula in the action variable, we obtain

$$\begin{aligned} & -\tau \int_{\mathbb{R}^d} \log \frac{\pi_t(a | s)}{p_s^{\pi_t}(a)} \partial_t \pi_t(a | s) da \\ &= -\tau^2 \int_{\mathbb{R}^d} \log \frac{\pi_t(a | s)}{p_s^{\pi_t}(a)} \nabla_a \cdot \left(\pi_t(a | s) \nabla_a \log \frac{\pi_t(a | s)}{p_s^{\pi_t}(a)} \right) da \\ &= \tau^2 \int_{\mathbb{R}^d} \left\| \nabla_a \log \frac{\pi_t(a | s)}{p_s^{\pi_t}(a)} \right\|^2 \pi_t(a | s) da \\ &= \tau^2 \mathcal{I}(\pi_t(\cdot | s) \| p_s^{\pi_t}). \end{aligned}$$

Thus the density-variation contribution is

$$g_t(s) := \tau^2 \mathcal{I}(\pi_t(\cdot | s) \| p_s^{\pi_t}).$$

Substituting the two evaluated contributions back into (32), we obtain

$$\dot{V}_t(s) = \gamma(P^{\pi_t} \dot{V}_t)(s) + g_t(s).$$

Equivalently,

$$(I - \gamma P^{\pi_t}) \dot{V}_t = g_t,$$

and hence

$$\dot{V}_t = (I - \gamma P^{\pi_t})^{-1} g_t.$$

Since P^{π_t} is positivity preserving and $\gamma \in (0, 1)$,

$$(I - \gamma P^{\pi_t})^{-1} = \sum_{n=0}^{\infty} \gamma^n (P^{\pi_t})^n$$

is also positivity preserving. Because $g_t(s) \geq 0$ for every s , we get

$$\dot{V}_t = g_t + \sum_{n=1}^{\infty} \gamma^n (P^{\pi_t})^n g_t \geq g_t \geq 0$$

pointwise on $\mathcal{S}$. This proves both the resolvent identity and the pointwise lower bound. ■

Appendix C. Uniform regularity bounds

This appendix verifies the quantitative bounds needed to use the core Bellman proof uniformly along WPGD. Pure measurability of kernels and conditional versions remains covered by the measurability convention in Section 2; Normalizability, Bellman differentiability, value bounds, drift bounds, moments, KL bounds, and LSI are proved below.

C.1 Explicit constants and feasible step sizes

For $V_{\max} \geq 0$, define

$$L_b(V_{\max}) := \beta + L_r + \gamma L_p V_{\max}, \quad G(V_{\max}) := G_r + \gamma G_p V_{\max}.$$

Let

$$U := \frac{R_{\max} + \tau \log Z_\beta}{1 - \gamma}, \quad L_\star := \frac{-R_{\max} + \tau \log Z_\beta}{1 - \gamma}, \quad \bar{E}_0 := \frac{2R_{\max} + \tau K_0}{1 - \gamma},$$

and set

$$\bar{V} := \max\{1, U, \bar{E}_0 - L_\star + 1\}. \quad (33)$$

The induced constants are

$$\bar{L}_b := L_b(\bar{V}), \quad \bar{G} := G(\bar{V}), \quad \bar{\alpha} := \frac{\beta}{\tau} \exp\left(-\frac{2(R_{\max} + \gamma \bar{V})}{\tau}\right), \quad c_\eta := 1 - e^{-\bar{\alpha} \tau \eta}.$$

With $M_0 = \sup_s \mathbb{E} \|A_0^s\|^2$, define

$$\bar{M}_\infty(\eta) := \frac{2}{\beta} \left(\frac{\bar{G}^2}{\beta} + 2\tau d \right) + \frac{4\bar{G}^2}{\beta} \eta, \quad B^2 := 2\beta^2 \max\{M_0 \bar{M}_\infty(\eta)\} + 2\bar{G}^2$$

The statewise one-step KL discretization error is

$$\bar{\delta}_\eta := \frac{\bar{L}_b^2 d}{2} \eta^2 + \frac{\bar{L}_b^2 B_\infty^2}{6\tau} \eta^3. \quad (34)$$

The auxiliary KL bounds are

$$\begin{aligned} \bar{K}_\eta &:= \max \left\{ K_0, 0, \frac{\beta M_\infty}{2\tau} + \log Z_\beta - \frac{d}{2} \log(4\pi e\tau\eta) \right\}, \\ \bar{H}_\eta &:= \bar{K}_\eta + \frac{2(R_{\max} + \gamma\bar{V})}{\tau}. \end{aligned}$$

The exact stability condition used in Proposition 4.1 is

$$0 < \eta \leq \frac{1}{4\beta}, \quad \bar{\alpha}\tau\eta \leq 1, \quad \frac{\tau}{(1-\gamma)^2} \frac{\bar{\delta}_\eta}{c_\eta} \leq 1. \quad (35)$$

For the simplified theorem statement, let

$$\begin{aligned} \bar{M} &:= \max \left\{ M_0, \frac{3\bar{G}^2}{\beta^2} + \frac{4\tau d}{\beta} \right\}, \quad \bar{B}^2 := 2\beta^2 \bar{M} + 2\bar{G}^2, \\ C_\delta &:= \frac{\bar{L}_b^2 d}{2} + \frac{\bar{L}_b^2 \bar{B}^2}{6\tau}, \end{aligned} \quad (36)$$

and

$$\eta_0 := \min \left\{ 1, \frac{1}{4\beta}, \frac{1}{\bar{\alpha}\tau}, \frac{\bar{\alpha}(1-\gamma)^2}{2C_\delta} \right\}. \quad (37)$$

For every $0 < \eta \leq \eta_0$, Lemma C.6 implies (35), and (34) satisfies $\bar{\delta}_\eta \leq C_\delta \eta^2$.

C.2 Bellman well-posedness and optimal Gibbs policies

Lemma C.1 (Soft optimality contraction and fixed point) *Under Assumptions 4.1 and 4.2, with the measurability convention of Section 2, the soft optimality operator $\mathcal{T}^*$ maps bounded measurable functions to bounded measurable functions, is monotone, and is a γ -contraction in $\|\cdot\|_\infty$:*

$$\|\mathcal{T}^*V - \mathcal{T}^*W\|_\infty \leq \gamma\|V - W\|_\infty.$$

Consequently, $\mathcal{T}^*$ has a unique bounded measurable fixed point V^* .

Proof Let $\mathcal{B}_b(\mathcal{S})$ denote the bounded measurable functions on $\mathcal{S}$ with the sup norm. The measurability part of the mapping property is covered by the convention in Section 2. The quantitative boundedness follows from the bounded-Gaussian representation used below: for bounded V ,

$$\exp(Q_V(s, a)/\tau) = Z_\beta \exp \left(\frac{r(s, a) + \gamma \int V(s') P(ds' | s, a)}{\tau} \right) \rho_\beta(a),$$

and the exponential factor multiplying ρ_β is uniformly bounded above and below by positive constants depending only on $\|V\|_\infty$.

Monotonicity is immediate from $V \leq W \Rightarrow Q_V \leq Q_W$. Since $Q_V(s, a) \leq Q_W(s, a) + \gamma \|V - W\|_\infty$ for all (s, a) ,

$$(\mathcal{T}^*V)(s) \leq (\mathcal{T}^*W)(s) + \gamma \|V - W\|_\infty.$$

Interchanging V and W gives $\|\mathcal{T}^*V - \mathcal{T}^*W\|_\infty \leq \gamma \|V - W\|_\infty$. The Banach fixed-point theorem on $\mathcal{B}_b(\mathcal{S})$ gives a unique bounded measurable fixed point V^* . $\blacksquare$

Lemma C.2 (Bellman well-posedness and WPG update) *Assume the bounded-reward condition in Assumption 4.1(i), the measurability convention in Section 2, and the action-regularity bounds in Assumption 4.2. Let $\pi \in \Pi$. Then the following statements hold.*

(i) *The policy-evaluation Bellman operator $\mathcal{T}^\pi$ is well-defined on bounded measurable functions, maps them to bounded functions, is a γ -contraction in $\|\cdot\|_\infty$, and has a unique bounded fixed point V^π .*

(ii) *For every state s , the action-value function*

$$Q^\pi(s, a) = r(s, a) - \frac{\beta}{2} \|a\|^2 + \gamma \int_S V^\pi(s') P(ds' | s, a)$$

is continuously differentiable in a , and

$$\nabla_a Q^\pi(s, a) = \nabla_a r(s, a) - \beta a + \gamma \int_S V^\pi(s') \nabla_a p(s' | s, a) \lambda(ds').$$

Moreover, if $\|V^\pi\|_\infty \leq V_{\max}$, then

$$\|\nabla_a Q^\pi(s, a) - \nabla_a Q^\pi(s, \bar{a})\| \leq (\beta + L_r + \gamma L_p V_{\max}) \|a - \bar{a}\|,$$

and

$$\nabla_a Q^\pi(s, a) = -\beta a + h^\pi(s, a), \quad \sup_{s, a} \|h^\pi(s, a)\| \leq G_r + \gamma G_p V_{\max}.$$

(iii) *For every bounded measurable $V : S \rightarrow \mathbb{R}$, the Gibbs kernel*

$$\mathcal{G}[V](a | s) = \frac{\exp(Q_V(s, a)/\tau)}{\int_{\mathbb{R}^d} \exp(Q_V(s, \tilde{a})/\tau) d\tilde{a}}$$

*is well-defined, belongs to Π , and attains the variational supremum defining $\mathcal{T}^*V$.*

(iv) *In particular, with V^* from C.1, the Gibbs kernel*

$$\pi^* := \mathcal{G}[V^*]$$

belongs to Π , satisfies $V^{\pi^} = V^*$, and is an optimal admissible policy.*

(v) *If $\pi_k \in \Pi$ and $b_k(s, a) := \nabla_a Q^{\pi_k}(s, a)$, then the WPG update has the statewise density*

$$\pi_{k+1}(y | s) = \int_{\mathbb{R}^d} \phi_{2\tau\eta}(y - a - \eta b_k(s, a)) \pi_k(a | s) da,$$

where

$$\phi_{2\tau\eta}(z) = (4\pi\tau\eta)^{-d/2} \exp\left(-\frac{\|z\|^2}{4\tau\eta}\right).$$

Equivalently,

$$A_{k+1}^s = A_k^s + \eta b_k(s, A_k^s) + \sqrt{2\tau\eta} \xi_{k+1}, \quad A_k^s \sim \pi_k(\cdot | s), \quad \xi_{k+1} \sim N(0, I_d).$$

For $t > 0$, the corresponding one step drift interpolation has density

$$q_t^s(y) = \int_{\mathbb{R}^d} \phi_{2\tau t}(y - a - tb_k(s, a)) \pi_k(a | s) da,$$

and its conditional drift is

$$u_t^s(y) = \frac{\int_{\mathbb{R}^d} b_k(s, a) \phi_{2\tau t}(y - a - tb_k(s, a)) \pi_k(a | s) da}{q_t^s(y)}$$

for $q_t^s(y)dy$ -a.e. y .

Finally, along the WPG sequence, admissibility of the updated densities is obtained from the moment and Gaussian-relative-entropy estimates in Proposition 4.1: under its step-size condition,

$$\sup_{k \geq 0} \sup_{s \in S} \text{KL}(\pi_k(\cdot | s) \| \rho_\beta) < \infty,$$

and hence all iterates remain in Π .

Proof [Proof of Lemma C.2] We prove the quantitative parts of the statement and leave only kernel measurability to the convention in Section 2.

First, for each state s , finite Gaussian-relative entropy gives $\pi(\cdot | s) \ll \rho_\beta \ll da$ and a finite second moment by Lemma E.1. Using

$$\log \rho_\beta(a) = -\log Z_\beta - \frac{\beta}{2\tau} \|a\|^2,$$

the one-step regularized reward can be written as the finite quantity

$$\begin{aligned} \bar{r}^\pi(s) &:= \int_{\mathbb{R}^d} \left(r(s, a) - \frac{\beta}{2} \|a\|^2 - \tau \log \pi(a | s) \right) \pi(a | s) da \\ &= \int_{\mathbb{R}^d} r(s, a) \pi(a | s) da + \tau \log Z_\beta - \tau \text{KL}(\pi(\cdot | s) \| \rho_\beta). \end{aligned}$$

Consequently, with $C_\pi := \sup_s \text{KL}(\pi(\cdot | s) \| \rho_\beta)$,

$$-R_{\max} + \tau \log Z_\beta - \tau C_\pi \leq \bar{r}^\pi(s) \leq R_{\max} + \tau \log Z_\beta. \quad (38)$$

For bounded V , the policy-evaluation operator is

$$(\mathcal{T}^\pi V)(s) = \bar{r}^\pi(s) + \gamma \int_{\mathbb{R}^d} \int_S V(s') p(s' | s, a) \lambda(ds') \pi(a | s) da.$$

The second term is bounded in absolute value by $\gamma \|V\|_\infty$, and (38) gives boundedness of $\mathcal{T}^\pi V$. Moreover,

$$\|\mathcal{T}^\pi V - \mathcal{T}^\pi W\|_\infty \leq \gamma \|V - W\|_\infty.$$

Thus $\mathcal{T}^\pi$ is a γ -contraction on the bounded measurable functions and has a unique bounded fixed point V^π .

Next, fix a bounded function V and define

$$\mathbf{P}_V(s, a) := \int_{\mathcal{S}} V(s') p(s' | s, a) \lambda(ds').$$

We verify the derivative-under-the-integral step. Fix s , a , and a direction $h \in \mathbb{R}^d$. For $\varepsilon \neq 0$, the fundamental theorem of calculus gives, for λ -a.e. s' ,

$$\frac{p(s' | s, a + \varepsilon h) - p(s' | s, a)}{\varepsilon} = \int_0^1 \langle \nabla_a p(s' | s, a + \theta \varepsilon h), h \rangle d\theta.$$

Subtracting the candidate directional derivative and integrating, we obtain

$$\begin{aligned} & \left| \frac{\mathbf{P}_V(s, a + \varepsilon h) - \mathbf{P}_V(s, a)}{\varepsilon} - \int_{\mathcal{S}} V(s') \langle \nabla_a p(s' | s, a), h \rangle \lambda(ds') \right| \\ & \leq \|V\|_\infty \int_0^1 \int_{\mathcal{S}} \|\nabla_a p(s' | s, a + \theta \varepsilon h) - \nabla_a p(s' | s, a)\| \lambda(ds') \|h\| d\theta \\ & \leq \frac{1}{2} \|V\|_\infty L_p |\varepsilon| \|h\|^2. \end{aligned}$$

Hence $a \mapsto \mathbf{P}_V(s, a)$ is differentiable and

$$\nabla_a \mathbf{P}_V(s, a) = \int_{\mathcal{S}} V(s') \nabla_a p(s' | s, a) \lambda(ds').$$

The same assumption gives the Lipschitz estimate

$$\|\nabla_a \mathbf{P}_V(s, a) - \nabla_a \mathbf{P}_V(s, \bar{a})\| \leq \|V\|_\infty L_p \|a - \bar{a}\|,$$

and therefore continuity of the derivative in a . Applying this with $V = V^\pi$ proves the displayed formula for $\nabla_a Q^\pi$.

If $\|V^\pi\|_\infty \leq V_{\max}$, then

$$\begin{aligned} \|b^\pi(s, a) - b^\pi(s, \bar{a})\| & \leq \beta \|a - \bar{a}\| + L_r \|a - \bar{a}\| \\ & \quad + \gamma \|V^\pi\|_\infty \int_{\mathcal{S}} \|\nabla_a p(s' | s, a) - \nabla_a p(s' | s, \bar{a})\| \lambda(ds') \\ & \leq (\beta + L_r + \gamma L_p V_{\max}) \|a - \bar{a}\|. \end{aligned}$$

Also

$$b^\pi(s, a) = -\beta a + h^\pi(s, a), \quad h^\pi(s, a) := \nabla_a r(s, a) + \gamma \int_{\mathcal{S}} V^\pi(s') \nabla_a p(s' | s, a) \lambda(ds'),$$

and the G_r, G_p bounds imply

$$\|h^\pi(s, a)\| \leq G_r + \gamma G_p V_{\max}.$$

We now turn to the soft optimality operator. For bounded V , set

$$\psi_V(s, a) := \frac{r(s, a) + \gamma \int_{\mathcal{S}} V(s') P(ds' | s, a)}{\tau}, \quad C_V := \frac{R_{\max} + \gamma \|V\|_{\infty}}{\tau}.$$

Then $|\psi_V(s, a)| \leq C_V$, and

$$\exp\left(\frac{r(s, a) - \frac{\beta}{2}\|a\|^2 + \gamma \int_{\mathcal{S}} V(s') P(ds' | s, a)}{\tau}\right) = Z_{\beta} e^{\psi_V(s, a)} \rho_{\beta}(a).$$

Thus the normalizer lies in

$$[Z_{\beta} e^{-C_V}, Z_{\beta} e^{C_V}],$$

so $\mathcal{T}^*V$ is finite and bounded. The Gibbs density can be written as

$$\mathcal{G}[V](a | s) = \frac{e^{\psi_V(s, a)}}{\int e^{\psi_V(s, \tilde{a})} \rho_{\beta}(\tilde{a}) d\tilde{a}} \rho_{\beta}(a).$$

Consequently,

$$\begin{aligned} \text{KL}(\mathcal{G}[V](\cdot | s) \| \rho_{\beta}) &= \int \psi_V(s, a) \mathcal{G}[V](a | s) da - \log \int e^{\psi_V(s, a)} \rho_{\beta}(a) da \\ &\leq 2C_V. \end{aligned}$$

Taking the supremum over s gives $\mathcal{G}[V] \in \Pi$. Finally, the Gibbs variational identity gives, for every admissible density μ ,

$$\int Q_V(s, a) \mu(a) da - \tau \int \mu(a) \log \mu(a) da = \tau \log \int e^{Q_V(s, a)/\tau} da - \tau \text{KL}(\mu \| \mathcal{G}[V](\cdot | s)),$$

so $\mathcal{G}[V]$ attains the one-step supremum.

For the WPG update, condition on $A_k^s = a$. The conditional distribution of $A_k^s + \eta b_k(s, A_k^s) + \sqrt{2\tau\eta} \xi_{k+1}$ is the Gaussian density $\varphi_{2\tau\eta}(y - a - \eta b_k(s, a))$. Integrating this conditional density against $\pi_k(a | s) da$ gives the displayed formula for $\pi_{k+1}(y | s)$, and Fubini's theorem gives

$$\int_{\mathbb{R}^d} \pi_{k+1}(y | s) dy = \int_{\mathbb{R}^d} \left[\int_{\mathbb{R}^d} \varphi_{2\tau\eta}(y - a - \eta b_k(s, a)) dy \right] \pi_k(a | s) da = 1.$$

The fixed-drift interpolation density is obtained by the same conditioning argument with η replaced by t . Since the Gaussian kernel is strictly positive, $q_t^s(y) > 0$ for every $t > 0$ and $y \in \mathbb{R}^d$. The numerator in the displayed formula for u_t^s is finite because $b_k(s, \cdot)$ has at most linear growth by the Lipschitz estimate above and $\pi_k(\cdot | s)$ has a finite second moment. Bayes' rule then gives

$$u_t^s(y) = \mathbb{E}[b_k(s, A_0^s) | A_t^s = y]$$

for $q_t^s(y) dy$ -a.e. y . This is the conditional-drift version used in the fixed-drift Fokker–Planck and entropy-differentiation arguments. $\blacksquare$

C.2.1 OPTIMAL GIBBS POLICY

Lemma C.3 (Optimal Gibbs policy) *Let $V^\star$ be the unique bounded fixed point of $\mathcal{T}^\star$, and define*

$$Q^\star(s, a) := \tilde{r}(s, a) + \gamma \int_S V^\star(s') P(ds' | s, a),$$

and

$$\pi^\star(a | s) := \frac{\exp(Q^\star(s, a)/\tau)}{\int_{\mathbb{R}^d} \exp(Q^\star(s, \tilde{a})/\tau) d\tilde{a}}. \quad (39)$$

Then $\pi^\star \in \Pi$,

$$\mathcal{T}^{\pi^\star} V^\star = \mathcal{T}^\star V^\star = V^\star,$$

and hence

$$V^{\pi^\star} = V^\star.$$

Moreover, for every admissible policy π ,

$$V^\pi \leq V^\star \quad \text{pointwise on } S.$$

Consequently, $\pi^\star$ is an optimal soft policy within the admissible class, and

$$J(\pi^\star) = \int_S V^\star(s) \rho_0(ds).$$

Proof [Proof of Lemma C.3] Since $V^\star$ is bounded and $|r| \leq R_{\max}$,

$$\left| r(s, a) + \gamma \int_S V^\star(s') P(ds' | s, a) \right| \leq R_{\max} + \gamma \|V^\star\|_\infty.$$

Write

$$\psi_s^\star(a) := \frac{r(s, a) + \gamma \int_S V^\star(s') P(ds' | s, a)}{\tau}.$$

Then

$$Q^\star(s, a) = -\frac{\beta}{2} \|a\|^2 + \tau \psi_s^\star(a),$$

and therefore

$$\exp(Q^\star(s, a)/\tau) = \exp(\psi_s^\star(a)) \exp\left(-\frac{\beta}{2\tau} \|a\|^2\right).$$

Let

$$C_\star := \frac{R_{\max} + \gamma \|V^\star\|_\infty}{\tau}.$$

Then $\|\psi_s^\star\|_\infty \leq C_\star$ for every s . Hence the normalizer in (39) satisfies

$$e^{-C_\star} Z_\beta \leq \int_{\mathbb{R}^d} \exp(Q^\star(s, a)/\tau) da \leq e^{C_\star} Z_\beta,$$

so it is finite and strictly positive. Moreover,

$$\pi^\star(a | s) = \frac{\exp(\psi_s^\star(a))}{\int_{\mathbb{R}^d} \exp(\psi_s^\star(\tilde{a})) \rho_\beta(\tilde{a}) d\tilde{a}} \rho_\beta(a).$$

Thus $\pi^*(\cdot | s)$ is a bounded perturbation of ρ_β . In particular,

$$\text{KL}(\pi^*(\cdot | s) \| \rho_\beta) = \int_{\mathbb{R}^d} \psi_s^*(a) \pi^*(a | s) da - \log \int_{\mathbb{R}^d} \exp(\psi_s^*(a)) \rho_\beta(a) da \leq 2C_\star.$$

Taking the supremum over s gives

$$\sup_{s \in S} \text{KL}(\pi^*(\cdot | s) \| \rho_\beta) \leq 2C_\star < \infty.$$

Therefore $\pi^* \in \Pi$. The required kernel measurability is part of the convention in Section 2.

We now show that π^* attains the soft optimality operator at V^* . For each fixed state s , let

$$Z_s^* := \int_{\mathbb{R}^d} \exp(Q^*(s, a)/\tau) da.$$

By definition,

$$\log \pi^*(a | s) = \frac{Q^*(s, a)}{\tau} - \log Z_s^*.$$

Therefore

$$Q^*(s, a) - \tau \log \pi^*(a | s) = \tau \log Z_s^*$$

for every a . Hence

$$\begin{aligned} (\mathcal{T}^{\pi^*} V^*)(s) &= \int_{\mathbb{R}^d} (Q^*(s, a) - \tau \log \pi^*(a | s)) \pi^*(a | s) da \\ &= \tau \log Z_s^*. \end{aligned}$$

On the other hand, the Gibbs variational formula gives

$$(\mathcal{T}^* V^*)(s) = \tau \log Z_s^*.$$

Since V^* is the fixed point of $\mathcal{T}^*$,

$$\mathcal{T}^{\pi^*} V^* = \mathcal{T}^* V^* = V^*.$$

Because $\pi^* \in \Pi$, the policy-evaluation operator $\mathcal{T}^{\pi^*}$ is a γ -contraction on bounded measurable functions and has a unique bounded fixed point, namely V^{π^*} . Since V^* is also a bounded fixed point of $\mathcal{T}^{\pi^*}$, uniqueness implies

$$V^{\pi^*} = V^*.$$

Finally, for any $\pi \in \Pi$ and any bounded measurable V ,

$$\mathcal{T}^\pi V \leq \mathcal{T}^* V$$

pointwise, because $\mathcal{T}^*$ is the supremum over admissible action densities in the one-step soft Bellman expression. Applying this with $V = V^*$ gives

$$\mathcal{T}^\pi V^* \leq \mathcal{T}^* V^* = V^*.$$

By monotonicity of $\mathcal{T}^\pi$,

$$(\mathcal{T}^\pi)^n V^\star \leq V^\star \quad \forall n \geq 1.$$

Since $\mathcal{T}^\pi$ is a γ -contraction, $(\mathcal{T}^\pi)^n V^\star$ converges uniformly to its unique bounded fixed point V^π . Passing to the limit yields

$$V^\pi \leq V^\star.$$

Thus $\pi^\star$ is optimal among admissible policies. Since $V^{\pi^\star} = V^\star$,

$$J(\pi^\star) = \int_S V^{\pi^\star}(s) \rho_0(ds) = \int_S V^\star(s) \rho_0(ds).$$

This completes the proof. ■

C.3 Value bounds and Bellman drift estimates

For every $\pi \in \Pi$, the one-step regularized reward obeys

$$\begin{aligned} \bar{r}^\pi(s) &:= \int_{\mathbb{R}^d} (\tilde{r}(s, a) - \tau \log \pi(a | s)) \pi(a | s) da \\ &= \int r(s, a) \pi(da | s) + \tau \log Z_\beta - \tau \text{KL}(\pi(\cdot | s) \| \rho_\beta). \end{aligned} \tag{40}$$

Consequently, under Assumption 4.1,

$$-R_{\max} + \tau \log Z_\beta - \tau C_\pi \leq \bar{r}^\pi(s) \leq R_{\max} + \tau \log Z_\beta, \quad C_\pi := \sup_s \text{KL}(\pi(\cdot | s) \| \rho_\beta).$$

This coercive identity is the basic reason admissibility relative to ρ_β makes Bellman evaluation finite on the unbounded action space.

Lemma C.4 (Basic value bounds) *Let*

$$U := \frac{R_{\max} + \tau \log Z_\beta}{1 - \gamma}, \quad L_\star := \frac{-R_{\max} + \tau \log Z_\beta}{1 - \gamma}, \quad \bar{E}_0 := \frac{2R_{\max} + \tau K_0}{1 - \gamma}.$$

Under Assumption 4.1, every admissible policy $\pi \in \Pi$ satisfies

$$V^\pi(s) \leq U \quad \forall s \in \mathcal{S}.$$

The optimal value satisfies

$$L_\star \leq V^\star(s) \leq U \quad \forall s \in \mathcal{S},$$

and the initial policy satisfies

$$V^{\pi_0}(s) \geq \frac{-R_{\max} + \tau \log Z_\beta - \tau K_0}{1 - \gamma}, \quad \|V^\star - V^{\pi_0}\|_\infty \leq \bar{E}_0.$$

Proof [Proof of Lemma C.4] For $\pi \in \Pi$, write $C_\pi = \sup_s \text{KL}(\pi(\cdot | s) \| \rho_\beta)$. The identity (40) gives, for every state s ,

$$-R_{\max} + \tau \log Z_\beta - \tau C_\pi \leq \bar{r}^\pi(s) \leq R_{\max} + \tau \log Z_\beta.$$

Thus $\mathcal{T}^\pi$ is an affine discounted Bellman operator with uniformly bounded one-step reward. For the constant function $U = (R_{\max} + \tau \log Z_\beta)/(1 - \gamma)$,

$$\mathcal{T}^\pi U \leq R_{\max} + \tau \log Z_\beta + \gamma U = U.$$

By monotonicity of $\mathcal{T}^\pi$ and contraction of policy evaluation, $(\mathcal{T}^\pi)^n U \rightarrow V^\pi$ uniformly and hence $V^\pi \leq U$.

For the lower bound on $V^\star$, use the stationary Gaussian policy $\pi^\beta(da | s) = \rho_\beta(a)da$. For this policy,

$$\int_{\mathbb{R}^d} (\tilde{r}(s, a) - \tau \log \rho_\beta(a)) \rho_\beta(a) da = \int r(s, a) \rho_\beta(a) da + \tau \log Z_\beta \geq -R_{\max} + \tau \log Z_\beta.$$

Therefore $V^{\pi^\beta} \geq L_\star$. Since $\mathcal{T}^\star V \geq \mathcal{T}^{\pi^\beta} V$ for every bounded V , monotone comparison of the discounted fixed points gives $V^\star \geq V^{\pi^\beta} \geq L_\star$. The upper bound $V^\star \leq U$ follows similarly from the log-partition form of $\mathcal{T}^\star$: for the constant function U ,

$$(\mathcal{T}^\star U)(s) \leq \tau \log \left(e^{(R_{\max} + \gamma U)/\tau} \int e^{-\beta \|a\|^2/(2\tau)} da \right) = R_{\max} + \gamma U + \tau \log Z_\beta = U,$$

and iterating $\mathcal{T}^\star$ from U yields $V^\star \leq U$.

Finally, Assumption 4.1(ii) and (40) imply

$$\bar{r}^{\pi_0}(s) \geq -R_{\max} + \tau \log Z_\beta - \tau K_0.$$

Applying the same monotone-comparison argument to $\mathcal{T}^{\pi_0}$ gives

$$V^{\pi_0}(s) \geq \frac{-R_{\max} + \tau \log Z_\beta - \tau K_0}{1 - \gamma}.$$

Since Lemma C.3 gives $V^{\pi_0} \leq V^\star$, the preceding initial lower bound and $V^\star \leq U$ yield

$$\|V^\star - V^{\pi_0}\|_\infty \leq \frac{R_{\max} + \tau \log Z_\beta}{1 - \gamma} - \frac{-R_{\max} + \tau \log Z_\beta - \tau K_0}{1 - \gamma} = \bar{E}_0.$$

■

Proof [Proof of Lemma 4.4] Lemma C.2 proves the Bellman action-gradient formula

$$b_k(s, a) = \nabla_a r(s, a) - \beta a + \gamma \int_{\mathcal{S}} V^{\pi_k}(s') \nabla_a p(s' | s, a) \lambda(ds').$$

Set

$$h_k(s, a) := \nabla_a r(s, a) + \gamma \int_{\mathcal{S}} V^{\pi_k}(s') \nabla_a p(s' | s, a) \lambda(ds').$$

Then $b_k(s, a) = -\beta a + h_k(s, a)$. If $\|V^{\pi_k}\|_\infty \leq V_{\max}$, Assumption 4.2 gives

$$\|h_k(s, a)\| \leq G_r + \gamma V_{\max} \int_{\mathcal{S}} \|\nabla_a p(s' \mid s, a)\| \lambda(ds') \leq G_r + \gamma G_p V_{\max}.$$

This proves the dissipative decomposition and (19). For the Lipschitz bound, subtract the displayed gradient formula at a and $\bar{a}$ and use the reward-gradient Lipschitz bound and the $L^1(\lambda)$ Lipschitz bound on $\nabla_a p$:

$$\begin{aligned} \|b_k(s, a) - b_k(s, \bar{a})\| &\leq \beta \|a - \bar{a}\| + L_r \|a - \bar{a}\| \\ &\quad + \gamma \|V^{\pi_k}\|_\infty \int_{\mathcal{S}} \|\nabla_a p(s' \mid s, a) - \nabla_a p(s' \mid s, \bar{a})\| \lambda(ds') \\ &\leq (\beta + L_r + \gamma L_p V_{\max}) \|a - \bar{a}\|. \end{aligned}$$

■

C.4 Moment bounds, step sizes, and uniform stability

Lemma C.5 (Second-moment bound under dissipativity) *Suppose that for all $0 \leq k \leq K-1$, Lemma 4.4 holds with $G(V_{\max}) = G$. Assume $M_0 := \sup_{s \in \mathcal{S}} \mathbb{E}\|A_0^s\|^2 < \infty$ and $0 < \eta \leq \frac{1}{4\beta}$. Then*

$$\sup_{0 \leq k \leq K} \sup_{s \in \mathcal{S}} \mathbb{E}\|A_k^s\|^2 \leq \max\{M_0, M_\infty(\eta)\}, \quad (41)$$

where one may take

$$M_\infty(\eta) := \frac{2}{\beta} \left(\frac{G^2}{\beta} + 2\tau d \right) + \frac{4G^2}{\beta} \eta.$$

Proof [Proof of Lemma C.5] Fix $s \in \mathcal{S}$ and write $A_k := A_k^s$ for brevity. Let

$$m_k := \mathbb{E}\|A_k\|^2, \quad 0 \leq k \leq K.$$

Recall the update

$$A_{k+1} = A_k + \eta b_k(s, A_k) + \sqrt{2\tau\eta} \xi_{k+1}, \quad \xi_{k+1} \sim \mathcal{N}(0, I_d),$$

with ξ_{k+1} independent of A_k . Taking squared norms and expectations gives

$$\begin{aligned} m_{k+1} &= \mathbb{E} \left\| A_k + \eta b_k(s, A_k) + \sqrt{2\tau\eta} \xi_{k+1} \right\|^2 \\ &= \mathbb{E} \left\| A_k + \eta b_k(s, A_k) \right\|^2 + 2\tau\eta \mathbb{E}\|\xi_{k+1}\|^2 \end{aligned}$$

because the cross term vanishes by independence and $\mathbb{E}[\xi_{k+1}] = 0$. Since $\mathbb{E}\|\xi_{k+1}\|^2 = d$, we have

$$m_{k+1} = \mathbb{E} \left\| A_k + \eta b_k(s, A_k) \right\|^2 + 2\tau\eta d. \quad (42)$$

Expanding the first term yields

$$\mathbb{E} \left\| A_k + \eta b_k(s, A_k) \right\|^2 = m_k + 2\eta \mathbb{E}\langle A_k, b_k(s, A_k) \rangle + \eta^2 \mathbb{E}\|b_k(s, A_k)\|^2. \quad (43)$$

By the assumed dissipativity bound (Lemma 4.4 with $G(V_{\max}) = G$), for each $0 \leq k \leq K - 1$ we may write

$$b_k(s, a) = -\beta a + h_k(s, a), \quad \sup_{s, a} \|h_k(s, a)\| \leq G.$$

Hence, for any $a \in \mathbb{R}^d$,

$$\langle a, b_k(s, a) \rangle = -\beta \|a\|^2 + \langle a, h_k(s, a) \rangle \leq -\beta \|a\|^2 + G \|a\| \leq -\frac{\beta}{2} \|a\|^2 + \frac{G^2}{2\beta},$$

where the last step uses Young's inequality $G \|a\| \leq \frac{\beta}{2} \|a\|^2 + \frac{G^2}{2\beta}$. Thus

$$\mathbb{E} \langle A_k, b_k(s, A_k) \rangle \leq -\frac{\beta}{2} m_k + \frac{G^2}{2\beta}. \quad (44)$$

Moreover, using $(x + y)^2 \leq 2x^2 + 2y^2$ and $\|h_k\| \leq G$,

$$\|b_k(s, a)\|^2 = \|- \beta a + h_k(s, a)\|^2 \leq 2\beta^2 \|a\|^2 + 2G^2,$$

so

$$\mathbb{E} \|b_k(s, A_k)\|^2 \leq 2\beta^2 m_k + 2G^2. \quad (45)$$

Plugging (44) and (45) into (42)-(43) gives, for $0 \leq k \leq K - 1$,

$$\begin{aligned} m_{k+1} &\leq m_k + 2\eta \left(-\frac{\beta}{2} m_k + \frac{G^2}{2\beta} \right) + \eta^2 (2\beta^2 m_k + 2G^2) + 2\tau\eta d \\ &= (1 - \beta\eta + 2\beta^2\eta^2) m_k + \eta \left(\frac{G^2}{\beta} + 2\tau d \right) + 2G^2\eta^2. \end{aligned}$$

Under the step-size condition $\eta \leq \frac{1}{4\beta}$,

$$2\beta^2\eta^2 \leq \frac{\beta\eta}{2} \quad \implies \quad 1 - \beta\eta + 2\beta^2\eta^2 \leq 1 - \frac{\beta\eta}{2}.$$

Define $\rho := 1 - \frac{\beta\eta}{2} \in (0, 1)$ and $\zeta_\eta := \eta \left(\frac{G^2}{\beta} + 2\tau d \right) + 2G^2\eta^2$.

Then we have the affine recursion

$$m_{k+1} \leq \rho m_k + \zeta_\eta, \quad 0 \leq k \leq K - 1. \quad (46)$$

Iterating (46) yields

$$\begin{aligned} m_k &\leq \rho^k m_0 + \zeta_\eta \sum_{j=0}^{k-1} \rho^j = \rho^k m_0 + \zeta_\eta \frac{1 - \rho^k}{1 - \rho} = \rho^k m_0 + (1 - \rho^k) \frac{\zeta_\eta}{1 - \rho}. \\ \frac{\zeta_\eta}{1 - \rho} &= \frac{\eta \left(\frac{G^2}{\beta} + 2\tau d \right) + 2G^2\eta^2}{\beta\eta/2} = \frac{2}{\beta} \left(\frac{G^2}{\beta} + 2\tau d \right) + \frac{4G^2}{\beta} \eta =: M_\infty(\eta). \end{aligned}$$

Therefore

$$m_k \leq \rho^k m_0 + (1 - \rho^k) M_\infty(\eta) \leq \max\{m_0, M_\infty(\eta)\} \quad \forall 0 \leq k \leq K.$$

Finally, taking $\sup_{0 \leq k \leq K}$ and then $\sup_{s \in \mathcal{S}}$, and using $m_0 = \mathbb{E}\|A_0^s\|^2 \leq M_0$, we obtain

$$\sup_{0 \leq k \leq K} \sup_{s \in \mathcal{S}} \mathbb{E}\|A_k^s\|^2 \leq \max\{M_0, M_\infty(\eta)\},$$

which is (41). ■

Lemma C.6 (Feasibility of the stability step-size condition) *Let the constants in (33)-(37) be defined as above. Then every $0 < \eta \leq \eta_0$ satisfies the exact stability condition (35). Moreover,*

$$\bar{\delta}_\eta \leq C_\delta \eta^2.$$

Proof [Proof of Lemma C.6] All constants in (37) are finite and positive under the standing assumptions, so $\eta_0 > 0$. Fix $0 < \eta \leq \eta_0$. The first two inequalities in (35) follow from

$$\eta \leq \frac{1}{4\beta}, \quad \eta \leq \frac{1}{\bar{\alpha}\tau}.$$

Since $\eta \leq 1/(4\beta)$,

$$\bar{M}_\infty(\eta) = \frac{2}{\beta} \left(\frac{\bar{G}^2}{\beta} + 2\tau d \right) + \frac{4\bar{G}^2}{\beta} \eta \leq \frac{3\bar{G}^2}{\beta^2} + \frac{4\tau d}{\beta}.$$

Therefore $\max\{M_0, \bar{M}_\infty(\eta)\} \leq \bar{M}$ and $B \leq \bar{B}$. Substituting this into (34) and using $\eta \leq 1$ gives

$$\bar{\delta}_\eta = \frac{\bar{L}_b^2 d}{2} \eta^2 + \frac{\bar{L}_b^2 B^2}{6\tau} \eta^3 \leq \left(\frac{\bar{L}_b^2 d}{2} + \frac{\bar{L}_b^2 \bar{B}^2}{6\tau} \right) \eta^2 = C_\delta \eta^2.$$

Finally, since $\bar{\alpha}\tau\eta \leq 1$ and $1 - e^{-x} \geq x/2$ for $x \in [0, 1]$,

$$c_\eta = 1 - e^{-\bar{\alpha}\tau\eta} \geq \frac{\bar{\alpha}\tau\eta}{2}.$$

Thus

$$\frac{\tau}{(1-\gamma)^2} \frac{\bar{\delta}_\eta}{c_\eta} \leq \frac{\tau}{(1-\gamma)^2} \frac{C_\delta \eta^2}{\bar{\alpha}\tau\eta/2} = \frac{2C_\delta}{\bar{\alpha}(1-\gamma)^2} \eta \leq 1,$$

where the last inequality uses the definition of η_0 . Hence (35) holds. ■

Proof [Proof of Proposition 4.1] By Lemma C.6, every $0 < \eta \leq \eta_0$ satisfies the exact stability condition (35). The proof below uses this exact condition. For any index k for which π_k is admissible, write

$$b_k(s, a) := \nabla_a Q^{\pi_k}(s, a).$$

For such k and each $s \in \mathcal{S}$, let

$$p_s^{\pi_k}(a) = \frac{\exp(Q^{\pi_k}(s, a)/\tau)}{\int_{\mathbb{R}^d} \exp(Q^{\pi_k}(s, \tilde{a})/\tau) d\tilde{a}}.$$

Define

$$W_k(s) := V^*(s) - V^{\pi_k}(s), \quad R_k(s) := (\mathcal{T}^* V^{\pi_k})(s) - V^{\pi_k}(s), \quad E_k := \|W_k\|_\infty.$$

By Lemma 2.1,

$$R_k(s) = \tau \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}).$$

Moreover, by Lemma C.1, $\mathcal{T}^*$ is monotone and a γ -contraction; since $V^* \geq V^{\pi_k}$ pointwise,

$$0 \leq V^*(s) - (\mathcal{T}^* V^{\pi_k})(s) = (\mathcal{T}^* V^*)(s) - (\mathcal{T}^* V^{\pi_k})(s) \leq \gamma E_k, \quad \forall s \in \mathcal{S}. \quad (47)$$

For each $m \geq 0$, let $\mathcal{I}_m$ denote the assertion that, for every $0 \leq j \leq m$,

$$\pi_j \in \Pi, \quad \|V^{\pi_j}\|_\infty \leq \bar{V}.$$

We prove $\mathcal{I}_m$ for all m by induction. The admissibility part ensures that Lemma C.2 applies recursively to the Bellman objects and to the WPG push-forward density; measurability is handled throughout by the convention in Section 2.

Base case. The admissibility of π_0 follows from Assumption 4.1(ii), while kernel measurability is covered by the convention in Section 2. By Lemma C.4, $\sup_s V^{\pi_0}(s) \leq U \leq \bar{V}$. By Lemma C.4,

$$V^{\pi_0}(s) \geq V^*(s) - E_0 \geq L_* - E_0.$$

By Lemma C.4, $E_0 \leq \bar{E}_0$. The definition (33) implies $\bar{V} \geq \bar{E}_0 - L_* + 1$, and hence $L_* - E_0 \geq L_* - \bar{E}_0 \geq -\bar{V}$. Thus $\|V^{\pi_0}\|_\infty \leq \bar{V}$, so $\mathcal{I}_0$ holds.

Induction step. Assume $\mathcal{I}_k$ holds. Then Lemma C.2 gives the Bellman objects $V^{\pi_j}, Q^{\pi_j}, b_j$ for $0 \leq j \leq k$ and the explicit WPG push-forward density for π_{k+1} . Lemma 4.4 implies that, for all $0 \leq j \leq k$,

$$\|b_j(s, a) - b_j(s, \bar{a})\| \leq \bar{L}_b \|a - \bar{a}\|, \quad b_j(s, a) = -\beta a + h_j(s, a), \quad \sup_{s, a} \|h_j(s, a)\| \leq \bar{G}.$$

Applying Lemma C.5 on the finite horizon $\{0, 1, \dots, k+1\}$ with $G = \bar{G}$ yields

$$\sup_{0 \leq j \leq k+1} \sup_{s \in \mathcal{S}} \mathbb{E} \|A_j^s\|^2 \leq \max\{M_0, \bar{M}_\infty(\eta)\} \leq \bar{M}. \quad (48)$$

Consequently, for every $0 \leq j \leq k$,

$$\sup_{s \in \mathcal{S}} \mathbb{E} \|b_j(s, A_j^s)\|^2 \leq 2\beta^2 \max\{M_0, \bar{M}_\infty(\eta)\} + 2\bar{G}^2 \leq \bar{B}^2.$$

At the current step k , Lemma 4.3 gives that $p_s^{\pi_k}$ satisfies an LSI with constant at least $\bar{\alpha}$. The Gibbs definition and the Bellman-gradient formula in Lemma C.2 also give $p_s^{\pi_k} > 0$, $\log p_s^{\pi_k} \in$

$C^1(\mathbb{R}^d)$, and the score identity $\tau \nabla_a \log p_s^{\pi_k} = b_k(s, \cdot)$. We next verify the finite-second-moment and finite-KL hypotheses in Lemma 4.5 quantitatively, and the admissibility/finite-KL hypotheses needed for Lemma 4.6.

For $j = 0$, Assumption 4.1(ii) gives

$$\sup_s \text{KL}(\pi_0(\cdot | s) \| \rho_\beta) \leq K_0 \leq \bar{K}_\eta.$$

For $1 \leq j \leq k+1$, the update writes

$$A_j^s = X_{j-1}^s + \sqrt{2\tau\eta}\xi_j, \quad X_{j-1}^s := A_{j-1}^s + \eta b_{j-1}(s, A_{j-1}^s),$$

with $\xi_j \sim \mathcal{N}(0, I_d)$ independent of X_{j-1}^s . Since (48) gives $\mathbb{E}\|A_j^s\|^2 \leq M_\infty$, Lemma E.3 with $\sigma^2 = 2\tau\eta$ yields, for $1 \leq j \leq k+1$,

$$\text{KL}(\pi_j(\cdot | s) \| \rho_\beta) \leq \frac{\beta M_\infty}{2\tau} + \log Z_\beta - \frac{d}{2} \log(4\pi e\tau\eta) \leq \bar{K}_\eta. \quad (49)$$

Combining the initialization bound for $j = 0$ with the smoothing bound for $j \geq 1$ gives the uniform finite-horizon estimate

$$\sup_{0 \leq j \leq k+1} \sup_{s \in \mathcal{S}} \text{KL}(\pi_j(\cdot | s) \| \rho_\beta) \leq \bar{K}_\eta. \quad (50)$$

Thus the Gaussian-KL bound holds for all policies generated up to time $k+1$, and in particular $\pi_j \in \Pi$ for all $j \leq k+1$. The remaining kernel measurability is covered by the convention in Section 2, so the admissibility part of the induction is closed.

Under the value bound, $p_s^{\pi_k}$ is a bounded perturbation of ρ_β :

$$p_s^{\pi_k}(a) = \frac{e^{\psi_{k,s}(a)}}{\int e^{\psi_{k,s}} d\rho_\beta} \rho_\beta(a), \quad \psi_{k,s}(a) := \frac{r(s, a) + \gamma \int V^{\pi_k}(s') P(ds' | s, a)}{\tau}.$$

Since $|r| \leq R_{\max}$ and $\|V^{\pi_k}\|_\infty \leq \bar{V}$, $\|\psi_{k,s}\|_\infty \leq (R_{\max} + \gamma\bar{V})/\tau$. The bounded-perturbation part of Lemma E.3, applied with the $j = k$ case of (50), gives

$$\text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) \leq \bar{K}_\eta + \frac{2(R_{\max} + \gamma\bar{V})}{\tau} = \bar{H}_\eta < \infty. \quad (51)$$

The finite-second-moment hypothesis of Lemma 4.5 follows from (48); the drift second-moment hypothesis follows from

$$\sup_{s \in \mathcal{S}} \mathbb{E}\|b_k(s, A_k^s)\|^2 \leq 2\beta^2 \max\{M_0, \bar{M}_\infty(\eta)\} + 2\bar{G}^2 = \bar{B}^2.$$

Applying Lemma 4.5 with $\alpha = \bar{\alpha}$, $L_b = \bar{L}_b$, and $B^2 = \bar{B}^2$ gives, for every s ,

$$\text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) \leq e^{-\bar{\alpha}\tau\eta} \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) + \bar{\delta}_\eta. \quad (52)$$

The right-hand side of (52) is finite by (51), and hence $\text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) < \infty$. Thus the admissibility and finite-KL hypotheses of Lemma 4.2 are verified for (π_k, π_{k+1}) . Applying Lemma 4.6 to (52) with $c_\eta := 1 - e^{-\bar{\alpha}\tau\eta}$ and $\delta = \bar{\delta}_\eta$ gives

$$g_k(s) \geq c_\eta R_k(s) - \tau \bar{\delta}_\eta.$$

In particular, since $R_k \geq 0$, $g_k(s) \geq -\tau\bar{\delta}_\eta$ and $\|g_k^-\|_\infty \leq \tau\bar{\delta}_\eta$.

The one-step resolvent identity gives

$$V^{\pi_{k+1}} - V^{\pi_k} = (I - \gamma P^{\pi_{k+1}})^{-1} g_k = \sum_{n \geq 0} \gamma^n (P^{\pi_{k+1}})^n g_k.$$

For $n \geq 1$, $(P^{\pi_{k+1}})^n g_k(s) \geq -\|g_k^-\|_\infty$, so

$$\begin{aligned} (V^{\pi_{k+1}} - V^{\pi_k})(s) &\geq g_k(s) - \sum_{n \geq 1} \gamma^n \|g_k^-\|_\infty \\ &\geq c_\eta R_k(s) - \tau\bar{\delta}_\eta - \frac{\gamma}{1-\gamma} \tau\bar{\delta}_\eta = c_\eta R_k(s) - \frac{\tau}{1-\gamma} \bar{\delta}_\eta. \end{aligned}$$

Equivalently,

$$W_{k+1}(s) \leq W_k(s) - c_\eta R_k(s) + \frac{\tau}{1-\gamma} \bar{\delta}_\eta. \quad (53)$$

Using (47),

$$R_k(s) = W_k(s) - (V^*(s) - (\mathcal{T}^* V^{\pi_k})(s)) \geq W_k(s) - \gamma E_k.$$

Combining this bound with (53) yields

$$W_{k+1}(s) \leq (1 - c_\eta) W_k(s) + c_\eta \gamma E_k + \frac{\tau}{1-\gamma} \bar{\delta}_\eta \leq (1 - (1 - \gamma) c_\eta) E_k + \frac{\tau}{1-\gamma} \bar{\delta}_\eta.$$

Taking the supremum over s gives

$$E_{k+1} \leq \kappa_\eta E_k + \frac{\tau}{1-\gamma} \bar{\delta}_\eta, \quad \kappa_\eta := 1 - (1 - \gamma) c_\eta. \quad (54)$$

Iterating (54) up to $k + 1$ gives

$$E_{k+1} \leq \kappa_\eta^{k+1} E_0 + \frac{\tau}{1-\gamma} \bar{\delta}_\eta \sum_{j=0}^k \kappa_\eta^j \leq E_0 + \frac{\tau}{(1-\gamma)^2 c_\eta} \bar{\delta}_\eta \leq \bar{E}_0 + 1,$$

where the last inequality is the stability step-size condition.

Finally, Lemma C.4 gives $V^{\pi_{k+1}}(s) \leq U \leq \bar{V}$, while Lemma C.4 gives

$$V^{\pi_{k+1}}(s) \geq V^*(s) - E_{k+1} \geq L_* - \bar{E}_0 - 1 \geq -\bar{V},$$

again by (33). Therefore $\|V^{\pi_{k+1}}\|_\infty \leq \bar{V}$. Together with the admissibility conclusion above, this proves $\mathcal{I}_{k+1}$ and closes the induction.

It remains to record the uniform consequences. The value bound just proved implies the drift Lipschitz bound and dissipative decomposition for every k via Lemma 4.4. Applying Lemma C.5 on an arbitrary finite horizon and then letting that horizon vary gives $\sup_s \mathbb{E} \|A_k^s\|^2 \leq \max\{M_0, \bar{M}_\infty(\eta)\}$ for every k . The drift second-moment bound follows from $\|b_k(s, a)\|^2 \leq 2\beta^2 \|a\|^2 + 2\bar{G}^2$. The quantitative Gaussian-KL bound (24) follows by applying (50) on arbitrary finite horizons; the $k = 0$ case is supplied by Assumption 4.1(ii), while every $k \geq 1$ case is supplied by the Gaussian-smoothing estimate (49). The kernel measurability assertion follows from the convention in Section 2. Once the value bound is

closed for all iterations, the bounded-perturbation argument above applies to every $p_s^{\pi_k}$ and gives (25). Finally, Lemma 4.3 with $V_{\max} = \bar{V}$ gives the uniform LSI constant $\bar{\alpha}$ for every $p_s^{\pi_k}$. $\blacksquare$

Proof [Proof of Lemma 4.3] Fix $s \in \mathcal{S}$. By the definition of $p_s^{\pi_k}$, write

$$p_s^{\pi_k}(a) \propto \exp\left(\frac{r(s, a) + \gamma \int_{\mathcal{S}} V^{\pi_k}(s') P(ds' | s, a)}{\tau}\right) \exp\left(-\frac{\beta}{2\tau} \|a\|^2\right).$$

Thus $p_s^{\pi_k}$ is a bounded perturbation of the Gaussian $\rho_\beta(a) \propto \exp(-\beta \|a\|^2 / (2\tau))$. If $\|V^{\pi_k}\|_\infty \leq V_{\max}$, then the perturbation

$$\psi_{k,s}(a) := \frac{r(s, a) + \gamma \int_{\mathcal{S}} V^{\pi_k}(s') P(ds' | s, a)}{\tau}$$

satisfies

$$|\psi_{k,s}(a)| \leq \frac{R_{\max} + \gamma V_{\max}}{\tau}, \quad \text{osc}(\psi_{k,s}) \leq \frac{2(R_{\max} + \gamma V_{\max})}{\tau}.$$

Since ρ_β satisfies an LSI with constant β/τ , the Holley-Stroock bounded perturbation lemma implies that $p_s^{\pi_k}$ satisfies an LSI with constant at least

$$\frac{\beta}{\tau} \exp\left(-\frac{2(R_{\max} + \gamma V_{\max})}{\tau}\right) = \alpha(V_{\max}).$$

This proves the claim. $\blacksquare$

Appendix D. Core Bellman Contraction Proof

This appendix proves the Bellman part of the argument. These are the nonstandard identities that replace the flat-convex gap-conversion step in mean-field Langevin analysis.

D.1 Bellman residual, residual-to-gap, and resolvent identities

Proof [Proof of Lemma 4.1] Recall

$$W_k(s) = V^*(s) - V^{\pi_k}(s), \quad E_k = \|W_k\|_\infty, \quad R_k(s) = (\mathcal{T}^* V^{\pi_k})(s) - V^{\pi_k}(s).$$

Decompose W_k by adding and subtracting $(\mathcal{T}^* V^{\pi_k})(s)$:

$$W_k(s) = (V^*(s) - (\mathcal{T}^* V^{\pi_k})(s)) + ((\mathcal{T}^* V^{\pi_k})(s) - V^{\pi_k}(s)) = (V^* - (\mathcal{T}^* V^{\pi_k}))(s) + R_k(s). \quad (55)$$

Since $V^* = \mathcal{T}^* V^*$ and $\mathcal{T}^*$ is a γ -contraction in $\|\cdot\|_\infty$ by Lemma C.1,

$$\|V^* - \mathcal{T}^* V^{\pi_k}\|_\infty = \|\mathcal{T}^* V^* - \mathcal{T}^* V^{\pi_k}\|_\infty \leq \gamma \|V^* - V^{\pi_k}\|_\infty = \gamma E_k.$$

In particular, for every $s \in \mathcal{S}$,

$$V^*(s) - (\mathcal{T}^* V^{\pi_k})(s) \leq \gamma E_k. \quad (56)$$

Using (55) and (56),

$$R_k(s) = W_k(s) - (V^\star - (\mathcal{T}^\star V^{\pi_k}))(s) \geq W_k(s) - \gamma E_k.$$

■

Proof [Proof of Lemma 4.2] Recall the policy evaluation operator $\mathcal{T}^\pi$ (2) and the induced state kernel $P^\pi(ds' | s)$. For a fixed policy π , define the regularized immediate reward

$$\bar{r}^\pi(s) := \int_{\mathcal{A}} (\tilde{r}(s, a) - \tau \log \pi(a | s)) \pi(a | s) da.$$

Then we can rewrite $\mathcal{T}^\pi$ as the affine map

$$(\mathcal{T}^\pi V)(s) = \bar{r}^\pi(s) + \gamma(P^\pi V)(s).$$

Since $V^{\pi_{k+1}}$ is the fixed point of $\mathcal{T}^{\pi_{k+1}}$, we have

$$V^{\pi_{k+1}} = \mathcal{T}^{\pi_{k+1}} V^{\pi_{k+1}}.$$

Subtract V^{π_k} from both sides and add and subtract $\mathcal{T}^{\pi_{k+1}} V^{\pi_k}$:

$$V^{\pi_{k+1}} - V^{\pi_k} = (\mathcal{T}^{\pi_{k+1}} V^{\pi_{k+1}} - \mathcal{T}^{\pi_{k+1}} V^{\pi_k}) + (\mathcal{T}^{\pi_{k+1}} V^{\pi_k} - V^{\pi_k}) = \gamma P^{\pi_{k+1}}(V^{\pi_{k+1}} - V^{\pi_k}) + g_k,$$

which rearranges to the claimed resolvent identity

$$(I - \gamma P^{\pi_{k+1}})(V^{\pi_{k+1}} - V^{\pi_k}) = g_k.$$

This proves (16).

Now compute $(\mathcal{T}^{\pi_{k+1}} V^{\pi_k})(s)$ using the definition of $\mathcal{T}^\pi$ and the definition of Q^{π_k} :

$$(\mathcal{T}^{\pi_{k+1}} V^{\pi_k})(s) = \int_{\mathcal{A}} (Q^{\pi_k}(s, a) - \tau \log \pi_{k+1}(a | s)) \pi_{k+1}(a | s) da$$

Similarly, since $V^{\pi_k} = \mathcal{T}^{\pi_k} V^{\pi_k}$,

$$V^{\pi_k}(s) = \int_{\mathcal{A}} (Q^{\pi_k}(s, a) - \tau \log \pi_k(a | s)) \pi_k(a | s) da.$$

Using the standard Gibbs identity

$$\int_{\mathcal{A}} f(a) \mu(a) da - \tau \int_{\mathcal{A}} \mu(a) \log \mu(a) da = \tau \log Z - \tau \text{KL}(\mu || p),$$

we have that

$$\begin{aligned} (\mathcal{T}^{\pi_{k+1}} V^{\pi_k})(s) &= \tau \log Z - \tau \text{KL}(\pi_{k+1}(\cdot | s) || p_s^{\pi_k}), \\ V^{\pi_k}(s) &= \tau \log Z - \tau \text{KL}(\pi_k(\cdot | s) || p_s^{\pi_k}). \end{aligned}$$

Subtracting gives

$$g_k(s) = (\mathcal{T}^{\pi_{k+1}} V^{\pi_k})(s) - V^{\pi_k}(s) = \tau \left(\text{KL}(\pi_k(\cdot | s) || p_s^{\pi_k}) - \text{KL}(\pi_{k+1}(\cdot | s) || p_s^{\pi_k}) \right),$$

which is exactly (17). ■

D.2 One-step KL contraction with Discretization Error

Lemma D.1 (Fokker–Planck with conditional drift) *Let $A_0 \sim \pi_k(\cdot \mid s)$ and let $(B_t)_{t \geq 0}$ be standard Brownian motion in $\mathbb{R}^d$, independent of A_0 . Let $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be Borel and assume $\mathbb{E}\|b(A_0)\| < \infty$. For $t \in [0, \eta]$ define*

$$A_t := A_0 + t b(A_0) + \sqrt{2\tau} B_t, \quad \mu_t := \text{Law}(A_t).$$

Define the conditional drift

$$u_t(a) := \mathbb{E}[b(A_0) \mid A_t = a].$$

Then, for $t > 0$, μ_t satisfies the Fokker–Planck equation

$$\partial_t \mu_t = -\nabla \cdot (u_t \mu_t) + \tau \Delta \mu_t$$

in the sense of distributions.

Proof [Proof of Lemma D.1] Condition on $A_0 = a_0$. Then

$$A_t = a_0 + t b(a_0) + \sqrt{2\tau} B_t,$$

so the conditional law of A_t has density

$$p_t(a \mid a_0) = \varphi_{2\tau t}(a - a_0 - t b(a_0)),$$

where $\varphi_{2\tau t}$ is the Gaussian density with covariance $2\tau t I_d$. Equivalently, $p_t(\cdot \mid a_0)$ solves the standard Fokker–Planck equation

$$\partial_t p_t(\cdot \mid a_0) = -\nabla \cdot (b(a_0) p_t(\cdot \mid a_0)) + \tau \Delta p_t(\cdot \mid a_0).$$

Now the marginal density of μ_t is the mixture

$$\mu_t(a) = \int p_t(a \mid a_0) \pi_k(da_0 \mid s).$$

Integrating the above Fokker–Planck equation over a_0 gives

$$\partial_t \mu_t(a) = -\nabla \cdot \left(\int b(a_0) p_t(a \mid a_0) \pi_k(da_0 \mid s) \right) + \tau \Delta \mu_t(a).$$

Define the flux

$$j_t(a) := \int b(a_0) p_t(a \mid a_0) \pi_k(da_0 \mid s).$$

By Bayes' rule, for μ_t -a.e. a ,

$$u_t(a) = \mathbb{E}[b(A_0) \mid A_t = a] = \frac{\int b(a_0) p_t(a \mid a_0) \pi_k(da_0 \mid s)}{\int p_t(a \mid a_0) \pi_k(da_0 \mid s)} = \frac{j_t(a)}{\mu_t(a)}.$$

Hence $j_t(a) = u_t(a) \mu_t(a)$, and substituting this into the previous display yields

$$\partial_t \mu_t = -\nabla \cdot (u_t \mu_t) + \tau \Delta \mu_t,$$

as claimed. ■

Proof [Proof of Lemma 4.5] Fix $k \geq 0$ and $s \in \mathcal{S}$. For brevity write

$$b(a) := b_k(s, a), \quad p(a) := p_s^{\pi_k}(a).$$

Let $A_0 \sim \pi_k(\cdot | s)$ and let $\{B_t\}_{t \geq 0}$ be standard Brownian motion in $\mathbb{R}^d$, independent of A_0 . For $t \in [0, \eta]$ define the one-step interpolation

$$A_t := A_0 + t b(A_0) + \sqrt{2\tau} B_t, \quad \mu_t := \text{Law}(A_t). \quad (57)$$

Then $\mu_0 = \pi_k(\cdot | s)$ and $\mu_\eta = \pi_{k+1}(\cdot | s)$. We denote the density of μ_t again by $\mu_t(a)$ and define

$$u_t(a) := \mathbb{E}[b(A_0) | A_t = a].$$

By Lemma D.1, μ_t solves

$$\partial_t \mu_t = -\nabla \cdot (u_t \mu_t) + \tau \Delta \mu_t.$$

Set

$$K(t) := \text{KL}(\mu_t \| p) = \int_{\mathbb{R}^d} \mu_t(a) \log \frac{\mu_t(a)}{p(a)} da.$$

Using the Fokker–Planck equation and differentiating under the integral sign,

$$\begin{aligned} K'(t) &= \frac{d}{dt} \int_{\mathbb{R}^d} \mu_t(a) \log \frac{\mu_t(a)}{p(a)} da \\ &= \int_{\mathbb{R}^d} \log \frac{\mu_t(a)}{p(a)} \partial_t \mu_t(a) da + \int_{\mathbb{R}^d} \partial_t \mu_t(a) da. \end{aligned}$$

Since $\int_{\mathbb{R}^d} \partial_t \mu_t(a) da = 0$, this becomes

$$K'(t) = \int_{\mathbb{R}^d} \log \frac{\mu_t(a)}{p(a)} [-\nabla \cdot (u_t(a) \mu_t(a)) + \tau \Delta \mu_t(a)] da.$$

Integrating by parts gives

$$\begin{aligned} K'(t) &= \int_{\mathbb{R}^d} \left\langle \nabla \log \frac{\mu_t}{p}(a), u_t(a) \right\rangle \mu_t(a) da - \tau \int_{\mathbb{R}^d} \left\langle \nabla \log \frac{\mu_t}{p}(a), \nabla \mu_t(a) \right\rangle da \\ &= \int_{\mathbb{R}^d} \left\langle \nabla \log \frac{\mu_t}{p}(a), u_t(a) \right\rangle \mu_t(a) da - \tau \int_{\mathbb{R}^d} \left\langle \nabla \log \frac{\mu_t}{p}(a), \nabla \log \mu_t(a) \right\rangle \mu_t(a) da. \end{aligned}$$

By the Gibbs-score identity,

$$\tau \nabla \log p = b.$$

Therefore

$$\nabla \log \mu_t = \nabla \log \frac{\mu_t}{p} + \nabla \log p = \nabla \log \frac{\mu_t}{p} + \frac{1}{\tau} b.$$

Substituting this identity into the preceding display yields

$$K'(t) = \mathbb{E}_{\mu_t} \left[\left\langle u_t(A) - b(A), \nabla \log \frac{\mu_t}{p}(A) \right\rangle \right] - \tau \mathcal{I}(\mu_t \| p). \quad (58)$$

Young's inequality gives

$$\mathbb{E}_{\mu_t} \left[\left\langle u_t(A) - b(A), \nabla \log \frac{\mu_t}{p}(A) \right\rangle \right] \leq \frac{1}{2\tau} \mathbb{E}_{\mu_t} \|u_t(A) - b(A)\|^2 + \frac{\tau}{2} \mathcal{I}(\mu_t \| p).$$

Combining this with (58),

$$K'(t) \leq -\frac{\tau}{2} \mathcal{I}(\mu_t \| p) + \frac{1}{2\tau} \mathbb{E}_{\mu_t} \|u_t(A) - b(A)\|^2.$$

By the LSI for p ,

$$\mathcal{I}(\mu_t \| p) \geq 2\alpha \text{KL}(\mu_t \| p) = 2\alpha K(t).$$

Hence

$$K'(t) \leq -\alpha\tau K(t) + \frac{1}{2\tau} \mathbb{E}_{\mu_t} \|u_t(A) - b(A)\|^2. \quad (59)$$

It remains to bound the interpolation error. By conditional Jensen's inequality,

$$\begin{aligned} \mathbb{E}_{\mu_t} \|u_t(A) - b(A)\|^2 &= \mathbb{E} \|\mathbb{E}[b(A_0) \mid A_t] - b(A_t)\|^2 \\ &= \mathbb{E} \|\mathbb{E}[b(A_0) - b(A_t) \mid A_t]\|^2 \\ &\leq \mathbb{E} [\mathbb{E} [\|b(A_0) - b(A_t)\|^2 \mid A_t]] \\ &= \mathbb{E} \|b(A_0) - b(A_t)\|^2. \end{aligned}$$

Using the L_b -Lipschitzness of b ,

$$\mathbb{E}_{\mu_t} \|u_t(A) - b(A)\|^2 \leq L_b^2 \mathbb{E} \|A_0 - A_t\|^2.$$

From (57),

$$A_t - A_0 = t b(A_0) + \sqrt{2\tau} B_t.$$

Since B_t is independent of A_0 and has mean zero,

$$\mathbb{E} \|A_t - A_0\|^2 = t^2 \mathbb{E} \|b(A_0)\|^2 + 2\tau d t \leq B^2 t^2 + 2\tau d t.$$

Thus

$$\mathbb{E}_{\mu_t} \|u_t(A) - b(A)\|^2 \leq L_b^2 B^2 t^2 + 2\tau L_b^2 d t.$$

Substituting this bound into (59), we get

$$K'(t) \leq -\alpha\tau K(t) + L_b^2 d t + \frac{L_b^2 B^2}{2\tau} t^2.$$

By Gronwall's inequality,

$$\begin{aligned} K(\eta) &\leq e^{-\alpha\tau\eta} K(0) + \int_0^\eta e^{-\alpha\tau(\eta-u)} \left(L_b^2 d u + \frac{L_b^2 B^2}{2\tau} u^2 \right) du \\ &\leq e^{-\alpha\tau\eta} K(0) + \frac{L_b^2 d}{2} \eta^2 + \frac{L_b^2 B^2}{6\tau} \eta^3. \end{aligned}$$

Recalling that

$$K(0) = \text{KL}(\pi_k(\cdot \mid s) \| p_s^{\pi_k}), \quad K(\eta) = \text{KL}(\pi_{k+1}(\cdot \mid s) \| p_s^{\pi_k}),$$

we obtain

$$\text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) \leq e^{-\alpha\tau\eta} \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) + \frac{L_b^2 d}{2} \eta^2 + \frac{L_b^2 B^2}{6\tau} \eta^3.$$

This proves (20). ■

D.3 From statewise KL contraction to Bellman improvement

Proof [Proof of Lemma 4.6] By Lemma 4.2,

$$g_k(s) = \tau \left(\text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) - \text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) \right).$$

Using (21),

$$\begin{aligned} g_k(s) &\geq \tau \left(\text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) - (1 - c) \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) - \delta \right) \\ &= c \tau \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) - \tau \delta = c R_k(s) - \tau \delta. \end{aligned}$$

This proves (22). The final sentence follows by substituting the KL contraction from Lemma 4.5. ■

D.4 Proof of the main theorem

Proof [Proof of Theorem 4.1] Define

$$W_k(s) := V^*(s) - V^{\pi_k}(s), \quad E_k := \|W_k\|_\infty, \quad R_k(s) := (\mathcal{T}^* V^{\pi_k})(s) - V^{\pi_k}(s).$$

By Proposition 4.1, the constants $\bar{L}_b$, $\bar{M}$, $\bar{B}^2$, $\bar{\alpha}$, $\bar{\delta}_\eta$, and c_η are valid uniformly over all iterates. The Gibbs targets are strictly positive bounded perturbations of ρ_β , the score identity (15) holds, and the finite-second-moment, drift-moment, admissibility, and $\text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) < \infty$ hypotheses needed for Lemmas 4.5 and 4.2 hold at every step. The one-step contraction below follows from the same argument used in the proof of Proposition 4.1; for completeness and readability, we spell out the derivation.

Lemma 4.5, applied with $(\alpha, L_b, B^2) = (\bar{\alpha}, \bar{L}_b, \bar{B}^2)$, gives

$$\text{KL}(\pi_{k+1}(\cdot | s) \| p_s^{\pi_k}) \leq e^{-\bar{\alpha}\tau\eta} \text{KL}(\pi_k(\cdot | s) \| p_s^{\pi_k}) + \bar{\delta}_\eta.$$

Applying Lemma 4.6 with $c_\eta = 1 - e^{-\bar{\alpha}\tau\eta}$ yields

$$g_k(s) \geq c_\eta R_k(s) - \tau \bar{\delta}_\eta.$$

Since $R_k \geq 0$, this also gives $g_k(s) \geq -\tau \bar{\delta}_\eta$. The one-step resolvent identity gives

$$V^{\pi_{k+1}} - V^{\pi_k} = (I - \gamma P^{\pi_{k+1}})^{-1} g_k = \sum_{n \geq 0} \gamma^n (P^{\pi_{k+1}})^n g_k,$$

and hence

$$(V^{\pi_{k+1}} - V^{\pi_k})(s) \geq c_\eta R_k(s) - \frac{\tau}{1-\gamma} \bar{\delta}_\eta.$$

Therefore

$$W_{k+1}(s) \leq W_k(s) - c_\eta R_k(s) + \frac{\tau}{1-\gamma} \bar{\delta}_\eta. \quad (60)$$

By Lemma 4.1, $R_k(s) \geq W_k(s) - \gamma E_k$. Substituting this into (60) and taking the supremum over s gives

$$E_{k+1} \leq \kappa_\eta E_k + \frac{\tau}{1-\gamma} \bar{\delta}_\eta, \quad \kappa_\eta := 1 - (1-\gamma)c_\eta.$$

Iterating,

$$E_k \leq \kappa_\eta^k E_0 + \frac{\tau}{(1-\gamma)^2} \frac{\bar{\delta}_\eta}{c_\eta}.$$

Since $0 < \eta \leq \eta_0$, we have $\bar{\alpha}\tau\eta \leq 1$, so $c_\eta = 1 - e^{-\bar{\alpha}\tau\eta} \geq \bar{\alpha}\tau\eta/2$, and $\kappa_\eta^k \leq \exp(-(1-\gamma)c_\eta k) \leq \exp(-\frac{1}{2}\bar{\alpha}\tau(1-\gamma)\eta k)$. Also, by (36) and (37), $\bar{\delta}_\eta \leq C_\delta \eta^2$. Thus

$$\|V^\star - V^{\pi_k}\|_\infty = E_k \leq \exp\left(-\frac{1}{2}\bar{\alpha}\tau(1-\gamma)\eta k\right) \|V^\star - V^{\pi_0}\|_\infty + \frac{2C_\delta}{\bar{\alpha}(1-\gamma)^2} \eta.$$

Finally, Lemma C.3 gives $V^{\pi_k} \leq V^\star$, and hence

$$J(\pi^\star) - J(\pi_k) = \int_{\mathcal{S}} (V^\star(s) - V^{\pi_k}(s)) \rho_0(ds) \leq \|V^\star - V^{\pi_k}\|_\infty.$$

■

Appendix E. Standard analytic tools

This appendix collects standard analytic inputs used in the fixed-drift Langevin estimate and the uniform bounds. These tools are stated separately so that the paper's main proof remains focused on the Bellman residual/resolvent mechanism.

E.1 Gaussian entropy, smoothing, and bounded perturbations

Lemma E.1 (Finite Gaussian-relative entropy gives a second moment) *Let μ be a probability measure on $\mathbb{R}^d$. If $\text{KL}(\mu\|\rho_\beta) < \infty$, then $\int \|a\|^2 \mu(da) < \infty$. More quantitatively, for every $c \in (0, \beta/(2\tau))$,*

$$c \int_{\mathbb{R}^d} \|a\|^2 \mu(da) \leq \text{KL}(\mu\|\rho_\beta) + \log \int_{\mathbb{R}^d} e^{c\|a\|^2} \rho_\beta(a) da < \infty.$$

Proof We use the entropy variational inequality: for any probability measures μ, ν and any measurable function f such that the right-hand side is well-defined,

$$\int f d\mu \leq \text{KL}(\mu\|\nu) + \log \int e^f d\nu.$$

Applying this inequality with $\nu = \rho_\beta$ and $f(a) = c\|a\|^2$ gives

$$c \int_{\mathbb{R}^d} \|a\|^2 \mu(da) \leq \text{KL}(\mu \| \rho_\beta) + \log \int_{\mathbb{R}^d} e^{c\|a\|^2} \rho_\beta(a) da.$$

It remains to check that the Gaussian exponential moment is finite. Since

$$\rho_\beta(a) = Z_\beta^{-1} \exp\left(-\frac{\beta}{2\tau} \|a\|^2\right),$$

we have

$$\int_{\mathbb{R}^d} e^{c\|a\|^2} \rho_\beta(a) da = Z_\beta^{-1} \int_{\mathbb{R}^d} \exp\left(-\left(\frac{\beta}{2\tau} - c\right) \|a\|^2\right) da < \infty$$

whenever $c \in (0, \beta/(2\tau))$. Therefore the displayed bound holds, and in particular $\int \|a\|^2 \mu(da) < \infty$. $\blacksquare$

Lemma E.2 (Holley–Stroock bounded perturbation) *Let μ be a probability measure on $\mathbb{R}^n$ satisfying $\text{LSI}(\alpha)$, namely,*

$$\mathcal{I}(\nu \| \mu) \geq 2\alpha \text{KL}(\nu \| \mu), \quad \forall \nu \ll \mu.$$

Let $\psi \in L^\infty(\mu)$ and define

$$d\tilde{\mu} := \frac{e^\psi}{\int e^\psi d\mu} d\mu, \quad \text{Osc}(\psi) := \text{ess sup } \psi - \text{ess inf } \psi.$$

Then $\tilde{\mu}$ satisfies $\text{LSI}(\tilde{\alpha})$ with

$$\tilde{\alpha} \geq e^{-\text{Osc}(\psi)} \alpha.$$

Equivalently,

$$\mathcal{I}(\nu \| \tilde{\mu}) \geq 2\alpha e^{-\text{Osc}(\psi)} \text{KL}(\nu \| \tilde{\mu}), \quad \forall \nu \ll \tilde{\mu}.$$

Lemma E.3 (Quantitative Gaussian KL after smoothing) *Let $\rho_\beta(a) = Z_\beta^{-1} \exp(-\beta\|a\|^2/(2\tau))$ on $\mathbb{R}^d$. Let $Y = X + \sigma\xi$, where $\sigma > 0$, $\xi \sim \mathcal{N}(0, I_d)$ is independent of X , and $\mathbb{E}\|Y\|^2 < \infty$. Then $\nu := \text{Law}(Y)$ has a density and*

$$\text{KL}(\nu \| \rho_\beta) \leq \frac{\beta}{2\tau} \mathbb{E}\|Y\|^2 + \log Z_\beta - \frac{d}{2} \log(2\pi e \sigma^2). \quad (61)$$

In particular, if $\sigma^2 = 2\tau\eta$ and $\mathbb{E}\|Y\|^2 \leq M$, then

$$\text{KL}(\nu \| \rho_\beta) \leq \frac{\beta M}{2\tau} + \log Z_\beta - \frac{d}{2} \log(4\pi e \tau \eta). \quad (62)$$

Moreover, if $p(a) = Z_p^{-1} e^{\psi(a)} \rho_\beta(a)$ with $\|\psi\|_\infty \leq C$, then every probability measure μ satisfying $\text{KL}(\mu \| \rho_\beta) < \infty$ obeys

$$\text{KL}(\mu \| p) \leq \text{KL}(\mu \| \rho_\beta) + 2C. \quad (63)$$

Proof [Proof of Lemma E.3] Let

$$g_\sigma(z) := (2\pi\sigma^2)^{-d/2} \exp\left(-\frac{\|z\|^2}{2\sigma^2}\right)$$

be the density of $\sigma\xi$. If $\mu_X = \text{Law}(X)$, then the law $\nu = \text{Law}(Y)$ has the Lebesgue density

$$q(y) = \int_{\mathbb{R}^d} g_\sigma(y-x) \mu_X(dx).$$

Since $g_\sigma > 0$, we have $q(y) > 0$ for every $y \in \mathbb{R}^d$.

We first justify the entropy lower bound without assuming that X has a density. The joint law of (X, Y) is

$$P_{X,Y}(dx, dy) = \mu_X(dx) g_\sigma(y-x) dy,$$

whereas $P_X \otimes \nu$ is

$$(P_X \otimes \nu)(dx, dy) = \mu_X(dx) q(y) dy.$$

Therefore $P_{X,Y} \ll P_X \otimes \nu$, and the mutual information satisfies

$$\begin{aligned} 0 &\leq I(X; Y) := \text{KL}(P_{X,Y} \| P_X \otimes \nu) \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \log\left(\frac{g_\sigma(y-x)}{q(y)}\right) g_\sigma(y-x) dy \mu_X(dx) \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} g_\sigma(y-x) \log g_\sigma(y-x) dy \mu_X(dx) - \int_{\mathbb{R}^d} q(y) \log q(y) dy \\ &= -h(\sigma\xi) + h(Y). \end{aligned}$$

Equivalently,

$$h(Y) \geq h(\sigma\xi) = \frac{d}{2} \log(2\pi e\sigma^2).$$

This argument is understood in the usual extended-real sense; the last display gives a finite lower bound. On the other hand, the Gaussian maximum-entropy inequality under the second-moment constraint gives a finite upper bound on $h(Y)$, since $\mathbb{E}\|Y\|^2 < \infty$. Hence $h(Y)$ is a finite real number.

Expanding the relative entropy to the Gaussian reference gives

$$\text{KL}(\nu \| \rho_\beta) = \int_{\mathbb{R}^d} q(y) \log q(y) dy - \int_{\mathbb{R}^d} q(y) \log \rho_\beta(y) dy = \frac{\beta}{2\tau} \mathbb{E}\|Y\|^2 - h(Y) + \log Z_\beta.$$

Using the entropy lower bound above proves

$$\text{KL}(\nu \| \rho_\beta) \leq \frac{\beta}{2\tau} \mathbb{E}\|Y\|^2 + \log Z_\beta - \frac{d}{2} \log(2\pi e\sigma^2),$$

which is (61). The bound (62) follows by substituting $\sigma^2 = 2\tau\eta$ and $\mathbb{E}\|Y\|^2 \leq M$.

For the bounded-perturbation claim, write $Z_p = \int e^\psi d\rho_\beta$. Since $\|\psi\|_\infty \leq C$, we have

$$e^{-C} \leq Z_p \leq e^C, \quad \log Z_p \leq C.$$

Moreover, if $\text{KL}(\mu\|\rho_\beta) < \infty$, then $\mu \ll \rho_\beta$, and since p is equivalent to ρ_β , the following computation is legitimate:

$$\text{KL}(\mu\|p) = \text{KL}(\mu\|\rho_\beta) - \int \psi d\mu + \log Z_p \leq \text{KL}(\mu\|\rho_\beta) + C + C.$$

This proves (63). ■