

Posterior contraction rates in Sobolev norms and Bayesian derivative estimation for infinite-dimensional exponential families

Emanuele Dolera*, Stefano Favaro^{*,†} and Matteo Giordano*

*University of Pavia, *University of Turin and [†]Collegio Carlo Alberto, Turin

Abstract

We study posterior contraction in positive-order Sobolev norms and Bayesian derivative estimation for infinite-dimensional exponential families. We embed the natural parameter in a Hilbert scale and model it via a standard Gaussian series prior expanded in the eigenbasis generating the scale. Under a two-sided link condition on the Fisher information and suitable local regularity assumptions, we show that smoothness-matching priors achieve minimax-optimal posterior contraction rates in any Hilbert scale norm up to the regularity of the ground truth. Our analysis builds on the novel approach to posterior contraction based on the Wasserstein distance recently introduced by [Dolera et al. \(2024b\)](#). It combines refined Laplace-type estimates for infinite-dimensional integrals associated to the posterior kernels with a mixed-geometry estimate controlling their stability under fluctuations in the data, itself resting on a tailored Poincaré inequality for posterior distributions conditioned on neighbourhoods of the truth. We apply the general theory to density estimation with a logistic parametrisation, Poisson intensity estimation with an exponential link, and the Gaussian white-noise model, yielding minimax contraction rates in Sobolev norms across all three settings. In particular, these yield optimal recovery of density score functions and derivatives of Poisson intensities.

Keywords. Derivative estimation; Frequentist analysis of posterior distributions; Gaussian priors; Infinite-dimensional exponential families; Wasserstein distance.

Contents

1	Introduction	2
1.1	Contributions	4
1.2	Basic notation and organisation of the paper	6
2	Infinite-dimensional exponential families in Hilbert scales	7
2.1	Hilbert scales	7
2.2	Infinite-dimensional exponential families	9
2.3	The Bayesian approach	11

3	Posterior contraction rates by Wasserstein dynamics	12
3.1	The Wasserstein dynamics approach to posterior contraction rates	12
3.2	A general posterior contraction rate theorem for Gaussian priors	14
3.2.1	Model assumptions	14
3.2.2	Main Result	16
4	Applications	18
4.1	Density estimation	18
4.2	Poisson Intensity estimation	21
4.3	The white noise model	24
5	Proofs	26
5.1	Bounds for the deterministic term	26
5.2	Bounds for the stochastic term	34
5.2.1	Construction of the localising event	37
5.2.2	Bounds for the conditioning errors	40
5.2.3	Transport techniques for conditioned posteriors	43

1 Introduction

Nonparametric Bayesian methods have become a standard approach to inference in infinite-dimensional statistical models. Within the Bayesian framework, a prior probability measure is placed on the unknown parameter and updated through the likelihood to produce the posterior distribution, from which point estimates, uncertainty quantification, and predictions are obtained. Even when the posterior is not available in closed form, it can often be explored numerically using powerful Markov chain Monte Carlo techniques. This unified treatment of the core tasks of statistical inference, together with the flexibility to encode structural information in the prior and the availability of advanced computational procedures, accounts for the widespread use of nonparametric Bayesian methods across the sciences. We refer to the monograph by [Ghosal and Van der Vaart \(2017\)](#), as well as to [Hjort et al. \(2010\)](#); [Müller et al. \(2015\)](#); [Castillo \(2024\)](#), for broad accounts of the nonparametric Bayesian approach, encompassing modelling, computation, applications, and theory.

The practical appeal of nonparametric Bayesian methods makes a rigorous assessment of their performance particularly important. In infinite-dimensional models, genuinely subjective prior beliefs are typically difficult to elicit, and the prior is often chosen partly for its regularising properties, controlling features of the parameter that are only weakly informed by the likelihood. A natural question is therefore whether, from the frequentist perspective in which the data are generated by a fixed ground truth, a poorly calibrated prior can lead to suboptimal inference for the true parameter or even prevent its recovery altogether. In regular finite-dimensional models, the Bernstein–von Mises theorem provides a broadly positive answer: as the sample size grows, the posterior contracts at the parametric rate, while credible sets are asymptotically valid confidence sets with optimal diameter ([Van der Vaart, 2000](#), Chapter 10). On the other hand, in the infinite-dimensional framework, no analogous general principle holds, as demonstrated by classical counterexamples of ([Diaconis and Freedman, 1986](#); [Cox, 1993](#); [Freedman, 1999](#)), among others.

This issue has motivated an extensive literature on the frequentist analysis of nonparametric Bayesian methods, with early foundational posterior consistency results established by [Doob \(1949\)](#) and [Schwartz \(1965\)](#). Their quantitative refinement, through the notion of contraction rates, was developed in the seminal works of [Ghosal et al. \(2000\)](#) and [Shen and Wasserman \(2001\)](#), which gave rise to an intensely active area of research that now encompasses a wide range of statistical models, prior distributions, loss functions, and analytical techniques ([Ghosal and van der Vaart, 2007](#); [Walker et al., 2007](#); [van der Vaart and van Zanten, 2008](#); [Giné and Nickl, 2011](#); [Rousseau and Mengersen, 2011](#)); see ([Giné and Nickl, 2016](#), Chapter 7.3); ([Ghosal and Van der Vaart, 2017](#), Chapters 8 and 9), and [Castillo \(2024\)](#), for a comprehensive treatment.

A defining feature of this line of work is that posterior contraction rates are typically established in statistical metrics determined by the testing geometry of the experiment under consideration, such as the Hellinger distance for independent and identically distributed (i.i.d.) observations ([Ghosal and Van der Vaart, 2017](#), Chapter 8). To obtain recovery guarantees for the parameter itself, these statistical metrics must then be related to the distance representing the inferential target. For several problems involving integrated losses, such as ones of L^1 - and L^2 -type, this comparison is rate-preserving and yields minimax-optimal contraction ([Ghosal and Van der Vaart, 2017](#), Chapter 9). By contrast, for stronger metrics, such as the supremum norm or positive-order Sobolev distances, this generally requires inversion inequalities or interpolation arguments and may lead to suboptimal results. In such cases, ad hoc studies tailored to specific model and prior combinations can still result in optimal contraction rates ([Castillo, 2014](#); [Yoo and Ghosal, 2016](#); [Shen and Ghosal, 2017](#), e.g.,). However, to the best of our knowledge, no comparably general techniques exists for frequentist analyses of posterior distributions in strong norms not directly controlled by the likelihood, for which, without additional structural restrictions, uniformly exponentially consistent tests at the desired scale are not guaranteed to exist.

Here, we are interested in posterior contraction in positive-order Sobolev norms, motivated by the task of inferring the derivatives of an unknown function. Such norms jointly control the distance between an estimate and its target, as well as the discrepancy between their mixed partial derivatives up to the corresponding order, thereby guaranteeing simultaneous recovery of the parameter and its differential structure. Derivatives are themselves central objects in many applications: they describe marginal effects and local rates of change in regression ([Härdle and Stoker, 1989](#)), determine modes and other shape features of probability density functions (p.d.f.’s) ([Genovese et al., 2016](#)), and underpin mean-shift methods for clustering, pattern recognition, and image processing ([Comaniciu and Meer, 2002](#)). Another prominent inferential target is the score function of an unknown p.d.f., whose estimation lies at the heart of score matching and modern score-based generative modelling, including diffusion models ([Song and Ermon, 2019](#)).

There is by now an extensive frequentist literature on derivative estimation, including foundational contributions by [Bhattacharya \(1967\)](#); [Singh \(1977\)](#); [Stone \(1982\)](#). More recently, minimax convergence rates under Sobolev norms were established by [Comte et al. \(2020\)](#) in density estimation and by [Fischer and Steinwart \(2020\)](#) in nonparametric regression models. See also [Wibisono et al. \(2024\)](#) for optimal results in score estimation problems. Theoretical results for the nonparametric Bayesian approach to the problem have developed mainly only over the past decade, and remain largely model- and prior-specific, reflecting

the aforementioned challenge in pursuing testing-based arguments for posterior contraction in stronger metrics than the Hellinger distance.

In multivariate nonparametric regression, [Yoo and Ghosal \(2016\)](#) obtained minimax pointwise, L^2 - and supremum-norm contraction rates for mixed partial derivatives using tensor-product B-spline priors. Adaptive rates were established by [Yoo et al. \(2018\)](#) under wavelet spike-and-slab priors and, recently, by [Liu and Li \(2026\)](#) for plug-in differentiated Gaussian-process posteriors. In the Gaussian white-noise model, [Dolera et al. \(2024a\)](#) proved optimal contraction in Sobolev norms and derivative recovery using Besov–Laplace series priors; see also earlier related results in [Castillo and Nickl \(2013\)](#). For density estimation, the only result we are aware of is due to [Shen and Ghosal \(2017\)](#), who derived minimax L^2 -contraction rates for the density derivatives under B-spline series priors. Lastly, we note that results for Gaussian sequence models and linear inverse problems (e.g., [Belitser and Ghosal, 2003](#); [Knapik et al., 2011](#); [Gugushvili et al., 2020](#)) can also accommodate derivative estimation by casting this problem in the framework of mildly ill-posed inverse problems.

Recently, [Dolera et al. \(2024b\)](#) introduced an alternative, non-testing-based route to the frequentist analysis of posterior distributions in general metric spaces, built on a characterisation of contraction rates through the Wasserstein distance ([Ambrosio et al., 2008](#)). They developed general bounding techniques that combine the Laplace methods for integral approximations [Wong \(2001\)](#) with the dynamic formulation of the Wasserstein distance ([Benamou and Brenier, 2000](#)) and applied the resulting framework to several applications, including an investigation of posterior contraction in Sobolev norms for infinite-dimensional exponential families and for density estimation under the logistic parametrisation. Their analysis, however, only applies to oracle Gaussian series priors whose covariance operators are required to be diagonal in an eigenbasis of the Fisher information at the ground truth cf. ([Dolera et al., 2024b](#), Section 3.1), and results in suboptimal contraction rates for the density derivatives, cf. ([Dolera et al., 2024b](#), Section 4.4).

1.1 Contributions

In the present paper, we build on the earlier investigation of [Dolera et al. \(2024b\)](#) to study posterior contraction in Sobolev norms and derivative estimation in infinite-dimensional exponential families ([Pistone and Sempì, 1995](#); [Canu and Smola, 2006](#); [Fukumizu, 2009](#)). These generalise classical parametric exponential families ([Brown, 1986](#)) by allowing the natural parameter to range over an infinite-dimensional Hilbert space and the data to enter the likelihood through a sufficient statistic taking values in its topological dual; see Section 2.2 for definitions and details. This class retains the convenient analytic structure of its parametric counterpart, yielding a convex and smooth log-partition function whose first two derivatives identify the mean map and the Fisher information, and whose associated Bregman divergence coincides with the Kullback–Leibler divergence. At the same time, infinite-dimensional exponential families accommodate several important statistical models through a suitable specification of their natural parameter and sufficient statistic, including density estimation, Poisson intensity estimation and Gaussian white noise, cf. Examples 2.3–2.5.

The asymptotic properties of frequentist estimators in infinite-dimensional exponential families have been investigated in several articles, notably in connection with score match-

ing and kernel methods (Fukumizu, 2009; Gretton et al., 2012; Sriperumbudur et al., 2017; Fukumizu, 2024). The nonparametric Bayesian literature on the topic is instead comparatively sparse, with available contributions including the Bernstein–von Mises-type Gaussian approximations of Shang and Cheng (2018), alongside the aforementioned posterior contraction analysis of Dolera et al. (2024b). Although a vast body of work provides contraction rates for the individual models encompassed by this framework, as reviewed above, general techniques yielding optimal rates in positive-order Sobolev norms remain unavailable. Our work aims to help fill this gap.

To formally encode the notion of Sobolev-type norms and smoothness within the considered infinite-dimensional exponential families, we embed the parameter space in a Hilbert scale $\{H_s\}_{s \in \mathbb{R}}$ (Krein and Petunin, 1966). This provides a unified framework to characterise the interplay among the strength of the loss, the regularity of the prior and the smoothness of the ground truth, and allows a concise presentation of the main model assumptions as comparisons between certain linear operators and fixed powers of the scale generator. Hilbert scales are standard tools in the analysis of ill-posed problems (e.g., Natterer, 1984) and are routinely employed to formulate regularity conditions in Bayesian inversion (Knapik et al., 2011, 2016; Gugushvili et al., 2020); here they play an analogous role. For the concrete families considered in Examples 2.3–2.5, the resulting Hilbert scales coincide, up to equivalent norms, with standard collections of fractional Sobolev spaces on bounded domains, while accommodating any required boundary or identifiability constraints, such as mean-zero conditions.

In this setting, we model the unknown parameter through Gaussian series priors built on the fixed eigenbasis generating the Hilbert scale, with vanishing coordinate variances; see Section 2.3. The rate of this decay is a central tuning parameter that directly controls the regularity of the prior draws. Priors of this kind form a canonical class for statistical models whose parameter space is a separable Hilbert space (e.g., Belitser and Ghosal, 2003; Knapik et al., 2011; Castillo and Nickl, 2013; Knapik et al., 2016; Gugushvili et al., 2020). In our main result, Theorem 3.5, we show that if the ground truth is in H_β , the posterior distribution resulting from an α -regular Gaussian series prior contracts in H_s -norm at rate $n^{-(\alpha \wedge \beta - s)/(2\alpha + d)}$, for all $0 \leq s < \alpha \wedge \beta$. Here, $d \in \mathbb{N}$ is an effective dimension index encoded by the asymptotic growth of the eigenvalues of the scale generator. For smoothness-matching priors (i.e., $\alpha = \beta$), the obtained H_s -rate is thus equal to $n^{-(\beta - s)/(2\beta + d)}$ for all $0 \leq s < \beta$, coinciding with the usual minimax rate of estimation in Sobolev norms of order s for β -smooth functions defined on d -dimensional domains (Stone, 1982).

Our proofs build on the novel approach to posterior contraction introduced by Dolera et al. (2024b), which bypasses testing arguments by directly controlling the expected Wasserstein distance between the posterior distribution and the Dirac mass at the ground truth, with transport cost induced by the metric of interest. This distance is upper bounded by the sum of a deterministic concentration term, governed by the local interaction between the likelihood and the prior near the true parameter, and a stochastic stability term that reflects the sensitivity of the posterior to fluctuations in the data; see Section 3.1. Such decomposition then allows these two summands to be treated via powerful analytical and probabilistic tools tailored to their structure. In particular, we control the deterministic term through Laplace-type integral bounds (Wong, 2001), refining the analysis of Dolera et al. (2024b) by replacing their oracle requirement that the prior covariance operator and the

Fisher information at the ground truth be simultaneously diagonalisable with a two-sided link condition comparing the latter with a fixed power of the scale generator, cf. Section 5.1. For the stochastic component, we prove a mixed-geometry stability estimate for the posterior kernels arising in infinite-dimensional exponential families; see Section 5.2. This result, which is of independent interest, rests on a Poincaré inequality for the posterior distribution conditioned on a neighbourhood of the truth, which we establish through Galerkin approximation and Brascamp–Lieb-type arguments (Bakry et al., 2008). The proof carefully decouples the strong geometry of the loss from the weaker geometry in which the sufficient statistic concentrates, thereby avoiding the algebraic loss in the rate incurred by the single-geometry strategy of Dolera et al. (2024b). See the discussion after Theorem 3.5 for further insights.

In Section 4, we apply our general theory to the three concrete statistical models introduced in Examples 2.3–2.5. Under smoothness-matching Gaussian series priors, we obtain minimax contraction rates in Sobolev norms of every order (up to the regularity of the ground truth) for density estimation with a logistic parametrisation (Theorem 4.1), Poisson intensity estimation with an exponential link (Theorem 4.2), and the Gaussian white-noise model (Theorem 4.3). Consequently, differentiated push-forward posteriors optimally recover the corresponding derivatives of the natural parameter, and, through the employed smooth parametrisation, also the derivatives of the target p.d.f. or intensity function. The analysis of the white noise model in fact hinges on simpler arguments, whereby conjugacy and coordinate-wise factorisation allow a direct treatment of the deterministic and stochastic terms, recovering known results from the Gaussian sequence and linear inverse problem literature (Belitser and Ghosal, 2003; Knapik et al., 2011; Gugushvili et al., 2020). For density estimation, the obtained results extend the B -spline analysis of Shen and Ghosal (2017) to Gaussian series priors expanded in standard bases generating the Sobolev scale, such as the Fourier and wavelet bases. In particular, the case $s = 1$ yields posterior contraction towards the true score function at the minimax rate $n^{-(\beta-1)/(2\beta+d)}$, or equivalently at the squared rate $n^{-2(\beta-1)/(2\beta+d)}$ relative to Fisher divergence (Wibisono et al., 2024). To the best of our knowledge, the application to Poisson processes provides the first optimal contraction rates for derivatives of a nonparametric intensity function.

1.2 Basic notation and organisation of the paper

For a given measure space $(\mathcal{X}, \mathfrak{X}, \mu)$ and a given normed vector space $(\mathcal{V}, \|\cdot\|_{\mathcal{V}})$, let $L^p(\mathcal{X}, \mu; \mathcal{V})$, $p \in [1, \infty]$, be the usual Lebesgue spaces of $\mathcal{V}$ -valued functions on $\mathcal{X}$ with integrable p^{th} -power, equipped with norm $\|h\|_{L^p(\mathcal{X}, \mu; \mathcal{V})}^p := \int_{\mathcal{X}} \|h(x)\|_{\mathcal{V}}^p d\mu(x)$, $h \in L^p(\mathcal{X}, \mu; \mathcal{V})$, replaced by the essential supremum of $\|h\|_{\mathcal{V}}$ over $\mathcal{X}$ if $p = \infty$. For open $\mathcal{X} \subseteq \mathbb{R}^m$, we denote by $C(\mathcal{X})$ the space of bounded continuous real-valued functions defined on $\mathcal{X}$, equipped with the supremum norm. For positive integers $\alpha \in \mathbb{N}$, let $C^\alpha(\mathcal{X})$ denote the space of α -times continuously differentiable real-valued functions defined on $\mathcal{X}$ whose partial derivatives up to order α are bounded, equipped with the norm $\|\cdot\|_{C^\alpha}$. Further, let $H^\alpha(\mathcal{X})$ be the usual Hilbert-Sobolev space of functions with square-integrable partial derivatives up to order α , with norm $\|\cdot\|_{H^\alpha}$. When no confusion may arise, we omit the dependence of the function spaces on the domain.

We denote the minimum and maximum between real numbers $a, b \in \mathbb{R}$ by $a \wedge b$ and $a \vee b$,

respectively, and use the symbols $\lesssim$, $\gtrsim$ and $\asymp$ for one- and two-sided inequalities holding up to multiplicative constants. For two real sequences $(a_n)_{n \geq 1}$, $(b_n)_{n \geq 1}$, we write $a_n = o(b_n)$ if $a_n/b_n \rightarrow 0$ as $n \rightarrow \infty$, and $a_n = O(b_n)$ if $a_n/b_n \lesssim 1$ for all sufficiently large n .

The rest of this article is organised as follows. Section 2 introduces Hilbert scales, infinite-dimensional exponential families and the Bayesian setup. Section 3 outlines the Wasserstein dynamic approach to posterior contraction rates, provides all model assumptions and states the main theorem. Section 4 is devoted to the applications to three concrete statistical models. Section 5 collects the proof of the main result.

2 Infinite-dimensional exponential families in Hilbert scales

In this section we recall the definition of infinite-dimensional exponential families (Pistone and Sempi, 1995; Canu and Smola, 2006; Fukumizu, 2009), casting it within the framework of Hilbert scales (Krein and Petunin, 1966; Natterer, 1984; Engl et al., 1996), where the notion of Sobolev norms and all model assumptions are made precise.

2.1 Hilbert scales

Throughout, we assume the existence of a reference separable Hilbert space H_0 , with inner product $\langle \cdot, \cdot \rangle_{H_0}$, norm $\| \cdot \|_{H_0}$ and orthonormal basis $\{e_k\}_{k=1}^\infty$, and fix a nondecreasing sequence of strictly positive numbers $\{\lambda_k\}_{k=1}^\infty$ with $\lambda_k \rightarrow \infty$. For $s \geq 0$, define the Hilbert space

$$H_s := \left\{ h = \sum_{k=1}^\infty h_k e_k : \|h\|_{H_s}^2 := \sum_{k=1}^\infty \lambda_k^{2s} h_k^2 < \infty \right\}, \quad \langle h, g \rangle_{H_s} := \sum_{k=1}^\infty \lambda_k^{2s} h_k g_k.$$

For $s < 0$ let H_s be the completion of H_0 under $\| \cdot \|_{H_s}$, with $\| \cdot \|_{H_s}$ formally defined as above; equivalently, H_s is the topological dual of H_{-s} under the $\langle \cdot, \cdot \rangle_{H_0}$ -pairing. The family $\{H_s\}_{s \in \mathbb{R}}$ is the *Hilbert scale* generated by the pairs $\{(e_k, \lambda_k)\}_{k=1}^\infty$; see, e.g., Krein and Petunin (1966); Natterer (1984); Engl et al. (1996). For every $s \in \mathbb{R}$, an orthonormal basis of H_s is given by $\{\lambda_k^{-s} e_k\}_{k=1}^\infty$. We recall standard properties used repeatedly in the following:

- for $s < r$, H_r is a dense and compactly embedded subspace of H_s ;
- for $s_1 \leq s \leq s_2$ with $s = (1-a)s_1 + as_2$, $a \in [0, 1]$, we have the interpolation inequality

$$\|h\|_{H_s} \leq \|h\|_{H_{s_1}}^{1-a} \|h\|_{H_{s_2}}^a, \quad h \in H_{s_2}. \quad (1)$$

Equivalently, the Hilbert scale is generated by the (unbounded, densely defined) self-adjoint linear operator

$$L e_k = \lambda_k e_k, \quad k \in \mathbb{N}, \quad (2)$$

on H_0 , with domain $D(L) = H_1$. The real powers of L are defined through spectral calculus: for $r \in \mathbb{R}$, $L^r e_k := \lambda_k^r e_k$, with domain $D(L^r) = \{h \in H_0 : \sum_{k=1}^\infty \lambda_k^{2r} h_k^2 < \infty\}$. Then, for any $s \in \mathbb{R}$, $L^r : H_s \rightarrow H_{s-r}$ is an isometric isomorphism; in particular, for $r > 0$, one has

$D(L^r) = H_r$ and $\|h\|_{H_r} = \|L^r h\|_{H_0}$, whereas the restriction of L^{-r} to H_0 defines a bounded (in fact, compact) operator. In the following, we assume the polynomial growth

$$\lambda_k \asymp k^{1/d}, \quad \text{as } k \rightarrow \infty, \quad \text{for some } d \in \mathbb{N}, \quad (3)$$

matching, in view of Weyl's law (e.g., [Taylor, 2011](#), Chapter 8), the eigenvalue asymptotics of positive elliptic (pseudo-)differential operators of order one on d -dimensional domains. This recovers canonical scales of Hilbert-Sobolev spaces ([Evans, 1998](#)), possibly under appropriate boundary conditions and up to equivalent norms.

Example 2.1 (Periodic Sobolev spaces). On the d -dimensional torus $\mathbb{T}^d$, interpreted as the unit hypercube $[0, 1]^d$ with opposite endpoints identified, we take as reference $H_0 = L^2(\mathbb{T}^d)$, the space of 1-periodic square-integrable functions, with orthonormal basis given by the Fourier system $e_\xi(\cdot) := e^{2\pi i \langle \xi, \cdot \rangle}$, $\xi \in \mathbb{Z}^d$, namely the eigenfunctions of the Laplacian on $\mathbb{T}^d$,

$$-\Delta e_\xi := -\sum_{j=1}^d \frac{\partial^2}{\partial x_j^2} e_\xi = (2\pi|\xi|)^2 e_\xi, \quad e_\xi(\cdot + m) = e_\xi(\cdot), \quad m \in \mathbb{Z}^d, \quad \xi \in \mathbb{Z}^d. \quad (4)$$

Noting that the constant eigenfunction e_0 lies in the kernel of $-\Delta$, we take the positive order-one operator $L = (I - \Delta)^{1/2}$ as the scale generator, with eigenvalues $(1 + (2\pi|\xi|)^2)^{1/2} \geq 1$. Enumerating the frequencies by increasing modulus as $\{\xi_k\}_{k \geq 0}$, the lattice-point count $|\{\xi \in \mathbb{Z}^d : |\xi| \leq u\}| \asymp u^d$ gives $\lambda_k = (1 + (2\pi|\xi_k|)^2)^{1/2} \asymp k^{1/d}$, so that the polynomial growth (3) holds with d the dimension of $\mathbb{T}^d$. The induced scale is the periodic Hilbert-Sobolev scale $H_s = H^s(\mathbb{T}^d)$, $s \in \mathbb{R}$, with

$$\|h\|_{H^s}^2 = \sum_{\xi \in \mathbb{Z}^d} (1 + (2\pi|\xi|)^2)^s |\hat{h}(\xi)|^2 = \|(I - \Delta)^{s/2} h\|_{L^2}^2, \quad \hat{h}(\xi) := \langle h, e_\xi \rangle_{L^2}.$$

For nonnegative integers $s \in \mathbb{N}_0$, $H^s(\mathbb{T}^d)$ coincides, with equivalent norms, with the standard periodic Sobolev space $W^{s,2}(\mathbb{T}^d)$ comprising all functions h with partial derivatives $D^\alpha h \in L^2(\mathbb{T}^d)$ for all multi-indices $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}_0^d$ such that $|\alpha| \leq s$, normed by $\|h\|_{W^{s,2}}^2 = \sum_{|\alpha| \leq s} \|D^\alpha h\|_{L^2}^2$. This follows from differentiating term by term the Fourier series representation of any function $h \in H^s(\mathbb{T}^d)$. In fact, for $s = 0, 1$, the norms coincide.

General boundary behaviours can be accommodated through other bases, for example ones of boundary-corrected wavelets.

Example 2.2 (Wavelet Sobolev scale on $[0, 1]^d$). Let $\{\psi_k\}_{k=1}^\infty$ be a single-index reordering by increasing resolution level of a tensor-product orthonormal wavelet basis of $L^2([0, 1]^d)$, built from S -regular ($S \in \mathbb{N}$), compactly supported, boundary-corrected Daubechies wavelets (see, e.g., [Giné and Nickl, 2016](#), Chapter 4.3). Setting $\lambda_k := k^{1/d}$, the induced scale $H_s = H^s([0, 1]^d)$, normed by

$$\|h\|_{H^s}^2 = \sum_{k=1}^\infty k^{2s/d} |\langle h, \psi_k \rangle_{L^2}|^2,$$

satisfies, in view of the wavelet characterisation of standard Sobolev spaces, $H^s([0, 1]^d) = W^{s,2}([0, 1]^d)$ and $\|\cdot\|_{H^s} \asymp \|\cdot\|_{W^{s,2}}$ for all integers $0 \leq s \leq S$, cf. ([Giné and Nickl, 2016](#), Chapter 4.3).

Depending on the statistical model at hand, further subspaces contained within the underlying Hilbert scale can be singled out through the series representation. For example, as the constant function is a basis element in both Examples 2.1 and 2.2, corresponding respectively to the zero-frequency mode e_0 and the coarsest scaling function ψ_1 , removing it imposes the mean-zero constraint and gives rise to the associated scales of homogeneous spaces $\dot{H}^s(\mathbb{T}^d)$ and $\dot{H}^s([0, 1]^d)$, $s \in \mathbb{R}$. This provides natural parameter spaces for models identifiable only up to an additive constant, such as density estimation under a logistic parametrisation; see Example 2.3.

2.2 Infinite-dimensional exponential families

Fix $p \geq 0$ and take as parameter space the Hilbert space $\Theta := H_p$. Throughout, we denote $\langle \cdot, \cdot \rangle_\Theta := \langle \cdot, \cdot \rangle_{H_p}$ and $\| \cdot \|_\Theta := \| \cdot \|_{H_p}$. Let $(\mathbb{X}, \mathcal{X}, \lambda)$ be a σ -finite measure space and let $\beta : \mathbb{X} \rightarrow \Theta^* = H_{-p}$ be a measurable map. Following Pistone and Sempi (1995); Canu and Smola (2006); Fukumizu (2009) (see also Definition 3 of Dolera et al., 2024b), the associated *regular infinite-dimensional exponential family* is the collection of λ -densities

$$f_\theta(x) := \exp \{ \beta_x(\theta) - M(\theta) \}, \quad M(\theta) := \log \int_{\mathbb{X}} e^{\beta_x(\theta)} d\lambda(x), \quad x \in \mathbb{X}, \quad \theta \in \Theta, \quad (5)$$

where $\beta_x(\theta)$ denotes the action of the linear operator $\beta_x \in \Theta^*$ on $\theta \in \Theta$, and the log-partition (or cumulant generating) function M is assumed to be finite for all $\theta \in \Theta$. The family (5) is expressed in natural form, with natural parameter space equal to Θ .

We assume we are given a random sample $X^{(n)} := (X_1, \dots, X_n)$ of independent and identically distributed (i.i.d.) $\mathbb{X}$ -valued random variables drawn from an unknown probability density function (p.d.f.) belonging to the exponential family,

$$X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f_\theta, \quad (6)$$

for some $\theta \in \Theta$ and f_θ as in (5). The goal is then to estimate θ from observations $X^{(n)}$. We denote by $d\mu_\theta := f_\theta d\lambda$, $\lambda^{(n)} := \lambda^{\otimes n}$, and write $\mu_\theta^{(n)} := \mu_\theta^{\otimes n}$ for the data distribution and $E_\theta^{(n)}$ for the expectation with respect to $\mu_\theta^{(n)}$. The likelihood equals

$$L^{(n)}(\theta) := \frac{d\mu_\theta^{(n)}}{d\lambda^{(n)}}(X^{(n)}) = \exp \left\{ n \left[\hat{T}_n(\theta) - M(\theta) \right] \right\}, \quad \theta \in \Theta, \quad (7)$$

with Θ^* -valued sufficient statistic

$$\hat{T}_n := \frac{1}{n} \sum_{i=1}^n \beta_{X_i}. \quad (8)$$

Infinite-dimensional exponential families provide a flexible yet analytically tractable framework for nonparametric inference, encompassing several widely used statistical models that are recovered through different choices of the map β and dominating measure λ . Below, we concretise our theoretical results on the following three canonical examples from statistical theory; see Section 4 for full details.

Example 2.3 (Density estimation on $[0, 1]^d$). We observe a random sample $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f$ for some unknown p.d.f. $f : [0, 1]^d \rightarrow (0, \infty)$, modelled through the logistic parametrisation $f_\theta = e^\theta / \int_{[0, 1]^d} e^{\theta(x)} dx$ for some $\theta : [0, 1]^d \rightarrow \mathbb{R}$. Since the latter is identifiable only up to an additive constant, we take Θ to be a mean-zero wavelet Sobolev space $\dot{H}^p([0, 1]^d)$ for some $p > d/2$, constructed as in the remark after Example 2.2. In view of the Sobolev embedding (e.g., [Giné and Nickl, 2016](#), Chapter 4.3), Θ is continuously embedded into the space of continuous functions $C([0, 1]^d)$. It follows that for each $x \in [0, 1]^d$ the point evaluation functional δ_x is linear and continuous on Θ , and we recover an infinite dimensional exponential family with $\beta_x = \delta_x$ and $M(\theta) = \log \int_{[0, 1]^d} e^{\theta(x)} dx$.

Example 2.4 (Poisson intensity estimation on $[0, 1]^d$). We observe n independent copies $N_1, \dots, N_n$ of an inhomogeneous Poisson point process N on $[0, 1]^d$ with unknown intensity function $\rho : [0, 1]^d \rightarrow (0, \infty)$, modelled through the exponential link $\rho = e^\theta$ for some $\theta : [0, 1]^d \rightarrow \mathbb{R}$. That is, $N = \sum_{j=1}^J \delta_{X_j}$, arising as

$$J \sim \text{Po} \left(\int_{[0, 1]^d} e^{\theta(x)} dx \right), \quad X_1, \dots, X_J | J \stackrel{\text{iid}}{\sim} \frac{e^\theta dx}{\int_{[0, 1]^d} e^{\theta(x)} dx}.$$

For any measurable and bounded function $\theta \in L^\infty([0, 1]^d)$, the law μ_θ of N is absolutely continuous with respect to the distribution μ_0 of the unit-intensity Poisson process (i.e., with $\theta \equiv 0$), with Radon-Nikodym derivative ([Kutoyants, 1998](#), Chapter 1),

$$\frac{d\mu_\theta}{d\mu_0}(N) = \exp \left\{ \sum_{j=1}^J \theta(X_j) - \int_{[0, 1]^d} [e^{\theta(x)} - 1] dx \right\}.$$

Here, θ is fully identifiable, and we take Θ to be a wavelet Sobolev space $H^p([0, 1]^d)$ from Example 2.2, with $p > d/2$. Arguing similarly to Example 2.3, we then recover an infinite-dimensional exponential family with $\beta_N := \sum_{j=1}^J \delta_{X_j} = N$ and $M(\theta) = \int_{[0, 1]^d} [e^{\theta(x)} - 1] dx$.

Example 2.5 (Signal in white noise on $[0, 1]^d$). We observe an unknown signal $\theta \in L^2([0, 1]^d)$ corrupted by additive Gaussian white noise at level $n^{-1/2}$, that is, a realisation of the Gaussian process $Y^{(n)}$ given by

$$Y^{(n)}(h) = \langle \theta, h \rangle_{L^2} + \frac{1}{\sqrt{n}} \mathbb{W}(h), \quad h \in L^2([0, 1]^d),$$

where $\mathbb{W}$ is a white noise process indexed by $L^2([0, 1]^d)$, with $\mathbb{W}(h) \sim \mathcal{N}(0, \|h\|_{L^2}^2)$ and $\text{Cov}(\mathbb{W}(h), \mathbb{W}(g)) = \langle h, g \rangle_{L^2}$ ([Giné and Nickl, 2016](#), Chapter 1). Equivalently, given the wavelet basis $\{\psi_k\}_{k=1}^\infty$ of Example 2.2, one observes the Gaussian sequence

$$Y_k^{(n)} := Y^{(n)}(\psi_k) = \langle \theta, \psi_k \rangle_{L^2} + n^{-1/2} Z_k, \quad Z_k \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1).$$

For any $\theta \in L^2([0, 1]^d)$, the law μ_θ of $Y^{(n)}$ is absolutely continuous with respect to the white noise distribution (corresponding to $\theta \equiv 0$), with Radon-Nikodym derivative ([Giné and Nickl, 2016](#), Section 6.1.1)

$$\frac{d\mu_\theta}{d\mu_0}(Y^{(n)}) = \exp \left\{ n \left[Y^{(n)}(\theta) - \frac{1}{2} \|\theta\|_{L^2}^2 \right] \right\}.$$

This formally recovers an infinite-dimensional exponential family, with parameter space $\Theta = L^2([0, 1]^d)$, sufficient statistic $\hat{T}_n = Y^{(n)}$ and $M(\theta) = \frac{1}{2}\|\theta\|_{L^2}^2$. Unlike the sampling models of Examples 2.3 and 2.4, we note that the white-noise experiment is not generated by i.i.d. draws, and that the sufficient statistic $\hat{T}_n$ only takes values in the dual Sobolev space $H^{-s}([0, 1]^d)$ for $s > d/2$ (Giné and Nickl, 2016, Section 4.4), and thus in particular not in Θ . The general techniques for the analysis of the infinite-dimensional exponential family (6) however readily extend to this case, cf. Section 4.

2.3 The Bayesian approach

We adopt the Bayesian approach to the problem of estimating the unknown parameter θ from data $X^{(n)}$ arising as in (6), modelling θ via a (prior) Borel probability measure Π on Θ . Note that the log-partition function M in (5) is convex and lower semicontinuous on Θ (cf. Brown, 1986, Theorems 1.13, 2.2 and 2.7); hence, M is Borel measurable, and so is the likelihood $L^{(n)}$ in (7). By Bayes' formula (e.g., Ghosal and Van der Vaart, 2017, p. 7), the resulting posterior distribution then equals

$$\Pi(A|X^{(n)}) = \frac{\int_A e^{n[\hat{T}_n(\theta) - M(\theta)]} d\Pi(\theta)}{\int_{\Theta} e^{n[\hat{T}_n(\theta) - M(\theta)]} d\Pi(\theta)}, \quad A \subseteq \Theta \text{ measurable.} \quad (9)$$

In writing (9) we implicitly assume the denominator to be finite, namely that

$$\int_{\Theta} e^{n[\tau(\theta) - M(\theta)]} d\Pi(\theta) < \infty, \quad \text{for all } n \in \mathbb{N} \text{ and } \tau \in \Theta^*. \quad (10)$$

This quantity is strictly positive in view of the model assumption $M(\theta) < \infty$ for every $\theta \in \Theta$; hence, $\Pi(\cdot|X^{(n)})$ in (9) is well defined under (10). For $\tau \in \Theta^*$, we use the notation $\Pi_n^*(\cdot|\tau)$ for the probability measure defined by the right hand side of (9) with $\hat{T}_n$ replaced by τ . Accordingly, $\Pi(\cdot|X^{(n)}) = \Pi_n^*(\cdot|\hat{T}_n)$.

In the following, we investigate the asymptotic behaviour of the posterior distribution as the sample size $n \rightarrow \infty$, under the frequentist assumption that the data have been generated by some fixed *ground truth* $\theta_0 \in \Theta$, namely $X^{(n)} \sim \mu_{\theta_0}^{(n)}$. We characterise the speed of concentration of $\Pi(\cdot|X^{(n)})$ around θ_0 according to the usual notion of *posterior contraction rates*, that is, positive real sequences $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ such that

$$E_{\theta_0}^{(n)} \left[\Pi(\theta \in \Theta : \ell(\theta, \theta_0) > M_n \varepsilon_n \mid X^{(n)}) \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (11)$$

for every $M_n \rightarrow \infty$, where ℓ is a (semi-)metric on Θ . See Ghosal and Van der Vaart (2017) for an in-depth treatment of the frequentist analysis of nonparametric Bayesian procedures. In particular, we are interested in posterior contraction rates relative to the Hilbert scale norms introduced in Section 2.1, taking $\ell(\theta_1, \theta_2) = \|\theta_1 - \theta_2\|_{H_s}$ for suitable $s \geq 0$. For the concrete scales of Examples 2.1 and 2.2, the resulting metrics are Sobolev norms, so that contraction in $\|\cdot\|_{H_s}$ quantifies the recovery of θ_0 jointly with its derivatives up to order s , the case $s = 0$ corresponding to the usual L^2 -distance.

The concentration in strong norms of Bayesian procedures for infinite-dimensional exponential families has been recently investigated by Dolera et al. (2024b). Their analysis

however hinges on oracle-type Gaussian priors calibrated to features of the unknown ground truth, and only results in suboptimal rates. An earlier contribution, developing Bernstein–von Mises-type approximations, is by [Shang and Cheng \(2018\)](#). Results for frequentist estimators and tests in this framework were derived by [Fukumizu \(2009\)](#); [Gretton et al. \(2012\)](#); [Sriperumbudur et al. \(2017\)](#), among the others.

There is an extensive nonparametric Bayesian literature for the concrete models introduced in Examples 2.3–2.5. Landmark contributions include [Scricciolo \(2006\)](#); [van der Vaart and van Zanten \(2008\)](#); [Rivoirard and Rousseau \(2012\)](#) for density estimation under the logistic parametrisation, [Belitser et al. \(2015\)](#); [Kirichenko and van Zanten \(2015\)](#); [Donnet et al. \(2017\)](#) for Poisson intensity estimation, and [Belitser and Ghosal \(2003\)](#); [Knapik et al. \(2011\)](#); [Castillo and Nickl \(2013\)](#) for the white noise model. We further point to the monograph by [Ghosal and Van der Vaart \(2017\)](#), where many further references can be found.

3 Posterior contraction rates by Wasserstein dynamics

We provide our main result concerning the concentration in Hilbert scale norms of the posterior distribution $\Pi(\cdot|X^{(n)})$ in (9) around the ground truth θ_0 generating the data (6). Our analysis is based on the novel *Wasserstein dynamics* approach to posterior contraction rates in general metric spaces recently developed by [Dolera et al. \(2024b\)](#), which we significantly refine in the context of infinite-dimensional exponential families. To clarify the model assumptions underlying the abstract theorem, we outline the proof argument in the next section.

3.1 The Wasserstein dynamics approach to posterior contraction rates

The starting point is a general upper bound for posterior contraction rates expressed in terms of the Wasserstein distance between probability measures. Recall $\Theta = H_p$ for some $p \geq 0$, and let $s \geq 0$ be the index of the target Hilbert scale norm for posterior contraction. Throughout, we assume that $\theta_0 \in H_\beta$ for some $\beta > s \vee p$, and that the prior (and hence the posterior) charge $H_s \cap \Theta = H_{s \vee p}$ with probability one. Let $\mathcal{P}_2(H_s)$ be the collection of all Borel probability measures ν on H_s satisfying

$$\int_{H_s} \|\theta - \theta^*\|_{H_s}^2 d\nu(\theta) < \infty, \quad \text{for some } \theta^* \in H_s, \quad (12)$$

and consider the 2-*Wasserstein distance*

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_1, \nu_2) := \sqrt{\inf_{\gamma \in \mathcal{F}(\nu_1, \nu_2)} \int_{H_s \times H_s} \|\theta_1 - \theta_2\|_{H_s}^2 d\gamma(\theta_1, \theta_2)}, \quad (13)$$

the infimum running over the class $\mathcal{F}(\nu_1, \nu_2)$ of all couplings of ν_1 and ν_2 , that is, Borel probability measures on $\Theta \times \Theta$ with marginals ν_1 and ν_2 ; see ([Ambrosio et al., 2008](#), Chapter 7) for details. Under the above requirements and a mild integrability condition ensuring that the prior and the posterior satisfy (12), Lemma 1 of [Dolera et al. \(2024b\)](#) shows by simple arguments that the quantity

$$\varepsilon_n := E_{\theta_0}^{(n)} [\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\Pi(\cdot|X^{(n)}), \delta_{\theta_0})], \quad (14)$$

where δ_{θ_0} denotes the Dirac mass at θ_0 , is an upper bound for the H_s -posterior contraction rate at θ_0 , in that (11) holds for every $M_n \rightarrow \infty$ as $n \rightarrow \infty$. We then proceed to bound ε_n in (14) through a combination of analytical and probabilistic techniques.

First, recall from Section 2.3 that $\Pi(\cdot|X^{(n)}) = \Pi_n^*(\cdot|\hat{T}_n)$ with $\hat{T}_n = \frac{1}{n} \sum_{i=1}^n \beta_{X_i}$ the sufficient statistic from (8). For all $\theta \in \Theta$ such that $E_\theta^{(1)}[\|\beta_{X_1}\|_{\Theta^*}] < \infty$, denote

$$T_\theta := E_\theta^{(1)}[\beta_{X_1}] = \int_{\mathbb{X}} \beta_x f_\theta(x) d\lambda(x), \quad (15)$$

where the above integral is in the Bochner sense and belongs to Θ^* . Then, provided that $E_{\theta_0}^{(1)}[\|\beta_{X_1}\|_{\Theta^*}] < \infty$, the strong law of large numbers in Hilbert spaces (Ledoux and Talagrand, 1991, Corollary 7.10) implies that $\|\hat{T}_n - T_{\theta_0}\|_{\Theta^*} \rightarrow 0$ $\mu_{\theta_0}^{(\infty)}$ -almost surely as $n \rightarrow \infty$. This suggests that the asymptotic behaviour of $\Pi_n^*(\cdot|\hat{T}_n)$ should be closely related to that of the deterministic measure $\Pi_n^*(\cdot|T_{\theta_0})$ obtained upon replacing $\hat{T}_n$ in (9) by its limit T_{θ_0} , as long as the Markov kernel $\Pi_n^*(\cdot|\tau)$, $\tau \in \Theta^*$, is stable with respect to its second argument in the relevant geometry. Based on this observation, we apply the triangle inequality for the Wasserstein distance to upper bound ε_n in (14) by

$$\varepsilon_n \leq \mathcal{W}_2^{\mathcal{P}_2(H_s)}(\Pi_n^*(\cdot|T_{\theta_0}), \delta_{\theta_0}) + E_{\theta_0}^{(n)} \left[\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\Pi_n^*(\cdot|T_{\theta_0}), \Pi_n^*(\cdot|\hat{T}_n)) \right]. \quad (16)$$

The first summand in (16) is a purely *deterministic term* that measures the speed of concentration of $\Pi_n^*(\cdot|T_{\theta_0})$ at θ_0 . Its analytical evaluation requires the study of the asymptotics of the ratio between two vanishing integrals, cf. (19) below, which we tackle via an infinite-dimensional extension of the classical Laplace methods for integral approximation developed by Dolera et al. (2024b). See Section 5.1. This hinges on rewriting $\Pi_n^*(\cdot|T_{\theta_0})$ in terms of the Kullback–Leibler divergence

$$K(\vartheta|\theta) := \int_{\mathbb{X}} \log \frac{f_\theta(x)}{f_\vartheta(x)} f_\theta(x) d\lambda(x), \quad \vartheta, \theta \in \Theta. \quad (17)$$

By (5), $\log[f_\theta(x)/f_\vartheta(x)] = \beta_x(\theta - \vartheta) + M(\vartheta) - M(\theta)$. Integrating against $f_\theta d\lambda$ and recalling the definition of T_θ in (15) yields

$$K(\vartheta|\theta) = M(\vartheta) - M(\theta) - \int_{\mathbb{X}} \beta_x(\vartheta - \theta) f_\theta(x) d\lambda(x) = M(\vartheta) - M(\theta) - T_\theta(\vartheta - \theta), \quad (18)$$

the properties of Bochner integrals allowing to exchange the order of integration and application of linear functionals in Θ^* . Thus, $-nK(\theta|\theta_0) = n[T_{\theta_0}(\theta) - M(\theta)] + n[M(\theta_0) - T_{\theta_0}(\theta_0)]$ for any $\theta \in \Theta$, and since the second term does not depend on θ , recalling (9), we obtain

$$d\Pi_n^*(\theta|T_{\theta_0}) = \frac{e^{n[T_{\theta_0}(\theta) - M(\theta)]}}{\int_{H_{s \vee p}} e^{n[T_{\theta_0}(\theta) - M(\theta)]} d\Pi(\theta)} d\Pi(\theta), \quad \theta \in H_{s \vee p},$$

and finally,

$$\begin{aligned} \mathcal{W}_2^{\mathcal{P}_2(H_s)}(\Pi_n^*(\cdot|T_{\theta_0}), \delta_{\theta_0}) &= \sqrt{\int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 d\Pi_n^*(\theta|T_{\theta_0})} \\ &= \sqrt{\frac{\int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{\int_{H_{s \vee p}} e^{-nK(\theta|\theta_0)} d\Pi(\theta)}}. \end{aligned} \quad (19)$$

The second summand in (16) is a *stochastic term* that measures the effect on the posterior of the fluctuation of $\hat{T}_n$ around T_{θ_0} . Its analysis rests on two ingredients. The first is a localisation: the expectation is split over an event of probability tending to one, on which the fluctuation of the sufficient statistic is small and the posterior charges a fixed neighbourhood of θ_0 up to a superpolynomially small remainder, and over its complement, where a crude second-moment bound suffices. The second is a stability estimate for the posterior kernel, obtained by connecting T_{θ_0} and $\hat{T}_n$ through the path $\tau_t = T_{\theta_0} + t(\hat{T}_n - T_{\theta_0})$, $t \in [0, 1]$, and using the dynamic formulation of the Wasserstein distance (Benamou and Brenier, 2000) to estimate the length in $\mathcal{P}_2(H_s)$ of the corresponding curve $\{\Pi_n^*(\cdot \mid \tau_t)\}_{t \in [0, 1]}$ of posterior kernels, conditioned on that neighbourhood. The resulting bound builds on the stability estimates of Dolera and Mainini (2020, 2023) and refines the analysis of Dolera et al. (2024b) by exploiting, within the transport problem underlying the Wasserstein dynamics, the joint structure of the exponential family and the Hilbert scale. See Section 5.2 for details.

3.2 A general posterior contraction rate theorem for Gaussian priors

We are now in a position to state and prove our main posterior contraction rate theorem. We assume that $\theta_0 \in H_\beta$ for some $\beta > p$, and consider the centred Gaussian series prior Π on the orthonormal basis $\{e_k\}_{k=1}^\infty$ of H_0 defined, for a given prior regularity parameter $\alpha > p$, as the law of

$$\theta = \sum_{k=1}^{\infty} \lambda_k^{-\alpha-d/2} t_k e_k, \quad t_k \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1). \quad (20)$$

The deterministic scaling in (20) calibrates the prior regularity within the Hilbert scale. Indeed, in view of the assumed eigenvalue asymptotics (3), $E[\|\theta\|_{H_s}^2] = \sum_{k=1}^{\infty} \lambda_k^{2(s-\alpha)-d} < \infty$ if and only if $s < \alpha$; accordingly, the draws from Π belong almost surely to H_s for every $s < \alpha$, while they almost surely belong to no H_s with $s \geq \alpha$. As customary, we then call Π an α -regular Gaussian series prior. In particular, Π is supported on $\Theta = H_p$ precisely by virtue of the requirement $\alpha > p$.

Equivalently, for L the scale generator from (2), Π defines a centred Gaussian Borel probability measure on the ambient space H_0 , with covariance operator $Q_\Pi = L^{-2\alpha-d}$. The associated reproducing kernel Hilbert space (RKHS, or Cameron–Martin space) is given by $H_\Pi = H_{\alpha+d/2}$, with $\|h\|_{H_\Pi}^2 = \langle h, Q_\Pi^{-1}h \rangle_{H_0} = \|h\|_{H_{\alpha+d/2}}^2$. We refer to (Giné and Nickl, 2016, Chapter 2) for definitions and background properties of Gaussian processes and measures used throughout.

For θ_0 and Π as above, we pursue H_s -posterior contraction rates via the Wasserstein dynamics approach outlined in Section 3.1. The following model assumptions formalise the conditions under which explicit, matching algebraic rates can be obtained for the deterministic and stochastic terms appearing in the upper bound (16). These are formulated relative to the abstract infinite-dimensional exponential family (5), and are later concretised in the three statistical models introduced in Examples 2.3–2.5.

3.2.1 Model assumptions

The main assumptions are formulated in terms of the *Fisher information* I , defined, for every $\theta \in \Theta$ for which the mean T_θ in (15) exists, as the nonnegative (extended) quadratic

form

$$I(\theta)[h] := \text{Var}_\theta^{(1)}(\beta_{X_1}(h)) = E_\theta^{(1)}\left[(\beta_{X_1}(h) - T_\theta(h))^2\right] \in [0, \infty], \quad h \in \Theta. \quad (21)$$

At any point $\theta \in \Theta$ where the log-partition function M from (5) is twice (Fréchet) differentiable, the identity (18) identifies T_θ and $I(\theta)$ with the first two derivatives of $M(\theta)$, respectively,

$$DM(\theta) = T_\theta, \quad D^2M(\theta)[h, h] = I(\theta)[h], \quad h \in \Theta,$$

whence, for $K(\vartheta|\theta)$, the Kullback–Leibler divergence from (17), $\theta, \vartheta \in \Theta$,

$$D_\vartheta K(\vartheta|\theta) = T_\vartheta - T_\theta, \quad D_\vartheta^2 K(\vartheta|\theta)[h, h] = I(\vartheta)[h]. \quad (22)$$

According to the second identity in (22), I is the curvature of the Kullback–Leibler divergence in its first argument, which regulates the asymptotics of the deterministic term in (16) via the representation (19). At the same time, through (21), $I(\theta)$ also governs the fluctuations of the sufficient statistic $\hat{T}_n$ from (8) around T_θ , which determine the decay of the stochastic term in (16), as

$$\text{Var}_\theta^{(n)}(\hat{T}_n(h)) = \frac{1}{n} I(\theta)[h], \quad h \in \Theta. \quad (23)$$

The following central assumption requires a pointwise upper bound and a local coercivity bound for the Fisher information, both expressed in the Hilbert-scale geometry.

Assumption 3.1 (Coercivity and boundedness of the Fisher information). There exists a constant $R_0 > 0$ such that $E_\theta^{(1)}[\|\beta_{X_1}\|_{\Theta^*}] < \infty$ for all $\theta \in B_{R_0}^\Theta(\theta_0) := \{\theta \in \Theta : \|\theta - \theta_0\|_\Theta \leq R_0\}$. Furthermore, there exist constants $0 < c_0 \leq C_0 < \infty$ such that

$$I(\theta_0)[h] \leq C_0 \|h\|_{H_0}^2, \quad \text{for all } h \in \Theta, \quad (24)$$

and

$$I(\theta)[h] \geq c_0 \|h\|_{H_0}^2, \quad \text{for all } h \in \Theta \text{ and } \theta \in B_{R_0}^\Theta(\theta_0). \quad (25)$$

Assumption 3.1 entails that $c_0 L^{-2p} \preceq I(\theta_0) \preceq C_0 L^{-2p}$ in the sense of quadratic forms on Θ , with the lower bound holding over a neighbourhood of θ_0 . Here $L^{-2p} : \Theta \rightarrow \Theta^*$ is understood as the map with associated form

$$h \in \Theta \mapsto \langle h, L^{-2p} h \rangle_\Theta = \|L^{-p} h\|_\Theta^2 = \|h\|_{H_0}^2.$$

The upper bound (24) guarantees that $I(\theta_0)$ is finite and bounded on Θ , since $\|h\|_{H_0} \leq \lambda_1^{-p} \|h\|_\Theta$. By polarisation it therefore extends to a bounded symmetric bilinear form on $\Theta \times \Theta$, induced by a unique bounded operator $I(\theta_0) : \Theta \rightarrow \Theta^*$ that coincides with the covariance operator of β_{X_1} under f_{θ_0} ,

$$(I(\theta_0)[h])(g) = \text{Cov}_{\theta_0}^{(1)}(\beta_{X_1}(g), \beta_{X_1}(h)), \quad g, h \in \Theta.$$

The two comparison operators are (multiples of) negative powers of L , and are diagonalised by the orthonormal basis $\{\lambda_k^{-p} e_k\}_{k=1}^\infty$ of Θ , with eigenvalues $\{c_0 \lambda_k^{-2p}\}_{k=1}^\infty$ and $\{C_0 \lambda_k^{-2p}\}_{k=1}^\infty$, respectively. In view of (20), the curvature of the Kullback–Leibler divergence is comparable,

up to the constants c_0, C_0 , to an operator that is diagonal in the same basis as the prior covariance. This is crucial for bounding both the deterministic and stochastic terms in (16), allowing for explicit evaluation of certain Gaussian quadratic form ratios appearing in the analysis.

The second assumption formalises the differentiability of the Kullback–Leibler divergence and its link to the Fisher information via (22), and imposes a second-order Taylor expansion around θ_0 with Hölder-type remainder.

Assumption 3.2 (Expansion of the Kullback–Leibler divergence). M is twice continuously Fréchet differentiable on Θ . Furthermore, there exists an index $m_K \in (d/2, p]$ and constants $q_K \in (0, 1]$ and $C_K, R_K > 0$ such that

$$|K(\theta|\theta_0) - \tfrac{1}{2}I(\theta_0)[\theta - \theta_0]| \leq C_K \|\theta - \theta_0\|_{H_{m_K}}^{2+q_K}, \quad \text{for all } \theta \in \Theta \cap B_{R_K}^{H_{m_K}}(\theta_0). \quad (26)$$

Next, we assume a (mild) global lower bound for the Kullback–Leibler divergence in terms of a negative-order Hilbert scale norm. Combined with properties of the employed Gaussian series prior, this underpins a basic localisation property for the posterior distribution around θ_0 as $n \rightarrow \infty$, allowing the asymptotic analysis to be restricted to neighbourhoods where (25) and (26) apply.

Assumption 3.3 (Global lower bound for the Kullback–Leibler divergence). There exist a nondecreasing function $\varphi : [0, \infty) \rightarrow [0, \infty)$ with $\varphi(u) > 0$ for $u > 0$ and $\varphi(u) \gtrsim u^2$ as $u \rightarrow 0^+$, and constants $c_\varphi > 0$ and $r_K \geq 0$, such that

$$K(\theta|\theta_0) \geq \varphi(c_\varphi \|\theta - \theta_0\|_{H_{-r_K}}), \quad \theta \in \Theta. \quad (27)$$

In the next assumption, we quantify the fluctuations of $\hat{T}_n$ around T_{θ_0} in a weak Hilbert scale geometry.

Assumption 3.4 (Concentration of the sufficient statistic). For the index m_K from Assumption 3.2, there exists a constant $A_T > 0$ such that

$$E_{\theta_0}^{(1)} \exp \left\{ \|\beta_{X_1} - T_{\theta_0}\|_{H_{-m_K}} / A_T \right\} < \infty. \quad (28)$$

The restriction $m_K \leq p$ guarantees, in view of the continuous embedding $H_{-m_K} \subseteq \Theta^*$, that $\|\hat{T}_n - T_{\theta_0}\|_{\Theta^*}$ is almost surely finite. On the other hand, (28) implies that $\|\beta_{X_1} - T_{\theta_0}\|_{H_{-m_K}}$ has sub-exponential tails, and hence moments of all orders. In view of (21), Assumption 3.1, the series representation of $\|\cdot\|_{H_{-m_K}}$ and the eigenvalue asymptotics (3), this necessarily requires $m_K > d/2$.

3.2.2 Main Result

Under the above prior specification and model assumptions, the following posterior concentration theorem in Hilbert scale norms is obtained. Its proof is deferred to Section 5.

Theorem 3.5. *Let $\{f_\theta, \theta \in H_p\}$, for some $p \geq 0$, be an infinite-dimensional exponential family as in (5). Let $\theta_0 \in H_\beta$, $\beta > p$, and let Π be an α -regular Gaussian series prior as in*

(20), for some $\alpha > p$. Let Assumptions 3.1–3.4 be satisfied for some $q_K \in (0, 1]$, $r_K \geq 0$, $m_K \in (d/2, p]$. Further assume that

$$q_K(\alpha \wedge \beta) > \max \{ (2 + q_K)p + 2r_K, (2 + q_K)m_K + d \}, \quad (29)$$

and that, in the oversmoothing case $\alpha > \beta$, in addition

$$\alpha < \frac{(2 + q_K)(\beta - m_K) - d}{2}. \quad (30)$$

Then, for every $0 \leq s < \alpha \wedge \beta$ and every $M_n \rightarrow \infty$,

$$E_{\theta_0}^{(n)} \left[\Pi(\theta \in \Theta : \|\theta - \theta_0\|_{H_s} > M_n n^{-\frac{(\alpha \wedge \beta) - s}{2\alpha + d}} \mid X^{(n)}) \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Conditions (29)–(30) are jointly equivalent to

$$q_K(\alpha \wedge \beta) > (2 + q_K)p + 2r_K, \quad (2 + q_K)((\alpha \wedge \beta) - m_K) > 2\alpha + d,$$

the second of which reduces to a lower bound on α when $\alpha \leq \beta$ and to the upper bound (30) when $\alpha > \beta$. In the concrete models studied in Section 4 one has $q_K = 1$ and may take m_K , p and r_K arbitrarily close to $d/2$, so that (29) reads $\alpha \wedge \beta > \frac{5}{2}d$ and (30) reads $\alpha < \frac{3}{2}\beta - \frac{5}{4}d$.

Theorem 3.5 substantially refines the results in Section 3.1 of Dolera et al. (2024b). Firstly, we note that their explicit bounds for infinite-dimensional exponential families (Dolera et al., 2024b, Propositions 2 and 5) require the prior covariance operator and the Fisher information $I(\theta_0)$ to be simultaneously diagonalised. Since outside linear models the eigenfunctions of $I(\theta_0)$ generally depends nontrivially on the unknown truth θ_0 , this leads to an oracle prior specification. Our two-sided link condition in Assumption 3.1 avoids this issue by comparing the Fisher information with fixed powers of the Hilbert-scale generator, thereby allowing Gaussian series priors in a standard, truth-independent basis.

Furthermore, our result also yields a quantitative improvement in the obtained rates. For the smoothness-matching case $\alpha = \beta$, Theorem 3.5 shows that the posterior concentrates around $\theta_0 \in H_\beta$ in H_s -norm at rate $n^{-(\beta-s)/(2\beta+d)}$, which represents the typical minimax rate for estimating β -smooth functions in H_s -losses (Stone, 1982). In Section 4, we show that this is indeed the case for the concrete models introduced in Examples 2.3–2.5, over the range $\beta > \frac{5}{2}d$.

For comparison, under the eigenvalue asymptotics (3), formally applying the argument of Dolera et al. (2024b) with $p = s$ as the index for the parameter space and target geometry yields (for the posterior arising from their oracle prior) the H_s -rate $n^{-\frac{\beta-2s+d/2}{2\beta+d}}$. For every $s > d/2$, which by the discussion following Assumption 3.4 is the range in which the choice $p = s$ is admissible, this is slower than the minimax rate by the algebraic factor $n^{\frac{s-d/2}{2\beta+d}}$. This suboptimality arises because their single-geometry stability argument uses H_s both as the target geometry and to control the sufficient-statistic fluctuation. By decoupling these two geometries and deriving in Section 5.2 a sharper mixed-geometry local stability estimate for the posterior kernel in Wasserstein distance, Theorem 3.5 instead recovers in applications the minimax benchmark.

4 Applications

In this section, we apply the general techniques and results to the three concrete models introduced in Examples 2.3–2.5, obtaining in each of them explicit posterior contraction rates in Sobolev norms.

4.1 Density estimation

Recall the setting from Example 2.3. Let $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f_{\theta_0}$ be i.i.d. random variables with values in $[0, 1]^d$ drawn from some unknown p.d.f.

$$f_{\theta_0}(x) = \frac{e^{\theta_0(x)}}{\int_{[0,1]^d} e^{\theta_0(x)} dx}, \quad x \in [0, 1]^d, \quad (31)$$

for some continuous function $\theta_0 : [0, 1]^d \rightarrow \mathbb{R}$. As described in Example 2.3, we ensure identifiability by imposing the zero-integral constraint, through the mean-zero wavelet Sobolev scale $\dot{H}^t([0, 1]^d)$, $t \in \mathbb{R}$, constructed from Example 2.2, and taking the parameter space $\Theta = \dot{H}^p([0, 1]^d)$ for any $p > d/2$. The latter restriction ensures the continuous embedding $\dot{H}^p([0, 1]^d) \subseteq C([0, 1]^d)$, and, consequently, the continuity of the point evaluation functionals δ_x , $x \in [0, 1]^d$, on $H^p([0, 1]^d)$. With these choices, model (31) defines an infinite-dimensional exponential family with $\beta_x = \delta_x$ and $M(\theta) = \log \int_{[0,1]^d} e^{\theta(x)} dx$. We then have, for every $h \in \dot{H}^p([0, 1]^d)$,

$$T_\theta(h) = E_\theta^{(1)}[h(X_1)] = \int_{[0,1]^d} h(x) f_\theta(x) dx$$

and

$$I(\theta)[h] = \text{Var}_\theta^{(1)}(h(X_1)) = \int_{[0,1]^d} h(x)^2 f_\theta(x) dx - \left(\int_{[0,1]^d} h(x) f_\theta(x) dx \right)^2.$$

We verify Assumptions 3.1–3.4, explicitly identifying all involved constants. We follow the exposition in (Dolera et al., 2024b, Section 4.4). First, note that

$$\begin{aligned} \sup_{x \in [0,1]^d} \|\delta_x\|_{\dot{H}^{-p}} &= \sup_{x \in [0,1]^d} \left\{ \sup_{h \in \dot{H}^p: \|h\|_{H^p}=1} |h(x)| \right\} \\ &\leq \sup_{h \in \dot{H}^p: \|h\|_{H^p}=1} \|h\|_{L^\infty} \lesssim \sup_{h \in \dot{H}^p: \|h\|_{H^p}=1} \|h\|_{H^p} = 1. \end{aligned}$$

This shows that $E_\theta^{(1)}[\|\delta_{X_1}\|_{\dot{H}^{-p}}]$ is uniformly bounded in $\dot{H}^p([0, 1]^d)$. Next, since, for every $\theta \in \dot{H}^p([0, 1]^d)$, $\inf_{x \in [0,1]^d} f_\theta(x) \geq e^{-2\|\theta\|_{L^\infty}}$, the triangle inequality gives

$$\inf_{x \in [0,1]^d} f_\theta(x) \geq \exp \left\{ -2(\|\theta_0\|_{L^\infty} + c_p R_0) \right\} =: c_0, \quad (32)$$

for every $\theta \in B_{R_0}^\Theta(\theta_0)$, where $c_p > 0$ is the multiplicative constant in the continuous embedding $\dot{H}^p([0, 1]^d) \subset C([0, 1]^d)$. Moreover, since every $h \in \dot{H}^p([0, 1]^d)$ has zero Lebesgue

integral, the variational characterisation of variances yields

$$\begin{aligned} I(\theta)[h] &= \inf_{a \in \mathbb{R}} \int_{[0,1]^d} (h(x) - a)^2 f_\theta(x) dx \\ &\geq \left(\inf_{x \in [0,1]^d} f_\theta(x) \right) \inf_{a \in \mathbb{R}} \int_{[0,1]^d} (h(x) - a)^2 dx \geq c_0 \|h\|_{L^2}^2. \end{aligned} \quad (33)$$

On the other hand,

$$I(\theta_0)[h] \leq \int_{[0,1]^d} h(x)^2 f_{\theta_0}(x) dx \leq \|f_{\theta_0}\|_{L^\infty} \|h\|_{L^2}^2. \quad (34)$$

Recalling that $\dot{L}^2([0,1]^d) = \dot{H}^0([0,1]^d)$, this shows that, for any $R_0 > 0$, Assumption 3.1 holds with c_0 as in (32) and $C_0 := \|f_{\theta_0}\|_{L^\infty}$.

Turning to Assumption 3.2, in view of the continuous embedding $\dot{H}^p([0,1]^d) \subseteq C([0,1]^d)$ and the chain rule, M is infinitely Fréchet differentiable (in fact, analytic) on $\dot{H}^p([0,1]^d)$. Fix any $m_K \in (d/2, p]$. Its first three derivatives in any direction $h \in \dot{H}^p([0,1]^d)$ satisfy $DM(\theta)[h] = E_\theta^{(1)}[h(X_1)]$, $D^2M(\theta)[h, h] = \text{Var}_\theta^{(1)}(h(X_1)) = I(\theta)[h]$ and

$$D^3M(\theta)[h, h, h] = E_\theta^{(1)} \left[\left(h(X_1) - E_\theta^{(1)}[h(X_1)] \right)^3 \right].$$

In particular, $|D^3M(\theta)[h, h, h]| \leq 8\|h\|_{L^\infty}^3 \leq 8c_{m_K}^3 \|h\|_{H^{m_K}}^3$, and Taylor's formula applied to $K(\theta \mid \theta_0) = M(\theta) - M(\theta_0) - T_{\theta_0}(\theta - \theta_0)$ therefore yields

$$\left| K(\theta \mid \theta_0) - \frac{1}{2} I(\theta_0)[\theta - \theta_0] \right| \leq \frac{4c_{m_K}^3}{3} \|\theta - \theta_0\|_{H^{m_K}}^3,$$

for all $\theta \in \dot{H}^p([0,1]^d)$, whence Assumption 3.2 is seen to hold with $q_K = 1$, $C_K = 4c_{m_K}^3/3$ and any $R_K > 0$.

We next derive the uniform lower bound in weak geometry for the Kullback-Leibler divergence required by Assumption 3.3. For $\theta \in \dot{H}^p([0,1]^d)$, write $h := \theta - \theta_0$ and set

$$g := h - E_{\theta_0}^{(1)}[h(X_1)], \quad T_{\theta_0}(|g|) := E_{\theta_0}^{(1)}[|g(X_1)|].$$

Then $E_{\theta_0}^{(1)}[g(X_1)] = 0$ and

$$K(\theta \mid \theta_0) = \log E_{\theta_0}^{(1)} \left[e^{g(X_1)} \right].$$

Since $E_{\theta_0}^{(1)}[g_+(X_1)] = T_{\theta_0}(|g|)/2$, the elementary inequality $e^u \geq 1 + u + u_+^2/2$, $u \in \mathbb{R}$, gives

$$E_{\theta_0}^{(1)} \left[e^{g(X_1)} \right] \geq 1 + \frac{T_{\theta_0}(|g|)^2}{8}.$$

This yields a quadratic lower bound when $T_{\theta_0}(|g|)$ is bounded. For $T_{\theta_0}(|g|) \geq 2$, if $q = P_{\theta_0}^{(1)}(g(X_1) \geq 0)$, conditional Jensen's inequality gives

$$\begin{aligned} E_{\theta_0}^{(1)} \left[e^{g(X_1)} \right] &\geq E_{\theta_0}^{(1)} \left[e^{g(X_1)} \mathbf{1}_{g(X_1) \geq 0} \right] \\ &\geq P_{\theta_0}^{(1)}(g(X_1) \geq 0) \exp \left\{ \frac{T_{\theta_0}(|g|)}{2P_{\theta_0}^{(1)}(g(X_1) \geq 0)} \right\} \geq e^{T_{\theta_0}(|g|)/2}. \end{aligned}$$

Combining the two preceding displays, we find that

$$K(\theta \mid \theta_0) \gtrsim T_{\theta_0}(|g|)^2 \wedge T_{\theta_0}(|g|).$$

Now since both θ and θ_0 have zero Lebesgue integral, $h = g - \int_{[0,1]^d} g(x) dx$, and hence

$$\|h\|_{L^1} \leq 2\|g\|_{L^1} \leq \frac{2}{\inf_{x \in [0,1]} f_{\theta_0}(x)} T_{\theta_0}(|g|) \leq \frac{2}{e^{-2\|\theta_0\|_{L^\infty}}} T_{\theta_0}(|g|).$$

Therefore, since for every $r_K > d/2$, the continuous embedding $\dot{H}^{r_K}([0,1]^d) \subset L^\infty([0,1]^d)$ and duality give $L^1([0,1]^d) \subset \dot{H}^{-r_K}([0,1]^d)$, so that

$$K(\theta \mid \theta_0) \gtrsim \varphi(c_\varphi \|\theta - \theta_0\|_{\dot{H}^{-r_K}}),$$

for some $c_\varphi > 0$ and where $\varphi(u) := c(u^2 \wedge u)$, $u \geq 0$. This proves that Assumption 3.3 is valid for every $r_K > d/2$.

We proceed studying the concentration properties of the sufficient statistic $\hat{T}_n$ from (8). For the index $m_K > d/2$ fixed above, $B_{m_K} := \sup_{x \in [0,1]^d} \|\delta_x\|_{\dot{H}^{-m_K}} < \infty$. Set $Z_i := \delta_{X_i} - T_{\theta_0}$, $i = 1, \dots, n$. These are independent, centred $\dot{H}^{-m_K}$ -valued random elements satisfying $\|Z_i\|_{\dot{H}^{-m_K}} \leq 2B_{m_K}$ almost surely, whence

$$E_{\theta_0}^{(1)} \exp \{ \|\delta_{X_1} - T_{\theta_0}\|_{\dot{H}^{-m_K}} / A_T \} \leq e^{2B_{m_K}/A_T} < \infty,$$

for every $A_T > 0$. This is precisely Assumption 3.4. As $p > d/2$, the index m_K may be chosen arbitrarily close to $d/2$ while still satisfying $m_K \leq p$. Thus, all assumptions of Theorem 3.5 are verified for the density estimation model (31). The following posterior contraction result is obtained as a consequence.

Theorem 4.1. *Let $\theta_0 \in \dot{H}^\beta([0,1]^d)$, and let Π be an α -regular Gaussian series prior as in (20), for some $\alpha > 5d/2$ with $\alpha \leq \beta$. Assume data $X^{(n)} = (X_1, \dots, X_n)$ with $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} f_{\theta_0}$ for f_{θ_0} as in (31). Then, for every $0 \leq s < \alpha \wedge \beta$ and every $M_n \rightarrow \infty$,*

$$E_{\theta_0}^{(n)} \left[\Pi(\theta \in \Theta : \|\theta - \theta_0\|_{H^s} > M_n n^{-\frac{(\alpha \wedge \beta) - s}{2\alpha + d}} \mid X^{(n)}) \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Theorem 4.1 yields, for smoothness matching priors, optimal recovery of the partial derivatives of the natural parameter θ , in view of the equivalence $\|\cdot\|_{H^s} \asymp \|\cdot\|_{W^{s,2}}$ holding for all $s \in \mathbb{N}$. The same conclusion transfers to the p.d.f. f_θ : for any $t > d/2$ the space $H^t([0,1]^d)$ is closed under multiplication, and the map $\theta \mapsto f_\theta$ in (31) is Lipschitz from bounded subsets of $\dot{H}^t([0,1]^d)$ into $H^t([0,1]^d)$. Since the events in Theorem 4.1 may be taken to lie in a fixed $\dot{H}^t$ -ball for all n large, the derivatives of f_θ of order at most s therefore contract around those of f_{θ_0} at the same rate $n^{-(\alpha \wedge \beta - s)/(2\alpha + d)}$, for every integer s with $d/2 < s < \alpha \wedge \beta$.

The case $s = 1$ deserves separate mention, as it concerns the *score function* of the model. Since $\log f_\theta = \theta - M(\theta)$ with $M(\theta)$ not depending on x , the score of f_θ is exactly the gradient of the natural parameter, $\nabla \log f_\theta = \nabla \theta$, so that Theorem 4.1 applied with $s = 1$ gives, for every $M_n \rightarrow \infty$,

$$E_{\theta_0}^{(n)} \left[\Pi \left(\theta : \|\nabla \log f_\theta - \nabla \log f_{\theta_0}\|_{L^2} > M_n n^{-\frac{(\alpha \wedge \beta) - 1}{2\alpha + d}} \mid X^{(n)} \right) \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

attaining for $\alpha = \beta$ the minimax rate $n^{-(\beta-1)/(2\beta+d)}$ for score estimation (Wibisono et al., 2024). Equivalently, since f_{θ_0} is continuous and strictly positive on $[0, 1]^d$, and hence bounded above and below, the *Fisher divergence* satisfies

$$D_F(f_{\theta_0} \| f_{\theta}) := \int_{[0,1]^d} \|\nabla \log f_{\theta_0}(x) - \nabla \log f_{\theta}(x)\|^2 f_{\theta_0}(x) dx \asymp \|\nabla \theta - \nabla \theta_0\|_{L^2}^2,$$

and therefore by Theorem 4.1, the posterior $\Pi(\cdot | X^{(n)})$ contracts about f_{θ_0} if Fisher divergence at the squared rate $n^{-2((\alpha \wedge \beta)-1)/(2\alpha+d)}$, equal to $n^{-2(\beta-1)/(2\beta+d)}$ when $\alpha = \beta$.

4.2 Poisson Intensity estimation

We next consider the nonparametric intensity-estimation model introduced in Example 2.4. Let $N_1, \dots, N_n$ be independent Poisson point processes on $[0, 1]^d$ with unknown common intensity

$$\rho_{\theta_0}(x) = e^{\theta_0(x)}, \quad x \in [0, 1]^d, \quad (35)$$

for some continuous function $\theta_0 : [0, 1]^d \rightarrow \mathbb{R}$. Here, we take the parameter space $\Theta = H^p([0, 1]^d)$ for any $p > d/2$, constructed as in Example 2.2. Then, $H^p([0, 1]^d) \subset C([0, 1]^d)$ continuously and, for every $r > d/2$, finite signed measures on $[0, 1]^d$ define elements of $H^{-r}([0, 1]^d) = (H^r([0, 1]^d))^*$. Recall from Example 2.4 that, relative to the law of a unit-intensity Poisson point process, this model is an exponential family with $\beta_N = N = \sum_{j=1}^J \delta_{X_j}$ and $M(\theta) = \int_{[0,1]^d} (e^{\theta(x)} - 1) dx$.

We verify the assumptions of Theorem 3.5. Since $p > d/2$, as argued at the beginning of Section 4.1, $B_p := \sup_{x \in [0,1]^d} \|\delta_x\|_{H^{-p}} < \infty$. Consequently,

$$\|N\|_{H^{-p}} \leq \sum_{j=1}^J \|\delta_{X_j}\|_{H^{-p}} \leq B_p J,$$

and therefore, recalling $J \sim \text{Po}(\Lambda_\theta)$, $\Lambda_\theta := \int_{[0,1]^d} e^{\theta(x)} dx$,

$$E_\theta^{(1)}[\|N\|_{H^{-p}}] \leq B_p \Lambda_\theta.$$

The right-hand side is uniformly bounded over any H^p -neighbourhood of $B_{R_0}^\Theta(\theta_0)$, $R_0 > 0$, of θ_0 , because $\Lambda_\theta \leq e^{\|\theta\|_{L^\infty}} \leq \exp\{\|\theta_0\|_{L^\infty} + c_p R_0\}$, where $c_p > 0$ denote the multiplicative constant in the continuous embedding $H^p([0, 1]^d) \subset C([0, 1]^d)$. Next, by the standard moment identities for Poisson stochastic integrals (Kutoyants, 1998, Section 1.1),

$$T_\theta(h) = E_\theta^{(1)}[N(h)] = \int_{[0,1]^d} h(x) e^{\theta(x)} dx, \quad \text{for all } h \in H^p([0, 1]^d),$$

and

$$I(\theta)[h] = \text{Var}_\theta^{(1)}[N(h)] = \int_{[0,1]^d} h(x)^2 e^{\theta(x)} dx.$$

Let $c_p > 0$ denote the multiplicative constant in the continuous embedding $H^p([0, 1]^d) \subset L^\infty([0, 1]^d)$. For every θ satisfying $\|\theta - \theta_0\|_{H^p} \leq R_0$, we have $\|\theta\|_{L^\infty} \leq \|\theta_0\|_{L^\infty} + c_p R_0$ and hence

$$I(\theta)[h] \geq \exp\{-\|\theta_0\|_{L^\infty} - c_p R_0\} \|h\|_{L^2}^2,$$

whereas, at the truth,

$$I(\theta_0)[h] \leq e^{\|\theta_0\|_{L^\infty}} \|h\|_{L^2}^2.$$

This shows that Assumption 3.1 holds with any $R_0 > 0$ and with $c_0 = \exp\{-\|\theta_0\|_{L^\infty} - c_p R_0\}$ and $C_0 = e^{\|\theta_0\|_{L^\infty}}$.

We proceed with the verification of Assumption 3.2. Similarly to Section 4.1, The log-partition function M is smooth (in fact, analytic) on $H^p([0, 1]^d)$, and its first three derivatives along any direction $h \in H^p([0, 1]^d)$ are $DM(\theta)[h] = T_\theta(h)$, $D^2M(\theta)[h, h] = I(\theta)[h]$ and

$$D^3M(\theta)[h, h, h] = \int_{[0, 1]^d} h(x)^3 e^{\theta(x)} dx.$$

Fix any $m_K \in (d/2, p]$ and $R_K > 0$, and write $h = \theta - \theta_0$. If $\|h\|_{H^{m_K}} \leq R_K$, then, uniformly over $t \in [0, 1]$,

$$\|\theta_0 + th\|_{L^\infty} \leq \|\theta_0\|_{L^\infty} + c_{m_K} R_K.$$

It follows that

$$\begin{aligned} |D^3M(\theta_0 + th)[h, h, h]| &\leq \exp\{\|\theta_0\|_{L^\infty} + c_{m_K} R_K\} \|h\|_{L^\infty} \|h\|_{L^2}^2 \\ &\lesssim \exp\{\|\theta_0\|_{L^\infty} + c_{m_K} R_K\} \|h\|_{H^{m_K}}^3. \end{aligned}$$

Taylor's theorem therefore gives

$$\left| M(\theta) - M(\theta_0) - DM(\theta_0)[h] - \frac{1}{2} I(\theta_0)[h] \right| \leq C_K \|h\|_{H^{m_K}}^3$$

for a constant C_K depending only on θ_0 , m_K and R_K . This verifies Assumption 3.2 with $q_K = 1$.

Moving to Assumption 3.3, For $h = \theta - \theta_0$, the Kullback–Leibler divergence admits the explicit representation

$$K(\theta \mid \theta_0) = M(\theta) - M(\theta_0) - T_{\theta_0}(h) = \int_{[0, 1]^d} e^{\theta_0(x)} (e^{h(x)} - 1 - h(x)) dx.$$

The function $\psi(u) = e^u - 1 - u$, $u \in \mathbb{R}$, satisfies $\psi(u) \geq c_\psi (u^2 \wedge |u|)$ for a universal constant $c_\psi > 0$. Since $\inf_{x \in [0, 1]^d} e^{\theta_0(x)} \geq e^{-\|\theta_0\|_{L^\infty}}$, we obtain

$$K(\theta \mid \theta_0) \geq c_\psi e^{-\|\theta_0\|_{L^\infty}} \int_{[0, 1]^d} (h(x)^2 \wedge |h(x)|) dx.$$

To relate the right-hand side to the L^1 -norm of h , note that

$$\|h\|_{L^1} = \int_{\{|h| \leq 1\}} |h(x)| dx + \int_{\{|h| > 1\}} |h(x)| dx,$$

so that, by the Cauchy–Schwarz inequality on the set $\{x : |h(x)| \leq 1\}$, whose Lebesgue measure is at most one,

$$\begin{aligned} \int_{[0, 1]^d} (h^2(x) \wedge |h(x)|) dx &= \int_{\{x: |h(x)| \leq 1\}} h(x)^2 dx + \int_{\{x: |h(x)| > 1\}} |h(x)| dx \\ &\geq \left(\int_{\{x: |h(x)| \leq 1\}} |h(x)| dx \right)^2 + \int_{\{x: |h(x)| > 1\}} |h(x)| dx. \end{aligned}$$

Considering separately the cases $\|h\|_{L^1} \leq 1$ and $\|h\|_{L^1} > 1$, one can lower bound the right-hand side in the last display by $\frac{1}{4}(\|h\|_{L^1}^2 \wedge \|h\|_{L^1})$. Consequently,

$$K(\theta \mid \theta_0) \gtrsim \|\theta - \theta_0\|_{L^1}^2 \wedge \|\theta - \theta_0\|_{L^1},$$

and by proceeding similarly to Section 4.1 we obtain that Assumption 3.3 is valid with the choice $\varphi(u) = u^2 \wedge u$, for every $r_K > d/2$, for some $c_\varphi > 0$.

Finally, for the index $m_K > d/2$ fixed above, let $B_{m_K} := \sup_{x \in [0,1]^d} \|\delta_x\|_{H^{-m_K}} < \infty$ and write $dT_{\theta_0}(x) = e^{\theta_0(x)} dx$, $Z_i = N_i - T_{\theta_0}$. If $J_i = N_i([0,1]^d)$ denotes the total number of points of N_i , then $J_i \sim \text{Poisson}(\Lambda_{\theta_0})$ and

$$\|Z_i\|_{H^{-m_K}} \leq B_{m_K}(J_i + \Lambda_{\theta_0}).$$

Hence, for every $\lambda > 0$,

$$\begin{aligned} E_{\theta_0}^{(1)} [\exp \{ \lambda \|Z_i\|_{H^{-m_K}} \}] &\leq e^{\lambda B_{m_K} \Lambda_{\theta_0}} E \left[e^{\lambda B_{m_K} J_i} \right] \\ &= \exp \left\{ \lambda B_{m_K} \Lambda_{\theta_0} + \Lambda_{\theta_0} (e^{\lambda B_{m_K}} - 1) \right\} < \infty. \end{aligned}$$

Since $\lambda > 0$ is arbitrary, taking $\lambda = 1/A_T$ for any $A_T > 0$ shows that $Z_1, \dots, Z_n$ are i.i.d. exponentially integrable, centred H^{-m_K} -valued random elements, with

$$E_{\theta_0}^{(1)} \exp \{ \|Z_1\|_{H^{-m_K}} / A_T \} < \infty.$$

This is precisely Assumption 3.4, and concludes the verification of Assumptions 3.1–3.4, yielding the following posterior contraction theorem.

Theorem 4.2. *Let $\theta_0 \in H^\beta([0,1]^d)$, and let Π be an α -regular Gaussian series prior as in (20), for some $\alpha > 5d/2$ with $\alpha \leq \beta$. Assume data $N^{(n)} = (N_1, \dots, N_n)$, where $N_1, \dots, N_n$ are i.i.d. copies of an inhomogeneous Poisson process on $[0,1]^d$ with intensity ρ_{θ_0} as in (35). Then, for every $0 \leq s < \alpha \wedge \beta$ and every $M_n \rightarrow \infty$,*

$$E_{\theta_0}^{(n)} \left[\Pi \left(\theta \in \Theta : \|\theta - \theta_0\|_{H^s} > M_n n^{-\frac{(\alpha \wedge \beta) - s}{2\alpha + d}} \mid N^{(n)} \right) \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

The remarks following Theorem 4.1 apply to the present model as well. Taking $s \in \mathbb{N}$ with $s < \alpha \wedge \beta$ and using again the equivalence $\|\cdot\|_{H^s} \asymp \|\cdot\|_{W^{s,2}}$, Theorem 4.2 yields, for smoothness-matching priors, optimal recovery of the partial derivatives of the natural parameter θ of order at most s . The transfer to the intensity function is immediate: since, for any $t > d/2$, the exponential map is Lipschitz from bounded subsets of $H^t([0,1]^d)$ into $H^t([0,1]^d)$, it follows that the induced posterior on the the partial derivatives of ρ_θ of order at most s contract around those of the true intensity function ρ_{θ_0} at the same rate $n^{-((\alpha \wedge \beta) - s)/(2\alpha + d)}$, for every integer s with $d/2 < s < \alpha \wedge \beta$. For $\alpha = \beta$, this yields, to the best of our knowledge, the first optimal posterior contraction rates for estimating the derivatives of the intensity function of a Poisson process.

4.3 The white noise model

We finally consider the Gaussian white-noise model introduced in Example 2.5. Given the wavelet basis $\{\psi_k\}_{k=1}^\infty$ of Example 2.2, the observation scheme is equivalent to the Gaussian sequence model

$$Y_k^{(n)} = \theta_{0,k} + n^{-1/2} Z_k, \quad \theta_{0,k} := \langle \theta_0, \psi_k \rangle_{L^2}, \quad Z_k \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1). \quad (36)$$

Here, both the likelihood and the prior factorise over the coordinates of the Hilbert scale. The posterior can therefore be computed explicitly and also factorises, allowing us to bound directly the deterministic and stochastic terms in (16), as we now illustrate.

Let Π be the α -regular Gaussian series prior (20), with $\alpha > 0$. For $\theta \sim \Pi$, writing $\theta_k = \langle \theta, \psi_k \rangle_{L^2}$, we have $\theta_k \stackrel{\text{ind}}{\sim} \mathcal{N}(0, \lambda_k^{-2\alpha-d})$, $k \geq 1$. We then have

$$\Pi(\cdot \mid Y^{(n)}) = \bigotimes_{k=1}^{\infty} \pi_{n,k}(\cdot \mid Y_k^{(n)}), \quad (37)$$

where the conditional distribution of the k th coordinate is given by

$$d\pi_{n,k}(\vartheta \mid y) \propto \exp \left\{ -\frac{n}{2}(\vartheta - y)^2 - \frac{\lambda_k^{2\alpha+d}}{2}\vartheta^2 \right\} d\vartheta, \quad \vartheta, y \in \mathbb{R},$$

whereby a standard conjugate computation gives

$$\pi_{n,k}(\vartheta \mid y) = N \left(\frac{ny}{n + \lambda_k^{2\alpha+d}}, \frac{1}{n + \lambda_k^{2\alpha+d}} \right). \quad (38)$$

Fix any $0 \leq s < \alpha \wedge \beta$. We proceed bounding directly the deterministic and stochastic term arising from the basic decomposition (16), with the choice $T_{\theta_0} = \theta_0$. The deterministic term equals, in view of the factorisation (37) and the series representation of the H^s -norm,

$$\begin{aligned} D_{n,s}^2 &:= \left[\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\Pi_n^*(\cdot \mid \theta_0), \delta_{\theta_0}) \right]^2 \\ &= \int_{H^{\alpha \vee \beta}} \|\theta - \theta_0\|_{H^s}^2 d\Pi(\theta \mid Y^{(n)}) \\ &= \sum_{k=1}^{\infty} \lambda_k^{2s} \frac{\int_{\mathbb{R}} (\vartheta - \theta_{0,k})^2 e^{-\frac{n}{2}(\vartheta - \theta_{0,k})^2 - \frac{\lambda_k^{2\alpha+d}}{2}\vartheta^2} d\vartheta}{\int_{\mathbb{R}} e^{-\frac{n}{2}(\vartheta - \theta_{0,k})^2 - \frac{\lambda_k^{2\alpha+d}}{2}\vartheta^2} d\vartheta}. \end{aligned}$$

Using the conjugate property (38), each ratio in the above series can be evaluated analytically. In particular, separating the bias and variance components, we obtain

$$D_{n,s}^2 = \sum_{k=1}^{\infty} \frac{\lambda_k^{2s}}{n + \lambda_k^{2\alpha+d}} + \sum_{k=1}^{\infty} \lambda_k^{2s} \theta_{0,k}^2 \left(\frac{\lambda_k^{2\alpha+d}}{n + \lambda_k^{2\alpha+d}} \right)^2.$$

Call $V_{n,s}$ and $B_{n,s}$ the first and second series in the last display, respectively. We upper bound the former. Let K_n be determined by $\lambda_{K_n}^{2\alpha+d} \lesssim n \lesssim \lambda_{K_n+1}^{2\alpha+d}$. By the eigenvalue asymptotics (3), $K_n \asymp n^{d/(2\alpha+d)}$. Splitting the series defining $V_{n,s}$ then gives

$$V_{n,s} \lesssim \frac{1}{n} \sum_{k \leq K_n} \lambda_k^{2s} + \sum_{k > K_n} \lambda_k^{-2(\alpha-s)-d} \lesssim n^{-\frac{2(\alpha-s)}{2\alpha+d}}.$$

For the bias term, since $\theta_0 \in H^\beta([0, 1]^d) \subseteq H^{\alpha \wedge \beta}([0, 1]^d)$,

$$B_{n,s} \leq \|\theta_0\|_{H^{\alpha \wedge \beta}}^2 \sup_{k \geq 1} \left\{ \lambda_k^{2(s-\alpha \wedge \beta)} \left(\frac{\lambda_k^{2\alpha+d}}{n + \lambda_k^{2\alpha+d}} \right)^2 \right\}.$$

Setting $\rho := \frac{\alpha \wedge \beta - s}{2\alpha+d} \in (0, 1)$, we have

$$\lambda_k^{s-\alpha \wedge \beta} \frac{\lambda_k^{2\alpha+d}}{n + \lambda_k^{2\alpha+d}} = n^{-\rho} \frac{(\lambda_k^{2\alpha+d}/n)^{1-\rho}}{1 + \lambda_k^{2\alpha+d}/n} \lesssim n^{-\rho},$$

whence

$$B_{n,s} \lesssim \|\theta_0\|_{H^{\alpha \wedge \beta}}^2 n^{-\frac{2(\alpha \wedge \beta - s)}{2\alpha+d}}.$$

Combining the obtained bounds for the variance and bias components shows that $D_{n,s} \lesssim n^{-\frac{\alpha \wedge \beta - s}{2\alpha+d}}$.

Moving to the stochastic term, note that the marginal posteriors $\pi_{n,k}(\cdot \mid Y_k^{(n)})$ and $\pi_{n,k}(\cdot \mid \theta_{0,k})$ have the same variance. Therefore, their one-dimensional Wasserstein distance equals the absolute difference between their means:

$$\mathcal{W}_2^{\mathcal{P}_2(\mathbb{R})} \left(\pi_{n,k}(\cdot \mid Y_k^{(n)}), \pi_{n,k}(\cdot \mid \theta_{0,k}) \right)^2 = \left(\frac{n}{n + \lambda_k^{2\alpha+d}} \right)^2 (Y_k^{(n)} - \theta_{0,k})^2.$$

The coordinatewise translation coupling is optimal for the product Gaussian measures, and therefore

$$\begin{aligned} S_{n,s}^2 &:= E_{\theta_0}^{(n)} \left[\mathcal{W}_2^{\mathcal{P}_2(H_s)} \left(\Pi(\cdot \mid Y^{(n)}), \Pi_n^*(\cdot \mid \theta_0) \right)^2 \right] \\ &= \sum_{k=1}^{\infty} \mathcal{W}_2^{\mathcal{P}_2(\mathbb{R})} \left(\pi_{n,k}(\cdot \mid Y_k^{(n)}), \pi_{n,k}(\cdot \mid \theta_{0,k}) \right)^2 \\ &= \sum_{k=1}^{\infty} \lambda_k^{2s} \left(\frac{n}{n + \lambda_k^{2\alpha+d}} \right)^2 E_{\theta_0}^{(n)} \left[(Y_k^{(n)} - \theta_{0,k})^2 \right] = n \sum_{k=1}^{\infty} \frac{\lambda_k^{2s}}{(n + \lambda_k^{2\alpha+d})^2}, \end{aligned}$$

where we used $E_{\theta_0}^{(n)}[(Y_k^{(n)} - \theta_{0,k})^2] = n^{-1}$. Splitting again at $K_n \asymp n^{d/(2\alpha+d)}$ gives

$$S_{n,s}^2 \lesssim \frac{1}{n} \sum_{k \leq K_n} \lambda_k^{2s} + n \sum_{k > K_n} \lambda_k^{2s-4\alpha-2d} \lesssim n^{-\frac{2(\alpha-s)}{2\alpha+d}}.$$

Thus, $S_{n,s} \lesssim n^{-\frac{\alpha-s}{2\alpha+d}}$, and by Jensen's inequality

$$E_{\theta_0}^{(n)} \left[\mathcal{W}_2^{\mathcal{P}_2(H_s)} \left(\Pi(\cdot | Y^{(n)}), \Pi_n^*(\cdot | \theta_0) \right) \right] \lesssim n^{-\frac{\alpha-s}{2\alpha+d}}.$$

Combined with the obtained bound for the deterministic term, this completes the proof of the following.

Theorem 4.3. *Let $\theta_0 \in H^\beta([0,1]^d)$ for some $\beta > 0$, and let Π be the α -regular Gaussian series prior (20), for some $\alpha > 0$. Assume that $Y^{(n)}$ arising from the Gaussian white-noise model (36). Then, for every $0 \leq s < \alpha \wedge \beta$ and every $M_n \rightarrow \infty$,*

$$E_{\theta_0}^{(n)} \left[\Pi \left(\theta : \|\theta - \theta_0\|_{H^s} > M_n n^{-\frac{(\alpha \wedge \beta) - s}{2\alpha+d}} \mid Y^{(n)} \right) \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

5 Proofs

Proof of Theorem 3.5. We follow the argument laid out in Section 3.1, and denote by $\nu_\tau := \Pi_n^*(\cdot | \tau)$, $\tau \in \Theta^*$. The kernel ν_τ is well defined by (10), the mean element T_{θ_0} exists by virtue of Assumption 3.1, and the expected Wasserstein distance ε_n in (14) is finite under the requirements stated at the beginning of Section 3.1. The first summand in the decomposition (16) is deterministic and is shown in Proposition 5.1 to be of order $n^{-\frac{(\alpha \wedge \beta) - s}{2\alpha+d}}$ as $n \rightarrow \infty$. The second summand is the expectation of a stochastic quantity, and is shown in Proposition 5.3 to be of the same or smaller order. Therefore

$$\varepsilon_n = O \left(n^{-\frac{(\alpha \wedge \beta) - s}{2\alpha+d}} \right), \quad \text{as } n \rightarrow \infty.$$

As recalled after (14), the convergence (11) holds with this choice of ε_n for every $M_n \rightarrow \infty$; since it is preserved upon replacing ε_n by any sequence of the same or larger order, the claim follows. $\square$

5.1 Bounds for the deterministic term

We start by deriving an upper bound for the deterministic term in (16). Recalling (19), this is given by the ratio of two Laplace integrals. We apply an infinite-dimensional extension to the classical Laplace method for integral approximation (Wong, 2001), building on ideas developed by Dolera et al. (2024b).

Proposition 5.1. *Under the assumptions of Theorem 3.5, for every $0 \leq s < \alpha \wedge \beta$, as $n \rightarrow \infty$,*

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{T_{\theta_0}}, \delta_{\theta_0}) = O \left(n^{-\frac{\alpha \wedge \beta - s}{2\alpha+d}} \right). \quad (39)$$

Proof. By (19),

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{T_{\theta_0}}, \delta_{\theta_0})^2 = \frac{\int_{H_{s \vee p}} \|\theta - \theta_0\|_{H^s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{\int_{H_{s \vee p}} e^{-nK(\theta|\theta_0)} d\Pi(\theta)}. \quad (40)$$

We use a localisation argument to compare this ratio with the one arising for the probability measure $\mu_{I,\theta_0}^{(n)}$ from Lemma 5.2, whose normalising constant is denoted by $Z_{I,\theta_0}^{(n)}$. In view of (29) and (30), we may fix γ such that

$$s \vee p < \gamma < \alpha \wedge \beta, \quad q_K \gamma > (2 + q_K)p + 2r_K, \quad (2 + q_K)(\gamma - m_K) > 2\alpha + d.$$

We then pick $a \in (1/(2 + q_K), (\gamma - m_K)/(2\alpha + d))$ and set $\eta_n := n^{-a}$. With this choice,

$$n\eta_n^{2+q_K} \rightarrow 0, \quad \eta_n^{-2} n^{-\frac{2((\alpha \wedge \beta) - m_K)}{2\alpha + d}} \rightarrow 0. \quad (41)$$

Next fix $R < R_0$ sufficiently small that $B_R(\theta_0) \subseteq B_{R_K}^{H_{m_K}}(\theta_0)$, for $R_0, R_K > 0$ the radii appearing in Assumptions 3.1 and 3.2 respectively. Integrating the local Fisher lower bound (25) along the segment joining θ_0 and θ gives

$$K(\theta|\theta_0) \geq \frac{c_0}{2} \|\theta - \theta_0\|_{H_0}^2, \quad \theta \in B_R(\theta_0). \quad (42)$$

Let $\mu_{c_0,\theta_0}^{(n)}$ be the probability measure obtained from (50) below by replacing $I(\theta_0)[\theta - \theta_0]$ with $c_0\|\theta - \theta_0\|_{H_0}^2$, and let $Z_{c_0,\theta_0}^{(n)}$ denote its normalising constant. Equivalently, this corresponds to taking $c = c_0/2$ in the final assertion of Lemma 5.2. Hence, the conclusions of that lemma also apply to $\mu_{c_0,\theta_0}^{(n)}$.

Note that, since Π is Gaussian, $\mu_{c_0,\theta_0}^{(n)}$ also is Gaussian. Denote by $\bar{\theta}_{c_0}^{(n)}$ its mean. The covariance operator of $\mu_{c_0,\theta_0}^{(n)}$ is given by $\Sigma_{c_0}^{(n)} = (nc_0I + Q_\Pi^{-1})^{-1}$, with I the identity operator. Write

$$\sigma_{n,m_K}^2 := \|L^{m_K} \Sigma_{c_0}^{(n)} L^{m_K}\|_{\text{op}} = \sup_{k \geq 1} \frac{\lambda_k^{2m_K}}{nc_0 + \lambda_k^{2\alpha+d}} \lesssim n^{-1 + \frac{2m_K}{2\alpha+d}}.$$

for the maximal variance of the centred Gaussian measure in the H_{m_K} -geometry, the last inequality following from the eigenvalue asymptotics (3). Lemma 5.2, applied with $t = m_K$, gives

$$E_{\mu_{c_0,\theta_0}^{(n)}} \left[\|\theta - \bar{\theta}_{c_0}^{(n)}\|_{H_{m_K}}^2 \right] + \|\bar{\theta}_{c_0}^{(n)} - \theta_0\|_{H_{m_K}}^2 = E_{\mu_{c_0,\theta_0}^{(n)}} \left[\|\theta - \theta_0\|_{H_{m_K}}^2 \right] \lesssim n^{-\frac{2((\alpha \wedge \beta) - m_K)}{2\alpha+d}},$$

so that by Jensen's inequality,

$$E_{\mu_{c_0,\theta_0}^{(n)}} \left[\|\theta - \bar{\theta}_{c_0}^{(n)}\|_{H_{m_K}} \right] \lesssim n^{-\frac{(\alpha \wedge \beta) - m_K}{2\alpha+d}}, \quad (43)$$

and

$$\|\bar{\theta}_{c_0}^{(n)} - \theta_0\|_{H_{m_K}} \lesssim n^{-\frac{(\alpha \wedge \beta) - m_K}{2\alpha+d}}. \quad (44)$$

Recall that $\eta_n = n^{-a}$ and that

$$a < \frac{\gamma - m_K}{2\alpha + d} < \frac{(\alpha \wedge \beta) - m_K}{2\alpha + d} \leq \frac{\alpha - m_K}{2\alpha + d}.$$

It follows that both the Gaussian fluctuation in (43) scale and the bias (44) are of smaller order than η_n . In particular, for all sufficiently large n ,

$$\eta_n - \|\bar{\theta}_{c_0}^{(n)} - \theta_0\|_{H_{m_K}} - E_{\mu_{c_0, \theta_0}^{(n)}} \left[\|\theta - \bar{\theta}_{c_0}^{(n)}\|_{H_{m_K}} \right] \geq \frac{\eta_n}{2}.$$

The Gaussian isoperimetric inequality (Giné and Nickl, 2016, Theorem 2.6.12) applied to the H_{m_K} -valued Gaussian random element $\theta - \bar{\theta}_{c_0}^{(n)}$, $\theta \sim \mu_{c_0, \theta_0}^{(n)}$ therefore gives, for some $c_1, c_2 > 0$,

$$\mu_{c_0, \theta_0}^{(n)} \left(\|\theta - \theta_0\|_{H_{m_K}} > \eta_n \right) \leq \exp \left\{ -\frac{c_1 \eta_n^2}{\sigma_{n, m_K}^2} \right\} \leq \exp \left\{ -c_2 n^{1 - \frac{2m_K}{2\alpha + d} - 2a} \right\}.$$

Then, by (42), the definition of $\mu_{c_0, \theta_0}^{(n)}$, and the Cauchy–Schwarz inequality,

$$\begin{aligned} & \int_{B_R(\theta_0) \cap B_{\eta_n}^{H_{m_K}}(\theta_0)^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) \\ & \leq Z_{c_0, \theta_0}^{(n)} \left[\mu_{c_0, \theta_0}^{(n)} \left(B_{\eta_n}^{H_{m_K}}(\theta_0)^c \right) + \left\{ \int \|\theta - \theta_0\|_{H_s}^4 d\mu_{c_0, \theta_0}^{(n)}(\theta) \right\}^{1/2} \mu_{c_0, \theta_0}^{(n)} \left(B_{\eta_n}^{H_{m_K}}(\theta_0)^c \right)^{1/2} \right] \\ & \lesssim Z_{c_0, \theta_0}^{(n)} \exp \left\{ -\frac{c_2}{2} n^{1 - \frac{2m_K}{2\alpha + d} - 2a} \right\}, \end{aligned}$$

where the last inequality follows from the preceding Gaussian concentration bound and the fourth moment estimate in Lemma 5.2 below, applied with $t = s$. Moreover, since $Z_{c_0, \theta_0}^{(n)} \leq 1$, the lower bound (51) gives

$$\frac{Z_{c_0, \theta_0}^{(n)}}{Z_{I, \theta_0}^{(n)}} \leq \exp \left\{ C n^{1 - \frac{2(\alpha \wedge \beta)}{2\alpha + d}} \right\}.$$

Since

$$1 - \frac{2m_K}{2\alpha + d} - 2a > 1 - \frac{2(\alpha \wedge \beta)}{2\alpha + d},$$

we may choose

$$1 - \frac{2(\alpha \wedge \beta)}{2\alpha + d} < \delta < 1 - \frac{2m_K}{2\alpha + d} - 2a,$$

to obtain, for some $c_3 > 0$ and all sufficiently large n ,

$$\int_{B_R(\theta_0) \cap B_{\eta_n}^{H_{m_K}}(\theta_0)^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) \leq e^{-c_3 n^\delta} Z_{I, \theta_0}^{(n)}. \quad (45)$$

There remains to bound the integral over $(B_R(\theta_0))^c$. By the interpolation inequality (1), for $h \in H_\gamma$, with γ chosen at the beginning of the proof,

$$\|h\|_{H_{-r_K}} \geq \|h\|_{H_p}^{\frac{\gamma + r_K}{\gamma - p}} \|h\|_{H_\gamma}^{-\frac{p + r_K}{\gamma - p}}. \quad (46)$$

Taking γ sufficiently close to $\alpha \wedge \beta$, the two strict regularity requirements in Theorem 3.5 allow us to choose

$$\frac{1}{2} \left(1 - \frac{2(\alpha \wedge \beta)}{2\alpha + d} \right) < b < \frac{(\alpha \wedge \beta)(\gamma - p)}{(2\alpha + d)(p + r_K)}.$$

We proceed splitting $(B_R(\theta_0))^c = \mathcal{B}_{n,1} \cup \mathcal{B}_{n,2}$ with $\mathcal{B}_{n,1} := \{\theta : \|\theta - \theta_0\|_{H_p} > R, \|\theta - \theta_0\|_{H_\gamma} \leq n^b\}$, and $\mathcal{B}_{n,2} := \{\theta : \|\theta - \theta_0\|_{H_p} > R, \|\theta - \theta_0\|_{H_\gamma} > n^b\}$. For every $\theta \in \mathcal{B}_{n,1}$, the interpolation inequality (46) gives,

$$\|\theta - \theta_0\|_{H_{-rK}} \geq \|\theta - \theta_0\|_{H_p}^{\frac{\gamma+rK}{\gamma-p}} \|\theta - \theta_0\|_{H_\gamma}^{-\frac{p+rK}{\gamma-p}} \geq R^{\frac{\gamma+rK}{\gamma-p}} n^{-\frac{b(p+rK)}{\gamma-p}}.$$

Combining this with the global KL lower bound (27) yields $K(\theta \mid \theta_0) \gtrsim n^{-\frac{2b(p+rK)}{\gamma-p}}$ for all $\theta \in \mathcal{B}_{n,1}$. Moreover, since $s < \gamma$, $\|\theta - \theta_0\|_{H_s} \lesssim \|\theta - \theta_0\|_{H_\gamma} \leq n^b$ for all $\theta \in \mathcal{B}_{n,1}$. It follows that

$$\begin{aligned} \int_{\mathcal{B}_{n,1}} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) &\lesssim (1 + n^{2b}) \exp \left\{ -c_3 n^{1 - \frac{2b(p+rK)}{\gamma-p}} \right\} \\ &\lesssim \exp \left\{ -c_4 n^{1 - \frac{2b(p+rK)}{\gamma-p}} \right\}, \end{aligned}$$

for constants $c_4, c_5 > 0$. For the integral over $\mathcal{B}_{n,2}$, we instead use the trivial upper bound $e^{-nK(\theta|\theta_0)} \leq 1$. By Cauchy–Schwarz inequality,

$$\begin{aligned} \int_{\mathcal{B}_{n,2}} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) \\ \leq \Pi \left(\|\theta - \theta_0\|_{H_\gamma} > n^b \right) + \left\{ \int \|\theta - \theta_0\|_{H_s}^4 d\Pi(\theta) \right\}^{1/2} \Pi \left(\|\theta - \theta_0\|_{H_\gamma} > n^b \right)^{1/2}. \end{aligned}$$

Since $\gamma < \alpha$, the prior Π can be regarded as centred Gaussian Borel probability measure on H_γ . Furthermore, $\theta_0 \in H_\gamma$, and Fernique’s theorem (Bogachev, 1998, Theorem 2.8.5) gives

$$\Pi \left(\theta : \|\theta - \theta_0\|_{H_\gamma} > n^b \right) \lesssim e^{-c_6 n^{2b}}.$$

for some $c_6 > 0$. The prior has a finite fourth moment in H_s , so the preceding display also implies

$$\int_{\mathcal{B}_{n,2}} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) \lesssim e^{-c_7 n^{2b}},$$

for some $c_7 > 0$. By the choice of b , we may find $\delta' > 0$ such that

$$1 - \frac{2(\alpha \wedge \beta)}{2\alpha + d} < \delta' < \min \left\{ 2b, 1 - \frac{2b(p+rK)}{\gamma-p} \right\}.$$

Thus, adding the two obtained bounds for the integral over $(B_R(\theta_0))^c$ gives, for some $c_8 > 0$ and all sufficiently large n ,

$$\int_{(B_R(\theta_0))^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) \leq e^{-c_8 n^{\delta'}}. \quad (47)$$

By (45), for every fixed $C > 0$,

$$\frac{\int_{B_R(\theta_0) \cap B_{\eta_n}^{H_m K}(\theta_0)^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{n^{-C} Z_{I, \theta_0}^{(n)}} \leq n^C e^{-c_3 n^\delta} \rightarrow 0.$$

Moreover, (47), the lower bound (51) and the choice of δ' imply

$$\frac{\int_{(B_R(\theta_0))^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{n^{-C} Z_{I,\theta_0}^{(n)}} \leq n^N \exp \left\{ -cn^{\delta'} + Cn^{1-\frac{2(\alpha \wedge \beta)}{2\alpha+d}} \right\} \rightarrow 0.$$

Conclude that, for every fixed $C > 0$,

$$\int_{B_R(\theta_0) \cap B_{\eta_n}^{H_{m_K}}(\theta_0)^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) = o \left(n^{-C} Z_{I,\theta_0}^{(n)} \right)$$

and

$$\int_{B_R(\theta_0)^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) = o \left(n^{-C} Z_{I,\theta_0}^{(n)} \right).$$

Combining the above shows

$$\int_{H_{s \vee p} \cap B_{\eta_n}^{H_{m_K}}(\theta_0)^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) = o \left(n^{-C} Z_{I,\theta_0}^{(n)} \right). \quad (48)$$

We next compare the two exponential weights inside $B_{\eta_n}^{H_{m_K}}(\theta_0)$. By (52), applied with $t = m_K$, and (41),

$$\mu_{I,\theta_0}^{(n)} \left(B_{\eta_n}^{H_{m_K}}(\theta_0)^c \right) \leq \eta_n^{-2} \int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_{m_K}}^2 d\mu_{I,\theta_0}^{(n)}(\theta) \lesssim \eta_n^{-2} n^{-\frac{2((\alpha \wedge \beta) - m_K)}{2\alpha+d}} = o(1).$$

Set

$$\zeta_n := \sup_{\theta \in H_{s \vee p} \cap B_{\eta_n}^{H_{m_K}}(\theta_0)} n \left| K(\theta | \theta_0) - \frac{1}{2} I(\theta_0)[\theta - \theta_0] \right|.$$

By Assumption 3.2 and the choice of η_n , $\zeta_n \lesssim n\eta_n^{2+q_K} = o(1)$. Therefore, for every $\theta \in H_{s \vee p} \cap B_{\eta_n}^{H_{m_K}}(\theta_0)$,

$$e^{-\zeta_n} \exp \left\{ -\frac{n}{2} I(\theta_0)[\theta - \theta_0] \right\} \leq e^{-nK(\theta|\theta_0)} \leq e^{\zeta_n} \exp \left\{ -\frac{n}{2} I(\theta_0)[\theta - \theta_0] \right\}.$$

It follows that

$$\int_{H_{s \vee p} \cap B_{\eta_n}^{H_{m_K}}(\theta_0)} e^{-nK(\theta|\theta_0)} d\Pi(\theta) \geq e^{-\zeta_n} Z_{I,\theta_0}^{(n)} \mu_{I,\theta_0}^{(n)} \left(B_{\eta_n}^{H_{m_K}}(\theta_0) \right) = (1 + o(1)) Z_{I,\theta_0}^{(n)}. \quad (49)$$

Similarly,

$$\int_{H_{s \vee p} \cap B_{\eta_n}^{H_{m_K}}(\theta_0)} \|\theta - \theta_0\|_{H_s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta) \leq e^{\zeta_n} Z_{I,\theta_0}^{(n)} \int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 d\mu_{I,\theta_0}^{(n)}(\theta).$$

Consequently,

$$\begin{aligned} \frac{\int_{H_{s \vee p} \cap B_{\eta_n}^{H_{mK}}(\theta_0)} \|\theta - \theta_0\|_{H_s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{\int_{H_{s \vee p} \cap B_{\eta_n}^{H_{mK}}(\theta_0)} e^{-nK(\theta|\theta_0)} d\Pi(\theta)} &\leq \frac{e^{2\zeta_n}}{\mu_{I, \theta_0}^{(n)}(B_{\eta_n}^{H_{mK}}(\theta_0))} \int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 d\mu_{I, \theta_0}^{(n)}(\theta) \\ &= (1 + o(1)) \int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 d\mu_{I, \theta_0}^{(n)}(\theta). \end{aligned}$$

In view of (48) and the preceding lower bound for the local denominator, an elementary decomposition of the numerator and denominator over $B_{\eta_n}^{H_{mK}}(\theta_0)$ and its complement gives, for every fixed $C > 0$,

$$\frac{\int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{\int_{H_{s \vee p}} e^{-nK(\theta|\theta_0)} d\Pi(\theta)} = \frac{\int_{H_{s \vee p} \cap B_{\eta_n}^{H_{mK}}(\theta_0)} \|\theta - \theta_0\|_{H_s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{\int_{H_{s \vee p} \cap B_{\eta_n}^{H_{mK}}(\theta_0)} e^{-nK(\theta|\theta_0)} d\Pi(\theta)} + o(n^{-C}).$$

Hence,

$$\frac{\int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{\int_{H_{s \vee p}} e^{-nK(\theta|\theta_0)} d\Pi(\theta)} \leq (1 + o(1)) \int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 d\mu_{I, \theta_0}^{(n)}(\theta) + o(n^{-C}).$$

Choose $C > \frac{2((\alpha \wedge \beta) - s)}{2\alpha + d}$. Lemma 5.2, applied with $t = s$, then yields

$$\frac{\int_{H_{s \vee p}} \|\theta - \theta_0\|_{H_s}^2 e^{-nK(\theta|\theta_0)} d\Pi(\theta)}{\int_{H_{s \vee p}} e^{-nK(\theta|\theta_0)} d\Pi(\theta)} \lesssim n^{-\frac{2((\alpha \wedge \beta) - s)}{2\alpha + d}}.$$

Taking square roots in (40) proves (39). □

Lemma 5.2. *Under the assumptions of Proposition 5.1, consider the probability measure on Θ defined by, for $\theta \in \Theta$,*

$$d\mu_{I, \theta_0}^{(n)}(\theta) := \frac{\exp\left\{-\frac{n}{2}I(\theta_0)[\theta - \theta_0]\right\}}{Z_{I, \theta_0}^{(n)}} d\Pi(\theta), \quad Z_{I, \theta_0}^{(n)} := \int_{H_{s \vee p}} \exp\left\{-\frac{n}{2}I(\theta_0)[\theta - \theta_0]\right\} d\Pi(\theta). \quad (50)$$

Then, there exists a constant $C_{I, \theta_0} > 0$ such that, for all n large enough,

$$Z_{I, \theta_0}^{(n)} \geq \exp\left\{-C_{I, \theta_0} n^{1 - \frac{2(\alpha \wedge \beta)}{2\alpha + d}}\right\}. \quad (51)$$

Furthermore, for every $0 \leq t < \alpha \wedge \beta$ and all n large enough,

$$\int \|\theta - \theta_0\|_{H_t}^2 d\mu_{I,\theta_0}^{(n)}(\theta) \lesssim n^{-\frac{2((\alpha \wedge \beta) - t)}{2\alpha + d}}; \quad \int \|\theta - \theta_0\|_{H_t}^4 d\mu_{I,\theta_0}^{(n)}(\theta) \lesssim n^{-\frac{4((\alpha \wedge \beta) - t)}{2\alpha + d}}. \quad (52)$$

These conclusions remain valid, with possibly different constants, if $I(\theta_0)[h]/2$ in (50) is replaced by $c\|h\|_{H_0}^2$, for any fixed $c > 0$.

Proof. The estimates follow by first considering finite-dimensional Galerkin spaces. For $K \in \mathbb{N}$, let $\Theta_K := \text{span}\{e_1, \dots, e_K\}$, with canonical projection $P_K : \Theta \rightarrow \Theta_K$. Set $\theta_{0,K} := P_K \theta_0$ and $\Pi_K := (P_K)_\# \Pi$, and define the probability measure $\mu_{I,\theta_0,K}^{(n)}$ on Θ_K by

$$d\mu_{I,\theta_0,K}^{(n)}(\theta) := \frac{\exp\left\{-\frac{n}{2}I(\theta_0)[\theta - \theta_{0,K}]\right\}}{Z_{I,\theta_0,K}^{(n)}} d\Pi_K(\theta),$$

with

$$Z_{I,\theta_0,K}^{(n)} := \int_{\Theta_K} \exp\left\{-\frac{n}{2}I(\theta_0)[\theta - \theta_{0,K}]\right\} d\Pi_K(\theta).$$

Identifying Θ_K with $\mathbb{R}^K$, write $I_K(\theta_0)$, $Q_{\Pi,K}$ and L_K for the restrictions to Θ_K of the Fisher information operator, the prior covariance operator Q_Π and the scale generator L from (2). Under the identification of Θ_K with $\mathbb{R}^K$, we have $L_K = \text{diag}[\{\lambda_k\}_{k=1}^K]$ and, in view of the identity $Q_\Pi = L^{-2\alpha-d}$, $Q_{\Pi,K} = \text{diag}[\{\lambda_k^{-2\alpha-d}\}_{k=1}^K]$. A standard Gaussian conjugate computation then shows that $\mu_{I,\theta_0,K}^{(n)} = \mathcal{N}_K(\theta_{0,K} + b_{n,K}, \Sigma_{n,K})$, where

$$b_{n,K} = -(nI_K(\theta_0) + Q_{\Pi,K}^{-1})^{-1} Q_{\Pi,K}^{-1} \theta_{0,K}, \quad \Sigma_{n,K} = (nI_K(\theta_0) + Q_{\Pi,K}^{-1})^{-1}.$$

By Assumption 3.1, the normalising constant satisfies

$$Z_{I,\theta_0,K}^{(n)} \geq \int_{\Theta_K} \exp\left\{-\frac{nC_0}{2}\|\theta - \theta_{0,K}\|_{H_0}^2\right\} d\Pi_K(\theta).$$

The integral on the right hand side can be evaluated analytically by another Gaussian conjugate computation (now with diagonal covariance matrices), leading to the upper bound

$$\begin{aligned} -\log Z_{I,\theta_0,K}^{(n)} &\lesssim \sum_{k=1}^K \log(1 + n\lambda_k^{-(2\alpha+d)}) + \sum_{k=1}^K \frac{n\lambda_k^{2\alpha+d}}{n + \lambda_k^{2\alpha+d}} \theta_{0,k}^2 \\ &\lesssim n^{\frac{d}{2\alpha+d}} + n^{\frac{(2\alpha+d-2\beta)+}{2\alpha+d}} \|\theta_0\|_{H_\beta}^2 \lesssim n^{1-\frac{2(\alpha \wedge \beta)}{2\alpha+d}}, \end{aligned}$$

holding uniformly in K , where the second to last inequality follows from the eigenvalue asymptotics (3).

Moving to the second moment, we have

$$\begin{aligned} \int_{\Theta_K} \|\theta - \theta_{0,K}\|_{H_t}^2 d\mu_{I,\theta_0,K}^{(n)}(\theta) &= E_{\mu_{I,\theta_0,K}^{(n)}} [\|\theta - (\theta_{0,K} + b_{n,K})\|_{H_t}^2] + \|b_{n,K}\|_{H_t}^2 \\ &= E_{\mu_{I,\theta_0,K}^{(n)}} [\|L_K^t[\theta - (\theta_{0,K} + b_{n,K})]\|_{H_0}^2] + \|b_{n,K}\|_{H_t}^2 \\ &= \text{Tr}[L_K^{2t} \Sigma_{n,K}] + \|b_{n,K}\|_{H_t}^2. \end{aligned}$$

We bound the two terms in the last line. By Assumption 3.1, and by the order-reversing property of inversion for positive-definite matrices (Horn and Johnson, 2012, Section 7.7),

$$\Sigma_{n,K} \preceq (nc_0 I_{\Theta_K} + Q_{\Pi,K}^{-1})^{-1} = (nc_0 I_{\Theta_K} + L_K^{2\alpha+d})^{-1},$$

where I_{Θ_K} is the identity matrix on Θ_K . We then obtain, for every $0 \leq t < \alpha$,

$$\text{Tr}[L_K^{2t} \Sigma_{n,K}] \lesssim \sum_{k=1}^K \frac{\lambda_k^{2t-(2\alpha+d)}}{1 + n\lambda_k^{-(2\alpha+d)}} \lesssim n^{-\frac{2(\alpha-t)}{2\alpha+d}}, \quad (53)$$

uniformly in K . To estimate the bias term, write $\bar{\theta}_{n,K} := \theta_{0,K} + b_{n,K}$ for the mean of $\mu_{I,\theta_0,K}^{(n)}$. By its variational characterisation, it is the unique minimiser over Θ_K of

$$u \in \Theta_K \mapsto nI(\theta_0)[u - \theta_{0,K}] + \|u\|_{H_{\alpha+d/2}}^2.$$

Let $K_n \in \mathbb{N}$ be the spectral cut-off determined by $\lambda_{K_n} \lesssim n^{1/(2\alpha+d)} \lesssim \lambda_{K_n+1}$. Since $\theta_0 \in H_{\alpha \wedge \beta}$, the above display and Assumption 3.1 give, uniformly in K ,

$$\begin{aligned} n\|\theta_{0,K} - \bar{\theta}_{n,K}\|_{H_0}^2 + \|\bar{\theta}_{n,K}\|_{H_{\alpha+d/2}}^2 &\lesssim n\|\theta_{0,K} - P_{K_n \wedge K} \theta_0\|_{H_0}^2 + \|P_{K_n \wedge K} \theta_0\|_{H_{\alpha+d/2}}^2 \\ &\lesssim n^{1-\frac{2(\alpha \wedge \beta)}{2\alpha+d}} \|\theta_0\|_{H_{\alpha \wedge \beta}}^2, \end{aligned}$$

the last inequality following standard projection estimates and the eigenvalue asymptotics (3). It follows that

$$\|\bar{\theta}_{n,K} - P_{K_n \wedge K} \theta_0\|_{H_0} \leq \|\bar{\theta}_{n,K} - \theta_{0,K}\|_{H_0} + \|P_{K_n \wedge K} \theta_0 - \theta_{0,K}\|_{H_0} \lesssim n^{-\frac{\alpha \wedge \beta}{2\alpha+d}} \|\theta_0\|_{H_{\alpha \wedge \beta}}$$

and, similarly,

$$\|\bar{\theta}_{n,K} - P_{K_n \wedge K} \theta_0\|_{H_{\alpha+d/2}} \lesssim n^{\frac{\alpha+d/2-(\alpha \wedge \beta)}{2\alpha+d}} \|\theta_0\|_{H_{\alpha \wedge \beta}}.$$

Interpolating between H_0 and $H_{\alpha+d/2}$ via the inequality (1),

$$\begin{aligned} \|\bar{\theta}_{n,K} - P_{K_n \wedge K} \theta_0\|_{H_t} &\lesssim \left(n^{-\frac{\alpha \wedge \beta}{2\alpha+d}}\right)^{1-\frac{t}{\alpha+d/2}} \left(n^{\frac{\alpha+d/2-(\alpha \wedge \beta)}{2\alpha+d}}\right)^{\frac{t}{\alpha+d/2}} \|\theta_0\|_{H_{\alpha \wedge \beta}} \\ &= n^{-\frac{(\alpha \wedge \beta)-t}{2\alpha+d}} \|\theta_0\|_{H_{\alpha \wedge \beta}}, \end{aligned}$$

and since also

$$\|\theta_{0,K} - P_{K_n \wedge K} \theta_0\|_{H_t} \lesssim n^{-\frac{(\alpha \wedge \beta)-t}{2\alpha+d}} \|\theta_0\|_{H_{\alpha \wedge \beta}},$$

we obtain from the triangle inequality, for all $0 \leq t < \alpha \wedge \beta$, uniformly in K ,

$$\|b_{n,K}\|_{H_t} = \|\bar{\theta}_{n,K} - \theta_{0,K}\|_{H_t} \lesssim n^{-\frac{(\alpha \wedge \beta)-t}{2\alpha+d}} \|\theta_0\|_{H_{\alpha \wedge \beta}}. \quad (54)$$

Combining (53) and (54) shows that

$$\int_{\Theta_K} \|\theta - \theta_{0,K}\|_{H_t}^2 d\mu_{I,\theta_0,K}^{(n)}(\theta) \lesssim n^{-\frac{2((\alpha \wedge \beta)-t)}{2\alpha+d}}.$$

Finally, the standard Gaussian fourth-moment identity gives

$$\begin{aligned} E_{\mu_{I,\theta_0,K}}^{(n)} [\|\theta - (\theta_{0,K} + b_{n,K})\|_{H_t}^4] &= \{\text{Tr}[L_K^{2t}\Sigma_{n,K}]\}^2 + 2\text{Tr}[(L_K^{2t}\Sigma_{n,K})^2] \\ &\leq 3\{\text{Tr}[L_K^{2t}\Sigma_{n,K}]\}^2 \lesssim n^{-\frac{4((\alpha\wedge\beta)-t)}{2\alpha+d}} \end{aligned}$$

where for the last inequality we have again applied (53). Thus, using (54) and the basic fact that $\|h + g\|_{H_t}^4 \leq 8\|h\|_{H_t}^4 + 8\|g\|_{H_t}^4$, for all $h, g \in H_t$ and all $t \in \mathbb{R}$,

$$\begin{aligned} \int_{\Theta_K} \|\theta - \theta_{0,K}\|_{H_t}^4 d\mu_{I,\theta_0,K}^{(n)}(\theta) &\leq 8E_{\mu_{I,\theta_0,K}}^{(n)} [\|\theta - (\theta_{0,K} + b_{n,K})\|_{H_t}^4] + 8\|b_{n,K}\|_{H_t}^4 \\ &\lesssim n^{-\frac{4((\alpha\wedge\beta)-t)}{2\alpha+d}}. \end{aligned}$$

To conclude, we pass to the limit as $K \rightarrow \infty$. Since $\Pi(H_{s\vee p}) = 1$, the normalising constant in (50) may equivalently be integrated over Θ . Moreover, by the definition of Π_K ,

$$Z_{I,\theta_0,K}^{(n)} = \int_{\Theta} \exp\left\{-\frac{n}{2}I(\theta_0)[P_K(\theta - \theta_0)]\right\} d\Pi(\theta).$$

Similarly, for $q \in \{2, 4\}$,

$$\int_{\Theta_K} \|\theta - \theta_{0,K}\|_{H_t}^q d\mu_{I,\theta_0,K}^{(n)}(\theta) = \frac{\int_{\Theta} \|P_K(\theta - \theta_0)\|_{H_t}^q \exp\left\{-\frac{n}{2}I(\theta_0)[P_K(\theta - \theta_0)]\right\} d\Pi(\theta)}{Z_{I,\theta_0,K}^{(n)}}.$$

For every $t < \alpha$, $P_K\theta \rightarrow \theta$ in H_t , Π -almost surely, while $P_K\theta_0 \rightarrow \theta_0$ in H_t for every $t < \beta$. Furthermore,

$$\|P_K(\theta - \theta_0)\|_{H_t}^q \leq \|\theta - \theta_0\|_{H_t}^q,$$

and the right-hand side is Π -integrable for $q \in \{2, 4\}$. Assumption 3.1 also implies that $I(\theta_0)$ is continuous on H_0 . Dominated convergence therefore gives

$$Z_{I,\theta_0,K}^{(n)} \rightarrow Z_{I,\theta_0}^{(n)}$$

and

$$\int_{\Theta_K} \|\theta - \theta_{0,K}\|_{H_t}^q d\mu_{I,\theta_0,K}^{(n)}(\theta) \longrightarrow \int_{\Theta} \|\theta - \theta_0\|_{H_t}^q d\mu_{I,\theta_0}^{(n)}(\theta).$$

Because all the preceding estimates are uniform in K , letting $K \rightarrow \infty$ proves (51) and (52). $\square$

5.2 Bounds for the stochastic term

We now bound the stochastic term in (16). Throughout this section, we write $\hat{D}_n := \hat{T}_n - T_{\theta_0}$ and $\ell_n(\theta) := \sum_{i=1}^n \log[f_{\theta}(X_i)/f_{\theta_0}(X_i)]$, $\theta \in \Theta$, for the centred sufficient statistic and log-likelihood, respectively. Note that by (5) and (18),

$$\ell_n(\theta) = n\hat{D}_n(\theta - \theta_0) - nK(\theta|\theta_0), \quad d\nu_{\hat{T}_n}(\theta) = \frac{e^{\ell_n(\theta)} d\Pi(\theta)}{\int_{H_{s\vee p}} e^{\ell_n(\theta)} d\Pi(\theta)}, \quad \theta \in \Theta. \quad (55)$$

Thus, the tilt of the posterior distribution $\nu_{\hat{T}_n}$ relative to the deterministic probability measure $\nu_{T_{\theta_0}}$ is given by the Θ^* -valued random element $n\hat{D}_n$, a structural feature of exponential families that we use repeatedly.

The following auxiliary definitions are necessary. For $R > 0$ to be fixed below, write shorthand $\mathcal{B} := B_R^\Theta(\theta_0) = \{\theta \in \Theta : \|\theta - \theta_0\|_\Theta \leq R\}$. Using that $\|h\|_{H_{m_K}} \leq \lambda_1^{m_K-p} \|h\|_{H_p} = \lambda_1^{m_K-p} \|h\|_\Theta$, holding since $m_K \leq p$, we may take sufficiently small $R < R_0$ such that

$$\mathcal{B} \subseteq B_{R_0}^\Theta(\theta_0) \cap B_{R_K}^{H_{m_K}}(\theta_0), \quad (56)$$

so that both the coercivity property (25) and the Taylor expansion (26) are available on $\mathcal{B}$, as is the local Kullback-Leibler lower bound (42) in view of the proof of Proposition 5.1. For any $\tau \in \Theta^*$, let

$$\tilde{\nu}_\tau(\cdot) := \nu_\tau(\cdot | \mathcal{B}) = \frac{\nu_\tau(\cdot \cap \mathcal{B})}{\nu_\tau(\mathcal{B})} \quad (57)$$

be the conditioning of ν_τ to $\mathcal{B}$, which is well defined since ν_τ and Π are equivalent and $\Pi(\mathcal{B}) > 0$.

Proposition 5.3. *Under the assumptions of Theorem 3.5, for every $0 \leq s < \alpha \wedge \beta$, as $n \rightarrow \infty$,*

$$E_{\theta_0}^{(n)} \left[\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}}) \right] = O \left(n^{-\frac{(\alpha \wedge \beta) - s}{2\alpha + d}} \right). \quad (58)$$

Proof. Let $\mathcal{E}_n$ be the event defined in Lemma 5.7 and split

$$E_{\theta_0}^{(n)} [\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}})] = E_{\theta_0}^{(n)} [\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}}) 1_{\mathcal{E}_n^c}] + E_{\theta_0}^{(n)} [\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}}) 1_{\mathcal{E}_n}]. \quad (59)$$

We upper bound the first term. Since M is convex, $D^2M(\theta) \succeq 0$ on Θ , so that the curvature of every ν_τ is bounded below globally by the prior precision Q_Π^{-1} . Applying the Bakry–Émery criterion on the finite-dimensional Galerkin spaces and passing to the limit as in the proof of Lemma 5.10 below (with $\Theta_{0,K}$ in place of $\mathcal{B} \cap \Theta_{0,K}$, which in fact simplifies the argument) gives, for all $\tau \in \Theta^*$,

$$\text{Var}_{\nu_\tau}[F(\theta)] \leq \bar{\kappa}_s \int_\Theta \|\nabla^{H_s} F\|_{H_s}^2 d\nu_\tau, \quad \text{Var}_{\nu_\tau}[F(\theta)] \leq \int_\Theta \|\nabla^{H_{\alpha+d/2}} F\|_{H_{\alpha+d/2}}^2 d\nu_\tau,$$

holding for all bounded smooth cylindrical functions of the form $F(\theta) = f(\theta_1, \dots, \theta_K)$ for some $K \in \mathbb{N}$ and some bounded and smooth $f : \mathbb{R}^K \rightarrow \mathbb{R}$ with bounded derivatives of all orders, and where, for $q \in \mathbb{R}$, $\nabla^{H_q} F = \sum_{k \leq K} \lambda_k^{-2q} \partial_k f e_k$ denotes the gradient of F in the H_q -geometry. Above, $\bar{\kappa}_s := \sup_{k \geq 1} \lambda_k^{2s-2\alpha-d} = \lambda_1^{2s-2\alpha-d} < \infty$ because $s < \alpha$.

Applying the transport techniques in the proof of Lemma 5.11 below with these two global inequalities in place of (81) yields the crude estimate

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_\tau, \nu_{\tau'}) \leq \bar{\kappa}_s^{1/2} n \|\tau - \tau'\|_{H_{-\alpha-d/2}}, \quad \text{for all } \tau, \tau' \in \Theta^*.$$

Then, $\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}})^2 \lesssim n^2 \|\hat{D}_n\|_{H_{-\alpha-d/2}}^2$ and since, by (23), (24) and Fubini's theorem,

$$E_{\theta_0}^{(n)} \|\hat{D}_n\|_{H_{-\alpha-d/2}}^2 = \frac{1}{n} \sum_{k=1}^{\infty} \lambda_k^{-2\alpha-d} I(\theta_0)[e_k] \leq \frac{C_0}{n} \sum_{k=1}^{\infty} \lambda_k^{-2\alpha-d} \lesssim \frac{1}{n},$$

the series converging in view of the eigenvalue asymptotics (3) for any $\alpha > 0$, we obtain $E_{\theta_0}^{(n)}[\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}})^2] \lesssim n$. By the Cauchy–Schwarz inequality and the exponential probability decay (70), this implies that

$$E_{\theta_0}^{(n)}[\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}})1_{\mathcal{E}_n^c}] \lesssim n^{1/2} \exp\{-\frac{c}{2}n^\varsigma\},$$

for some $c, \varsigma > 0$. The right hand side vanishes faster than any negative power of n , and hence is negligible relative to the target rate in (58).

Turning to the second term in (59), we introduce the further split

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \nu_{T_{\theta_0}}) \leq \mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \tilde{\nu}_{\hat{T}_n}) + \mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{T_{\theta_0}}, \nu_{T_{\theta_0}}) + \mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}}), \quad (60)$$

where $\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}}$ are defined as in (57) with $\tau = \hat{T}_n$ and $\tau = T_{\theta_0}$, respectively. Lemma 5.9 then shows that the expectations of the first two terms in (60), restricted to $\mathcal{E}_n$, are of lower order than the rate in (58). Lastly, by Lemma 5.11 and the Cauchy–Schwarz inequality,

$$E_{\theta_0}^{(n)}[\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}})1_{\mathcal{E}_n}] \leq n\sqrt{\kappa_{n,s}} \sqrt{E_{\theta_0}^{(n)}[\hat{D}_n(\mathcal{C}_n \hat{D}_n)]},$$

where $\mathcal{C}_n : \Theta^* \rightarrow \Theta$ is the linear diagonal operator defined by (80) below, and where the constant $\kappa_{n,s}$ is given by (79) and satisfies $\kappa_{n,s} \lesssim n^{-1+2s/(2\alpha+d)}$. By (23), the definition of $\mathcal{C}_n$, (24) and Fubini’s theorem,

$$E_{\theta_0}^{(n)}[\hat{D}_n(\mathcal{C}_n \hat{D}_n)] = \frac{1}{n} \sum_{k=1}^{\infty} \frac{I(\theta_0)[e_k]}{nc_0 + \lambda_k^{2\alpha+d}} \leq \frac{C_0}{n} \sum_{k=1}^{\infty} \frac{1}{nc_0 + \lambda_k^{2\alpha+d}} \lesssim n^{-2+\frac{d}{2\alpha+d}},$$

the last inequality following from (3) upon splitting the series at the spectral cut-off $\lambda_k^{2\alpha+d} \asymp n$, with both resulting pieces being of order $n^{-1+d/(2\alpha+d)}$. It follows that

$$E_{\theta_0}^{(n)}[\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}})] \lesssim n n^{-\frac{1}{2}+\frac{s}{2\alpha+d}} n^{-1+\frac{d/2}{2\alpha+d}} = n^{-\frac{\alpha-s}{2\alpha+d}}.$$

Combining the obtained upper bounds proves (58). $\square$

The rest of this section is devoted to proving the auxiliary results for the proof of Proposition 5.3. To that end, we introduce the following notation and select suitable exponents that appear repeatedly in the argument. Recall that $\alpha \wedge \beta > p$ and, from Assumption 3.1–3.4, that $m_K \in (d/2, p]$ and $r_K \geq 0$. Define

$$\rho := \frac{\alpha \wedge \beta - m_K}{2(\alpha - m_K) + d} \in (0, 1/2), \quad \delta := 1 - 2\rho = \frac{2(\alpha - \alpha \wedge \beta) + d}{2(\alpha - m_K) + d} \in (0, 1), \quad \epsilon_n := n^{-\rho}, \quad (61)$$

so that $n\epsilon_n^2 = n^\delta$, and for $\gamma \in (p, \alpha \wedge \beta]$ set

$$C_*(\gamma) := \frac{\gamma + p + 2r_K}{\gamma - p}. \quad (62)$$

The compatibility conditions (29) and (30) of Theorem 3.5 imply $\delta(1 + C_*(\alpha \wedge \beta)) < 1$. Since C_* is decreasing and continuous, we can therefore choose $\gamma \in (s \vee p, \alpha \wedge \beta)$, $a < 1/2$ and $b > \delta/2$ such that

$$b C_*(\gamma) < a < \frac{1 - \delta}{2}. \quad (63)$$

Furthermore, for these choices, the quantity $\omega := 2b(p + r_K)/(\gamma - p)$ satisfies

$$a > b + \omega, \quad 1 - \omega > \delta. \quad (64)$$

The preceding choices reflect the three Hilbert-scale geometries used in the proof. The conditioning set $\mathcal{B}$ appearing in (57) is a fixed-radius ball in the parameter space $\Theta = H_p$, on which the Fisher information is coercive; see Assumption 3.1. The index $m_K \in (d/2, p]$ corresponds to a weaker geometry, relative to which the sufficient-statistic fluctuations are controlled and the prior places sufficient mass on the shrinking ball $B_{\epsilon_n}^{H_{m_K}}(\theta_0)$; see Lemma 5.4. Finally, $\gamma \in (s \vee p, \alpha \wedge \beta)$ indexes a stronger geometry in which the complement of the growing ball $B_{n^b}^{H_\gamma}(\theta_0)$ has negligible prior probability; see the proof of Lemma 5.7.

5.2.1 Construction of the localising event

We first record a small-ball estimate for the prior Π in the Hilbert scale. Its verification is standard in Bayesian nonparametrics (Ghosal and Van der Vaart, 2017, Chapter 11). It is included for completeness and the convenience of the reader.

Lemma 5.4. *Let $\theta_0 \in H_\beta$ for some $\beta > 0$, and let Π be the α -regular Gaussian series prior (20), for some $\alpha > 0$. Then, for any $0 \leq m < \alpha \wedge \beta$ and all sufficiently small $\epsilon > 0$,*

$$-\log \Pi(B_\epsilon^{H_m}(\theta_0)) \lesssim \epsilon^{-\frac{d}{\alpha-m}} + \epsilon^{-\frac{(2\alpha+d-2\beta)_+}{\beta-m}}. \quad (65)$$

In particular, for $m_K \in (d/2, p]$ from Assumption 3.2 for ρ, δ and ϵ_n as in (61),

$$-\log \Pi(B_{\epsilon_n}^{H_{m_K}}(\theta_0)) \lesssim n^\delta = n\epsilon_n^2. \quad (66)$$

Proof. In the orthonormal basis $\{\lambda_k^{-m} e_k\}_{k \geq 1}$ of H_m , the coordinates of a draw from Π are independent centred Gaussian random variables with variances $\lambda_k^{-2(\alpha-m)-d} \asymp k^{-1-2(\alpha-m)/d}$, $k \in \mathbb{N}$. Classical small-deviation asymptotics for weighted sums of squared Gaussians (Bogachev, 1998, Section 4.3) give, as $\epsilon \rightarrow 0$,

$$-\log \Pi(B_\epsilon^{H_m}(0)) = -\log \Pr \left(\sum_{k=1}^{\infty} \lambda_k^{-2(\alpha-m)-d} Z_k^2 \leq \epsilon^2 \right) \asymp -\epsilon^{-d/(\alpha-m)},$$

yielding the first term of (65). Next, recall from Section 3.2 that the RKHS of Π is $H_{\alpha+d/2}$. Let P_K denote the projection onto $\Theta_K = \text{span}\{e_1, \dots, e_K\}$ and take $K(\epsilon) \in \mathbb{N}$ to be the smallest integer such that $\lambda_{K(\epsilon)} \geq ((1 + \|\theta_0\|_{H_\beta})/\epsilon)^{1/(\beta-m)}$. Then, $\lambda_{K(\epsilon)} \asymp ((1 + \|\theta_0\|_{H_\beta})/\epsilon)^{1/(\beta-m)}$ and

$$\|\theta_0 - P_{K(\epsilon)}\theta_0\|_{H_m} \leq \lambda_{K(\epsilon)}^{-(\beta-m)} \|\theta_0\|_{H_\beta} \leq \epsilon, \quad \|P_{K(\epsilon)}\theta_0\|_{H_{\alpha+d/2}}^2 \leq \lambda_{K(\epsilon)}^{(2\alpha+d-2\beta)_+} \|\theta_0\|_{H_\beta}^2.$$

The first claim follows from the second inequality in the last display and an application of the Cameron–Martin theorem as in van der Vaart and van Zanten (2008, Lemma 5.3). The second claim follows from the first upon choosing $m = m_K$ and $\epsilon = \epsilon_n$, since $\epsilon_n \rightarrow 0$. $\square$

The next lemma converts the exponential moment condition of Assumption 3.4 into two deviation inequalities for $\hat{D}_n$: one in the H_{-m_K} -norm and the other when $\hat{D}_n$ is evaluated at a fixed element of H_{m_K} .

Lemma 5.5. *Let Assumption 3.4 hold for some $A_T > 0$. Then, there exists a constant $v_T > 0$ such that, for all $u > 0$ and $n \in \mathbb{N}$,*

$$\mu_{\theta_0}^{(n)}(\|\hat{D}_n\|_{H_{-m_K}} \geq u) \leq 2 \exp \left\{ -\frac{nu^2}{2(v_T + A_T u)} \right\}, \quad (67)$$

and, for every fixed $h \in H_{m_K}$, $h \neq 0$, and all $u > 0$,

$$\mu_{\theta_0}^{(n)}(|\hat{D}_n(h)| \geq u) \leq 2 \exp \left\{ -\frac{nu^2}{2(v_T \|h\|_{H_{m_K}}^2 + A_T \|h\|_{H_{m_K}} u)} \right\}. \quad (68)$$

Proof. Write $\xi_i := \beta_{X_i} - T_{\theta_0}$. These are independent and identically distributed centred random elements of the separable Hilbert space H_{-m_K} . By (28) and the elementary bound $w^k \leq k! A_T^k e^{w/A_T}$, valid for any $w \geq 0$,

$$E_{\theta_0}^{(1)} \left[\|\xi_1\|_{H_{-m_K}}^k \right] \leq k! A_T^k E_{\theta_0}^{(1)} \left[e^{\|\xi_1\|_{H_{-m_K}}/A_T} \right] = \frac{k!}{2} v_T A_T^{k-2}, \quad k \geq 2,$$

with $v_T := 2A_T^2 E_{\theta_0}^{(1)}[e^{\|\xi_1\|_{H_{-m_K}}/A_T}] < \infty$. This is the Bernstein moment condition, and, recalling $\hat{D}_n = n^{-1} \sum_{i=1}^n \xi_i$, (67) is the corresponding Bernstein inequality for sums of independent random elements of a separable Hilbert space, e.g. (Pinelis, 1994, Theorem 3.3). For (68), note that $|\xi_i(h)| \leq \|\xi_i\|_{H_{-m_K}} \|h\|_{H_{m_K}}$, so the real random variables $\xi_i(h)$ are centred and satisfy the same moment condition with $v_T \|h\|_{H_{m_K}}^2$ and $A_T \|h\|_{H_{m_K}}$ in place of v_T and A_T ; the classical scalar Bernstein inequality then applies. $\square$

Finally, we note that Assumptions 3.1 and 3.2 yield the following Kullback–Leibler upper bound in H_{m_K} .

Lemma 5.6. *Let Assumptions 3.1 and 3.2 hold. Then, with $C_U := \frac{1}{2} C_0 \lambda_1^{-2m_K} + C_K R_K^{q_K}$,*

$$K(\theta|\theta_0) \leq C_U \|\theta - \theta_0\|_{H_{m_K}}^2, \quad \text{for all } \theta \in \Theta \cap B_{R_K}^{H_{m_K}}(\theta_0). \quad (69)$$

Proof. Fix $\theta \in \Theta \cap B_{R_K}^{H_{m_K}}(\theta_0)$ and write $h = \theta - \theta_0$. By (26) and (24),

$$K(\theta|\theta_0) \leq \frac{1}{2} I(\theta_0)[h] + C_K \|h\|_{H_{m_K}}^{2+q_K} \leq \frac{1}{2} C_0 \|h\|_{H_0}^2 + C_K R_K^{q_K} \|h\|_{H_{m_K}}^2,$$

and $\|h\|_{H_0} \leq \lambda_1^{-m_K} \|h\|_{H_{m_K}}$ because $m_K \geq 0$ and $\lambda_k \geq \lambda_1$ for all k . $\square$

We are now in a position to state and prove the main result of this subsection of the proof, identifying the localising event for the basic stochastic term split (59).

Lemma 5.7. *Let $\gamma \in (s \vee p, \alpha \wedge \beta)$, $a \in (0, 1/2)$ and $b > 0$, and let δ and ϵ_n be as in (61). For $C_1, C_2 > 0$, define the events $\mathcal{E}_n^{(1)} := \{\|\hat{D}_n\|_{H_{-m_K}} \leq n^{-a}\}$, $\mathcal{E}_n^{(2)} := \{\int_{H_{s \vee p}} e^{\ell_n(\theta)} d\Pi(\theta) \geq e^{-C_1 n^\delta}\}$,*

$$\mathcal{E}_n^{(3)} := \left\{ \int_{H_{s \vee p}} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{\ell_n(\theta)} d\Pi(\theta) \leq e^{n^\delta} \right\}$$

and

$$\mathcal{E}_n^{(4)} := \left\{ \int_{(B_{n^b}^{H_\gamma}(\theta_0))^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{\ell_n(\theta)} d\Pi(\theta) \leq e^{-C_2 n^{2b/2}} \right\},$$

Set $\mathcal{E}_n := \bigcap_{j=1}^4 \mathcal{E}_n^{(j)}$. Then, there exist constants $C_1, C_2 > 0$, depending only on the model, the prior and the above exponents, such that, for some $C > 0$ and all n large enough,

$$\mu_{\theta_0}^{(n)}(\mathcal{E}_n^c) \leq \exp\{-C n^\varsigma\}, \quad \varsigma := \min\{1 - 2a, \delta, 2b\} > 0. \quad (70)$$

Proof. We upper bound the probability of each event individually. For $\mathcal{E}_n^{(1)}$, we apply (67) with $u = n^{-a}$ obtaining $\mu_{\theta_0}^{(n)}((\mathcal{E}_n^{(1)})^c) \leq 2e^{-c_1 n^{1-2a}}$ for some $c_1 > 0$, where $1 - 2a > 0$ since $a < 1/2$.

The estimate for the second event is a standard argument in Bayesian nonparametrics and follows from the small ball lower bound in Lemma 5.4, cf. (Ghosal and Van der Vaart, 2017, Lemma 8.10). Indeed, restricting the integral to $B_{\epsilon_n}^{H_{m_K}}(\theta_0)$ and applying Jensen's inequality to the probability measure $\Pi(\cdot|B_{\epsilon_n}^{H_{m_K}}(\theta_0))$,

$$\begin{aligned} \int_{H_{s \vee p}} e^{\ell_n(\theta)} d\Pi(\theta) &\geq \Pi(B_{\epsilon_n}^{H_{m_K}}(\theta_0)) \exp \left\{ \int_{B_{\epsilon_n}^{H_{m_K}}(\theta_0)} \ell_n d\Pi(\theta|B_{\epsilon_n}^{H_{m_K}}(\theta_0)) \right\} \\ &= \Pi(B_{\epsilon_n}^{H_{m_K}}(\theta_0)) e^{n\hat{D}_n(\bar{h}_n) - n\bar{K}_n}, \end{aligned}$$

by the identity (55), where $\bar{h}_n := \Pi(B_{\epsilon_n}^{H_{m_K}}(\theta_0))^{-1} \int_{B_{\epsilon_n}^{H_{m_K}}(\theta_0)} (\theta - \theta_0) d\Pi(\theta)$ is a Bochner integral in H_{m_K} and $\bar{K}_n := \Pi(B_{\epsilon_n}^{H_{m_K}}(\theta_0))^{-1} \int_{B_{\epsilon_n}^{H_{m_K}}(\theta_0)} K(\theta|\theta_0) d\Pi(\theta)$. Here, the interchange of $\hat{D}_n$, which takes values in H_{-m_K} almost surely, with the Bochner integral is legitimate by the properties of the latter. Since $B_{\epsilon_n}^{H_{m_K}}(\theta_0)$ is closed and convex in H_{m_K} , its barycentre lies in $B_{\epsilon_n}^{H_{m_K}}(\theta_0)$, and it follows that $\|\bar{h}_n\|_{H_{m_K}} \leq \epsilon_n$. On the other hand, since $\epsilon_n \leq R_K$ for n large, Lemma 5.6 applies pointwise on $B_{\epsilon_n}^{H_{m_K}}(\theta_0)$ and gives $\bar{K}_n \leq C_U \epsilon_n^2$, whence $n\bar{K}_n \lesssim n\epsilon_n^2 = n^\delta$. If $\bar{h}_n = 0$ the required stochastic bound follows immediately; otherwise, applying (68) with $h = \bar{h}_n$ (which is deterministic) and $u = \epsilon_n^2$ yields

$$\begin{aligned} \mu_{\theta_0}^{(n)}(n|\hat{D}_n(\bar{h}_n)| \geq n^\delta) &= \mu_{\theta_0}^{(n)}(|\hat{D}_n(\bar{h}_n)| \geq \epsilon_n^2) \\ &\leq 2 \exp \left\{ -\frac{n\epsilon_n^4}{2(v_T \|\bar{h}_n\|_{H_{m_K}}^2 + A_T \|\bar{h}_n\|_{H_{m_K}} \epsilon_n^2)} \right\} \\ &\leq 2 \exp \left\{ -\frac{n\epsilon_n^2}{2(v_T + A_T \epsilon_n)} \right\} \leq 2e^{-c_2 n^\delta}, \end{aligned}$$

for some $c_2 > 0$ and all sufficiently large n , where we used $\|\bar{h}_n\|_{H_{m_K}} \leq \epsilon_n$ and $n\epsilon_n^2 = n^\delta$. Combining the above with $-\log \Pi(B_{\epsilon_n}^{H_{m_K}}(\theta_0)) \lesssim n^\delta$ from (66) and taking sufficiently large $C_1 > 0$ then yields $\mu_{\theta_0}^{(n)}((\mathcal{E}_n^{(2)})^c) \leq 2e^{-c_2 n^\delta}$.

Turning to the events $\mathcal{E}_n^{(3)}$ and $\mathcal{E}_n^{(4)}$, note that for every fixed θ , $E_{\theta_0}^{(n)}[e^{\ell_n(\theta)}] = E_\theta^{(1)}[1] = 1$. Fubini's theorem therefore gives, for any Borel set $A \subseteq H_{s \vee p}$,

$$E_{\theta_0}^{(n)} \left[\int_A (1 + \|\theta - \theta_0\|_{H_s}^2) e^{\ell_n(\theta)} d\Pi(\theta) \right] = \int_A (1 + \|\theta - \theta_0\|_{H_s}^2) d\Pi(\theta). \quad (71)$$

Taking $A = H_{s \vee p}$, we obtain $E_{\theta_0}^{(n)}[\int_{H_{s \vee p}} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{\ell_n(\theta)} d\Pi(\theta)] = 1 + E_\Pi[\|\theta - \theta_0\|_{H_s}^2] < \infty$ since $s < \alpha \wedge \beta$. Then, by Markov's inequality, $\mu_{\theta_0}^{(n)}((\mathcal{E}_n^{(3)})^c) = O(e^{-n^\delta})$. Lastly, taking $A = (B_{n^b}^{H_\gamma}(\theta_0))^c$ in (71), Cauchy–Schwarz inequality gives

$$\int_{(B_{n^b}^{H_\gamma}(\theta_0))^c} (1 + \|\theta - \theta_0\|_{H_s}^2) d\Pi(\theta) \leq (E_\Pi[(1 + \|\theta - \theta_0\|_{H_s}^2)^2])^{1/2} \Pi\left((B_{n^b}^{H_\gamma}(\theta_0))^c\right)^{1/2}.$$

The expectation on the right hand side is finite by Fernique's theorem (Bogachev, 1998, Theorem 2.8.5). Furthermore, since $\gamma < \alpha$, Π is supported on H_γ , and the Borell–Sudakov–Tsirelson inequality (Giné and Nickl, 2016, Theorem 2.5.8) gives

$$\Pi((B_{n^b}^{H_\gamma}(\theta_0))^c) \leq \Pi(\theta : \|\theta\|_{H_\gamma} > n^b - \|\theta_0\|_{H_\gamma}) \leq e^{-c_3 n^{2b}}$$

for some $c_3 > 0$ and all sufficiently large n . Another application of Markov's inequality implies that upon taking $C_2 > 0$ small enough, $\mu_{\theta_0}^{(n)}((\mathcal{E}_n^{(4)})^c) \leq e^{-c_4 n^{2b}/2}$ for some $c_4 > 0$. Adding the four upper bounds proves (70). $\square$

5.2.2 Bounds for the conditioning errors

The next preparatory step for the proof of Proposition 5.3 is to control, over the localising event $\mathcal{E}_n$ constructed in Lemma 5.7, the Wasserstein distance between the probability measures $\nu_{\hat{T}_n}, \nu_{T_{\theta_0}}$ and, respectively, their conditioned versions $\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}}$ over the set (56). We first show that on $\mathcal{E}_n$, the posterior masses and the weighted posterior second moments outside the conditioning set $\mathcal{B}$ are exponentially negligible. This ensures that the conditioning operation introduces an error smaller than any polynomial order.

Lemma 5.8. *Let b and ω be as chosen in (63) and (64), and let $\mathcal{E}_n$ be the event from Lemma 5.7. Then, there exists a constant $C > 0$ such that, on $\mathcal{E}_n$ and for all n large enough,*

$$\int_{\mathcal{B}^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{\ell_n(\theta)} d\Pi(\theta) \leq e^{-Cn^{\zeta'}}, \quad \zeta' = \min\{1 - \omega, 2b\} > \delta. \quad (72)$$

Consequently, on $\mathcal{E}_n$, both $\nu_{\hat{T}_n}(\mathcal{B}^c)$ and $\int_{\mathcal{B}^c} \|\theta - \theta_0\|_{H_s}^2 d\nu_{\hat{T}_n}(\theta)$ are bounded by $e^{C_1 n^\delta - Cn^{\zeta'}}$, and hence decay faster than any negative power of n .

Proof. Split $\mathcal{B}^c = (\mathcal{B}^c \cap B_{n^b}^{H_\gamma}(\theta_0)) \cup (\mathcal{B}^c \cap (B_{n^b}^{H_\gamma}(\theta_0))^c)$. The integral over $\mathcal{B}^c \cap (B_{n^b}^{H_\gamma}(\theta_0))^c$ is bounded by $e^{-C_2 n^{2b}/2}$ on $\mathcal{E}_n^{(4)}$. Hence, it suffices to treat $\mathcal{B}^c \cap B_{n^b}^{H_\gamma}(\theta_0)$, on which $h := \theta - \theta_0$

satisfies $\|h\|_{H_p} > R$ and $\|h\|_{H_\gamma} \leq n^b$, upper bounding, in view of the likelihood identity (55),

$$\int_{\mathcal{B}^c \cap B_{n^b}^{H_\gamma}(\theta_0)} (1 + \|h\|_{H_s}^2) \exp \left\{ n\hat{D}_n(h) - nK(\theta | \theta_0) \right\} d\Pi(\theta). \quad (73)$$

Fix any $\theta \in \mathcal{B}^c \cap B_{n^b}^{H_\gamma}(\theta_0)$. Since $m_K \leq p < \gamma$ by our choice of γ , we have $\|h\|_{H_{m_K}} \leq \lambda_1^{m_K - \gamma} \|h\|_{H_\gamma}$, so on $\mathcal{E}_n^{(1)}$,

$$n|\hat{D}_n(h)| \leq n\|\hat{D}_n\|_{H_{-m_K}} \|h\|_{H_{m_K}} \lesssim n^{1-a+b}.$$

Next, writing $p = -r_K(1-z) + \gamma z$ with $z = (p+r_K)/(\gamma+r_K)$, the interpolation inequality (1) gives $\|h\|_{H_p} \leq \|h\|_{H_{-r_K}}^{1-z} \|h\|_{H_\gamma}^z$ and hence

$$\|h\|_{H_{-r_K}} \geq \|h\|_{H_p}^{\frac{\gamma+r_K}{\gamma-p}} \|h\|_{H_\gamma}^{-\frac{p+r_K}{\gamma-p}} \geq R^{\frac{\gamma+r_K}{\gamma-p}} n^{-\omega/2},$$

with ω as in (64). The right-hand side tends to 0, so, by Assumption 3.3, $nK(\theta|\theta_0) \gtrsim n^{1-\omega}$. Finally, since $s \leq \gamma$, $1 + \|h\|_{H_s}^2 \leq 1 + \lambda_1^{2(s-\gamma)} \|h\|_{H_\gamma}^2 \lesssim n^{2b}$. Combining the three obtained estimates and recalling $a > b + \omega$ from (64), which ensures $1 - a + b < 1 - \omega$, we obtain for all sufficiently large n that (73) is upper bounded by

$$n^{2b} \exp \{ c_1 n^{1-a+b} - c_2 n^{1-\omega} \} \leq e^{-\frac{c_2}{2} n^{1-\omega}},$$

for some $c_1, c_2 > 0$. This concludes the verification of (72). The final assertion follows upon dividing (72) by the denominator bound of $\mathcal{E}_n^{(2)}$, using that $1 \vee \|\theta - \theta_0\|_{H_s}^2 \leq 1 + \|\theta - \theta_0\|_{H_s}^2$, and noting that $\varsigma' > \delta$. $\square$

The preceding exterior-mass estimates imply that, on $\mathcal{E}_n$, conditioning either measure $\nu_{\hat{T}_n}$ or $\nu_{T_{\theta_0}}$ on $\mathcal{B}$ introduces a negligible error in Wasserstein distance, as we now show.

Lemma 5.9. *For every $C > 0$ and all n large enough,*

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{T_{\theta_0}}, \tilde{\nu}_{T_{\theta_0}}) \lesssim n^{-C}. \quad (74)$$

Moreover, on the event $\mathcal{E}_n$ from Lemma 5.7, with $\varsigma' > 0$ the exponent from Lemma 5.8, for some constant $C' > 0$ and for all n large enough,

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \tilde{\nu}_{\hat{T}_n}) \leq e^{-C' n^{\varsigma'}}. \quad (75)$$

Proof. Fix any Borel probability measure $\nu \in \mathcal{P}_2(H_s)$ such that $\nu(\mathcal{B}) > 0$, and let

$$\tilde{\nu}(\cdot) := \nu(\cdot | \mathcal{B}) = \frac{\nu|_{\mathcal{B}}(\cdot)}{\nu(\mathcal{B})} = \frac{\nu|_{\mathcal{B}}(\cdot)}{1 - \nu(\mathcal{B}^c)}.$$

A simple coupling between ν and $\tilde{\nu}$ is given by

$$\pi := (\text{Id}, \text{Id})_{\#}(\nu|_{\mathcal{B}}) + \frac{1}{1 - \nu(\mathcal{B}^c)}(\nu|_{\mathcal{B}^c}) \otimes (\nu|_{\mathcal{B}}).$$

Indeed, its first marginal is given by, for any measurable $A \subseteq H_s$,

$$\begin{aligned}\pi_1(A) &= [(\text{Id}, \text{Id})_{\#}(\nu|_{\mathcal{B}})](A \times H_s) + \frac{1}{1 - \nu(\mathcal{B}^c)} [(\nu|_{\mathcal{B}^c}) \otimes (\nu|_{\mathcal{B}})](A \times H_s) \\ &= \nu(A \cap \mathcal{B}) + \frac{\nu(A \cap \mathcal{B}^c)\nu(\mathcal{B})}{1 - \nu(\mathcal{B}^c)} = \nu(A \cap \mathcal{B}) + \nu(A \cap \mathcal{B}^c) = \nu(A),\end{aligned}$$

and similarly it holds that the second marginal $\pi_2 = \tilde{\nu}$. Consequently, since there is no transport associated to the diagonal part of π ,

$$\begin{aligned}\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu, \tilde{\nu})^2 &\leq \int \|\theta - \eta\|_{H_s}^2 d\pi(\theta, \eta) \\ &= \frac{1}{1 - \nu(\mathcal{B}^c)} \int_{\mathcal{B}^c} \int_{\mathcal{B}} \|\theta - \eta\|_{H_s}^2 d\nu(\eta) d\nu(\theta) \\ &\leq 2 \int_{\mathcal{B}^c} \|\theta - \theta_0\|_{H_s}^2 d\nu(\theta) + \frac{2\nu(\mathcal{B}^c)}{1 - \nu(\mathcal{B}^c)} \int_{\mathcal{B}} \|\eta - \theta_0\|_{H_s}^2 d\nu(\eta) \\ &= 2 \int_{\mathcal{B}^c} \|\theta - \theta_0\|_{H_s}^2 d\nu(\theta) + 2\nu(\mathcal{B}^c) \int_{\mathcal{B}} \|\eta - \theta_0\|_{H_s}^2 d\tilde{\nu}(\eta).\end{aligned}$$

Apply the preceding inequality with $\nu = \nu_{T_{\theta_0}}$. The estimates (47) and (51) established in the proof of Proposition 5.1 imply that, for every $M > 0$,

$$\int_{\mathcal{B}^c} (1 + \|\theta - \theta_0\|_{H_s}^2) e^{-nK(\theta|\theta_0)} d\Pi(\theta) \leq n^{-M} Z_{I, \theta_0}^{(n)}, \quad \int_{H_s \vee p} e^{-nK(\theta|\theta_0)} d\Pi(\theta) \geq \frac{1}{2} Z_{I, \theta_0}^{(n)},$$

for all sufficiently large n , where $Z_{I, \theta_0}^{(n)}$ is the normalising constant in (50). Taking ratios shows that

$$\nu_{T_{\theta_0}}(\mathcal{B}^c) \vee \int_{\mathcal{B}^c} \|\theta - \theta_0\|_{H_s}^2 d\nu_{T_{\theta_0}}(\theta) = O(n^{-M})$$

for every $M > 0$. In particular, $\nu_{T_{\theta_0}}(\mathcal{B}^c) \leq 1/2$ for all sufficiently large n . Proposition 5.1 also gives

$$\int_{\mathcal{B}} \|\theta - \theta_0\|_{H_s}^2 d\tilde{\nu}_{T_{\theta_0}}(\theta) \leq 2 \int_{H_s \vee p} \|\theta - \theta_0\|_{H_s}^2 d\nu_{T_{\theta_0}}(\theta) = O(1).$$

Applying the coupling bound once more yields $\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{T_{\theta_0}}, \tilde{\nu}_{T_{\theta_0}})^2 = o(n^{-M})$ for every $M > 0$. Since M is arbitrary, replacing M by $2C$ and taking square roots proves (74).

There remains to consider $\nu = \nu_{\hat{T}_n}$. By Lemma 5.8, on $\mathcal{E}_n$,

$$\nu_{\hat{T}_n}(\mathcal{B}^c) \vee \int_{\mathcal{B}^c} \|\theta - \theta_0\|_{H_s}^2 d\nu_{\hat{T}_n}(\theta) \leq \exp \left\{ C_1 n^\delta - C n^{\varsigma'} \right\}.$$

Since $\varsigma' > \delta$, the right-hand side converges to zero. In particular, $\nu_{\hat{T}_n}(\mathcal{B}^c) \leq 1/2$ for all sufficiently large n . We next control the second moment of $\tilde{\nu}_{\hat{T}_n}$ which equals

$$\int_{\mathcal{B}} \|\theta - \theta_0\|_{H_s}^2 d\tilde{\nu}_{\hat{T}_n}(\theta) = \frac{\int_{\mathcal{B}} \|\theta - \theta_0\|_{H_s}^2 e^{\ell_n(\theta)} d\Pi(\theta)}{\int_{\mathcal{B}} e^{\ell_n(\theta)} d\Pi(\theta)}.$$

On $\mathcal{E}_n$,

$$\int_{\mathcal{B}} e^{\ell_n(\theta)} d\Pi(\theta) = \nu_{\hat{T}_n}(\mathcal{B}) \int_{H_{s \vee p}} e^{\ell_n(\theta)} d\Pi(\theta) \geq \frac{1}{2} e^{-C_1 n^\delta}, \quad \int_{\mathcal{B}} \|\theta - \theta_0\|_{H_s}^2 e^{\ell_n(\theta)} d\Pi(\theta) \leq e^{n^\delta},$$

whence $\int_{\mathcal{B}} \|\theta - \theta_0\|_{H_s}^2 d\tilde{\nu}_{\hat{T}_n}(\theta) \leq 2e^{(1+C_1)n^\delta}$. Thus, on $\mathcal{E}_n$, the coupling bound gives

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\hat{T}_n}, \tilde{\nu}_{\hat{T}_n})^2 \leq 2e^{C_1 n^\delta - C n^{\varsigma'}} + 4e^{(1+2C_1)n^\delta - C n^{\varsigma'}} \leq e^{-2C' n^{\varsigma'}},$$

holding for some $C' > 0$ and for all sufficiently large n , having used that $\varsigma' > \delta$. Taking square roots proves (75). $\square$

5.2.3 Transport techniques for conditioned posteriors

We now come to the estimate of the Wasserstein distance between the conditioned probability measures $\tilde{\nu}_{\hat{T}_n}(\cdot) = \nu_{\hat{T}_n}(\cdot | \mathcal{B})$ and $\tilde{\nu}_{T_{\theta_0}}(\cdot) = \nu_{T_{\theta_0}}(\cdot | \mathcal{B})$. A key technical tool is the Poincaré inequality derived below, which provides an upper bound, uniformly for $\tau \in \Theta^*$, for the variances $\text{Var}_{\tilde{\nu}_\tau}[F(\theta)]$ of *regular cylindrical functions* $F : \Theta \rightarrow \mathbb{R}$. These take the form $F(\theta) = f(\theta_1, \dots, \theta_K)$ for some $K \in \mathbb{N}$ and some bounded and smooth $f : \mathbb{R}^K \rightarrow \mathbb{R}$ with bounded derivatives of all orders. For any $q \in \mathbb{R}$, let

$$\nabla^{H_q} F(\theta) = \sum_{k=1}^K \lambda_k^{-2q} \partial_k f(\theta_1, \dots, \theta_K) e_k, \quad \theta \in \Theta,$$

be the gradient of F in the H_q -geometry, where $\partial_k f$, $k = 1, \dots, K$ are the standard partial derivatives of f .

The proof of the aforementioned Poincaré inequality hinges on an approximation argument based on the finite-dimensional Galerkin subspaces $\Theta_K := \text{span}\{e_1, \dots, e_K\}$, $K \in \mathbb{N}$, and on the associated affine spaces $\Theta_{0,K} := \theta_0 + \Theta_K$. We denote by $P_K : \Theta \rightarrow \Theta_K$ the canonical projection operator and let $P_{0,K}\theta := P_K\theta + (I - P_K)\theta_0$, which satisfies

$$P_{0,K}\theta - \theta_0 = P_K(\theta - \theta_0), \quad \theta \in \Theta. \quad (76)$$

Throughout, for any fixed $q \in [0, \alpha + d/2]$, we regard $\Theta_{0,K}$ as a K -dimensional Euclidean affine space equipped with the metric induced by H_q . Its tangent space coincides with Θ_K , equipped with the H_q -geometry, for which $\{\lambda_k^{-q} e_k\}_{k=1}^K$ is an orthonormal basis. By definition $P_{0,K}$, every $\theta \in \Theta_{0,K}$ admits the unique representation $\theta = (I - P_K)\theta_0 + \sum_{k=1}^K \theta_k e_k$.

Next, define the push-forward prior distribution $\Pi_{0,K} := (P_{0,K})_\# \Pi$ on $\Theta_{0,K}$. For $\tau \in \Theta^*$, denote by $\nu_{\tau,K}$ the τ -induced Galerkin posterior on $\Theta_{0,K}$, given by

$$d\nu_{\tau,K}(\theta) = \frac{\exp\{n[\tau(\theta) - M(\theta)]\}}{\int_{\Theta_{0,K}} \exp\{n[\tau(\theta) - M(\theta)]\} d\Pi_{0,K}(\theta)} d\Pi_{0,K}(\theta), \quad \theta \in \Theta_{0,K}. \quad (77)$$

Further set $\mathcal{B}_K := \mathcal{B} \cap \Theta_{0,K}$ and consider the conditioned Galerkin posterior $\tilde{\nu}_{\tau,K}(\cdot) := \nu_{\tau,K}(\cdot | \mathcal{B}_K)$ on $\Theta_{0,K}$. By (76), $\mathcal{B}_K = \{\theta \in \Theta_{0,K} : \|P_K(\theta - \theta_0)\|_{H_p} \leq R\}$ is a closed, bounded and convex subset of the finite-dimensional affine space $\Theta_{0,K}$ containing θ_0 ; in particular, $\Pi_{0,K}(\mathcal{B}_K) > 0$ and $\tilde{\nu}_{\tau,K}$ is well defined.

Lastly, we introduce two diagonal operators that control the reciprocal of the curvature of the negative log-posterior through a directional lower bound. For $q \in [0, \alpha + d/2]$ and $n \in \mathbb{N}$, let $\mathcal{A}_{n,q} : H_q \rightarrow H_q$ be the linear diagonal operator

$$\mathcal{A}_{n,q} e_k := \frac{\lambda_k^{2q}}{nc_0 + \lambda_k^{2\alpha+d}} e_k, \quad k \in \mathbb{N}, \quad (78)$$

with $c_0 > 0$ the local coercivity constant of Assumption 3.1, and let

$$\kappa_{n,q} := \|\mathcal{A}_{n,q}\|_{\mathcal{L}(H_q)} = \sup_{\|h\|_{H_q}=1} \|\mathcal{A}_{n,q} h\|_{H_q} = \sup_{k \in \mathbb{N}} \frac{\lambda_k^{2q}}{nc_0 + \lambda_k^{2\alpha+d}} \lesssim n^{-1+\frac{2q}{2\alpha+d}}, \quad (79)$$

the upper bound following by maximising over the spectral variable. The restriction $q \leq \alpha + d/2$ ensures that the displayed supremum is finite. Define also the positive linear operator $\mathcal{C}_n : \Theta^* \rightarrow \Theta$ by

$$\mathcal{C}_n \chi := \sum_{k=1}^{\infty} \frac{\chi(e_k)}{nc_0 + \lambda_k^{2\alpha+d}} e_k, \quad \chi \in \Theta^*, \quad (80)$$

for which we have, since $\alpha > p$,

$$\|\mathcal{C}_n \chi\|_{H_p}^2 = \sum_{k=1}^{\infty} \frac{\lambda_k^{2p} \chi(e_k)^2}{(nc_0 + \lambda_k^{2\alpha+d})^2} \lesssim \sum_{k=1}^{\infty} \lambda_k^{-2p} \chi(e_k)^2 = \|\chi\|_{H_{-p}}^2.$$

Lemma 5.10. *Let Assumption 3.1 be satisfied and suppose that M is twice continuously Fréchet differentiable on Θ . Then, for every $q \in [0, \alpha + d/2]$, $n \in \mathbb{N}$, $\tau \in \Theta^*$, and any regular cylindrical function $F : \Theta \rightarrow \mathbb{R}$,*

$$\text{Var}_{\tilde{\nu}_\tau} [F(\theta)] \leq \int_{\mathcal{B}} \langle \mathcal{A}_{n,q} \nabla^{H_q} F(\theta), \nabla^{H_q} F(\theta) \rangle_{H_q} d\tilde{\nu}_\tau(\theta) \leq \kappa_{n,q} \int_{\mathcal{B}} \|\nabla^{H_q} F(\theta)\|_{H_q}^2 d\tilde{\nu}_\tau(\theta). \quad (81)$$

Proof. We first establish the corresponding inequality for the conditioned Galerkin measures $\tilde{\nu}_{\tau,K}$, namely that for every $K \in \mathbb{N}$ and any bounded and smooth function $F : \Theta_{0,K} \rightarrow \mathbb{R}$ with bounded derivatives,

$$\begin{aligned} \text{Var}_{\tilde{\nu}_{\tau,K}} [F(\theta)] &\leq \int_{\mathcal{B}_K} \langle \mathcal{A}_{n,q} \nabla^{H_q} F(\theta), \nabla^{H_q} F(\theta) \rangle_{H_q} d\tilde{\nu}_{\tau,K}(\theta) \\ &\leq \kappa_{n,q} \int_{\mathcal{B}_K} \|\nabla^{H_q} F(\theta)\|_{H_q}^2 d\tilde{\nu}_{\tau,K}(\theta). \end{aligned} \quad (82)$$

Let $q \in [0, \alpha + d/2]$, $n, K \in \mathbb{N}$, $\tau \in \Theta^*$ and F be fixed. Identifying $\Theta_{0,K}$ with $\mathbb{R}^K$, the negative log-density of the unconditioned measure $\nu_{\tau,K}$ from (77) with respect to Lebesgue measure equals, up to an additive constant,

$$V_{\tau,K}(\theta) := n\{M(\theta) - \tau(\theta)\} + \frac{1}{2} \sum_{k=1}^K \lambda_k^{2\alpha+d} \theta_k^2, \quad \theta \in \Theta_{0,K}. \quad (83)$$

Crucially, τ enters $V_{\tau,K}$ linearly, and we have, for all $\theta \in \Theta_{0,K}$,

$$DV_{\tau,K}(\theta)[h] = n\{DM(\theta)[h] - \tau(h)\} + \sum_{k=1}^K \lambda_k^{2\alpha+d} \theta_k h_k, \quad h \in \Theta_K,$$

and

$$D^2V_{\tau,K}(\theta)[h, g] = nD^2M(\theta)[h, g] + \langle h, g \rangle_{H_{\alpha+d/2}}, \quad h, g \in \Theta_K,$$

the latter being independent of τ . For $\theta \in \Theta_{0,K}$, let $Hess^{H_q}V_{\tau,K}(\theta) : \Theta_K \rightarrow \Theta_K$ denote the H_q -Hessian of $V_{\tau,K}$ at θ , namely the unique self-adjoint operator satisfying

$$\langle Hess^{H_q}V_{\tau,K}(\theta)h, g \rangle_{H_q} = D^2V_{\tau,K}(\theta)[h, g], \quad h, g \in \Theta_K,$$

given by

$$Hess^{H_q}V_{\tau,K}(\theta)h = \sum_{k=1}^K \lambda_k^{-2q} \left[nD^2M(\theta)[h, e_k] + \lambda_k^{2\alpha+d} h_k \right] e_k, \quad h \in \Theta_K.$$

The set $\mathcal{B}_K = \mathcal{B} \cap \Theta_{0,K}$ is convex and contained in $B_{R_0}^\Theta(\theta_0)$ by (56), so the local coercivity condition (25) gives, for $\theta \in \mathcal{B}_K$ and $h \in \Theta_K$,

$$D^2V_{\tau,K}(\theta)[h, h] \geq nc_0 \|h\|_{H_0}^2 + \|h\|_{H_{\alpha+d/2}}^2 = \sum_{k=1}^K \lambda_k^{-2q} (nc_0 + \lambda_k^{2\alpha+d}) \lambda_k^{2q} h_k^2.$$

This implies that, as self-adjoint operators on Θ_K equipped with the H_q -inner product,

$$\mathcal{A}_{n,q,K}^{1/2} Hess^{H_q}V_{\tau,K}(\theta) \mathcal{A}_{n,q,K}^{1/2} \succeq I, \quad \text{for all } \theta \in \mathcal{B}_K,$$

where $\mathcal{A}_{n,q,K}$ the restriction of (78) to Θ_K (i.e., to the first K coordinates). The linear change of variables $v = \mathcal{A}_{n,q,K}^{-1/2}(\theta - \theta_0)$ transforms $\mathcal{B}_K$ into the convex set $\mathcal{D}_{n,q,K} := \mathcal{A}_{n,q,K}^{-1/2}(\mathcal{B}_K - \theta_0) \subset \Theta_K$. Let $\bar{\nu}_{\tau,K}$ denote the push-forward of $\tilde{\nu}_{\tau,K}$ on Θ_K under this transformation. As the constant Jacobian of the change of variables is absorbed into the normalising constant, we have

$$d\bar{\nu}_{\tau,K}(v) = \frac{\mathbf{1}_{\mathcal{D}_{n,q,K}}(v) e^{-W_{\tau,K}(v)}}{\int_{\mathcal{D}_{n,q,K}} e^{-W_{\tau,K}(v)} dv} dv, \quad v \in \Theta_K,$$

where $W_{\tau,K}(v) := V_{\tau,K}(\theta_0 + \mathcal{A}_{n,q,K}^{1/2}v)$. Since $\mathcal{A}_{n,q,K}^{1/2}$ is self-adjoint in the H_q -inner product, the chain rule and the preceding curvature bound give

$$Hess^{H_q}W_{\tau,K}(v) = \mathcal{A}_{n,q,K}^{1/2} Hess^{H_q}V_{\tau,K}(\theta_0 + \mathcal{A}_{n,q,K}^{1/2}v) \mathcal{A}_{n,q,K}^{1/2} \succeq I, \quad \text{for all } v \in \mathcal{D}_{n,q,K}.$$

Thus, $\bar{\nu}_{\tau,K}$ has unit lower curvature on a convex domain. The Bakry–Émery criterion on convex domains (Kolesnikov and Milman, 2016, Theorem 2.1) therefore yields the log-Sobolev inequality

$$Ent_{\bar{\nu}_{\tau,K}} [\bar{F}^2(v)] \leq 2 \int_{\mathcal{D}_{n,q,K}} \|\nabla^{H_q} \bar{F}(v)\|_{H_q}^2 d\bar{\nu}_{\tau,K}(v)$$

holding for every bounded smooth function $\bar{F} : \Theta_K \rightarrow \mathbb{R}$ with bounded derivatives, where $Ent_{\bar{\nu}_{\tau,K}}(\bar{F}^2(v)) := \int_{\mathcal{D}_{n,q,K}} \bar{F}^2(v) \log \left(\bar{F}^2(v) / \int_{\mathcal{D}_{n,q,K}} \bar{F}^2(v) d\bar{\nu}_{\tau,K}(v) \right) \bar{\nu}_{\tau,K}(v)$. The usual linearisation step for log-Sobolev inequalities then yields the Poincaré inequality

$$Var_{\bar{\nu}_{\tau,K}} [\bar{F}(v)] \leq \int_{\mathcal{D}_{n,q,K}} \|\nabla^{H_q} \bar{F}(v)\|_{H_q}^2 d\bar{\nu}_{\tau,K}(v).$$

In particular, for $\bar{F}(v) := F(\theta_0 + \mathcal{A}_{n,q,K}^{1/2} v)$, with F fixed at the beginning of the proof, we have

$$\nabla^{H_q} \bar{F}(v) = \mathcal{A}_{n,q,K}^{1/2} \nabla^{H_q} F(\theta_0 + \mathcal{A}_{n,q,K}^{1/2} v), \quad v \in \Theta_K,$$

whence applying the inverse change of variable in the $\theta = \theta_0 + \mathcal{A}_{n,q,K}^{1/2} v$ within the Poincaré inequality for $\bar{\nu}_{\tau,K}$ gives

$$Var_{\tilde{\nu}_{\tau,K}}[F(\theta)] \leq \int_{\mathcal{B}_K} \langle \mathcal{A}_{n,q,K} \nabla^{H_q} F(\theta), \nabla^{H_q} F(\theta) \rangle_{H_q} d\tilde{\nu}_{\tau,K}(\theta).$$

Since $\mathcal{A}_{n,q,K}$ is the restriction of $\mathcal{A}_{n,q}$ to Θ_K , this proves the first inequality in (82). The second follows upon noting that

$$\kappa_{n,q,K} := \sup_{\theta \in \Theta_K : \|\theta\|_{H_q} = 1} \|\mathcal{A}_{n,q,K} \theta\|_{H_q} = \max_{k=1,\dots,K} \frac{\lambda_k^{2q}}{nc_0 + \lambda_k^{2\alpha+d}} \leq \kappa_{n,q},$$

and that in view of the positivity and self-adjointness of $\mathcal{A}_{n,q,K}$,

$$\langle \mathcal{A}_{n,q,K} \nabla^{H_q} F, \nabla^{H_q} F \rangle_{H_q} \leq \kappa_{n,q,K} \|\nabla^{H_q} F\|_{H_q}^2 \leq \kappa_{n,q} \|\nabla^{H_q} F\|_{H_q}^2.$$

There remains to prove (81) by letting $K \rightarrow \infty$. We have $P_{0,K}\theta \rightarrow \theta$ in $\Theta = H_p$ for every $\theta \in \Theta$ and, by (76), $\|P_K(\theta - \theta_0)\|_{\Theta} \rightarrow \|\theta - \theta_0\|_{\Theta}$. Thus, since $\tau \in \Theta^*$ and $M : \Theta \rightarrow \mathbb{R}$ are continuous,

$$1_{\{\|P_K(\theta - \theta_0)\|_{\Theta} \leq R\}} e^{n[\tau(P_{0,K}\theta) - M(P_{0,K}\theta)]} \rightarrow 1_{\mathcal{B}}(\theta) e^{n[\tau(\theta) - M(\theta)]}.$$

Convexity of M gives

$$\tau(P_{0,K}\theta) - M(P_{0,K}\theta) \leq \tau(\theta_0) - M(\theta_0) + [\tau - DM(\theta_0)](P_{0,K}\theta - \theta_0).$$

Whenever the indicator is nonzero, (76) implies that $P_{0,K}\theta \in \mathcal{B}$, and hence the last is bounded above by $R\|\tau - DM(\theta_0)\|_{\Theta^*}$, whence, for every $\theta \in \Theta^*$,

$$1_{\{\|P_K(\theta - \theta_0)\|_{\Theta} \leq R\}} e^{n[\tau(P_{0,K}\theta) - M(P_{0,K}\theta)]} \leq e^{n[\tau(\theta_0) - M(\theta_0) + R\|\tau - DM(\theta_0)\|_{\Theta^*}]},$$

which is a finite constant independent of K . Thus, by the dominated convergence theorem,

$$\int_{\Theta} 1_{\{\|P_K(\theta - \theta_0)\|_{\Theta} \leq R\}} e^{n[\tau(P_{0,K}\theta) - M(P_{0,K}\theta)]} d\Pi(\theta) \rightarrow \int_{\Theta} 1_{\mathcal{B}}(\theta) e^{n[\tau(\theta) - M(\theta)]} d\Pi(\theta).$$

Let now $F : \Theta \rightarrow \mathbb{R}$ be any regular cylindrical function $J \in \mathbb{N}$ coordinates. The preceding convergence and another application of the dominated convergence theorem yields

$$\begin{aligned} \int_{\mathcal{B}_K} F(\theta) d\tilde{\nu}_{\tau,K}(\theta) &= \frac{\int_{\Theta} F(P_{0,K}\theta) 1_{\{\|P_K(\theta-\theta_0)\|_{\Theta} \leq R\}} e^{n[\tau(P_{0,K}\theta)-M(P_{0,K}\theta)]} d\Pi(\theta)}{\int_{\Theta} 1_{\{\|P_K(\theta-\theta_0)\|_{\Theta} \leq R\}} e^{n[\tau(P_{0,K}\theta)-M(P_{0,K}\theta)]} d\Pi(\theta)} \\ &\rightarrow \frac{\int_{\Theta} F(\theta) 1_{\mathcal{B}}(\theta) e^{n[\tau(\theta)-M(\theta)]} d\Pi(\theta)}{\int_{\Theta} 1_{\mathcal{B}}(\theta) e^{n[\tau(\theta)-M(\theta)]} d\Pi(\theta)} = \int_{\mathcal{B}} F(\theta) d\tilde{\nu}_{\tau}(\theta). \end{aligned}$$

Apply this conclusion to $F, F^2, \langle \mathcal{A}_{n,q,K} \nabla^{H_q} F, \nabla^{H_q} F \rangle_{H_q}$ (which equals $\langle \mathcal{A}_{n,q} \nabla^{H_q} F, \nabla^{H_q} F \rangle_{H_q}$ when $K > J$) and $\|\nabla^{H_q} F\|_{H_q}^2$, which are all regular cylindrical and depend only on the first J coordinates shows that $\text{Var}_{\tilde{\nu}_{\tau,K}}[F(\theta)] \rightarrow \text{Var}_{\tilde{\nu}_{\tau}}[F(\theta)]$ as well as

$$\int_{\mathcal{B}_K} \langle \mathcal{A}_{n,q,K} \nabla^{H_q} F(\theta), \nabla^{H_q} F(\theta) \rangle_{H_q} d\tilde{\nu}_{\tau,K}(\theta) \rightarrow \int_{\mathcal{B}} \langle \mathcal{A}_{n,q} \nabla^{H_q} F(\theta), \nabla^{H_q} F(\theta) \rangle_{H_q} d\tilde{\nu}_{\tau}(\theta),$$

and

$$\int_{\mathcal{B}_K} \|\nabla^{H_q} F(\theta)\|_{H_q}^2 d\tilde{\nu}_{\tau,K} \rightarrow \int_{\mathcal{B}} \|\nabla^{H_q} F(\theta)\|_{H_q}^2 d\tilde{\nu}_{\tau}.$$

Taking $K \rightarrow \infty$ in (82) therefore proves (81). $\square$

The final key lemma combines the above Poincaré inequality for conditioned posteriors with an interpolation argument to estimate the Wasserstein distance between $nu_{\hat{T}_n}(\cdot)$ and $\tilde{\nu}_{T_{\theta_0}}$ through a preconditioned quadratic measure of the empirical fluctuation $\hat{D}_n = \hat{T}_n - T_{\theta_0}$.

Lemma 5.11. *Under the assumptions of Proposition 5.3, almost surely,*

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}}) \leq n \sqrt{\kappa_{n,s} \hat{D}_n(\mathcal{C}_n \hat{D}_n)}, \quad (84)$$

where $\mathcal{C}_n : \Theta^* \rightarrow \Theta$ is the linear operator from (80) and $\kappa_{n,s} > 0$ is defined as in (79).

Proof. Similarly to the proof of Lemma 5.10, we first establish the finite-dimensional analogue of the transport estimate (84) for the conditioned Galerkin posteriors $\tilde{\nu}_{\hat{T}_n,K}$ and $\tilde{\nu}_{T_{\theta_0},K}$, defined as after (77) with $\tau = \hat{T}_n$ and $\tau = T_{\theta_0}$, respectively. The proof is based on the dynamic formulation of the Wasserstein distance (Benamou and Brenier, 2000). For any $t \in [0, 1]$, define the Θ^* -valued random element $\hat{\tau}_t := T_{\theta_0} + t\hat{D}_n$, and let $\tilde{\nu}_{\hat{\tau}_t,K}$ be the associated conditioned Galerkin posterior. Then $\{\tilde{\nu}_{\hat{\tau}_t,K}\}_{t \in [0,1]}$ defines a curve of Borel probability measures in $\mathcal{P}_2(H_s)$ that are supported on the common convex set $\mathcal{B}_K$, and joins $\tilde{\nu}_{\hat{\tau}_0,K} = \tilde{\nu}_{T_{\theta_0},K}$ to $\tilde{\nu}_{\hat{\tau}_1,K} = \tilde{\nu}_{\hat{T}_n,K}$. An admissible velocity field for this curve is a measurable vector field $v_{t,K} : \mathcal{B}_K \rightarrow \Theta_K$, $t \in [0, 1]$ such that

$$\int_0^1 \left(\int_{\mathcal{B}_K} \|v_{t,K}(\theta)\|_{H_s}^2 d\tilde{\nu}_{\hat{\tau}_t,K}(\theta) \right)^{1/2} dt < \infty,$$

and satisfying the weak continuity equation

$$\frac{d}{dt} \int_{\mathcal{B}_K} \phi(\theta) d\tilde{\nu}_{\hat{\tau}_t,K}(\theta) = \int_{\mathcal{B}_K} \langle \nabla^{H_s} \phi(\theta), v_{t,K}(\theta) \rangle_{H_s} d\tilde{\nu}_{\hat{\tau}_t,K}(\theta) \quad (85)$$

for almost every $t \in (0, 1)$ and every bounded smooth test function $\phi : \Theta_{0,K} \rightarrow \mathbb{R}$ of the form $\phi(\theta) = f(\theta_1, \dots, \theta_K)$, $\theta = (I - P_K)\theta_0 + \sum_{k=1}^K \theta_k e_k \in \Theta_{0,K}$, for some bounded and smooth $f : \mathbb{R}^K \rightarrow \mathbb{R}$ with bounded derivatives of all orders. The dynamic formulation then gives (Ambrosio et al., 2008, Theorem 8.3.1)

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n, K}, \tilde{\nu}_{T_{\theta_0}, K}) \leq \int_0^1 \left(\int_{\mathcal{B}_K} \|v_{t,K}(\theta)\|_{H_s}^2 d\tilde{\nu}_{\hat{T}_t, K}(\theta) \right)^{1/2} dt. \quad (86)$$

We move to the construction of an admissible velocity field satisfying the continuity equation (85) and obtain a uniform bound for the integrated norm appearing in (86). Using that $\frac{d}{dt}\hat{\tau}_t(\theta) = \hat{D}_n(\theta)$ for all $t \in [0, 1]$ and $\theta \in \Theta$, differentiating under the integral sign the left hand side of (85) gives

$$\begin{aligned} \frac{d}{dt} \int_{\mathcal{B}_K} \phi(\theta) d\tilde{\nu}_{\hat{\tau}_t, K}(\theta) &= n \int_{\mathcal{B}_K} \phi(\theta) \hat{D}_n(\theta) d\tilde{\nu}_{\hat{\tau}_t, K}(\theta) \\ &\quad - n \left(\int_{\mathcal{B}_K} \phi(\theta) d\tilde{\nu}_{\hat{\tau}_t, K}(\theta) \right) \left(\int_{\mathcal{B}_K} \hat{D}_n(\theta) d\tilde{\nu}_{\hat{\tau}_t, K}(\theta) \right) \\ &= n \text{Cov}_{\tilde{\nu}_{\hat{\tau}_t, K}}(\phi(\theta), \hat{D}_n(\theta)). \end{aligned}$$

Consequently, by Cauchy–Schwarz inequality

$$\left| \frac{d}{dt} \int_{\mathcal{B}_K} \phi(\theta) d\tilde{\nu}_{\hat{\tau}_t, K}(\theta) \right| \leq n \sqrt{\text{Var}_{\tilde{\nu}_{\hat{\tau}_t, K}}[\phi(\theta)]} \sqrt{\text{Var}_{\tilde{\nu}_{\hat{\tau}_t, K}}[\hat{D}_n(\theta)]}. \quad (87)$$

We next estimate the two variances using the Poincaré inequality from Lemma 5.10. Applying the second inequality in (82) with $q = s$, we directly obtain

$$\text{Var}_{\tilde{\nu}_{\hat{\tau}_t, K}}[\phi(\theta)] \leq \kappa_{n,s} \int_{\mathcal{B}_K} \|\nabla^{H_s} \phi(\theta)\|_{H_s}^2 d\tilde{\nu}_{\hat{\tau}_t, K}(\theta).$$

For the second variance in (87), we apply the first inequality in (82) with $q = s$ to the restriction of the linear functional $\hat{D}_n$ to $\Theta_{0,K}$. Its H_s -gradient is the constant vector field

$$\nabla^{H_s} \hat{D}_n(\theta) = \sum_{k=1}^K \lambda_k^{-2s} \hat{D}_n(e_k) e_k.$$

Consequently,

$$\left\langle \mathcal{A}_{n,s,K} \nabla^{H_s} \hat{D}_n(\theta), \nabla^{H_s} \hat{D}_n(\theta) \right\rangle_{H_s} = \sum_{k=1}^K \lambda_k^{2s} \frac{\hat{D}_n(e_k)}{nc_0 + \lambda_k^{2\alpha+d}} \lambda_k^{-2s} \hat{D}_n(e_k) = \sum_{k=1}^K \frac{\hat{D}_n(e_k)^2}{nc_0 + \lambda_k^{2\alpha+d}},$$

which we note is independent of s (and θ). Since the preceding integrand is constant in θ , the first inequality in (82) yields

$$\text{Var}_{\tilde{\nu}_{\hat{\tau}_t, K}}[\hat{D}_n(\theta)] \leq \sum_{k=1}^K \frac{\hat{D}_n(e_k)^2}{nc_0 + \lambda_k^{2\alpha+d}}. \quad (88)$$

Strictly speaking, (82) is stated for bounded functions, whereas $\hat{D}_n$ is not. This is immaterial here, since $\mathcal{B}_K$ is bounded, $|\hat{D}_n(\theta)| \leq \|\hat{D}_n\|_{\Theta^*}(\|\theta_0\|_{\Theta} + R)$ on $\theta \in \mathcal{B}_K$, and it suffices to apply the first inequality in (82) to $H \circ \hat{D}_n$ for a smooth bounded functions $H : \mathbb{R} \rightarrow \mathbb{R}$ with bounded derivatives of all orders such that $H(z) = z$ for all $|z| \leq 1 + \|\hat{D}_n\|_{\Theta^*}(\|\theta_0\|_{\Theta} + R)$. Then $H \circ \hat{D}_n = \hat{D}_n$ and $\nabla^{H_s}(H \circ \hat{D}_n) = \nabla^{H_s} \hat{D}_n$ on $\mathcal{B}_K$, which carries all the mass of $\tilde{\nu}_{\hat{\tau}_t, K}$. Combining the obtained bounds with (87) yields

$$\left| \frac{d}{dt} \int_{\mathcal{B}_K} \phi(\theta) d\tilde{\nu}_{\hat{\tau}_t, K}(\theta) \right| \leq n \left(\sum_{k=1}^K \frac{\hat{D}_n(e_k)^2}{nc_0 + \lambda_k^{2\alpha+d}} \right)^{1/2} \left(\kappa_{n,s} \int_{\mathcal{B}_K} \|\nabla^{H_s} \phi\|_{H_s}^2 d\tilde{\nu}_{\hat{\tau}_t, K} \right)^{1/2}.$$

We now use the preceding bound to construct the admissible velocity field $v_{t,K}$ for (86). By identifying the affine space $\Theta_{0,K}$ with $\mathbb{R}^K$ as done previously, the set $\mathcal{B}_K = \{\theta \in \Theta_{0,K} : \|\theta - \theta_0\|_{H_p} \leq R\}$ is a bounded ellipsoid in $\mathbb{R}^K$, with smooth boundary. Let $H^1(\mathcal{B}_K)$ denote the standard Hilbert-Sobolev space of square integrable functions with square integrable first-order weak derivatives, equipped with the usual inner product and norm, and denote by

$$\dot{H}^1(\mathcal{B}_K) := \left\{ \varphi \in H^1(\mathcal{B}_K) : \int_{\mathcal{B}_K} \varphi(\theta) d\theta = 0 \right\}.$$

For each fixed K and $t \in [0, 1]$, the density of $\tilde{\nu}_{\hat{\tau}_t, K}$ with respect to the Lebesgue measure on $\mathcal{B}_K$ is proportional to $\exp\{-V_{\hat{\tau}_t, K}(\theta)\}$, $\theta \in \mathcal{B}_K$, where $V_{\hat{\tau}_t, K}$ is the potential from (83) with $\tau = \hat{\tau}_t$. Since $[0, 1] \times \mathcal{B}_K$ is compact, such density is bounded and bounded away from zero by constants that do not depend on t . Furthermore, since for all differentiable $\varphi : \mathcal{B}_K \rightarrow \mathbb{R}$

$$\|\nabla^{H_s} \varphi(\theta)\|_{H_s}^2 = \sum_{k=1}^K \lambda_k^{-2s} |\partial_k \varphi(\theta)|^2,$$

whence $\lambda_K^{-2s} \|\nabla \varphi(\theta)\|_{\mathbb{R}^K}^2 \leq \|\nabla^{H_s} \varphi(\theta)\|_{H_s}^2 \leq \lambda_1^{-2s} \|\nabla \varphi(\theta)\|_{\mathbb{R}^K}^2$, for all $\theta \in \Theta$, the norms

$$\|\varphi\|_{t,K} := \left(\int_{\mathcal{B}_K} \|\nabla^{H_s} \varphi(\theta)\|_{H_s}^2 d\tilde{\nu}_{\hat{\tau}_t, K}(\theta) \right)^{1/2}, \quad t \in [0, 1],$$

are uniformly equivalent to the usual gradient norm on $\dot{H}^1(\mathcal{B}_K)$. For smooth $\phi \in C^\infty(\overline{\mathcal{B}_K}) \cap \dot{H}^1(\mathcal{B}_K)$, define

$$L_{t,K}(\phi) := \frac{d}{dt} \int_{\mathcal{B}_K} \phi(\theta) d\tilde{\nu}_{\hat{\tau}_t, K}(\theta).$$

The preceding covariance bound shows that $L_{t,K}$ extends to a bounded linear functional on $\dot{H}^1(\mathcal{B}_K)$ satisfying

$$|L_{t,K}(\phi)| \leq n \left(\kappa_{n,s} \sum_{k=1}^K \frac{\hat{D}_n(e_k)^2}{nc_0 + \lambda_k^{2\alpha+d}} \right)^{1/2} \|\phi\|_{t,K}, \quad (89)$$

and the Riesz representation theorem therefore gives a unique $u_{t,K} \in \dot{H}^1(\mathcal{B}_K)$ such that

$$L_{t,K}(\phi) = \int_{\mathcal{B}_K} \langle \nabla^{H_s} \phi(\theta), \nabla^{H_s} u_{t,K}(\theta) \rangle_{H_s} d\tilde{\nu}_{\hat{\tau}_t, K}(\theta), \quad \text{for all } \phi \in \dot{H}^1(\mathcal{B}_K). \quad (90)$$

Since $L_{t,K}$ annihilates constants and gradients are unchanged by adding constants, this identity also holds for every bounded smooth test function on $\Theta_{0,K}$. Moreover, the continuous dependence on t of the density of $\tilde{\nu}_{\hat{\tau}_t,K}$ and of the functional $L_{t,K}$ readily imply that the map $t \in [0, 1] \mapsto u_{t,K}$ is continuous. Hence, $v_{t,K} := \nabla^{H_s} u_{t,K}$ may be chosen jointly measurable in t and θ . In view of (90), $v_{t,K}$ solves the continuity equation (85), and therefore is an admissible velocity field for the curve $\{\tilde{\nu}_{\hat{\tau}_t,K}\}_{t \in [0,1]}$. Finally, taking $\phi = u_{t,K}$ in (89) and (90) gives

$$\int_{\mathcal{B}_K} \|\nabla^{H_s} u_{t,K}(\theta)\|_{H_s}^2 d\tilde{\nu}_{\hat{\tau}_t,K}(\theta) \leq n^2 \kappa_{n,s} \sum_{k=1}^K \frac{\hat{D}_n(e_k)^2}{nc_0 + \lambda_k^{2\alpha+d}},$$

and, via (86),

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n,K}, \tilde{\nu}_{T_{\theta_0},K}) \leq n \left(\kappa_{n,s} \sum_{k=1}^K \frac{\hat{D}_n(e_k)^2}{nc_0 + \lambda_k^{2\alpha+d}} \right)^{1/2}. \quad (91)$$

There remains to lift the result to $\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}})$ by letting $K \rightarrow \infty$. The argument used at the end of the proof of Lemma 5.10 applies without changes to give $\tilde{\nu}_{\tau,K} \rightarrow \tilde{\nu}_\tau$ weakly in H_s . Thus, to deduce convergence in Wasserstein distance, we only have to check convergence of the H_s -second moments (Ambrosio et al., 2008, Chapter 7). For all $\theta \in H_s$, $P_{0,K}\theta \rightarrow \theta$ in H_s . Further, by (76), $\|P_{0,K}\theta - \theta_0\|_{H_s} \leq \|\theta - \theta_0\|_{H_s}$. Since the latter is Π -integrable as $s < \alpha$, the same dominated-convergence argument yields as required

$$\int_{\mathcal{B}_K} \|\theta - \theta_0\|_{H_s}^2 d\tilde{\nu}_{\tau,K}(\theta) \longrightarrow \int_{\mathcal{B}} \|\theta - \theta_0\|_{H_s}^2 d\tilde{\nu}_\tau(\theta).$$

Conclude that $\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\tau,K}, \tilde{\nu}_\tau) \rightarrow 0$ for all $\tau \in \Theta^*$. Applying this with $\hat{\tau}_0 = T_{\theta_0}$ and $\hat{\tau}_1 = \hat{T}_n$ and combining (91) with the triangle inequality for the Wasserstein distance gives, letting $K \rightarrow \infty$,

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\tilde{\nu}_{\hat{T}_n}, \tilde{\nu}_{T_{\theta_0}}) \leq n \left(\kappa_{n,s} \sum_{k=1}^{\infty} \frac{\hat{D}_n(e_k)^2}{nc_0 + \lambda_k^{2\alpha+d}} \right)^{1/2} = n \sqrt{\kappa_{n,s} \hat{D}_n(\mathcal{C}_n \hat{D}_n)},$$

concluding the proof. $\square$

Remark 5.12 (Global transport estimate). The same argument, without conditioning, justifies the global transport estimate used on $\mathcal{E}_n^c$ in the proof of Proposition 5.3. Indeed, global convexity of M gives the two Poincaré inequalities stated there, first on the Galerkin spaces and then on Θ by the limiting argument of Lemma 5.10. Although the linear functional $\chi := \tau - \tau'$ is now unbounded, the Galerkin measures $\nu_{\tau,K}$ have Gaussian tails by global strong log-concavity. Hence, applying the Poincaré inequality to smooth truncations with increasing resolution extends it to χ . Repeating the preceding finite-dimensional transport argument therefore gives

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\tau,K}, \nu_{\tau',K}) \leq n \sqrt{\kappa_s} \left(\sum_{k=1}^K \lambda_k^{-2\alpha-d} (\tau - \tau')(e_k)^2 \right)^{1/2}.$$

Finally, the pullback argument and dominated convergence used above, now combining convexity of M with Fernique’s theorem, shows that $\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\tau,K}, \nu_{\tau}) \rightarrow 0$ for every $\tau \in \Theta^*$. Passing to the limit at both endpoints yields

$$\mathcal{W}_2^{\mathcal{P}_2(H_s)}(\nu_{\tau}, \nu_{\tau'}) \leq n\sqrt{\kappa_s} \|\tau - \tau'\|_{H_{-\alpha-d/2}},$$

which is precisely the employed crude global bound.

References

- L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows: in metric spaces and in the space of probability measures*. Springer Science Business & Media, 2008.
- D. Bakry, F. Barthe, P. Cattiaux, and A. Guillin. A simple proof of the Poincaré inequality for a large class of probability measures including the log-concave case. *Electronic Communications in Probability*, 13:60–66, 2008. doi: 10.1214/ECP.v13-1352.
- E. Belitser and S. Ghosal. Adaptive Bayesian inference on the mean of an infinite-dimensional normal distribution. *The Annals of Statistics*, 31(2):536–559, 2003. doi: 10.1214/aos/1051027880.
- E. N. Belitser, P. Serra, and H. van Zanten. Rate-optimal Bayesian intensity smoothing for inhomogeneous Poisson processes. *Journal of Statistical Planning and Inference*, 166: 24–35, 2015. doi: 10.1016/j.jspi.2014.03.009.
- J.-D. Benamou and Y. Brenier. A computational fluid mechanics solution to the monge-kantorovich mass transfer problem. *Numerische Mathematik*, 84(3):375–393, 2000.
- P. K. Bhattacharya. Estimation of a probability density function and its derivatives. *Sankhyā: The Indian Journal of Statistics, Series A*, 29(4):373–382, 1967.
- V. I. Bogachev. *Gaussian Measures*, volume 62 of *Mathematical Surveys and Monographs*. American Mathematical Society, 1998.
- L. D. Brown. *Fundamentals of Statistical Exponential Families with Applications in Statistical Decision Theory*, volume 9 of *IMS Lecture Notes–Monograph Series*. Institute of Mathematical Statistics, Hayward, CA, 1986.
- S. Canu and A. Smola. Kernel methods and the exponential family. *Neurocomputing*, 69 (7–9):714–720, 2006.
- I. Castillo. On Bayesian supremum norm contraction rates. *The Annals of Statistics*, 42(5): 2058–2091, 2014. doi: 10.1214/14-AOS1253.
- I. Castillo. *Bayesian Nonparametric Statistics: École d’Été de Probabilités de Saint-Flour LI – 2023*, volume 2358 of *Lecture Notes in Mathematics*. Springer, Cham, 2024. doi: 10.1007/978-3-031-74035-0.
- I. Castillo and R. Nickl. Nonparametric Bernstein–von Mises theorem in Gaussian white noise. *The Annals of Statistics*, 41(4):1999–2028, 2013.

- D. Comaniciu and P. Meer. Mean shift: A robust approach toward feature space analysis. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 24(5):603–619, 2002. doi: 10.1109/34.1000236.
- F. Comte, C. Duval, and O. Sacko. Optimal adaptive estimation on $\mathbb{R}$ or $\mathbb{R}^+$ of the derivatives of a density. *Mathematical Methods of Statistics*, 29(1):1–31, 2020.
- D. D. Cox. An analysis of Bayesian inference for nonparametric regression. *The Annals of Statistics*, 21(2):903–923, 1993.
- P. Diaconis and D. Freedman. On the consistency of Bayes estimates. *The Annals of Statistics*, 14(1):1–26, 1986.
- E. Dolera and E. Mainini. On uniform continuity of posterior distributions. *Statistics & Probability Letters*, 157:108627, 2020.
- E. Dolera and E. Mainini. Lipschitz continuity of probability kernels in the optimal transport framework. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 59(4):1778–1812, 2023. doi: 10.1214/23-AIHP1389.
- E. Dolera, S. Favaro, and M. Giordano. On strong posterior contraction rates for Besov–Laplace priors in the white noise model. *arXiv preprint arXiv:2411.06981*, 2024a.
- E. Dolera, S. Favaro, and E. Mainini. Strong posterior contraction rates via Wasserstein dynamics. *Probability Theory and Related Fields*, 189(1):659–720, 2024b.
- S. Donnet, V. Rivoirard, J. Rousseau, and C. Scricciolo. Posterior concentration rates for counting processes with Aalen multiplicative intensities. *Bayesian Analysis*, 12(1):53–87, 2017.
- J. L. Doob. Application of the theory of martingales. *Le calcul des probabilités et ses applications*, pages 23–27, 1949.
- H. W. Engl, M. Hanke, and A. Neubauer. *Regularization of Inverse Problems*, volume 375 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 1996.
- L. C. Evans. *Partial Differential Equations*. Graduate studies in mathematics. American Mathematical Society, 1998. ISBN 9780821807729. URL <https://books.google.it/books?id=YHJfvwEACAAJ>.
- S. Fischer and I. Steinwart. Sobolev norm learning rates for regularized least-squares algorithms. *Journal of Machine Learning Research*, 21(205):1–38, 2020. URL <https://jmlr.org/papers/v21/19-734.html>.
- D. Freedman. On the Bernstein–von Mises theorem with infinite-dimensional parameters. *The Annals of Statistics*, 27(4):1119–1140, 1999.
- K. Fukumizu. Exponential manifold by reproducing kernel Hilbert spaces. In P. Gibilisco, E. Riccomagno, M. P. Rogantin, and H. P. Wynn, editors, *Algebraic and Geometric Methods in Statistics*, pages 291–306. Cambridge University Press, Cambridge, 2009. doi: 10.1017/CBO9780511642401.019.

- K. Fukumizu. Estimation with infinite-dimensional exponential family and Fisher divergence. *Information Geometry*, 7:609–622, 2024. doi: 10.1007/s41884-023-00122-z.
- C. R. Genovese, M. Perone-Pacifico, I. Verdinelli, and L. Wasserman. Non-parametric inference for density modes. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 78(1):99–126, 2016.
- S. Ghosal and A. van der Vaart. Convergence rates of posterior distributions for noniid observations. *The Annals of Statistics*, 35(1):192–223, 2007.
- S. Ghosal and A. W. Van der Vaart. *Fundamentals of nonparametric Bayesian inference*, volume 44. Cambridge University Press, 2017.
- S. Ghosal, J. K. Ghosh, and A. W. van der Vaart. Convergence rates of posterior distributions. *The Annals of Statistics*, 28(2):500–531, 2000.
- E. Giné and R. Nickl. Rates of contraction for posterior distributions in L^r -metrics, $1 \leq r \leq \infty$. *The Annals of Statistics*, 39(6):2883–2911, 2011.
- E. Giné and R. Nickl. *Mathematical Foundations of Infinite-Dimensional Statistical Models*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, New York, 2016.
- A. Gretton, K. M. Borgwardt, M. J. Rasch, B. Schölkopf, and A. Smola. A kernel two-sample test. *Journal of Machine Learning Research*, 13(1):723–773, 2012.
- S. Gugushvili, A. W. van der Vaart, and D. Yan. Bayesian linear inverse problems in regularity scales. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 56(3):2081–2107, 2020.
- W. Härdle and T. M. Stoker. Investigating smooth multiple regression by the method of average derivatives. *Journal of the American Statistical Association*, 84(408):986–995, 1989. doi: 10.1080/01621459.1989.10478863.
- N. L. Hjort, C. Holmes, P. Müller, and S. G. Walker, editors. *Bayesian Nonparametrics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 2010.
- R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge university press, 2012.
- A. Kirichenko and H. van Zanten. Optimality of Poisson processes intensity learning with Gaussian processes. *Journal of Machine Learning Research*, 16:2909–2919, 2015.
- B. T. Knapik, A. W. van der Vaart, and J. H. van Zanten. Bayesian inverse problems with Gaussian priors. *The Annals of Statistics*, 39(5):2626–2657, 2011.
- B. T. Knapik, B. T. Szabó, A. W. van der Vaart, and J. H. van Zanten. Bayes procedures for adaptive inference in inverse problems for the white noise model. *Probability Theory and Related Fields*, 164(3–4):771–813, 2016. doi: 10.1007/s00440-015-0619-7.

- A. V. Kolesnikov and E. Milman. Riemannian metrics on convex sets with applications to Poincaré and log-Sobolev inequalities. *Calculus of Variations and Partial Differential Equations*, 55(4):77, 2016. doi: 10.1007/s00526-016-1018-3.
- S. G. Krein and J. I. Petunin. Scales of Banach spaces. *Russian Mathematical Surveys*, 21(2):85–159, 1966.
- Y. A. Kutoyants. *Statistical Inference for Spatial Poisson Processes*, volume 134 of *Lecture Notes in Statistics*. Springer-Verlag, New York, 1998.
- M. Ledoux and M. Talagrand. *Probability in Banach Spaces: isoperimetry and processes*, volume 23. Springer, 1991.
- Z. Liu and M. Li. Optimal plug-in Gaussian processes for modeling derivatives. *Journal of the American Statistical Association*, 2026. doi: 10.1080/01621459.2026.2612778. Published online.
- P. Müller, F. A. Quintana, A. Jara, and T. Hanson. *Bayesian Nonparametric Data Analysis*. Springer Series in Statistics. Springer, Cham, 2015. doi: 10.1007/978-3-319-18968-0.
- F. Natterer. Error bounds for Tikhonov regularization in Hilbert scales. *Applicable Analysis*, 18(1–2):29–37, 1984.
- I. Pinelis. Optimum bounds for the distributions of martingales in Banach spaces. *The Annals of Probability*, 22(4):1679–1706, 1994. doi: 10.1214/aop/1176988477.
- G. Pistone and C. Sempì. An infinite-dimensional geometric structure on the space of all the probability measures equivalent to a given one. *The Annals of Statistics*, 23(5):1543–1561, 1995.
- V. Rivoirard and J. Rousseau. Posterior concentration rates for infinite dimensional exponential families. *Bayesian Analysis*, 7(2):311–334, 2012. doi: 10.1214/12-BA710.
- J. Rousseau and K. Mengersen. Asymptotic behaviour of the posterior distribution in overfitted mixture models. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 73(5):689–710, 2011.
- L. Schwartz. On Bayes procedures. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 4(1):10–26, 1965.
- C. Scricciolo. Convergence rates for Bayesian density estimation of infinite-dimensional exponential families. *The Annals of Statistics*, 34(6):2897–2920, 2006.
- Z. Shang and G. Cheng. Gaussian approximation of general non-parametric posterior distributions. *Information and Inference: A Journal of the IMA*, 7(3):509–529, 2018. doi: 10.1093/imaiai/iax017.
- W. Shen and S. Ghosal. Posterior contraction rates of density derivative estimation. *Sankhyā: The Indian Journal of Statistics, Series A*, 79(2):336–354, 2017. doi: 10.1007/s13171-017-0105-7.

- X. Shen and L. Wasserman. Rates of convergence of posterior distributions. *The Annals of Statistics*, 29(3):687–714, 2001.
- R. S. Singh. Improvement on some known nonparametric uniformly consistent estimators of derivatives of a density. *The Annals of Statistics*, 5(2):394–399, 1977. doi: 10.1214/aos/1176343805.
- Y. Song and S. Ermon. Generative modeling by estimating gradients of the data distribution. In *Advances in Neural Information Processing Systems*, volume 32, 2019.
- B. Sriperumbudur, K. Fukumizu, A. Gretton, A. Hyvärinen, and R. Kumar. Density estimation in infinite dimensional exponential families. *Journal of Machine Learning Research*, 18(57):1–59, 2017.
- C. J. Stone. Optimal global rates of convergence for nonparametric regression. *The Annals of Statistics*, 10(4):1040–1053, 1982. doi: 10.1214/aos/1176345969.
- M. E. Taylor. *Partial differential equations II. Qualitative Studies of Linear Equations*. Springer, New York, 2011.
- A. W. Van der Vaart. *Asymptotic statistics*, volume 3. Cambridge university press, 2000.
- A. W. van der Vaart and J. H. van Zanten. Rates of contraction of posterior distributions based on Gaussian process priors. *The Annals of Statistics*, 36(3):1435–1463, 2008.
- S. G. Walker, A. Lijoi, and I. Prünster. On rates of convergence for posterior distributions in infinite-dimensional models. *The Annals of Statistics*, 35(2):738–746, 2007. doi: 10.1214/009053606000001361.
- A. Wibisono, Y. Wu, and K. Y. Yang. Optimal score estimation via empirical Bayes smoothing. In *Proceedings of the Thirty-Seventh Conference on Learning Theory*, volume 247 of *Proceedings of Machine Learning Research*, pages 4958–4991. PMLR, 2024. URL <https://proceedings.mlr.press/v247/wibisono24a.html>.
- R. Wong. *Asymptotic approximations of integrals*. SIAM, 2001.
- W. W. Yoo and S. Ghosal. Supremum norm posterior contraction and credible sets for nonparametric multivariate regression. *The Annals of Statistics*, 44(3):1069–1102, 2016. doi: 10.1214/15-AOS1398.
- W. W. Yoo, V. Rivoirard, and J. Rousseau. Adaptive supremum norm posterior contraction: Wavelet spike-and-slab and anisotropic Besov spaces. *arXiv preprint arXiv:1708.01909*, 2018.