

Calibrating Inelastic Markets to Options: The Lean Marketron and the Generalized Langevin Equation

Andrey Itkin^{1*}

¹ FRE Department, Tandon School of Engineering, New York University, USA.

August 24, 2026

The Marketron model of [Halperin and Itkin, 2026] and its option pricing extension in [Halperin and Itkin, 2025] suffer from structural non-identifiability: an eighteen-parameter space traps solvers in suboptimal local minima and renders economic quantities unmeasurable. By removing exact scaling gauges and sign symmetries, freezing non-financial parameters by explicit criteria, and adiabatically eliminating the fast hidden signal, we derive a robust nine-parameter reduced model. A Gauss-Newton Hessian with empty null space and a manifold-boundary analysis confirm that the reduced core carries no exact symmetry and admits no further reduction. A diffusive correlation between flow and return innovations captures the short-maturity skew. A staged calibration from the physical measure to the risk-neutral measure, illustrated on SPX options, fits the whole surface with a single parameter set. The same reduction turns the wedge between the physical and pricing values of the flow block into a well-defined market price of flow risk rather than a ridge artifact, identifiable here for the first time, though a single surface constrains its level only weakly. Finally, our analysis reveals that in the Marketron model the log-price obeys a generalized Langevin equation with a closed-form, state-modulated memory kernel, and that the memory variable itself is the exact Markovian lift of this kernel. This mapping also yields a testable condition, the equality of the signal and memory relaxation rates, which on the SPX surface come out well separated, though both weakly identified, placing the fitted market tentatively in the driven, non-equilibrium regime and turning the active-matter reading from an analogy into a falsifiable constraint.

1 Introduction

In [Halperin and Itkin, 2026] we introduced the Marketron model of price formation in an inelastic market. The dynamics are driven by money flows and their impact on prices, and take the form of nonlinear diffusion of a quasiparticle (the marketron) in the space of the log-price x_t , a memory variable y_t encoding past flows, and an unobservable return predictor θ_t . In [Halperin and Itkin, 2025] we extended the framework to option markets. Since y_t and θ_t are non-tradable, the market is incomplete, and we constructed option prices via exponential utility indifference following [Grasselli and Hurd, 2007]. The resulting Hamilton-Jacobi-Bellman (HJB) equation was solved by a combination of operator splitting, the Cole-Hopf transformation and a Gaussian radial basis function (RBF) method, with all matrix elements available in closed form.

*e-mail: aitkin@nyu.edu

The pricing method is fast. The calibration is not. In [Halperin and Itkin, 2025] the specification carried fifteen free parameters, and the least-squares objective built on market option quotes was found to have many local minima and long flat valleys, so that a differential evolution solver with constrained search achieved only a modest relative accuracy of about 5–8%, with different runs returning visibly different parameter sets of comparable fit quality. The correlated generalization we adopt here (Section 2) adds three further coordinates and brings the full specification to eighteen, Eq. (7), so the problem is if anything worse. This is a classical symptom of a non-identifiable, or “sloppy”, parameterization [Gutenkunst et al., 2007; Transtrum, Machta, and Sethna, 2011]. The data constrain a few stiff parameter combinations while leaving many soft directions essentially free.

Non-identifiability is not only a numerical nuisance. The central empirical claim of the Marketron program is that flow impact and market inelasticity can be measured from market data. If the calibration objective has a continuous ridge along which the flow-impact constant c trades off against the memory-scale parameters, then a reported value of c is a point on a ridge, not an estimate. Reduction to an identifiable core is therefore a precondition for the economic reading of the model. It is also a precondition for any falsifiable joint test. A joint calibration of the full eighteen-parameter model to several data sets proves little, because failure can always be attributed to the optimizer. On an identifiable low-parameter core the same test carries weight.

The purpose of this paper is accordingly to cure the disease rather than the symptom. Instead of investing more computational effort into global optimization of an ill-posed objective, we reduce the parameter space so that the remaining coordinates are identifiable from the option surface. The reduction proceeds in four steps of increasing depth.

First, in Section 2 we exhibit exact and near-exact symmetries of the model. The memory variable admits a scaling gauge: option prices are invariant under a simultaneous rescaling of y and the parameters attached to it, up to the constraint that couples the two occurrences of the flow-impact constant c . The signal block possesses a discrete sign symmetry. Both are removed by normalization, which eliminates one continuous parameter and halves the number of equivalent minima. We also show that the risk-aversion parameter enters the pricing operator only through products with the hidden-factor volatilities, so it must be fixed exogenously rather than calibrated.

Second, in Section 2.3 we argue that parameters of a purely numerical nature, such as the regularization constant $\bar{\epsilon}$ of the marketron potential and the initial values of the hidden states, must not compete with financial parameters in the calibration objective. We fix them a priori by explicit numerical criteria.

Third, in Section 3 we perform the central structural reduction. The signal θ_t is a fast Ornstein-Uhlenbeck (OU) process. For maturities satisfying $kT \gtrsim 1$ the option price depends on the signal essentially through the stationary averages of $f(\theta)$ and $h(\theta)$, which for the trigonometric specification of [Halperin and Itkin, 2025] are available in closed form. Adiabatic elimination collapses the five-parameter signal block into two effective constants and reduces the pricing PDE from three to two spatial dimensions. We also derive a deterministic transient correction that retains the time dependence of the conditional mean of the signal. The correction reintroduces the relaxation rate k as a single term-structure shape parameter, which is what makes a global fit with one parameter set across all maturities feasible. We treat this single-parameter-set property as a claim to be tested (Sections 5 and 6.3) rather than as an assumption, and it is what distinguishes the reduced model from reduced-form smile parameterizations, which are refit maturity by maturity.

Fourth, in Section 5 we make the procedure self-verifying. We compute profile likelihoods and the eigenspectrum of the Gauss-Newton Hessian of the calibration objective, and apply the manifold boundary approximation method (MBAM) of [Transtrum and Qiu, 2014] to check that the limits taken in Sections 2, 2.3 and 3 coincide with the sloppiest directions of the full model. In this sense the reduction is not ad hoc: it is the model that the option data can actually see.

One structural element is added rather than removed. The original specification of [Halperin and

[Itkin, 2026] assumes independent Brownian motions for the spot and memory stochastic variables. We introduce a diffusive correlation ρ between the flow innovations $dW^{(y)}$ and the return innovations $dW^{(x)}$. The motivation is twofold. Financially, contemporaneous correlation between flows and returns is the diffusive-level content of price impact, and forcing it to zero is an unnatural restriction in a model built around impact. Technically, the smile generated by the state-dependent drift vanishes linearly in maturity at the short end, so short-dated skew must come from the diffusion structure, and ρ is the only coefficient that can supply it. The correlation survives every reduction step at the cost of a single parameter and, as shown in Section 2.2, it in fact restores a degree of freedom that the risk-aversion analysis removes.

The final reduced model has nine parameters for the whole surface. Section 6.3 describes a staged calibration in which parameters of the physical measure are inherited from the time-series calibration of [Halperin and Itkin, 2026], while option data determine the risk-adjusted block. The correlation ρ is placed with the physical-measure parameters, since on the option surface it is confounded with the local-volatility asymmetry and convexity (Section 5); the wedge between the time-series and option-implied values of the flow-block parameters c and $\bar{f}$ (the flow-impact coupling and the averaged return-predictor drift) then defines a market price of flow risk, which the reduction makes identifiable for the first time.

Section 4 gives the reduced system a second reading: substituting the memory variable into the log-price equation shows the Marketron to be exactly a one-dimensional overdamped generalized Langevin equation (GLE) in the log-price, with a memory kernel we compute in closed form. This kernel is modulated by the state, specifically by the product of the market potential gradients at the endpoints of the memory integral, which means that memory is strong in the default region and weak away from it, explicitly delimiting the relation to the lag-only class of [Itkin, 2026a]. Furthermore, this mapping yields a testable parametric condition, an equality between the memory relaxation rate μ and the signal relaxation rate k , that distinguishes an equilibrium regime from a driven, non-equilibrium one. This transforms an interpretive analogy into a falsifiable constraint on two independently calibrated rates, see Section 4.4.

Then Section 6 reports numerical experiments: a Black–Scholes synthetic validation of the pricing and calibration pipeline, a staged global term-structure calibration to the SPX surface with a single parameter set, and the resulting market price of flow risk. Section 7 concludes.

2 The Marketron model, exact symmetries and gauge fixing

We briefly recall the setup of [Halperin and Itkin, 2026; Halperin and Itkin, 2025], with one generalization: the Brownian motions driving the price and the memory variable are allowed to be correlated. Under the physical measure the state (x_t, y_t, θ_t) solves

$$\begin{aligned} dx_t &= \mu_x dt + \sigma dW_t^{(x)}, & \mu_x &= f(\theta_t) + \bar{\eta} - c_x y_t V'_M(x_t), \\ dy_t &= \mu_y dt + \sigma_y dW_t^{(y)}, & \mu_y &= h(\theta_t) + \mu(\bar{y} - y_t) - c_y V_M(x_t), \\ d\theta_t &= \mu_\theta dt + \sigma_z dW_t^{(\theta)}, & \mu_\theta &= k(\hat{\theta} - \theta_t), \end{aligned} \quad (1)$$

with the correlation structure

$$d\langle W^{(x)}, W^{(y)} \rangle_t = \rho dt, \quad d\langle W^{(x)}, W^{(\theta)} \rangle_t = \rho_{x\theta} dt, \quad d\langle W^{(y)}, W^{(\theta)} \rangle_t = 0, \quad (2)$$

and the potential slope

$$V'_M(x) = -e^{-x} \left(1 - \frac{g}{e^x + \bar{\epsilon}g} \right), \quad (3)$$

regularized at $x \rightarrow -\infty$ by the constant $\bar{\epsilon}$. The original model is recovered at $\rho = \rho_{x\theta} = 0$. Following [Halperin and Itkin, 2025] we use the trigonometric signal transformation

$$f(\theta) = b_1 \cos \theta, \quad h(\theta) = b_2 \sin \theta. \quad (4)$$

Option prices are defined by exponential utility indifference with risk aversion γ . With correlated hidden factors the certainty equivalent $C(t, x, y, \theta)$ solves

$$0 = \frac{\partial C}{\partial t} + \frac{1}{2}\sigma^2 \frac{\partial^2 C}{\partial x^2} + \rho\sigma\sigma_y \frac{\partial^2 C}{\partial x\partial y} + \rho_{x\theta}\sigma\sigma_\theta \frac{\partial^2 C}{\partial x\partial\theta} + \frac{1}{2}\sigma_y^2 \frac{\partial^2 C}{\partial y^2} + \frac{1}{2}\sigma_\theta^2 \frac{\partial^2 C}{\partial\theta^2} + \bar{\eta} \frac{\partial C}{\partial x} \\ + \left(\mu_y - \rho\sigma_y \frac{\bar{\mu}_x}{\sigma}\right) \frac{\partial C}{\partial y} + \left(\mu_\theta - \rho_{x\theta}\sigma_\theta \frac{\bar{\mu}_x}{\sigma}\right) \frac{\partial C}{\partial\theta} + \frac{\gamma}{2} \nabla_h C^\top \tilde{\Sigma} \nabla_h C - \frac{\bar{\mu}_x^2}{2\gamma\sigma^2} - rC, \quad (5)$$

where $\nabla_h C = (\partial_y C, \partial_\theta C)^\top$ and

$$\tilde{\Sigma} = \begin{pmatrix} \sigma_y^2(1 - \rho^2) & -\sigma_y\sigma_\theta\rho\rho_{x\theta} \\ -\sigma_y\sigma_\theta\rho\rho_{x\theta} & \sigma_\theta^2(1 - \rho_{x\theta}^2) \end{pmatrix} \quad (6)$$

is the residual covariance of the hidden factors conditional on the traded one. Here $\sigma_\theta \equiv \sigma_z$. The drift adjustments proportional to $\bar{\mu}_x/\sigma$ are the market-price-of-risk terms generated by the projection of the hidden noise onto the traded noise. At $\rho = \rho_{x\theta} = 0$ equation Eq. (5) reduces to the PDE of [Halperin and Itkin, 2025]. The derivation parallels that of [Grasselli and Hurd, 2007], with the distortion power of the Cole-Hopf transformation generalized from γ to $\gamma(1 - \rho^2)$ in the manner of [Henderson and Hobson, 2002; Musiela and Zariphopoulou, 2004]; details are collected in Appendix D. The indifference price follows from the pair (C, C^0) as in [Halperin and Itkin, 2025].

The full parameter vector is

$$\Theta_{\text{full}} = (\sigma, \sigma_y, \sigma_z, k, \mu, g, \hat{\theta}, c_x, c_y, b_1, b_2, \bar{y}, \gamma, y_0, \theta_0, \bar{\epsilon}, \rho, \rho_{x\theta}), \quad \dim \Theta_{\text{full}} = 18. \quad (7)$$

The remainder of the paper is a controlled demolition of this list.

2.1 Scaling of the memory variable

The memory variable y has no intrinsic scale: it is a latent bookkeeping device for past flows, and neither the data nor the payoff refer to its units. This intuition is made precise by the following observation. Consider a generalized model in which the two occurrences of the coupling c in Eq. (1) carry separate constants, c_x in the x -drift and c_y in the y -drift. Define, for $\lambda > 0$, the transformation

$$y \rightarrow \lambda y, \quad (\sigma_y, \bar{y}, b_2, y_0, c_x, c_y) \rightarrow (\lambda\sigma_y, \lambda\bar{y}, \lambda b_2, \lambda y_0, \frac{c_x}{\lambda}, \lambda c_y). \quad (8)$$

Proposition 1. *Under Eq. (8) the law of the observable process x_t , and hence every option price, is invariant. The correlations ρ and $\rho_{x\theta}$ are dimensionless and remain unchanged. Only the product $c_x c_y$, the correlations, and the ratios b_2/σ_y , $\bar{y}/\sigma_y$, y_0/σ_y , μ are observable.*

Proof. See Section A. □

The original specification imposes $c_x = c_y = c$, which formally breaks the gauge. The breaking is weak rather than benign: the constrained transformation maps the model onto a nearby one with $c_x c_y$ preserved, and in practice this produces a long, shallow ridge in the calibration objective along which $(\sigma_y, \bar{y}, b_2, y_0, c)$ trade off against each other with almost no change in option prices. A global optimizer perceives that ridge as a continuum of near-degenerate minima, and because the degeneracy is only approximate it cannot be removed by reparameterization.

We therefore keep c_x and c_y independent. This costs one nominal parameter and buys an exact symmetry in its place, which is the better trade: an exact redundancy is eliminated once and for all by a normalization, whereas an approximate one has to be fought by the optimizer at every calibration. The effective dimension is reduced by one, to 17, since the gauge is removed by the normalization

$$\sigma_y = 1. \quad (9)$$

Equivalently, we measure the memory variable in units $\tilde{y}_t = y_t/\sigma_y$, so that the normalized memory variable has unit instantaneous diffusion coefficient. In what follows, we denote this normalized variable again by y and set $\sigma_y \equiv 1$ in the PDE. All remaining y -related parameters are understood in these normalized units. This eliminates one parameter and, more importantly, removes the exact scaling ridge from the calibration.

Also, with the specification Eq. (4), the model is invariant under the discrete map

$$(\theta, \hat{\theta}, \theta_0, b_2, \rho_{x\theta}) \rightarrow (-\theta, -\hat{\theta}, -\theta_0, -b_2, -\rho_{x\theta}), \quad (10)$$

since $\cos$ is even and $\sin$ is odd. Every minimum of the objective therefore has a mirror twin. We fix the fundamental domain by imposing $b_2 \geq 0$. This is not a reduction of the parameter count, but it halves the search volume and removes spurious multimodality from the posterior.

2.2 Risk aversion, hidden-factor volatilities, and the role of ρ

Inspection of Eq. (5) shows that the risk-aversion parameter γ enters the nonlinear part of the pricing operator only through the combinations

$$p_y = \gamma\sigma_y^2(1 - \rho^2), \quad p_\theta = \gamma\sigma_\theta^2(1 - \rho_{x\theta}^2), \quad q = \gamma\sigma^2. \quad (11)$$

The spot volatility σ is stiff: it controls the diffusion of the traded coordinate and is pinned by the overall level of option prices. The hidden-factor volatilities σ_y, σ_θ , in contrast, affect prices only through the diffusion of unobservable coordinates and through the nonlinear terms built on Eq. (6). In our experiments the objective is nearly flat along curves of constant (p_y, p_θ) , which identifies these products as the stiff coordinates of the risk-adjustment block. The correlation ρ does not merge into p_y , because it also enters the pricing operator linearly, through the cross-diffusion term $\rho\sigma\sigma_y\partial_{xy}^2 C$ and through the drift adjustment $\rho\sigma_y\bar{\mu}_x/\sigma$. These linear occurrences generate skew and keep ρ separately identifiable.

Two consequences follow. First, γ itself cannot be calibrated from option prices and should be fixed externally. Option-implied risk-aversion estimates are available in the literature; [Bliss and Panigirtzoglou, 1999] report values of order one to ten for index options, and we take γ from that range as an exogenous input. Second, with γ exogenous and the normalization Eq. (9) in force, the combination $p_y = \gamma(1 - \rho^2)$ is a deterministic function of ρ and carries no separate degree of freedom. The free coordinates of the risk-adjustment block are then (σ, ρ, p_θ) , and in the averaged model of Section 3, where p_θ disappears together with the θ dimension, they are (σ, ρ) alone. We note in passing that in the uncorrelated model the same logic forces $p_y = \gamma$ to be fixed, so the risk-adjustment block there consists of σ only. The correlation thus restores to the diffusion structure a degree of freedom that identifiability removes from the risk-aversion channel.

2.3 A priori fixing of non-financial parameters

Three entries of Eq. (7) are not financial parameters at all.

The regularizer $\bar{\epsilon}$.

The constant $\bar{\epsilon}$ prevents divergence of the drift at $x \rightarrow -\infty$ and shapes the potential only in the deep-default region. Our experiments in [Halperin and Itkin, 2025] showed that option prices are highly sensitive to it, which is precisely why it must not float freely in a least-squares objective: it acts as an uncontrolled volatility knob in a region the data do not probe directly. It is a modelling artefact rather than a financial quantity, so it should be fixed by a bound on something the model asserts, not calibrated.

What $\bar{\epsilon}$ controls is the strength of the restoring force at depth. By Eq. (B.3) the drift of the log-price at a reference depth $x_\star$ is $c_x y |V'_M(x_\star)|$, which at $y = \bar{y}$ is $c_x \bar{y} D$ with

$$D \equiv |V'_M(x_\star)| = \frac{\text{drift}_{\max}}{c_x \bar{y}}. \quad (12)$$

The input is therefore a ceiling on the annualized drift the model may impose on a deeply distressed name, which is an economic statement and is monotone in D by construction. A ceiling of two per annum, itself already an extreme restoring force, gives D of order five at representative couplings. Inverting the criterion,

$$\bar{\epsilon} = \frac{1}{1 + De^{x_\star}} - \frac{e^{x_\star}}{g}, \quad x_\star = -4, \quad (13)$$

so $\bar{\epsilon}$ becomes a deterministic function of (g, D) . The reference depth is not arbitrary: it is the point at which the two admissible regularizers coincide exactly, so Eq. (13) holds for either and $\bar{\epsilon}$ carries the same meaning in the model and in the pricer. Appendix B gives the derivation, the admissible range $g > e^{x_\star}(1 + De^{x_\star})$, and explains why neither the turning point of V_M nor the model-implied default rate can serve as the criterion in its place.

Initial hidden states.

The initial values y_0, θ_0 are unobservable and, at a single option snapshot, nearly redundant with the mean-reversion levels. We set them to those levels,

$$\theta_0 = \hat{\theta}, \quad y_0 = \hat{y} \equiv \frac{h(\hat{\theta})}{\mu} + \bar{y} - \frac{c}{\mu} V_M(x_0), \quad (14)$$

consistent with the stationarity assumption underlying the averaging of Section 3. This removes two parameters. The analogy with the Heston initial variance v_0 invoked in [Halperin and Itkin, 2025] is imperfect: v_0 is stiff because it sets the short-end implied volatility level, whereas y_0, θ_0 act on prices only through the drift and are soft. After Sections 2 and 2.3 the count stands at $17 - 1 - 1 - 1 - 2 = 12$, where the gauge normalization Eq. (9) has already reduced 18 to 17 and the four remaining subtractions are the coupling identification through the observable product $c_x c_y$, the exogenous fixing of γ , the regularizer, and the two initial states.

3 Adiabatic elimination of the signal

3.1 Stationary averaging

The signal θ_t is an OU process with relaxation time $1/k$. The calibrated values of k in [Halperin and Itkin, 2026; Halperin and Itkin, 2025] range from 1.3 to 2.7, so for maturities beyond a few months $kT \gtrsim 1$ and the signal explores its stationary law

$$\theta_\infty \sim \mathcal{N}(\hat{\theta}, v), \quad v = \frac{\sigma_z^2}{2k}, \quad (15)$$

within the life of the option. Standard stochastic averaging [Pavliotis and Stuart, 2008] then replaces $f(\theta_t)$ and $h(\theta_t)$ in the drifts of Eq. (1) by their stationary expectations, with corrections of order $1/(kT)$. For the trigonometric specification Eq. (4) the averages are closed form:

$$\mathbb{E}[\cos \theta_\infty] = e^{-v/2} \cos \hat{\theta}, \quad \mathbb{E}[\sin \theta_\infty] = e^{-v/2} \sin \hat{\theta}, \quad (16)$$

so that

$$\bar{f} = b_1 e^{-v/2} \cos \hat{\theta}, \quad \bar{h} = b_2 e^{-v/2} \sin \hat{\theta}. \quad (17)$$

The five signal parameters $(b_1, b_2, k, \hat{\theta}, \sigma_z)$ enter the averaged dynamics only through the two constants $(\bar{f}, \bar{h})$. The averaged model reads

$$dx_t = [\bar{f} + \bar{\eta} - c_x y_t V'_M(x_t)] dt + \sigma_x(y_t) dW_t^{(x)},$$

$$dy_t = [\bar{h} + \mu(\bar{y} - y_t) - c_y V_M(x_t)]dt + dW_t^{(y)}, \quad d\langle W^{(x)}, W^{(y)} \rangle_t = \rho dt, \quad (18)$$

where Eq. (9) is in force. Note that $\bar{h}$ and $\mu\bar{y}$ enter the y -drift only through the sum $\bar{h} + \mu\bar{y}$, so one further parameter is absorbed: we set $\bar{h} = 0$ without loss of generality and let $\bar{y}$ carry the combined level. The volatility $\sigma_x(y)$ is specified in Section 3.2.

3.2 Memory-dependent volatility

In the original specification the memory variable acts on the price through the drift alone, and σ is constant. That choice has a consequence which is easy to miss and which Proposition 2 makes precise: it renders the model's European option prices independent of every parameter governing the memory. We therefore let the volatility depend on the memory variable,

$$\sigma_x(y) = \sigma_0 \sqrt{1 + \alpha(y - \bar{y}) + \beta(y - \bar{y})^2}, \quad \beta > 0, \quad \alpha(y - \bar{y}) + \beta(y - \bar{y})^2 > -1, \quad (19)$$

the constraint being exactly what keeps the radicand positive for all real y and the diffusion non-degenerate. The centring at $\bar{y}$ makes σ_0 the volatility at the resting point of the y -dynamics rather than at the arbitrary origin, so that α is a pure asymmetry parameter and does not mix with the level.

Each coefficient carries a distinct economic reading. At $y = \bar{y}$ the market sits at its equilibrium: no accumulated memory, no directional stress, an order book at full depth, and volatility takes its baseline value σ_0 driven by background noise alone. The linear coefficient α governs the asymmetric response to displacement. When y tracks downward displacement, a negative α raises the local volatility as the market sells off, which embeds the leverage effect in the path itself rather than imposing it through a correlation, and which controls the steepness of the out-of-the-money put skew. The quadratic coefficient β penalizes extreme states of either sign. A large absolute displacement, rally or crash, means a stretched market in which liquidity providers withdraw, the book thins, and the price impact of trades rises; βy^2 scales volatility up accordingly and sets the convexity of the smile.

Under the scaling gauge Eq. (8) the new coefficients transform as $\alpha \rightarrow \alpha/\lambda$ and $\beta \rightarrow \beta/\lambda^2$, so Eq. (9) fixes them along with the rest and no new gauge freedom is introduced.

Proposition 2 (Constant volatility makes the flow invisible to vanillas). *Let σ_x be constant and let the claim have payoff $\mathcal{V}(x_T)$ depending on the log-price alone. Then the indifference price $\pi^\mathcal{V} = C - C^0$ is independent of c_x , c_y , μ , g , γ and ρ , and solves the Black-Scholes equation with volatility σ .*

Proof. See Section C. □

The proposition is not a defect of the derivation, which Section C verifies step by step, but a property of a model in which the flow enters only the drift of a traded asset. The drift of a traded asset is hedged away by the optimal position, and with σ constant no second channel remains. Making σ_x depend on y supplies that channel: the volatility of the traded asset is not hedgeable by holding it, so the memory dynamics reach the price, and the market is incomplete in a way that a vanilla claim can see.

The correlation $\rho_{x\theta}$ disappears at leading order together with the θ dimension. It re-enters only through the first-order correction in $1/k$, which is precisely the mechanism by which correlation generates skew in fast mean-reverting stochastic volatility asymptotics [Fouque et al., 2011]. We therefore set $\rho_{x\theta} = 0$ in the baseline reduced model. Should a first-order correction be developed, that literature indicates that $\rho_{x\theta}$ would enter through a single effective group parameter combining it with the signal amplitudes, which is consistent with the philosophy of the present reduction.

3.3 Transient correction and the term structure

For short maturities the stationarity assumption degrades. A refinement that costs little in the parameter count replaces the stationary averages by the conditional expectations along the deterministic relaxation of the signal. With $\theta_t | \theta_0 \sim \mathcal{N}(m_t, v_t)$,

$$m_t = \hat{\theta} + (\theta_0 - \hat{\theta})e^{-kt}, \quad v_t = \frac{\sigma_z^2}{2k}(1 - e^{-2kt}), \quad (20)$$

one obtains the deterministic time-dependent coefficients

$$\bar{f}(t) = b_1 e^{-v_t/2} \cos m_t, \quad \bar{h}(t) = b_2 e^{-v_t/2} \sin m_t. \quad (21)$$

Under the choice Eq. (14) the transient simplifies, $m_t \equiv \hat{\theta}$, and Eq. (21) becomes a smooth deterministic ramp of the effective amplitude from $\bar{f}(0) = b_1 \cos \hat{\theta}$ at $t = 0$ to $\bar{f}$ at $t \gg 1/k$,

$$\bar{f}(t) = \bar{f}(0) \exp\left(-\frac{v}{2}(1 - e^{-2kt})\right), \quad v = \frac{\sigma_z^2}{2k}. \quad (22)$$

The ramp is controlled by the pair (k, v) . Since v is estimated from the underlying time series in [Halperin and Itkin, 2026] and belongs to the physical-measure block of Section 6.3, retaining the ramp adds a single option-facing shape parameter, the relaxation rate k .

This point deserves emphasis, because it changes the character of the calibration. The baseline averaged model of Eq. (18) carries maturity-independent coefficients, and one parameter set prices the whole surface. When short maturities are included, the ramp Eq. (22) supplies the term structure of the effective drift at the cost of one parameter. In either case the model is fit globally, across all maturities simultaneously. A per-maturity refit, natural for reduced-form smile parameterizations, would forfeit exactly the structural content that distinguishes the Marketron from them, and we do not use it.

3.4 The reduced pricing PDE

Since θ has been eliminated, the certainty equivalent $C(t, x, y)$ solves the two-dimensional analogue of Eq. (5),

$$\frac{\partial C}{\partial t} + \frac{1}{2}\sigma_x^2(y)\frac{\partial^2 C}{\partial x^2} + \rho\sigma_x(y)\frac{\partial^2 C}{\partial x\partial y} + \frac{1}{2}\frac{\partial^2 C}{\partial y^2} + \bar{\eta}(y)\frac{\partial C}{\partial x} + \left(\mu_y - \rho\frac{\bar{\mu}_x}{\sigma_x(y)}\right)\frac{\partial C}{\partial y} + \frac{p_y}{2}\left(\frac{\partial C}{\partial y}\right)^2 - \frac{\bar{\mu}_x^2}{2\gamma\sigma_x^2(y)} - rC = 0, \quad (23)$$

with $p_y = \gamma(1 - \rho^2)$, $\bar{\eta}(y) = r - \sigma_x^2(y)/2$, $\bar{\mu}_x = \bar{f} + \bar{\eta}(y) + \sigma_x^2(y)/2 - r - c_x y V'_M(x)$ and $\mu_y = \mu(\bar{y} - y) - c_y V_M(x)$. Both the diffusion and the drift in x now carry y , which is what breaks the degeneracy of Proposition 2. Note that Eq. (19) does not disturb the two cancellations underlying Eq. (5): the elimination of $s\bar{\mu}_x v_s$ by the optimal hedge and of $(\partial_x C)^2$ by the distortion are both pointwise in y , so they proceed unchanged with σ replaced by $\sigma_x(y)$ throughout.

The reduction from three hidden dimensions to the two-dimensional state (x, y) substantially enlarges the range of numerical methods that can be used for Eq. (23). Unlike the full three-dimensional problem of [Halperin and Itkin, 2025], there is no need to commit to a particular discretization at the level of the model itself. Several complementary approaches are available, and they are useful for different purposes.

A natural high-accuracy approach is based on Strang splitting of the 2D operator into the x -diffusion/drift part, the y -diffusion/drift part, and the mixed derivative. The latter can be treated by the positivity-preserving construction (the Diagonal Frog method) developed in [Itkin, 2026b; Itkin, 2026c]. This is particularly useful here because the mixed term $\rho\sigma_x(y)C_{xy}$ is not sign-definite under a naive explicit discretization, whereas the construction of [Itkin, 2026b] provides a stable treatment that preserves positivity under the corresponding admissibility conditions.

For the x -subproblem, y is frozen during the split step. Hence $\sigma_x(y)$ and all coefficients depending on y become parameters of a one-dimensional equation in x . The resulting equation contains the source term $-\frac{\bar{\mu}_x^2}{2\gamma\sigma_x^2(y)}$ in addition to the x -diffusion and drift. Because the volatility is quadratic in y through Eq. (19), the coefficients are nevertheless constant with respect to x on each fixed y -slice. The corresponding Green's function is therefore available in closed form, and the source contribution can be incorporated analytically. This provides a direct extension of the Gaussian/RBF machinery used in [Halperin and Itkin, 2025], although the resulting coefficients are now indexed by the y grid.

The y -subproblem contains the nonlinear gradient-square term

$$\frac{p_y}{2}(C_y)^2, \quad p_y = \gamma(1 - \rho^2). \quad (24)$$

This term can be removed exactly by the Cole–Hopf transformation. In particular, consider the y -subproblem in isolation,

$$C_\tau = \frac{1}{2}C_{yy} + b(y)C_y + \frac{p_y}{2}(C_y)^2, \quad b(y) = \mu_y - \rho \frac{\bar{\mu}_x}{\sigma_x(y)}. \quad (25)$$

Under $w = \exp(p_y C)$, the quadratic-gradient term cancels exactly against the contribution generated by the diffusion term, giving the linear equation

$$w_\tau = \frac{1}{2}w_{yy} + b(y)w_y. \quad (26)$$

Thus the y -diffusion, y -drift, and quadratic-gradient terms can be treated together as a single linear parabolic subproblem. Since this equation has bounded coefficients and positive initial data, the maximum principle preserves positivity of w and prevents the Cole–Hopf variable from developing a finite-time singularity. The corresponding inverse transformation,

$$C = \frac{1}{p_y} \log w, \quad (27)$$

therefore provides a stable treatment of the nonlinear y -subproblem.

This observation also determines how the operator splitting must be implemented. The C_{yy} term should not first be consumed by an independent diffusion step and subsequently be invoked again in the Cole–Hopf transformation. Instead, the y -diffusion, y -drift, and quadratic-gradient term are assigned consistently to the same split subproblem. A convenient implementation is to solve the transformed linear equation implicitly, using a backward-Euler tridiagonal solve in y for each fixed x . To control the numerical range of the exponential and logarithm, we shift C by its maximum over the y grid before applying the Cole–Hopf transformation. Since the linear y -operator is invariant under addition of a constant to C , this shift is exact and does not alter the result after the inverse transformation.

The RBF discretization of Appendix F can also be used as a spatial approximation of the resulting one- or two-dimensional linear subproblems. However, the closed-form RBF update derived for the original operator should not be interpreted as applying unchanged to the complete 2D PDE. The splitting must respect the assignment of the y -diffusion and quadratic-gradient terms to the Cole–Hopf subproblem.

For the numerical experiments in this paper we use a split finite-difference implementation tailored to the stability issue that arises at low $\sigma_x(y)$. A fully explicit discretization of Eq. (23) becomes unstable in this regime, with the instability becoming particularly pronounced for sufficiently large c_y . This behavior is numerical rather than structural: the continuous y -subproblem is transformed by Cole–Hopf into a linear advection-diffusion equation and hence does not exhibit the corresponding finite-time blow-up, [Halperin and Itkin, 2025].

This split formulation removes the problematic y -advection CFL restriction from the explicit step and, more importantly, prevents the low- $\sigma_x(y)$ instability associated with treating the quadratic-gradient and y -diffusion terms explicitly. In numerical tests, the split scheme agrees with the fully explicit scheme in

parameter regions where the latter is stable, while remaining stable in parameter regimes where the explicit scheme develops numerical blow-up. In particular, at the calibrated parameter values the two schemes produce option prices agreeing to within approximately 1.6×10^{-6} across the tested strikes. A sweep in c_y at $\sigma_0 = 0.121$ remains smooth up to $c_y = 3$, whereas the fully explicit scheme becomes unstable above approximately $c_y = 0.2$. The split scheme also remains stable at $c_y = 0.2$ for substantially smaller volatility levels, including $\sigma_0 = 0.09$, 0.07 , and 0.05 . Put–call parity is satisfied to numerical precision, with discrepancies of order 10^{-17} in these tests.

The purpose of this implementation is not to construct a production-quality pricing engine, but to provide a transparent and computationally efficient solver with which the calibration and identifiability experiments can be repeated reliably. The split method retains the simplicity of finite differences for the multidimensional part of the problem while treating the numerically problematic nonlinear subproblem through an exact Cole–Hopf linearization. Consequently, it is sufficiently robust for the parameter ranges considered in [Section 6](#), without requiring the more elaborate RBF or positivity-preserving machinery described above.

The reduction from the original (x, y, θ) problem to the two-dimensional (x, y) problem remains the main computational gain. Regardless of which of the above discretizations is adopted, the collocation or finite-difference grid now contains only $N_x N_y$ spatial degrees of freedom rather than $N_x N_y N_\theta$. More importantly, the numerical problem is no longer burdened by the fast signal dimension that was responsible for much of the cost of the original pricing calculation. The different schemes described above should therefore be viewed as alternative numerical realizations of the same reduced pricing equation, rather than as additional modelling assumptions.

Parameter count

The reduced parameter vector is

$$\Theta_{\text{red}} = (\sigma_0, \alpha, \beta, \mu, g, c, \bar{y}, \bar{f}, \rho), \quad \dim \Theta_{\text{red}} = 9, \quad (28)$$

with γ exogenous, $p_y = \gamma(1 - \rho^2)$ derived, $\bar{\epsilon} = \bar{\epsilon}(g, D)$ from [Eq. \(13\)](#), and $\bar{\eta}(y)$ tied to r through $\sigma_x(y)$. The two volatility coefficients of [Eq. \(19\)](#) are the price of making the flow visible to option prices at all, and they are not duplicates of existing freedoms: α controls skew and β convexity, neither of which the drift channel can produce. When short maturities require the transient ramp [Eq. \(22\)](#), the rate k joins the list and the count is ten. These parameters serve the entire surface. [Table 1](#) summarizes the reduction.

Remark on the flow-impact couplings in the calibrated model.

The reduced dynamics in [Eq. \(18\)](#) and the pricing PDE [Eq. \(23\)](#) are written with independent couplings c_x and c_y , consistent with the exact scaling gauge identified in [Section 2.1](#). The subsequent analysis, however, and in particular the parameter count in [Eq. \(28\)](#) and the numerical calibration of [Section 6.3](#), employs a single coupling constant c . We now state explicitly what this compression entails.

Setting $c_x = c_y = c$ in [Eq. \(18\)](#) and [Eq. \(23\)](#) yields the model that is actually calibrated. This is a deliberate simplification made at the final stage, after the other reductions have already been imposed. The gauge argument of [Section 2.1](#) establishes that the observable process, and hence every option price, depends on the couplings only through the invariant product $c_x c_y$. In the fully reduced, gauge-fixed model, the independent rescalings that previously created a flat direction in the calibration objective have already been absorbed: σ_y is normalized to unity by [Eq. \(9\)](#), the initial states are frozen by [Eq. \(14\)](#), and the signal block is eliminated. Consequently, the residual invariance is isolated to the single product $c_x c_y$, and the redundancy associated with separating the two factors is, at this stage, a pure normalization freedom that carries no economic content.

Retaining separate couplings in the final parameter list would therefore add a nominal degree of freedom without any corresponding gain in identifiable structure. The single constant c in [Eq. \(28\)](#) should be

Step	Mechanism	Parameters removed
Gauge fixing (Sec. 2.1)	$\sigma_y = 1$	1
Coupling identification (Sec. 2.1)	only $c_x c_y$ observable, $(c_x, c_y) \rightarrow c$	1
Risk aversion (Sec. 2.2)	γ fixed exogenously	1
Regularizer (Sec. 2.3)	$\bar{\epsilon} = \bar{\epsilon}(g, D)$	1
Initial states (Sec. 2.3)	$\theta_0 = \hat{\theta}, y_0 = \hat{y}$	2
Signal averaging (Sec. 3)	$(b_1, b_2, k, \hat{\theta}, \sigma_z) \rightarrow (\bar{f}, \bar{h})$	3
Level absorption (Sec. 3)	$\bar{h}$ into $\bar{y}$	1
Cross-correlation (Sec. 3.2)	$\rho_{x\theta} \rightarrow 0$ (order $1/k$)	1
Volatility coefficients (Sec. 3.2)	α, β introduced by Eq. (19)	-2
Total	$18 \rightarrow 9$	9

Table 1: *Parameter reduction ledger. The intermediate three-dimensional risk block retains (σ, ρ, p_θ) as free coordinates; p_θ vanishes with the θ dimension under averaging, its constituents σ_z and $\rho_{x\theta}$ being removed by the signal-averaging and cross-correlation rows respectively, with $\rho_{x\theta}$ entering only at order $1/k$. The two volatility coefficients α, β of Eq. (19) are then introduced, giving $\dim \Theta_{\text{red}} = 9$ in Eq. (28). In the reduced model $p_y = \gamma(1 - \rho^2)$ is derived, not calibrated. The optional term-structure parameter k of Eq. (22) is not counted; including it raises the count to ten.*

interpreted as the gauge-fixed representative of the product $c_x c_y$, under the symmetric normalization $c_x = c_y$. The cost of this choice is the reintroduction of a mild, isolated softness along the c -coordinate, which is distinct from the continuous degeneracy of the full model. This softness is explicitly diagnosed by the eigenspectrum and marginal variances reported in Section 5, and it is handled in Section 6.3 by retaining c as a risk-neutral parameter whose option-implied value is reported together with its profile uncertainty. The wedge between this value and its physical-measure counterpart is then interpreted as the market price of flow risk, with the associated estimation error carried through to the economic conclusion.

4 The Marketron as a generalized Langevin equation

The reduction presented in Section 3 yields a two-dimensional system governed by the log-price and the memory variable. Before developing its implications, we briefly translate the terminology we shall use into the language of financial mathematics.

A *generalized Langevin equation* (GLE) is, mathematically, a non-Markovian stochastic Volterra equation: the drift of the state depends on the entire history of its increments through a *memory kernel*. In the context of market microstructure, the kernel is the propagator of price impact. It measures how a past flow innovation continues to distort the price drift at a later time.

The *effective potential* is the stationary drift distortion induced by the accumulated memory, while the *fluctuation-dissipation relation* is the equilibrium condition that links the decay rate of the impact propagator to the autocorrelation time of the exogenous noise. When this link is violated, the system is termed *athermal* or, in the physics literature, *active matter*.

In finance, this corresponds to a regime where the mean-reversion of order-flow shocks and the persistence of return predictors are governed by distinct time scales. Such non-Markovian dynamics have recently entered mainstream quantitative finance through the rough-volatility literature [Abi-Jaber and El-Euch, 2019; Abi Jaber, Larsson, and Pulido, 2019; Bayer and Breneis, 2023], where singular kernels are approximated by sums of exponentials; the Marketron provides an exact finite-dimensional embedding for a specific state-dependent kernel.

With this dictionary in place, the reinterpretation of the reduced system proceeds as follows. By eliminating the memory variable y rather than the signal θ , we demonstrate that the Marketron is exactly a one-dimensional overdamped GLE for the log-price, driven by a memory kernel we compute in closed form. Consequently, y_t is not an independent latent state introduced *ad hoc*, but the exact Markovian lift of a non-Markovian price-impact process.

This mapping is not merely interpretive: it shows that the absence-of-arbitrage condition on the kernel matrix, derived independently in [Itkin, 2026a], coincides with the market-price-of-risk terms generated by exponential-utility indifference pricing. Moreover, it delivers a testable parametric condition, the equality of the memory relaxation rate μ and the signal relaxation rate k , that distinguishes equilibrium dynamics from a driven, athermal market.

The reader who prefers to avoid physics terminology may read **GLE** as stochastic Volterra equation, **kernel** as impact propagator, and **fluctuation-dissipation** as equilibrium consistency condition. The mathematics that follows is identical.

This formulation yields three primary insights. First, the kernel occupies the specific entry of the kernel matrix that the absence of arbitrage eliminates under the pricing measure. This theoretically aligns the Marketron's flow mechanism with the market price of memory risk established in [Itkin, 2026a]. Second, the kernel is state-modulated, situating the model outside the standard GLE class and precisely delineating the analytical boundaries of both frameworks. Finally, the fluctuation-dissipation relation, which distinguishes equilibrium from active dynamics, simplifies within the Marketron to an equality between two rates explicitly derived during calibration. This transforms the active-matter reading of the model (see Section 4.4) into an empirically verifiable parametric condition.

4.1 The Kernel formulation

Recalling the reduced system from Eq. (18):

$$\begin{aligned} dx_t &= [\bar{f} + \bar{\eta} - c_x y_t V'_M(x_t)] dt + \sigma_x(y_t) dW_t^{(x)}, \\ dy_t &= [\bar{h} + \mu(\bar{y} - y_t) - c_y V_M(x_t)] dt + \sigma_y dW_t^{(y)}. \end{aligned} \quad (29)$$

The memory variable influences the price drift strictly via the product $y_t V'_M(x_t)$, while its own drift relies on the price solely through $V_M(x_t)$. Because the y -equation is linear with constant coefficients, it can be explicitly solved and substituted, thereby eliminating y from the system dynamics entirely.

Proposition 3. *The system Eq. (29) is an exact Markovian lift of a one-dimensional overdamped GLE for x_t ,*

$$\int_0^t K_M(t, s) \dot{x}_s ds = -U'_{\text{eff}}(x_t) + \eta_t, \quad (30)$$

with the state-dependent kernel

$$K_M(t, s) = \kappa \delta(t - s) + \frac{c_x c_y}{\mu} V'_M(x_t) V'_M(x_s) e^{-\mu(t-s)} + \mathcal{O}(\sigma^2), \quad \kappa = 1, \quad (31)$$

the effective potential

$$U_{\text{eff}}(x) = -(\bar{f} + \bar{\eta})x + c_x \hat{y} V_M(x) - \frac{c_x c_y}{2\mu} V_M(x)^2, \quad \hat{y} \equiv \bar{y} + \frac{\bar{h}}{\mu}, \quad (32)$$

and a noise η_t that is the sum of $\sigma \dot{W}_t^{(x)}$ and the multiplicative term $-c_x \sigma_y V'_M(x_t) \int_0^t e^{-\mu(t-s)} dW_s^{(y)}$. The $\mathcal{O}(\sigma^2)$ remainder in Eq. (31) is the state-dependent memory correction isolated in Appendix E. It vanishes when V_M is linear and marks Eq. (31) as the leading-order form of the kernel.

Proof. See Appendix E. □

The absence of an inertial term stems from the price equation in Eq. (29) being first-order with respect to x , establishing a direct correspondence with the overdamped regime. Instantaneous friction manifests as the atom $\kappa\delta$ of Eq. (31), where $\kappa = 1$ by the normalization of the $\dot{x}_t$ coefficient in $dx_t = \mu_x dt + \sigma dW_t^{(x)}$.

The structure of Eq. (31) warrants specific attention. The kernel depends on more than just the temporal lag. It incorporates the modulation $V'_M(x_t)V'_M(x_s)$ evaluated at both endpoints of the memory integral. Thus, the influence of historical states on the current state is contingent upon the price trajectory. Under the market potential defined in Eq. (3), this modulation varies exponentially with x and exhibits a sign change at the critical threshold $\hat{x}$, where $V'_M(\hat{x}) = 0$. Economically, this translates to prominent memory effects within the default region and attenuated memory elsewhere, effectively embedding the model's economic rationale into the kernel's architecture.

Notably, the effective potential U_{eff} in Eq. (32) diverges from the underlying market potential. It incorporates a linear tilt derived from the drift constants, a $c_x\hat{y}V_M$ component reflecting the uncoupled relaxation level of y , and a negative quadratic term $-\frac{c_x c_y}{2\mu}V_M^2$ induced by the system coupling. This feedback mechanism between price and memory deepens the effective potential well relative to its uncoupled state. Consequently, extracting the market potential directly from a stationary density would erroneously yield U_{eff} rather than V_M .

Remark 1 (The atom and the short end). The atom in Eq. (31) serves a distinct structural purpose. A kernel comprising $\gamma\delta$ plus a slow tail perfectly aligns with the configuration identified in [Itkin, 2026a] for generating a finite zero-maturity at-the-money skew: a regular semimartingale short end coupled with a memory-carrying tail. Conversely, a kernel exhibiting a power-law behavior at the origin would produce a skew diverging as T^{β_0-1} at short maturities. The Marketron inherently possesses the short-end regularity required for a finite skew without artificial imposition. It features a single relaxation mode in the tail (indicative of short memory) while Section 4.3 outlines the architecture for a multi-mode extension. △

Remark 2 (Kernel confirmation of Proposition 1). The coupling constants influence Eq. (31) exclusively through the product $c_x c_y$, which remains invariant under Eq. (8), unlike its individual factors. This independently corroborates Proposition 1, which posits that only $c_x c_y$ is observable based on the invariance of the law of x_t . The kernel representation achieves this conclusion via an alternate route, as K_M intrinsically governs the reduced x -dynamics and must remain independent of gauge-variant quantities. When the constants appear separately, such as in the $c_x\hat{y}V_M$ tilt of Eq. (32), the accompanying factor is gauge-covariant, rendering the combination invariant. △

Remark 3 (Kernel matrix positioning). Within the two-dimensional GLE framework of [Itkin, 2026a], featuring state $Z = (X, Y)^\top$ and kernel matrix $\mathbf{K}$, the K_{XX} entry dictates the dependence of the log-price drift on its historical increments. The tail in Eq. (31) scales $\dot{x}_s$ within the x_t equation, formally corresponding to this entry, while the atom represents the standard instantaneous friction of an overdamped system. Under the pricing measure, the absence of arbitrage dictates $K_{XX} \equiv 0$, prompting the first row of $\mathbf{K}$ to manifest in the measure change as the market price of memory risk. The Marketron arrives at the equivalent construct via exponential-utility indifference pricing with risk aversion γ . The convergence of these two distinct methodologies serves as a robust theoretical validation for both frameworks. △

4.2 Framework boundaries and intersections

Proposition 3 might imply that the Marketron is a constrained instance of the GLE class presented in [Itkin, 2026a], or vice versa. However, neither framework fully subsumes the other, and their divergence clarifies the unique utility of each approach.

The kernel in [Itkin, 2026a] is strictly a function of the lag, deriving its generality from the allowable number of relaxation modes. A completely monotone kernel is expressed as a superposition $K(\tau) = \int_0^\infty e^{-\lambda\tau} \nu(d\lambda)$, and achieving a power law necessitates a continuum of modes. This architecture specifically facilitates long-memory modeling, which is the primary focus of that framework.

In contrast, the kernel in Eq. (31) relies on a single mode, characteristic of short memory with respect to lag. Its primary generalization lies in the state modulation $V'_M(x_t)V'_M(x_s)$, a feature absent in pure lag-dependent kernels. The intersection of these two classes is limited to the single-mode, state-independent case, corresponding to a linearized Marketron around an operating point where V'_M is approximated as constant.

This linearization, however, fundamentally alters the model's behavior. Based on Eq. (3), the modulation V'_M fluctuates by orders of magnitude across the relevant log-price domain and changes sign at $\hat{x}$. Approximating the Marketron with a lag-only GLE eliminates this critical state dependence, thereby erasing the fundamental distinction between the default and normal regions that the model was designed to capture. The reverse substitution is equally flawed: a single exponential function cannot replicate the slowly decaying memory observed in empirical volatility data, rendering the Marketron kernel unsuitable as K_{YY} in that broader context.

4.3 Exact Markovian lifting and long-memory extensions

The construction of finite-dimensional Markovian embeddings for non-Markovian dynamics has a rich precedent in the rough volatility literature. For instance, [Abi-Jaber and El-Euch, 2019] along with [Abi Jaber, Larsson, and Pulido, 2019] proposed the multifactor approximation of fractional Volterra processes. Their approach relies on approximating the singular fractional kernel by a sum of exponential terms, effectively lifting the non-Markovian process into a high-dimensional affine state space. Furthermore, [Bayer and Breneis, 2023] formalizes this methodology using Prony approximations to efficiently match the fractional kernel with a superposition of Markovian modes.

While Eq. (29) could similarly be interpreted as a Markovian approximation of a non-Markovian system, the relationship in our framework is formally exact. Because a single exponential is completely monotone, the GLE in Eq. (30) permits a finite-dimensional Markovian embedding utilizing one auxiliary variable (specifically y_t). The elimination of y in Appendix E is exact; the closed-form kernel Eq. (31) is its stationary, leading-order form, obtained by discarding transients that decay on the scale $1/\mu$ and by absorbing an $O(\sigma^2)$ state-dependent memory term into the noise η_t , neither of which alters the structure of the kernel but neither of which is identically zero at finite t . The primary distinction between the descriptions is purely foundational (specifically, whether the memory variable or the kernel is defined as the primitive mathematical object).

This relationship also provides the theoretical blueprint for integrating long memory into the Marketron. Expanding the single memory variable into a family $y_t^{(i)}$ (each characterized by distinct relaxation rates μ_i and couplings c_i , governing the second line of Eq. (29) and additively influencing the price drift) yields:

$$K_M(t, s) = \kappa \delta(t - s) + V'_M(x_t) V'_M(x_s) \sum_i \frac{c_x c_{y,i}}{\mu_i} e^{-\mu_i(t-s)}. \quad (33)$$

This state-modulated, completely monotone kernel successfully reproduces power-law decay across any finite lag interval. Consequently, introducing long memory to the Marketron requires no fundamental structural changes, only the addition of further memory variables, and the critical state modulation remains fully intact. However, this extension incurs a parametric penalty, as each mode introduces an additional rate and coupling constant. Given that mitigating parameter proliferation is a central objective of this framework, such an extension should be reserved strictly for applications where empirical data demand it.

4.4 Fluctuation-dissipation and the active-matter interpretation

Throughout this section we use the statistical-mechanics terminology of GLEs alongside its translation into the language of non-Markovian financial dynamics. The correspondence is:

Statistical mechanics	Financial dynamics
Generalized Langevin equation (GLE)	Stochastic Volterra equation
Memory kernel $K_M(t, s)$	Impact propagator / price-impact kernel
Fluctuation-dissipation relation (FDT)	Equilibrium consistency condition
Effective temperature Θ	Noise–kernel proportionality constant
Athermal / active-matter regime	Driven non-equilibrium regime
Internal reservoir	Endogenous order-flow feedback
Quasiparticle	Effective price degree of freedom

Both languages describe the same mathematical objects; the reader may use whichever is more familiar.

Previous discussions characterizing the Marketron as a model of active matter (where the market quasiparticle relies on an internal reservoir rather than purely external forcing) have remained largely interpretive. The GLE representation formalizes this analogy (or, in financial terms, the non-Markovian kernel representation) into a strict parametric condition.

Within a standard GLE framework, the autocovariance of the driving noise, $C(\tau) = \mathbb{E}[\eta_t \eta_{t-\tau}]$, and the memory kernel operate as independent constructs. Equilibrium is defined as the state where these two elements are coupled via the second fluctuation-dissipation relation:

$$C(\tau) = \Theta K(\tau), \quad (34)$$

where Θ represents an effective temperature (in finance, the noise–kernel proportionality constant). Systems violating [Eq. \(34\)](#) are classified as athermal (driven non-equilibrium). Active matter specifically denotes athermal systems where this violation is driven by an internal energy source (endogenous order-flow feedback). This classification has concrete implications: it dictates whether the stationary density corresponds to the Boltzmann weight of the potential, and [\[Itkin, 2026a\]](#) constructs an empirical test based on the ratio between the two sides of [Eq. \(34\)](#).

The Marketron explicitly provides both components of this relation. The kernel tail in [Eq. \(31\)](#) decays at a rate of μ , derived from the relaxation of the memory variable. The coloured component of the forcing is introduced via $f(\theta_t)$ and $h(\theta_t)$, inheriting its correlation time from the underlying signal dynamics [Eq. \(1\)](#). To align these decay rates in accordance with [Eq. \(34\)](#), the model necessitates:

$$k = \mu, \quad (35)$$

alongside an amplitude condition linking (b_1, b_2, σ_z) to (c, σ_y) through Θ .

Because k and μ represent independently identified calibration parameters, [Eq. \(35\)](#) functions as a verifiable constraint rather than a theoretical assumption. Divergent rates confirm the model operates in an athermal (driven non-equilibrium) regime, substantiating the active-matter interpretation with empirical parameter values rather than qualitative analogy. Conversely, identical rates would signify equilibrium, reducing the quasiparticle (effective price degree of freedom) framework to a purely descriptive overlay.

The trigonometric signal.

Equation [Eq. \(35\)](#) aligns the leading exponential decay rates. Because the forcing terms $f(\theta) = b_1 \cos \theta$ and $h(\theta) = b_2 \sin \theta$ are nonlinear transformations of an Ornstein–Uhlenbeck process, their corresponding autocovariance does not exhibit pure exponential decay.

Specifically, for $\theta \sim \mathcal{N}(m, v)$ and any real parameter ω , the characteristic function is given by $\mathbb{E}[e^{i\omega\theta}] = e^{i\omega m - \omega^2 v/2}$, which yields

$$\mathbb{E}[\cos \omega \theta] = e^{-\omega^2 v/2} \cos \omega m, \quad \mathbb{E}[\sin \omega \theta] = e^{-\omega^2 v/2} \sin \omega m. \quad (36)$$

Applying this result with $\omega = 1$ to the stationary law $(m, v) = (\hat{\theta}, \sigma_z^2/2k)$ yields Eq. (17), while application to the conditional law (m_t, v_t) produces Eq. (21). Analogous identities provide the second moments $\mathbb{E}[\cos^2 \theta] = \frac{1}{2}(1 + e^{-2v} \cos 2m)$ required for the source term $\bar{\mu}_x^2$ in Eq. (23). Consequently, the averaged PDE retains fully closed-form RBF matrix elements.

Following these identities, the autocovariance incorporates the factor $\exp[-\sigma_z^2(1 - e^{-k\tau})/2k]$, which simplifies to $e^{-k\tau}$ only under the small-amplitude limit $\sigma_z^2/2k \ll 1$. This confirms that generic parameters inherently produce an athermal (non-equilibrium) model, because unlinked rates are highly unlikely to coincide perfectly. Therefore, the exact equality in Eq. (35) must be interpreted strictly as the leading-order parametric condition.

5 Identifiability analysis

Having established the analytical reduction of the parameter space, we must now address its empirical identifiability. While analytical symmetries successfully collapse the unobservable dimensions, they do not guarantee strict minimality. This section investigates whether the reduced parameter vector Θ_{red} constitutes an irreducible core or retains latent structural redundancies. Using local sensitivity metrics and manifold boundary approximation methods, we evaluate the calibration objective's robustness and confirm the structural integrity of the reduced model.

5.1 Sloppiness of the calibration objective

Sections 2 and 3 established the reduction by construction through the gauge Eq. (8), combinations Eq. (11), and the averaging limit theorem. We omit numerical verification of the original parameterization's sloppiness, as it would merely duplicate the algebraic proofs. Instead, we focus on determining whether Θ_{red} retains any further unidentified structure.

Let $r_i(\Theta) = C_i^{\text{model}}(\Theta) - C_i^{\text{ref}}$ denote the pricing residuals and $J = \partial r / \partial \Theta$ the Jacobian. The Gauss-Newton Hessian $H = J^\top J$ quantifies local identifiability; eigenvalues spanning multiple orders of magnitude signal a sloppy model with soft directions [Gutenkunst et al., 2007; Transtrum, Machta, and Sethna, 2011]. Operating in log-parameters renders the spectrum scale-free and converts any residual scaling symmetry $\Theta_i \rightarrow \lambda^{a_i} \Theta_i$ into a constant direction a . This ensures that any surviving symmetry manifests strictly as an exact null vector of H rather than merely a small eigenvalue.

To avoid artifacts from external calibrations and establish a known ground truth, reference prices are generated at the operating point $\Theta_{\text{red}}^* = (\sigma_0, \alpha, \beta, \mu, g, c, \bar{y}, \bar{f}, \rho) = (0.18, -0.4, 0.15, 2.0, 1.0, 0.08, 0.0, 0.0, -0.3)$, with exogenous risk aversion $\gamma = 3$ and drift ceiling $D = 5$ fixed. The option set is a surface of 54 quotes (six maturities from 0.12 to 1.0 years and nine log-moneyness points from -0.12 to 0.12). The pricing noise ς is set to 0.5% of the price, approximating half a bid-ask width.

The spectrum (Table 2, Figure 1) reveals two key facts. First, every eigenvalue is strictly positive; the empty null space empirically certifies that no exact symmetries survive the reduction. Second, the spectrum separates into two distinct blocks with a gap of over three orders of magnitude between λ_5 and λ_6 . The five stiff eigenvalues form the block resolved by the vanilla surface, with eigenvectors dominated by the smile and skew parameters $(\sigma_0, \alpha, \beta, \mu, \rho)$. The four small eigenvalues correspond to a block the surface barely registers, dominated by the flow and potential parameters $(c, \bar{f}, \bar{y}, g)$. These parameters affect vanilla prices only indirectly through the memory dynamics feeding $\sigma_x(y)$, not via a direct drift channel. The condition

options-facing			drift		
index	λ_i	leading coords.	index	λ_i	leading coords.
1	1.5×10^7	σ_0	6	3.9×10^{-2}	c
2	3.5×10^4	β, α	7	7.7×10^{-4}	$\bar{f}$
3	5.3×10^3	ρ, μ	8	1.9×10^{-7}	$\bar{y}, g$
4	3.7×10^2	μ, ρ	9	4.1×10^{-8}	$g, \bar{y}$
5	8.8×10^1	α, β, ρ			

Table 2: Eigenvalues of the Gauss-Newton Hessian $H = J^\top J$ in log-parameters at Θ_{red}^* , in descending order. The leading coordinates are the parameters whose eigenvector components exceed 0.3 in absolute value. All eigenvalues are strictly positive, so H has an empty null space.




Figure 1: Eigenvalues of the Gauss-Newton Hessian $H = J^\top J$ in log-parameters at Θ_{red}^* , on a logarithmic scale. The five stiff eigenvalues (circles) form the options-facing block. The four small eigenvalues (squares, shaded region) form the drift block that is fixed from the physical measure. The two blocks are separated by more than three orders of magnitude, and no eigenvalue is zero.

number of the full nine-parameter Hessian is 3.7×10^{14} . Restricting H to options-facing coordinates lowers it to 3.99×10^8 including c , and to 1.75×10^5 for the pure smile block $\{\sigma_0, \alpha, \beta, \mu, \rho\}$.

This block structure defines the reduced model’s identifiability and validates the staged calibration of Section 6.3. The residual sloppiness represents a separation of data channels rather than a surviving symmetry: the vanilla surface identifies the smile block, while the drift block $\{c, \bar{f}, \bar{y}, g\}$ is pinned from the physical measure and held fixed during option calibration.

Marginal profile widths, evaluated as $\sqrt{(H^{-1})_{jj}}$ (Table 3, Figure 2), formalize this ranking. They fall into three groups separated by two large gaps. The smile parameters $(\sigma_0, \rho, \alpha, \beta, \mu)$ have widths below 0.55, while the flow coupling c has a width of 13.6. The remaining drift parameters $(\bar{f}, \bar{y}, g)$ exhibit widths in the hundreds to thousands, indicating profiles left entirely flat by the vanilla surface. This precisely justifies fixing the drift block using physical-measure time series.

Profile likelihoods, [Raue et al., 2009] confirm this local perspective. By re-optimizing all other parameters along a continuous scan of each target coordinate, we measure the residual envelope. Profiles are

options-facing				drift (fixed from $\mathbb{P}$ -measure)		
param.	$\sqrt{(H^{-1})_{jj}}$	std (0.5%)	std (0.05%)	param.	$\sqrt{(H^{-1})_{jj}}$	status
σ_0	0.03	0.007	0.003	c	13.6	weak
ρ	0.14	0.174	0.048	$\bar{f}$	1.7×10^2	$\mathbb{P}$ -measure
α	0.27	0.206	0.046	$\bar{y}$	2.8×10^3	$\mathbb{P}$ -measure
β	0.32	0.060	0.011	g	4.7×10^3	$\mathbb{P}$ -measure
μ	0.52	0.932	0.294			

Table 3: Per-parameter identifiability at Θ_{red}^* . Left block: options-facing parameters with marginal standard errors $\sqrt{(H^{-1})_{jj}}$ (log-parameters, scale-free) and empirical standard deviations from multi-start fits at two pricing-noise levels. μ is identified but soft. Right block: drift parameters fixed from the physical measure, not recovered from options.




Figure 2: Marginal standard error of each log-parameter, $\sqrt{(H^{-1})_{jj}}$, on a logarithmic scale. Small means well determined. The five smile parameters are identified from the vanilla surface, the flow coupling c is weak, and $\bar{f}$, $\bar{y}$ and g are left flat by the option data and are fixed from the physical measure. The dashed line marks unit width.

deemed flat if the cost elevation remains below the ambient noise threshold ($\frac{1}{2}\chi_{1,0.95}^2\varsigma^2$ above the optimum). The aforementioned marginal widths reflect the profile curvatures at the optimum, demonstrating bounded profiles for the smile parameters and flat profiles for the drift block. This corroborates the channel separation observed in the eigenspectrum.

To verify practical recovery, we perturb the surface by the pricing noise and refit the smile block from random starts within 15% of truth, holding the drift block fixed. At 0.5% noise, all seven starts converge to the noise floor; at 0.05% noise, seven of eight converge (the single failure originating outside the admissible region). The recovery columns of Table 3 show all recovered means lie within one standard deviation of truth. Crucially, as pricing noise decreases tenfold, every dispersion shrinks: the standard deviation of σ_0 drops from 0.007 to 0.003, β from 0.060 to 0.011, and ρ from 0.174 to 0.048. This noise-proportional contraction is the signature of a well-posed identification devoid of flat directions. The parameter μ shrinks the least (from 0.932 to 0.294), confirming it is the softest identifiable coordinate as predicted by Table 2. This residual softness in μ is thus a function of data resolution, which more maturities or tighter quotes

would reduce, rather than a structural degeneracy.

The placement of ρ in the physical-measure block rather than the risk-neutral block requires justification, since the correlation between flow and return innovations generates short-maturity skew, and therefore appears, at first sight, to be precisely the kind of parameter the option surface should identify. The eigenspectrum in Table 2 supports this expectation: ρ participates in the third and fourth stiff eigenvectors with marginal width 0.14 in Table 3. The practical recovery experiments, however, tell a more nuanced story. When ρ is calibrated jointly with the smile parameters, its posterior is systematically biased toward the boundary of the admissible region, and the bias does not shrink as pricing noise decreases. This is the signature of a parameter that is formally identifiable but economically underdetermined: the option surface constrains ρ only through the skew it generates in combination with α and β , and the three parameters trade off along a shallow direction that the local Hessian spectrum, computed at a single operating point, does not fully expose.

The resolution is to recognize that ρ carries two distinct economic contents. In the physical measure, ρ is the contemporaneous correlation between order-flow and return innovations, a quantity with a direct time-series interpretation that is estimable from high-frequency data in which both series are observed jointly. In the risk-neutral measure, ρ enters only through the cross-diffusion term and the drift adjustment, where it generates skew but is confounded with the local-volatility asymmetry α and the convexity β . The physical-measure estimate is the cleaner of the two: it is identified by data that observes the correlation directly, whereas the option-implied estimate is identified only through a second-order shape effect that is nearly degenerate with two other parameters. Fixing ρ from the physical measure therefore removes a degree of freedom that the surface cannot independently pin down, while leaving the parameters the surface does identify, namely σ_0 , α , β , and the drift block, sharper than they would be in a joint fit. This treatment is consistent with the broader staged-calibration philosophy: parameters that are stiff in the physical measure and soft in the risk-neutral measure belong to the physical block, and ρ falls squarely into that category. The option surface is then used to identify the parameters it can actually see, and the market price of flow risk in Section 6.5 is computed as the wedge in the parameters identifiable in both measures, namely the flow couplings c and $\bar{f}$, rather than as a wedge in ρ , which the surface cannot independently resolve.

5.2 Manifold boundary approximation and irreducibility verification

The manifold boundary approximation method (MBAM) detailed in [Transtrum and Qiu, 2014] directly addresses the irreducibility question. The procedure follows a geodesic of the Fisher metric from the operating point along the sloppiest eigendirection until it reaches the model manifold boundary. This boundary corresponds to a structural limit, such as a parameter approaching 0 or ∞ , or two parameters merging into a product or ratio. Such limits can then be imposed exactly to yield a model with one fewer parameter. When applied to a model that is already irreducible, the geodesic instead runs to the edge of the admissible region without causing the Fisher metric to degenerate, indicating that no further reduction is available.

We execute the MBAM on the reduced model using synthetic vanilla European option prices generated across a grid of six maturities and denser log-moneynesses. Computing model prices and parameter sensitivities through the full pricing pipeline requires a high-performance FD option pricing solver. To make this computationally feasible, the framework is implemented using JAX with global double-precision FLOAT64 arithmetic. Automatic differentiation JAX.JACFWD computes exact parameter Jacobians through the pricing pipeline. To maintain high performance (≈ 76.6 ms per ODE call), the connection coefficients are evaluated using a Gauss-Newton projection approximation that avoids expensive full Hessian-vector product passes. The resulting geodesic ODEs are integrated using SCIPY hybrid explicit/implicit LSODA solver, paired with a terminal event function that monitors the minimum eigenvalue of the regularized

Parameter	Initial ($\tau=0$)	$r_{\text{tol}}=10^{-3}$	$r_{\text{tol}}=10^{-4}$
<i>Integration summary</i>			
Final Time (τ_{final})	0.0	3.0000	3.0000
Steps Taken	0	63	63
Status	Starting	Completed (τ_{max})	Completed (τ_{max})
<i>Final parameter values</i>			
σ_0	0.180	1.800×10^{-1}	1.800×10^{-1}
α	-0.400	-4.000×10^{-1}	-4.000×10^{-1}
β	0.150	1.500×10^{-1}	1.500×10^{-1}
μ	2.000	$2.000 \times 10^{+0}$	$2.000 \times 10^{+0}$
g	1.000	$4.201 \times 10^{+0}$	$4.201 \times 10^{+0}$
c	0.080	8.026×10^{-2}	8.026×10^{-2}
$\bar{y}$	0.000	-4.893×10^{-2}	-4.893×10^{-2}
$\bar{f}$	0.000	2.510×10^{-3}	2.510×10^{-3}
ρ	-0.300	-3.000×10^{-1}	-3.000×10^{-1}
Step-size check: $\ u(r_{\text{tol}}=10^{-3}) - u(r_{\text{tol}}=10^{-4})\ = 2.56 \times 10^{-4}$			

Table 4: MBAM geodesic integration results for the nine-parameter reduced model from initial operating point Θ_{red}^* across different relative error tolerances (r_{tol}) using a grid of six maturities and denser log-moneyness spacing. Shaded cells indicate parameters ($\sigma_0, \alpha, \beta, \mu, \rho$) that remain unchanged (within integration tolerance) from initial values, whereas $g = \theta_5$ actively evolves along the geodesic trajectory.

Fisher Information Matrix.

Starting from the baseline operating point:

$$\theta^* = (0.18 \quad -0.4 \quad 0.15 \quad 2.0 \quad 1.0 \quad 0.08 \quad 0.0 \quad 0.0 \quad -0.3), \quad (37)$$

with an initial sloppiest eigendirection corresponding to a regularized minimum eigenvalue of $\lambda_1 \approx 14.4$, the geodesic integrates smoothly out to the maximum horizon $\tau_{\text{final}} = 3.0000$ (in 63 steps) across both relative error tolerances ($r_{\text{tol}} = 10^{-3}$ and $r_{\text{tol}} = 10^{-4}$). As detailed in Table 4, the options-facing smile parameters remain invariant (shaded), while $g = \theta_5$ (along with minor adjustments in the drift-block parameters) actively traverses the geodesic trajectory up to τ_{final} without encountering a premature finite-time singularity or metric degeneracy. Two procedural requirements are fully satisfied:

1. *Nondegenerate Origin.* The geodesic originates from a strictly nondegenerate metric (any underlying exact symmetries having been removed analytically during the prior reduction step to Θ_{red}).
2. *Step-Size Independence.* Verification across tighter error tolerances confirms robust resolution of the underlying manifold. Specifically, the parameter distance between the trajectories at the two tested tolerances is minimal ($\|u(r_{\text{tol}} = 10^{-3}) - u(r_{\text{tol}} = 10^{-4})\| = 2.56 \times 10^{-4}$), indicating that the trajectory is accurately resolved and free of numerical artifacts.

Furthermore, employing synthetic vanilla European option prices rather than empirical market data does not compromise or alter these qualitative conclusions. The MBAM probes the intrinsic geometric properties of the model manifold, specifically the rank deficiency of the Fisher Information Matrix governed by the Jacobian mapping of the model. Because structural limits, parameter boundaries, and manifold degeneracies are intrinsic characteristics of the model's functional architecture rather than artifacts of a specific price realization, the verification of the irreducible core remains robust across comprehensive option surfaces.

Two qualifications bound what this analysis establishes, and both matter for the calibration that follows. First, the method identifies a single direction, the sloppiest one, and along that direction it is g that runs to its boundary while the shaded parameters stay fixed. The result therefore ranks g as the most removable coordinate. It does not order the remaining parameters among themselves, and it does not certify that each shaded parameter is stiff in an absolute sense, which would require the full Fisher spectrum along the trajectory rather than the first eigendirection alone. Second, the computation uses a grid of six maturities. It speaks to identifiability across a term structure, and not to identifiability within a single maturity band, where the same parameters can behave very differently.

The memory reversion rate μ illustrates the second qualification. On a multi-maturity surface μ is identifiable, in the sense that it moves away from its physical value when it is calibrated. Its leverage on the observable is nonetheless slight. Sweeping μ across its full admissible range shifts the at-the-money term-structure slope by less than a tenth of a volatility point. On a single maturity band it is degenerate with the local-volatility parameters σ_0, α, β and does not move at all. We therefore do not treat μ as a free risk-neutral coordinate on its own, and we calibrate it only jointly with the term-structure shape parameter k introduced below.

This weak leverage reflects a structural feature of the reduced model that bears directly on the scope of a single-parameter-set fit. In its stationary form, that is, without the transient correction, the model produces an at-the-money term structure that is flat to mildly downward sloping, and this holds for every admissible value of μ, β , and γ . A representative SPX surface over maturities from roughly one month to six months instead shows an at-the-money term structure that rises by about three and a half volatility points. A single stationary parameter set can match the interior of this range while missing both ends, with the largest error at the shortest maturities, precisely where the transient is strongest. The property that one parameter set spans the entire surface is therefore not yet established for this model, and it cannot be established from the stationary block alone. We treat it as a claim to be tested rather than an assumption, and we return to it in Section 6.3.

The transient correction is what closes this gap. It retains the time dependence of the conditional mean of the signal and reintroduces the relaxation rate k as a term-structure shape parameter. Because k controls the maturity dependence that the stationary block lacks, it is the parameter that carries the term structure rather than a small correction to it. We therefore include k in the calibrated set, and we assess the single-parameter-set property only with k active. A natural question is then whether the drift block $\{c, \bar{f}, \bar{y}, g\}$ can be identified under the physical measure $\mathbb{P}$. While vanilla option surfaces are largely insensitive to these slow parameters, rendering them flat in the risk-neutral calibration, their primary statistical footprint resides in time-series dynamics, driven by historical drift, persistence, and stationary moments of the underlying asset paths.

A rigorous dual-measure identifiability analysis for both measures simultaneously falls outside the intended scope of this pricing framework. Instead, consistent with standard practice in stochastic volatility modeling, the slow drift parameters $\bar{y}$ and g are reliably pinned from historical time-series data using standard estimation techniques, where unconditional empirical moments and long-term trends effectively constrain them prior to calibrating the options-facing block. The two flow parameters c and $\bar{f}$ are treated differently. The market price of flow risk is defined as the wedge between their physical and risk-neutral values, so for that wedge to be an important output rather than an identity, c and $\bar{f}$ must belong to the calibrated risk-neutral set. We adopt this in Section 6.3, where c and $\bar{f}$ are calibrated alongside the smile parameters and the term-structure parameter k .

6 Numerical experiments

The preceding sections established the reduced model by analytical reduction and proved its identifiability through the geometry of the calibration manifold. The purpose of this section is not to present a

comprehensive empirical asset-pricing study, but to verify that the theoretical properties survive contact with market data. Specifically, we show that the reduced parameter vector can be calibrated stably to a standard option surface, and that the staged separation of physical and risk-neutral measure parameters yields a measurable wedge (the market price of flow risk), rather than an ill-defined difference between two ridge points.

To this end, a single representative SPX snapshot suffices. The exercise has two goals. First, to confirm that the global term-structure fit with one parameter set across all maturities is feasible, and that the transient ramp supplies the necessary short-end shape without resorting to maturity-wise refitting. Second, to illustrate that the flow-impact parameters, calibrated on the risk-neutral measure, produce option-implied values whose difference from their time-series counterparts is an estimate with a well-defined standard error derived from the Gauss-Newton Hessian. The results below should be read as an existence proof for the practical identifiability of the reduced core, not as a trading manual or a backtested strategy.

6.1 Implementation

The reduced pricing PDE [Eq. \(23\)](#) is solved in the present numerical experiments by an explicit finite-difference scheme implemented in JAX. The implementation is deliberately lightweight rather than a production pricing engine: its purpose is to provide a fast, stable, and fully differentiable numerical solver for the calibration and identifiability experiments. The more elaborate splitting and RBF constructions discussed in [Section 3.4](#), together with the positivity-preserving treatment of the mixed derivative of [\[Itkin, 2026b\]](#), provide alternative routes for a production implementation.

Explicit pricing scheme.

The two-dimensional state space (x, y) is discretized on a tensor grid with $N_x = 70$ log-price nodes and $N_y = 50$ memory nodes. The computational domain is centered around the initial state and uses half-widths $h_x = 1$, $h_y = 1.5$. Time is discretized with $N_\tau = 500$ steps. The relatively fine grid is chosen to make the explicit scheme robust over the parameter ranges explored during calibration rather than to optimize the cost of a single pricing run. The time step is restricted by the corresponding CFL conditions for the state-dependent diffusion, the memory diffusion, and the mixed derivative. Within this stability region the scheme is numerically stable for all parameter sets retained in the experiments.

The spatial derivatives in [Eq. \(23\)](#) are evaluated directly on the finite-difference grid. The state-dependent volatility in [Eq. \(19\)](#) is evaluated at the memory nodes, while the drift, diffusion, mixed-derivative and nonlinear terms are advanced explicitly. Boundary values are chosen consistently with the asymptotic option behavior. The resulting implementation contains no RBF interpolation or matrix factorization and therefore avoids the ill-conditioning associated with the flat Gaussian bases used in the earlier implementation.

Differentiability and calibration.

The complete pricing and calibration pipeline is implemented in JAX, with double precision enabled. Grid construction, coefficient evaluation, finite-difference time stepping, option-price extraction, and the conversion from prices to implied volatilities are all part of the differentiable computational graph. The objective and its gradient are compiled with `jax.jit` and evaluated using `jax.value_and_grad`. Thus, unlike a finite-difference approximation to the calibration gradient, the optimizer receives derivatives obtained by automatic differentiation through the numerical pricing scheme.

Calibration is performed in implied-volatility space. For a parameter vector ϑ , the model implied volatility for quote i is denoted by $\sigma_{IV,i}^{\text{mod}}(\vartheta)$, and the objective is

$$\mathcal{J}(\vartheta) = \frac{1}{2M} \sum_{i=1}^M \left[\sigma_{IV,i}^{\text{mod}}(\vartheta) - \sigma_{IV,i}^{\text{mkt}} \right]^2. \quad (38)$$

Working in implied-volatility rather than price space prevents options with large dollar prices from dominating the objective and makes residuals comparable across strikes and maturities.

The six calibrated parameters are $\vartheta = (\sigma_0, \alpha, \beta, \mu, c, k)$. The physical-measure quantities $(g, \bar{y}, \sigma_z, \rho, \bar{f})$ and the exogenous quantities (γ, D) are held fixed at the values specified in [Section 6.3](#). The parameter c is retained in the option calibration despite its relatively weak identifiability. This is a deliberate compromise: the identifiability analysis shows that $\bar{f}$ is the strongest physical-measure parameter and c is the next strongest, but moving c entirely into the physical-measure block would eliminate the option-implied wedge needed to define the market price of flow risk. We therefore calibrate c on the option surface while explicitly recognizing that its estimate is substantially softer than those of the principal option-facing parameters. The parameter k plays a different role: it is retained primarily to describe the short-maturity term-structure effect associated with the transient signal correction in [Section 3.3](#).

The optimization uses the bound-constrained SLSQP algorithm from SCIPY. The physical-measure parameters $(g, \bar{y}, \sigma_z, \rho, \bar{f})$ are partly inherited from the time-series calibration of [\[Halperin and Itkin, 2026\]](#), while γ and the regularization parameter D are fixed exogenously. The option-facing parameters are initialized and bounded as shown in [Table 5](#).

Parameter	Initial value	Lower bound	Upper bound
σ_0	0.200	0.050	0.990
α	0.000	-2.000	2.000
β	0.100	0.010	25.000
μ	0.500	0.010	7.000
c	0.900	0.001	4.000
k	1.000	0.010	5.000

Table 5: Initial values and parameter bounds used in the option calibration.

The values of the fixed parameters are as follows:

$$(g, \bar{y}, \sigma_z, \rho, \bar{f}) = (0.3, 1.5, 0.5, -0.5, 0.1), \quad \gamma = 3, \quad D = 5. \quad (39)$$

The risk aversion parameter is fixed exogenously at $\gamma = 3$. Because the state variable x represents the log-price, the exponential utility specification effectively behaves as a power utility function over the arithmetic price level, rendering γ a dimensionless parameter that governs relative risk aversion [\[Henderson, 2002\]](#). This structural equivalence circumvents the scale-dependence typical of exponential utility and aligns our calibration with standard macro-finance benchmarks, where empirically supported relative risk aversion coefficients are strongly anchored between 2 and 4 [\[Friend and Blume, 1975; Mehra and Prescott, 1985\]](#).

The positivity requirement for the volatility radicand is imposed through a smooth penalty based on the minimum of the quadratic form in [Eq. \(19\)](#) over the computational memory interval. This replaces the nonlinear constraint used in the earlier implementation while keeping the objective fully differentiable. The penalty is negligible when the radicand remains positive and increases smoothly as the minimum approaches or crosses the admissible boundary.

Market data and stratification.

The calibration does not use every raw option quote. The objective is intended to represent the information content of the surface rather than the microstructure noise of individual observations. Market quotes are therefore first filtered for basic validity and then stratified across the term structure and moneyness. Maturities that are sufficiently close are assigned to the same maturity group, after which a representative set of strikes is selected within each group so that both the term structure and the smile are covered.

For each maturity group the solver evaluates the required strikes together, so that common intermediate quantities in the pricing calculation can be reused. The resulting workflow is therefore

raw quotes $\rightarrow$ **filtered/stratified surface** $\rightarrow$ **explicit 2D pricing** $\rightarrow$ **implied volatilities** $\rightarrow \mathcal{J}(\vartheta), \nabla \mathcal{J}(\vartheta)$

with the last two quantities supplied directly to the constrained optimizer. This combination of surface stratification, a CFL-stable explicit solver, and automatic differentiation makes repeated calibrations sufficiently fast for the identifiability and robustness experiments of [Section 6](#).

6.2 Black–Scholes synthetic-data experiment

As a controlled validation of the pricing and calibration machinery, we first consider a synthetic option surface generated from the Black–Scholes model. This experiment provides a useful benchmark because the data-generating parameters are known exactly, while the calibration is nevertheless performed with the full nonlinear two-dimensional pricing model rather than with the Black–Scholes formula. Thus, the experiment tests whether the proposed pricing–calibration pipeline can recover a surface generated by a substantially simpler model without introducing material pricing errors.

The synthetic market is initialized with $S_0 = 100, r = 0.05, q = 0.02, \sigma_{BS} = 0.25$ and covers the period from January 1 through December 31, 2024. For each of the four maturities $T \in \{0.25, 0.50, 0.75, 1.00\}$ options are generated at five strikes $K \in \{90, 95, 100, 105, 110\}$. Both calls and puts are included where applicable, producing a deliberately small but representative cross-section spanning near- and moderately out-of-the-money options and maturities from three months to one year. The resulting synthetic quotes have a constant Black–Scholes implied volatility of 25%, while prices and Greeks vary across strike and maturity in the usual way.

The synthetic prices are subsequently treated as market observations and the parameters of the proposed model are calibrated without imposing the Black–Scholes restriction. In particular, the six risk-neutral calibration parameters $(\sigma_0, \alpha, \beta, \mu, c, k)$ are optimized, while the physical parameters and exogenous parameters are held fixed at their prescribed values. The experiment therefore constitutes a stringent consistency check: the calibration must reproduce a flat implied-volatility surface using a nonlinear two-dimensional model with state-dependent volatility and memory effects.

The calibration converges successfully, with the final parameter vector $\sigma_0 = 0.24763, \alpha = -0.00034, \beta = 0.05349, \mu = 0.90734, c = 0.07690, k = 0.88869$. The resulting implied-volatility residuals are extremely small. Across the 20 calibration points, the mean residual is approximately zero, the standard deviation is 2×10^{-4} , and the root-mean-square error is $\text{RMSE}_{IV} = 2 \times 10^{-4}$, with residuals ranging from approximately -4×10^{-4} to 4×10^{-4} . These errors are negligible relative to the 25% input volatility and demonstrate that the numerical pricing scheme can reproduce the synthetic Black–Scholes surface to high accuracy. The calibrated parameters themselves need not coincide with a unique Black–Scholes representation: the purpose of the experiment is instead to verify that the richer model contains a parameter configuration capable of reproducing the benchmark surface.

[Figure 3a](#) shows the resulting residual surface in strike–maturity space, while [Fig. 3b](#) presents the same residuals in two dimensions together with the individual calibration errors. The residuals remain uniformly close to zero over the entire calibration region, with no visible systematic dependence on either strike or maturity. This is particularly important because systematic patterns would indicate a numerical or implementation bias rather than mere optimization error.

Overall, the experiment confirms both components of the numerical procedure: the explicit two-dimensional pricer is sufficiently accurate for calibration, and the JAX-based automatic differentiation combined with the bound-constrained optimizer is able to identify a parameter configuration that reproduces the benchmark surface with essentially negligible implied volatility error. The experiment is therefore used as a numerical validation rather than as an economic calibration exercise.




Figure 3: Black-Scholes synthetic-data experiment: implied-volatility residual a) surface as a function of strike and maturity; b) 2D and calibration errors. The residuals are uniformly close to zero across the calibration domain.

In other words, the Black-Scholes test confirms that the calibration does not manufacture identification where none exists. The reduced Marketron model reproduces a flat 25% implied-volatility surface with an RMSE of only 0.02 volatility points, while the fitted parameters (β, μ, c, k) remain free to move substantially along a nearly flat objective valley. The asymmetric coefficient α is driven to zero, as expected for a surface with no skew. Thus the test separates numerical accuracy from parameter identification: an arbitrarily accurate fit does not imply that the underlying structural parameters are identified.

6.3 Staged calibration to SPX options

We test the reduced model on SPX options data from January 18, 2017, filtered to the maturity band $T \in [0.04, 0.5]$ years. The data processing pipeline described in Section 6.1 yields 14 maturity groups (grouped by minimum distance of 0.01 year) and a calibration set of 138 options. This market data is presented in Fig. 4. Calls and puts are retained where available so that the calibration contains information from both sides of the smile. The stratification is performed once before the optimization; the resulting quotes are then held fixed throughout the calibration. This prevents changes in the numerical sample from being mistaken for changes in the calibrated parameter vector.

With the physical block frozen as in Eq. (39), the model calibrates Θ_Q to the full option surface using the implied-volatility objective and the SLSQP optimizer described in Section 6.1. The regularizer $\bar{\epsilon}$ is fixed via Eq. (13).

The results are reported in Table 6 and the overall fit is illustrated in Fig. 5: the model reproduces the level, skew, and convexity of the SPX surface across all maturity groups with a single parameter set. The quantitative residual statistics provide a more detailed measure of the fit. The residual statistics are computed on the (K, T) residual grid. On this grid, quotes that share a strike and maturity, such as a call and a put, are averaged into a single cell, so the 138 calibration quotes reduce to 110 distinct cells. Across these 110 cells, the mean implied-volatility residual is 0.0017, with a standard deviation of 0.0183 and $\text{RMSE}_{\text{IV}} = 0.0184$. The residuals range from approximately -4.23 to 4.06 vol points. Thus, the calibrated model achieves an RMS error of approximately two vol-points while fitting simultaneously across strikes and maturities.




Figure 4: Stratified/clustered input option data of SPX used for calibration.

Parameter	Initial			Calibrated		
	P	Endog.	Q	P	Endog.	Q
g	0.3	—	—	0.3	—	—
$\bar{y}$	1.5	—	—	1.5	—	—
σ_z	0.5	—	—	0.5	—	—
ρ	-0.5	—	—	-0.5	—	—
$\bar{f}$	0.1	—	—	0.1	—	—
γ	—	3.0	—	—	3.0	—
D	—	5.0	—	—	5.0	—
σ_0	—	—	0.2	—	—	0.05823
α	—	—	0.0	—	—	2.00000
β	—	—	0.1	—	—	1.92349
μ	—	—	0.5	—	—	0.01000
c	—	—	0.9	—	—	0.62824
k	—	—	1.0	—	—	1.00276

Table 6: Initial and calibrated parameter values.

The calibrated parameters also provide useful information about the role and identifiability of the model components. In particular, α and β are calibrated to 2.0 and 1.92349, respectively. Both parameters are substantially different from zero. This contrasts with the Black–Scholes limiting case, in which these parameters would theoretically vanish, and indicates that the additional dynamics associated with them are relevant for the observed SPX surface. Their nonzero values are therefore consistent with the pronounced skew and curvature present in the market data.

The calibration also highlights differences in parameter identifiability. While α and β move materially from their initial values, α resides at the boundary which may indicate its weak identifiability. Indeed, based on Eq. (19), same volatility can be obtained by varying σ_0, α, β simultaneously. The parameter k remains essentially unchanged: its initial value of 1.0 moves only to 1.00276. This suggests that k is weakly identified by the available SPX cross-section over the maturity range considered. In contrast, the substantial movement in α and β indicates that the option surface contains considerably more information about these parameters. This distinction is important when interpreting the calibrated parameter vector: successful optimization and a good surface fit do not imply that all model parameters are equally well identified by the available option data.




Figure 5: SPX market data experiment: implied-volatility residual a) surface as a function of strike and maturity; b) 2D and calibration errors. The residuals are uniformly small across the calibration domain.

Note that the fitted instantaneous volatility scale should not be interpreted directly as the level of market-implied volatility. The latter is an option-pricing quantity that incorporates the full future dynamics of the memory state and the nonlinear pricing operator. It is therefore not appropriate to assess the consistency of the calibrated baseline volatility with the observed SPX implied-volatility level by comparing σ_0 , or $\sigma_x(y)$ at a particular value of y , directly with the market IVs. Instead, consistency with the observed volatility surface should be assessed through the model-implied option prices and the corresponding implied volatilities. The calibrated value of σ_0 should thus be interpreted as the baseline instantaneous volatility scale of the state-dependent diffusion, rather than as a direct estimate of the market-implied volatility level.

The estimate of the mean-reversion parameter μ also sits at the low boundary, while similar values obtained in the option calibration of [Halperin and Itkin, 2025] are $\mu \approx 4.6$. The difference is material and most likely can be attributed to how we do adiabatic averaging in Section 3.1, i.e. by setting $\bar{h} = 0$, and hence only the product $\mu\bar{y}$ does matter, while in [Halperin and Itkin, 2025] $\bar{h}$ is negative. The high value of μ in [Halperin and Itkin, 2025] provides fast mean-reversion of the memory variable, but to a stochastic level while here the mean-reversion level is deterministic. Anyway, this disagreement is particularly noteworthy because the present model is a reduced two-dimensional formulation with a different volatility specification and a substantially different treatment of parameter identifiability. The result suggests that the mean-reversion timescale associated with the memory variable is unstable across the two formulations.

The flow-impact parameter c behaves quite differently. In [Halperin and Itkin, 2025] its calibrated value was approximately 0.9, whereas in the present experiment the optimizer yields $c = 0.62824$. The identifiability analysis in Section 5 shows that vanilla option prices contain relatively little information about the absolute level of c , which is substantially less tightly identified than the principal smile parameters. Nevertheless, conditional on the physical-measure block and the other option-facing parameters used here, the SPX surface favors a lower risk-neutral value of c than the initial value inherited from the physical measure. The fact that the optimum differs from this initial value suggests that the vanilla surface contains some information about c^Q beyond the physical-measure specification, although the weak identification of c implies that this distinction should not be interpreted as strong evidence for a precisely identified risk-neutral value. Finally, the comparison with [Halperin and Itkin, 2025] shows that the two calibrated

values of c are of the same order of magnitude despite the differences between the underlying models.

The residual surface in Fig. 5a and its projection in Fig. 5b provide a further diagnostic of the calibration. They show how the remaining pricing errors are distributed across strike and maturity rather than only reporting their aggregate magnitude. In particular, systematic residual patterns would indicate regions of the empirical SPX surface that are not fully captured by the reduced model, whereas residuals that remain centered around zero across the calibration domain would support the adequacy of the model specification for the observed surface.

From a pricing perspective, the IV residual surface reveals a clear spatial pattern in the model's fit across the observed domain of OTM calls and puts, with moneyness $M \in [0.1, 1.01]$. Residuals are closest to zero in a band surrounding the approximate at-the-money region ($K \approx 2250\text{--}2260$), indicating that the model fits near-ATM options most closely and that the quality of the fit deteriorates toward the wings. The largest positive residuals, corresponding to model over-prediction of IV and reaching approximately $+0.0225$, concentrate at higher strikes and short maturities, i.e., in the OTM-call region. The largest negative residuals, corresponding to model under-prediction and reaching approximately -0.040 , occur at lower strikes across the maturity range, i.e., in the OTM-put region.

Consequently, the model provides a substantially better fit for near-ATM options than for deep-OTM puts and, to a somewhat lesser extent, short-dated OTM calls. The persistence of the short-maturity call residuals is notable because the parameter k is retained in the model specifically to control short-time maturity behavior. The residual surface therefore suggests that this mechanism, while important for the short-time asymptotics, is not sufficient to capture the observed short-dated upside wing. The asymmetric wing behavior points instead to two distinct directions for subsequent model refinement: a better representation of the downside skew in OTM puts and an additional mechanism for the short-term upside wing in OTM calls, precisely where the absolute residuals are largest within the available market data.

6.4 Another SPX experiment

To examine the sensitivity of these conclusions to the calibration region, we performed a second calibration with the same SPX data and the same exogenous parameters, but with a wider admissible domain for the risk-neutral parameters. In this experiment we used

$$(g, \bar{y}, \sigma_z, \rho, \bar{f}) = (0.3, 1.5, 0.8, -0.5, 0.1), \quad (\gamma, D) = (3, 5), \quad (40)$$

and initialized $(\sigma_0, \alpha, \beta, \mu, c, k) = (0.2, 0, 0.1, 2, 5, 1)$. The bounds were enlarged to

$$\sigma_0 \in [0.05, 0.99], \quad \alpha \in [-2, 2], \quad \beta \in [0.01, 5], \quad \mu \in [10^{-4}, 10], \quad c \in [0.01, 15], \quad k \in [0.01, 5]. \quad (41)$$

The resulting calibration was $(\sigma_0, \alpha, \beta, \mu, c, k) = (0.05004, 0.34809, 0.50984, 0.00606, 1.64334, 0.86631)$, with $\text{RMSE}_{\text{IV}} = 0.0165$. Thus, the fit is somewhat better than that of the baseline calibration, for which $\text{RMSE}_{\text{IV}} = 0.0184$, although the improvement should not be interpreted as a pure optimizer comparison because the physical volatility parameter was changed from $\sigma_z = 0.5$ in the baseline experiment to $\sigma_z = 0.8$ in the second experiment. More importantly, the two calibrations converge to materially different parameter vectors, indicating the presence of multiple favorable regions, or basins, in the parameter-to-option-price map. The computed residuals are also presented in Fig. 6.

The difference between the two solutions also illustrates the limited identifiability of the parameters governing the state-dependent instantaneous volatility. In the baseline calibration, $(\alpha, \beta) = (2.0000, 1.9235)$, with α at its upper bound, whereas the wider calibration gives $(\alpha, \beta) = (0.3481, 0.5098)$, with both parameters well inside their bounds. Since the option-pricing operator observes the volatility function $\sigma(y)$ rather than σ_0 , α , and β separately, different combinations of these parameters can generate similar effective volatility profiles over the region of the memory state visited by the pricing dynamics. The movement




Figure 6: SPX market data *second* experiment: implied-volatility residual a) surface as a function of strike and maturity; b) 2D and calibration errors. The residuals are uniformly small across the calibration domain.

between these two parameterizations therefore provides empirical evidence for the approximate $(\sigma_0, \alpha, \beta)$ redundancy discussed above. The fact that the two calibrations nevertheless produce similar overall pricing errors shows that the option surface constrains the resulting volatility structure more strongly than it constrains its particular parametric decomposition. This also qualifies the local identifiability analysis of Section 5, whose Hessian and profile widths are evaluated at the operating point Θ_{red}^* , which is far from either SPX optimum: the $(\sigma_0, \alpha, \beta)$ direction that reads as stiff there is, at the fitted surface, a shallow one along which α and β trade off, much as ρ does in Section 5.

The memory parameter μ provides a particularly clear result. In both experiments the optimizer drives μ to the lower edge of the admissible region: $\mu = 0.01$ in the baseline calibration and $\mu = 0.0061$ in the wider calibration. Thus, the option data do not appear to support appreciable mean-reversion of the memory variable through the explicit term $\mu(\bar{y} - y)$. Instead, within the present reduced specification, the effective stabilization of the memory dynamics appears to be generated primarily through its coupling to the price and the nonlinear potential. This conclusion should nevertheless be treated cautiously because the two experiments use different σ_z and therefore do not isolate the effect of μ alone.

The flow-impact coefficient c moves from 0.6282 to 1.6433 across the two calibrations. This spread is expected rather than troubling: Section 5 identifies c as the soft option-facing coordinate, so its point value is not sharply pinned by a single surface. What both basins share is that $c^{\mathbb{Q}}$ does not collapse to the inherited physical value $c^{\mathbb{P}} = 0.9$, so the staged calibration produces a nonzero $\mathbb{Q}$ – $\mathbb{P}$ separation in the flow block in either basin. The market price of flow risk is not this scalar difference, however, but the wedge in the market price of risk $\lambda_t^{(x)}$ of Section 6.5, into which c enters together with $\bar{f}$ and the state-dependent volatility. The two-basin spread should therefore be read as evidence that a flow-risk premium is identifiable and nonzero, rather than as a sharp estimate of its level.

By contrast, k moves from 1.0028 to 0.8663, a considerably smaller relative change. This suggests that the signal relaxation timescale may be more stable across calibrations, although the available data are not sufficient to claim strong identification.

6.5 The market price of risk

In the original Marketron model of [Halperin and Itkin, 2026], the three Brownian motions $W_t^{(x)}$, $W_t^{(y)}$, and $W_t^{(\theta)}$ are mutually independent. The utility-based market price of risk (MPR) is therefore a three-component vector,

$$\lambda_t = \left(\lambda_t^{(x)}, \lambda_t^{(y)}, \lambda_t^{(\theta)} \right), \quad (42)$$

with one component per stochastic factor. The component associated with the traded Brownian motion $W_t^{(x)}$ takes the same functional form as in a complete market,

$$\lambda_t^{(x)} = \frac{\mu_x(t, x_t, y_t, \theta_t) - r}{\sigma}, \quad (43)$$

while the components $\lambda_t^{(y)}$ and $\lambda_t^{(\theta)}$ correspond to the unobservable memory and signal variables and are not directly measurable.

The reduced model changes this structure in two ways. First, adiabatic elimination of the signal removes θ from the state space, so the third component $\lambda_t^{(\theta)}$ disappears entirely. Since the reduced model sets $\rho_{x\theta} = 0$ at leading order (the correlation enters only through the first-order $1/k$ correction discussed in Section 3.3), there is no residual signal contribution to the MPR:

$$\lambda_t^{(\theta)} \equiv 0. \quad (44)$$

Second, the introduction of the diffusive correlation ρ between $W_t^{(x)}$ and $W_t^{(y)}$ generates a non-zero second component $\lambda_t^{(y)}$. Because the memory variable y_t is now partially correlated with the traded asset, it is partially hedgeable, and the market demands compensation for its residual risk. The MPR vector in the reduced model is therefore two-dimensional:

$$\lambda_t = \left(\lambda_t^{(x)}, \lambda_t^{(y)} \right), \quad (45)$$

where the first component follows from the indifference pricing framework and takes the same functional form as in a complete market:

$$\lambda_t^{(x)} = \frac{\bar{\mu}_x(t, x_t, y_t)}{\sigma_x(y_t)}, \quad (46)$$

with

$$\bar{\mu}_x(t, x_t, y_t) = \bar{f}(t) + \bar{\eta}(y_t) + \frac{\sigma_x^2(y_t)}{2} - r - c_x y_t V'_M(x_t), \quad (47)$$

and $\bar{\eta}(y_t) = r - \sigma_x^2(y_t)/2$. The expression simplifies to

$$\lambda_t^{(x)} = \frac{\bar{f}(t) - c_x y_t V'_M(x_t)}{\sigma_x(y_t)}. \quad (48)$$

The second component $\lambda_t^{(y)}$ is generated by the correlation ρ through the market-price-of-risk terms in the pricing PDE Eq. (23). From the drift adjustment in the y -substep,

$$\mu_y^{\mathbb{Q}} = \mu_y - \rho \frac{\bar{\mu}_x}{\sigma_x(y)}, \quad (49)$$

the component is

$$\lambda_t^{(y)} = \frac{\rho \bar{\mu}_x(t, x_t, y_t)}{\sigma_x(y_t)} = \rho \lambda_t^{(x)}. \quad (50)$$

Thus, the two components are proportional, with the correlation ρ as the proportionality constant. When $\rho = 0$, the MPR reduces to the single component $\lambda_t^{(x)}$ of the uncorrelated model, consistent with [Halperin and Itkin, 2025]. When $\rho \neq 0$, the market demands an additional premium for bearing the partially hedgeable memory risk, and this premium is exactly ρ times the traded-asset premium.

In contrast to the complete-market setting, both components of the MPR in the reduced Marketron are *stochastic*: they depend on the memory variable y_t both directly through the term $c_x y_t V'_M(x_t)$ and indirectly through the state-dependent volatility $\sigma_x(y_t)$. The memory variable encodes the accumulated flow impact, and its fluctuations generate a time-varying, state-dependent risk premium.

Numerical computation.

Since y_t is unobservable, the empirical MPR is computed by Monte Carlo simulation of the reduced system Eq. (18) under the physical measure. Along each simulated path, we evaluate Eq. (48) and Eq. (50) and take the expectation over the ensemble,

$$\bar{\lambda}_t^{(x)} = \mathbb{E}[\lambda_t^{(x)}] = \mathbb{E}\left[\frac{\bar{f}(t) - c_x y_t V'_M(x_t)}{\sigma_x(y_t)}\right], \quad \bar{\lambda}_t^{(y)} = \rho \bar{\lambda}_t^{(x)}. \quad (51)$$

Defaulted paths, where x_t crosses the threshold $\hat{x}$ into the default region, are excluded from the expectation, consistent with the treatment in [Halperin and Itkin, 2025]. The standard error of the Monte Carlo estimate is computed from the cross-sectional dispersion of the MPR across paths.

The wedge between the physical and risk-neutral measures is then quantified through the *level* of the MPR. The physical-measure MPR $\bar{\lambda}_t^{(x),\mathbb{P}}$ is computed from the time-series calibration of [Halperin and Itkin, 2026], while the risk-neutral MPR $\bar{\lambda}_t^{(x),\mathbb{Q}}$ is obtained from the option-implied parameters. The difference,

$$\Delta\lambda_t \equiv \bar{\lambda}_t^{(x),\mathbb{Q}} - \bar{\lambda}_t^{(x),\mathbb{P}}, \quad (52)$$

represents the market price of flow risk as a function of time.

Remark. The MPR level and the wedge Eq. (52) were computed numerically for the full Marketron in [Halperin and Itkin, 2025], and we do not reproduce those figures here. In isolation, with no second measure or model to compare against, the reduced-model values would add length without information. What the reduced core contributes is not a new number but the status of the wedge: on the sloppy full model the $\mathbb{Q}$ – $\mathbb{P}$ separation is a ridge artifact, whereas on the reduced core it is a well-defined object, subject to the weak identification of c noted in Section 6.3.

7 Discussion

The central message of this paper is epistemological rather than merely computational. The full Marketron, like many structural models in quantitative finance, carries more parameters than the option surface can resolve. This is not a numerical inconvenience to be overcome with better optimizers. It is a structural defect that renders the model's economic claims unfalsifiable. By removing exact symmetries, freezing non-financial parameters, and adiabatically eliminating the fast signal, we have reduced the model to an identifiable core in which every parameter is either estimated from the data or fixed by an explicit economic criterion. The calibration objective is no longer a heuristic search over a sloppy landscape but a far better-posed estimation problem, even though, as the SPX fits show, individual coordinates such as c and μ remain softly determined and the optimum can sit at a bound.

This change of status has immediate economic consequences. In the full 18-parameter model Eq. (7), the flow-impact constant $c = c_x = c_y$ sits on a continuous ridge along which it trades off against the memory scale, the hidden volatility, and the initial state. A reported value of c is therefore a point on a ridge,

not an estimate. On the reduced core [Eq. \(28\)](#), the ridge is destroyed by the gauge fixing [Eq. \(9\)](#) and the level absorption $\bar{h} \rightarrow \bar{y}$, and c becomes an identifiable coordinate of the drift block, the softest of the option-facing set but no longer a ridge direction. The market price of flow risk, the wedge $\Delta\lambda_t$ of [Eq. \(52\)](#) between the physical and pricing values of the flow block, is consequently a well-defined object rather than a point on a ridge: a compensation for bearing non-hedgeable order-flow exposure that is identifiable rather than asserted, though [Section 6.3](#) shows its level is only weakly pinned by a single surface. To our knowledge, this is the first such wedge made identifiable within a structural model of inelastic markets.

The correlation ρ plays a subtle but essential role in this reduction. In the uncorrelated model, risk aversion γ and the hidden volatility σ_y collapse into the single product $p_y = \gamma$, removing one degree of freedom from the diffusion structure. The correlation ρ restores it by entering the pricing operator linearly through the cross-term $\partial_{xy}^2 C$ and the drift adjustment, generating short-maturity skew that the state-dependent drift alone cannot supply. Financially, ρ is the diffusive signature of contemporaneous price impact; technically, it is the unique coefficient that breaks the maturity degeneracy of [Proposition 2](#) at the short end. Together with the effective constants $(\bar{f}, \bar{h})$, which encode the risk-adjusted drift wedge between the physical and pricing measures, ρ completes the economic reading of the reduced surface. In the calibration itself ρ is not free: it is fixed from the physical measure ([Section 5](#)), where it is directly estimable, because on the option surface its skew is confounded with α and β . Its role here is structural, restoring the degree of freedom that γ removes, rather than serving as a calibrated coordinate.

The reduction is not without cost. Adiabatic elimination of θ replaces the rich signal-driven term structure of the full model with the deterministic ramp [Eq. \(22\)](#), which captures only the relaxation of the conditional mean. When the signal autocorrelation is strong ($\sigma_z^2/2k \ll 1$) or when option maturities span the relaxation time $1/k$ densely, the full three-dimensional model with the transient correction of [Section 3.3](#) should be preferred. Similarly, the fluctuation-dissipation relation [Eq. \(35\)](#) reduces to the equality $k = \mu$ only at leading order; for strongly nonlinear forcing, the higher harmonics of the trigonometric signal introduce memory timescales that the single-mode kernel [Eq. \(31\)](#) cannot resolve. These are not failures of the reduced model but boundaries of its domain of validity, and they are precisely the sort of falsifiable conditions that an identifiable framework makes possible.

An important finding of the paper is that substituting the memory variable into the log-price equation yields a generalized Langevin equation with a closed-form, state-modulated memory kernel, while the memory variable itself provides an exact Markovian lift of this kernel. Thus, the representation in [Section 4](#) points to a broader class of models. The kernel [Eq. \(31\)](#) is separable and state-dependent, of the form $\mathbf{K}(Z_t, Z_s, \tau) = \Phi(Z_t) \mathbf{k}(\tau) \Phi(Z_s)^\top$ whereas, e.g., the framework of [\[Itkin, 2026a\]](#) assumes $\Phi \equiv \mathbf{I}$. The smallest class containing both is therefore the separable state-dependent GLE. Four questions must be settled before this broader class can be used as broadly as the lag-only class:

- Whether the Markovian embedding remains exact and finite-dimensional for a general Φ ? In financial-mathematics terms (FMT), whether the corresponding non-Markovian stochastic Volterra equation with a state-dependent impact propagator admits an exact finite-dimensional Markovian realization?
- What replaces [Eq. \(34\)](#) when the system–bath coupling is weighted by Φ , since the driving noise then becomes state-dependent and multiplicative? In FMT, what equilibrium-consistency condition replaces the usual relation between the impact-propagator decay and the exogenous-flow correlation time when the return innovation has state-dependent volatility?
- What the stationary law is, given that the Boltzmann form is already broken here, as [Eq. \(32\)](#) illustrates through the modulation-induced contribution to the effective potential? In FMT, what invariant distribution, if any, is induced by the state-dependent drift and diffusion of the reduced price–memory system, and how it differs from the equilibrium stationary distribution?
- Whether the triangular structure imposed on the kernel matrix by the absence of arbitrage survives state modulation, since the corresponding Girsanov density then acquires a state-dependent factor? In FMT, whether the no-arbitrage restriction on the impact-propagator matrix remains triangular

under a state-dependent market price of memory risk, and whether the resulting change of measure remains compatible with the pricing dynamics?

A fifth question is empirical: the separable form is not identified without a normalization, because $\Phi \mapsto \alpha\Phi$ and $\mathbf{k} \mapsto \alpha^{-2}\mathbf{k}$ leave the kernel unchanged, so the class inherits a gauge freedom of exactly the kind [Section 2](#) removes here.

Two empirical directions are immediate. First, the model-implied VIX inherits state dependence from the nonlinear indifference pricing operator and from the default-region contribution of the potential, despite the absence of a separate stochastic volatility factor, with instantaneous volatility given by the state-dependent function $\sigma_x(y)$. Computing this index and comparing it with the CBOE time series is a strict out-of-sample falsification test: if the two regimes cannot lift the model VIX materially away from the level implied by $\sigma_x(y)$, the Marketron in its present form cannot support volatility derivatives without a stochastic spot-volatility extension.

Second, joint calibration to VIX options would constrain the soft μ -direction identified in [Section 5](#) and sharpen the premium estimates in [Eq. \(52\)](#). Both tests are only meaningful on the identifiable core; on the full model, a mismatch could always be attributed to a poor optimum.

The asymmetric wing behavior reported in [Section 6](#) points to two distinct directions for subsequent model refinement: a better representation of the downside skew in OTM puts and an additional mechanism for the short-term upside wing in OTM calls. In particular, introducing a separate stochastic volatility factor could provide additional flexibility in the joint strike–maturity dynamics of implied volatility and may help account for the residual short-maturity call bias that is not eliminated by the current k -dependent short-time mechanism. Such an extension would, however, need to preserve the model’s no-arbitrage structure and would introduce additional parameters and a corresponding calibration trade-off. We reserve this extension for future investigation.

A further limitation is that the present calibration fixes the P -parameters rather than estimating them jointly with the endogenous parameters. In a full calibration, these parameters should themselves be inferred from the relevant market data, and their estimation may alter both the fitted values and the implied residual structure reported above. A natural direction for future research is therefore to investigate the sensitivity of the model and its pricing performance to both the P -parameters and the endogenous parameters, including their joint calibration.

Finally, the computational cost of the reduced pricer could be further reduced by constructing a neural surrogate trained on solutions of the reduced PDE [Eq. \(23\)](#). Such a surrogate could bring pricing times to the millisecond range, making the model more suitable for live trading and intraday recalibration. Because the reduced parameter space is low-dimensional and the calibration problem exhibits substantially improved identifiability, the training set could be generated on a regular lattice using standard supervised learning methods. The surrogate would then approximate the pricing map of the structurally constrained reduced pricer, while the latter would remain the source of the model’s arbitrage and boundary-condition guarantees. We also leave this extension to future work.

Taken together with the analysis in [\[Itkin, 2026a\]](#), the results of this paper point to a broader interpretation of the Marketron as an economically structured realization of generalized Langevin dynamics. The memory variable in the reduced Marketron corresponds to a single exponential memory mode, whereas the generalized Langevin formulation permits a general memory kernel or a superposition of exponential modes, allowing the temporal structure of memory to be determined empirically rather than imposed through a single relaxation scale. Conversely, the Marketron provides an economic interpretation of the memory and price-impact mechanisms that are introduced more phenomenologically in the generalized Langevin framework. This connection suggests a natural next step: to combine the economic structure of the Marketron with the more general GLE memory specification and investigate its implications in a joint calibration to SPX and VIX data, thereby testing whether a common framework can consistently capture the short-, intermediate-, and long-horizon dynamics of volatility and leverage.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

No funding was received.

Disclaimer

Opinions expressed here are author's own, and do not represent views of their employers. A standard disclaimer applies.

Acknowledgments

I thank my long-time co-author Igor Halperin for our joint work on the Marketron model and valuable discussions, as well as Michael Isichenko for his constructive feedback regarding the Marketron model's dimensionality.

References

- Abi-Jaber, E and O. El-Euch (2019). "Multifactor approximation of rough volatility models". In: *SIAM Journal on Financial Mathematics* **10**, 2, pp. 309–349.
- Abi Jaber, E., M. Larsson, and S. Pulido (2019). "Affine Volterra processes". In: *Annals of Applied Probability* **29**, 5, pp. 3155–3200.
- Bayer, C. and S. Breneis (2023). "Markovian approximations of stochastic Volterra equations with the fractional kernel". In: *Quantitative Finance* **23**, 1, pp. 53–70.
- Bliss, R. R. and N. Panigirtzoglou (1999). *Testing the stability of implied probability density functions*. Tech. rep. Bank of England.
- Fouque, J.-P. et al. (2011). *Multiscale Stochastic Volatility for Equity, Interest Rate, and Credit Derivatives*. Cambridge: Cambridge University Press.
- Friend, I. and M. E. Blume (1975). "The demand for risky assets". In: *The American Economic Review* **65**, 5, pp. 900–922.
- Grasselli, M. and T. R. Hurd (2007). "Indifference pricing and hedging in stochastic volatility models". In: *Applied Mathematical Finance* **14**, 4, pp. 303–317.
- Gutenkunst, R. N. et al. (2007). "Universally Sloppy Parameter Sensitivities in Systems Biology Models". In: *PLoS Computational Biology* **3**, 10, e189.
- Halperin, I. and A. Itkin (Feb. 2026). "Marketron Games: Self-Propelling Stocks vs Dumb Money and Metastable Dynamics of the Good, Bad and Ugly Markets". In: *Communications in Nonlinear Science and Numerical Simulation* **153**, p. 109529.
- Halperin, I. and A. Itkin (2025). *Marketron Through the Looking Glass: From Equity Dynamics to Option Pricing in Incomplete Markets*. arXiv: [2508.09863](https://arxiv.org/abs/2508.09863).
- Henderson, V. and J. Hobson (2002). "Substitute hedging". In: *Risk* **15**, 5, pp. 71–75.
- Henderson, V. (2002). "Valuation of claims on nontraded assets using utility maximization". In: *Mathematical Finance* **12**, 4, pp. 351–373.
- Itkin, A. (2026a). "Beyond rough volatility: decoupling memory and scaling via a generalized Langevin equation". In: *SSRN Electronic Journal*. URL: <https://ssrn.com/abstract=7202798>.

- Itkin, A. (2026b). “Diagonal Frog: High-order positivity-preserving FD schemes for anisotropic Fokker-Planck equations”. In: arXiv: [2606.23980 \[math.NA\]](#).
- Itkin, A. (2026c). “Flux-Corrected Diagonal Frog: second order and positivity at all time steps”. In: arXiv: [2607.20415 \[math.NA\]](#).
- Mehra, R. and E. C. Prescott (1985). “The equity premium: A puzzle”. In: *Journal of monetary economics* **15**, 2, pp. 145–161.
- Musiela, M. and T. Zariphopoulou (2004). “An example of indifference prices under exponential preferences”. In: *Finance and Stochastics* **8**, 2, pp. 229–239.
- Pavliotis, G. A. and A. M. Stuart (2008). *Multiscale Methods: Averaging and Homogenization*. Vol. 53. Texts in Applied Mathematics. New York: Springer.
- Raue, A. et al. (2009). “Structural and Practical Identifiability Analysis of Partially Observed Dynamical Models by Exploiting the Profile Likelihood”. In: *Bioinformatics* **25**, 15, pp. 1923–1929.
- Transtrum, M. K., B. B. Machta, and J. P. Sethna (2011). “Geometry of Nonlinear Least Squares with Applications to Sloppy Models and Optimization”. In: *Physical Review E* **83**, p. 036701.
- Transtrum, M. K. and P. Qiu (2014). “Model Reduction by Manifold Boundaries”. In: *Physical Review Letters* **113**, p. 098116.

Appendices

A Proof of Proposition 1

Proof. Let $\tilde{y}_t = y_t/\lambda$ denote the transformed process under the gauge transformation Eq. (8). Applying Itô’s lemma to $\tilde{y}_t$, we obtain $d\tilde{y}_t = \frac{1}{\lambda} dy_t$. Under this mapping, the spatial parameters scale proportionally, meaning $\bar{y} \rightarrow \bar{y}/\lambda$, $y_0 \rightarrow y_0/\lambda$, and $\sigma_y \rightarrow \sigma_y/\lambda$, while the mean-reversion rate μ remains invariant. Consequently, the ratios b_2/σ_y , $\bar{y}/\sigma_y$, and y_0/σ_y are invariant, as the scaling factor cancels out. The correlation parameters ρ and $\rho_{x\theta}$ are dimensionless and therefore unaffected by the state scaling. Assuming the product $c_x c_y$ is preserved under the transformation, the pair $(x_t, \tilde{y}_t)$ exactly solves the stochastic system with the transformed parameters, leaving the law of the observable process x_t unchanged.

To demonstrate that the option prices are preserved, we apply the spatial scaling to the indifference-pricing PDE Eq. (5). By the chain rule, the partial derivatives with respect to the scaled variable transform as $\partial_y C \rightarrow \frac{1}{\lambda} \partial_y C$, $\partial_y^2 C \rightarrow \frac{1}{\lambda^2} \partial_y^2 C$, and $\partial_{xy}^2 C \rightarrow \frac{1}{\lambda} \partial_{xy}^2 C$. Substituting these alongside the transformed volatility $\sigma_y \rightarrow \lambda \sigma_y$, we evaluate the key PDE components:

$$\begin{aligned} \gamma(\lambda\sigma_y)^2(1-\rho^2) \left(\frac{1}{\lambda} \partial_y C \right)^2 &= \gamma\sigma_y^2(1-\rho^2)(\partial_y C)^2, \\ (\lambda\sigma_y)^2 \left(\frac{1}{\lambda^2} \partial_y^2 C \right) &= \sigma_y^2 \partial_y^2 C, \\ \rho\sigma(\lambda\sigma_y) \left(\frac{1}{\lambda} \partial_{xy}^2 C \right) &= \rho\sigma\sigma_y \partial_{xy}^2 C. \end{aligned}$$

Since these terms are homogeneous of degree zero under the joint transformation $y \rightarrow \lambda y$ and $\sigma_y \rightarrow \lambda \sigma_y$, the PDE Eq. (5) is strictly form-invariant. Because both the observable system dynamics and the pricing equation are invariant, it follows that every option price remains unchanged. $\square$

B Fixing the regularizer

The regularizer $\bar{\epsilon}$ is not a financial parameter. This appendix fixes it, and in doing so settles which criterion can do the job, since the obvious one cannot.

Two admissible regularizers.

Write $V'_M(x) = -e^{-x}[1 - R(x)]$. The original specification of [Halperin and Itkin, 2026] takes

$$R_1(x) = \frac{g}{e^x + \bar{\epsilon}g}, \quad (\text{B.1})$$

whereas the pricer of [Halperin and Itkin, 2025] takes

$$R_2(x) = \frac{1}{2\bar{\epsilon}}[1 - \text{Erf}(x + b)], \quad b = 4 - \text{Erf}^{-1}\left[1 - \frac{2}{1 + e^4 g \bar{\epsilon}}\right], \quad (\text{B.2})$$

because the quadrature underlying the Gaussian RBF matrix elements closes in elementary functions for R_2 and does not for R_1 . Both share the limits $R \rightarrow 0$ as $x \rightarrow +\infty$ and $R \rightarrow 1/\bar{\epsilon}$ as $x \rightarrow -\infty$, so both reduce the divergence of V'_M in the default region from e^{-2x} to e^{-x} , and in both

$$V'_M(x) \sim \left(\frac{1}{\bar{\epsilon}} - 1\right)e^{-x}, \quad x \rightarrow -\infty. \quad (\text{B.3})$$

We adopt R_2 throughout, so that the model as analyzed and the model as priced are the same object.

The constant b anchors R_2 to R_1 .

The definition of b in Eq. (B.2) looks arbitrary but is not. Substituting $x = -4$ gives $\text{Erf}(b - 4) = -[1 - 2/(1 + e^4 g \bar{\epsilon})]$, whence

$$R_2(-4) = \frac{ge^4}{1 + \bar{\epsilon}ge^4} = R_1(-4). \quad (\text{B.4})$$

The two regularizers agree exactly at $x_\star = -4$, and b is precisely the constant that makes them do so. This is a matching condition in the default region, which is where regularization is needed, and it is the reason the substitution is legitimate.

The turning point cannot fix $\bar{\epsilon}$.

Under R_1 the condition $V'_M(\hat{x}) = 0$ is linear in $\bar{\epsilon}$ once the denominator is cleared, giving $\bar{\epsilon} = 1 - e^{\hat{x}}/g$ and inviting the criterion that the mean-reversion range extend to a chosen $\hat{x}$. Under R_2 this fails. The turning point solves $\text{Erf}(\hat{x} + b) = 1 - 2\bar{\epsilon}$, that is

$$\hat{x} = \text{Erf}^{-1}(1 - 2\bar{\epsilon}) - b(g, \bar{\epsilon}), \quad (\text{B.5})$$

and because b carries its own dependence on $\bar{\epsilon}$ the two contributions very nearly cancel. At $g = 1$, $\bar{\epsilon} = 0.1$ we find $\partial\hat{x}/\partial\log\bar{\epsilon} = -0.014$ against $\partial\hat{x}/\partial\log g = +0.389$, so the turning point is some twenty-five times more responsive to the coupling than to the regularizer, and it is not even monotone in $\bar{\epsilon}$. Over $\bar{\epsilon} \in (0.01, 0.45)$ and $g \in (0.1, 10)$ the reachable range of $\hat{x}$ is roughly $(-3.5, -1.6)$, so a target such as $\hat{x} = x_0 - 2$ is in general not attainable at all. Under R_2 the turning point is a property of g , not of $\bar{\epsilon}$, and the criterion must be sought elsewhere.

The default rate cannot fix it either.

Since the regularizer governs the deep region, it is natural to ask whether $\bar{\epsilon}$ can be calibrated to an observed default rate. It cannot, for two reasons. The first is the turning point again: with the barrier top drifting non-monotonically over $\hat{x} \in (-2.73, -2.38)$ as D ranges over two orders of magnitude, a fixed default

threshold lies sometimes inside the trapping region and sometimes outside it, and the resulting first-passage probability is not monotone in $\bar{\epsilon}$. In simulations of Eq. (18) at $\sigma = 1$ with a threshold at $x = -2.5$ we obtain one-year rates of 0.047%, 0.068%, 0.090% and 0.045% as D increases through 50, 200, 800, 1500, which admits no inversion. The second reason is structural and would remain even if the first were repaired: the regularizer exists in order to make the deep region unreachable, so it suppresses precisely the events one would calibrate against. At $\sigma = 0.6$ the same experiment returns a default rate indistinguishable from zero for every D we tried. A quantity whose purpose is to prevent defaults is a poor instrument for setting their frequency.

The criterion, and why it is imposed at x_* .

What $\bar{\epsilon}$ does control, monotonically and by construction, is the strength of the restoring force at depth, which is the quantity that must be bounded if the model is not to impose unrealistic drifts on distressed names. We therefore impose

$$|V'_M(x_*)| = D \quad (\text{B.6})$$

at a reference depth x_* , with D a chosen bound. Taking $x_* = -4$ makes the criterion independent of which regularizer is used, by Eq. (B.4), and has the further advantage that the inversion remains closed form even though R_2 is transcendental. Writing $A = ge^{-x_*}$ and $B = 1 + De^{x_*}$, condition Eq. (B.6) reads $A/(1 + \bar{\epsilon}A) = B$, so that $\bar{\epsilon} = 1/B - 1/A$, which is Eq. (13). Note that a restoring force at depth requires $R(x_*) > 1$, so D is attained on the branch $V'_M(x_*) > 0$. We have verified this against direct evaluation of V'_M under R_2 to fourteen digits.

The admissible range follows from $\bar{\epsilon} > 0$, which is needed to keep R_1 pole-free and to keep R_2 decreasing, and reads

$$g > e^{x_*}(1 + De^{x_*}). \quad (\text{B.7})$$

At $x_* = -4$ this is a weak restriction: $g > 0.018$ for $D = 0$, $g > 0.153$ for $D = 400$, and $g > 0.354$ for $D = 1000$. Setting $D = 0$ recovers $\bar{\epsilon} = 1 - e^{x_*}/g$, which is the turning-point formula with $\hat{x} = x_*$; this is the one turning point that both regularizers share, and it is available only because it sits at the anchor.

The potential and the sign structure.

Integrating R_2 gives V_M in closed form,

$$V_M(x) = \frac{1}{2\bar{\epsilon}} \left\{ \text{Erfc}(b) + e^{-x} [\text{Erf}(b+x) - 1] + e^{b+1/4} [\text{Erf}(b + \frac{1}{2}) - \text{Erf}(b+x + \frac{1}{2})] \right\}, \quad (\text{B.8})$$

up to an additive constant, which is immaterial since only V'_M enters the price drift and only differences of V_M enter the effective potential Eq. (32). For $x > \hat{x}$ the bracket $1 - R(x)$ is positive and $V'_M < 0$; for $x < \hat{x}$ it is negative and $V'_M > 0$. The turning point is therefore a maximum of V_M , the top of the barrier separating the default region from the normal one, rather than a resting point. Consequently the modulation $V'_M(x_t)V'_M(x_s)$ in the kernel Eq. (31) is negative when the two times sit on opposite sides of $\hat{x}$ and positive when they sit on the same side: memory reverses sign across the barrier, which is the sense in which the default region carries its own dynamics.

C Proof of Proposition 2

We first establish that Eq. (5) is correctly derived from the indifference principle, ensuring the proposition addresses the model's fundamental properties rather than an artifact of the derivation. Subsequently, we demonstrate the implications of this equation for a contingent claim written exclusively on x_T .

Validation of the derivation.

Applying the transformation $u = -e^{-\gamma z}v$, the bracket maximized over the hedge h forms a quadratic expression with a negative leading coefficient, yielding the maximizer $h^* = v_s/(\gamma v) + \bar{\mu}_x/(\gamma s\sigma^2)$. By

substituting h^* and dividing by $-e^{-\gamma z}$, the term $s\bar{\mu}_x v_s$ emerges twice with opposite signs—once originating from the uncontrolled generator and once from the maximized bracket—resulting in an exact cancellation. This reflects the standard financial argument that the drift of a traded asset is entirely hedged away, restricting the survival of $\bar{\mu}_x$ strictly to the quadratic source term. Transitioning to the certainty equivalent $C = \gamma^{-1} \log v$, the component $\frac{1}{2}\sigma^2 s^2 \gamma C_s^2$ induced by v_{ss} cancels exactly against $-\frac{1}{2}\gamma\sigma^2 s^2 C_s^2$ arising from v_s^2/v . Consequently, Eq. (5) features quadratic gradient terms exclusively in the hidden factors and none in x , confirming that the x -direction is spanned by the traded asset and thus carries no residual risk. Finally, applying the coordinate transformations $s\partial_s = \partial_x$ and $s^2\partial_{ss} = \partial_{xx} - \partial_x$ converts $\frac{1}{2}\sigma^2 s^2 C_{ss}$ into $\frac{1}{2}\sigma^2(C_{xx} - C_x)$, yielding the discounted drift coefficient $\bar{\eta} = r - \sigma^2/2$. The integrity of these cancellations and the final differential equation have been verified via symbolic algebra.

Implications for an x -only payoff.

Defining the pricing difference as $P = C - C^0$ and subtracting the respective PDEs, we observe that the source term $-\bar{\mu}_x^2/(2\gamma\sigma^2)$ is common to both equations and additive in neither C nor C^0 . Thus, it cancels identically irrespective of the terminal data. Expanding the difference of the quadratic terms yields

$$(C_y)^2 - (C_y^0)^2 = 2C_y^0 P_y + P_y^2, \quad (\text{C.1})$$

with a symmetric expansion holding for the θ derivatives, such that

$$P_t + \frac{1}{2}\sigma^2 P_{xx} + \bar{\eta} P_x - rP + \frac{1}{2}\sigma_y^2 P_{yy} + \mu_y P_y + \gamma\sigma_y^2 C_y^0 P_y + \frac{1}{2}\gamma\sigma_y^2 P_y^2 + (\theta\text{-terms}) = 0, \quad (\text{C.2})$$

subject to the terminal condition $P(T) = \mathcal{V}(x_T)$. Crucially, every term capable of inducing dependence on the hidden factors is strictly proportional to P_y , P_θ , or their corresponding derivatives. Because the terminal payoff $\mathcal{V}$ depends solely on x_T , it follows that $P_y(T) = P_\theta(T) = 0$. Therefore, the trivial solution $P_y \equiv P_\theta \equiv 0$ is consistent with Eq. (C.2), and by uniqueness, it is the exact solution. Eliminating these terms reduces the system to $P_t + \frac{1}{2}\sigma^2 P_{xx} + \bar{\eta} P_x - rP = 0$, recovering the classic Black-Scholes equation.

From a numerical perspective using a Strang splitting scheme, an equivalent argument applies. After applying the Cole-Hopf transformation $w = e^{\gamma C}$, the hidden-factor substep becomes linear in w . The terminal ratio $w/w^0 = e^{\gamma\mathcal{V}(x)}$ is uniform in y and θ , allowing it to commute with any fractional step containing only hidden-factor derivatives, regardless of how those coefficients depend on x . Consequently, the ratio passes through the hidden-factor substep unchanged. The subsequent x -substep, featuring y -independent coefficients, structurally preserves this difference. Neither fractional step can introduce hidden-factor dependence into P .

Scope and limitations of the proposition.

It is important to emphasize that this proposition does not imply $C = C^0$, nor does it suggest that either certainty equivalent is independent of the memory parameters. On the contrary, both exhibit strong dependence on these parameters through the source term. In our numerical experiments using the parameters outlined in Section 6, the gradient C_y spans a range of ± 8 across the computational grid, whereas the difference $C_y - C_y^0$ remains on the order of 10^{-15} within the interior domain. This indicates that the two certainty equivalents differ exclusively by a function of (t, x) . It is this shared y -profile (rather than a flat spatial gradient) that renders the final option price blind to the order flow. Furthermore, this degenerate behavior breaks down if σ becomes a function of y , as the x -substep coefficients would then introduce y -dependence, making $P_y \equiv 0$ inconsistent. Similarly, the proposition ceases to hold if the terminal payoff explicitly incorporates y_T , which would force a non-zero terminal gradient $P_y(T) \neq 0$.

D Indifference pricing with correlated hidden factors

This appendix derives Eq. (5)–Eq. (6). The uncorrelated case is in [Halperin and Itkin, 2025]; what follows carries ρ and $\rho_{x\theta}$ through the same argument. Throughout, s_t is the discounted price, Z_t the discounted

wealth, and the hidden factors are $h = (y, \theta)^\top$.

Hamilton-Jacobi-Bellman equation.

With the correlation structure Eq. (2), the wealth increment $dZ_t = h_t ds_t$ now covaries with both hidden factors, so the generator acquires two cross terms in z that are absent when the Brownian motions are independent. The value function $u(t, z, s, y, \theta)$ satisfies

$$0 = \frac{\partial u}{\partial t} + \frac{1}{2}\sigma^2 s^2 \frac{\partial^2 u}{\partial s^2} + \frac{1}{2}\sigma_y^2 \frac{\partial^2 u}{\partial y^2} + \frac{1}{2}\sigma_\theta^2 \frac{\partial^2 u}{\partial \theta^2} + \rho \sigma \sigma_y s \frac{\partial^2 u}{\partial s \partial y} + \rho_{x\theta} \sigma \sigma_\theta s \frac{\partial^2 u}{\partial s \partial \theta} + s \bar{\mu}_x \frac{\partial u}{\partial s} + \mu_y \frac{\partial u}{\partial y} + \mu_\theta \frac{\partial u}{\partial \theta} + \max_h \left[\frac{1}{2} h^2 s^2 \sigma^2 \frac{\partial^2 u}{\partial z^2} + h \mathcal{B}[u] \right], \quad (\text{D.1})$$

where

$$\mathcal{B}[u] = s^2 \sigma^2 \frac{\partial^2 u}{\partial z \partial s} + s \sigma \sigma_y \rho \frac{\partial^2 u}{\partial z \partial y} + s \sigma \sigma_\theta \rho_{x\theta} \frac{\partial^2 u}{\partial z \partial \theta} + s \bar{\mu}_x \frac{\partial u}{\partial z}. \quad (\text{D.2})$$

The last two terms of Eq. (D.2) beyond the first and fourth are new; setting $\rho = \rho_{x\theta} = 0$ recovers the bracket of [Halperin and Itkin, 2025].

Optimal hedge.

The bracket in Eq. (D.1) is quadratic in h with negative leading coefficient, so the maximizer is $h^* = -\mathcal{B}[u]/(s^2 \sigma^2 u_{zz})$ and the maximized value is $-\mathcal{B}[u]^2/(2s^2 \sigma^2 u_{zz})$. Exponential utility factorizes the value function as $u = -e^{-\gamma z} v(t, s, y, \theta)$, so every z -derivative brings down a factor of γ and the exponential cancels, giving

$$h^* = \frac{1}{\gamma} \frac{\partial_s v}{v} + \frac{\rho \sigma_y}{\gamma s \sigma} \frac{\partial_y v}{v} + \frac{\rho_{x\theta} \sigma_\theta}{\gamma s \sigma} \frac{\partial_\theta v}{v} + \frac{\bar{\mu}_x}{\gamma s \sigma^2}. \quad (\text{D.3})$$

In terms of the certainty equivalent $C = \gamma^{-1} \log v$ and the log-price, this reads

$$s h^* = \underbrace{\frac{\partial C}{\partial x}}_{\text{delta}} + \underbrace{\frac{1}{\sigma} \left(\rho \sigma_y \frac{\partial C}{\partial y} + \rho_{x\theta} \sigma_\theta \frac{\partial C}{\partial \theta} \right)}_{\text{correlation hedge}} + \underbrace{\frac{\bar{\mu}_x}{\gamma \sigma^2}}_{\text{Merton}}. \quad (\text{D.4})$$

The middle term is the whole effect of the correlations on the hedge: the traded asset is used to hedge the part of the hidden noise that it spans. It vanishes with ρ and $\rho_{x\theta}$, and Eq. (D.4) then reduces to delta plus Merton.

Reduction to the certainty equivalent.

Substituting h^* back and writing $q \equiv \rho \sigma_y \partial_y C + \rho_{x\theta} \sigma_\theta \partial_\theta C$, the terms in the resulting equation for v that are quadratic in derivatives organize as $-(P + \gamma v q + R)^2/(2v)$ with $P = \sigma \partial_x v$ and $R = \bar{\mu}_x v/\sigma$. Two cancellations then occur when $v = e^{\gamma C}$ is inserted.

The first is the classical Cole-Hopf cancellation. The term $\frac{1}{2} \sigma^2 \partial_x^2 v$ contributes $\frac{\gamma}{2} \sigma^2 (\partial_x C)^2$, which is removed exactly by $-P^2/(2v)$. The second is new and is what makes the correlated problem tractable: the cross contributions $\rho \sigma \sigma_y \partial_{xy}^2 v$ and $\rho_{x\theta} \sigma \sigma_\theta \partial_{x\theta}^2 v$ generate $\gamma \sigma \partial_x C q$, which is removed exactly by the cross product $-2Pq\gamma v/(2v)$. No term mixing $\partial_x C$ with the hidden gradients survives.

What remains of the quadratic terms is

$$\frac{\gamma}{2} \left[\sigma_y^2 (\partial_y C)^2 + \sigma_\theta^2 (\partial_\theta C)^2 - q^2 \right] = \frac{\gamma}{2} \nabla_h C^\top \tilde{\Sigma} \nabla_h C, \quad (\text{D.5})$$

with $\tilde{\Sigma}$ as in Eq. (6), since expanding q^2 produces $\rho^2 \sigma_y^2 (\partial_y C)^2 + \rho_{x\theta}^2 \sigma_\theta^2 (\partial_\theta C)^2 + 2\rho \rho_{x\theta} \sigma_y \sigma_\theta \partial_y C \partial_\theta C$. The cross product of q with R supplies the two drift adjustments $-\rho \sigma_y \bar{\mu}_x/\sigma$ and $-\rho_{x\theta} \sigma_\theta \bar{\mu}_x/\sigma$ appearing in

Eq. (5), and R^2 supplies the term $-\bar{\mu}_x^2/(2\gamma\sigma^2)$ unchanged. The drift and discount terms in x are unaffected by the correlations and carry over from [Halperin and Itkin, 2025]. This establishes Eq. (5).

The residual covariance is a Schur complement.

Write the full instantaneous covariance of (x, h) in block form, with $\Sigma_{hh} = \text{diag}(\sigma_y^2, \sigma_\theta^2)$ because $\rho_{y\theta} = 0$, and $\Sigma_{hx} = (\rho\sigma\sigma_y, \rho_{x\theta}\sigma\sigma_\theta)^\top$. Then

$$\tilde{\Sigma} = \Sigma_{hh} - \frac{1}{\sigma^2} \Sigma_{hx} \Sigma_{hx}^\top, \quad (\text{D.6})$$

which is precisely Eq. (6). The interpretation is immediate: $\tilde{\Sigma}$ is the conditional covariance of the hidden factors given the traded one, so the quadratic penalty in Eq. (5) charges the investor only for the hidden risk that the traded asset cannot span. The off-diagonal entry is negative whenever ρ and $\rho_{x\theta}$ share a sign, because hedging through the common traded asset induces a negative residual dependence between the two hidden factors even though their driving noises are uncorrelated.

Limiting cases.

At $\rho = \rho_{x\theta} = 0$ we have $q \equiv 0$, so Eq. (D.4) loses its middle term, the drift adjustments vanish, and $\tilde{\Sigma} = \text{diag}(\sigma_y^2, \sigma_\theta^2)$, which returns the quadratic gradient terms $\frac{\gamma}{2}\sigma_y^2(\partial_y C)^2 + \frac{\gamma}{2}\sigma_\theta^2(\partial_\theta C)^2$ of [Halperin and Itkin, 2025].

In the scalar case, with θ absent, $\tilde{\Sigma}$ collapses to $\sigma_y^2(1 - \rho^2)$, so the quadratic term is $\frac{1}{2}\gamma(1 - \rho^2)\sigma_y^2(\partial_y C)^2$ and the effective distortion exponent is $\gamma(1 - \rho^2)$, in agreement with [Henderson and Hobson, 2002; Musiela and Zariphopoulou, 2004]. This is the origin of $p_y = \gamma(1 - \rho^2)$ in the reduced parameter list: the derived combination, not γ itself, is what the pricing equation sees.

The reduced equation.

Adiabatic elimination of θ replaces $f(\theta)$ and $h(\theta)$ by their stationary averages and removes θ from the state. In Eq. (5) this deletes the row and column of $\tilde{\Sigma}$ carrying θ , so the quadratic term becomes $\frac{1}{2}\gamma(1 - \rho^2)\sigma_y^2(\partial_y C)^2$, deletes the θ drift and diffusion terms, and leaves the x and y blocks untouched. The result is Eq. (23). Note that $\rho_{x\theta}$ does not disappear silently: it enters the reduced description only through the transient correction of Section 3.3, at order $1/k$, which is why the baseline reduced model sets it to zero.

E Proof of Proposition 3

The second line of Eq. (29) is linear in y with constant coefficients, so it integrates in closed form,

$$y_t = e^{-\mu t} y_0 + \hat{y}(1 - e^{-\mu t}) - c_y \int_0^t e^{-\mu(t-s)} V_M(x_s) ds + \sigma_y \int_0^t e^{-\mu(t-s)} dW_s^{(y)}, \quad (\text{E.1})$$

with $\hat{y} = \bar{y} + \bar{h}/\mu$ the level to which y relaxes in the absence of coupling. The memory variable is thus an exponentially weighted average of $V_M(x_s)$ along the realized path, with forgetting rate μ .

Substituting Eq. (E.1) into the drift of x gives

$$\bar{f} + \bar{\eta} - c_x \hat{y} V_M'(x_t) + c_x c_y V_M'(x_t) \int_0^t e^{-\mu(t-s)} V_M(x_s) ds - c_x \sigma_y V_M'(x_t) \int_0^t e^{-\mu(t-s)} dW_s^{(y)}, \quad (\text{E.2})$$

where terms carrying $e^{-\mu t}$ have been suppressed; they decay on the scale $1/\mu$ and do not affect the stationary structure. The first two terms are local in the state and supply the first two terms of Eq. (32). The last is the multiplicative noise named in the proposition. The memory sits in the third term, which acts on the level of $V_M(x_s)$ rather than on a velocity and so is not yet in GLE form.

Integrate that term by parts. With $u = V_M(x_s)$ and $dv = e^{-\mu(t-s)} ds$, so that $v = e^{-\mu(t-s)}/\mu$,

$$\int_0^t e^{-\mu(t-s)} V_M(x_s) ds = \frac{V_M(x_t)}{\mu} - \frac{V_M(x_0)e^{-\mu t}}{\mu} - \frac{1}{\mu} \int_0^t e^{-\mu(t-s)} dV_M(x_s). \quad (\text{E.3})$$

By Itô's lemma $dV_M(x_s) = V'_M(x_s) dx_s + \frac{1}{2} V''_M(x_s) \sigma^2 ds$, so

$$\int_0^t e^{-\mu(t-s)} V_M(x_s) ds = \frac{V_M(x_t)}{\mu} - \frac{1}{\mu} \int_0^t e^{-\mu(t-s)} V'_M(x_s) \dot{x}_s ds - \frac{\sigma^2}{2\mu} \int_0^t e^{-\mu(t-s)} V''_M(x_s) ds, \quad (\text{E.4})$$

again dropping the transient. Multiplying by $c_x c_y V'_M(x_t)$, the first term on the right gives

$$\frac{c_x c_y}{\mu} V'_M(x_t) V_M(x_t) = \frac{c_x c_y}{2\mu} \frac{d}{dx} [V_M(x)^2] \Big|_{x=x_t}, \quad (\text{E.5})$$

which is the third term of Eq. (32) and is again local. The second term gives

$$- \frac{c_x c_y}{\mu} V'_M(x_t) \int_0^t e^{-\mu(t-s)} V'_M(x_s) \dot{x}_s ds, \quad (\text{E.6})$$

which is the friction term of an overdamped GLE with the tail part of Eq. (31). The third term of Eq. (E.4) contributes

$$- \frac{c_x c_y \sigma^2}{2\mu} V'_M(x_t) \int_0^t e^{-\mu(t-s)} V''_M(x_s) ds, \quad (\text{E.7})$$

a further memory term acting on the level of V''_M . It is $O(\sigma^2)$, vanishes identically when V_M is linear, and we absorb it into η_t ; it does not affect the structure of the kernel.

Collecting the local contributions into U_{eff} and the noise sources into η_t , the drift equation for x reads

$$\dot{x}_t + \frac{c_x c_y}{\mu} V'_M(x_t) \int_0^t e^{-\mu(t-s)} V'_M(x_s) \dot{x}_s ds = -U'_{\text{eff}}(x_t) + \eta_t. \quad (\text{E.8})$$

Writing the bare term as $\dot{x}_t = \int_0^t \delta(t-s) \dot{x}_s ds$ collects the left-hand side into a single convolution with the kernel K_M of Eq. (31), which is Eq. (30). $\square$

F RBF matrix elements of the reduced model

Appendix A of [Halperin and Itkin, 2025] evaluates the matrix elements of the Volterra kernel in closed form. Its derivation rests on the Green's function being a product of three independent one-dimensional Gaussians, which factorizes every integral into three scalar ones with

$$\int_{-\infty}^{\infty} \frac{e^{-\varepsilon(\xi-x_k)^2 - (x-\xi)^2/(2\sigma^2\Delta\tau)}}{\sigma\sqrt{2\pi\Delta\tau}} d\xi = \frac{1}{a(\sigma)} e^{-\varepsilon(x-x_k)^2/a^2(\sigma)}, \quad a^2(\sigma) = 1 + 2\varepsilon\sigma^2\Delta\tau. \quad (\text{F.1})$$

Under the reduction the θ direction disappears, but the surviving pair (x, y) is correlated through ρ , so the Green's function no longer factorizes and Eq. (F.1) cannot be applied coordinate by coordinate.

One route is to rotate to the eigenbasis of the covariance. This works, because the Gaussian RBF is isotropic and therefore invariant under orthogonal transformations, but it is not necessary: the entire appendix generalizes by promoting the scalar $a^2(\sigma)$ to a matrix, and the resulting formulas contain the uncorrelated ones as the diagonal case.

The master identity.

Write $\mathbf{p} = (\xi, \eta)^\top$ for the integration variable, $\mathbf{u} = (x, y)^\top$ for the evaluation point, $\mathbf{p}_k = (x_k, y_j)^\top$ for an RBF centre, and

$$\Sigma = \begin{pmatrix} \sigma^2 & \rho\sigma\sigma_y \\ \rho\sigma\sigma_y & \sigma_y^2 \end{pmatrix} \quad (\text{F.2})$$

for the diffusion covariance of the reduced model. With Eq. (19) in force $\Sigma = \Sigma(y)$, since σ_x appears in three of its four entries; the formulas below hold verbatim with Σ evaluated at the y -node of the substep, at the cost of one 2×2 inverse per node rather than one overall. The Green's function of the homogeneous part of Eq. (23) is the bivariate Gaussian with covariance $\Sigma\Delta\tau$. Then, for the isotropic Gaussian RBF $e^{-\varepsilon|\mathbf{p}-\mathbf{p}_k|^2}$,

$$\int_{\mathbb{R}^2} e^{-\varepsilon|\mathbf{p}-\mathbf{p}_k|^2} G(\Delta\tau; \mathbf{u} | \mathbf{p}) d\mathbf{p} = \frac{1}{\sqrt{\det \mathbf{A}}} \exp\left[-\varepsilon (\mathbf{u} - \mathbf{p}_k)^\top \mathbf{A}^{-1} (\mathbf{u} - \mathbf{p}_k)\right], \quad \mathbf{A} \equiv \mathbf{I} + 2\varepsilon \Sigma \Delta\tau. \quad (\text{F.3})$$

This follows from the convolution of two Gaussians, the RBF being an unnormalized Gaussian of covariance $(2\varepsilon)^{-1}\mathbf{I}$. When $\rho = 0$ the matrix $\mathbf{A}$ is diagonal with entries $a^2(\sigma)$ and $a^2(\sigma_y)$, and Eq. (F.3) splits into the product of two copies of Eq. (F.1). The reduction therefore amounts to the substitution

$$\prod_{\text{coords}} \frac{1}{a(\cdot)} e^{-\varepsilon(\cdot)^2/a^2(\cdot)} \mapsto \frac{1}{\sqrt{\det \mathbf{A}}} e^{-\varepsilon(\cdot)^\top \mathbf{A}^{-1}(\cdot)}, \quad (\text{F.4})$$

applied wherever Appendix A of [Halperin and Itkin, 2025] produces a product of a -factors, together with the deletion of the $a(\sigma_\theta)$ factor.

Linear terms.

The terms $\mathcal{A}_1$ and $\mathcal{A}_2$ of [Halperin and Itkin, 2025] carry a linear prefactor $(\mathbf{p} - \mathbf{p}_k)$ multiplying the RBF, produced by differentiating the Gaussian. Since $\partial_{\mathbf{p}_k} e^{-\varepsilon|\mathbf{p}-\mathbf{p}_k|^2} = 2\varepsilon(\mathbf{p} - \mathbf{p}_k)e^{-\varepsilon|\mathbf{p}-\mathbf{p}_k|^2}$, these integrals follow from Eq. (F.3) by differentiation with respect to the centre,

$$\int_{\mathbb{R}^2} (\mathbf{p} - \mathbf{p}_k) e^{-\varepsilon|\mathbf{p}-\mathbf{p}_k|^2} G(\Delta\tau; \mathbf{u} | \mathbf{p}) d\mathbf{p} = \mathbf{A}^{-1}(\mathbf{u} - \mathbf{p}_k) \times \frac{e^{-\varepsilon(\mathbf{u}-\mathbf{p}_k)^\top \mathbf{A}^{-1}(\mathbf{u}-\mathbf{p}_k)}}{\sqrt{\det \mathbf{A}}}. \quad (\text{F.5})$$

In the uncorrelated case $\mathbf{A}^{-1}$ is diagonal and each component reduces to the corresponding scalar result. With $\rho \neq 0$ the off-diagonal entries of $\mathbf{A}^{-1}$ mix the two directions, so the x -component of Eq. (F.5) acquires a contribution proportional to $(y - y_j)$ and conversely. This mixing is the only structural change: no new special functions appear and everything remains closed form.

Quadratic terms.

The term $\bar{\Psi}_2$ contains products of two RBFs centred at $\mathbf{p}_k$ and $\mathbf{p}_{k^*}$. Because the RBF is isotropic, such a product is again an isotropic Gaussian,

$$e^{-\varepsilon|\mathbf{p}-\mathbf{p}_k|^2} e^{-\varepsilon|\mathbf{p}-\mathbf{p}_{k^*}|^2} = e^{-\frac{\varepsilon}{2}|\mathbf{p}_k-\mathbf{p}_{k^*}|^2} e^{-2\varepsilon|\mathbf{p}-\bar{\mathbf{p}}|^2}, \quad \bar{\mathbf{p}} = \frac{1}{2}(\mathbf{p}_k + \mathbf{p}_{k^*}), \quad (\text{F.6})$$

so the quadratic matrix elements are given by Eq. (F.3) with ε replaced by 2ε , the centre replaced by the midpoint $\bar{\mathbf{p}}$, and an overall factor $\exp[-\frac{\varepsilon}{2}|\mathbf{p}_k - \mathbf{p}_{k^*}|^2]$. Explicitly, with $\mathbf{A}_2 \equiv \mathbf{I} + 4\varepsilon\Sigma\Delta\tau$,

$$\int_{\mathbb{R}^2} e^{-\varepsilon|\mathbf{p}-\mathbf{p}_k|^2} e^{-\varepsilon|\mathbf{p}-\mathbf{p}_{k^*}|^2} G(\Delta\tau; \mathbf{u} | \mathbf{p}) d\mathbf{p} = \frac{e^{-\frac{\varepsilon}{2}|\mathbf{p}_k-\mathbf{p}_{k^*}|^2}}{\sqrt{\det \mathbf{A}_2}} \exp\left[-2\varepsilon(\mathbf{u} - \bar{\mathbf{p}})^\top \mathbf{A}_2^{-1}(\mathbf{u} - \bar{\mathbf{p}})\right]. \quad (\text{F.7})$$

Linear prefactors in the quadratic block are handled by differentiating Eq. (F.7) with respect to $\bar{\mathbf{p}}$, exactly as in Eq. (F.5).

Distinguishing the covariance matrices.

It is critical to distinguish between the two contexts in which a covariance matrix enters the formulation, as they represent distinct mathematical objects. The Green's function, and consequently the integral representations Eq. (F.3)–Eq. (F.7), incorporates the primary diffusion covariance Σ defined in Eq. (F.2).

Conversely, the quadratic gradient term within the pricing equation is governed by the residual covariance $\tilde{\Sigma}$ of Eq. (6), which simplifies in the reduced model to the scalar $\sigma_y^2(1 - \rho^2)$. As a result, the coefficient scaling C_η^2 inside $\bar{\Phi}$ becomes $\frac{1}{2}\gamma(1 - \rho^2)\sigma_y^2 = \frac{1}{2}p_y\sigma_y^2$, where p_y is the derived parameter introduced in Section 2.2. Meanwhile, the kernel of the integral operator retains σ_y^2 alongside the full correlation structure. Conflating these two matrices would erroneously rescale the Green's function instead of the intended nonlinearity.

Memory-dependent volatility.

The introduction of Eq. (19) preserves the preceding structural arguments. During the x -substep of the Strang splitting scheme, y acts as a frozen parameter. Consequently, the differential operator $\frac{1}{2}\sigma_x^2(y)\partial_{xx} + \bar{\eta}(y)\partial_x - r$ exhibits constant coefficients with respect to x , yielding a Gaussian Green's function with variance $\sigma_x^2(y)\Delta\tau$ and mean shift $\bar{\eta}(y)\Delta\tau$. The corresponding one-dimensional element is therefore evaluated as

$$\int_{\mathbb{R}} e^{-\varepsilon(\xi - x_k)^2} G(\Delta\tau; x | \xi) d\xi = \frac{e^{-r\Delta\tau}}{a(\sigma_x(y))} \exp\left[-\frac{\varepsilon(x - x_k - \bar{\eta}(y)\Delta\tau)^2}{a^2(\sigma_x(y))}\right], \quad a^2 = 1 + 2\varepsilon\sigma_x^2(y)\Delta\tau, \quad (\text{F.8})$$

a result we have verified against direct quadrature to machine precision. Relative to the constant-volatility regime, two computational adjustments are necessary: the scaling factor a becomes an array indexed by the y -nodes rather than remaining a scalar, and the mean shift $\bar{\eta}(y)\Delta\tau$ —previously an absorbable constant—must be explicitly tracked per node. Notably, neither modification compromises the closed-form tractability of the integrals. This highlights a fundamental advantage of the operator splitting framework: if applied directly to the unsplit operator, a y -dependent diffusion coefficient would fundamentally destroy the Gaussian structure of the Green's function.

Summary of the model reduction.

To adapt the framework presented in Appendix A of [Halperin and Itkin, 2025] to the present reduced model, the following systematic modifications are required: first, eliminate the θ integral and the associated scaling factor $a(\sigma_\theta)$ entirely. Second, replace each surviving product of a -factors with the matrix formulation Eq. (F.4), setting $\mathbf{A} = \mathbf{I} + 2\varepsilon\Sigma\Delta\tau$ for the linear terms, or $\mathbf{A}_2 = \mathbf{I} + 4\varepsilon\Sigma\Delta\tau$ within the quadratic block. Third, substitute each scalar linear-prefactor result with the generalized form in Eq. (F.5). Finally, fix the coefficient of the quadratic gradient term at $\frac{1}{2}p_y\sigma_y^2$. Under these modifications, every matrix element remains analytically tractable in closed form. The computational complexity per element is strictly preserved, with the sole addition being the inversion of a 2×2 matrix, an operation performed only once per time step as it is uniformly applicable across all elements.

The final reduced model retains nine parameters for the whole surface. The physical-measure parameters $\bar{f}$ and c are inherited from the time-series calibration, except that c is deliberately released in the option calibration. This is a compromise between statistical identifiability and economic identification: the eigenanalysis shows that $\bar{f}$ is a stiff direction whereas c is substantially softer, but fixing c at its physical-measure value would eliminate the option-implied flow-risk premium altogether. We therefore retain c as a Q-measure calibration parameter and explicitly report its weak identification through profile likelihoods and Hessian eigenvalues. The difference between its Q- and P-measure estimates is consequently interpreted as a market price of flow risk, with the associated uncertainty carried into the economic interpretation.

The SPX surface contains limited information about the absolute level of the flow-impact coefficient. Nevertheless, retaining c in the Q-measure calibration is necessary for identifying the Q–P wedge interpreted as the market price of flow risk. We therefore report the point estimate together with its profile uncertainty and do not interpret the point estimate as a sharply identified structural coefficient.