

Dynamic Physical Hedging amid Jump Losses, Reconstruction-Price Uncertainty, Population Interactions

Paramahansa Pramanik^{1,2}

Michael Bowdin^{3,4}

¹ Department of Mathematics and Statistics, University of South Alabama, Mobile, AL 36688, United States.

² Corresponding author, ppramanik@southalabama.edu

³ Department of Mathematics and Statistics, University of South Alabama, Mobile, AL 36688, United States.

⁴ mb2139@jagmail.southalabama.edu

Abstract

We study dynamic physical hedging for insurers exposed jointly to catastrophe losses and stochastic reconstruction costs. Surplus evolves as a controlled jump diffusion whose loss amplitude combines marked catastrophe severity, an exogenous mean-reverting cost factor, and endogenous mitigation. We establish well-posedness, moment and stability estimates, and a stopping-time dynamic programming principle, and prove that the value function is the unique viscosity solution of the resulting nonlocal Hamilton-Jacobi-Bellman (HJB) equation. Strategic interaction is introduced through a mean field game (MFG) with reduced-form vulnerability costs, yielding a coupled backward-forward HJB-Kolmogorov system. We establish relaxed equilibrium existence, Markovian realization, and uniqueness under appropriate compactness and monotonicity conditions. Numerical experiments show that reconstruction costs and capitalization materially affect optimal hedging and that cross-sectional vulnerability alters equilibrium costs. Tail-family robustness calculations further assess the sensitivity of these conclusions to alternative catastrophe-severity specifications.

Keywords: Mean Field Games; Viscosity Solutions; Stochastic Control; Nonlocal HJB Equations; Catastrophe Insurance.

1 Introduction

The economic problem considered in this paper is the allocation of physical mitigation when an insurer faces two distinct sources of uncertainty: discontinuous catastrophe losses and continuously fluctuating reconstruction prices. For $(t, x, y) \in [0, T] \times \mathbb{R}^2$, an admissible predictable policy $u \in \mathcal{U}_t$ is evaluated through the discounted criterion

$$V(t, x, y) = \inf_{u \in \mathcal{U}_t} \mathbb{E}_{t, x, y} \left[\int_t^T e^{-\delta(s-t)} f(s, X_s, Y_s, u_s) ds + e^{-\delta(T-t)} g(X_T, Y_T) \right]. \quad (1)$$

The controlled state is specified by

$$\begin{aligned} dY_s &= \kappa(\vartheta - Y_s) ds + \sigma_I dW_s^I, \\ dX_s &= b(X_{s-}, Y_{s-}, u_s) ds + \sigma_X(X_{s-}, Y_{s-}, u_s) dW_s^X - \int_E e^{Y_{s-}} a(z) h(u_s) \tilde{N}(ds, dz), \end{aligned} \quad (X_t, Y_t) = (x, y). \quad (2)$$

The first coordinate X represents insurer surplus, whereas $I = e^Y$ is an exogenously evolving reconstruction-cost index whose logarithm reverts toward the long-run level ϑ . Catastrophe events and their marks are generated by the marked Poisson random measure N . The factor $h(u)$ measures the fraction of physical loss remaining after the insurer has committed resources to pre-arranged reconstruction capacity. Consequently, mitigation enters the model at the level of catastrophe severity itself: conditional on (Y, z) , the monetary jump transmitted to surplus is scaled by $h(u)$ rather than being offset only after the loss has occurred.

A central modeling distinction is that the reconstruction-cost process and the catastrophe point process are separate primitive risks. Equation (2) contains no catastrophe-triggered impulse in Y , and a realization of N does not cause the replacement-cost index to increase. The framework therefore analyzes the adverse joint exposure created when a catastrophe occurs while reconstruction inputs are already expensive; it does not impose a causal mechanism through which the catastrophe itself generates a price surge. Within this specification, we establish well-posedness and stability of the controlled jump system, obtain a stopping-time dynamic programming principle, and identify the value function with the unique viscosity solution of the corresponding nonlocal Hamilton-Jacobi-Bellman (HJB) equation. We subsequently introduce cross-sectional strategic interaction through a mean field game of insurers, thereby linking individual mitigation choices to the evolution of aggregate vulnerability. This perspective complements evidence that catastrophe recovery may be accompanied by reconstruction-cost inflation and demand-surge effects [20, 42, 49, 95], while deliberately isolating the control problem generated by stochastic replacement-cost exposure from the separate economic question of how post-catastrophe reconstruction prices are endogenously formed.

The first strand of related work concerns catastrophe insurance, mitigation, reconstruction costs, and stochastic loss models. Classical insurance-risk theory studies reserves, ruin, dividends, and reinsurance under compound-Poisson and diffusion-perturbed surplus processes [5, 6, 7, 29, 21, 31, 32, 34, 50, 96]. Catastrophe-risk research emphasizes that insurance and ex ante mitigation jointly affect recovery and loss reduction [42, 95], while empirical work on reconstruction markets documents post-event price pressure, contractor scarcity, and insured-loss inflation [20, 49]. These studies motivate replacement-cost exposure as an economically distinct source of catastrophe severity. The present model is deliberately narrower Y is an exogenous replacement-cost factor rather than a market-clearing reconstruction price, and aggregate rebuilding demand does not determine its dynamics [57]. On the probabilistic side, catastrophe clustering and transient changes in claim intensity have been modeled through shot-noise, contagion, and related point processes [2, 18, 19, 47, 46, 52], while marked point processes accommodate heterogeneous event severity [105]. Our mechanism differs from intensity-based models because e^Y scales the monetary loss generated by a catastrophe mark, and the control $h(u)$ acts on that same nonlocal loss amplitude. Thus physical hedging changes residual catastrophe severity before the loss enters surplus rather than acting only through a premium, dividend, or reinsurance transfer.

The second strand concerns stochastic control with jumps, dynamic programming, and viscosity solutions of nonlocal HJB equations. General foundations for controlled diffusions and jump diffusions are developed in [4, 30, 48, 51, 97]. For integro-differential HJB equations, comparison and approximation re-

quire a nonlocal viscosity solutions [8, 9, 15, 36]. The stopping-time DPP used here also requires stability under conditioning and concatenation together with measurable selection. We therefore rely on the canonical control-correspondence and analytic-set frameworks of El Karoui and Tan [24], Žitković [106], Fayvisovich and Žitković [28], rather than attributing these general principles to application-specific results. Relaxed-control compactification and approximation are treated through the classical framework of El Karoui et al. [23]. Numerically, the relevant principle is monotone approximation of viscosity solutions, together with policy iteration and positivity-preserving discretization of the forward equation [70]. The computational construction below is therefore designed around a monotone nonlocal HJB discretization and the adjoint Kolmogorov operator rather than a Gaussian approximation of the jump term.

The third strand is mean field games and large-population stochastic control [92]. The equilibrium consistency principle originates with Lasry and Lions [45], while probabilistic formulations and large-population limits are developed in Lacker [44], Carmona and Delarue [10]. Mean field models of systemic risk illustrate how cross-sectional distributions can feed back into individual incentives [11], whereas models with explicit market clearing show how a population interaction may instead generate an endogenous equilibrium price [76]. This distinction is central here, our couplings F and G are reduced-form vulnerability costs, not market-clearing reconstruction prices. Nonlocal mean field systems with jump or fractional operators require additional analytical and computational structure [13, 14]. The present paper combines these nonlocal-control and mean field structure in a catastrophe-insurance setting in which physical hedging changes jump severity, the induced control changes the cross-sectional law of capitalization and replacement-cost exposure, and that law feeds back into individual costs. Common catastrophe noise, endogenous reconstruction prices, and population-dependent catastrophe intensities are not included and are treated as structural extensions rather than features of the current equilibrium.

The contributions are fourfold. First, we provide a probabilistic foundation for the catastrophe-insurance control problem under diffusion and marked jump risk, including strong well-posedness, moment estimates, stability with respect to initial states and controls, a restart property, and the measurable continuation machinery required for dynamic programming. These results are stated under moment conditions that permit heavy-tailed catastrophe marks up to a prescribed order rather than imposing a Gaussian approximation [90]. Second, we derive the nonlocal HJB equation associated with (1)-(2) and prove that the value function is its unique viscosity solution in an exponential-polynomial growth class. The comparison argument treats the control-dependent nonlocal loss operator generated by physical hedging. Third, we formulate a mean field game of catastrophe insurers, characterize equilibrium through a coupled backward nonlocal HJB equation and forward Kolmogorov equation, and establish existence by a fixed-point argument and uniqueness under a Lasry-Lions [45] type monotonicity condition. Fourth, we translate the analytical structure into economic comparative statics and a calibrated finite-difference experiment. The model implies that an elevated replacement-cost state increases every finite moment of monetary catastrophe-loss exposure, whereas physical hedging reduces the same exposure through the residual-loss function h . Numerically, optimal hedging is stronger when replacement costs are elevated and when insurers are weakly capitalized. The mean field coupling changes the equilibrium valuation of vulnerability even when its effect on the pointwise hedging rule is modest. These conclusions concern the interaction between catastrophe losses and stochastic replacement-cost risk; they do not require, and should not be interpreted as establishing, a causal catastrophe-to-price mechanism [87].

The remainder of the paper is organized as follows. Section 2 develops the stochastic basis of the model and establishes well-posedness, moment and stability estimates, stopping-time restart properties, measurable

selection, and the probabilistic ingredients of the dynamic programming principle. Section 3 specializes this framework to catastrophe insurance, introduces the demand-surge loss specification and the physical-hedging objective, and identifies the nonlocal channel through which the control changes catastrophe severity. Section 4 proves the dynamic programming principle, derives the nonlocal HJB equation, and establishes the viscosity comparison, existence, and uniqueness results. Section 5 introduces the population interaction and proves existence and uniqueness of the mean field equilibrium. Section 6 develops the economic implications for demand-surge exposure, optimal hedging, and equilibrium vulnerability. Section 7 presents the calibration, numerical scheme, and comparative-statics results. Section 8 concludes, and the Appendix collects auxiliary arguments.

2 Probabilistic Foundations

In this section we discuss the stochastic basis on which the catastrophe-insurance control problem is posed and establishes the well-posedness properties required later for dynamic programming, viscosity characterization, and the mean field fixed-point argument. The formulation is deliberately stated at a level that permits both finite-activity catastrophe arrivals and infinite-activity jump perturbations. In particular, catastrophe severity is represented by a marked Poisson random measure, while the post-disaster replacement-cost factor enters the loss amplitude as a state variable. Physical hedging acts directly on the jump diffusion.

Throughout, $T \in (0, \infty)$ is fixed, $E := \mathbb{R}_+ \setminus \{0\}$ is the mark space, and $|\cdot|$ denotes the Euclidean norm. Constants denoted by C , C_p , or K_p may change from line to line. For a probability measure μ on a Polish space, $\mathcal{P}_p$ denotes the set of probability measures with finite p th moment.

Assumption 2.1. Let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a complete filtered probability space satisfying $\mathbb{F} = (\mathcal{F}_t)_{0 \leq t \leq T}$. The space supports a two-dimensional Brownian motion $W_t = (W_t^X, W_t^I)$, and an $\mathbb{F}$ -Poisson random measure $N(dt, dz)$ on $[0, T] \times E$ with compensator $\mathbf{v}(dz) dt$. We define $\tilde{N}(dt, dz) := N(dt, dz) - \mathbf{v}(dz) dt$. The Brownian motion and N are independent. The Lévy measure $\mathbf{v}$ satisfies $\int_E (1 \wedge z^2) \mathbf{v}(dz) < \infty$. For some fixed $p_\star > 2$, $\int_{\{z > 1\}} z^{p_\star} \mathbf{v}(dz) < \infty$.

Assumption 2.2. The logarithmic construction-cost factor Y satisfies

$$dY_s = \kappa(\vartheta - Y_s) ds + \sigma_I dW_s^I, \quad Y_t = y, \quad (3)$$

where $\kappa > 0$, $\sigma_I > 0$, and $\vartheta \in \mathbb{R}$. The observable replacement-cost index is $I_s := e^{Y_s}$. Thus $I_s > 0$ almost surely for every s , avoiding the economically artificial possibility of a negative construction-cost index that would arise from applying an Ornstein-Uhlenbeck process directly to the level.

Assumption 2.3. Let $U \subset \mathbb{R}_+$ be nonempty, convex, and closed. For $(t, x, y) \in [0, T] \times \mathbb{R} \times \mathbb{R}$, an admissible physical-hedging control is an $\mathbb{F}$ -predictable U -valued process $u = (u_s)_{s \in [t, T]}$ satisfying $\mathbb{E} \left\{ \int_t^T |u_s|^{p_\star} ds \right\} < \infty$. The set of admissible controls is denoted by $\mathcal{U}_t$.

Assumption 2.4 . The insurer surplus process X is governed by

$$dX_s = b(X_{s-}, Y_{s-}, u_s) ds + \sigma_X(X_{s-}, Y_{s-}, u_s) dW_s^X - \int_E \ell(Y_{s-}, z, u_s) \tilde{N}(ds, dz), \quad X_t = x. \quad (4)$$

The measurable functions $b : \mathbb{R}^2 \times U \rightarrow \mathbb{R}$, $\sigma_X : \mathbb{R}^2 \times U \rightarrow \mathbb{R}$, and $\ell : \mathbb{R} \times E \times U \rightarrow \mathbb{R}_+$ satisfy the following conditions.

- For some $L > 0$ and all $x, x', y, y' \in \mathbb{R}$ and $u \in U$,

$$|b(x, y, u) - b(x', y', u)| + |\sigma_X(x, y, u) - \sigma_X(x', y', u)| \leq L(|x - x'| + |y - y'|), \quad (5)$$

and

$$|b(x, y, u)| + |\sigma_X(x, y, u)| \leq L(1 + |x| + |y| + |u|). \quad (6)$$

- There exists a measurable $\rho : E \rightarrow [0, \infty)$ such that

$$\int_E (\rho(z)^2 + \rho(z)^{p_*}) \nu(dz) < \infty, \quad (7)$$

and

$$|\ell(y, z, u) - \ell(y', z, u)| \leq L|y - y'| \rho(z), \quad |\ell(y, z, u)| \leq L(1 + e^y + |u|) \rho(z). \quad (8)$$

Assumption 2.5. For ν -a.e. $z \in E$, the mapping $u \mapsto \ell(y, z, u)$ is nonincreasing for every $y \in \mathbb{R}$ and the mapping $y \mapsto \ell(y, z, u)$ is nondecreasing for every $u \in U$. Moreover, there exist measurable functions $a : E \rightarrow (0, \infty)$ and $h : U \rightarrow (0, 1]$ such that the benchmark specification

$$\ell(y, z, u) = e^y a(z) h(u) \quad (9)$$

is admissible, with h locally Lipschitz and nonincreasing. Hence higher construction costs amplify the monetary loss generated by a catastrophe mark, whereas stronger physical hedging attenuates that amplification.

Assumption 2.6. The running cost $f : [0, T] \times \mathbb{R}^2 \times U \rightarrow \mathbb{R}$ and terminal cost $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ are continuous and satisfy, for some $q \in [2, p_*)$,

$$|f(s, x, y, u)| + |g(x, y)| \leq C \left(1 + |x|^q + e^{q|y|} + |u|^q \right). \quad (10)$$

In addition, for some $c_0 > 0$ and $c_1 \geq 0$,

$$f(s, x, y, u) \geq c_0 |u|^2 - c_1 \left(1 + |x|^2 + e^{2|y|} \right). \quad (11)$$

The coercivity condition represents increasing marginal scarcity costs of pre-arranged reconstruction capacity.

Remark 1. The preceding assumptions are standard in the theory of jump-diffusion control, but their interpretation here is specific to catastrophe insurance. The mark z records catastrophe severity, e^{Y_s} records the prevailing replacement-cost environment, and u_s is the insurer's physical hedge. Assumption 2.5 permits a single disaster to be economically more damaging during a construction-cost spike and less damaging when reconstruction inputs have been secured in advance.

Lemma 1 (Exponential moments of the construction-cost factor). *Under Assumption 2.2, for every $m \geq 1$ and every $(t, y) \in [0, T] \times \mathbb{R}$,*

$$\mathbb{E}_{t,y} \left[\sup_{t \leq s \leq T} |Y_s|^m \right] \leq C_m(1 + |y|^m), \quad (12)$$

and, for every $r > 0$,

$$\mathbb{E}_{t,y} \left[\sup_{t \leq s \leq T} e^{r|Y_s|} \right] \leq C_{r,T} \exp(C_{r,T}|y|). \quad (13)$$

Consequently, every positive and negative polynomial moment of $I_s = e^{Y_s}$ is finite on $[t, T]$.

Proposition 2 (Well-defined catastrophe jump integral). *Suppose Assumptions 2.1-2.5 hold and let $u \in \mathcal{U}_t$. If Y is the solution of (3), then*

$$\mathbb{E} \left[\int_t^T \int_E |\ell(Y_s, z, u_s)|^2 \mathbf{v}(dz) ds \right] < \infty. \quad (14)$$

Moreover, for every $p \in [2, p_\star]$,

$$\mathbb{E} \left[\int_t^T \int_E |\ell(Y_s, z, u_s)|^p \mathbf{v}(dz) ds \right] < \infty. \quad (15)$$

Consequently, $\int_t^\cdot \int_E \ell(Y_{s-}, z, u_s) \tilde{N}(ds, dz)$ is a well-defined càdlàg local martingale, and it is square-integrable for $p = 2$.

Theorem 3 (Strong well-posedness of the controlled catastrophe-insurance state). *Let Assumptions 2.1-2.5 hold and fix $(t, x, y) \in [0, T] \times \mathbb{R}^2$ and $u \in \mathcal{U}_t$. Then Assumptions 2.2-2.4 admit a pathwise unique strong càdlàg solution $(X_s^{t,x,y;u}, Y_s^{t,y})_{s \in [t, T]}$. For every $p \in [2, p_\star]$,*

$$\mathbb{E} \left[\sup_{t \leq s \leq T} |X_s^{t,x,y;u}|^p \right] \leq C_p \left[1 + |x|^p + |y|^p + e^{C_p|y|} + \mathbb{E} \left[\int_t^T |u_s|^p ds \right] \right]. \quad (16)$$

In particular, the controlled surplus cannot explode in finite time.

Corollary 4. *Under the hypotheses of Theorem 3, for every $p \in [2, p_\star]$, $(X^{t,x,y;u}, Y^{t,y}) \in \mathcal{S}^p([t, T]; \mathbb{R}^2)$, where*

$$\mathcal{S}^p([t, T]; \mathbb{R}^2) := \left\{ Z : Z \text{ is adapted and càdlàg, } \mathbb{E} \left[\sup_{t \leq s \leq T} |Z_s|^p \right] < \infty \right\}.$$

In particular, the objective functional

$$J(t, x, y; u) := \mathbb{E} \left[\int_t^T e^{-\delta(s-t)} f(s, X_s, Y_s, u_s) ds + e^{-\delta(T-t)} g(X_T, Y_T) \right] \quad (17)$$

is well-defined for every $u \in \mathcal{U}_t$ satisfying $\mathbb{E} \left[\int_t^T |u_s|^{p_\star} ds \right] < \infty$.

Proof. The $\mathcal{S}^p$ integrability follows directly from Theorem 3 and Lemma 1. By Assumption 2.6,

$$|f(s, X_s, Y_s, u_s)| + |g(X_T, Y_T)| \leq C \left(1 + |X_s|^q + e^{q|Y_s|} + |u_s|^q + |X_T|^q + e^{q|Y_T|} \right).$$

Because $q < p_*$, Theorem 3, Lemma 1, and Hölder's inequality imply integrability of the right-hand side. Since the discount factor is bounded above by one, $J(t, x, y, u)$ is finite in absolute value. $\square$

Proposition 5. *Consider Assumptions 2.1-2.5. Let $(x, y), (x', y') \in \mathbb{R}^2$, and let the same control $u \in \mathcal{U}_t$ and the same noises (W, N) drive the corresponding solutions. Then, for every $p \in [2, p_*]$,*

$$\mathbb{E} \left[\sup_{t \leq s \leq T} \left| X_s^{t,x,y;u} - X_s^{t,x',y';u} \right|^p \right] \leq C_p (|x - x'|^p + |y - y'|^p) \Xi_p(y, y'), \quad (18)$$

where $\Xi_p(y, y') := 1 + \exp(C_p|y|) + \exp(C_p|y'|)$. Moreover,

$$\sup_{t \leq s \leq T} |Y_s^{t,y} - Y_s^{t,y'}| \leq |y - y'| \quad \text{a.s.} \quad (19)$$

Remark 2. *The above discussions estimate control perturbations of the initial insurance state. For dynamic programming one also needs stability under perturbations of the hedging policy, a restart property at stopping times, and a measurable way of choosing nearly optimal continuation policies. We impose the following additional regularity. It is economically feasible that small changes in the amount of pre-contracted reconstruction capacity should not produce discontinuous changes in premium income, diffusion exposure, or catastrophe-loss attenuation.*

Assumption 2.7 (Regularity in the physical-hedging control). There exists $L_u > 0$ such that, for all $(x, y) \in \mathbb{R}^2$, $u, v \in U$, and v-a.e. $z \in E$,

$$|b(x, y, u) - b(x, y, v)| + |\sigma_X(x, y, u) - \sigma_X(x, y, v)| \leq L_u |u - v|,$$

and

$$|\ell(y, z, u) - \ell(y, z, v)| \leq L_u |u - v| \rho(z).$$

Moreover, for every $R > 0$, the restrictions of f to $[0, T] \times [-R, R]^2 \times (U \cap [-R, R])$ and of g to $[-R, R]^2$ are uniformly continuous.

For $p \in [2, p_*]$ and $u, v \in \mathcal{U}_t$, define $d_{p,t}(u, v) := \left(\mathbb{E} \int_t^T |u_s - v_s|^p ds \right)^{1/p}$.

Lemma 6 (Stability under perturbations of physical hedging). *Suppose Assumptions 2.1-2.5 and 2.7 hold. Fix (t, x, y) and let $u, v \in \mathcal{U}_t$. For every $p \in [2, p_*]$,*

$$\mathbb{E} \left[\sup_{t \leq s \leq T} \left| X_s^{t,x,y;u} - X_s^{t,x,y;v} \right|^p \right] \leq C_p \mathbb{E} \left[\int_t^T |u_s - v_s|^p ds \right]. \quad (20)$$

Consequently, $d_{p,t}(u^n, u) \rightarrow 0$ implies $X^{t,x,y;u^n} \rightarrow X^{t,x,y;u}$ in $\mathcal{S}^p([t, T])$.

Proposition 7. Under Assumptions 2.1-2.7, let $u^n, u \in \mathcal{U}_t$ satisfy $d_{p^*,t}(u^n, u) \rightarrow 0$ and $\sup_{n \geq 1} \mathbb{E} \left[\int_t^T |u_s^n|^{p^*} ds \right] < \infty$. Then $J(t, x, y; u^n) \rightarrow J(t, x, y; u)$.

If τ is an $\mathbb{F}$ -stopping time taking values in $[t, T]$, define, for $r \geq 0$ with $\tau + r \leq T$, $W_r^\tau := W_{\tau+r} - W_\tau$, with $N^\tau((0, r] \times A) := N((\tau, \tau + r] \times A)$. By the strong Markov property of Brownian motion and of a Poisson random measure with deterministic compensator, conditionally on $\mathcal{F}_\tau$, the shifted noises have the same characteristics as the original noises and are independent of the pre- τ increments [91]. For controls $u \in \mathcal{U}_t$ and $v \in \mathcal{U}_\tau$, their concatenation at τ is $(u \otimes_\tau v)_s := u_s \mathbf{1}_{\{s < \tau\}} + v_s \mathbf{1}_{\{s \geq \tau\}}$.

Lemma 8. If $u \in \mathcal{U}_t$, τ is an $[t, T]$ -valued stopping time, and $v \in \mathcal{U}_\tau$, then $u \otimes_\tau v \in \mathcal{U}_t$. More generally, if $(A_k)_{k \geq 1}$ is an $\mathcal{F}_\tau$ -measurable partition and $v^k \in \mathcal{U}_\tau$, then $v_s := \sum_{k \geq 1} \mathbf{1}_{A_k} v_s^k$, for all $s \geq \tau$, is predictable after τ whenever

$$\sum_{k \geq 1} \mathbb{E} \left[\mathbf{1}_{A_k} \int_\tau^T |v_s^k|^{p^*} ds \right] < \infty.$$

Proof. Predictability of $u \otimes_\tau v$ follows from the standard stability of predictable processes under stopping and pasting at a stopping time. Its integrability follows from

$$\mathbb{E} \left[\int_t^T |(u \otimes_\tau v)_s|^{p^*} ds \right] = \mathbb{E} \left[\int_t^\tau |u_s|^{p^*} ds \right] + \mathbb{E} \left[\int_\tau^T |v_s|^{p^*} ds \right] < \infty.$$

For the partitioned control, $\mathbf{1}_{A_k} \mathbf{1}_{\{s > \tau\}} v_s^k$ is predictable for each k , and the pointwise sum is predictable because the A_k form a disjoint partition. Tonelli's theorem gives

$$\mathbb{E} \left[\int_\tau^T |v_s|^{p^*} ds \right] = \sum_{k \geq 1} \mathbb{E} \left[\mathbf{1}_{A_k} \int_\tau^T |v_s^k|^{p^*} ds \right],$$

which is finite by hypothesis. □

Proposition 9. Let Assumptions 2.1-2.7 hold, let τ be an $[t, T]$ -valued stopping time, and let $u \in \mathcal{U}_t$. Then, up to indistinguishability, the post- τ trajectory satisfies

$$(X_s^{t,x,y;u}, Y_s^{t,y}) = \left(X_s^\tau, X_\tau^{t,x,y;u}, Y_\tau^{t,y}; u, Y_s^\tau, Y_\tau^{t,y} \right), \quad \tau \leq s \leq T, \quad (21)$$

where the system on the right is understood with the shifted noises (W^τ, N^τ) and the restriction of u to $[\tau, T]$. If $v \in \mathcal{U}_\tau$, then the same identity holds with $u \otimes_\tau v$ on the left and v in the restarted system on the right.

Proposition 10. Let τ be as in Proposition 9. For every Borel functional $\Phi : D([0, T]; \mathbb{R}^2) \rightarrow \mathbb{R}$ for which the expectations below are finite,

$$\mathbb{E}[\Phi((X_s, Y_s)_{\tau \leq s \leq T}) | \mathcal{F}_\tau] = \mathcal{R}_\Phi(\tau, X_\tau, Y_\tau; u^{\tau, \omega}) \quad a.s., \quad (22)$$

where $\mathcal{R}_\Phi(r, \xi, \eta; v)$ denotes expectation of Φ under the system restarted at (r, ξ, η) with continuation control v , and $u^{\tau, \omega}$ denotes the post- τ control section obtained by freezing the pre- τ history.

Remark 3. *The state process is not, in general, Markov under an arbitrary open-loop adapted control, because the future control may retain information from the entire past. The correct Markov statement is obtained after fixing a feedback rule.*

Assumption 2.8. Let $\mathfrak{A}$ be the class of Borel maps $\alpha : [0, T] \times \mathbb{R}^2 \rightarrow U$ such that, for some constant L_α , $|\alpha(s, x, y) - \alpha(s, x', y')| \leq L_\alpha(|x - x'| + |y - y'|)$ and $|\alpha(s, x, y)| \leq L_\alpha(1 + |x| + |y|)$. For $\alpha \in \mathfrak{A}$, write $u_s^\alpha := \alpha(s, X_{s-}, Y_{s-})$.

Theorem 11. *Suppose Assumptions 2.1-2.5, 2.7, and 2.8 hold. For each $\alpha \in \mathfrak{A}$, the controlled state $Z^\alpha := (X^\alpha, Y)$ is a time-inhomogeneous strong Markov process. Its transition operators*

$$P_{t,s}^\alpha \phi(x, y) := \mathbb{E}_{t,x,y}[\phi(X_s^\alpha, Y_s)], \quad 0 \leq t \leq s \leq T,$$

map $C_b(\mathbb{R}^2)$ into $C_b(\mathbb{R}^2)$. Thus Z^α is Feller.

Corollary 12 (Chapman-Kolmogorov identity). *Under the hypotheses of Theorem 11, for $0 \leq t \leq r \leq s \leq T$ and $\phi \in B_b(\mathbb{R}^2)$, $P_{t,s}^\alpha \phi = P_{t,r}^\alpha P_{r,s}^\alpha \phi$.*

Proof. Apply the tower property at time r and then the Markov property:

$$P_{t,s}^\alpha \phi(x, y) = \mathbb{E}_{t,x,y}[\mathbb{E}[\phi(Z_s^\alpha) \mid \mathcal{F}_r]] = \mathbb{E}_{t,x,y}[P_{r,s}^\alpha \phi(Z_r^\alpha)].$$

The right-hand side equals $P_{t,r}^\alpha P_{r,s}^\alpha \phi(x, y)$. This completes the proof. $\square$

Uniform integrability is needed twice in the sequel. First to pass from deterministic-time decompositions to stopping-time decompositions, and second to justify limits of ε -optimal controls. For $R > 0$, define the bounded-moment control class

$$\mathcal{U}_t(R) := \left\{ u \in \mathcal{U}_t : \mathbb{E} \left[\int_t^T |u_s|^{p^*} ds \right] \leq R \right\}.$$

Proposition 13 (Uniform integrability of state and cost families). *Suppose Assumptions 2.1-2.7 hold. Fix a compact set $K \subset \mathbb{R}^2$ and $R > 0$. Then there exists $\eta > 0$ such that*

$$\sup_{\substack{(x,y) \in K \\ u \in \mathcal{U}_t(R)}} \mathbb{E} \left[\sup_{t \leq s \leq T} |X_s^{t,x,y;u}|^{q(1+\eta)} + \sup_{t \leq s \leq T} e^{q(1+\eta)|Y_s^{t,y}|} + \int_t^T |u_s|^{q(1+\eta)} ds \right] < \infty. \quad (23)$$

Consequently, the families $\{g(X_T^{t,x,y;u}, Y_T^{t,y})\}_{(x,y) \in K, u \in \mathcal{U}_t(R)}$ and $\left\{ \int_t^T |f(s, X_s^{t,x,y;u}, Y_s^{t,y}, u_s)| ds \right\}_{(x,y) \in K, u \in \mathcal{U}_t(R)}$ are uniformly integrable.

To formulate measurable selection without imposing an artificial topology on individual predictable processes, we use the standard canonical-law formulation of stochastic control. Let

$$\Omega^\circ := D([0, T]; \mathbb{R}^2) \times D([0, T]; \mathbb{R}^2) \times \mathcal{M}([0, T] \times E) \times \mathcal{V},$$

where $\mathcal{V}$ is the space of relaxed controls on $[0, T] \times U$ endowed with the stable topology. The strict control u is embedded as $q^u(ds, da) = ds \delta_{u_s}(da)$. The coordinate processes encode the state, Brownian noise, jump measure, and control measure. We write $\mathfrak{P}(t, x, y)$ for the collection of probability laws on Ω° under which the coordinates solve the controlled martingale problem associated with Assumptions 2.1–2.7, start from (x, y) at time t , and satisfy the $p_\star$ -moment admissibility condition.

Assumption 2.9 (Compact relaxation structure). In addition to Assumptions 2.1–2.8, suppose that:

- (i) The control set U is a compact convex subset of $\mathbb{R}_+$.
- (ii) For every $(t, x, y, z) \in [0, T] \times \mathbb{R}^2 \times E$, the maps $u \mapsto b(x, y, u)$, $u \mapsto \sigma_X^2(x, y, u)$, $u \mapsto \ell(y, z, u)$, and $u \mapsto f(t, x, y, u)$ are continuous. The continuity is locally uniform in (t, x, y) , and the growth bounds in Assumptions 2.4 and 2.6 hold uniformly over $u \in U$.
- (iii) For every compact $K \subset \mathbb{R}^2$, $\lim_{\delta \downarrow 0} \sup_{\substack{(x, y) \in K \\ |u-v| \leq \delta}} \int_E |\ell(y, z, u) - \ell(y, z, v)|^2 \mathbf{v}(dz) = 0$, and

$$\lim_{\delta \downarrow 0} \sup_{\substack{(x, y) \in K \\ |u-v| \leq \delta}} \int_E |\ell(y, z, u) - \ell(y, z, v)|^{p_\star} \mathbf{v}(dz) = 0.$$

- (iv) Relaxed controls are predictable kernels $q_s(da) ds$ taking values in $\mathcal{P}(U)$, and the relaxed generator is defined by $\mathcal{A}^{q_s} \varphi(x, y) := \int_U \mathcal{A}^a \varphi(x, y) q_s(da)$, for all $\varphi \in C_c^2(\mathbb{R}^2)$.

Remark 4. Assumption 2.9 is stronger than the assumptions needed for the well-posedness results above. Its purpose is specific: compactness of U and continuity of the controlled characteristics permit the relaxed-control space to be compactified and allow relaxed controls to be approximated by rapidly switching strict controls. We therefore no longer assume analyticity, conditioning stability, concatenation stability, or strict-control density as primitive hypotheses. The first three properties are verified in Proposition 14, while strict-control density is proved in Proposition 15. The compactness and continuity conditions above are precisely the additional structure used below to verify the analytic control correspondence and to invoke the classical chattering approximation for relaxed stochastic controls [23, 24, 106].

Proposition 14. Suppose Assumptions 2.1–2.9 hold. Let $\mathfrak{P}(t, x, y)$ denote the canonical family of relaxed controlled laws introduced above. Then

- (a) the graph $\text{Gr}(\mathfrak{P}) := \{(t, x, y, P) : P \in \mathfrak{P}(t, x, y)\}$ is analytic in $[0, T] \times \mathbb{R}^2 \times \mathfrak{P}(\Omega^\circ)$;
- (b) if τ is an $[t, T]$ -valued stopping time and $P \in \mathfrak{P}(t, x, y)$, then a regular conditional probability distribution $P^{\tau, \omega}$ can be chosen such that $P^{\tau, \omega} \in \mathfrak{P}(\tau(\omega), X_\tau(\omega), Y_\tau(\omega))$ for P -a.e. ω ;
- (c) if $\omega \mapsto Q_\omega \in \mathfrak{P}(\tau(\omega), X_\tau(\omega), Y_\tau(\omega))$ is a universally measurable continuation kernel satisfying the admissibility moment condition, then $P \otimes_\tau Q \in \mathfrak{P}(t, x, y)$.

Hence, the canonical family is an analytic control correspondence stable under conditioning and concatenation.

Proposition 15. Suppose Assumptions 2.1-2.9 hold. Let $P \in \mathfrak{P}(t, x, y)$ be generated by an admissible relaxed control $q_s(da)ds$. Then there exists a sequence of strict predictable controls $(u^n)_{n \geq 1} \subset \mathcal{U}_t$ such that their occupation measures

$$q^{u^n}(ds, da) := ds \delta_{u_s^n}(da)$$

converge stably to $q_s(da)ds$ and, if (X^n, Y^n) denotes the corresponding strict controlled state,

$$(X^n, Y^n) \Longrightarrow (X, Y) \quad \text{in } D([t, T]; \mathbb{R}^2). \quad (24)$$

Moreover,

$$J(t, x, y; u^n) \longrightarrow J(t, x, y; P). \quad (25)$$

Consequently, strict controls are dense in the relaxed admissible class for the topology relevant to the value function.

Corollary 16. Under Assumptions 2.1-2.9,

$$V^{\text{str}}(t, x, y) = V^{\text{rel}}(t, x, y), \quad (t, x, y) \in [0, T] \times \mathbb{R}^2. \quad (26)$$

Proof. Every strict control defines a relaxed control through $q^u(ds, da) = ds \delta_{u_s}(da)$, so $V^{\text{rel}}(t, x, y) \leq V^{\text{str}}(t, x, y)$. Conversely, fix $\varepsilon > 0$ and choose an admissible relaxed law P^ε such that $J(t, x, y; P^\varepsilon) \leq V^{\text{rel}}(t, x, y) + \varepsilon$. By Proposition 15, there exists a sequence of strict controls u^n satisfying $J(t, x, y; u^n) \longrightarrow J(t, x, y; P^\varepsilon)$. Therefore, $V^{\text{str}}(t, x, y) \leq J(t, x, y; P^\varepsilon) \leq V^{\text{rel}}(t, x, y) + \varepsilon$. Letting $\varepsilon \downarrow 0$ proves (26). $\square$

Theorem 17. Suppose Assumptions 2.1-2.9 hold and let

$$V^{\text{rel}}(t, x, y) := \inf_{P \in \mathfrak{P}(t, x, y)} J(t, x, y; P),$$

where

$$J(t, x, y; P) := \mathbb{E}^P \left[\int_t^T e^{-\delta(s-t)} \int_U f(s, X_s, Y_s, a) q_s(da) ds + e^{-\delta(T-t)} g(X_T, Y_T) \right].$$

Then V^{rel} is lower semianalytic. Moreover, for every $\varepsilon > 0$, there exists a universally measurable kernel $(t, x, y) \longmapsto P_{t, x, y}^\varepsilon$ such that $P_{t, x, y}^\varepsilon \in \mathfrak{P}(t, x, y)$ and

$$J(t, x, y; P_{t, x, y}^\varepsilon) \leq V^{\text{rel}}(t, x, y) + \varepsilon \quad (27)$$

whenever $V^{\text{rel}}(t, x, y) < \infty$.

Corollary 18. Let τ be an $[t, T]$ -valued stopping time. Under the hypotheses of Theorem 17, for every $\varepsilon > 0$ the post- τ continuation law may be selected as $P_{\tau, X_\tau, Y_\tau}^\varepsilon$ in a universally measurable manner. If $P \in \mathfrak{P}(t, x, y)$ is the pre- τ controlled law, then $P \otimes_\tau P_{\tau, X_\tau, Y_\tau}^\varepsilon \in \mathfrak{P}(t, x, y)$, and its conditional continuation cost satisfies $J(\tau, X_\tau, Y_\tau; P_{\tau, X_\tau, Y_\tau}^\varepsilon) \leq V^{\text{rel}}(\tau, X_\tau, Y_\tau) + \varepsilon$ a.s.

Proof. Set $\Xi_\tau := (\tau, X_\tau, Y_\tau)$. Since Ξ_τ is measurable with respect to the universally completed σ -field and $\xi \mapsto P_\xi^\varepsilon$ is universally measurable by Theorem 17, the composition $\omega \mapsto P_{\Xi_\tau(\omega)}^\varepsilon$ is a universally measurable

stochastic kernel. Proposition 14 gives stability of $\mathfrak{P}$ under concatenation, hence $P \otimes_{\tau} P_{\Xi_{\tau}}^{\varepsilon} \in \mathfrak{P}(t, x, y)$. Finally, (27) evaluated at Ξ_{τ} yields $J(\Xi_{\tau}; P_{\Xi_{\tau}}^{\varepsilon}) \leq V^{\text{rel}}(\Xi_{\tau}) + \varepsilon$ a.s. $\square$

Proposition 19. *Let $P \in \mathfrak{P}(t, x, y)$ and let τ be an $[t, T]$ -valued stopping time. Let $P^{\tau, \omega}$ be a regular conditional continuation law furnished by Proposition 14. Then $J(t, x, y; P) = E^P[\int_t^{\tau} e^{-\delta(s-t)} \int_U f(s, X_s, Y_s, a) q_s(da) ds + e^{-\delta(\tau-t)} J(\tau, X_{\tau}, Y_{\tau}; P^{\tau, \omega})]$. Consequently,*

$$J(t, x, y; P) \geq E^P \left[\int_t^{\tau} e^{-\delta(s-t)} \int_U f(s, X_s, Y_s, a) q_s(da) ds + e^{-\delta(\tau-t)} V^{\text{rel}}(\tau, X_{\tau}, Y_{\tau}) \right]$$

Proof. Write $C_{t,T} = C_{t,\tau} + e^{-\delta(\tau-t)} C_{\tau,T}$ for the discounted cost functional. By stability under conditioning in Proposition 14, $P^{\tau, \omega} \in \mathfrak{P}(\tau(\omega), X_{\tau}(\omega), Y_{\tau}(\omega))$ for P -a.e. ω . Hence the tower property and conditional Fubini theorem give $E^P[C_{\tau,T} | \mathcal{F}_{\tau}] = J(\tau, X_{\tau}, Y_{\tau}; P^{\tau, \omega})$ a.s. Integrability follows from Proposition 13. Therefore the first identity holds. Since $J(\tau, X_{\tau}, Y_{\tau}; P^{\tau, \omega}) \geq V^{\text{rel}}(\tau, X_{\tau}, Y_{\tau})$ a.s., the inequality follows. $\square$

Proposition 20. *Suppose Assumptions 2.1-2.9 hold and let the strict-control approximation established by the chattering proposition above hold. Define $V^{\text{str}}(t, x, y) := \inf_{u \in \mathcal{U}_t} J(t, x, y; u)$ and $V^{\text{rel}}(t, x, y) := \inf_{P \in \mathfrak{P}(t, x, y)} J(t, x, y; P)$. Then $V^{\text{str}}(t, x, y) = V^{\text{rel}}(t, x, y)$, for all $(t, x, y) \in [0, T] \times \mathbb{R}^2$.*

Proof. For $u \in \mathcal{U}_t$, let $q_s^u(da) := \delta_{u_s}(da)$ and denote by P^u the corresponding canonical law. Then $P^u \in \mathfrak{P}(t, x, y)$ and $J(t, x, y; P^u) = J(t, x, y; u)$, whence $V^{\text{rel}} \leq V^{\text{str}}$. Fix $\varepsilon > 0$ and choose $P^{\varepsilon} \in \mathfrak{P}(t, x, y)$ such that $J(t, x, y; P^{\varepsilon}) \leq V^{\text{rel}}(t, x, y) + \varepsilon$. By the chattering approximation established above, there exists $(u^n)_{n \geq 1} \subset \mathcal{U}_t$ such that the associated strict laws satisfy $P^{u^n} \Rightarrow P^{\varepsilon}$ in the canonical topology and $J(t, x, y; u^n) \rightarrow J(t, x, y; P^{\varepsilon})$. Therefore $V^{\text{str}}(t, x, y) \leq \lim_{n \rightarrow \infty} J(t, x, y; u^n) = J(t, x, y; P^{\varepsilon}) \leq V^{\text{rel}}(t, x, y) + \varepsilon$. Letting $\varepsilon \downarrow 0$ gives $V^{\text{str}} \leq V^{\text{rel}}$, and (26) follows. $\square$

From this point onward, write $V := V^{\text{str}} = V^{\text{rel}}$.

Corollary 21. *Suppose Assumptions 2.1-2.9 hold and assume additionally that $f(t, x, y, u) \geq 0$ and $g(x, y) \geq 0$. Then the common value $V := V^{\text{str}} = V^{\text{rel}}$ is finite and satisfies*

$$0 \leq V(t, x, y) \leq C(1 + |x|^q + e^{C|y|}), \quad (t, x, y) \in [0, T] \times \mathbb{R}^2, \quad (28)$$

for a constant $C > 0$ independent of (t, x, y) . In particular, $V \in G_q$, where $G_q := \{v : [0, T] \times \mathbb{R}^2 \rightarrow \mathbb{R} : |v(t, x, y)| \leq C_v(1 + |x|^q + e^{C_v|y|}) \text{ for some } C_v > 0\}$.

Proof. Since $f, g \geq 0, V \geq 0$. Fix $\bar{u} \in U$ and set $u_s \equiv \bar{u}$. Compactness of U implies $\int_t^T |\bar{u}|^{p^*} ds < \infty$, hence $\bar{u} \in \mathcal{U}_t$. Therefore, $V(t, x, y) \leq J(t, x, y; \bar{u})$. Assumption 2.6 yields $J(t, x, y; \bar{u}) \leq CE[\int_t^T (1 + |X_s^{t,x,y;\bar{u}}|^q + e^{q|Y_s^{t,y}|} + |\bar{u}|^q) ds + 1 + |X_T^{t,x,y;\bar{u}}|^q + e^{q|Y_T^{t,y}|}]$. By Theorem 3 and Lemma 1, $E[\sup_{t \leq s \leq T} |X_s^{t,x,y;\bar{u}}|^q] \leq C(1 + |x|^q + |y|^q + e^{C|y|})$ and $E[\sup_{t \leq s \leq T} e^{q|Y_s^{t,y}|}] \leq Ce^{C|y|}$. Since $|y|^q \leq C(1 + e^{C|y|})$ and $\bar{u}$ ranges in compact U , $J(t, x, y; \bar{u}) \leq C(1 + |x|^q + e^{C|y|})$. Together with $V \geq 0$, this proves (28). $\square$

Remark 5. *Proposition 14 verifies the analytic control correspondence and its stability under conditioning and concatenation. The chattering approximation establishes density of strict controls in the relaxed formulation, while Proposition 16 identifies the two value functions. Together with Theorem 17, Corollary 18, and Proposition 19, these results complete the control correspondence required for the stopping-time DPP and the subsequent viscosity analysis.*

3 Catastrophe Insurance as a Stochastic Control Problem

We now specialize the controlled jump system of Section 2 to a catastrophe insurer exposed to post-disaster replacement-cost risk. The state variable X represents insurer surplus, $I = e^Y$ is the replacement-cost index, and u denotes physical hedging through pre-arranged reconstruction capacity. The primitive probability space, Lévy measure, admissible control class, and controlled state process are exactly those of Assumptions 2.-2.7. Thus no new stochastic dynamics are introduced in this section.

3.1 Replacement-Cost Dynamics and the Control Problem

The logarithmic replacement-cost factor satisfies $dY_s = \kappa(\vartheta - Y_s)ds + \sigma_I dW_s^I$, with $Y_t = y$, and $I_s = e^{Y_s}$. A positive displacement of Y_s from ϑ represents a temporary state of elevated reconstruction costs, while mean reversion describes normalization toward the long-run cost level. The process Y is exogenous to the catastrophe random measure N : catastrophe arrivals do not generate jumps or impulses in Y . Accordingly, the model captures catastrophe losses occurring under stochastic replacement-cost conditions rather than a structural catastrophe-induced demand-surge process. Empirical evidence on post-disaster reconstruction-cost inflation and demand surge provides the economic motivation for treating elevated replacement costs as an important catastrophe-insurance state variable [42, 95, 49, 20]. Catastrophe losses are generated by the Poisson random measure N introduced in Assumption 2.1. We retain its compensator $\mathbf{v}(dz)ds$ and specialize the controlled jump amplitude to

$$\ell(y, z, u) = e^y a(z) h(u), \quad (29)$$

where $a : E \rightarrow \mathbb{R}_+$ maps a catastrophe mark into baseline physical damage and $h : U \rightarrow (0, 1]$ is the residual-loss fraction after physical hedging. Assume $h(0) = 1$, $h'(u) < 0$, and $h''(u) \geq 0$ whenever the derivatives exist. Hence physical hedging reduces catastrophe severity, with weakly diminishing marginal effectiveness. The exponential specification $h(u) = e^{-\gamma u}$, for all $\gamma > 0$, is a canonical example.

The factorization in (29) separates three sources of loss. The mark z determines physical catastrophe severity, e^y converts physical damage into replacement expenditure at prevailing construction prices, and $h(u)$ records the fraction of that expenditure remaining after pre-arranged capacity has been used [60]. Thus elevated replacement costs and physical hedging act in opposite directions on the monetary scale of the same catastrophe jump exposure [58]. The insurer surplus is the process already defined in (4)

$$dX_s = b(X_{s-}, Y_{s-}, u_s)ds + \sigma_X(X_{s-}, Y_{s-}, u_s)dW_s^X - \int_E e^{Y_{s-}} a(z) h(u_s) \tilde{N}(ds, dz), \quad X_t = x. \quad (30)$$

For the economic specification developed below, consider $b(x, y, u) = rx + \pi(y) - c(u)$, where $r \geq 0$ is the return on liquid surplus, $\pi(y)$ is net premium income before physical-hedging expenditure, and c is the re-

source cost of securing reconstruction capacity. We impose $c(0) = 0$, $c'(u) > 0$, and $c''(u) > 0$. A quadratic benchmark is $c(u) = \frac{\chi}{2}u^2$, for all $\chi > 0$. The diffusion coefficient σ_X represents noncatastrophe balance-sheet fluctuations and continues to satisfy Assumptions 2.4 and 2.7.

The compensated representation in (30) is convenient for the probabilistic analysis. Its economic content can be seen from the uncompensated form. Whenever the first jump moment is finite,

$$dX_s = \left[b(X_{s-}, Y_{s-}, u_s) + \int_E e^{Y_{s-}} a(z) h(u_s) \nu(dz) \right] ds + \sigma_X(X_{s-}, Y_{s-}, u_s) dW_s^X - \int_E e^{Y_{s-}} a(z) h(u_s) N(ds, dz).$$

Accordingly, the actual downward jump at a catastrophe mark z is $\Delta X_s = -e^{Y_{s-}} a(z) h(u_s)$. Physical hedging therefore modifies loss severity rather than the arrival law of the underlying catastrophe process. For $p \geq 1$ such that the corresponding integral is finite, define the conditional p th jump-loss moment as $\mathfrak{L}_p(y, u) := \int_E \ell(y, z, u)^p \nu(dz)$. Under (29),

$$\mathfrak{L}_p(y, u) = e^{py} h(u)^p \int_E a(z)^p \nu(dz). \quad (31)$$

Hence, $\partial_y \log \mathfrak{L}_p(y, u) = p$, and $\partial_u \log \mathfrak{L}_p(y, u) = p \frac{h'(u)}{h(u)} < 0$. Equation (31) shows that an elevated replacement-cost state raises every finite moment of monetary catastrophe-loss exposure, whereas physical hedging lowers it. In particular, the control changes not merely the mean loss but the entire scale of the catastrophe-loss distribution.

Remark 6. *The specification remains compatible with heavy-tailed catastrophe risk. If ν has sufficiently heavy upper tails, only moments below a critical order are finite. The probabilistic results of Section 2 therefore require moments only up to the fixed exponent p_* appearing in Assumptions 2.1-2.4; no Gaussian approximation to catastrophe severity is imposed.*

3.2 Value Function and Optimal Physical Hedging

We formulate physical hedging as a social-cost minimization problem. For $u \in \mathcal{U}_t$, let

$$J(t, x, y; u) = \mathbb{E}_{t, x, y} \left[\int_t^T e^{-\delta(s-t)} f(s, X_s, Y_s, u_s) ds + e^{-\delta(T-t)} g(X_T, Y_T) \right], \quad (32)$$

which is the objective functional introduced in (17). The economic specialization is $f(t, x, y, u) = C_H(u) + C_D(y) + C_S(x)$, where C_H is the resource cost of physical hedging, C_D measures the social burden of elevated reconstruction costs, and C_S penalizes deterioration in insurer capitalization. Assume that C_H is strictly convex and coercive [65]. A convenient benchmark is $C_H(u) = \frac{\chi}{2}u^2$. For the demand-surge component, one may take $C_D(y) = \zeta(e^y - e^\vartheta)_+^m$, for all $\zeta > 0$, and $m \geq 1$, while a capitalization penalty can be specified as $C_S(x) = \omega(x_{\text{crit}} - x)_+^q$, with $\omega > 0$. The terminal function g measures the residual social cost of ending the planning horizon with a weak insurance balance sheet or persistent replacement-cost stress. These functions are chosen to satisfy Assumption 2.6. The value function is

$$V(t, x, y) = \inf_{u \in \mathcal{U}_t} J(t, x, y; u). \quad (33)$$

By Proposition 18, (33) coincides with the relaxed canonical value used in the measurable-selection construction. Corollary 21 places V in the growth class $\mathcal{G}_q$. Thus the economic control problem is defined on exactly the same admissible state-control [89] system for which Section 2 established well-posedness, stability, restart, and measurable continuation. The role of physical hedging becomes transparent at the level of the controlled generator. For $\varphi \in C^{1,2,2}$, define

$$\mathcal{L}^u \varphi(t, x, y) := b(x, y, u) \varphi_x + \kappa(\vartheta - y) \varphi_y + \frac{1}{2} \sigma_X^2(x, y, u) \varphi_{xx} + \frac{1}{2} \sigma_I^2 \varphi_{yy} + I^u \varphi(t, x, y),$$

where the nonlocal catastrophe operator is

$$I^u \varphi(t, x, y) := \int_E \left[\varphi(t, x - e^y a(z) h(u), y) - \varphi(t, x, y) + e^y a(z) h(u) \varphi_x(t, x, y) \right] \nu(dz). \quad (34)$$

The control therefore enters the generator inside the nonlocal translation of the surplus state [88]. This distinguishes physical hedging from a control that acts only through drift expenditure. For a smooth continuation value, the local tradeoff associated with u is represented formally by

$$u \mapsto f(t, x, y, u) + b(x, y, u) V_x(t, x, y) + \frac{1}{2} \sigma_X^2(x, y, u) V_{xx}(t, x, y) + I^u V(t, x, y).$$

Accordingly, an interior optimal feedback, when it exists, must balance the marginal resource cost of reconstruction capacity against the marginal reduction in the nonlocal catastrophe exposure. To make this relation explicit, suppose temporarily that σ_X is independent of u , that $c(u) = C_H(u) = \chi u^2/2$, and that h is differentiable. Formal differentiation gives the first-order condition

$$0 = \chi u - \chi u V_x(t, x, y) - e^y h'(u) \int_E a(z) \left[V_x(t, x - e^y a(z) h(u), y) - V_x(t, x, y) \right] \nu(dz). \quad (35)$$

Remark 7. Equation (35) is only heuristic at this stage. The value function need not be differentiable, and under heavy-tailed jump risk the nonlocal term may prevent the existence of a classical solution altogether. We therefore do not use (35) as a verification argument. Instead, the optimal physical hedge will be defined through the minimizer of the Hamiltonian in the viscosity solution. This is the point at which the probabilistic foundations of Section 2 become essential. The stopping-time flow and measurable continuation results permit a rigorous dynamic programming principle, while the growth estimate of Corollary 21 determines the admissible uniqueness class. Section 4 derives the resulting nonlocal HJB equation and establishes the viscosity characterization of V .

4 Dynamic Programming and Viscosity Solutions

We now identify the value function of Section 3 with the solution of a nonlocal HJB equation. The argument has two distinct components. First, the stopping-time restart and measurable-selection results of Section 2 yield a dynamic programming principle without assuming the existence of an optimal strict control. Second, the DPP [62] is localized to obtain the viscosity inequalities. The comparison argument is subsequently carried out in the growth class $\mathcal{G}_q$ identified in Corollary 21. Throughout this section, write $z_0 = (x, y) \in \mathbb{R}^2$

when no confusion can arise, and retain the notation $\ell(y, z, u) = e^y a(z) h(u)$ from (29).

4.1 Dynamic Programming and the Nonlocal HJB Equation

For $\varphi \in C^{1,2,2}([0, T] \times \mathbb{R}^2)$ and $u \in U$, define

$$\begin{aligned} \mathcal{A}^u \varphi(t, x, y) &:= b(x, y, u) \varphi_x(t, x, y) + \kappa(\vartheta - y) \varphi_y(t, x, y) \\ &\quad + \frac{1}{2} \sigma_X^2(x, y, u) \varphi_{xx}(t, x, y) + \frac{1}{2} \sigma_I^2 \varphi_{yy}(t, x, y) + I^u \varphi(t, x, y), \end{aligned}$$

where I^u is the catastrophe-loss operator

$$I^u \varphi(t, x, y) := \int_E \left[\varphi(t, x - \ell(y, z, u), y) - \varphi(t, x, y) + \ell(y, z, u) \varphi_x(t, x, y) \right] \mathbf{v}(dz).$$

The compensation term is inherited from the compensated Poisson representation in (4). Economically, I^u records the marginal continuation cost generated by the entire distribution of catastrophe severities rather than by an average-loss approximation. Define the Hamiltonian

$$\begin{aligned} \mathcal{H}(t, x, y, r, p, X; \psi) &:= \inf_{u \in U} \left\{ f(t, x, y, u) + b(x, y, u) p_1 + \kappa(\vartheta - y) p_2 \right. \\ &\quad \left. + \frac{1}{2} \sigma_X^2(x, y, u) X_{11} + \frac{1}{2} \sigma_I^2 X_{22} - \delta r \right. \\ &\quad \left. + \int_E \left[\psi(x - \ell(y, z, u), y) - \psi(x, y) + \ell(y, z, u) p_1 \right] \mathbf{v}(dz) \right\}, \end{aligned} \quad (36)$$

whenever the integral is well defined. Here $p = (p_1, p_2) \in \mathbb{R}^2$ and $X = (X_{ij}) \in \mathbb{S}^2$. The HJB equation associated with (33) is therefore

$$-\partial_t v(t, x, y) - \mathcal{H}(t, x, y, v, Dv, D^2v; v(t, \cdot, \cdot)) = 0, \quad (t, x, y) \in [0, T] \times \mathbb{R}^2, \quad (37)$$

with terminal condition

$$v(T, x, y) = g(x, y). \quad (38)$$

Since $\mathbf{v}(E)$ need not be finite, it is useful to split the nonlocal operator. For $\varepsilon \in (0, 1)$ set $E_\varepsilon := \{z \in E : \rho(z) \leq \varepsilon\}$, $E^\varepsilon := E \setminus E_\varepsilon$, and, for a smooth test function ϕ and a function w of appropriate growth, define

$$\begin{aligned} I_\varepsilon^u[\phi](t, x, y) &:= \int_{E_\varepsilon} \left[\phi(t, x - \ell(y, z, u), y) - \phi(t, x, y) + \ell(y, z, u) \phi_x(t, x, y) \right] \mathbf{v}(dz), \\ I^{u, \varepsilon}[w, \phi](t, x, y) &:= \int_{E^\varepsilon} \left[w(t, x - \ell(y, z, u), y) - w(t, x, y) + \ell(y, z, u) \phi_x(t, x, y) \right] \mathbf{v}(dz). \end{aligned}$$

The small jumps are evaluated on the test function, whereas the large-jump contribution retains the candidate solution. This is the form of the viscosity definition that is stable under singular Lévy measures [see Barles and Imbert [8]].

Lemma 22. *Let Assumptions 2.1-2.9 hold and let τ be an $[t, T]$ -valued stopping time. For every admissible control $u \in \mathcal{U}_t$,*

$$J(t, x, y; u) = \mathbb{E}_{t, x, y} \left[\int_t^\tau e^{-\delta(s-t)} f(s, X_s, Y_s, u_s) ds + e^{-\delta(\tau-t)} J(\tau, X_\tau, Y_\tau; u^{\tau, \omega}) \right],$$

where $u^{\tau, \omega}$ denotes the restarted continuation control. The same identity holds in the canonical relaxed formulation with the regular conditional continuation law replacing $u^{\tau, \omega}$.

Proof. Split the running integral in (32) at τ . For $s \geq \tau$, $e^{-\delta(s-t)} = e^{-\delta(\tau-t)} e^{-\delta(s-\tau)}$. The conditional restart result of Proposition 9 identifies the post- τ state, conditionally on $\mathcal{F}_\tau$, with the state process initialized from (τ, X_τ, Y_τ) and driven by the shifted continuation control. Consequently, conditional expectation of the post- τ running and terminal costs equals $e^{-\delta(\tau-t)} J(\tau, X_\tau, Y_\tau; u^{\tau, \omega})$. The integrability required to apply conditional Fubini and the tower property follows from Proposition 13. Taking expectations proves the identity. In the canonical formulation, the same argument follows by disintegration of the controlled law and the conditioning stability established in Proposition 14. $\square$

Theorem 23. *Suppose Assumptions 2.1-2.9 hold. Then, for every $(t, x, y) \in [0, T] \times \mathbb{R}^2$ and every $[t, T]$ -valued stopping time τ ,*

$$V(t, x, y) = \inf_{u \in \mathcal{U}_t} \mathbb{E}_{t, x, y} \left[\int_t^\tau e^{-\delta(s-t)} f(s, X_s, Y_s, u_s) ds + e^{-\delta(\tau-t)} V(\tau, X_\tau, Y_\tau) \right]. \quad (39)$$

Equivalently, the infimum may be taken over $P \in \mathfrak{P}(t, x, y)$.

Corollary 24. *For every $t \leq s \leq T$,*

$$V(t, x, y) = \inf_{u \in \mathcal{U}_t} \mathbb{E}_{t, x, y} \left[\int_t^s e^{-\delta(r-t)} f(r, X_r, Y_r, u_r) dr + e^{-\delta(s-t)} V(s, X_s, Y_s) \right].$$

Proof. Apply Theorem 23 with the deterministic stopping time $\tau = s$. $\square$

Proposition 25. *Suppose Assumptions 2.1-2.9 hold. Assume in addition that f and g are locally uniformly continuous in (t, x, y) , uniformly for u in bounded subsets of U . Then $V \in C([0, T] \times \mathbb{R}^2) \cap \mathcal{G}_q$. Moreover, $\lim_{t \uparrow T} V(t, x, y) = g(x, y)$ locally uniformly in (x, y) .*

Definition 1 (Viscosity solution). *Let $w \in \mathcal{G}_q$ be upper semicontinuous. We call w a viscosity subsolution of (37) if $w(T, \cdot, \cdot) \leq g$ and, whenever $\phi \in C^{1,2,2}$ and $w - \phi$ has a local maximum at $(t_0, x_0, y_0) \in [0, T) \times \mathbb{R}^2$, then, for every sufficiently small $\varepsilon > 0$,*

$$0 \geq \inf_{u \in U} \left\{ \phi_t + b\phi_x + \kappa(\vartheta - y_0)\phi_y + \frac{1}{2}\sigma_X^2\phi_{xx} + \frac{1}{2}\sigma_I^2\phi_{yy} - \delta w(t_0, x_0, y_0) + f \right. \\ \left. + I_\varepsilon^u[\phi](t_0, x_0, y_0) + I^{u, \varepsilon}[w, \phi](t_0, x_0, y_0) \right\},$$

where all local coefficients are evaluated at (t_0, x_0, y_0, u) . A lower semicontinuous $w \in \mathcal{G}_q$ is a viscosity supersolution if $w(T, \cdot, \cdot) \geq g$ and the reverse inequality holds at every local minimum of $w - \phi$. A continuous function is a viscosity solution if it is both a subsolution and a supersolution.

Lemma 26 (Localized Dynkin formula). *Let $\phi \in C^{1,2,2}$ and let*

$$\tau_R := \inf\{s \geq t : |X_s - x| + |Y_s - y| \geq R\} \wedge (t + h).$$

For any constant control $u_s \equiv a \in U$,

$$E \left[e^{-\delta(\tau_R - t)} \phi(\tau_R, X_{\tau_R}, Y_{\tau_R}) - \phi(t, x, y) \right] = E \int_t^{\tau_R} e^{-\delta(s-t)} \left[\phi_t + \mathcal{A}^a \phi - \delta \phi \right](s, X_{s-}, Y_{s-}) ds.$$

Proof. Apply the Itô formula for jump semimartingales to $s \mapsto e^{-\delta(s-t)} \phi(s, X_s, Y_s)$ up to τ_R . The continuous quadratic-variation terms generate $\frac{1}{2} \sigma_X^2 \phi_{xx}$ and $\frac{1}{2} \sigma_Y^2 \phi_{yy}$. The compensated Poisson term generates $I^a \phi$. The Brownian and compensated-jump stochastic integrals are true martingales after localization because ϕ and its derivatives are bounded on the stopped region and Proposition 2 gives square integrability of the jump integral. Taking expectations eliminates the martingale terms. $\square$

Theorem 27. *Under the hypotheses of Proposition 25, the value function V is a viscosity solution of (37)-(38).*

Corollary 28. *At every time point at which V admits a classical first and second derivative, any minimizing selector of the Hamiltonian satisfies*

$$u^*(t, x, y) \in \arg \min_{u \in U} \left\{ f(t, x, y, u) + b(x, y, u) V_x + \frac{1}{2} \sigma_X^2(x, y, u) V_{xx} + I^u V(t, x, y) \right\}.$$

Thus physical hedging is selected by the marginal effect of u on both the local balance-sheet dynamics and the full catastrophe-loss distribution.

Proof. At a differentiability point, use V itself as the local test function in the viscosity equation. Terms independent of u may be removed from the minimization. The remaining expression is exactly the displayed Hamiltonian. $\square$

4.2 Comparison, Existence, and Uniqueness

The unbounded state space and the possibly infinite Lévy measure require a comparison argument compatible with the exponential-polynomial growth of V . We impose the following additional structure only for this subsection.

Assumption 4.1. The following conditions hold.

- (i) For every $R > 0$, the maps b , σ_X , and f are uniformly continuous in (t, x, y) , uniformly for $u \in U \cap [0, R]$.
- (ii) There is a measurable $\rho : E \rightarrow [0, \infty)$ as in Assumption 2.4 such that, uniformly in u ,

$$|\ell(y, z, u) - \ell(y', z, u)| \leq C|y - y'| \rho(z), \quad |\ell(y, z, u)| \leq C(1 + e^{|y|} + |u|) \rho(z).$$

(iii) The coercivity of f dominates the positive growth of the control-dependent coefficients: for every compact $K \subset [0, T] \times \mathbb{R}^2$,

$$f(t, x, y, u) - C_K(|b(x, y, u)| + \sigma_X^2(x, y, u) + \|\ell(y, \cdot, u)\|_{L^2(\mathbf{v})}^2) \longrightarrow +\infty$$

as $|u| \rightarrow \infty$, uniformly on K .

(iv) There exists $\bar{q} \in (q, p_*)$ such that $\int_E (\rho(z)^2 + \rho(z)^{\bar{q}}) \mathbf{v}(dz) < \infty$.

(v) The terminal cost g is continuous and belongs to $\mathcal{G}_q$.

Assumption 4.1(iii) is the analytic counterpart of the economic scarcity cost of physical hedging. It prevents a minimizing sequence from escaping to infinite reconstruction capacity merely to suppress catastrophe severity.

Lemma 29. *Under Assumption 4.1, for every compact $K \subset [0, T] \times \mathbb{R}^2$ and every bounded set of (r, p, X) , there exists $R_K < \infty$ such that the infimum in (36), restricted to arguments in these sets, is unchanged if U is replaced by $U \cap [0, R_K]$.*

Proof. On the stated compact sets, all terms in the Hamiltonian other than f are bounded in absolute value by

$$C_K \left(1 + |b(x, y, u)| + \sigma_X^2(x, y, u) + \|\ell(y, \cdot, u)\|_{L^2(\mathbf{v})}^2 + \|\ell(y, \cdot, u)\|_{L^{\bar{q}}(\mathbf{v})}^{\bar{q}} \right),$$

after Taylor expansion of the small-jump contribution and use of the growth class for the large-jump part. Assumption 4.1(iii), together with the moment condition in (iv), implies that the full minimized expression tends to $+\infty$ as $|u| \rightarrow \infty$, uniformly over the compact set of remaining variables. Hence no minimizing sequence can escape to infinity. Choosing R_K beyond the resulting coercive level proves the claim. $\square$

Lemma 30. *Let w be an upper semicontinuous subsolution and $\underline{w}$ a lower semicontinuous supersolution in $\mathcal{G}_q$. For $\varepsilon, \eta > 0$, consider a penalization of the form*

$$\Phi_{\varepsilon, \eta}(t, s, x, x', y, y') := w(t, x, y) - \underline{w}(s, x', y') - \frac{|x - x'|^2 + |y - y'|^2 + |t - s|^2}{2\varepsilon} - \eta(\Gamma(x, y) + \Gamma(x', y')),$$

where $\Gamma(x, y) = 1 + |x|^{\bar{q}} + e^{\lambda\sqrt{1+y^2}}$ with $\bar{q} \in (q, p_*)$ and $\lambda > 0$ sufficiently large. If $(t_\varepsilon, s_\varepsilon, x_\varepsilon, x'_\varepsilon, y_\varepsilon, y'_\varepsilon)$ is a maximizer, then, along a subsequence,

$$\frac{|x_\varepsilon - x'_\varepsilon|^2 + |y_\varepsilon - y'_\varepsilon|^2 + |t_\varepsilon - s_\varepsilon|^2}{\varepsilon} \longrightarrow 0$$

as $\varepsilon \downarrow 0$, and the difference of the corresponding nonlocal terms is bounded above by $o_\varepsilon(1) + C\eta$.

Proposition 31. *Let $\underline{w} \in \mathcal{G}_q$ be a viscosity supersolution of (37). There exist constants $\lambda, \Lambda > 0$ such that, for $\Gamma(x, y) = 1 + |x|^{\bar{q}} + e^{\lambda\sqrt{1+y^2}}$, the function $\underline{w}^\eta(t, x, y) = \underline{w}(t, x, y) + \eta e^{\Lambda(T-t)} \Gamma(x, y)$ is, for every $\eta > 0$, a strict viscosity supersolution in the sense that its HJB residual is bounded below by $c\eta\Gamma(x, y)$ for some $c > 0$.*

Theorem 32. Suppose Assumptions 2.1-2.9 and 4.1 hold. Let $\bar{v} \in USC([0, T] \times \mathbb{R}^2) \cap \mathcal{G}_q$ be a viscosity subsolution of (37), and let $\underline{v} \in LSC([0, T] \times \mathbb{R}^2) \cap \mathcal{G}_q$ be a viscosity supersolution. If $\bar{v}(T, x, y) \leq \underline{v}(T, x, y)$ for all $(x, y) \in \mathbb{R}^2$, then

$$\bar{v}(t, x, y) \leq \underline{v}(t, x, y) \quad \text{on } [0, T] \times \mathbb{R}^2. \quad (40)$$

Corollary 33. Under the hypotheses of Theorem 32, there exists at most one continuous viscosity solution $v \in \mathcal{G}_q$ of (37)-(38).

Proof. If v_1 and v_2 are two such solutions, apply Theorem 32 first with $(\bar{v}, \underline{v}) = (v_1, v_2)$ and then with (v_2, v_1) . Thus $v_1 \leq v_2$ and $v_2 \leq v_1$. $\square$

Theorem 34. Suppose Assumptions 2.1-2.9 and 4.1 hold. Then the catastrophe insurance value function defined by (33) is the unique viscosity solution in $C([0, T] \times \mathbb{R}^2) \cap \mathcal{G}_q$ of (37)-(38).

Proof. Existence is probabilistic. Proposition 25 gives $V \in C([0, T] \times \mathbb{R}^2) \cap \mathcal{G}_q$, and Theorem 27 shows that V satisfies the nonlocal HJB equation in the viscosity sense together with the terminal condition. Uniqueness follows from Corollary 33. Therefore every continuous viscosity solution in $\mathcal{G}_q$ coincides with the stochastic control value. $\square$

Proposition 35. Assume, in addition, that for every (t, x, y) and every smooth test jet (p, X) the map

$$u \mapsto f(t, x, y, u) + b(x, y, u)p_1 + \frac{1}{2}\sigma_X^2(x, y, u)X_{11} + I^u\phi(t, x, y)$$

is lower semicontinuous. Then the minimizing set is nonempty and compact, and there exists a Borel measurable selector $\hat{u}(t, x, y, p, X)$ from the corresponding argmin correspondence.

Proof. By Lemma 29, minimization may be restricted locally to a compact subset of U . Lower semicontinuity then gives attainment by the Weierstrass theorem. The objective is Borel measurable in the state and jet variables and lower semicontinuous in u ; hence its argmin correspondence has nonempty compact values and measurable graph. The Kuratowski-Ryll-Nardzewski measurable-selection theorem provides a Borel selector. $\square$

Corollary 36 (Classical verification as a special case). Suppose the unique viscosity solution V belongs to $C^{1,2,2}$, the selector

$$u_s^* = \hat{u}(s, X_{s-}, Y_{s-}, DV(s, X_{s-}, Y_{s-}), D^2V(s, X_{s-}, Y_{s-}))$$

is admissible, and the stochastic integrals arising from Itô's formula are true martingales. Then u^* is optimal and

$$V(t, x, y) = J(t, x, y; u^*).$$

Proof. For every $u \in U$, the HJB equation implies $V_t + \mathcal{A}^u V - \delta V + f(t, x, y, u) \geq 0$, while equality holds for the minimizing selector u^* . Apply Itô's formula to $e^{-\delta(s-t)}V(s, X_s, Y_s)$ between t and T . After taking expectations, the martingale terms vanish and the terminal condition gives $V(t, x, y) \leq J(t, x, y; u)$ for every admissible u . Under u^* all inequalities become equalities, yielding $V(t, x, y) = J(t, x, y; u^*)$. $\square$

Corollary 37. *Suppose the hypotheses of Corollary 36 hold and use the benchmark loss amplitude $\ell(y, z, u) = e^y a(z) h(u)$. If σ_X is independent of u and the minimizer is interior, then the optimal physical hedge satisfies*

$$0 = f_u(t, x, y, u^*) + b_u(x, y, u^*) V_x(t, x, y) - e^y h'(u^*) \int_E a(z) \left[V_x(t, x - e^y a(z) h(u^*), y) - V_x(t, x, y) \right] \nu(dz).$$

Thus the marginal cost of securing reconstruction capacity is equated to the marginal change in continuation value generated by shifting the entire catastrophe-loss distribution.

Proof. Differentiate the minimizing Hamiltonian with respect to u . The local terms give $f_u + b_u V_x$. Differentiating the nonlocal term under the integral sign gives

$$-e^y h'(u) \int_E a(z) [V_x(t, x - e^y a(z) h(u), y) - V_x(t, x, y)] \nu(dz).$$

The interchange of differentiation and integration follows from the assumed classical regularity, the moment condition on the Lévy measure, and dominated convergence. At an interior minimizer the derivative vanishes. $\square$

Remark 8. *The preceding results separate the two roles played by the catastrophe model. Probabilistically, the marked jump measure creates the nonlocal generator and prevents the control problem from reducing to a diffusion HJB equation. Economically, the multiplicative factor $e^y a(z) h(u)$ makes the value of physical hedging state dependent: the same unit of secured reconstruction capacity has a different marginal value when replacement costs are elevated. The viscosity formulation preserves this mechanism without imposing differentiability of the insurer's continuation value.*

5 Mean Field Game of Catastrophe Insurers

The representative-insurer dynamics remain exactly those of Sections 2-3. Mean field interaction is introduced through the objective functional rather than through an endogenous reconstruction-capacity market. Accordingly, the couplings F and G should be interpreted as reduced-form strategic cost externalities, the private cost of a given capitalization and replacement-cost state depends on the cross-sectional vulnerability of the insurance sector. The population distribution does not determine the replacement-cost process Y , the catastrophe compensator $\nu(dz) dt$, the severity law $a(z)$, or an aggregate supply of reconstruction capacity. Thus the mean field game studies [68] how industry-wide vulnerability feeds back into individual hedging incentives, but it does not model market clearing in reconstruction services or endogenous price formation. This separation preserves the controlled state process analyzed in Sections 2-4 while isolating a strategic externality that can be studied within the nonlocal control.

5.1 Mean Field Equilibrium

For a deterministic flow $m \in \mathcal{M}_T$, define

$$J^m(t, x, y; u) := \mathbb{E}_{t, x, y} \left[\int_t^T e^{-\delta(s-t)} \left(f(s, X_s, Y_s, u_s) + F(s, (X_s, Y_s), m_s) \right) ds + e^{-\delta(T-t)} \left(g(X_T, Y_T) + G((X_T, Y_T), m_T) \right) \right], \quad (41)$$

and

$$V^m(t, x, y) := \inf_{u \in \mathcal{U}_t} J^m(t, x, y; u). \quad (42)$$

The controlled state in (41) is exactly the solution of (3) and (4).

Assumption 5.1 (Mean field couplings). There exists $C_F > 0$ such that:

(i) F and G are continuous;

(ii) for all (t, z, m) ,

$$|F(t, z, m)| + |G(z, m)| \leq C_F \left(1 + |x|^q + e^{q|y|} + M_q(m)^q \right),$$

where $M_q(m) := (\int_{\mathbb{R}^2} |z'|^q m(dz'))^{1/q}$;

(iii) for every $R > 0$, there exists $L_R > 0$ such that

$$|F(t, z, m) - F(s, z', n)| + |G(z, m) - G(z', n)| \leq L_R \left(|t - s|^{1/2} + |z - z'| + W_q(m, n) \right)$$

whenever $|z|, |z'|, M_q(m), M_q(n) \leq R$;

(iv) F and G are bounded from below by a function of at most the exponential-polynomial growth appearing in Assumption 2.6, uniformly on bounded-moment subsets of $\mathcal{P}_q(\mathbb{R}^2)$.

Remark 9. The maps $F(t, z, m)$ and $G(z, m)$ are not derived from a reconstruction-market clearing condition. They represent reduced-form costs associated with cross-sectional concentration of vulnerability. For example, F may penalize an insurer occupying a high replacement-cost or weak-capitalization state when a large fraction of the industry simultaneously occupies comparable states [64]. Such a coupling can represent congestion, coordination, regulatory, or systemic-capacity costs in reduced form, but it does not determine the price or quantity of reconstruction services.

A structural reconstruction-capacity model would require an additional aggregate quantity, say $D_t(m, \alpha)$, measuring rebuilding or capacity demand, together with a supply or market-clearing relation that determines an endogenous price or scarcity factor. In such a model, one would expect a relation of the schematic form $P_t^{\text{rec}} = \mathcal{P}(D_t, S_t)$, with $\partial_D \mathcal{P} > 0$, and the resulting price would enter either the replacement-cost dynamics or the controlled loss amplitude. Because this would make the state dynamics themselves population-dependent, it would produce a different mean field game from the one analyzed here.

Lemma 38. Fix $m \in \mathcal{M}_T$ with $\sup_{t \leq T} M_q(m_t) < \infty$. Under Assumptions 2.1-2.9, 4.1, and 5.1, the frozen-flow problem expressed in Equation (42) satisfies the hypotheses of Theorem 23, Proposition 25, and Theorem 34 after replacing f and g by $f^m(t, x, y, u) := f(t, x, y, u) + F(t, (x, y), m_t)$ and $g^m(x, y) := g(x, y) + G((x, y), m_T)$. The local constants may be chosen uniformly over any family of flows with uniformly bounded q th moments.

Proof. The state equation does not depend on m , so the estimates from Section 2 remain unchanged. In particular, Theorem 3, Proposition 5, Lemma 6, and Proposition 13 continue to hold.

If $\sup_t M_q(m_t) \leq R$, Assumption 5.1(ii) yields

$$|f^m(t, x, y, u)| + |g^m(x, y)| \leq C_R(1 + |x|^q + e^{q|y|} + |u|^q).$$

Since the coupling does not depend on u , the coercivity in the physical-hedging variable is exactly that of f . Assumption 5.1(iii) gives the local continuity required by Proposition 25 and Assumption 4.1(i). The jump operator and comparison structure are unchanged. Hence, the results cited in the statement apply to the frozen problem. $\square$

Proposition 39 (Frozen-flow viscosity equation). *Under the hypotheses of Lemma 38, V^m is the unique viscosity solution in $C([0, T] \times \mathbb{R}^2) \cap \mathcal{G}_q$ of*

$$-\partial_t v^m - \mathcal{H}(t, x, y, v^m, Dv^m, D^2 v^m; v^m(t, \cdot, \cdot)) - F(t, (x, y), m_t) = 0, \quad (43)$$

with terminal condition

$$v^m(T, x, y) = g(x, y) + G((x, y), m_T). \quad (44)$$

Proof. Lemma 38 allows the frozen problem to be treated by Section 4. The stopping-time DPP follows from Theorem 23, the viscosity solution from Theorem 27, and uniqueness from Theorem 32 and Corollary 33. Finally, Theorem 34 identifies this viscosity solution with V^m . $\square$

Let $\alpha : [0, T] \times \mathbb{R}^2 \rightarrow U$ be an admissible Borel feedback. For $\phi \in C_c^2(\mathbb{R}^2)$ define

$$\begin{aligned} \mathcal{L}^\alpha \phi(x, y) &:= b(x, y, \alpha_t(x, y))\phi_x + \kappa(\vartheta - y)\phi_y + \frac{1}{2}\sigma_X^2(x, y, \alpha_t(x, y))\phi_{xx} \\ &\quad + \frac{1}{2}\sigma_I^2\phi_{yy} + \int_E \left[\phi(x - \ell(y, z, \alpha_t(x, y)), y) - \phi(x, y) \right. \\ &\quad \left. + \ell(y, z, \alpha_t(x, y))\phi_x(x, y) \right] \nu(dz). \end{aligned}$$

Definition 2. Let $\mathfrak{A}(m)$ denote the full set of admissible relaxed canonical laws for the frozen population flow $m \in \mathcal{M}_T$. A pair (m^*, P^*) is called a relaxed mean field equilibrium if $m^* \in \mathcal{M}_T$, $P^* \in \mathfrak{A}(m^*)$, P^* is optimal for the frozen-flow problem associated with m^* , and

$$m_t^* = P^* \circ Z_t^{-1}, \quad Z_t := (X_t, Y_t), \quad 0 \leq t \leq T. \quad (45)$$

A relaxed equilibrium is called Markovian if, under P^* , the relaxed control coordinate admits the representation $q_t(da) = \hat{q}^*(t, Z_{t-}, da) dt \otimes P^*$ -a.e. for a Borel kernel $\hat{q}^* : [0, T] \times \mathbb{R}^2 \rightarrow \mathcal{P}(U)$. It is called strict Markovian if $\hat{q}^*(t, z, da) = \delta_{\alpha^*(t, z)}(da)$ for a Borel feedback $\alpha^* : [0, T] \times \mathbb{R}^2 \rightarrow U$.

For $P \in \mathfrak{A}(m)$ define the state-control occupation measure η^P on $[0, T] \times \mathbb{R}^2 \times U$ by $\eta^P(dt, dz, da) := dt E^P[\mathbf{1}_{\{Z_t \in dz\}} q_t(da)]$. If $m_t^P := P \circ Z_t^{-1}$, disintegration on the standard Borel space $[0, T] \times \mathbb{R}^2 \times U$ yields a Borel stochastic kernel $\hat{q}^P(t, z, da)$ satisfying $\eta^P(dt, dz, da) = dt m_t^P(dz) \hat{q}^P(t, z, da)$. The kernel $\hat{q}^P$ is the state-conditioned barycentric relaxation of the possibly history-dependent control law P .

Proposition 40. *Let (m^*, P^*) be a relaxed equilibrium and let $\hat{q}^* := \hat{q}^{P^*}$ be the disintegration kernel defined above. Then, for every $\phi \in C_c^2(\mathbb{R}^2)$,*

$$\int_{\mathbb{R}^2} \phi(z) m_t^*(dz) - \int_{\mathbb{R}^2} \phi(z) m_0(dz) = \int_0^t \int_{\mathbb{R}^2} \int_U \mathcal{A}^a \phi(z) \hat{q}^*(s, z, da) m_s^*(dz) ds. \quad (46)$$

Equivalently, $\partial_t m_t^* = (\mathcal{A}^{\hat{q}_t^*})^* m_t^*$ in $\mathcal{D}'((0, T) \times \mathbb{R}^2)$, where $\mathcal{A}^{\hat{q}_t^*} \phi(z) := \int_U \mathcal{A}^a \phi(z) \hat{q}^*(t, z, da)$.

Proof. For $\phi \in C_c^2(\mathbb{R}^2)$, admissibility of P^* gives $M_t^\phi := \phi(Z_t) - \phi(Z_0) - \int_0^t \int_U \mathcal{A}^a \phi(Z_{s-}) q_s(da) ds$ as a local P^* -martingale. Proposition 2 and the moment estimates of Theorem 3 make the stopped family uniformly integrable, hence $E^{P^*}[M_t^\phi] = 0$. Therefore $\int \phi dm_t^* - \int \phi dm_0 = \int_0^t E^{P^*}[\int_U \mathcal{A}^a \phi(Z_{s-}) q_s(da)] ds$. By the defining disintegration of η^{P^*} , $E^{P^*}[\int_U \mathcal{A}^a \phi(Z_s) q_s(da)] = \int_{\mathbb{R}^2} \int_U \mathcal{A}^a \phi(z) \hat{q}^*(s, z, da) m_s^*(dz)$ for ds -a.e. s . Substitution gives (46). $\square$

Corollary 41 (Relaxed nonlocal HJB-Kolmogorov system). *Let (m^*, P^*) be a relaxed equilibrium and set $v^* := V^{m^*}$. Then $(v^*, m^*, \hat{q}^*)$ satisfies*

$$\begin{cases} -\partial_t v^* - \mathcal{H}(t, x, y, v^*, Dv^*, D^2 v^*; v^*(t, \cdot, \cdot)) - F(t, (x, y), m_t^*) = 0, \\ \partial_t m_t^* = (\mathcal{A}^{\hat{q}_t^*})^* m_t^*, \\ v^*(T, x, y) = g(x, y) + G((x, y), m_T^*), \\ m_0^* = m_0. \end{cases} \quad (47)$$

The backward equation is understood in the viscosity sense and the forward equation in the distributional sense. If the equilibrium is strict Markovian, then $\hat{q}^*(t, z, da) = \delta_{\alpha^*(t, z)}(da)$ and the second equation reduces to $\partial_t m_t^* = (\mathcal{L}^{\alpha_t^*})^* m_t^*$.

Proof. Proposition 39, applied with $m = m^*$, gives the backward equation and terminal condition. Proposition 40 gives the forward equation. The strict Markov reduction follows from $\mathcal{A}^{\delta_{\alpha^*(t, z)}} = \mathcal{L}^{\alpha_t^*}$. $\square$

Remark 10. *The equilibrium feedback in this section is strategic but not a market-clearing feedback. A change in the hedging rule changes the jump amplitude through $h(u)$ in (29), thereby changing the population law m_t ; the changed law alters the reduced-form coupling F and hence the subsequent best response. In contrast, m_t does not feed back into the catastrophe intensity, the primitive severity law, or the replacement-cost dynamics. The equilibrium therefore captures an endogenous distributional externality among insurers, not an endogenous reconstruction-price or reconstruction-capacity mechanism [75].*

5.2 Existence and Uniqueness of Equilibrium

Existence is established in the relaxed formulation and transferred to strict feedback controls whenever the minimizing control can be selected uniquely.

Assumption 5.2 (Compact relaxed best responses). There exist $p \in (q, \bar{q}]$, $R_0 < \infty$, and a compact convex set $U_0 \subset U$ such that:

- (i) $m_0 \in \mathcal{P}_p(\mathbb{R}^2)$;
- (ii) for every $m \in \mathcal{M}_T$ with $\sup_{t \leq T} M_p(m_t) \leq R_0$, the frozen problem admits an optimal relaxed canonical law $P \in \mathfrak{A}(m)$ whose control coordinate is supported on U_0 $dt \otimes P$ -a.e.;
- (iii) if $m^n \rightarrow m$ in $\mathbf{W}_q$, $P^n \in \mathfrak{A}(m^n)$ are optimal, and $P^n \Rightarrow P$ in the canonical topology, then $P \in \mathfrak{A}(m)$ and P is optimal for the frozen problem associated with m ;
- (iv) the coefficients restricted to U_0 satisfy Assumptions 2.1-2.9 and 4.1 uniformly in $a \in U_0$.

Remark 11. No Markovianity is imposed in Assumption 5.2. In particular, the compactness and fixed-point argument below is carried out over the full convex set of relaxed canonical laws [63]. Markovianity is addressed only after existence of the relaxed equilibrium has been established.

For $R > 0$ and $\gamma \in (0, 1/2]$, retain $\mathcal{K}_{R,\gamma} := \{m \in \mathcal{M}_T : m_0 = m_0, \sup_{t \leq T} M_p(m_t) \leq R, W_q(m_t, m_s) \leq R|t - s|^\gamma\}$. Lemma 42 remains valid with “relaxed control” interpreted as an arbitrary relaxed canonical law supported on U_0 . For $m \in \mathcal{K}_{R,\gamma}$ define the full relaxed best-response set $\mathfrak{BR}(m) := \{P \in \mathfrak{A}(m) : J^m(P) = \inf_{Q \in \mathfrak{A}(m)} J^m(Q)\}$, and define $\Phi(m) := \{(P \circ Z_t^{-1})_{0 \leq t \leq T} : P \in \mathfrak{BR}(m)\}$. Thus $\mathfrak{BR}(m)$ contains all optimal relaxed laws, not only those admitting a Markov representation.

Lemma 42. Under Assumptions 2.1-2.5 and 5.2, there exist $R < \infty$ and $\gamma > 0$ such that every state-law flow induced by a relaxed control supported on U_0 belongs to $\mathcal{K}_{R,\gamma}$. Moreover, $\mathcal{K}_{R,\gamma}$ is compact in $C([0, T]; \mathcal{P}_q(\mathbb{R}^2))$ endowed with $\mathbf{W}_q$.

Proof. Uniform support of the controls in U_0 and Theorem 3 yield

$$\sup_u \mathbb{E} \left[\sup_{0 \leq t \leq T} |X_t|^p \right] + \mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t|^p \right] \leq C(1 + M_p(m_0)^p).$$

Hence all induced state laws have uniformly bounded p th moments [84]. For $0 \leq s < t \leq T$, the state equations, the BDG inequality, and the Bichteler-Jacod inequality yields $\mathbb{E}|Y_t - Y_s|^q \leq C|t - s|^{q/2}$, and $\mathbb{E}|X_t - X_s|^q \leq C(|t - s|^{q/2} + |t - s|)$. Therefore, for some $\gamma > 0$, we have $W_q(\mathcal{L}(X_t, Y_t), \mathcal{L}(X_s, Y_s)) \leq C|t - s|^\gamma$. Choosing R sufficiently large gives membership in $\mathcal{K}_{R,\gamma}$. Since $p > q$, the uniform p th-moment estimate implies uniform q -integrability and relative compactness of each time marginal in W_q . The common modulus of continuity gives equicontinuity. The metric Arzelà-Ascoli theorem then yields relative compactness in $C([0, T]; \mathcal{P}_q)$. The defining moment and modulus constraints are closed, so $\mathcal{K}_{R,\gamma}$ is compact. $\square$

Lemma 43. Suppose Assumptions 2.1-2.9, 4.1, 5.1, and 5.2 hold. If $m^n \rightarrow m$ in $\mathbf{W}_q$, $\mu^n \in \Phi(m^n)$, and $\mu^n \rightarrow \mu$ in $\mathbf{W}_q$, then $\mu \in \Phi(m)$.

Proof. Choose $P^n \in \mathfrak{BR}(m^n)$ with $\mu_t^n = P^n \circ Z_t^{-1}$. Compactness of U_0 , Lemma 42, the moment bounds of Theorem 3, and tightness of the canonical relaxed-control coordinate imply tightness of (P^n) on Ω° . Passing to a subsequence, $P^n \Rightarrow P$. Assumption 5.2(iii) gives $P \in \mathfrak{A}(m)$ and optimality of P for the frozen flow m .

For every t at which the canonical evaluation map is P -a.s. continuous, $P^n \circ Z_t^{-1} \Rightarrow P \circ Z_t^{-1}$. The uniform p -moment bound with $p > q$ upgrades this convergence to W_q . Since $\mu^n \rightarrow \mu$ in $\mathbf{W}_q$, $\mu_t = P \circ Z_t^{-1}$ on a dense set of times. Both $t \mapsto \mu_t$ and $t \mapsto P \circ Z_t^{-1}$ are W_q -continuous by Lemma 42; hence the equality holds for all $t \in [0, T]$. Therefore $P \in \mathfrak{BR}(m)$ and $\mu \in \Phi(m)$. $\square$

Lemma 44. *For every $m \in \mathcal{K}_{R,\gamma}$, $\mathfrak{BR}(m)$ is convex. Consequently, $\Phi(m)$ is convex.*

Proposition 45. *Under the hypotheses of Lemma 43, $\Phi : \mathcal{K}_{R,\gamma} \rightrightarrows \mathcal{K}_{R,\gamma}$ has nonempty compact convex values and is upper hemicontinuous.*

Proof. Nonemptiness follows from Assumption 5.2(ii). Convexity follows from Lemma 44. Let $m \in \mathcal{K}_{R,\gamma}$ and $(\mu^n) \subset \Phi(m)$. Compactness of $\mathcal{K}_{R,\gamma}$ gives a subsequence $\mu^{n_k} \rightarrow \mu$ in $\mathbf{W}_q$. Since $m^{n_k} \equiv m$, Lemma 43 yields $\mu \in \Phi(m)$; hence $\Phi(m)$ is closed in the compact set $\mathcal{K}_{R,\gamma}$ and is therefore compact. Lemma 43 also gives $\text{Gr}(\Phi)$ closed in $\mathcal{K}_{R,\gamma} \times \mathcal{K}_{R,\gamma}$. A compact-valued correspondence with closed graph from a compact metric space into a Hausdorff space is upper hemicontinuous. Thus Φ has all the properties claimed. $\square$

Theorem 46 (Existence of a relaxed mean field equilibrium). *Suppose Assumptions 2.1-2.9, 4.1, 5.1, and 5.2 hold. Then there exist $m^* \in \mathcal{K}_{R,\gamma}$ and $P^* \in \mathfrak{BR}(m^*)$ such that $m_t^* = P^* \circ Z_t^{-1}$ for every $t \in [0, T]$. Equivalently, (m^*, P^*) is a relaxed mean field equilibrium in the sense of Definition 2.*

Proof. Lemma 42 gives a nonempty compact convex set $\mathcal{K}_{R,\gamma}$. Proposition 45 gives a correspondence $\Phi : \mathcal{K}_{R,\gamma} \rightrightarrows \mathcal{K}_{R,\gamma}$ with nonempty compact convex values and closed graph. The Kakutani-Fa-Glicksberg theorem yields $m^* \in \mathcal{K}_{R,\gamma}$ such that $m^* \in \Phi(m^*)$. By the definition of Φ , there exists $P^* \in \mathfrak{BR}(m^*)$ with $m_t^* = P^* \circ Z_t^{-1}$ for all t . The first statement is optimality; the second is the consistency condition (45). $\square$

Assumption 5.3 (Markov realization). For every admissible relaxed law P supported on U_0 , let $\hat{q}^P$ be the disintegration kernel determined by $\eta^P(dt, dz, da) = dt m_t^P(dz) \hat{q}^P(t, z, da)$. Assume that the controlled martingale problem with the Markov relaxed kernel $\hat{q}^P$ admits a solution $\hat{P}$ satisfying $\hat{P} \circ Z_t^{-1} = P \circ Z_t^{-1}$ for every $t \in [0, T]$ and $J^m(\hat{P}) = J^m(P)$ whenever P is evaluated in the frozen environment m .

Remark 12. *Assumption 5.3 is a Markovianization hypothesis, not a convexity hypothesis. It is separated from the fixed-point argument because convexity of Φ is already obtained at the full relaxed-law level in Lemma 44. Markovian realization theorems for controlled martingale problems are developed in Kurtz and Stockbridge [43]. In applications, Assumption 5.3 may be verified by identifying the state-conditioned kernel $\hat{q}^P(t, z, \cdot)$ and proving well-posedness of the associated Markov martingale problem.*

Proposition 47. *Suppose the hypotheses of Theorem 46 and Assumption 5.3 hold. Then every relaxed equilibrium (m^*, P^*) admits an optimal relaxed Markov realization $(m^*, \hat{P}^*)$ with $\hat{P}^* \circ Z_t^{-1} = m_t^*$ for all t and $q_t(da) = \hat{q}^*(t, Z_{t-}, da) dt \otimes \hat{P}^*$ -a.e.*

Proof. Let (m^*, P^*) be furnished by Theorem 46. Disintegrate its occupation measure as $\eta^{P^*}(dt, dz, da) = dt m_t^*(dz) \hat{q}^*(t, z, da)$. Assumption 5.3 provides a Markov relaxed law $\hat{P}^*$ generated by $\hat{q}^*$ such that $\hat{P}^* \circ Z_t^{-1} = m_t^*$ for every t and $J^{m^*}(\hat{P}^*) = J^{m^*}(P^*)$. Since $P^* \in \mathfrak{BR}(m^*)$, $J^{m^*}(P^*) = V^{m^*, \text{rel}}$; therefore $J^{m^*}(\hat{P}^*) = V^{m^*, \text{rel}}$ and $\hat{P}^* \in \mathfrak{BR}(m^*)$. The marginal identity preserves (45), so $(m^*, \hat{P}^*)$ is a relaxed Markov equilibrium. $\square$

Corollary 48. *Suppose the hypotheses of Proposition 47 hold. Assume additionally that the optimal Markov relaxed kernel is $dt \otimes m_t^*$ -a.e. Dirac, i.e., $\hat{q}^*(t, z, da) = \delta_{\alpha^*(t, z)}(da)$ for a Borel selector $\alpha^* : [0, T] \times \mathbb{R}^2 \rightarrow U_0$. Then (m^*, α^*) is a strict Markov equilibrium. In particular, the conclusion applies when the frozen Hamiltonian has a unique pointwise minimizer and the hypotheses of Proposition 35 hold [74].*

Proof. By Proposition 47, $\hat{P}^*$ is optimal for the frozen flow m^* and has state marginals m^* . The Dirac representation gives the strict feedback control $u_t^* = \alpha^*(t, Z_{t-})$. Hence the optimality and consistency conditions in Definition 2 hold with a strict Markov control. $\square$

For uniqueness, we impose the monotonicity structure of Lasry and Lions [45].

Assumption 5.4 (Lasry-Lions monotonicity). For every $t \in [0, T]$ and $m, n \in \mathcal{P}_q(\mathbb{R}^2)$,

$$\int_{\mathbb{R}^2} (F(t, z, m) - F(t, z, n))(m - n)(dz) \geq 0, \quad (48)$$

and

$$\int_{\mathbb{R}^2} (G(z, m) - G(z, n))(m - n)(dz) \geq 0. \quad (49)$$

Equality in both monotonicity relations for an entire equilibrium flow implies equality of the two flows.

Assumption 5.5. For every $m \in \mathcal{K}_{R, \gamma}$, the frozen problem admits a unique optimal state-control law. A sufficient condition is strict convexity of the frozen Hamiltonian in the control variable on the effective compact set of Lemma 29, together with uniqueness in law of the controlled state equation.

Lemma 49. *Let (m^1, P^1) and (m^2, P^2) be two relaxed equilibria. Then*

$$\begin{aligned} & \int_0^T e^{-\delta t} \int_{\mathbb{R}^2} (F(t, z, m_t^1) - F(t, z, m_t^2))(m_t^1 - m_t^2)(dz) dt \\ & + e^{-\delta T} \int_{\mathbb{R}^2} (G(z, m_T^1) - G(z, m_T^2))(m_T^1 - m_T^2)(dz) \leq 0. \end{aligned} \quad (50)$$

Proof. Optimality yields $J^{m^1}(P^1) \leq J^{m^1}(P^2)$, and $J^{m^2}(P^2) \leq J^{m^2}(P^1)$. The cross-comparisons are legitimate because the controlled dynamics do not depend on the mean field. Adding the inequalities cancels the private running cost f , terminal cost g , and all control-dependent state costs. Using the equilibrium consistency relation and collecting the remaining coupling terms yields (50). $\square$

Theorem 50. *Suppose Assumptions 2.1-2.9, 4.1, and 5.1-5.5 hold. Then the mean field equilibrium is unique in law. If the frozen optimal control is generated by a unique Markov feedback, then the equilibrium feedback is unique $dt \otimes m_t^*$ -a.e.*

Proof. Let (m^1, P^1) and (m^2, P^2) be two equilibria. By Lemma 49, the sum of the two monotonicity expressions is nonpositive. Assumption 5.3 makes each expression nonnegative. Hence, both must vanish. The strictness clause in Assumption 5.3 implies $m_t^1 = m_t^2$ for every $t \in [0, T]$, where continuity upgrades almost-everywhere equality in time to pointwise equality. Let the common flow be m^* . Both P^1 and P^2 are optimal for the same frozen problem [61]. Assumption 5.4 therefore gives $P^1 = P^2$. If this law is induced by a unique Markov feedback, the feedbacks agree $dt \otimes m_t^*$ -a.e. $\square$

Corollary 51 (Uniqueness of the coupled HJB–Kolmogorov solution). *Under the hypotheses of Theorem 50, the equilibrium pair*

$$(v^*, m^*) \in (C([0, T] \times \mathbb{R}^2) \cap \mathcal{G}_q) \times \mathcal{K}_{R, \gamma}$$

solving (47) is unique within the class generated by admissible equilibrium controls.

Proof. Theorem 50 gives uniqueness of m^* . For this fixed flow, Proposition 39 and Corollary 33 give uniqueness of the backward value function. Proposition 40 identifies the forward component with the unique equilibrium law. $\square$

Proposition 52. *Let $\psi_D, \psi_S : \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous functions of at most q th-order growth and define $M_D(m) := \int \psi_D(z) m(dz)$, and $M_S(m) := \int \psi_S(z) m(dz)$. For $\gamma_D, \gamma_S \geq 0$, set*

$$F(t, z, m) = \gamma_D \psi_D(z) M_D(m) + \gamma_S \psi_S(z) M_S(m). \quad (51)$$

Then

$$\int (F(t, z, m) - F(t, z, n))(m - n)(dz) = \gamma_D (M_D(m) - M_D(n))^2 + \gamma_S (M_S(m) - M_S(n))^2 \geq 0.$$

Hence (51) satisfies Assumption 5.3.

Proof. By linearity,

$$\int \psi_D(z)(m - n)(dz) = M_D(m) - M_D(n),$$

and the same identity holds for ψ_S . Substitution into (51) gives the stated sum of squares. $\square$

Corollary 53. *Let $\psi_D(x, y)$ be an increasing transformation of replacement-cost exposure and let $\psi_S(x, y)$ be an increasing penalty for weak capitalization. Then the coupling (51) assigns a larger private cost to states whose exposure is aligned with a larger industry-wide concentration of the same exposure, while preserving the monotonicity required for Theorem 50.*

Proof. The conclusion follows from Proposition 52. The chosen functions alter only the economic interpretation of M_D and M_S , not the sum-of-squares monotonicity calculation. $\square$

Remark 13. *For every frozen population flow, Theorem 34 supplies the unique viscosity value function; the population law is propagated by the same jump-diffusion as in (4); and equilibrium requires that the frozen flow coincide with the law generated by its own optimal response [3]. No state-dependent catastrophe intensity, common noise, or population-dependent jump kernel is added in this section.*

6 Economic Implications

The preceding sections separate physical catastrophe severity, stochastic replacement-cost conditions, and the insurer's endogenous mitigation decision. In the present model these objects enter surplus multiplicatively through (29). An elevated realization of Y therefore rescales the monetary consequence of every catastrophe mark occurring in that state. Physical hedging acts on the same multiplicative channel through $h(u)$ [25]. The

model should not be interpreted as implying that catastrophe arrivals cause changes in Y ; rather, it quantifies the exposure created when catastrophe losses and high reconstruction costs coincide. Section 5 adds a second layer by allowing the resulting state distribution to feed back into individual incentives through the mean field coupling [26].

In classical ruin theory, subexponential asymptotics imply that extreme loss probabilities are governed by the heaviest-tailed component of the risk structure. The present objective is a finite-horizon stochastic-control functional rather than an infinite-horizon ruin probability, so those asymptotic results do not transfer directly. Nevertheless, they motivate examining whether the optimal hedge and capitalization outcomes are robust to alternative severity families.

6.1 Replacement-Cost Exposure and Optimal Hedging

For $p \in [2, p_\star]$ with $A_p := \int_E a(z)^p \mathbf{v}(dz) < \infty$, define

$$\Lambda_p(y, u) := \int_E \ell(y, z, u)^p \mathbf{v}(dz) = e^{py} h(u)^p A_p. \quad (52)$$

This is the instantaneous p th-moment intensity of the catastrophe jump amplitude.

From $\Lambda_p(y, u) = e^{py} h(u)^p A_p$, one obtains directly $\partial_y \log \Lambda_p(y, u) = p$ and, whenever h is differentiable, $\partial_u \log \Lambda_p(y, u) = ph'(u)/h(u) \leq 0$. Thus replacement-cost exposure has elasticity p with respect to the log replacement-cost state, while physical hedging weakly reduces every finite loss moment. In particular, if $2 \leq p_1 < p_2 \leq p_\star$, then the higher-order exposure is proportionally more sensitive to the replacement-cost state, since $\partial_y \log \Lambda_{p_2} / \partial_y \log \Lambda_{p_1} = p_2 / p_1 > 1$.

Corollary 54. *If $2 \leq p_1 < p_2 \leq p_\star$, then the proportional sensitivity of the higher finite moment to demand surge is larger $\frac{\partial_y \log \Lambda_{p_2}}{\partial_y \log \Lambda_{p_1}} = \frac{p_2}{p_1} > 1$.*

Variation in the replacement-cost factor does not change the tail index of the catastrophe mark law, the Lévy measure $\mathbf{v}$ is unchanged [11]. Instead, e^Y changes the monetary scale associated with a given catastrophe mark. Hence the heavy-tail structure remains that of the primitive catastrophe process, while the economic severity of its realizations depends on the contemporaneous replacement-cost state. For a smooth test function define

$$\mathfrak{C}[\varphi](t, x, y, u) := \int_E [\varphi(t, x - e^y a(z) h(u), y) - \varphi(t, x, y) + e^y a(z) h(u) \varphi_x(t, x, y)] \mathbf{v}(dz). \quad (53)$$

Proposition 55. *Suppose, for this comparative-static calculation, that $\mathbf{v}$ is C^2 in x , $h \in C^1$, and differentiation under the Lévy integral is valid. Then*

$$\partial_u \mathfrak{C}[\mathbf{v}] = -e^y h'(u) \int_E a(z) [\mathbf{v}_x(t, x - e^y a(z) h(u), y) - \mathbf{v}_x(t, x, y)] \mathbf{v}(dz). \quad (54)$$

If $\mathbf{v}$ is convex in x , then $\partial_u \mathfrak{C}[\mathbf{v}] \leq 0$.

Proof. Differentiate (53). Convexity makes $\mathbf{v}_x$ nondecreasing, while $x - e^y a(z) h(u) \leq x$, so the bracket in (54) is nonpositive. Since $h'(u) \leq 0$, the prefactor $-e^y h'(u)$ is nonnegative. $\square$

Corollary 56. *Holding the local curvature profile of v and u fixed, the absolute scale of the nonlocal catastrophe-risk reduction from a marginal increase in hedging is amplified by the factor e^y .*

Proof. This is immediate from (54). The qualification is necessary because the displaced derivative also changes with y . $\square$

This is a local mechanism, not a global theorem that the optimal feedback is increasing in y . The rigorous policy remains the measurable Hamiltonian selector of Proposition 35. For the quadratic benchmark $c(u) = C_H(u) = \chi u^2/2$ with control-independent σ_X , the formal first-order condition (35) becomes

$$\chi u(1 - v_x) = e^y h'(u) \int_E a(z) [v_x(t, x - e^y a(z) h(u), y) - v_x(t, x, y)] \nu(dz). \quad (55)$$

The left side is the marginal resource-and-drift cost of hedging evaluated at the shadow value of surplus; the right side is the marginal reduction in nonlocal continuation cost.

Proposition 57. *If the pointwise Hamiltonian is strictly convex in u and its derivative changes sign in the interior of U , then the pointwise hedge is unique and is characterized by (55). This is compatible with Corollary 48.*

Proof. Strict convexity makes the derivative strictly increasing, so a sign change gives a unique zero and hence the unique minimizer. Corollary 48 gives the strict Markov equilibrium representation under the corresponding selector hypotheses. $\square$

The mechanism is consistent with empirical work documenting post-disaster increases in reconstruction costs and labor scarcity, commonly termed demand surge. See Olsen and Porter [49], Döhrmann et al. [20], and Kousky [42]. Mean reversion in (3) makes the value of pre-arranged capacity state and horizon dependent: hedging is valuable when catastrophe exposure and temporary reconstruction scarcity overlap [11].

6.2 Systemic Effects of Equilibrium Hedging

By Definition 2, an equilibrium feedback is optimal against the population flow that it generates. Theorem 46 guarantees a relaxed equilibrium, Corollary 48 gives a strict Markov representation under the selector hypothesis, and Theorem 50 rules out competing equilibrium laws under the monotonicity assumptions.

For a feedback α define the aggregate p th catastrophe-loss exposure

$$\mathcal{A}_p(t; m, \alpha) := A_p \int_{\mathbb{R}^2} e^{py} h(\alpha(t, x, y))^p m_t(dx, dy). \quad (56)$$

For a fixed cross-sectional law m_t , the monotonicity of h gives $\alpha_1 \geq \alpha_2$ m_t -a.e. $\Rightarrow \mathcal{A}_p(t; m, \alpha_1) \leq \mathcal{A}_p(t; m, \alpha_2)$. Hence an industry-wide increase in physical hedging reduces every finite aggregate catastrophe-loss exposure for which $A_p < \infty$. This is a direct loss-scale effect and should be distinguished from the endogenous change in m_t generated by equilibrium behavior.

For a fixed cross-sectional state law, an industry-wide increase in physical hedging reduces every finite aggregate catastrophe-loss exposure (56). It does so without changing catastrophe arrival intensity. In equilibrium the state law is not fixed. Corollary 41 propagates the endogenous distributional response [17]. Hence

the total systemic effect decomposes into direct attenuation through $h(\alpha)$ and a distributional effect through m ; only the first has an unconditional sign under the present assumptions. For the coupling in Proposition 52,

$$F(t, z, m) = \gamma_D \Psi_D(z) M_D(m) + \gamma_S \Psi_S(z) M_S(m).$$

Corollary 53 interprets these moments as demand-surge exposure and weak capitalization. The following calculation concerns the reduced-form strategic externality encoded by F ; it should not be interpreted as an endogenously determined reconstruction-market price.

Proposition 58. *For fixed individual state z , a change in aggregate moments changes the running coupling by $\Delta F(t, z) = \gamma_D \Psi_D(z) \Delta M_D + \gamma_S \Psi_S(z) \Delta M_S$. If $\Psi_D(z), \Psi_S(z) \geq 0$, an increase in either aggregate vulnerability moment weakly raises the private cost of occupying z .*

Proof. Subtract the coupling evaluated at the two population laws and use $\gamma_D, \gamma_S \geq 0$. □

Under Assumption 5.4, changes in aggregate vulnerability alter the representative insurer's unique frozen best response; the resulting feedback changes jump amplitudes through (29), changes the forward law through Proposition 40, and is closed by the consistency condition (45). The equilibrium externality is strategic rather than physical contagion. One insurer's surplus does not enter another insurer's state equation, and Section 5 introduces neither common noise nor a population-dependent catastrophe intensity. The cost of occupying a vulnerable capitalization demand-surge state nevertheless depends on the industry's concentration in comparable states. Mean field models have been used to isolate analogous endogenous systemic channels in financial systems [11]. Under Theorem 50, distinct high-exposure and low-exposure population flows cannot both be self-consistent equilibrium regimes for the same primitives. Under Theorem 50, any measurable equilibrium statistic of (m^*, α^*) , including (56), $M_D(m_t^*)$, $M_S(m_t^*)$, and the surplus distribution, is single-valued for fixed model primitives. Physical hedging should not be interpreted as a mechanical substitute for reinsurance or liquid capital. It operates earlier in the loss-production mechanism by reducing the residual fraction $h(u)$ before the catastrophe loss enters surplus [78]. This is consistent with the disaster-insurance literature emphasizing the interaction between insurance, recovery, and ex ante mitigation [42]. A subsidy to pre-arranged capacity, for example, changes the private Hamiltonian immediately, but its total equilibrium effect is determined only after the induced feedback is propagated through the Kolmogorov equation and the mean field fixed point.

7 Calibration and Numerical Analysis

We calibrate the numerical model in normalized annual units. The calibration is designed to preserve the empirical features motivating the stochastic model while remaining consistent with the moment and regularity conditions imposed in Sections 2 -5. More broadly, recent simulation-based work emphasizes the value of stochastic scenario analysis for financial decisions under volatile price environments [12]. We cite this work only as a recent example of simulation-based analysis under price uncertainty. In particular, the empirically estimated heavy-tail [99] index is retained, but the far tail of the catastrophe mark measure is tempered so that the p_* -moment assumption required by Assumption 2.1 remains valid. Let one unit of surplus represent the insurer's initial capitalization and normalize the long-run replacement-cost index to one. Thus, $X_0 = 1$, $\vartheta = 0$, and $I_0 = e^{Y_0} = 1$. The time unit is one year. The numerical horizon is five years, which is sufficiently long

relative to the estimated mean-reversion time scale of the construction-cost process. For the catastrophe marks, we use the tempered Pareto Lévy density

$$\mathbf{v}(dz) = \lambda_J \alpha z_{\min}^{\alpha} z^{-1-\alpha} e^{-\tau_J(z-z_{\min})} \mathbf{1}_{\{z \geq z_{\min}\}} dz. \quad (57)$$

The tempered Pareto specification is used as the baseline rather than as a claim that catastrophe severities belong uniquely to this parametric family. Heavy-tailed insurance losses may be represented by a substantially broader class of subexponential distributions, including regularly varying, lognormal, and heavy-tailed Weibull models; see Embrechts et al. [27] and Klüppelberg and Mikosch [41]. This distinction is relevant because tail-equivalent behavior [35], finite-sample loss quantiles, and ruin-type functionals may differ materially across subexponential families. The analytical results of Sections 2-5 depend on the stated moment and continuity conditions on the jump measure and not on the tempered Pareto form itself [80]. The parametric choice becomes consequential in the numerical analysis, and we therefore supplement the baseline calibration with a tail-family [67] robustness experiment below. The power-law component reproduces the empirical tail estimate $\hat{\alpha} = 1.71$, while the exponential tempering parameter $\tau_J > 0$ guarantees $\int_E z^p \mathbf{v}(dz) < \infty$ for every $p > 0$. Hence, the numerical jump specification is compatible with the $p_* > 2$ integrability imposed in Assumption 2.1. Over moderate and large losses below the tempering scale [56], the model retains the heavy-tailed behavior implied by the empirical estimate. The construction-cost factor is simulated from $dY_t = \kappa(\vartheta - Y_t) dt + \sigma_I dW_t^I$, with the empirical mean-reversion estimate $\hat{\kappa} = 7.07$. Its corresponding half-life is

$$t_{1/2} = \frac{\log 2}{\kappa} \approx 0.098 \text{ years}, \quad (58)$$

or approximately 35.8 days. Thus the calibrated construction-cost shock is economically sharp but transitory, as required by the demand-surge interpretation. The remaining coefficients are structural benchmark values rather than direct statistical estimates [66]. They are chosen in normalized units so that neither the diffusion component nor the control cost dominates the catastrophe jump term mechanically. The baseline calibration is reported in Table 1.

Table 1: Baseline calibration for the stochastic control and mean field simulations

Parameter	Baseline value	Role	Calibration status
T	5	Time horizon, years	Numerical
X_0	1	Initial normalized insurer surplus	Normalization
Y_0	0	Initial log replacement-cost factor	Normalization
ϑ	0	Long-run log replacement-cost level	Normalization
κ	7.07	Mean-reversion speed	Empirical
σ_I	0.20	Construction-cost volatility	Benchmark
α	1.71	Catastrophe-loss tail exponent	Empirical
λ_J	0.20	Catastrophe arrival intensity, annual	Benchmark
$z_{\min}$	0.01	Minimum normalized catastrophe mark	Normalization
τ_J	0.10	Exponential tail-tempering parameter	Regularization
r	0.03	Surplus accumulation rate	Benchmark
π_0	0.08	Baseline net premium inflow	Benchmark
σ_X	0.10	Noncatastrophe surplus volatility	Benchmark
γ	1.00	Hedging effectiveness in $h(u) = e^{-\gamma u}$	Normalization
χ	0.20	Quadratic physical-hedging cost	Benchmark
δ	0.03	Social discount rate	Benchmark
x_{crit}	0.50	Capitalization threshold	Benchmark
ω	2.00	Solvency-penalty weight	Benchmark
ζ	0.50	Demand-surge cost weight	Benchmark
m	2	Demand-surge penalty exponent	Benchmark
q	2	Solvency-loss exponent	Structural
$p_\star$	3	Moment order used in numerical bounds	Structural
γ_D	0.25	Mean field demand-surge interaction	Benchmark
γ_S	0.50	Mean field capitalization interaction	Benchmark

For the drift specification introduced in Section 3, we use $b(x, y, u) = rx + \pi_0 - \frac{\chi}{2}u^2$, and take $\sigma_X(x, y, u) \equiv \sigma_X$. The physical-hedging response is $h(u) = e^{-\gamma u}$, which satisfies the monotonicity and convexity requirements imposed in Section 3, $h'(u) < 0$, with $h''(u) > 0$. The running social cost is taken to be

$$f(t, x, y, u) = \frac{\chi}{2}u^2 + \zeta(e^y - 1)_+^m + \omega(x_{\text{crit}} - x)_+^q. \quad (59)$$

This specification separates the resource cost of physical hedging, the social cost of demand-surge inflation, and the cost of weak insurer capitalization [54]. It also preserves the coercivity in u required in Assumption 2.6. For the mean field component, we use the monotone coupling introduced in Proposition 52,

$$F(t, z, \mu) = \gamma_D \psi_D(z) M_D(\mu) + \gamma_S \psi_S(z) M_S(\mu), \quad (60)$$

with $\psi_D(x, y) = (e^y - 1)_+$, and $\psi_S(x, y) = (x_{\text{crit}} - x)_+$. By Proposition 52, (60) satisfies the Lasry-Lions monotonicity condition used in Theorem 50. The distinction between estimated and benchmark parameters

is important. The values $\hat{\alpha} = 1.71$, and $\hat{\kappa} = 7.07$ are inherited from the empirical calibration motivating the model. The remaining parameters in Table 1 define the baseline numerical economy and will subsequently be varied over economically meaningful ranges. The numerical conclusions will therefore be based primarily on comparative statics rather than on interpreting the benchmark normalizations as point estimates.

7.1 Discretization and Computational Scheme

We solve the calibrated control and mean field problems by combining exact time stepping for the mean-reverting replacement-cost factor [22], Monte Carlo generation of the tempered catastrophe marks, a monotone discretization of the nonlocal HJB equation [79], policy iteration for the physical-hedging feedback, and a damped fixed-point iteration for the mean field equilibrium. The computational procedure is constructed to respect the probabilistic model of Sections 2 and 3 rather than replacing the jump process by a Gaussian approximation [86].

Let

$$0 = t_0 < t_1 < \dots < t_{N_t} = T, \quad \Delta t = T/N_t,$$

and introduce rectangular state grids

$$x_i = x_{\min} + i\Delta x, \quad i = 0, \dots, N_x,$$

and

$$y_j = y_{\min} + j\Delta y, \quad j = 0, \dots, N_y.$$

The numerical domain is chosen sufficiently wide that the probability of reaching its artificial boundaries under the baseline calibration is negligible. Grid enlargement is subsequently used as a robustness check.

Exact discretization of the replacement-cost factor. Because the process in (3) is Ornstein-Uhlenbeck, its transition law is known explicitly. We therefore do not introduce Euler discretization error into the construction-cost state [55]. Conditional on $Y_{t_n} = Y_n$,

$$Y_{n+1} = \vartheta + (Y_n - \vartheta)e^{-\kappa\Delta t} + \sigma_I \sqrt{\frac{1 - e^{-2\kappa\Delta t}}{2\kappa}} \xi_{n+1}, \quad \xi_{n+1} \sim N(0, 1). \quad (61)$$

Consequently,

$$Y_{n+1} \mid Y_n \sim N\left(\vartheta + (Y_n - \vartheta)e^{-\kappa\Delta t}, \frac{\sigma_I^2}{2\kappa}(1 - e^{-2\kappa\Delta t})\right).$$

The replacement-cost index is then recovered as $I_{n+1} = e^{Y_{n+1}}$. Equation (61) is used both in the Monte Carlo experiments and in the validation of the finite-difference solution.

Simulation of catastrophe arrivals and severities. Under (57), the total jump intensity is

$$\lambda_v := v(E) = \lambda_J \int_{z_{\min}}^{\infty} \alpha z_{\min}^{\alpha} z^{-1-\alpha} e^{-\tau_J(z-z_{\min})} dz.$$

Over a time interval of length Δt , the number of catastrophe marks is sampled as $K_n \stackrel{iid}{\sim} \text{Poisson}(\lambda_v \Delta t)$. Conditional on an arrival, the mark density is

$$f_Z(z) = \frac{\alpha z_{\min}^\alpha z^{-1-\alpha} e^{-\tau_J(z-z_{\min})}}{\int_{z_{\min}}^{\infty} \alpha z_{\min}^\alpha r^{-1-\alpha} e^{-\tau_J(r-z_{\min})} dr}, \quad z \geq z_{\min}. \quad (62)$$

Marks from (62) are generated by acceptance-rejection [73]. We first draw $Z^{(0)} = z_{\min}(1-U)^{-1/\alpha}$, where $U \sim \text{Unif}(0,1)$, from the Pareto envelope and accept it with probability $\exp\{-\tau_J(Z^{(0)} - z_{\min})\}$. The accepted value has density (62). Given state (Y_n, u_n) , the associated monetary loss is $L_n = e^{Y_n} a(Z_n) h(u_n)$. Thus the simulation preserves exactly the multiplicative loss mechanism used in (29). For Monte Carlo simulation [72], the surplus is advanced according to

$$X_{n+1} = X_n + b(X_n, Y_n, u_n) \Delta t + \sigma_X \sqrt{\Delta t} \varepsilon_{n+1} - \sum_{k=1}^{K_n} e^{Y_n} a(Z_{n,k}) h(u_n) + \Delta t \int_E e^{Y_n} a(z) h(u_n) \nu(dz), \quad (63)$$

where $\varepsilon_{n+1} \sim N(0,1)$. The final term is the compensator required by the compensated jump representation in (4).

Discretization of the nonlocal HJB equation. Let $V_{i,j}^n \approx V(t_n, x_i, y_j)$. We solve backward from $V_{i,j}^{N_t} = g(x_i, y_j)$. For the local terms, the second derivatives are approximated by centered differences,

$$D_{xx} V_{i,j}^n = \frac{V_{i+1,j}^n - 2V_{i,j}^n + V_{i-1,j}^n}{(\Delta x)^2},$$

and

$$D_{yy} V_{i,j}^n = \frac{V_{i,j+1}^n - 2V_{i,j}^n + V_{i,j-1}^n}{(\Delta y)^2}.$$

The first-order drift terms are discretized by an upwind operator, denoted by $D_x^u V_{i,j}^n$, and $D_y V_{i,j}^n$, with the upwind direction chosen according to the signs of $b(x_i, y_j, u)$ and $\kappa(\vartheta - y_j)$. For each control value u , define the discrete nonlocal operator

$$I_\Delta^u V_{i,j}^n := \sum_{k=1}^{N_z} w_k \left[\mathcal{I}_x V^n(x_i - e^{y_j} a(z_k) h(u), y_j) - V_{i,j}^n + e^{y_j} a(z_k) h(u) D_x^u V_{i,j}^n \right],$$

where $\{z_k, w_k\}_{k=1}^{N_z}$ is a positive quadrature rule for $\nu(dz)$ and $\mathcal{I}_x$ denotes monotone linear interpolation in the surplus direction. Positive quadrature weights and monotone interpolation are used so that the discrete jump operator preserves the order structure required by the viscosity solution.

For a fixed policy $u_{i,j}^n$, the backward equation is discretized implicitly as

$$\begin{aligned} \frac{V_{i,j}^n - V_{i,j}^{n+1}}{\Delta t} + f(t_n, x_i, y_j, u_{i,j}^n) - \delta V_{i,j}^n \\ + b(x_i, y_j, u_{i,j}^n) D_x^u V_{i,j}^n + \kappa(\vartheta - y_j) D_y V_{i,j}^n + \frac{1}{2} \sigma_X^2 D_{xx} V_{i,j}^n + \frac{1}{2} \sigma_I^2 D_{yy} V_{i,j}^n + I_{\Delta}^{u_{i,j}^n} V_{i,j}^n = 0. \end{aligned} \quad (64)$$

The fully implicit treatment of the local diffusion terms improves stability when κ is large, as in the empirical calibration $\kappa = 7.07$. The jump quadrature is truncated at $z_{\max}$ and selected so that $\int_{z_{\max}}^{\infty} (1 + z^q) \nu(dz)$ is numerically negligible [72]. Because the calibrated measure is exponentially tempered, the truncation error decays exponentially in $z_{\max}$.

Policy iteration for physical hedging. The nonlinear minimization in the HJB equation is solved by Howard-type policy iteration. Let $u^{(r)}$ denote the policy at iteration r . Given $u^{(r)}$, solve the linear system $\mathcal{L}_{\Delta}^{u^{(r)}} V^{(r+1)} + f^{u^{(r)}} = 0$ backward in time, where $\mathcal{L}_{\Delta}^u$ denotes the discrete operator in (64). The policy is then updated pointwise according to

$$u_{i,j}^{(r+1),n} \in \arg \min_{u \in U_{\Delta}} \left\{ f(t_n, x_i, y_j, u) + b(x_i, y_j, u) D_x V_{i,j}^{(r+1),n} + I_{\Delta}^u V_{i,j}^{(r+1),n} \right\}, \quad (65)$$

where $U_{\Delta} = \{0, \Delta u, 2\Delta u, \dots, u_{\max}\}$. If σ_X depends on u , its corresponding discrete diffusion term is included in the minimization in (65). The iteration terminates when

$$\max_{n,i,j} \left| u_{i,j}^{(r+1),n} - u_{i,j}^{(r),n} \right| \leq \varepsilon_{\text{PI}}. \quad (66)$$

For the baseline computations we use $\varepsilon_{\text{PI}} = 10^{-7}$. The monotone discretization and policy iteration are consistent with the viscosity solution of Section 4; numerical methods of this type for HJB and nonlocal mean field equations are developed, for example, in Achdou [1] and Chowdhury et al. [13].

Forward discretization of the population law. For a fixed feedback policy $\alpha_{i,j}^n$, the population distribution is advanced using the discrete adjoint of the controlled generator. Let $m_{i,j}^n$ denote the probability mass at (x_i, y_j) at time t_n . The forward step is

$$m^{n+1} = m^n + \Delta t \left(\mathcal{L}_{\Delta}^{\alpha^n} \right)^* m^n, \quad (67)$$

subject to $m_{i,j}^{n+1} \geq 0$, $\sum_{i,j} m_{i,j}^{n+1} = 1$. A positivity-preserving flux discretization is used for the local drift terms, while the jump redistribution is implemented with the same interpolation weights used in the backward nonlocal operator [69]. This backward-forward consistency is important: the discrete Kolmogorov operator is the adjoint of the discrete controlled generator appearing in the HJB equation [59].

Mean field fixed-point iteration. Let $m^{(k)}$ denote the population-flow iterate. Starting from the uncontrolled law $m^{(0)} = \mathcal{L}(X^{u \equiv 0}, Y)$, the k th fixed-point step consists of

$$m^{(k)} \longrightarrow F(\cdot, \cdot, m^{(k)}) \longrightarrow V^{(k)} \longrightarrow \alpha^{(k)} \longrightarrow \widehat{m}^{(k+1)}.$$

The backward HJB equation is first solved using the frozen coupling $m^{(k)}$. Policy iteration yields the corresponding best-response feedback $\alpha^{(k)}$. Equation (67) is then solved from m_0 to produce the implied population flow $\widehat{m}^{(k+1)}$. To stabilize the forward-backward iteration, we use relaxation

$$m^{(k+1)} = (1 - \omega_{\text{MFG}})m^{(k)} + \omega_{\text{MFG}}\widehat{m}^{(k+1)}, \quad 0 < \omega_{\text{MFG}} \leq 1. \quad (68)$$

The baseline value is $\omega_{\text{MFG}} = 0.25$. The iteration is stopped when

$$\max_{0 \leq n \leq N_t} W_1(m_n^{(k+1)}, m_n^{(k)}) \leq \varepsilon_{\text{MFG}}, \quad \varepsilon_{\text{MFG}} = 10^{-6}. \quad (69)$$

The iteration (68) is used as a computational fixed-point procedure; uniqueness of the continuous mean field equilibrium does not by itself imply convergence of this iteration [82]. Numerical convergence is therefore assessed independently through the residual criterion (69), robustness to the relaxation parameter, and convergence from multiple initial population flows.

Numerical consistency checks. We evaluate numerical reliability using three diagnostics. First, the HJB residual associated with (64) is computed over the full interior grid. Second, the total probability mass in (67) is monitored at every time step. Third, the equilibrium residual is defined by

$$\mathcal{R}_{\text{MFG}} := \max_{0 \leq n \leq N_t} W_1(m_n^*, \widehat{m}_n^*). \quad (70)$$

A converged numerical equilibrium must satisfy simultaneously $\|\mathcal{R}_{\text{HJB}}\|_{\infty} \mathbf{1}$, $\mathcal{R}_{\text{MFG}} \mathbf{1}$, and $\max_n \left| \sum_{i,j} m_{i,j}^n - 1 \right| \mathbf{1}$. Finally, all reported comparative statics are repeated after halving Δt , Δx , Δy , and Δu , and after increasing z_{max} . The numerical conclusions [38] are regarded as stable only when the resulting changes in the value function, equilibrium hedge, and aggregate exposure (56) are negligible relative to the reported economic effects.

7.2 Numerical Experiments and Comparative-Statics Design

We organize the numerical experiments around the two economic mechanisms identified in Sections 6 and 5: the interaction between demand-surge exposure and physical hedging at the individual-insurer level, and the endogenous change in aggregate catastrophe exposure generated by equilibrium hedging [103]. The baseline economy is the calibration reported in Table 1, and the numerical solution is computed using the backward-forward procedure of Section 7.1. All comparative statics alter one primitive at a time unless otherwise stated. The computational domain and baseline grid are

$$(t, x, y, u) \in [0, 5] \times [-1, 3] \times [-0.75, 0.75] \times [0, 2], \quad (N_t, N_x, N_y, N_u) = (500, 400, 150, 100). \quad (71)$$

Hence, $\Delta t = 0.01$, $\Delta x = 0.01$, $\Delta y = 0.01$, and $\Delta u = 0.02$. The interval for y contains economically large replacement-cost shocks $e^{-0.75} \approx 0.472$, and $e^{0.75} \approx 2.117$. Thus the upper boundary corresponds to reconstruction costs exceeding their long-run level by more than 100% [33]. The surplus domain permits both severe undercapitalization and substantial positive capitalization [37]. The catastrophe integral is evaluated on $[z_{\min}, z_{\max}] = [0.01, 20]$, using positive quadrature weights. The exponentially tempered tail outside $z_{\max}$ is monitored separately [68]. The baseline number of quadrature nodes is $N_z = 200$. To separate capitalization effects from demand-surge effects, the optimal feedback is evaluated at the benchmark states

$$\mathcal{S} = \left\{ \begin{array}{l} (0.40, -0.20), (0.40, 0), (0.40, 0.20), \\ (1.00, -0.20), (1.00, 0), (1.00, 0.20), \\ (1.60, -0.20), (1.60, 0), (1.60, 0.20) \end{array} \right\}. \quad (72)$$

The three surplus levels represent weakly capitalized, baseline, and strongly capitalized insurers. The three demand-surge states imply replacement-cost multipliers $e^{-0.20} \approx 0.819$, 1, and $e^{0.20} \approx 1.221$. The resulting 3×3 design makes it possible to identify whether the response of physical hedging to demand surge depends materially on the insurer's capitalization [81].

The first experiment varies the initial demand-surge state while holding all remaining primitives at their baseline values:

$$Y_0 \in \{-0.30, -0.20, -0.10, 0, 0.10, 0.20, 0.30, 0.40\}. \quad (73)$$

These values correspond to initial reconstruction-cost multipliers between approximately 0.741 and 1.492. For each Y_0 , we record $u^*(0, X_0, Y_0)$, $V(0, X_0, Y_0)$, together with the expected discounted hedging expenditure, the probability of weak capitalization [104], and the aggregate catastrophe-loss exposure defined in (56). The principal numerical elasticity is

$$\mathcal{E}_{u,y} := \frac{u^*(0, X_0, y + \Delta y) - u^*(0, X_0, y - \Delta y)}{2\Delta y}. \quad (74)$$

Its sign is therefore a numerical equilibrium implication rather than an assumption of the model. To determine whether stronger tail risk induces greater physical hedging, we vary

$$\alpha \in \{1.50, 1.60, 1.71, 1.80, 2.00, 2.25\}, \quad (75)$$

holding τ_J fixed at its baseline value. Lower α places more probability on large catastrophe marks over the empirically relevant range [53]. For each value of α , the jump density is renormalized according to (62). We determine the optimal initial hedge, the value function, and $\mathcal{A}_p(t; m^*, \alpha^*)$ for the admissible moment orders used in the numerical analysis. Because exponential tempering remains in force, all experiments remain consistent with the moment conditions of Section 2.

Heavy-tail family robustness. To distinguish sensitivity to the tail exponent from sensitivity to the parametric tail family, we repeat the benchmark computation under three severity specifications: the tempered Pareto baseline, a lognormal distribution, and a heavy-tailed Weibull distribution. The alternative distributions are normalized to have the same reference upper quantile as the baseline severity law, so that differences in the resulting control are driven primarily by tail shape rather than by an arbitrary rescaling of losses. Let

F_{TP} , F_{LN} , and F_{W} denote the three severity distribution functions [83]. For a fixed reference level p_0 , the scale parameters of the alternatives are selected so that $F_{\text{LN}}^{-1}(p_0) = F_{\text{W}}^{-1}(p_0) = F_{\text{TP}}^{-1}(p_0)$. We use $p_0 = 0.95$ in the baseline robustness exercise. For each family, the positive quadrature rule is reconstructed and the HJB problem is resolved without changing the remaining economic parameters [98]. The comparison is based on the initial optimal hedge $u^*(0, X_0, Y_0)$, the initial value $V(0, X_0, Y_0)$, and the weak-capitalization probability P_{weak} . This experiment is not intended to select a universally preferred catastrophe distribution. Its purpose is to measure how strongly the economic conclusions depend on the assumed shape of the severity tail within the broader heavy-tailed class.

Strictly speaking, exponential tempering removes regular variation in the far tail and produces finite moments of every order. We therefore use the term “tempered heavy-tail benchmark” for the numerical specification: it preserves Pareto-type behavior over the empirically relevant range while regularizing the extreme asymptotic tail so that the simulation remains compatible with the moment assumptions of the analytical model [16].

The annual catastrophe intensity is varied over

$$\lambda_J \in \{0.05, 0.10, 0.20, 0.30, 0.40, 0.50\}. \quad (76)$$

The corresponding expected interarrival times under the unnormalized benchmark interpretation range from twenty years to two years. This experiment separates the effect of catastrophe frequency from the severity effect studied through α . For each intensity, the relevant policy response is summarized by $\Delta_\lambda u^* = u_{\lambda_J}^*(0, X_0, Y_0) - u_{\lambda_{\text{base}}}^*(0, X_0, Y_0)$. The mean-reversion coefficient is varied according to

$$\kappa \in \{1, 2, 4, 7.07, 10, 15\}. \quad (77)$$

The associated half-life $t_{1/2}(\kappa) = \frac{\log 2}{\kappa}$ ranges from approximately 253 days at $\kappa = 1$ to approximately 17 days at $\kappa = 15$. This experiment distinguishes the magnitude of a demand-surge shock from its persistence. For a fixed initial $Y_0 > 0$, a smaller κ implies that elevated reconstruction costs are expected to remain economically relevant for longer. We therefore compare $u^*(0, X_0, Y_0; \kappa)$ and the expected cumulative hedging expenditure across the values in (77). The quadratic resource-cost parameter is varied over

$$\chi \in \{0.05, 0.10, 0.20, 0.30, 0.50, 0.75, 1.00\}. \quad (78)$$

This experiment measures the shadow value of access to pre-arranged reconstruction capacity. For each χ , we calculate the optimal policy and the percentage reduction in catastrophe exposure relative to the uncontrolled insurer:

$$\mathfrak{R}_p(t) := 100 \left[1 - \frac{\mathcal{A}_p(t; m^*, \alpha^*)}{\mathcal{A}_p(t; m^0, 0)} \right]. \quad (79)$$

Here m^0 denotes the state law generated under the no-hedging policy $u \equiv 0$. A positive value of $\mathfrak{R}_p$ therefore measures the percentage reduction in aggregate p th-order catastrophe-loss exposure attributable to optimal physical hedging. The initial surplus is varied over

$$X_0 \in \{0.30, 0.40, 0.50, 0.75, 1.00, 1.25, 1.50, 2.00\}. \quad (80)$$

The capitalization threshold remains $x_{\text{crit}} = 0.50$. We record the policy $u^*(0, X_0, Y_0)$ and the probability

$$P_{\text{weak}} := \mathbb{P} \left(\inf_{0 \leq t \leq T} X_t < x_{\text{crit}} \right). \quad (81)$$

The statistic in (81) is estimated from the controlled Monte Carlo paths generated using (61) and (63). To isolate the systemic channel, we vary the interaction coefficients in (60). The demand-surge interaction takes values $\gamma_D \in \{0, 0.10, 0.25, 0.50, 0.75, 1.00\}$, while the capitalization interaction takes values $\gamma_S \in \{0, 0.10, 0.25, 0.50, 0.75, 1.00\}$. The case $(\gamma_D, \gamma_S) = (0, 0)$ is the nonstrategic benchmark: insurers solve independent stochastic control problems and the population distribution does not enter their objective. The systemic effect of the mean field interaction is measured by

$$\mathfrak{S}_p := 100 \left[1 - \frac{\int_0^T \mathcal{A}_p(t; m^*, \alpha^*) dt}{\int_0^T \mathcal{A}_p(t; m^{\text{ind}}, \alpha^{\text{ind}}) dt} \right], \quad (82)$$

where $(m^{\text{ind}}, \alpha^{\text{ind}})$ denotes the population law and optimal policy when $\gamma_D = \gamma_S = 0$. Thus $\mathfrak{S}_p > 0$ means that strategic equilibrium interaction reduces cumulative aggregate catastrophe exposure relative to the independent-insurer benchmark [40]. We also report the equilibrium change in weak capitalization,

$$\Delta_{\text{sys}} P_{\text{weak}} := P_{\text{weak}}^{\text{MFG}} - P_{\text{weak}}^{\text{ind}}. \quad (83)$$

A negative value indicates that equilibrium hedging reduces the probability of entering the weak-capitalization region. The economically most severe states arise when a heavy catastrophe mark coincides with elevated reconstruction costs [77]. We therefore conduct a two-dimensional stress experiment over

$$(\alpha, Y_0) \in \{1.50, 1.71, 2.00\} \times \{0, 0.20, 0.40\}. \quad (84)$$

For each pair we compute the optimal hedge, expected social cost, weak-capitalization probability, and cumulative catastrophe exposure. To measure the economic value of allowing the hedge to respond dynamically to demand surge, define

$$\mathfrak{G} := 100 \frac{J(u^{\text{static}}) - J(u^*)}{|J(u^{\text{static}})|}, \quad (85)$$

where u^{static} is the optimal constant hedge obtained by restricting the admissible policy class to time- and state-independent controls. Positive $\mathfrak{G}$ measures the percentage reduction in social catastrophe cost obtained from the dynamic feedback policy. This comparison is particularly informative because both strategies operate through the same physical-hedging technology. Hence (85) isolates the economic value of state contingency rather than the value of hedging itself. For each baseline or counterfactual specification, the computed feedback is evaluated using N_{MC} independent simulated paths. The baseline choice is

$$N_{\text{MC}} = 100000. \quad (86)$$

For a generic path-dependent statistic H , its Monte Carlo estimate and standard error are $\hat{H} = \frac{1}{N_{\text{MC}}} \sum_{r=1}^{N_{\text{MC}}} H(r)$,

and $\text{SE}(\hat{H}) = \frac{\hat{\sigma}_H}{\sqrt{N_{\text{MC}}}}$. Reported Monte Carlo quantities are accompanied by 95% confidence intervals $\hat{H} \pm 1.96 \text{SE}(\hat{H})$. Common random numbers are used when comparing two policies under the same primitive calibration. This coupling reduces simulation noise in policy differences and makes the estimated comparative statics more precise [39]. Numerical convergence is assessed on three nested grids. If Q_h denotes a reported numerical statistic on the baseline grid and $Q_{h/2}$ the corresponding statistic after simultaneous refinement, we define

$$\mathcal{E}_{\text{grid}}(Q) := \frac{|Q_{h/2} - Q_h|}{1 + |Q_{h/2}|}. \quad (87)$$

The principal economic conclusions are reported only if $\mathcal{E}_{\text{grid}}(Q)$ is small relative to the comparative effect being interpreted. The same requirement is imposed on the equilibrium computation. Specifically, the final population flow must satisfy (69), while the equilibrium residual (70) and the HJB residual must remain below their prescribed tolerances on the refined grid. The experiments therefore distinguish four economically different sources of variation: catastrophe severity through α , catastrophe frequency through λ_J , demand-surge persistence through κ , and endogenous mitigation incentives through χ , γ_D , and γ_S [93]. This decomposition allows the numerical results to identify whether changes in equilibrium loss exposure arise from the primitive risk environment, the private physical-hedging response, or the mean field feedback generated by industry-wide vulnerability.

7.3 Numerical Results

We report only the numerical quantities directly related to the two principal mechanisms of the model: state-dependent physical hedging and the equilibrium effect of strategic interaction. The computations use the calibrated model of Table 1 together with the numerical scheme of Section 7.1. During implementation we found that the original benchmark pair $\chi = 0.20$, and $\lambda_J = 0.20$, generated the degenerate policy $u^* \equiv 0$ over the benchmark state region. Because this configuration is numerically valid but uninformative for comparative-statics analysis [38], the simulation benchmark used in this subsection adopts $\chi = 0.04$, $\lambda_J = 0.30$, $z_{\min} = 0.05$, and $\omega = 4$, while retaining the empirical values $\alpha = 1.71$, and $\kappa = 7.07$. All other coefficients are kept at their calibrated baseline values [102]. The resulting policy is strongly state dependent. At time zero, the optimal hedge over representative capitalization and demand-surge states is approximately

Table 2: Optimal physical hedge $u^*(0, x, y)$ under the simulation benchmark

	$y = -0.20$	$y = 0$	$y = 0.20$	$y = 0.40$
$x = 0.40$	0.30	0.30	0.40	0.50
$x = 1.00$	0.00	0.10	0.10	0.20
$x = 1.60$	0.00	0.00	0.00	0.00

Two features are immediate. First, holding capitalization fixed, optimal hedging is larger in elevated demand-surge states. Second, holding demand surge fixed, weakly capitalized insurers hedge more aggressively. The first effect is consistent with the local amplification mechanism in Corollary 56; the second arises

from the larger marginal cost of entering the weak-capitalization region [85]. For $x = 1$, the corresponding value function increases with the replacement-cost state:

Table 3: Value function at $t = 0$ and $x = 1$

y	Independent control	Mean field objective
-0.20	0.00685	0.01106
0	0.00716	0.01163
0.20	0.01006	0.01579
0.40	0.01979	0.02712

The monotone increase in the value function quantifies the higher expected social cost associated with elevated replacement costs. The discrete policy also yields a positive demand-surge response over the central state region. In particular, $u^*(0, 1, 0.40) - u^*(0, 1, 0) = 0.10$. Since the control grid is discrete, this difference should be interpreted as a policy-level comparative static rather than as a smooth elasticity. The mean field coupling raises the equilibrium value function because the representative insurer internalizes aggregate vulnerability through the coupling term F . Over the benchmark states, however, the equilibrium feedback remains close to the independent-insurer policy [100]. This indicates that, at the baseline interaction intensities, the principal equilibrium effect is a change in social cost rather than a large displacement of the pointwise physical-hedging rule.

The numerical solution therefore supports three conclusions. First, the optimal physical hedge responds jointly to capitalization and demand-surge exposure. Second, the value of insurance protection rises sharply when replacement costs are elevated. Third, the mean field interaction changes the equilibrium objective even when its effect on the local hedge is modest. These findings are consistent with the nonlocal control mechanism of Section 4 and the equilibrium feedback structure of Section 5. Complete implementation details, including the numerical primitives, boundary treatment, jump quadrature, monotonicity and positivity conditions, initialization, stopping rules, and reproducibility protocol, are collected in Appendix B.

8 Conclusion

This paper develops a nonlocal stochastic-control framework for catastrophe insurance when monetary catastrophe losses are exposed to stochastic replacement-cost variation. Catastrophe severity is generated by a marked jump process, replacement costs evolve through an exogenous mean-reverting factor, and physical hedging reduces the residual loss transmitted to insurer surplus. We establish the probabilistic foundations of the controlled state process, derive the dynamic programming principle, and characterize the value function as the unique viscosity solution of the associated nonlocal HJB equation. Extending the individual problem to interacting insurers yields a coupled HJB-Kolmogorov mean field system for which existence and, under the stated monotonicity conditions, uniqueness of equilibrium are obtained.

The analytical and numerical results show that elevated replacement costs magnify monetary catastrophe-loss exposure, while physical hedging attenuates the same channel. The numerical solution assigns greater hedging intensity to insurers facing high reconstruction costs and, particularly, to weakly capitalized insurers

[101]. Mean field interaction changes the equilibrium valuation of vulnerability even when its effect on the pointwise hedge is comparatively modest [71]. These findings concern the adverse coincidence of catastrophe losses with elevated reconstruction costs; the model does not imply that the catastrophe process itself generates the replacement-cost shock.

These distinctions also identify two related limitations of the present framework. First, the replacement-cost factor is exogenous: catastrophe arrivals do not generate jumps or impulses in Y , and the model contains neither common catastrophe noise nor a shared reconstruction-cost shock. Second, the mean field interaction is reduced form. Aggregate insurer behavior enters the individual problem through the cost couplings F and G , but it does not determine reconstruction prices, available physical capacity, catastrophe intensity, or the severity distribution [94]. Consequently, the mean field equilibrium should be interpreted as an equilibrium in strategic vulnerability costs rather than as an equilibrium of a reconstruction-capacity market. A structural extension could introduce aggregate rebuilding demand and a finite supply of contractors, materials, or pre-arranged capacity, together with a market-clearing reconstruction price increasing in aggregate demand. Embedding that price in the replacement-cost process or the catastrophe-loss amplitude would make reconstruction scarcity endogenous to the population distribution and would provide a direct mean field representation of post-catastrophe congestion. Such an extension would require a population-dependent controlled generator and potentially an additional aggregate state variable. Further extensions could incorporate common catastrophe shocks, heterogeneous geographic exposure, and interactions between physical hedging, reinsurance, and capital regulation.

Declarations.

Ethics approval and consent to participate.

Not applicable.

Consent for publication.

Not applicable.

Availability of data and material.

No data has been used.

Competing interests.

No potential conflict of interest was reported by the authors.

Funding.

Not applicable.

A Auxiliary Probabilistic Proofs

Proof of Lemma 1

Proof. The explicit representation of (3) is

$$Y_s = \vartheta + (y - \vartheta)e^{-\kappa(s-t)} + \sigma_I \int_t^s e^{-\kappa(s-r)} dW_r^I. \quad (88)$$

Define $M_s := \int_t^s e^{\kappa r} dW_r^I$. Then $\int_t^s e^{-\kappa(s-r)} dW_r^I = e^{-\kappa s} M_s$. Burkholder-Davis-Gundy (BDG) inequality yields

$$\mathbb{E} \left[\sup_{t \leq s \leq T} |M_s|^m \right] \leq C_m \mathbb{E} \left[\left(\int_t^T e^{2\kappa r} dr \right)^{m/2} \right] < \infty.$$

Using (88), the elementary inequality $(a + b + c)^m \leq C_m(a^m + b^m + c^m)$, and $e^{-\kappa(s-t)} \leq 1$ yields

$$\mathbb{E}_{t,y} \left[\sup_{t \leq s \leq T} |Y_s|^m \right] \leq C_m(1 + |y|^m),$$

which proves (12). For the exponential estimate, the Gaussian process $G_s := \sigma_I \int_t^s e^{-\kappa(s-r)} dW_r^I$ is centered and has variance bounded by

$$\sup_{t \leq s \leq T} \text{Var}(G_s) = \sup_{t \leq s \leq T} \sigma_I^2 \int_t^s e^{-2\kappa(s-r)} dr \leq \frac{\sigma_I^2}{2\kappa}.$$

Since G is a continuous Gaussian process on the compact interval $[t, T]$, Fernique's theorem implies the existence of $\eta > 0$ such that

$$\mathbb{E} \left[\exp \left(\eta \sup_{t \leq s \leq T} |G_s|^2 \right) \right] < \infty.$$

For arbitrary $r > 0$, Young's inequality implies $r \sup_s |G_s| \leq \eta \sup_s |G_s|^2 + \frac{r^2}{4\eta}$. Hence, $\mathbb{E}[\exp(r \sup_s |G_s|)] < \infty$. Combining this with (88) yields $\sup_{t \leq s \leq T} |Y_s| \leq |\vartheta| + |y - \vartheta| + \sup_{t \leq s \leq T} |G_s|$, and therefore,

$$\mathbb{E}_{t,y} \left[\sup_{t \leq s \leq T} e^{r|Y_s|} \right] \leq C_{r,T} \exp(C_{r,T}|y|).$$

Finally, because $I_s^\alpha = e^{\alpha Y_s}$ for every $\alpha \in \mathbb{R}$, the last assertion follows immediately. $\square$

Proof of Proposition 2

Proof. By Assumption 2.4, $|\ell(Y_s, z, u_s)|^p \leq C_p [1 + e^{pY_s} + |u_s|^p] \rho(z)^p$. Integrating with respect to $\mathbf{v}(dz) ds$ and applying Tonelli's theorem yields

$$\mathbb{E} \left[\int_t^T \int_E |\ell(Y_s, z, u_s)|^p \mathbf{v}(dz) ds \right] \leq C_p \left[\int_E \rho(z)^p \mathbf{v}(dz) \right] \cdot \mathbb{E} \left[\int_t^T (1 + e^{pY_s} + |u_s|^p) ds \right].$$

The first factor is finite by (7). The exponential moment in the second factor is finite by Lemma 1. Finally, $\mathbb{E} \left[\int_t^T |u_s|^p ds \right] < \infty$ for $p \leq p_*$ by Assumption 2.3 and Hölder's inequality on the finite interval $[t, T]$. This proves (15), and (14) is the special case $p = 2$. Standard stochastic integration with respect to compensated Poisson random measures then yields the claimed local-martingale and square-integrability properties [4, 97, 48]. $\square$

Proof of Theorem 3

Proof. The equation for Y is autonomous and has the unique strong solution (88). It therefore suffices to establish existence and pathwise uniqueness for X conditional on the already constructed process Y . Fix $n \in \mathbb{N}$ and define the stopping time $\tau_n := \inf \{s \geq t : |Y_s| + |X_s| \geq n\} \wedge T$. On $[t, \tau_n]$, the random coefficients $(x, s) \mapsto b(x, Y_s, u_s)$, and $(x, s) \mapsto \sigma_X(x, Y_s, u_s)$ are globally Lipschitz in x uniformly in (s, ω) , and the jump amplitude is Lipschitz in Y_s with $\mathbf{v}$ -square-integrable envelope ρ . Standard Picard iteration for stochastic differential equations with jumps therefore yields a unique strong solution up to τ_n ; cf. Situ [97] and Applebaum [4]. To extend the solution globally, it remains to prove the a priori estimate (16) independently of n . For $s \in [t, T]$,

$$X_{s \wedge \tau_n} = x + \int_t^{s \wedge \tau_n} b(X_{r-}, Y_{r-}, u_r) dr + \int_t^{s \wedge \tau_n} \sigma_X(X_{r-}, Y_{r-}, u_r) dW_r^X - \int_t^{s \wedge \tau_n} \int_E \ell(Y_{r-}, z, u_r) \tilde{N}(dr, dz).$$

For $p \geq 2$, by invoking $|a_1 + \dots + a_4|^p \leq C_p \sum_{j=1}^4 |a_j|^p$ we obtain

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq r \leq s} |X_{r \wedge \tau_n}|^p \right] &\leq C_p |x|^p + C_p \mathbb{E} \left[\int_t^{s \wedge \tau_n} |b(X_{r-}, Y_{r-}, u_r)| dr \right]^p \\ &\quad + C_p \cdot \mathbb{E} \left[\sup_{t \leq v \leq s} \left| \int_t^{v \wedge \tau_n} \sigma_X(X_{r-}, Y_{r-}, u_r) dW_r^X \right|^p \right] \\ &\quad + C_p \cdot \mathbb{E} \left[\sup_{t \leq v \leq s} \left| \int_t^{v \wedge \tau_n} \int_E \ell(Y_{r-}, z, u_r) \tilde{N}(dr, dz) \right|^p \right]. \end{aligned} \quad (89)$$

Hölder's inequality and (6) yield

$$\mathbb{E} \left[\int_t^{s \wedge \tau_n} |b_r| dr \right]^p \leq C_p \cdot \int_t^s \mathbb{E} \left[1 + \sup_{t \leq v \leq r} |X_{v \wedge \tau_n}|^p + |Y_r|^p + |u_r|^p \right] dr. \quad (90)$$

The Brownian BDG inequality gives

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq v \leq s} \left| \int_t^{v \wedge \tau_n} \sigma_X dW_r^X \right|^p \right] &\leq C_p \mathbb{E} \left[\int_t^{s \wedge \tau_n} |\sigma_X|^2 dr \right]^{p/2} \\ &\leq C_p \int_t^s \mathbb{E} \left[1 + \sup_{t \leq v \leq r} |X_{v \wedge \tau_n}|^p + |Y_r|^p + |u_r|^p \right] dr. \end{aligned} \quad (91)$$

For the compensated jump martingale, the Bichteler–Jacod inequality yields

$$\begin{aligned} \mathbb{E} \left[\sup_{t \leq v \leq s} \left| \int_t^{v \wedge \tau_n} \int_E \ell(Y_{r-}, z, u_r) \tilde{N}(dr, dz) \right|^p \right] \\ \leq C_p \mathbb{E} \left(\int_t^{s \wedge \tau_n} \int_E |\ell(Y_{r-}, z, u_r)|^2 \nu(dz) dr \right)^{p/2} + C_p \cdot \mathbb{E} \left[\int_t^{s \wedge \tau_n} \int_E |\ell(Y_{r-}, z, u_r)|^p \nu(dz) dr \right]. \end{aligned} \quad (92)$$

Using (7), Hölder’s inequality, and Lemma 1, the right-hand side of (92) is bounded by

$$C_p \int_t^s \left[1 + \mathbb{E} e^{p|Y_r|} + \mathbb{E} |u_r|^p \right] dr. \quad (93)$$

Substituting these bounds into (89) gives

$$\mathbb{E} \left[\sup_{t \leq r \leq s} |X_{r \wedge \tau_n}|^p \right] \leq C_p \left[1 + |x|^p + |y|^p + e^{C_p|y|} + \mathbb{E} \int_t^T |u_r|^p dr \right] + C_p \int_t^s \mathbb{E} \left[\sup_{t \leq v \leq r} |X_{v \wedge \tau_n}|^p \right] dr. \quad (94)$$

Gronwall’s inequality therefore yields

$$\sup_{n \geq 1} \mathbb{E} \left[\sup_{t \leq s \leq T} |X_{s \wedge \tau_n}|^p \right] \leq C_p \left[1 + |x|^p + |y|^p + e^{C_p|y|} + \mathbb{E} \int_t^T |u_s|^p ds \right]. \quad (95)$$

By Markov’s inequality,

$$\mathbb{P}(\tau_n < T) \leq \mathbb{P} \left(\sup_{t \leq s \leq T} |X_{s \wedge \tau_n}| \geq n/2 \right) + \mathbb{P} \left(\sup_{t \leq s \leq T} |Y_s| \geq n/2 \right),$$

and the right-hand side converges to zero as $n \rightarrow \infty$ by (95) and Lemma 1. Hence, $\tau_n \uparrow T$ almost surely and the local solution extends to all of $[t, T]$. Moreover, Fatou’s lemma applied to (95) gives (16). For pathwise uniqueness, let X and $\hat{X}$ be two solutions driven by the same (W, N) with the same initial state and control. Since the jump amplitude in (4) does not depend on X , the jump terms cancel in $X - \hat{X}$. Using (5), the BDG inequality, and Gronwall’s inequality gives $\mathbb{E} \left[\sup_{t \leq s \leq T} |X_s - \hat{X}_s|^2 \right] = 0$. Therefore, X and $\hat{X}$ are indistinguishable. Strong existence and pathwise uniqueness follow. $\square$

Proof of Proposition 5

Proof. From the explicit representation (88), $Y_s^{t,y} - Y_s^{t,y'} = (y - y')e^{-\kappa(s-t)}$, which proves (19). Define $\Delta X_s := X_s^{t,x,y;u} - X_s^{t,x',y';u}$, and $\Delta Y_s := Y_s^{t,y} - Y_s^{t,y'}$. Then

$$\Delta X_s = x - x' + \int_t^s \Delta b_r dr + \int_t^s \Delta \sigma_r dW_r^X - \int_t^s \int_E \Delta \ell_r(z) \tilde{N}(dr, dz),$$

where $|\Delta b_r| + |\Delta \sigma_r| \leq L(|\Delta X_r| + |\Delta Y_r|)$ and, by (8), $|\Delta \ell_r(z)| \leq L|\Delta Y_r|\rho(z)$. Applying Hölder's inequality, the Brownian BDG inequality, and the Bichteler-Jacod inequality as in the proof of Theorem 3, we obtain

$$\mathbb{E} \left[\sup_{t \leq v \leq s} |\Delta X_v|^p \right] \leq C_p |x - x'|^p + C_p \int_t^s \mathbb{E} \left[\sup_{t \leq q \leq r} |\Delta X_q|^p \right] dr + C_p \int_t^s \mathbb{E} [|\Delta Y_r|^p] dr.$$

Using (19), $\int_t^s \mathbb{E} [|\Delta Y_r|^p] dr \leq T|y - y'|^p$. Gronwall's inequality therefore gives the stronger bound

$$\mathbb{E} \left[\sup_{t \leq s \leq T} |\Delta X_s|^p \right] \leq C_p (|x - x'|^p + |y - y'|^p).$$

This estimate already implies (18); the factor $\Xi_p(y, y') \geq 1$ is retained because later extensions in which ℓ is locally Lipschitz in the level $I = e^Y$ naturally produce this exponential weight. The stated form is therefore stable under the economically important multiplicative specification (9). $\square$

Proof of Lemma 6

Proof. Since the construction-cost factor is uncontrolled, both systems have the same process $Y^{t,y}$. Set $\Delta X_s := X_s^{t,x,y;u} - X_s^{t,x,y;v}$. Subtracting the two surplus equations gives

$$\Delta X_s = \int_t^s \Delta b_r dr + \int_t^s \Delta \sigma_r dW_r^X - \int_t^s \int_E \Delta \ell_r(z) \tilde{N}(dr, dz),$$

where Assumptions 2.4 and 2.7 imply $|\Delta b_r| + |\Delta \sigma_r| \leq C(|\Delta X_r| + |u_r - v_r|)$, and $|\Delta \ell_r(z)| \leq L_u |u_r - v_r| \rho(z)$. Hölder's inequality and the Brownian BDG inequality therefore yield

$$\mathbb{E} \sup_{t \leq q \leq s} \left| \int_t^q \Delta b_r dr \right|^p + \mathbb{E} \sup_{t \leq q \leq s} \left| \int_t^q \Delta \sigma_r dW_r^X \right|^p \leq C_p \int_t^s \mathbb{E} \left[\sup_{t \leq a \leq r} |\Delta X_a|^p + |u_r - v_r|^p \right] dr.$$

For the jump martingale, the Bichteler-Jacod inequality and Assumption 2.4 yield

$$\begin{aligned} & \mathbb{E} \left[\sup_{t \leq q \leq s} \left| \int_t^q \int_E \Delta \ell_r(z) \tilde{N}(dr, dz) \right|^p \right] \\ & \leq C_p \cdot \mathbb{E} \left(\int_t^s \int_E |\Delta \ell_r(z)|^2 \mathbf{v}(dz) dr \right)^{p/2} + C_p \cdot \mathbb{E} \left[\int_t^s \int_E |\Delta \ell_r(z)|^p \mathbf{v}(dz) dr \right] \\ & \leq C_p \cdot \mathbb{E} \left[\int_t^s |u_r - v_r|^p dr \right], \end{aligned}$$

where finiteness of the ρ^2 - and ρ^p -moments follows from Assumption 2.4. Combining the preceding estimates,

$$\mathbb{E} \left[\sup_{t \leq q \leq s} |\Delta X_q|^p \right] \leq C_p \cdot \int_t^s \mathbb{E} \left[\sup_{t \leq a \leq r} |\Delta X_a|^p \right] dr + C_p \cdot \mathbb{E} \left[\int_t^s |u_r - v_r|^p dr \right].$$

Gronwall's inequality proves (20). The final assertion is immediate from the definition of the $\mathcal{S}^p$ norm. $\square$

Proof of Proposition 7

Proof. Lemma 6 guarantees convergence of X^{u^n} to X^u in $\mathcal{S}^{p*}$, hence in probability uniformly on $[t, T]$. Since Y is common to all controls, the vector $(s, X_s^{u^n}, Y_s, u_s^n)$ converges to (s, X_s^u, Y_s, u_s) in measure on the product space $([t, T] \times \Omega, ds \otimes d\mathbb{P})$. After passage to a subsequence, convergence holds $ds \otimes d\mathbb{P}$ -a.e. Continuity of f then gives pointwise convergence of the running costs along that subsequence. It remains to justify passage under the expectation. Choose $\varepsilon > 0$ such that $q(1 + \varepsilon) < p_*$, which is possible by Assumption 2.6. The growth bound on f , Theorem 3, Lemma 1, and the assumed uniform p_* -bound on the controls imply $\sup_n \mathbb{E} \left[\int_t^T |f(s, X_s^{u^n}, Y_s, u_s^n)|^{1+\varepsilon} ds \right] < \infty$. Thus the running-cost family is uniformly integrable. The same argument, without the control term, shows uniform integrability of $\{g(X_T^{u^n}, Y_T)\}_{n \geq 1}$. Vitali's theorem yields convergence of both the running and terminal terms in L^1 . Since every subsequence admits a further subsequence with the same limit, the full sequence converges, proving the claim. $\square$

Proof of Proposition 9

Proof. Integrate the original state equations from τ to s . For the cost factor,

$$Y_s = Y_\tau + \int_\tau^s \kappa(\vartheta - Y_r) dr + \sigma_I(W_s^I - W_\tau^I).$$

For the surplus,

$$X_s = X_\tau + \int_\tau^s b(X_{r-}, Y_{r-}, u_r) dr + \int_\tau^s \sigma_X(X_{r-}, Y_{r-}, u_r) dW_r^X - \int_\tau^s \int_E \ell(Y_{r-}, z, u_r) \tilde{N}(dr, dz).$$

After the time change $r = \tau + a$, these are exactly the integral equations defining the system restarted from (X_τ, Y_τ) and driven by (W^τ, N^τ) . Theorem 3 gives pathwise uniqueness. Hence the original post- τ trajectory

and the restarted trajectory are indistinguishable, proving (21). If the original control is replaced by $u \otimes_{\tau} v$, its restriction to $[\tau, T]$ equals v , and the same argument applies. $\square$

Proof of Proposition 10

Proof. First suppose that τ takes finitely many deterministic values and that Φ is a bounded cylinder functional. On each event $\{\tau = t_j\}$, Proposition 9 identifies the future state with the solution driven by increments of (W, N) after t_j . These increments are independent of $\mathcal{F}_{t_j}$ and have the same law as fresh Brownian and Poisson noises. Conditioning therefore gives (22). For a general stopping time, choose stopping times τ_n taking values in the dyadic grid such that $\tau_n \downarrow \tau$. Càdlàg paths, the state stability estimates, and right-continuity of the filtration permit passage from τ_n to τ for bounded continuous cylinder functionals. A monotone-class argument extends the identity to bounded Borel Φ . Finally, truncation and uniform integrability extend the result to every integrable Φ covered by the statement. This is the controlled jump-diffusion version of the standard restart argument underlying the DPP [30, 51], and the measurable-selection formulations in El Karoui and Tan [24] and Žitković [106]. $\square$

Proof of Theorem 11

Proof. Substituting $u_s = \alpha(s, X_{s-}, Y_{s-})$ into the coefficients produces a time-inhomogeneous jump SDE whose coefficients are Borel in time and globally Lipschitz in the state, with linear growth. Indeed, Assumptions 2.4, 2.7, and 2.8 imply

$$|b(x, y, \alpha(s, x, y)) - b(x', y', \alpha(s, x', y'))| \leq C_{\alpha}(|x - x'| + |y - y'|),$$

and the same estimate holds for σ_X ; the jump coefficient satisfies the corresponding $L^2(v) \cap L^p(v)$ estimate. Hence Theorem 3 applies to the feedback system, and pathwise uniqueness holds from every deterministic initial state. The flow property of Proposition 9, now with a feedback control depending only on the current time and state, implies the Markov property at deterministic times. At a stopping time τ , the continuation rule is $s \mapsto \alpha(s, X_{s-}, Y_{s-})$, which depends on the past only through the restarted state. Proposition 10, together with pathwise uniqueness, therefore yields $\mathbb{E}_{t,x,y}[\varphi(Z_{\tau+r}^{\alpha}) \mid \mathcal{F}_{\tau}] = P_{\tau, \tau+r}^{\alpha} \varphi(Z_{\tau}^{\alpha})$ a.s. for bounded Borel φ , proving the strong Markov property. For the Feller property, let $(x_n, y_n) \rightarrow (x, y)$. Repeating the proof of Proposition 5 with the feedback controls generated by the two state trajectories and using the Lipschitz property of α gives, for every $p \in [2, p_{\star}]$,

$$\mathbb{E} \left[\sup_{t \leq r \leq s} |Z_r^{t, x_n, y_n; \alpha} - Z_r^{t, x, y; \alpha}|^p \right] \longrightarrow 0.$$

Hence $Z_s^{t, x_n, y_n; \alpha} \rightarrow Z_s^{t, x, y; \alpha}$ in probability. If $\varphi \in C_b(\mathbb{R}^2)$, boundedness and convergence in probability imply $P_{t,s}^{\alpha} \varphi(x_n, y_n) \rightarrow P_{t,s}^{\alpha} \varphi(x, y)$, so $P_{t,s}^{\alpha} \varphi$ is continuous. Boundedness is immediate from $\|P_{t,s}^{\alpha} \varphi\|_{\infty} \leq \|\varphi\|_{\infty}$. $\square$

Proof of Proposition 13

Proof. Since $q < p_{\star}$, choose $\eta > 0$ so small that $q(1 + \eta) < p_{\star}$. Apply Theorem 3 with $p = q(1 + \eta)$. Because K is compact, $|x|$, $|y|$, and $\exp(C|y|)$ are uniformly bounded on K . Moreover, by Hölder's inequality on the

finite measure interval $[t, T]$,

$$\mathbb{E} \left[\int_t^T |u_s|^{q(1+\eta)} ds \right] \leq (T-t)^{1-q(1+\eta)/p_*} \left(\mathbb{E} \int_t^T |u_s|^{p_*} ds \right)^{q(1+\eta)/p_*},$$

which is uniformly bounded over $\mathcal{U}_t(R)$. Lemma 1 supplies the exponential moment of Y . This proves (23).

By Assumption 2.6,

$$|g(X_T, Y_T)|^{1+\eta} \leq C_\eta \left(1 + |X_T|^{q(1+\eta)} + e^{q(1+\eta)|Y_T|} \right),$$

whose expectations are uniformly bounded by (23). Hence the terminal-cost family is uniformly integrable by the de la Vallée-Poussin criterion. Similarly, Jensen's inequality yields

$$\left(\int_t^T |f_s| ds \right)^{1+\eta} \leq (T-t)^\eta \int_t^T |f_s|^{1+\eta} ds,$$

and the growth condition on f together with (23) yields a uniform bound on the expectation of the right-hand side. The running-cost family is therefore uniformly integrable as well. $\square$

Proof of Proposition 14

Proof. Since U is compact metric, the relaxed-control space equipped with the stable topology is a standard Borel space. Hence the canonical product space Ω° is standard Borel, and $\mathfrak{P}(\Omega^\circ)$ endowed with the weak topology is likewise standard Borel. Regular conditional probabilities and analytic-set arguments may therefore be used on the canonical space. Let $Z = (X, Y)$ denote the canonical state coordinate and let q denote the relaxed-control coordinate. For $\varphi \in C_c^2(\mathbb{R}^2)$, define

$$\begin{aligned} \mathcal{A}^a \varphi(x, y) &= b(x, y, a) \varphi_x(x, y) + \kappa(\vartheta - y) \varphi_y(x, y) + \frac{1}{2} \sigma_X^2(x, y, a) \varphi_{xx}(x, y) + \frac{1}{2} \sigma_I^2 \varphi_{yy}(x, y) \\ &\quad + \int_E \left[\varphi(x - \ell(y, z, a), y) - \varphi(x, y) + \ell(y, z, a) \varphi_x(x, y) \right] \nu(dz). \end{aligned}$$

For a relaxed control q , put $\mathcal{A}^{q_s} \varphi(x, y) = \int_U \mathcal{A}^a \varphi(x, y) q_s(da)$. The canonical martingale problem requires

$$M_s^\varphi := \varphi(Z_s) - \varphi(Z_t) - \int_t^s \mathcal{A}^{q_r} \varphi(Z_{r-}) dr$$

to be a local martingale, together with the initial-state and admissibility conditions. We first prove analyticity. Choose a countable family $\{\varphi_k\}_{k \geq 1} \subset C_c^2(\mathbb{R}^2)$ dense in the local C^2 topology, rational times $r < s$, a countable generating algebra of bounded cylinder functions measurable at time r , and a sequence of localization stopping times. The martingale problem is equivalent to the resulting countable collection of identities

$$E^P \left[(M_{s \wedge \tau_n}^{\varphi_k} - M_{r \wedge \tau_n}^{\varphi_k}) H \right] = 0.$$

Assumption 2.9(ii)-(iii), together with the growth conditions of Assumptions 2.4 and 2.6, implies that the canonical integrands are Borel measurable. The initial-state constraint and the required moment conditions are likewise Borel after truncation. Consequently the set of probability measures satisfying the localized martingale identities and admissibility conditions is Borel in the canonical probability space. Its projection onto (t, x, y, P) is therefore analytic. This proves (a). For (b), fix $P \in \mathfrak{P}(t, x, y)$ and an $[t, T]$ -valued stopping time τ . For every localized martingale, optional sampling gives

$$E^P [M_{s \wedge \tau_n}^{\phi_k} - M_{\tau \wedge \tau_n}^{\phi_k} \mid \mathcal{F}_\tau] = 0.$$

Disintegrating P with respect to $\mathcal{F}_\tau$ shows that, for P -a.e. ω , the conditional law satisfies the same localized martingale identities after $\tau(\omega)$ and starts from $(X_\tau(\omega), Y_\tau(\omega))$. The conditional admissibility requirement follows from conditional Tonelli's theorem. Hence, $P^{\tau, \omega} \in \mathfrak{P}(\tau(\omega), X_\tau(\omega), Y_\tau(\omega))$ for P -a.e. ω . For (c), let Q_ω be a universally measurable admissible continuation kernel. Under $P \otimes_\tau Q$, every localized martingale identity holds before τ because it holds under P and after τ because it holds under Q_ω for P -a.e. ω . Splitting an increment at τ and applying the tower property therefore yields the martingale identity on the full time interval. The moment admissibility condition is preserved by the corresponding decomposition at τ . Therefore, $P \otimes_\tau Q \in \mathfrak{P}(t, x, y)$. This proves (b)-(c). The argument is the model-specific verification of the abstract conditioning and concatenation structure used in El Karoui and Tan [24], Žitković [106], and Fayvisovich and Žitković [28]. $\square$

A.1 Proof of Proposition 15

Proof. Since, U is compact metric, the space of relaxed control measures with time marginal equal to Lebesgue measure is compact under the stable topology. The chattering lemma yields predictable U -valued controls u^n such that, for every bounded measurable $\psi : [t, T] \times \Omega \times U \rightarrow \mathbb{R}$ that is continuous in the control variable,

$$\int_t^T \psi(s, \omega, u_s^n) ds \longrightarrow \int_t^T \int_U \psi(s, \omega, a) q_s(da) ds \quad (96)$$

in probability, after passage to a subsequence when necessary. We apply this approximation simultaneously to the controlled characteristics of the martingale problem. By Assumption 2.9(ii)-(iii), the maps $a \mapsto b(x, y, a)$, $a \mapsto \sigma_X^2(x, y, a)$, and

$$a \mapsto \int_E \left[\varphi(x - \ell(y, z, a), y) - \varphi(x, y) + \ell(y, z, a) \varphi_x(x, y) \right] \nu(dz)$$

are continuous for every $\varphi \in C_c^2(\mathbb{R}^2)$. For the jump term, continuity follows from Taylor's theorem on the small-jump region, Assumption 2.9(iii), and dominated convergence on the complementary region. Hence (96) implies convergence of the integrated characteristics associated with every test function in the countable martingale-problem core. The moment estimates of Theorem 3 are uniform over strict controls because U is compact. Together with the increment estimates used in the proof of Proposition 13, they imply tightness of $\{(X^n, Y^n)\}_{n \geq 1}$ in $D([t, T]; \mathbb{R}^2)$. “The compactness of U , the continuity hypotheses in Assumption 2.9(ii)–(iii), and the stable convergence of the occupation measures place the present controlled martingale problem within the classical compactification framework for relaxed stochastic controls. The chattering

approximation theorem of El Karoui et al. [23] therefore implies that the strict controlled laws associated with (u^n) converge, along the approximating sequence, to the relaxed controlled law P in the canonical weak topology. In particular,

$$(X^n, Y^n) \Longrightarrow (X, Y) \quad \text{in } D([t, T]; \mathbb{R}^2). \quad (97)$$

We stress that this is a weak convergence statement for the martingale formulation. It does not identify the relaxed dynamics with the strict SDE of Theorem 3; rather, the relaxed generator averages the controlled local and jump characteristics with respect to q_s . This is precisely the role of the compactification procedure.

It remains to prove convergence of the objective functionals. Assumption 2.9(ii), stable convergence of the controls, and (97) imply convergence in probability of the truncated running costs. The uniform moment estimates established in Proposition 13, together with the polynomial growth bound of Assumption 2.6, give uniform integrability. Vitali's theorem therefore yields

$$E \int_t^T e^{-\delta(s-t)} f(s, X_s^n, Y_s^n, u_s^n) ds \rightarrow E^P \int_t^T e^{-\delta(s-t)} \int_U f(s, X_s, Y_s, a) q_s(da) ds.$$

The same weak convergence, continuity of g , and uniform integrability imply $E[g(X_T^n, Y_T^n)] \rightarrow E^P[g(X_T, Y_T)]$. Consequently,

$$J(t, x, y; u^n) \rightarrow J(t, x, y; P). \quad (98)$$

□

A.2 Proof of Theorem 17

Proof. Set $E := [0, T] \times \mathbb{R}^2$, $P := \mathfrak{P}(\Omega^\circ)$, and $\mathcal{G} := \text{Gr}(\mathfrak{P}) \subset E \times P$. Write $\xi = (t, x, y) \in E$. By Proposition 14, $\mathcal{G} \in \mathcal{A}(E \times P)$, where $\mathcal{A}$ denotes the class of analytic sets. For $(\xi, P) \in \mathcal{G}$, define $\Gamma(\xi, P) := J(t, x, y; P)$, i.e., $\Gamma(\xi, P) = E^P[\int_t^T e^{-\delta(s-t)} \int_U f(s, X_s, Y_s, a) q_s(da) ds + e^{-\delta(T-t)} g(X_T, Y_T)]$. For $N \in \mathbb{N}$, let $f_N := (-N) \vee (f \wedge N)$ and $g_N := (-N) \vee (g \wedge N)$, and define Γ_N by replacing (f, g) with (f_N, g_N) in Γ . The canonical evaluation maps and the relaxed-control coordinate are Borel, hence Γ_N is Borel on $\mathcal{G}$. Assumption 2.6 and Proposition 13 imply $\Gamma_N(\xi, P) \rightarrow \Gamma(\xi, P)$ on $\mathcal{G}$, with uniformly integrable truncation error on bounded-moment subsets. Therefore $\Gamma \in \text{LSA}(\mathcal{G})$.

Extend Γ to $E \times P$ by $\bar{\Gamma}(\xi, P) := \Gamma(\xi, P)$ on $\mathcal{G}$ and $\bar{\Gamma}(\xi, P) := +\infty$ otherwise. Since $\mathcal{G}$ is analytic, $\bar{\Gamma} \in \text{LSA}(E \times P)$. Consequently, $V^{\text{rel}}(\xi) = \inf_{P \in P} \bar{\Gamma}(\xi, P) = \inf_{P \in \mathfrak{P}(\xi)} \Gamma(\xi, P)$. For every $c \in \mathbb{R}$, $\{\xi : V^{\text{rel}}(\xi) < c\} = \text{proj}_E\{(\xi, P) : \bar{\Gamma}(\xi, P) < c\}$. The set on the right is analytic, and analytic sets are stable under projection; hence $\{\xi : V^{\text{rel}}(\xi) < c\} \in \mathcal{A}(E)$. Thus $V^{\text{rel}} \in \text{LSA}(E)$. Fix $\varepsilon > 0$ and set $\mathcal{G}_\varepsilon := \{(\xi, P) \in \mathcal{G} : \Gamma(\xi, P) \leq V^{\text{rel}}(\xi) + \varepsilon\}$. Using rational approximation, $\mathcal{G}_\varepsilon = \bigcap_{n \geq 1} \bigcup_{r \in \mathbb{Q}} (\{\xi : V^{\text{rel}}(\xi) < r\} \times P) \cap \{(\xi, P) \in \mathcal{G} : \Gamma(\xi, P) < r + \varepsilon + n^{-1}\}$. Hence $\mathcal{G}_\varepsilon \in \mathcal{A}(E \times P)$. Let $D_V := \{\xi \in E : V^{\text{rel}}(\xi) < \infty\}$. For every $\xi \in D_V$, the section $(\mathcal{G}_\varepsilon)_\xi := \{P \in P : (\xi, P) \in \mathcal{G}_\varepsilon\}$ is nonempty by definition of the infimum. The Jankov-von Neumann selection theorem therefore yields a universally measurable selector $\xi \mapsto P_\xi^\varepsilon$ satisfying $(\xi, P_\xi^\varepsilon) \in \mathcal{G}_\varepsilon$ for all $\xi \in D_V$. Hence $P_\xi^\varepsilon \in \mathfrak{P}(\xi)$ and $\Gamma(\xi, P_\xi^\varepsilon) \leq V^{\text{rel}}(\xi) + \varepsilon$. Returning to $\xi = (t, x, y)$ gives $J(t, x, y; P_{t,x,y}^\varepsilon) \leq V^{\text{rel}}(t, x, y) + \varepsilon$, which is (27). □

Proof of Theorem 23

Proof. By Proposition 16 the strict and relaxed values coincide, so it is enough to prove the identity in the canonical formulation, where measurable continuation is available. Fix $P \in \mathfrak{P}(t, x, y)$. By Lemma 22 and the definition of the value function, the conditional continuation cost at τ is bounded below by $V(\tau, X_\tau, Y_\tau)$. Hence,

$$J(t, x, y; P) \geq E^P \left[\int_t^\tau e^{-\delta(s-t)} f_s ds + e^{-\delta(\tau-t)} V(\tau, X_\tau, Y_\tau) \right].$$

Taking the infimum over P yields the “ $\geq$ ” inequality in (39). This is the inequality already anticipated by the pre-DPP decomposition of Proposition 19. For the reverse inequality, fix $\varepsilon > 0$. The universally measurable selection result of Theorem 17 provides a continuation kernel $(s, \xi, \eta) \mapsto Q_{s, \xi, \eta}^\varepsilon \in \mathfrak{P}(s, \xi, \eta)$ such that $J(s, \xi, \eta; Q_{s, \xi, \eta}^\varepsilon) \leq V(s, \xi, \eta) + \varepsilon$. Let P be any admissible law up to τ . By Proposition 14, the concatenation $P \otimes_\tau Q_{\tau, X_\tau, Y_\tau}^\varepsilon$ is admissible from (t, x, y) . Applying Lemma 22 to this concatenated law gives

$$V(t, x, y) \leq E^P \left[\int_t^\tau e^{-\delta(s-t)} f_s ds + e^{-\delta(\tau-t)} (V(\tau, X_\tau, Y_\tau) + \varepsilon) \right].$$

Since $e^{-\delta(\tau-t)} \leq e^{|\delta|T}$, the error is bounded by $C\varepsilon$. Taking the infimum over P and then letting $\varepsilon \downarrow 0$ proves the opposite inequality. The strict form follows again from Proposition 16. $\square$

Proof of Proposition 25

Proof. Membership in $\mathcal{G}_q$ follows from Corollary 21. Fix a compact $K \subset \mathbb{R}^2$. Coercivity of f , together with the growth bound for V , permits restriction, up to an arbitrarily small error, to controls whose $p_\star$ -moments are uniformly bounded on K . For such controls, Proposition 7 gives local uniform stability of the state with respect to the initial condition. Local uniform continuity of f and g , followed by truncation outside a large compact set and the uniform-integrability estimate of Proposition 13, therefore yields $\sup_u |J(t, x, y; u) - J(t, x', y'; u)| \rightarrow 0$ as $(x', y') \rightarrow (x, y)$, locally uniformly in $(x, y) \in K$. Taking infima preserves this modulus. For time continuity, apply Corollary 24 over an interval of length h . The running-cost contribution tends to zero uniformly on compact sets by uniform integrability, while the stochastic continuity of (X, Y) and the already established spatial continuity of V imply $E[V(t+h, X_{t+h}, Y_{t+h})] \rightarrow V(t, x, y)$. This proves continuity for $t < T$. Finally, use the DPP with $s = T$. The running cost over $[t, T]$ vanishes as $t \uparrow T$, while $(X_T, Y_T) \rightarrow (x, y)$ in probability uniformly over the truncated near-optimal control family. Uniform integrability and continuity of g then give the terminal limit. $\square$

Proof of Theorem 27

Proof. We prove the two inequalities separately.

Subsolution property. Let $\phi \in C^{1,2,2}$ and suppose that $V - \phi$ has a strict local maximum equal to zero at (t_0, x_0, y_0) , with $t_0 < T$. Fix $a \in U$ and choose $R, h > 0$ small enough that $V \leq \phi$ on the relevant local cylinder.

Let τ_R be as in Lemma 26, under the constant control a . The DPP gives

$$V(t_0, x_0, y_0) \leq E \left[\int_{t_0}^{\tau_R} e^{-\delta(s-t_0)} f(s, X_s, Y_s, a) ds + e^{-\delta(\tau_R-t_0)} V(\tau_R, X_{\tau_R}, Y_{\tau_R}) \right].$$

For jumps remaining inside the local cylinder we may replace V by ϕ . For jumps leaving it, the large-jump contribution is retained on V ; this is precisely the split appearing in Definition 1. Combining the preceding inequality with Lemma 26, dividing by $E[\tau_R - t_0]$, and sending $h \downarrow 0$ and then $R \downarrow 0$ yields

$$\begin{aligned} 0 \leq & f(t_0, x_0, y_0, a) + \phi_t(t_0, x_0, y_0) + b(x_0, y_0, a)\phi_x(t_0, x_0, y_0) \\ & + \kappa(\vartheta - y_0)\phi_y(t_0, x_0, y_0) + \frac{1}{2}\sigma_X^2(x_0, y_0, a)\phi_{xx}(t_0, x_0, y_0) + \frac{1}{2}\sigma_I^2\phi_{yy}(t_0, x_0, y_0) \\ & - \delta V(t_0, x_0, y_0) + I_\varepsilon^a[\phi](t_0, x_0, y_0) + I^{a, \varepsilon}[V, \phi](t_0, x_0, y_0). \end{aligned}$$

Since $a \in U$ was arbitrary, taking the infimum gives the subsolution inequality in the sign convention of (37).

Supersolution property. Suppose now that $V - \phi$ has a strict local minimum equal to zero at (t_0, x_0, y_0) . If the supersolution inequality failed, there would exist $\eta > 0$ and a sufficiently small neighborhood of (t_0, x_0, y_0) such that

$$\inf_{u \in U} \{f + \phi_t + \mathcal{A}^u \phi - \delta \phi\} \geq \eta,$$

with the nonlocal term interpreted through the small/large jump splitting. By coercivity, controls outside a sufficiently large compact subset of U cannot decrease the Hamiltonian on this neighborhood. Choose an ε_h -optimal control in the DPP, with $\varepsilon_h = o(h)$, and stop at the first exit from the neighborhood or at $t_0 + h$. Since $V \geq \phi$ locally, the DPP and the localized Dynkin formula imply $0 \geq \eta E[\tau_R - t_0] - o(h)$, where the exit contribution is controlled by the growth estimate and uniform integrability. The state moment bounds imply $E[\tau_R - t_0] = h + o(h)$ as $h \downarrow 0$. Dividing by h gives a contradiction. The terminal inequalities follow from Proposition 25. Hence V is both a subsolution and a supersolution. $\square$

Proof of Lemma 30

Proof. Because $\bar{q} > q$ and the exponential factor in Γ dominates the y -growth defining G_q , the penalized function tends to $-\infty$ as either spatial variable escapes to infinity. Therefore a maximizer exists. The standard doubling-variable argument gives boundedness, for fixed η , of the maximizing sequence and $|x_\varepsilon - x'_\varepsilon| + |y_\varepsilon - y'_\varepsilon| + |t_\varepsilon - s_\varepsilon| \rightarrow 0$. The sharper quotient convergence follows from the usual comparison of the maximum value with its diagonal competitor. For the jump terms, use maximality of $\Phi_{\varepsilon, \eta}$ after translating the two surplus variables by $-\ell(y_\varepsilon, z, u)$ and $-\ell(y'_\varepsilon, z, u)$. The quadratic penalization produces $\frac{|\ell(y_\varepsilon, z, u) - \ell(y'_\varepsilon, z, u)|^2}{\varepsilon}$, which is bounded by $C \frac{|y_\varepsilon - y'_\varepsilon|^2}{\varepsilon} \rho(z)^2$. Its v -integral tends to zero by Assumption 4.1(ii) and the doubling estimate. The Lyapunov penalization contributes at most $C\eta(\rho(z)^2 + \rho(z)^{\bar{q}})$, whose integral is finite by Assumption 4.1(iv). Dominated convergence therefore yields $I^u[w] - I^u[\underline{w}] \leq o_\varepsilon(1) + C\eta$, uniformly over controls in the compact set furnished by Lemma 29. This is the required estimate. The argument is the finite-dimensional specialization of the nonlocal Jensen-Ishii method of Barles and Imbert [8]; related semicontinuous maximum principles for integro-PDEs are developed in Jakobsen and Karlsen [36]. $\square$

Proof of Proposition 31

Proof. The derivatives of Γ satisfy $|D\Gamma| \leq C\Gamma$, and $|D^2\Gamma| \leq C\Gamma$. For the jump part, Taylor's theorem on $\{\rho \leq 1\}$ and the polynomial inequality $|x - r|^{\bar{q}} \leq C_{\bar{q}}(1 + |x|^{\bar{q}} + |r|^{\bar{q}})$ on $\{\rho > 1\}$ imply $|I^u\Gamma(x, y)| \leq C\Gamma(x, y) + C(1 + |u|^{\bar{q}})$ on compact state sets, with constants controlled by Assumption 4.1(iv). The coercivity reduction of Lemma 29 permits restriction to bounded controls when testing the equation locally. Hence, $\sup_u |\mathcal{A}^u\Gamma| \leq C\Gamma$ on the relevant minimizing set. The time derivative of the perturbation contributes $-\partial_t (\eta e^{\Lambda(T-t)}\Gamma) = \Lambda\eta e^{\Lambda(T-t)}\Gamma$. Choosing Λ larger than the constants arising from the local and nonlocal spatial terms, as well as the discount term, leaves a strict positive remainder of order $\eta\Gamma$. Stability of viscosity supersolutions under addition of a smooth strict barrier completes the proof. $\square$

Proof of Theorem 32

Proof. Assume, to the contrary, that $M := \sup_{[0, T] \times \mathbb{R}^2} (\bar{v} - \underline{v}) > 0$. Fix $\eta > 0$ sufficiently small and replace $\underline{v}$ by the strict supersolution $\underline{v}^\eta$ of Proposition 31. For $\varepsilon > 0$, maximize

$$\Phi_\varepsilon(t, s, x, x', y, y') := \bar{v}(t, x, y) - \underline{v}^\eta(s, x', y') - \frac{|x - x'|^2 + |y - y'|^2}{2\varepsilon} - \frac{|t - s|^2}{2\varepsilon}.$$

The strict growth barrier forces the maximizing points into a compact set independent of ε . The terminal ordering implies that, for sufficiently small ε , the maximizing times are strictly smaller than T . Apply the parabolic nonlocal theorem of sums at the maximizing points. There exist jets $(a_\varepsilon, p_\varepsilon, X_\varepsilon) \in \overline{\mathcal{P}}^{2,+}_{\bar{v}}$ and $(b_\varepsilon, p'_\varepsilon, Y_\varepsilon) \in \overline{\mathcal{P}}^{2,-}_{\underline{v}^\eta}$ such that $a_\varepsilon - b_\varepsilon = \frac{t_\varepsilon - s_\varepsilon}{\varepsilon} - \frac{t_\varepsilon - s_\varepsilon}{\varepsilon} = 0$, and the matrix inequality supplied by the theorem of sums holds.

By Lemma 29, both Hamiltonians may be restricted to a common compact control set. Fix a control u_ε that is ε -optimal for the supersolution Hamiltonian. Use the same control in the subsolution inequality. Subtracting the two viscosity inequalities gives

$$c\eta \leq [f(t_\varepsilon, x_\varepsilon, y_\varepsilon, u_\varepsilon) - f(s_\varepsilon, x'_\varepsilon, y'_\varepsilon, u_\varepsilon)] + \text{drift}_\varepsilon + \text{diffusion}_\varepsilon + \text{jump}_\varepsilon - \delta(\bar{v}(t_\varepsilon, x_\varepsilon, y_\varepsilon) - \underline{v}^\eta(s_\varepsilon, x'_\varepsilon, y'_\varepsilon)).$$

The uniform continuity in Assumption 4.1(i) and the matrix inequality show that the local drift, diffusion, and running-cost differences are $o_\varepsilon(1)$. The nonlocal contribution is $o_\varepsilon(1) + C\eta$ Lemma 30. If $\delta \geq 0$, the zero-order term has the favorable sign at a positive maximum. If $\delta = 0$, the strict supersolution perturbation alone supplies the contradiction. More generally, an exponential time change $v \mapsto e^{-\gamma(T-t)}v$ makes the equation proper. Sending $\varepsilon \downarrow 0$ therefore yields $c\eta \leq C\eta$ with the strictness parameter in Proposition 31 chosen so that $c > C$. This is impossible. Hence $\sup(\bar{v} - \underline{v}) \leq 0$, proving (40). $\square$

Proof of Lemma 44

Proof. Fix $P^1, P^2 \in \mathfrak{B}\mathfrak{R}(m)$ and $\lambda \in [0, 1]$, and set $P^\lambda := \lambda P^1 + (1 - \lambda)P^2$ as a probability measure on the canonical space Ω° . Since P^1 and P^2 have the same prescribed initial law m_0 , so does P^λ . For $\varphi \in C_c^2(\mathbb{R}^2)$, rational times $0 \leq r < s \leq T$, and every bounded canonical $\mathcal{F}_r$ -measurable cylinder functional H , admissibility of P^i gives $E^{P^i}[(M_s^\varphi - M_r^\varphi)H] = 0$, $i = 1, 2$. Hence $E^{P^\lambda}[(M_s^\varphi - M_r^\varphi)H] = \lambda E^{P^1}[(M_s^\varphi - M_r^\varphi)H] +$

$(1 - \lambda)E^{P^2}[(M_s^\Phi - M_r^\Phi)H] = 0$. The moment constraint is preserved because $E^{P^\lambda}[\int_0^T \int_{U_0} |a|^{P^*} q_t(da) dt] = \lambda E^{P^1}[\int_0^T \int_{U_0} |a|^{P^*} q_t(da) dt] + (1 - \lambda)E^{P^2}[\int_0^T \int_{U_0} |a|^{P^*} q_t(da) dt] < \infty$. Thus $P^\lambda \in \mathfrak{A}(m)$. The relaxed objective is affine in the canonical law: $J^m(P^\lambda) = \lambda J^m(P^1) + (1 - \lambda)J^m(P^2)$. Since P^1 and P^2 attain the same minimum $V^{m,\text{rel}}$, one has $J^m(P^\lambda) = V^{m,\text{rel}}$, so $P^\lambda \in \mathfrak{BR}(m)$. This proves convexity of $\mathfrak{BR}(m)$. Let $\mu^i \in \Phi(m)$ be generated by P^i . Then $P^\lambda \circ Z_t^{-1} = \lambda(P^1 \circ Z_t^{-1}) + (1 - \lambda)(P^2 \circ Z_t^{-1}) = \lambda\mu_t^1 + (1 - \lambda)\mu_t^2$ for every t . Hence $\lambda\mu^1 + (1 - \lambda)\mu^2 \in \Phi(m)$, proving convexity of $\Phi(m)$. No auxiliary time-zero randomization or Markov-control representation is used. $\square$

B Computational Specification and Reproducibility

This appendix specifies the numerical implementation used for Tables 2 and 3. All quantities not listed here retain the values in Table 1.

Numerical primitives. The numerical loss map is $a(z) = z$, and the physical-hedging technology is $h(u) = e^{-\gamma u}$. The drift and diffusion coefficients are $b(x, y, u) = rx + \pi_0 - \chi u^2/2$ and $\sigma_X(x, y, u) = \sigma_X$. The running cost is $f(t, x, y, u) = \chi u^2/2 + \zeta(e^y - 1)_+^2 + \omega(x_{\text{crit}} - x)_+^2$, and the terminal cost used in the computation is $g(x, y) = 2(x_{\text{crit}} - x)_+^2 + \frac{1}{2}(e^y - 1)_+^2$. For the mean field computation, $\psi_D(x, y) = (e^y - 1)_+$ and $\psi_S(x, y) = (x_{\text{crit}} - x)_+$.

The baseline initial state is $(X_0, Y_0) = (1, 0)$. For the population calculation we take $m_0 = \delta_{(1,0)}$. Thus the individual and population computations begin from the same normalized capitalization and replacement-cost state. If a nondegenerate initial distribution is used in a robustness experiment, its law is reported separately with the corresponding result.

The computational domain is $\mathcal{D}_h = [x_{\min}, x_{\max}] \times [y_{\min}, y_{\max}] = [-1, 3] \times [-0.75, 0.75]$, with $T = 5$. The baseline mesh is $\Delta t = 0.01$, $\Delta x = 0.01$, $\Delta y = 0.01$, and $\Delta u = 0.02$, with $U_\Delta = \{0, \Delta u, \dots, 2\}$.

Jump measure and quadrature. The numerical Lévy measure is $\nu(dz) = \lambda_J \alpha z_{\min}^\alpha z^{-1-\alpha} e^{-\tau_J(z-z_{\min})} \mathbf{1}_{\{z \geq z_{\min}\}} dz$. The nonlocal integral is truncated to $[z_{\min}, z_{\max}]$, with $z_{\max} = 20$. Let $\{z_k, \tilde{w}_k\}_{k=1}^{N_z}$ be the positive quadrature rule for the unnormalized density $\alpha z_{\min}^\alpha z^{-1-\alpha} e^{-\tau_J(z-z_{\min})}$. The weights entering the generator are $w_k = \lambda_J \tilde{w}_k$ and satisfy $\sum_{k=1}^{N_z} w_k \approx \nu([z_{\min}, z_{\max}])$. We use $N_z = 200$. No renormalization to unit mass is performed in the HJB operator: the quadrature approximates the Lévy measure itself. The omitted tail mass and moment are monitored through $\nu((z_{\max}, \infty))$ and $\int_{z_{\max}}^\infty z^q \nu(dz)$. For a shifted point $\xi = x_i - e^{y_j} a(z_k) h(u)$ satisfying $x_\ell \leq \xi \leq x_{\ell+1}$, monotone interpolation is $\mathcal{J}_x V(\xi, y_j) = (1 - \theta)V_{\ell,j} + \theta V_{\ell+1,j}$, where $\theta = (\xi - x_\ell)/\Delta x \in [0, 1]$. Hence the interpolation weights are nonnegative and sum to one. If $\xi < x_{\min}$, we set $\mathcal{J}_x V(\xi, y_j) = V(x_{\min}, y_j)$. If $\xi > x_{\max}$, we set $\mathcal{J}_x V(\xi, y_j) = V(x_{\max}, y_j)$. Thus jumps leaving the truncated surplus domain are projected onto the nearest boundary node. The domain-enlargement experiment reported below checks that this truncation does not materially affect Tables 2 and 3.

Boundary treatment. At $x = x_{\min}$ and $x = x_{\max}$ we use zero-normal-gradient ghost values, $V_{-1,j} = V_{0,j}$ and $V_{N_x+1,j} = V_{N_x,j}$. At $y = y_{\min}$ and $y = y_{\max}$ we similarly impose $V_{i,-1} = V_{i,0}$ and $V_{i,N_y+1} = V_{i,N_y}$. The same projection convention is used by the forward operator. These are numerical truncation conditions, not

economic boundary conditions on the original unbounded-state problem; their effect is therefore assessed by enlarging $\mathcal{D}_h$.

Monotone backward operator. For a fixed control u , write $\ell_{ijk}(u) = e^{y_j} a(z_k) h(u)$ and $\bar{\ell}_{ij}(u) = \sum_k w_k \ell_{ijk}(u)$. The compensated first-order coefficient is $\beta_{ij}(u) = b(x_i, y_j, u) + \bar{\ell}_{ij}(u)$. We discretize the first-order terms through the generator-form upwind operator $\beta_{ij}^+(u)(V_{i+1,j} - V_{i,j})/\Delta x + \beta_{ij}^-(u)(V_{i-1,j} - V_{i,j})/\Delta x$, where $\beta^+ = \max(\beta, 0)$ and $\beta^- = \max(-\beta, 0)$. For the y -drift, set $\mu_j = \kappa(\vartheta - y_j)$ and use the analogous decomposition $\mu_j^+(V_{i,j+1} - V_{i,j})/\Delta y + \mu_j^-(V_{i,j-1} - V_{i,j})/\Delta y$. The diffusion coefficients are $d_x = \sigma_X^2/(2\Delta x^2)$ and $d_y = \sigma_I^2/(2\Delta y^2)$. The quadrature-interpolation contribution has nonnegative off-diagonal coefficients because $w_k \geq 0$ and the interpolation weights belong to $[0, 1]$. Consequently, for each frozen policy, the spatial discrete generator L_h^u has nonnegative off-diagonal entries and satisfies $L_h^u \mathbf{1} = 0$ at interior nodes. The implicit backward matrix is $A_h^u = (I + \delta\Delta t)I - \Delta t L_h^u$. Its diagonal entries are positive, its off-diagonal entries are non-positive, and $A_h^u \mathbf{1} = (1 + \delta\Delta t)\mathbf{1}$. Hence A_h^u is strictly diagonally dominant and is a nonsingular M-matrix. In particular, $(A_h^u)^{-1} \geq 0$, which gives monotonicity of the fixed-policy backward step.

The linear system at each policy-evaluation step is solved to relative residual tolerance 10^{-10} . Policy iteration is terminated when $\max_{n,i,j} |u_{i,j}^{(r+1),n} - u_{i,j}^{(r),n}| \leq 10^{-7}$.

Positivity of the forward step. The forward discretization uses the transpose of the same generator: $m^{n+1} = (I + \Delta t (L_h^{\alpha^n})^\top) m^n$. Since the off-diagonal entries of $L_h^{\alpha^n}$ are nonnegative, this explicit step preserves nonnegativity provided $\Delta t \Lambda_h \leq 1$, where $\Lambda_h := \max_{n,i,j} [-(L_h^{\alpha^n})_{(i,j),(i,j)}]$. Equivalently, a sufficient CFL condition is $\Delta t \max_{n,i,j} \{|\beta_{ij}(\alpha_{ij}^n)|/\Delta x + |\mu_j|/\Delta y + \sigma_X^2/\Delta x^2 + \sigma_I^2/\Delta y^2 + \sum_k w_k\} \leq 1$. The implementation checks this inequality before the forward sweep. If it fails on a refined spatial grid, the forward equation is substepped with $\Delta t_F \leq \Lambda_h^{-1}$ while the backward HJB time grid is left unchanged. After every forward sweep we verify $\min_{n,i,j} m_{i,j}^n \geq -10^{-14}$ and $\max_n |\sum_{i,j} m_{i,j}^n - 1| \leq 10^{-12}$.

Mean field initialization and fixed-point diagnostics. The baseline fixed-point iteration starts from the uncontrolled flow $m^{(0)} = \mathcal{L}(X^{u \equiv 0}, Y)$. To test dependence on initialization, the computation is repeated from three additional flows: the state law generated by the maximal control $u \equiv u_{\max}$, the midpoint control $u \equiv u_{\max}/2$, and the convex midpoint of the uncontrolled and maximally controlled flows. For each initialization, the same damping parameter $\omega_{\text{MFG}} = 0.25$ is used. The iteration is terminated when $\max_n W_1(m_n^{(k+1)}, m_n^{(k)}) \leq 10^{-6}$ and the consistency residual $\max_n W_1(m_n^{(k)}, \hat{m}_n^{(k)}) \leq 10^{-6}$. No theoretical convergence claim for this iteration is made. Uniqueness of the continuous equilibrium does not imply convergence of the discrete fixed-point algorithm. Numerical convergence is accepted only when all four initializations reach the same discrete solution within the reported tolerance.

Grid and domain verification. Tables 2 and 3 are recomputed on at least three nested discretizations. For every reported scalar Q , we record $\mathcal{E}_{\text{grid}}(Q) = |Q_{h/2} - Q_h|/(1 + |Q_{h/2}|)$. The calculation is also repeated after increasing $x_{\max}$, decreasing $x_{\min}$, increasing $|y_{\min}|$ and $y_{\max}$, and increasing $z_{\max}$. Numerical conclusions are retained only when the discretization and domain effects are smaller than the economic comparative-static effect being reported.

Algorithm. The complete computation is summarized by the following pseudocode.

- 1: Set model parameters, $\mathcal{D}_h$, U_Δ , and the positive Lévy quadrature.
- 2: Set $V_{i,j}^{N_t} = g(x_i, y_j)$.
- 3: **for** $n = N_t - 1, \dots, 0$ **do**
- 4: Initialize the policy $u^{(0),n}$.
- 5: **repeat**
- 6: Assemble the M-matrix $A_h^{u^{(r),n}}$.
- 7: Solve the fixed-policy linear system for $V^{(r+1),n}$.
- 8: Minimize the discrete Hamiltonian over U_Δ nodewise to obtain $u^{(r+1),n}$.
- 9: **until** the policy tolerance is satisfied
- 10: **end for**
- 11: Propagate m_0 with the positivity-preserving adjoint forward operator, using CFL substepping when necessary.
- 12: Update the population flow by damped fixed-point iteration.
- 13: Repeat the backward and forward sweeps until both MFG residuals are below tolerance.
- 14: Repeat from all prescribed initial population-flow guesses.
- 15: Repeat on refined grids and enlarged state/jump domains.
- 16: Generate Tables 2 and 3.

Computational environment. The numerical calculations underlying Tables 2 and 3 were implemented in Python using the NumPy scientific-computing library. Randomized calculations use seed 42. The model parameters, computational domain, quadrature rule, interpolation procedure, boundary treatment, and stopping criteria are documented in this Appendix. Executable Python code reproducing Tables 2 and 3 is provided as Supplementary File.

References

- [1] Achdou, Y. (2013). Finite difference methods for mean field games. In *Hamilton-Jacobi Equations: Approximations, Numerical Analysis and Applications*. Springer.
- [2] Albrecher, H. and Asmussen, S. (2006). Ruin probabilities and aggregate claims distributions for shot noise cox processes. *Journal of Applied Probability*, 43(1):1–12.
- [3] Anderson, V., Pramanik, P., and Robinson, H. K. (2026). Obesity and sociodemographic factors in luminal breast cancer. *arXiv preprint arXiv:2605.30591*.
- [4] Applebaum, D. (2009). *Lévy Processes and Stochastic Calculus*, volume 116 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2 edition.
- [5] Asmussen, S. (2000). *Ruin Probabilities*. World Scientific.
- [6] Avanzi, B. (2009). Strategies for dividend distribution: A review. *North American Actuarial Journal*.

- [7] Azcue, P. and Muler, N. (2010). Optimal reinsurance and dividend distribution policies with transaction costs. *Insurance: Mathematics and Economics*.
- [8] Barles, G. and Imbert, C. (2008). Second-order elliptic integro-differential equations: Viscosity solutions' theory revisited. *Annales de l'Institut Henri Poincaré C, Analyse Non Linéaire*, 25(3):567–585.
- [9] Biswas, I. H., Jakobsen, E. R., and Karlsen, K. H. (2010). Viscosity solutions for a system of integro-pdes and connections to optimal switching and control of jump-diffusion processes. *Applied Mathematics and Optimization*, 62(1):47–80.
- [10] Carmona, R. and Delarue, F. (2018). *Probabilistic Theory of Mean Field Games with Applications I: Mean Field FBSDEs, Control, and Games*, volume 83 of *Probability Theory and Stochastic Modelling*. Springer, Cham.
- [11] Carmona, R., Fouque, J.-P., and Sun, L.-H. (2015). Mean field games and systemic risk. *Communications in Mathematical Sciences*.
- [12] Choupi, L., Margaritis, V., and Angelidis, G. (2026). Scenario-based financial planning in gold mining under commodity price uncertainty. *Commodities*, 5(2):12.
- [13] Chowdhury, I., Ersland, O., and Jakobsen, E. R. (2023). On numerical approximations of fractional and nonlocal mean field games. *Foundations of Computational Mathematics*.
- [14] Chowdhury, I., Jakobsen, E. R., and Krupski, M. (2024). On fully nonlinear parabolic mean field games with nonlocal and local diffusions. *SIAM Journal on Mathematical Analysis*.
- [15] Crandall, M. G., Ishii, H., and Lions, P.-L. (1992). User's guide to viscosity solutions of second order partial differential equations. *Bulletin of the American Mathematical Society*, 27(1):1–67.
- [16] Dasgupta, S., Acharya, S., Khan, M. A., Pramanik, P., Marbut, S. M., Yunus, F., Galeas, J. N., Singh, S., Singh, A. P., and Dasgupta, S. (2023). Frequent loss of cacna1c, a calcium voltage-gated channel subunit is associated with lung adenocarcinoma progression and poor prognosis. *Cancer Research*, 83(7_Supplement):3318–3318.
- [17] Dasgupta, S., Galappaththi, S. L., Banerjee, R., Alsatari, E. S., Pramanik, P., Marbut, S. M., Yunus, F., Galeas, J. N., and Dasgupta, S. (2026). Frequent loss of cacna1c is associated with poor prognosis in non-small cell lung cancer. *The FASEB Journal*, 40(4):e71614.
- [18] Dassios, A. and Jang, J.-W. (2003). Pricing of catastrophe reinsurance and derivatives using the cox process with shot noise intensity. *Insurance: Mathematics and Economics*.
- [19] Dassios, A. and Zhao, H. (2012). Ruin by dynamic contagion claims.
- [20] Döhrmann, D., Gürtler, M., and Hibbeln, M. (2017). Insured loss inflation: How natural catastrophes affect reconstruction costs. *Journal of Risk and Insurance*.
- [21] Dufresne, F. and Gerber, H. U. (1991). Risk theory for the compound poisson process that is perturbed by diffusion. *Insurance: Mathematics and Economics*.

- [22] Dunbar, B., Pramanik, P., and Robinson, H. K. (2026). Modeling age-adjusted mortality in the united states. *arXiv preprint arXiv:2601.13504*.
- [23] El Karoui, N., Nguyen, D. H., and Jeanblanc-Picqué, M. (1987). Compactification methods in the control of degenerate diffusions: Existence of an optimal control. *Stochastics*, 20(3):169–219.
- [24] El Karoui, N. and Tan, X. (2013). Capacities, measurable selection and dynamic programming part ii: Application in stochastic control problems. *arXiv preprint arXiv:1310.3364*.
- [25] Ellington, C., Pramanik, P., and Robinson, H. K. (2025a). Metascorelens: Evaluating user feedback across digital entertainment systems. *arXiv preprint arXiv:2601.11523*.
- [26] Ellington, C., Pramanik, P., and Robinson, H. K. (2025b). Playmydata: A statistical analysis of a video game dataset on review scores and gaming platforms. *Analytics*, 4(4):31.
- [27] Embrechts, P., Klüppelberg, C., and Mikosch, T. (1997). *Modelling Extremal Events for Insurance and Finance*. Springer, Berlin.
- [28] Fayvisovich, R. and Žitković, G. (2021). A framework for the dynamic programming principle and martingale-generated control correspondences. *Applied Mathematics & Optimization*, 83:2437–2472.
- [29] Finetti, B. D. (1957). Su un’impostazione alternativa della teoria collettiva del rischio. *Giornale dell’Istituto Italiano degli Attuari*.
- [30] Fleming, W. H. and Soner, H. M. (2006). *Controlled Markov Processes and Viscosity Solutions*, volume 25 of *Stochastic Modelling and Applied Probability*. Springer, New York, 2 edition.
- [31] Garrido, J. (1989). Stochastic differential equations in the analysis of ruin. *Insurance: Mathematics and Economics*.
- [32] Gerber, H. U. (1969). Entscheidungskriterien fuer den zusammengesetzten poisson-prozess. *Mitteilungen der Vereinigung Schweizerischer Versicherungsmathematiker*.
- [33] Hertweck, K. L., Vikramdeo, K. S., Galeas, J. N., Marbut, S. M., Pramanik, P., Yunus, F., Singh, S., Singh, A. P., and Dasgupta, S. (2023). Clinicopathological significance of unraveling mitochondrial pathway alterations in non-small-cell lung cancer. *The FASEB Journal*, 37(7):e23018.
- [34] Høgaard, B. and Taksar, M. (1999). Optimal dynamic reinsurance policies for large insurance portfolios. *Applied Mathematics and Optimization*.
- [35] Hua, L., Polansky, A., and Pramanik, P. (2019). Assessing bivariate tail non-exchangeable dependence. *Statistics & Probability Letters*, 155:108556.
- [36] Jakobsen, E. R. and Karlsen, K. H. (2006). A “maximum principle for semicontinuous functions” applicable to integro-partial differential equations. *NoDEA Nonlinear Differential Equations and Applications*, 13(2):137–165.

- [37] Kakkat, S., Pramanik, P., Singh, S., Singh, A. P., Sarkar, C., and Chakroborty, D. (2023). Cardiovascular complications in patients with prostate cancer: Potential molecular connections. *International Journal of Molecular Sciences*, 24(8):6984.
- [38] Kakkat, S., Suman, P., Goswami, S., Kola, B., Bruno, K. A., Frankel, W. L., Basu, S., Turbat-Herrera, E. A., Ramirez-Alcantara, V., Heslin, M. J., et al. (2026). Angiotensin ii-angiotensin ii receptor type 1 signaling facilitates gastric cancer metastasis via kruppel-like factor 4 suppression and tight junction breakdown. *bioRxiv*, pages 2026–01.
- [39] Khan, M. A., Acharya, S., Anand, S., Sameeta, F., Pramanik, P., Keel, C., Singh, S., Carter, J. E., Dasgupta, S., and Singh, A. P. (2023). Myb exhibits racially disparate expression, clinicopathologic association, and predictive potential for biochemical recurrence in prostate cancer. *Iscience*, 26(12).
- [40] Khan, M. A., Acharya, S., Kreitz, N., Anand, S., Sameeta, F., Pramanik, P., Keel, C., Singh, S., Carter, J., Dasgupta, S., et al. (2024). Mp60-05 myb exhibits racially disparate expression and clinicopathologic association and is a promising predictor of biochemical recurrence in prostate cancer. *Journal of Urology*, 211(5S):e1000.
- [41] Klüppelberg, C. and Mikosch, T. (1997). Large deviations of heavy-tailed random sums with applications in insurance and finance. *Journal of Applied Probability*, 34(2):293–308.
- [42] Kousky, C. (2019). The role of natural disaster insurance in recovery and risk reduction. *Annual Review of Resource Economics*, 11:399–418.
- [43] Kurtz, T. G. and Stockbridge, R. H. (1998). Existence of markov controls and characterization of optimal markov controls. *SIAM Journal on Control and Optimization*, 36(2):609–653.
- [44] Lackner, D. (2016). A general characterization of the mean field limit for stochastic differential games. *Probability Theory and Related Fields*, 165(3–4):581–648.
- [45] Lasry, J.-M. and Lions, P.-L. (2007). Mean field games. *Japanese Journal of Mathematics*, 2(1):229–260.
- [46] Liu and Cadenillas (2023). Optimal insurance contracts for a shot-noise cox claim process and persistent insured’s actions.
- [47] Macci, C. and Torrisi, G. L. (2011). Risk processes with shot noise cox claim number process and reserve dependent premium rate. *Insurance: Mathematics and Economics*, 48(2):285–298.
- [48] Øksendal, B. and Sulem, A. (2019). *Applied Stochastic Control of Jump Diffusions*. Springer, Cham, 3 edition.
- [49] Olsen, A. H. and Porter, K. A. (2011). What we know about demand surge: Brief summary. *Natural Hazards Review*.
- [50] Panjer, H. H. and Willmot, G. E. (1992). *Insurance Risk Models*. Society of Actuaries.

- [51] Pham, H. (2009). *Continuous-time Stochastic Control and Optimization with Financial Applications*, volume 61 of *Stochastic Modelling and Applied Probability*. Springer, Berlin.
- [52] Pojer, J. and Thonhauser, S. (2023). Ruin probabilities in a markovian shot-noise environment. *Applied Mathematics and Optimization*.
- [53] Polansky, A. M. and Pramanik, P. (2021). A motif building process for simulating random networks. *Computational Statistics & Data Analysis*, 162:107263.
- [54] Powell, A. and Pramanik, P. (2025). Genomic influence of a key transcription factor in male glandular malignancy. *arXiv preprint arXiv:2510.11959*.
- [55] Powell, A. and Pramanik, P. (2026). The role of myb in prostate cancer: A statistical analysis. *European Journal of Statistics*, 6:1–1.
- [56] Pramanik, P. (2016). *Tail non-exchangeability*. Northern Illinois University.
- [57] Pramanik, P. (2020). Optimization of market stochastic dynamics. In *SN Operations Research Forum*, volume 1, page 31. Springer.
- [58] Pramanik, P. (2021a). Consensus as a nash equilibrium of a stochastic differential game. *arXiv preprint arXiv:2107.05183*.
- [59] Pramanik, P. (2021b). Effects of water currents on fish migration through a feynman-type path integral approach under $8/3$ liouville-like quantum gravity surfaces. *Theory in Biosciences*, 140(2):205–223.
- [60] Pramanik, P. (2021c). *Optimization of dynamic objective functions using path integrals*. PhD thesis, Northern Illinois University.
- [61] Pramanik, P. (2022a). On lock-down control of a pandemic model. *arXiv preprint arXiv:2206.04248*.
- [62] Pramanik, P. (2022b). Stochastic control of a sir model with non-linear incidence rate through euclidean path integral. *arXiv preprint arXiv:2209.13733*.
- [63] Pramanik, P. (2023a). Optimal lock-down intensity: A stochastic pandemic control approach of path integral. *Computational and Mathematical Biophysics*, 11(1):20230110.
- [64] Pramanik, P. (2023b). Path integral control in infectious disease modeling. *arXiv preprint arXiv:2311.02113*.
- [65] Pramanik, P. (2023c). Path integral control of a stochastic multi-risk sir pandemic model. *Theory in Biosciences*, 142(2):107–142.
- [66] Pramanik, P. (2024a). Dependence on tail copula. *J*, 7(2):127–152.
- [67] Pramanik, P. (2024b). Measuring asymmetric tails under copula distributions. *European Journal of Statistics*, 4:7–7.

- [68] Pramanik, P. (2024c). On estimation of function-on-function regression kernels with brownian berkson errors.
- [69] Pramanik, P. (2024d). Stochastic control in determining a soccer player's performance. *J. Compr. Pure Appl. Math*, 2:111.
- [70] Pramanik, P. (2025a). Construction of an optimal strategy: An analytic insight through path integral control driven by a mckean–vlasov opinion dynamics. *Mathematics*, 13(17):2842.
- [71] Pramanik, P. (2025b). Construction of an optimal strategy: An analytic insight through path integral control driven by a mckean–vlasov opinion dynamics. *Mathematics*, 13(17):2842.
- [72] Pramanik, P. (2025c). Optimal feedback control in social networks in a mckean-vlasov-friedkin-johnsen system. *arXiv preprint arXiv:2508.17138*.
- [73] Pramanik, P. (2025d). Stubbornness as control in professional soccer games: A bppsde approach. *Mathematics*, 13(3):475.
- [74] Pramanik, P. (2026a). The quantum structure of markets: Linking hamiltonian-jacobi-bellman dynamics to schrodinger equation through feynman action. *arXiv preprint arXiv:2603.25086*.
- [75] Pramanik, P. (2026b). Stochastic choice with distribution-dependent preferences. *arXiv preprint arXiv:2608.06152*.
- [76] Pramanik, P. (2026c). Strategic dynamics of firms via path integral control. *International Game Theory Review*, page 2650006.
- [77] Pramanik, P., Boone, E. L., and Ghanam, R. A. (2024a). Parametric estimation in fractional stochastic differential equation. *Stats*, 7(3):745–760.
- [78] Pramanik, P. and Dong, L. (2025a). Impact of random monetary shock: a keynesian case. *arXiv preprint arXiv:2505.00800*.
- [79] Pramanik, P. and Dong, L. (2025b). Strategic complementarities due to monetary shock under sticky price. *European Journal of Statistics*, 5:9–9.
- [80] Pramanik, P., Graff, J., and DeCaro, M. (2024b). Analysis of a tiered pricing model for ecw clients. In *International Conference on Mathematical Modeling in Physical Sciences*, pages 515–535. Springer.
- [81] Pramanik, P., Graff, J., and Decaro, M. (2025a). Dissecting multi-level pricing schemes in the context of ecw client engagement. *arXiv preprint arXiv:2509.22669*.
- [82] Pramanik, P., Graff, J., and Decaro, M. (2025b). On factors influencing consumer preference in pipeline stages: an experiment. *arXiv preprint arXiv:2501.03418*.
- [83] Pramanik, P., Graff, J., and Decaro, M. (2025c). Strategies to increase pipeline status: A case study from eclinical data. *European Journal of Statistics*, 5:3–3.

- [84] Pramanik, P. and Johora, F. (2026). Optimal harvesting under stochastic control: Hjb equation and feynman-kac representation. *arXiv preprint arXiv:2606.08428*.
- [85] Pramanik, P. and Maity, A. K. (2024). Bayes factor of zero inflated models under jeffreys prior. *arXiv preprint arXiv:2401.03649*.
- [86] Pramanik, P., Maity, A. K., Mandal, A., and Robinson, H. K. (2026). A bayesian discrete framework for enhancing decision-making processes in clinical trial designs and evaluations. *arXiv preprint arXiv:2601.10615*.
- [87] Pramanik, P. and Polansky, A. M. (2020). Motivation to run in one-day cricket. *arXiv preprint arXiv:2001.11099*.
- [88] Pramanik, P. and Polansky, A. M. (2021). Optimal estimation of brownian penalized regression coefficients. *arXiv preprint arXiv:2107.02291*.
- [89] Pramanik, P. and Polansky, A. M. (2023a). Optimization of a dynamic profit function using euclidean path integral. *SN Business & Economics*, 4(1):8.
- [90] Pramanik, P. and Polansky, A. M. (2023b). Scoring a goal optimally in a soccer game under liouville-like quantum gravity action. In *Operations Research Forum*, volume 4, page 66. Springer.
- [91] Pramanik, P. and Polansky, A. M. (2024a). Motivation to run in one-day cricket. *Mathematics*, 12(17):2739.
- [92] Pramanik, P. and Polansky, A. M. (2024b). Semicoooperation under curved strategy spacetime. *The Journal of Mathematical Sociology*, 48(2):172–206.
- [93] Reed, B. and Pramanik, P. (2026a). Modeling educational performance using school demographics and teacher characteristics. *arXiv preprint arXiv:2606.27654*.
- [94] Reed, B. and Pramanik, P. (2026b). Structural inequities and mathematics achievement in alabama public schools. *Analytics*, 5(3):23.
- [95] Roth, Richard J., S. and Kunreuther, H., editors (1998). *Paying the Price: The Status and Role of Insurance Against Natural Disasters in the United States*. Joseph Henry Press, Washington, DC.
- [96] Schmidli, H. (2008). *Stochastic Control in Insurance*. Probability and Its Applications. Springer, London.
- [97] Situ, R. (2005). *Theory of Stochastic Differential Equations with Jumps and Applications*. Mathematical and Analytical Techniques with Applications to Engineering. Springer, New York.
- [98] Valdez, I. and Pramanik, P. (2025a). Association between obesity, race, and luminal subtypes of breast cancer. *European Journal of Statistics*, 5:12–12.
- [99] Valdez, I. and Pramanik, P. (2025b). Exploring the interplay of adiposity, ethnicity, and hormone receptor profiles in breast cancer subtypes. *arXiv preprint arXiv:2507.21348*.

- [100] Vikramdeo, K., Anand, S., Sudan, S., Pramanik, P., Singh, S., Godwin, A., Singh, A., and Dasgupta, S. (2024a). Abstract po3-16-05: Mitochondrial dna mutation detection in tumors and circulating extracellular vesicles of triple negative breast cancer patients for biomarker development. *Cancer Research*, 84(9_Supplement):PO3–16.
- [101] Vikramdeo, K., Anand, S., Sudan, S., Pramanik, P., Singh, S., Godwin, A., Singh, A., and Dasgupta, S. (2024b). Mitochondrial dna mutation detection in tumors and circulating extracellular vesicles of triple negative breast cancer patients for biomarker development. In *CANCER RESEARCH*, volume 84. AMER ASSOC CANCER RESEARCH 615 CHESTNUT ST, 17TH FLOOR, PHILADELPHIA, PA
- [102] Vikramdeo, K. S., Anand, S., Sudan, S. K., Pramanik, P., Singh, S., Godwin, A. K., Singh, A. P., and Dasgupta, S. (2023). Profiling mitochondrial dna mutations in tumors and circulating extracellular vesicles of triple-negative breast cancer patients for potential biomarker development. *FASEB BioAdvances*, 5(10):412.
- [103] Yusuf, A. and Pramanik, P. (2025a). Predictive significance of cd276/b7-h3 expression in baseline biopsies of advanced prostate carcinoma. *arXiv preprint arXiv:2508.09373*.
- [104] Yusuf, A. and Pramanik, P. (2025b). Prognostic role of b7-h3 (cd276) expression in initial biopsies of metastatic prostate cancer. *Onco*, 5(3):38.
- [105] Zeller, G. and Scherer, M. (2022). A comprehensive model for cyber risk based on marked point processes. *Insurance: Mathematics and Economics*.
- [106] Žitković, G. (2014). Dynamic programming for controlled markov families: Abstractly and over martingale measures. *SIAM Journal on Control and Optimization*, 52(3):1597–1621.