

The Mathematics of Heuristic Portfolio Optimization (HPO)

Miquel Noguer i Alonso
Artificial Intelligence Finance Institute

June 12, 2026

Abstract

Practitioners often allocate capital with forecast-light rules such as equal weight, inverse volatility, risk parity, HRP, and return-adjusted HRP (RA-HRP). This paper develops *Heuristic Portfolio Optimization* (HPO): portfolio construction as an information-restricted projection of the Markowitz/tangency solution onto a stable rule class. The implied-return principle, $\mathbf{w}$ is maximum-Sharpe iff $\boldsymbol{\mu}_e \propto \boldsymbol{\Sigma}\mathbf{w}$, yields closed-form optimality sets for the main heuristics and exposes the Schur-complement substitutions behind HRP. For RA-HRP we add fixed-tree cluster-Sharpe recursion, unit-free HRP–RA-HRP interpolation, nonlinear tangency-coincidence conditions, Schur-RA-HRP conditional-risk splits, and pathwise/KL decompositions of HRP–RA-HRP weight distortion; a first-order Sharpe calculus along the unit-free homotopy expresses the marginal value of return information as a sum of nodewise alphas against HRP and certifies a KL trust budget linear in the interpolation weight. We then formalize generic HPO maps, define the implied-return defect, prove that it equals squared Sharpe inefficiency, characterize tree-HPO coincidence by nodewise mass ratios, and give an exact bias–variance decomposition for estimated heuristic rules. The final contribution embeds HPO into Reinforcement Learning Portfolio Optimization (RLPO): every HPO map induces a deterministic stationary policy; static HPO is the $\gamma = 0$ no-friction face of the Bellman problem; RA-HRP provides a hierarchical policy prior through node-level split probabilities; and dynamic policy improvement is justified exactly when continuation value exceeds myopic HPO defect and trading frictions; a performance-difference identity prices the total value gap of acting myopically, yielding an $\varepsilon/(1 - \gamma)$ myopia bound, and the nodewise alphas reappear as the myopic policy-gradient coordinates of the hierarchical actor at its prior. The resulting framework treats HPO as the static optimality layer and RLPO as the dynamic control layer. The conditions remain GRS-testable, transfer to mean–CVaR and expected utility under ellipticity, and become Kelly-growth conditions in diffusion limits.

Keywords: heuristic portfolio optimization, HPO, reinforcement learning portfolio optimization, RLPO, hierarchical risk parity, return-adjusted HRP, Schur complement, implied returns, Sharpe efficiency, policy priors, model ambiguity.

Contents

1	Introduction	4
2	Scope, Standing Assumptions, and Paper Map	6
2.1	Standing assumptions	6
2.2	Contribution map	7
2.3	What the theory deliberately does not assert	7
3	Setup, the Implied-Return Principle, and the Geometry of Optimality	7
3.1	Notation	7
3.2	The first-order characterization	8
3.3	Efficiency is an angle	9
3.4	The catalogue	10
4	Exact Conditions: Equal Weight, Inverse Volatility, Minimum Variance, Maximum Diversification	10
4.1	Equal weight	10
4.2	Inverse volatility	11
4.3	Minimum variance	11
4.4	Maximum diversification	11
5	One-Factor Structure: Conditions in Observables	12
6	Risk Parity: Existence, Uniqueness, and Optimality as Performance Parity	12
6.1	Existence and uniqueness	13
6.2	Optimality	13
7	Hierarchical and Return-Adjusted Hierarchical Risk Parity	14
7.1	The algorithm	14
7.2	What the optimizer actually does at a split	14
7.3	Exact coincidence results	15
7.4	Return-adjusted hierarchical risk parity	16
7.5	First-order Sharpe calculus along the HRP–RA–HRP homotopy	21
8	The Mathematics of Heuristic Portfolio Optimization (HPO)	23
8.1	HPO maps as information-restricted portfolio rules	24
8.2	The implied-return defect of a heuristic rule	24
8.3	Tree HPO: local scores as stochastic kernels	25
8.4	The exact Schur-tangency split	26
8.5	Sampling error, approximation error, and the HPO bias–variance identity	27
8.6	The master HPO objective	28
9	From HPO to Reinforcement Learning Portfolio Optimization (RLPO)	28
9.1	Controlled market formulation	29
9.2	Continuation value versus myopic HPO defect	30
9.3	The exact value gap of an HPO policy	31
9.4	HPO-regularized RLPO objectives	32
9.5	RA–HRP as a hierarchical policy prior	32
9.6	Tree topology, regime shifts, and the dynamic open problem	33
9.7	Operational blueprint	34

10 Transfer: Elliptical Risk Measures and Growth Optimality	35
10.1 Elliptical transfer	35
10.2 Growth-optimal transfer	35
11 Symmetry and Bayes Optimality: When the Heuristic Is the Optimizer	36
12 Ambiguity, Robustness, and Estimation	37
13 Constrained, Costly, and Implementable Optimality	38
13.1 Normal-cone optimality under mandate constraints	38
13.2 Transaction-cost wedges	39
13.3 Benchmark-relative and factor-neutral mandates	40
14 Estimation Geometry, High-Dimensional Risk, and Empirical Diagnostics	40
14.1 First-order sensitivity of tangency weights	40
14.2 Projection onto a heuristic optimality set	41
14.3 GRS and spanning tests as optimality tests	41
14.4 High-dimensional covariance and the inversion tax	42
14.5 A reproducible empirical protocol	42
14.6 Computational profile of the allocation ladder	43
15 Non-Elliptical Tails, Downside Risk, and Where the Conditions Break	43
16 Synthesis: The Optimality Map	44
17 Limitations and Open Problems	48
18 Conclusion	48
A Technical Appendix: Identities Used Repeatedly	50
A.1 Block inverse and Schur complement	50
A.2 Ray projection	50
A.3 Tree products and binary KL	50

1 Introduction

Seventy years after Markowitz (1952), a striking share of institutional capital is allocated by rules that refuse to optimize. The equal-weight portfolio ignores both means and covariances; inverse-volatility weighting uses only marginal volatilities; equal-risk-contribution (ERC, “risk parity”) portfolios use the full covariance matrix but no means (Maillard et al., 2010; Roncalli, 2013; Qian, 2006); maximum diversification uses volatilities and covariances in a specific ratio (Choueifaty and Coignard, 2008); hierarchical risk parity (HRP) uses the covariance matrix only through a clustering tree and a sequence of within-cluster variances (López de Prado, 2016); and return-adjusted HRP (RA-HRP) keeps the tree but lets local cluster-Sharpe scores tilt the recursive budget (Noguer i Alonso, 2026). The key refinement is not merely that RA-HRP uses expected returns, but that return information is consumed locally, after clustering, through dimensionless split fractions that can be interpolated continuously from HRP without mixing inverse variances and Sharpe ratios. The standard defense of these rules is empirical: optimized portfolios inherit estimation error in μ at first order (Merton, 1980; Michaud, 1989; Best and Grauer, 1991; Chopra and Ziemba, 1993), and naive rules are hard to beat out of sample (DeMiguel et al., 2009).

We use the term *Heuristic Portfolio Optimization* (HPO) for this discipline: the mathematical study of portfolio maps that deliberately consume less information than the full Markowitz optimizer, together with the exact optimality, efficiency, estimation, and belief conditions under which that deletion of information is harmless or beneficial. In this sense HPO is not the opposite of optimization. It is optimization after restricting the admissible decision maps to those that are stable, interpretable, low-inversion, low-turnover, or compatible with a reduced information set.

This paper develops the theory behind that practice at five levels of depth.

Level 1: exact optimality conditions. The implied-return principle (Theorem 3.3) states that a fully invested portfolio with positive expected excess return is maximum-Sharpe if and only if expected excess returns are proportional to $\Sigma \mathbf{w}$. Consequently *every* heuristic is optimal for *some* parameter configuration, and the substantive object is the exact shape of its *optimality set*

$$\mathcal{O}(\mathbf{w}^\circ) = \{(\mu_e, \Sigma) : \mu_e = a \Sigma \mathbf{w}^\circ, a > 0\},$$

because adopting the rule is behaviorally equivalent to asserting membership. We compute this set in closed form for every rule in common use (Sections 4–7): zero alpha against itself for equal weight (Theorem 4.1); Sharpe ratios proportional to correlation row sums for inverse volatility (Theorem 4.3); equal premia for minimum variance (Theorem 4.5); equal Sharpe ratios, with no restriction on correlations, for maximum diversification (Theorem 4.7); and, for risk parity, the new characterization that the equal *risk* budget must double as an equal *performance* budget, $w_i \mu_{e,i}$ constant (Theorem 6.2), of which the equal-Sharpe/constant-correlation condition of Maillard et al. (2010) is a corollary and demonstrably far from necessary. For HRP we first derive the exact Schur-complement bisection identity satisfied by the true minimum-variance portfolio (Theorem 7.2), which isolates the three substitutions the algorithm makes — raw block for Schur complement, flat budget for tilted budget, inverse variance for within-cluster minimum variance — and then prove the two regimes in which all three are simultaneously costless: diagonal covariance under *any* dendrogram (Theorem 7.5) and uncorrelated exchangeable blocks under balanced tree-aligned bisection (Proposition 7.6), together with the minimal two-asset counterexample marking the boundary (Remark 7.7). For RA-HRP we show that adding returns changes the optimality set from a ray to a nonlinear fixed-point set: the exact condition is $\mu_e = a \Sigma \Phi_{\mathcal{T}, \varepsilon}^{\text{RA}}(\mu_e, \Sigma)$, equivalently node-by-node equality between the RA-HRP floored cluster-Sharpe split and the exact Schur-tangency split (Theorem 7.15). We also prove a two-asset boundary result, a balanced equal-volatility diagonal exactness result, and a pathwise/KL decomposition of the

distortion from HRP to RA-HRP (Corollary 7.20, Proposition 7.21, Theorem 7.22). A first-order Sharpe calculus along the homotopy (Section 7.5) then computes the derivative of efficiency in the interpolation weight as a sum of nodewise alphas against HRP itself (Theorem 7.27), making the marginal value of return information a GRS-estimable quantity, and bounds the induced reallocation by a KL trust budget linear in λ (Proposition 7.29). The interpolation from HRP to RA-HRP is made at the level of local branch probabilities $r_v, s_v \in (0, 1)$, not by adding inverse-variance scores to Sharpe scores; this unit-free homotopy is the mathematically clean way to control how much local return information enters the hierarchy. We then add Schur-RA-HRP, which replaces standalone cluster risk by conditional Schur risk and therefore targets exactly the cross-cluster covariance gap diagnosed by the HRP Schur identity.

Level 2: the geometry and cost of suboptimality. Conditions that hold on a measure-zero set would be a curiosity unless violations were cheap. Both halves are theorems. The Sharpe ratio of *any* portfolio obeys the exact angular identity $S(\mathbf{w}) = S_{\max} \cos \theta_{\Sigma}(\mathbf{w}, \mathbf{w}_T)$ in the Σ -inner product (Theorem 3.7): inefficiency is purely angular. Each optimality set is a ray — Lebesgue-null in $\mathbb{R}^N$ — so under any diffuse belief, exact optimality of a fixed heuristic is a probability-zero event (Proposition 3.8). Yet the efficiency map is *stationary* on the set: perturbing μ_e off the coincidence ray by εv degrades squared Sharpe efficiency by an explicitly computable $O(\varepsilon^2)$ term whose coefficient is a generalized residual variance (Theorem 3.9). Never optimal, rarely costly — with the rate and the constant. Each condition is moreover a zero-intercept restriction in time-series regressions on the heuristic itself, hence directly testable by the F -test of Gibbons et al. (1989, C) (Remark 3.6). Under one-factor structure, every condition becomes a statement about observable betas and idiosyncratic variances (Section 5), including a self-contained derivation of the minimum-variance threshold-beta representation of Clarke et al. (2011) and the result that maximum diversification is optimal in a priced one-factor world precisely when all assets are equally correlated with the factor.

Level 3: in what sense the heuristics are right. Two transfer theorems show the conditions are not artifacts of the quadratic objective: under elliptical returns they govern mean-CVaR optimality at every confidence level and expected-utility optimality for every risk-averse investor (Theorem 10.2), and in the continuous-time limit they are simultaneously the conditions for (fractional) Kelly growth optimality (Proposition 10.3). The deepest reading then inverts the question: each heuristic is the *exactly optimal* policy of an investor whose information set is smaller than (μ_e, Σ) , with ignorance resolved by symmetry. We prove Bayes-optimality theorems at three rungs (Section 11): full exchangeability of beliefs forces the Bayes portfolio to be $1/N$ (Theorem 11.3); exchangeability of beliefs about *standardized* returns, volatilities known, forces inverse volatility (Theorem 11.4); and conjugate mean uncertainty with an uninformative premium prior forces minimum variance (Proposition 11.5). The $1/N$ rule is also the exact limit of distributionally robust optimization as the ambiguity radius diverges (Pflug et al., 2012, C), and constraint-induced shrinkage (Jagannathan and Ma, 2003, C) plays the analogous role for minimum variance. Comparable foundations for ERC and HRP are, to our knowledge, open; we say so explicitly rather than assert them.

Level 4: implementation and statistical diagnostics. The population theory is extended to mandate constraints and trading frictions. For any closed convex feasible set, exact optimality becomes a normal-cone implied-return condition: residual alphas must be shadow prices of active constraints (Theorem 13.1); with quadratic transaction costs, implied returns acquire an explicit turnover wedge (Proposition 13.4). We then derive the first differential of the tangency map (Theorem 14.1), showing exactly why mean errors dominate covariance errors, convert every optimality condition into a projection diagnostic against its implied-return ray (Proposition 14.3), state the usable GRS formula, and add the high-dimensional random-matrix inversion tax for

plug-in GMV (Proposition 14.6). Finally, we isolate the exact place where the elliptical transfer fails: nonconstant standardized tail shape adds a first-order downside-risk wedge.

Level 5: HPO as the static policy face of RLPO. The final layer connects the static theory to Reinforcement Learning Portfolio Optimization (RLPO). An HPO rule Φ is a statewise portfolio map; in a Markov decision problem it therefore induces the deterministic stationary policy $\pi_\Phi(\cdot | s) = \delta_{\Phi(s)}$. When the discount factor is zero, trading frictions are absent, and the one-period reward is the same static objective used by HPO, the Bellman problem collapses exactly to the HPO problem. For $\gamma > 0$, state-dependent costs, regime transitions, and feedback effects, RLPO is the dynamic policy-improvement operator over the HPO prior. The paper formalizes this bridge in Section 9: the HPO implied-return defect becomes a reward-shaping penalty, RA-HRP becomes a hierarchical actor prior through node-level branch probabilities, and dynamic deviations from HPO are justified precisely when continuation value exceeds myopic HPO inefficiency and turnover costs. A performance-difference identity (Proposition 9.7) prices the total value gap of the myopic policy as an occupancy-weighted HPO advantage with an $\varepsilon/(1 - \gamma)$ myopia bound (Corollary 9.8), and the nodewise alphas of the homotopy calculus reappear as the myopic policy-gradient coordinates of the hierarchical actor at its prior (Corollary 9.10).

Claim tiering. Following the discipline used throughout our research program, every result is labeled. **P** marks results proved in this paper with self-contained proofs; **C** marks results proved in the cited literature and stated without proof; **E** marks empirical regularities reported in cited work. We make no conjectural claims and report no numerics of our own.

2 Scope, Standing Assumptions, and Paper Map

This section fixes the boundary of the results. The paper does not claim that any single heuristic dominates out of sample, nor that a neural policy can replace portfolio theory. It claims something narrower and stronger: for each information-restricted allocation map, one can write the exact population condition under which the rule is optimal, the exact geometric loss when it is not, the statistical diagnostics that test the condition, and the dynamic Bellman inequality that decides whether an RL policy should deviate from it.

2.1 Standing assumptions

Unless explicitly stated otherwise, the paper uses the following standing assumptions.

- A1. Finite asset universe.** There are $N < \infty$ risky assets and a riskless asset with rate r_f . Expected excess returns are $\mu_e = \mu - r_f \mathbf{1}$ and the covariance matrix satisfies $\Sigma \succ 0$.
- A2. Population versus implementation.** Population statements use the true pair (μ_e, Σ) . Estimated rules replace this pair by statewise estimates $(\hat{\mu}_{e,t}, \hat{\Sigma}_t)$ and are analyzed separately through sampling, turnover, and diagnostic terms.
- A3. Feasibility.** The baseline tangency and mean–variance results are stated on the fully invested affine hyperplane. Long-only constraints, caps, benchmark bands, and transaction-cost wedges are introduced explicitly in Section 13 through normal cones and cost-adjusted implied returns.
- A4. Rule versus family.** A fixed heuristic is a map Φ from a reduced information statistic to the simplex. A tunable heuristic family $\{\Phi_\lambda\}$ is selected by projecting the full optimizer onto a lower-complexity image set, not by pretending that the heuristic image set equals the full feasible set.

A5. Tree stability. HRP, RA-HRP, and Schur-RA-HRP are first analyzed for a fixed tree. Rolling-window implementations with re-estimated dendrograms add a discrete topology layer; the relevant quantities are merge preservation, cophenetic perturbation, and pathwise split distortion.

A6. Claim tiering. A label **P** in a result header denotes a result proved here, **C** denotes a cited theorem or classical result used without reproving it, and **E** denotes an empirical regularity reported in the literature. Open problems are explicitly marked as open.

2.2 Contribution map

Layer	Question answered	Mathematical object produced
Static optimality	When is a named heuristic exactly Markowitz/tangency optimal?	Implied-return sets $\{\boldsymbol{\mu}_e = a\boldsymbol{\Sigma}\mathbf{w}\}$ and rule-specific coincidence restrictions.
Approximate efficiency	If exact optimality is generically false, how costly is misspecification?	Angular Sharpe identity and second-order efficiency expansion around each coincidence ray.
Hierarchical HPO	What information does HRP delete, and what does RA-HRP restore?	Schur-bisection identity, score-tree kernels, pathwise/KL split attribution, and Schur-RA-HRP conditional-risk scores.
Statistical implementation	Why can a biased heuristic beat a plug-in optimizer?	Exact mean–variance bias–variance identity, tangency-map differential, GRS/Wald diagnostics, and high-dimensional inversion tax.
Dynamic policy improvement	How should a learned policy use a heuristic prior?	HPO-induced stationary policies, deviation inequality, performance-difference identity, and nodewise hierarchical actor gradients.

Table 1: Contribution map: each layer converts a practitioner question into a theorem-level object.

2.3 What the theory deliberately does not assert

The framework is falsifiable. A heuristic can fail its zero-alpha or implied-return condition, exhibit a large angular loss, generate excessive turnover after realistic costs, or break down under non-elliptical tail-shape heterogeneity. A learned RLPO policy can also fail if its continuation-value gain is not large enough to pay for added HPO defect and trading frictions. These are not exceptions to the theory; they are exactly the diagnostics the theory prescribes.

3 Setup, the Implied-Return Principle, and the Geometry of Optimality

3.1 Notation

There are N risky assets with return vector $\mathbf{r}$, mean $\boldsymbol{\mu} = \mathbb{E}[\mathbf{r}]$, and covariance matrix $\boldsymbol{\Sigma} \succ 0$. A riskless rate r_f is available; $\boldsymbol{\mu}_e = \boldsymbol{\mu} - r_f \mathbf{1}$ denotes expected excess returns, assumed nonzero. Volatilities are $\sigma_i = \sqrt{\Sigma_{ii}}$, $D = \text{diag}(\sigma_1, \dots, \sigma_N)$, and the correlation matrix is $R = D^{-1}\boldsymbol{\Sigma}D^{-1}$, so $\boldsymbol{\Sigma} = DRD$. Sharpe ratios are $S_i = \mu_{e,i}/\sigma_i$. A portfolio is $\mathbf{w} \in \mathbb{R}^N$, fully invested if $\mathbf{1}^\top \mathbf{w} = 1$. Portfolio variance is $\sigma_p^2(\mathbf{w}) = \mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}$, the Sharpe ratio is $S(\mathbf{w}) = \mathbf{w}^\top \boldsymbol{\mu}_e / \sigma_p(\mathbf{w})$, and $\langle \mathbf{u}, \mathbf{v} \rangle_{\boldsymbol{\Sigma}} = \mathbf{u}^\top \boldsymbol{\Sigma} \mathbf{v}$, $\|\mathbf{u}\|_{\boldsymbol{\Sigma}} = \langle \mathbf{u}, \mathbf{u} \rangle_{\boldsymbol{\Sigma}}^{1/2}$ denote the $\boldsymbol{\Sigma}$ -inner product and norm.

Definition 3.1 (Tangency coincidence). Assume $\mathbf{1}^\top \Sigma^{-1} \mu_e > 0$. The tangency (maximum-Sharpe) portfolio is

$$\mathbf{w}_T = \frac{\Sigma^{-1} \mu_e}{\mathbf{1}^\top \Sigma^{-1} \mu_e}, \quad S_{\max} := S(\mathbf{w}_T) = \sqrt{\mu_e^\top \Sigma^{-1} \mu_e}. \quad (3.1)$$

A fully invested $\mathbf{w}^\circ$ coincides with the tangency portfolio if $\mathbf{w}^\circ = \mathbf{w}_T$. This is the canonical notion of optimality in this paper: with a riskless asset, every mean–variance investor holds cash plus $\mathbf{w}_T$, so a heuristic equal to $\mathbf{w}_T$ is optimal for all risk aversions simultaneously.

Definition 3.2 (Frontier membership). A fully invested $\mathbf{w}^\circ$ is frontier-efficient if it solves $\max_{\mathbf{w}} \{\mathbf{w}^\top \mu - \frac{\lambda}{2} \mathbf{w}^\top \Sigma \mathbf{w} : \mathbf{1}^\top \mathbf{w} = 1\}$ for some $\lambda > 0$.

3.2 The first-order characterization

Theorem 3.3 (Implied-return principle; **P**). Let $\mathbf{w}^\circ$ be fully invested.

(i) $\mathbf{w}^\circ = \mathbf{w}_T$ if and only if there exists $a > 0$ such that

$$\mu_e = a \Sigma \mathbf{w}^\circ. \quad (3.2)$$

(ii) $\mathbf{w}^\circ$ is frontier-efficient if and only if there exist $a > 0$ and $b \in \mathbb{R}$ such that

$$\mu = a \Sigma \mathbf{w}^\circ + b \mathbf{1}. \quad (3.3)$$

Proof. (i) If $\mu_e = a \Sigma \mathbf{w}^\circ$ with $a > 0$, then $\Sigma^{-1} \mu_e = a \mathbf{w}^\circ$, and normalizing by $\mathbf{1}^\top \Sigma^{-1} \mu_e = a > 0$ (using $\mathbf{1}^\top \mathbf{w}^\circ = 1$) gives $\mathbf{w}_T = \mathbf{w}^\circ$. Conversely, if $\mathbf{w}^\circ = \mathbf{w}_T$ then $\mu_e = (\mathbf{1}^\top \Sigma^{-1} \mu_e) \Sigma \mathbf{w}^\circ$ with positive scalar.

(ii) The program in Definition 3.2 is strictly concave with one affine constraint; the Karush–Kuhn–Tucker conditions $\mu - \lambda \Sigma \mathbf{w} - \gamma \mathbf{1} = 0$, $\mathbf{1}^\top \mathbf{w} = 1$ are necessary and sufficient. Identify $a = \lambda$, $b = \gamma$. $\square$

Corollary 3.4 (Reverse optimization; **P**, classical). Every fully invested $\mathbf{w}^\circ$ with $\mathbf{w}^{\circ\top} \Sigma \mathbf{w}^\circ > 0$ is the tangency portfolio of the implied excess returns $\mu_e^{\text{imp}} = \Sigma \mathbf{w}^\circ$ (up to positive scaling), and frontier-efficient for $\Sigma \mathbf{w}^\circ + b \mathbf{1}$, any b .

This is the reverse-optimization step of [Black and Litterman \(1992\)](#) and the sensitivity logic of [Best and Grauer \(1991\)](#). Its consequence is worth stating bluntly: *no heuristic can be unconditionally suboptimal*. The substantive question is the shape of the optimality set, because adopting the rule $\mathbf{w}^\circ$ is behaviorally equivalent to asserting $(\mu_e, \Sigma) \in \mathcal{O}(\mathbf{w}^\circ)$.

Remark 3.5 (Beta form). Condition (3.2) says expected excess returns are proportional to covariances with the candidate: $\mu_{e,i} = a \text{Cov}(r_i, \mathbf{w}^{\circ\top} \mathbf{r})$. Dividing by the candidate’s own excess return gives the exact one-factor pricing relation

$$\mu_{e,i} = \beta_i(\mathbf{w}^\circ) (\mathbf{w}^{\circ\top} \mu_e), \quad \beta_i(\mathbf{w}^\circ) = \frac{\text{Cov}(r_i, \mathbf{w}^{\circ\top} \mathbf{r})}{\text{Var}(\mathbf{w}^{\circ\top} \mathbf{r})}. \quad (3.4)$$

A heuristic is maximum-Sharpe iff no asset has alpha against it — it prices the cross-section as if it were the market portfolio of a one-factor CAPM ([Sharpe, 1964](#)). All conditions below specialize (3.4).

Remark 3.6 (Testability; **C**). Relation (3.4) is the null hypothesis $\alpha = 0$ in the system of time-series regressions $r_{e,i,t} = \alpha_i + \beta_i r_{e,t}^\circ + \epsilon_{i,t}$ where $r_{e,t}^\circ$ is the heuristic’s realized excess return. Hence every optimality condition in this paper is directly testable by the finite-sample F -statistic of [Gibbons et al. \(1989\)](#) for the mean–variance efficiency of a given portfolio: “is equal weight

optimal in this universe” is a GRS test with the equal-weight portfolio as the candidate factor, and identically for IV, ERC, MD, GMV, HRP, and any fixed realization of RA-HRP. If the RA-HRP weights are estimated from the same return panel used in the test, the clean interpretation requires a split-sample or out-of-sample implementation; conditionally on the realized candidate weights, the GRS null is still the same zero-alpha restriction. The asymptotic and finite-sample distribution theory is otherwise standard.

3.3 Efficiency is an angle

Coincidence conditions classify the boundary; the angular identity measures everything else.

Theorem 3.7 (Angular efficiency identity; **P**). *For any portfolio $\mathbf{w} \neq 0$,*

$$S(\mathbf{w}) = S_{\max} \cos \theta_{\Sigma}(\mathbf{w}, \mathbf{w}_T), \quad \cos \theta_{\Sigma}(\mathbf{w}, \mathbf{w}_T) := \frac{\langle \mathbf{w}, \mathbf{w}_T \rangle_{\Sigma}}{\|\mathbf{w}\|_{\Sigma} \|\mathbf{w}_T\|_{\Sigma}}. \quad (3.5)$$

In particular the Sharpe efficiency $\eta(\mathbf{w}) := S(\mathbf{w})/S_{\max} = \cos \theta_{\Sigma}$, and $1 - \eta(\mathbf{w})^2 = \sin^2 \theta_{\Sigma}$ exactly. A portfolio attains the maximum Sharpe ratio iff it is Σ -parallel to $\Sigma^{-1}\boldsymbol{\mu}_e$, recovering Theorem 3.3(i).

Proof. Let $\tilde{\mathbf{w}} = \Sigma^{-1}\boldsymbol{\mu}_e$ (unnormalized tangency direction). Then $\mathbf{w}^{\top}\boldsymbol{\mu}_e = \mathbf{w}^{\top}\Sigma\tilde{\mathbf{w}} = \langle \mathbf{w}, \tilde{\mathbf{w}} \rangle_{\Sigma}$ and $\|\tilde{\mathbf{w}}\|_{\Sigma}^2 = \boldsymbol{\mu}_e^{\top}\Sigma^{-1}\boldsymbol{\mu}_e = S_{\max}^2$. Hence

$$S(\mathbf{w}) = \frac{\langle \mathbf{w}, \tilde{\mathbf{w}} \rangle_{\Sigma}}{\|\mathbf{w}\|_{\Sigma}} = \|\tilde{\mathbf{w}}\|_{\Sigma} \frac{\langle \mathbf{w}, \tilde{\mathbf{w}} \rangle_{\Sigma}}{\|\mathbf{w}\|_{\Sigma} \|\tilde{\mathbf{w}}\|_{\Sigma}} = S_{\max} \cos \theta_{\Sigma}(\mathbf{w}, \tilde{\mathbf{w}}),$$

and $\theta_{\Sigma}(\mathbf{w}, \tilde{\mathbf{w}}) = \theta_{\Sigma}(\mathbf{w}, \mathbf{w}_T)$ since $\mathbf{w}_T \propto \tilde{\mathbf{w}}$ with positive constant. $\square$

The identity converts every question about heuristic performance into a question about an angle in the metric defined by risk itself. Two structural consequences follow: optimality sets are thin, and the efficiency surface is flat across them.

Proposition 3.8 (Genericity: exact optimality is null; **P**). *Fix $\Sigma \succ 0$ and a fully invested $\mathbf{w}^{\circ}$. The tangency-coincidence set $\mathcal{O}_{\Sigma}(\mathbf{w}^{\circ}) = \{\boldsymbol{\mu}_e : \boldsymbol{\mu}_e = a\Sigma\mathbf{w}^{\circ}, a > 0\}$ is an open ray — a one-dimensional, Lebesgue-null subset of $\mathbb{R}^N$ for $N \geq 2$. The frontier set $\{\boldsymbol{\mu} : \boldsymbol{\mu} = a\Sigma\mathbf{w}^{\circ} + b\mathbf{1}, a > 0\}$ is an open half-plane, null for $N \geq 3$. Consequently, under any belief about $\boldsymbol{\mu}_e$ (or $\boldsymbol{\mu}$) absolutely continuous with respect to Lebesgue measure, the probability that a fixed heuristic is exactly maximum-Sharpe (exactly frontier-efficient) is zero.*

Proof. The sets are images of $(0, \infty)$ and $(0, \infty) \times \mathbb{R}$ under injective affine maps, hence smooth manifolds of dimension 1 and 2; sets of dimension $< N$ have Lebesgue measure zero, and null sets have probability zero under absolutely continuous laws. $\square$

Theorem 3.9 (Second-order robustness; **P**). *Fix $\Sigma \succ 0$, a fully invested $\mathbf{w}^{\circ}$ with $A := \mathbf{w}^{\circ\top}\Sigma\mathbf{w}^{\circ} > 0$, and a point on the coincidence ray $\boldsymbol{\mu}_e^0 = a\Sigma\mathbf{w}^{\circ}$, $a > 0$. Perturb in an arbitrary direction $v \in \mathbb{R}^N$: $\boldsymbol{\mu}_e(\varepsilon) = a\Sigma\mathbf{w}^{\circ} + \varepsilon v$. Then the squared Sharpe efficiency of $\mathbf{w}^{\circ}$ satisfies, exactly,*

$$\eta^2(\varepsilon) = \frac{(aA + \varepsilon b)^2}{A(a^2A + 2a\varepsilon b + \varepsilon^2 c)}, \quad b := v^{\top}\mathbf{w}^{\circ}, \quad c := v^{\top}\Sigma^{-1}v, \quad (3.6)$$

and therefore

$$1 - \eta^2(\varepsilon) = \frac{\varepsilon^2}{a^2} \cdot \frac{cA - b^2}{A^2} + O(\varepsilon^3), \quad cA - b^2 \geq 0, \quad (3.7)$$

with $cA - b^2 = 0$ iff v is itself on the ray ($v \propto \Sigma\mathbf{w}^{\circ}$). In particular $\frac{d}{d\varepsilon}\eta^2|_{\varepsilon=0} = 0$ for every direction v : the efficiency map is stationary on the coincidence set, and the leading loss is quadratic with coefficient equal to the Σ^{-1} -residual of v after projection on the ray.

Proof. $S(\mathbf{w}^\circ)^2 = (\mathbf{w}^\circ{}^\top \boldsymbol{\mu}_e)^2/A = (aA + \varepsilon b)^2/A$ and $S_{\max}^2 = \boldsymbol{\mu}_e^\top \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_e = a^2 A + 2a\varepsilon b + \varepsilon^2 c$, giving (3.6). Expanding,

$$\eta^2 = \frac{a^2 A^2 + 2a\varepsilon bA + \varepsilon^2 b^2}{a^2 A^2 + 2a\varepsilon bA + \varepsilon^2 cA} = 1 - \varepsilon^2 \frac{cA - b^2}{a^2 A^2} + O(\varepsilon^3).$$

Nonnegativity of $cA - b^2$ is Cauchy–Schwarz in the $\boldsymbol{\Sigma}^{-1}$ -inner product applied to v and $\boldsymbol{\Sigma} \mathbf{w}^\circ$: $\langle v, \boldsymbol{\Sigma} \mathbf{w}^\circ \rangle_{\boldsymbol{\Sigma}^{-1}} = v^\top \mathbf{w}^\circ = b$ and $\|\boldsymbol{\Sigma} \mathbf{w}^\circ\|_{\boldsymbol{\Sigma}^{-1}}^2 = A$, so $b^2 \leq cA$ with equality iff $v \parallel \boldsymbol{\Sigma} \mathbf{w}^\circ$. $\square$

Remark 3.10 (Never optimal, rarely costly). *Proposition 3.8 and Theorem 3.9 jointly formalize the folklore. Exact optimality of any fixed rule is a probability-zero event; but because the efficiency surface has a vanishing gradient on the entire coincidence ray, an order- ε misspecification of premia costs only order- ε^2 in squared Sharpe efficiency. The same flatness, viewed from the optimizer’s side, is the source of mean–variance instability — the optimum moves at order ε while the objective moves at order ε^2 (Best and Grauer, 1991; Michaud, 1989; Chopra and Ziemba, 1993) — so a single geometric fact explains simultaneously why heuristics are cheap and why optimizers are fragile.*

3.4 The catalogue

Definition 3.11 (Allocation rules). *All rules are fully invested and long-only by construction.*

$$\text{Equal weight (EW): } w_i = 1/N. \quad (3.8)$$

$$\text{Inverse volatility (IV): } w_i = \sigma_i^{-1} / \sum_j \sigma_j^{-1}. \quad (3.9)$$

$$\text{Inverse variance (IVar): } w_i = \sigma_i^{-2} / \sum_j \sigma_j^{-2}. \quad (3.10)$$

$$\text{Global minimum variance (GMV): } \mathbf{w}_{\text{GMV}} = \boldsymbol{\Sigma}^{-1} \mathbf{1} / \mathbf{1}^\top \boldsymbol{\Sigma}^{-1} \mathbf{1}. \quad (3.11)$$

$$\text{Equal risk contribution (ERC): } w_i (\boldsymbol{\Sigma} \mathbf{w})_i = \frac{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}}{N} \quad \forall i, \quad \mathbf{w} > 0, \quad \mathbf{1}^\top \mathbf{w} = 1. \quad (3.12)$$

$$\text{Maximum diversification (MD): } \mathbf{w}_{\text{MD}} = \arg \max_{\mathbf{1}^\top \mathbf{w} = 1} \frac{\mathbf{w}^\top \boldsymbol{\sigma}}{\sqrt{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}}}. \quad (3.13)$$

$$\text{Hierarchical risk parity (HRP): } \text{Definition 7.1 below.} \quad (3.14)$$

The MD portfolio satisfies $\mathbf{w}_{\text{MD}} \propto \boldsymbol{\Sigma}^{-1} \boldsymbol{\sigma}$ by the argument of Theorem 3.3(i) with $\boldsymbol{\sigma}$ in place of $\boldsymbol{\mu}_e$; when $\boldsymbol{\Sigma}^{-1} \boldsymbol{\sigma}$ has negative entries the long-only MD portfolio differs, and our statements concern the unconstrained solution. Existence and uniqueness of ERC is Theorem 6.1 below.

4 Exact Conditions: Equal Weight, Inverse Volatility, Minimum Variance, Maximum Diversification

4.1 Equal weight

Theorem 4.1 (P). *The equal-weight portfolio coincides with the tangency portfolio if and only if*

$$\boldsymbol{\mu}_e \propto \boldsymbol{\Sigma} \mathbf{1} \quad (\text{positive constant}), \quad (4.1)$$

i.e. iff every asset’s expected excess return is proportional to its covariance with the equal-weight portfolio — equivalently, every asset has zero alpha in (3.4) with $\mathbf{w}^\circ = \mathbf{1}/N$. It is frontier-efficient iff $\boldsymbol{\mu} \in \{a\boldsymbol{\Sigma} \mathbf{1} + b\mathbf{1} : a > 0, b \in \mathbb{R}\}$.

Proof. Theorem 3.3 with $\mathbf{w}^\circ = \mathbf{1}/N$. For the beta form: with $\boldsymbol{\mu}_e = c\boldsymbol{\Sigma} \mathbf{1}$, $c > 0$, $\beta_i \cdot (\mathbf{1}^\top \boldsymbol{\mu}_e / N) = \frac{N(\boldsymbol{\Sigma} \mathbf{1})_i}{\mathbf{1}^\top \boldsymbol{\Sigma} \mathbf{1}} \cdot \frac{c\mathbf{1}^\top \boldsymbol{\Sigma} \mathbf{1}}{N} = c(\boldsymbol{\Sigma} \mathbf{1})_i = \mu_{e,i}$. $\square$

Corollary 4.2 (Symmetry suffices; **P**). If $\boldsymbol{\mu}_e = m\mathbf{1}$ with $m > 0$ and $\boldsymbol{\Sigma}\mathbf{1} = c\mathbf{1}$ (equal row sums; e.g. equal volatilities with constant correlation, or any $(\boldsymbol{\mu}_e, \boldsymbol{\Sigma})$ invariant under all permutations of assets), then EW is the tangency portfolio.

Proof. $\boldsymbol{\mu}_e = m\mathbf{1} = (m/c)\boldsymbol{\Sigma}\mathbf{1}$, $c = \mathbf{1}^\top \boldsymbol{\Sigma}\mathbf{1}/N > 0$. Permutation invariance forces both hypotheses. $\square$

Note what is *not* required: volatilities need not be equal and correlations need not be constant. The exact requirement couples means to covariance row sums; an asset may carry twice the covariance with the market of another provided it also carries twice the risk premium. Section 11 upgrades the symmetry observation from a sufficient condition on parameters to a Bayes-optimality theorem on beliefs.

4.2 Inverse volatility

Theorem 4.3 (**P**). Write $\boldsymbol{\Sigma} = DRD$. The inverse-volatility portfolio coincides with the tangency portfolio if and only if

$$S_i \propto (R\mathbf{1})_i = \sum_{j=1}^N \rho_{ij} \quad \forall i \quad (\text{positive constant}) : \quad (4.2)$$

each asset's Sharpe ratio must be proportional to the row sum of its correlations.

Proof. $\mathbf{w}_{IV} \propto D^{-1}\mathbf{1}$; coincidence iff $\boldsymbol{\mu}_e \propto \boldsymbol{\Sigma}D^{-1}\mathbf{1} = DRD D^{-1}\mathbf{1} = DR\mathbf{1}$, i.e. $\mu_{e,i} \propto \sigma_i(R\mathbf{1})_i$, i.e. $S_i \propto (R\mathbf{1})_i$, positive constants throughout. $\square$

Corollary 4.4 (**P**). If $S_i \equiv S > 0$ and R has equal row sums, $R\mathbf{1} = \kappa\mathbf{1}$ — in particular under constant pairwise correlation $\rho > -1/(N-1)$, where $\kappa = 1 + (N-1)\rho > 0$ — then IV is the tangency portfolio.

4.3 Minimum variance

Theorem 4.5 (**P**). The GMV portfolio coincides with the tangency portfolio if and only if expected excess returns are equal across assets: $\boldsymbol{\mu}_e = m\mathbf{1}$, $m > 0$.

Proof. $\boldsymbol{\Sigma}^{-1}\boldsymbol{\mu}_e \propto \boldsymbol{\Sigma}^{-1}\mathbf{1}$ (positive constant) iff $\boldsymbol{\mu}_e \propto \mathbf{1}$, since $\boldsymbol{\Sigma}^{-1}$ is a bijection. $\square$

Remark 4.6. GMV is always frontier-efficient — the vertex of the frontier — so the content of Theorem 4.5 concerns coincidence with the portfolio every investor holds when a riskless asset exists. The equal-premium condition is the formal statement of the agnosticism motivating minimum-variance investing (Clarke et al., 2011): GMV is the optimizer of an investor who believes risk is forecastable and premia are not. Proposition 11.5 makes the belief statement exact.

4.4 Maximum diversification

Theorem 4.7 (**P**). The (unconstrained) maximum-diversification portfolio coincides with the tangency portfolio if and only if all Sharpe ratios are equal: $\boldsymbol{\mu}_e \propto \boldsymbol{\sigma}$, positive constant. No restriction on R is required.

Proof. $\mathbf{w}_{MD} \propto \boldsymbol{\Sigma}^{-1}\boldsymbol{\sigma}$, $\mathbf{w}_T \propto \boldsymbol{\Sigma}^{-1}\boldsymbol{\mu}_e$; equality up to positive scaling iff $\boldsymbol{\mu}_e \propto \boldsymbol{\sigma}$. At the level of objectives: if $\boldsymbol{\mu}_e = S\boldsymbol{\sigma}$ then $S(\mathbf{w}) = S \cdot \mathbf{w}^\top \boldsymbol{\sigma} / \sigma_p(\mathbf{w})$, so the Sharpe functional and the diversification ratio are positive multiples and share a maximizer. $\square$

This was the design premise of Choueifaty and Coignard (2008): MD is the bet that risk is priced uniformly per unit of volatility. Comparing Theorems 4.3 and 4.7 quantifies the gap between IV and MD: both encode equal Sharpe ratios, but IV additionally requires equal correlation row sums, because it ignores the correlation structure MD inverts.

5 One-Factor Structure: Conditions in Observables

The conditions above restrict the unobservable pair $(\boldsymbol{\mu}_e, \boldsymbol{\Sigma})$. Under one-factor structure they become statements about betas and idiosyncratic variances — quantities the practitioner actually estimates.

Assumption 5.1 (Priced one-factor model). $r_{e,i} = \beta_i f_e + u_i$ with $\mathbb{E}[f_e] = \lambda_m > 0$, $\text{Var}(f_e) = \sigma_m^2$, $\mathbb{E}[u_i] = 0$, $\text{Cov}(u_i, u_j) = \delta_i^2 \mathbf{1}\{i = j\}$, $\text{Cov}(f_e, u_i) = 0$. Hence $\boldsymbol{\mu}_e = \lambda_m \boldsymbol{\beta}$ and $\boldsymbol{\Sigma} = \sigma_m^2 \boldsymbol{\beta} \boldsymbol{\beta}^\top + \Delta$, $\Delta = \text{diag}(\delta_1^2, \dots, \delta_N^2) \succ 0$. Exact pricing ($\boldsymbol{\alpha} = 0$) is part of the assumption.

Lemma 5.2 (Sherman–Morrison; $\mathbf{P}$). Under Assumption 5.1, with $q := \boldsymbol{\beta}^\top \Delta^{-1} \boldsymbol{\beta}$,

$$\boldsymbol{\Sigma}^{-1} = \Delta^{-1} - \frac{\sigma_m^2 \Delta^{-1} \boldsymbol{\beta} \boldsymbol{\beta}^\top \Delta^{-1}}{1 + \sigma_m^2 q}, \quad \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta} = \frac{\Delta^{-1} \boldsymbol{\beta}}{1 + \sigma_m^2 q}.$$

Proof. The first display is the Sherman–Morrison identity for the rank-one update $\boldsymbol{\Sigma} = \Delta + (\sigma_m \boldsymbol{\beta})(\sigma_m \boldsymbol{\beta})^\top$. Applying it to $\boldsymbol{\beta}$: $\boldsymbol{\Sigma}^{-1} \boldsymbol{\beta} = \Delta^{-1} \boldsymbol{\beta} - \sigma_m^2 \Delta^{-1} \boldsymbol{\beta} q / (1 + \sigma_m^2 q) = \Delta^{-1} \boldsymbol{\beta} / (1 + \sigma_m^2 q)$. $\square$

Proposition 5.3 (Tangency and GMV in observables; $\mathbf{P}$; GMV form after [Clarke et al., \(2011\)](#)). Under Assumption 5.1:

- (i) The tangency portfolio is $\mathbf{w}_T \propto \Delta^{-1} \boldsymbol{\beta}$, i.e. $w_{T,i} \propto \beta_i / \delta_i^2$.
- (ii) The GMV portfolio admits the threshold-beta representation

$$w_{\text{GMV},i} \propto \frac{1}{\delta_i^2} \left(1 - \frac{\beta_i}{\beta_L}\right), \quad \beta_L = \frac{1 + \sigma_m^2 \boldsymbol{\beta}^\top \Delta^{-1} \boldsymbol{\beta}}{\sigma_m^2 \boldsymbol{\beta}^\top \Delta^{-1} \mathbf{1}},$$

(assuming $\boldsymbol{\beta}^\top \Delta^{-1} \mathbf{1} > 0$): assets with $\beta_i > \beta_L$ are shorted, and long-only GMV concentrates in low-beta, low-idiosyncratic-risk names.

- (iii) Coincidence conditions in observables: EW is tangency iff $\beta_i \propto \delta_i^2$; IVar is tangency iff $\beta_i \propto \delta_i^2 / \sigma_i^2$; IV is tangency iff $\beta_i \propto \delta_i^2 / \sigma_i$; GMV is tangency iff β_i is constant (consistent with (ii): constant betas give $\mathbf{w}_{\text{GMV}} \propto \Delta^{-1} \mathbf{1} \propto \mathbf{w}_T$); MD is tangency iff $\beta_i \propto \sigma_i$, i.e. iff all assets have the same correlation with the factor, $\text{Corr}(r_i, f) = \beta_i \sigma_m / \sigma_i$ constant.

Proof. (i) $\mathbf{w}_T \propto \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_e = \lambda_m \boldsymbol{\Sigma}^{-1} \boldsymbol{\beta} \propto \Delta^{-1} \boldsymbol{\beta}$ by Lemma 5.2. (ii) Apply Lemma 5.2 to $\mathbf{1}$:

$$\boldsymbol{\Sigma}^{-1} \mathbf{1} = \Delta^{-1} \mathbf{1} - \frac{\sigma_m^2 (\boldsymbol{\beta}^\top \Delta^{-1} \mathbf{1})}{1 + \sigma_m^2 q} \Delta^{-1} \boldsymbol{\beta} \implies (\boldsymbol{\Sigma}^{-1} \mathbf{1})_i = \frac{1}{\delta_i^2} \left(1 - \frac{\beta_i}{\beta_L}\right)$$

with β_L as displayed. (iii) Each statement is Theorem 3.3(i): the heuristic must be proportional to $\Delta^{-1} \boldsymbol{\beta}$. EW: $\mathbf{1} \propto \Delta^{-1} \boldsymbol{\beta} \iff \beta_i \propto \delta_i^2$. IVar: $D^{-2} \mathbf{1} \propto \Delta^{-1} \boldsymbol{\beta} \iff \beta_i \propto \delta_i^2 / \sigma_i^2$. IV: $D^{-1} \mathbf{1} \propto \Delta^{-1} \boldsymbol{\beta} \iff \beta_i \propto \delta_i^2 / \sigma_i$. GMV: $\boldsymbol{\mu}_e \propto \mathbf{1}$ (Theorem 4.5) iff β_i constant. MD: $\boldsymbol{\mu}_e \propto \boldsymbol{\sigma}$ (Theorem 4.7) iff $\lambda_m \beta_i \propto \sigma_i$. $\square$

Statement (iii) is the practical payoff: in a priced single-factor world, the question “which heuristic should be closest to optimal in this universe” reduces to inspecting the cross-sectional relation between β_i , δ_i^2 , and σ_i — no premium estimates required, because exact pricing has already collapsed $\boldsymbol{\mu}_e$ onto $\boldsymbol{\beta}$.

6 Risk Parity: Existence, Uniqueness, and Optimality as Performance Parity

Risk contributions $\text{RC}_i(\mathbf{w}) = w_i (\boldsymbol{\Sigma} \mathbf{w})_i$ satisfy the Euler identity $\sum_i \text{RC}_i = \sigma_p^2$, which underwrites their interpretation as a decomposition of risk ([Qian, 2006](#)). Define analogously the *performance contribution* $\text{PC}_i(\mathbf{w}) = w_i \mu_{e,i}$, with $\sum_i \text{PC}_i = \mathbf{w}^\top \boldsymbol{\mu}_e$.

6.1 Existence and uniqueness

Theorem 6.1 (**P**; cf. Maillard et al., 2010; Roncalli, 2013). *For every $\Sigma \succ 0$ there exists exactly one fully invested $\mathbf{w} > 0$ with equal risk contributions.*

Proof. Fix $c > 0$ and consider $F_c(\mathbf{w}) = \frac{1}{2}\mathbf{w}^\top \Sigma \mathbf{w} - c \sum_i \ln w_i$ on the open orthant $\mathbb{R}_{++}^N$. F_c is strictly convex ($\Sigma \succ 0$ plus convexity of $-\ln$) and coercive: $F_c \rightarrow \infty$ as $\|\mathbf{w}\| \rightarrow \infty$ (the quadratic dominates the logarithm) and as any $w_i \downarrow 0$ (the barrier diverges; on bounded sets the quadratic and the remaining logarithms are bounded below). Hence a unique minimizer $\mathbf{w}(c) \in \mathbb{R}_{++}^N$ exists, characterized by the first-order condition $\Sigma \mathbf{w} = c \mathbf{w}^{-1}$ (elementwise reciprocal), i.e. $w_i(\Sigma \mathbf{w})_i = c$ for all i : every barrier solution is an unnormalized ERC portfolio. The family is a single ray: if $\Sigma \mathbf{w} = c \mathbf{w}^{-1}$ then $\Sigma(t\mathbf{w}) = (t^2 c)(t\mathbf{w})^{-1}$, so $\mathbf{w}(t^2 c) = t \mathbf{w}(c)$. Normalizing to the simplex gives existence. For uniqueness, suppose $\mathbf{u}, \mathbf{v}$ are both fully invested ERC portfolios with constants $c_u = \mathbf{u}^\top \Sigma \mathbf{u}/N$, c_v . Then $\mathbf{u} = \mathbf{w}(c_u)$ and $\mathbf{v} = \mathbf{w}(c_v)$ by uniqueness of barrier minimizers, and by the ray property $\mathbf{v} = t \mathbf{u}$ with $t = \sqrt{c_v/c_u}$; full investment of both forces $t = 1$. $\square$

6.2 Optimality

Theorem 6.2 (ERC optimality = performance parity; **P**). *Let $\mathbf{w}$ be the ERC portfolio. Then $\mathbf{w}$ coincides with the tangency portfolio if and only if performance contributions are also equal:*

$$w_i \mu_{e,i} = \frac{\mathbf{w}^\top \boldsymbol{\mu}_e}{N} \quad \forall i. \quad (6.1)$$

Proof. Since $\mathbf{w}$ is ERC, $(\Sigma \mathbf{w})_i = \sigma_p^2/(Nw_i)$, $w_i > 0$. By Theorem 3.3(i), coincidence holds iff $\boldsymbol{\mu}_e = a \Sigma \mathbf{w}$, $a > 0$, i.e. iff $\mu_{e,i} = a \sigma_p^2/(Nw_i)$ for all i , i.e. iff $w_i \mu_{e,i}$ is constant; summing identifies the constant, and $a > 0$ iff $\mathbf{w}^\top \boldsymbol{\mu}_e > 0$. $\square$

The economic reading is sharp: a risk parity portfolio is mean–variance optimal precisely when the equal risk budget doubles as an equal *return* budget — each line contributes the same expected excess return in currency terms. When risk budgets and performance budgets disagree under the true $\boldsymbol{\mu}_e$, ERC is on neither the tangency ray nor, generically, the frontier.

Corollary 6.3 (Frontier version; **P**). *The ERC portfolio is frontier-efficient iff there exist $\kappa > 0$, $b \in \mathbb{R}$ with $w_i \mu_i = \kappa + b w_i$ for all i : performance contributions affine in weights, with the tangency case $b = r_f \cdot 1$ -adjusted to (6.1).*

Proof. Theorem 3.3(ii) with $(\Sigma \mathbf{w})_i = \sigma_p^2/(Nw_i)$: $\mu_i = a \sigma_p^2/(Nw_i) + b \iff w_i \mu_i = a \sigma_p^2/N + b w_i$. $\square$

Theorem 6.4 (Budgeted generalization; **P**). *Let $b_i > 0$, $\sum_i b_i = 1$, and let $\mathbf{w}$ satisfy $\text{RC}_i(\mathbf{w}) = b_i \sigma_p^2$. Then $\mathbf{w}$ coincides with the tangency portfolio iff $\text{PC}_i(\mathbf{w}) = b_i \mathbf{w}^\top \boldsymbol{\mu}_e$ for all i : performance must be budgeted in the same proportions as risk.*

Proof. As in Theorem 6.2 with σ_p^2/N replaced by $b_i \sigma_p^2$. $\square$

Proposition 6.5 (Constant correlation collapses ERC to IV; **P**, after Maillard et al., 2010). *If $R = (1 - \rho)I + \rho \mathbf{1}\mathbf{1}^\top$ with $\rho > -1/(N - 1)$, the ERC portfolio is the inverse-volatility portfolio.*

Proof. Take $w_i = \sigma_i^{-1}/T$, $T = \sum_j \sigma_j^{-1}$. Then $(\Sigma \mathbf{w})_i = \sigma_i \sum_j \rho_{ij} \sigma_j w_j = \frac{\sigma_i}{T} (1 + (N - 1)\rho)$, so $w_i(\Sigma \mathbf{w})_i = (1 + (N - 1)\rho)/T^2$ is constant. Conclude by Theorem 6.1. $\square$

Corollary 6.6 (Sufficient condition of Maillard et al., 2010; **P**). *Equal Sharpe ratios and constant pairwise correlation imply that ERC coincides with the tangency portfolio.*

Proof. Proposition 6.5 gives $\text{ERC} = \text{IV}$; Corollary 4.4 gives $\text{IV} = \text{tangency}$. $\square$

Remark 6.7 (The sufficient condition is far from necessary; **P**). By Corollary 3.4, take any $\Sigma \succ 0$ — in particular one with wildly non-constant correlations — compute its ERC portfolio $\mathbf{w}$, and set $\mu_e = \Sigma \mathbf{w}$. Then ERC is exactly the tangency portfolio while the Maillard et al. (2010) hypotheses fail. The exact optimality set is the performance-parity condition (6.1), a codimension- $(N - 1)$ restriction; equal-Sharpe-plus-constant-correlation is a thin, interpretable slice of it.

Proposition 6.8 (Variance ordering and interpolation; **C**). On the long-only simplex one has the bracketing $\sigma_p(\mathbf{w}_{\text{GMV}}^+) \leq \sigma_p(\mathbf{w}_{\text{ERC}}) \leq \sigma_p(\mathbf{w}_{\text{EW}})$, where $\mathbf{w}_{\text{GMV}}^+$ is long-only GMV. The proof of Maillard et al. (2010) runs through the variational problem of Theorem 6.1 in constrained form, $\min\{\sqrt{\mathbf{w}^\top \Sigma \mathbf{w}} : \sum_i \ln w_i \geq c, \mathbf{1}^\top \mathbf{w} = 1, \mathbf{w} \geq 0\}$: as c increases to its maximal feasible value $-N \ln N$ (attained only at EW, by strict concavity of $\sum_i \ln w_i$ on the simplex), the feasible set shrinks from the full simplex to $\{\mathbf{w}_{\text{EW}}\}$ and the minimized volatility increases monotonically; ERC arises at an intermediate level. See also Roncali (2013).

The interpolation reading anticipates Section 12: ERC sits between the rule that trusts Σ completely (GMV) and the rule that trusts nothing (EW), exactly as an estimator shrunk between a sample-efficient and a maximally robust target.

7 Hierarchical and Return-Adjusted Hierarchical Risk Parity

7.1 The algorithm

Definition 7.1 (HRP; López de Prado, 2016). Given Σ with correlation matrix R :

- (1) Tree clustering. Compute $d_{ij} = \sqrt{(1 - \rho_{ij})/2}$ and build a dendrogram $\mathcal{T}$ by hierarchical (in the original, single-linkage) clustering.
- (2) Quasi-diagonalization. Reorder assets so the dendrogram's leaves are contiguous.
- (3) Recursive bisection. Initialize $w_i = 1$. Recursively split each cluster C into A and B ; assign

$$\alpha_A = \frac{\tilde{V}_B}{\tilde{V}_A + \tilde{V}_B}, \quad \alpha_B = 1 - \alpha_A, \quad \tilde{V}_K = \tilde{\mathbf{w}}_K^\top \Sigma_K \tilde{\mathbf{w}}_K, \quad \tilde{w}_{K,i} = \frac{\sigma_i^{-2}}{\sum_{j \in K} \sigma_j^{-2}}, \quad (7.1)$$

multiply the running weights in A by α_A and in B by α_B , and recurse to singletons.

We call the bisection tree-aligned if every split occurs at a node of $\mathcal{T}$, and balanced if the two sub-clusters at every split have equal cardinality. (The original implementation splits the quasi-diagonalized list in half; common variants split at dendrogram nodes. Our exactness results hold for whichever scheme satisfies the stated hypotheses.)

7.2 What the optimizer actually does at a split

To see precisely which information HRP discards, write the minimum-variance solution of a two-block system in split form.

Theorem 7.2 (Exact Schur bisection identity; **P**). Consider a partition $C = A \cup B$ with $\Sigma_C = \begin{pmatrix} \Sigma_A & \Sigma_{AB} \\ \Sigma_{BA} & \Sigma_B \end{pmatrix} \succ 0$. Define the Schur complements and tilted budget vectors

$$\begin{aligned} \mathbf{S}_A &= \Sigma_A - \Sigma_{AB} \Sigma_B^{-1} \Sigma_{BA}, & \mathbf{S}_B &= \Sigma_B - \Sigma_{BA} \Sigma_A^{-1} \Sigma_{AB}, \\ \mathbf{b}_A &= \mathbf{1}_A - \Sigma_{AB} \Sigma_B^{-1} \mathbf{1}_B, & \mathbf{b}_B &= \mathbf{1}_B - \Sigma_{BA} \Sigma_A^{-1} \mathbf{1}_A. \end{aligned}$$

Then the unnormalized GMV vector splits exactly as

$$(\Sigma_C^{-1} \mathbf{1}_C)_A = \mathbf{S}_A^{-1} \mathbf{b}_A, \quad (\Sigma_C^{-1} \mathbf{1}_C)_B = \mathbf{S}_B^{-1} \mathbf{b}_B, \quad (7.2)$$

so the GMV mass allocated to A is $\mathbf{1}_A^\top \mathbf{S}_A^{-1} \mathbf{b}_A / (\mathbf{1}_A^\top \mathbf{S}_A^{-1} \mathbf{b}_A + \mathbf{1}_B^\top \mathbf{S}_B^{-1} \mathbf{b}_B)$ and the within- A composition is $\mathbf{S}_A^{-1} \mathbf{b}_A$ renormalized.

Proof. By the block-inverse formula for $\Sigma_C \succ 0$ (both Schur complements are positive definite),

$$(\Sigma_C^{-1})_{AA} = \mathbf{S}_A^{-1}, \quad (\Sigma_C^{-1})_{AB} = -\mathbf{S}_A^{-1} \Sigma_{AB} \Sigma_B^{-1},$$

hence $(\Sigma_C^{-1} \mathbf{1}_C)_A = \mathbf{S}_A^{-1} \mathbf{1}_A - \mathbf{S}_A^{-1} \Sigma_{AB} \Sigma_B^{-1} \mathbf{1}_B = \mathbf{S}_A^{-1} \mathbf{b}_A$; symmetrically for B . $\square$

Corollary 7.3 (The three substitutions; **P**). *At every split, HRP computes its allocation from the triple $(\Sigma_K, \mathbf{1}_K, \text{inverse-variance weights})$ where the exact recursion (7.2) requires $(\mathbf{S}_K, \mathbf{b}_K, \text{minimum-variance weights})$. The algorithm therefore commits exactly three substitutions: (i) raw block for Schur complement, $\Sigma_K \leftarrow \mathbf{S}_K$ — discarding the variance reduction available from cross-hedging against the sibling cluster; (ii) flat budget for tilted budget, $\mathbf{1}_K \leftarrow \mathbf{b}_K$ — discarding the demand tilt induced by cross-cluster covariance; (iii) inverse variance for within-cluster minimum variance — discarding within-cluster correlations in the composition and in the cluster-variance functional used by the split rule. All three are exact when $\Sigma_{AB} = 0$ and the within-cluster IV portfolio is the within-cluster GMV; the theorems below identify when that happens, and Cotton (2024) constructs the one-parameter family that reinstates (i)–(ii) continuously, interpolating between HRP and exact GMV, with the recursive Schur reduction developed further by Mograby (2025).*

7.3 Exact coincidence results

Lemma 7.4 (Harmonic aggregation; **P**). *If Σ_K is diagonal, the cluster variance in (7.1) equals $\tilde{V}_K = (\sum_{i \in K} \sigma_i^{-2})^{-1} = 1/T_K$, $T_K := \sum_{i \in K} \sigma_i^{-2}$, and the inverse-variance weights are exactly the GMV weights of Σ_K , with $\tilde{V}_K = (\mathbf{1}^\top \Sigma_K^{-1} \mathbf{1})^{-1}$.*

Proof. $\tilde{V}_K = \sum_{i \in K} \tilde{w}_{K,i}^2 \sigma_i^2 = \sum_{i \in K} \sigma_i^{-2} / T_K^2 = 1/T_K$; and $\Sigma_K^{-1} \mathbf{1} = (\sigma_i^{-2})_{i \in K}$ gives the rest. $\square$

Theorem 7.5 (Diagonal exactness; **P**). *If Σ is diagonal, then for every dendrogram and every bisection scheme HRP returns the inverse-variance portfolio, which is the GMV portfolio: $w_i^{\text{HRP}} = \sigma_i^{-2} / \sum_j \sigma_j^{-2} = w_i^{\text{GMV}}$.*

Proof. By Lemma 7.4, at any split of C into A, B , $\alpha_A = \frac{1/T_B}{1/T_A + 1/T_B} = \frac{T_A}{T_A + T_B} = \frac{T_A}{T_C}$. Asset i 's final weight is the product of fractions along its root-to-leaf path, which telescopes: $w_i^{\text{HRP}} = \prod \frac{T_{\text{child}}}{T_{\text{parent}}} = \frac{T_{\{i\}}}{T_{\text{root}}} = \frac{\sigma_i^{-2}}{\sum_j \sigma_j^{-2}}$, independent of the tree; GMV for diagonal Σ is the same vector. Equivalently: $\Sigma_{AB} = 0$ everywhere and IV = GMV in every cluster, so all three substitutions of Corollary 7.3 are identities. $\square$

Proposition 7.6 (Uncorrelated exchangeable blocks; **P**). *Let Σ be block-diagonal with two blocks, block $K \in \{A, B\}$ exchangeable: $\Sigma_K = \sigma_K^2 [(1 - \rho_K)I + \rho_K \mathbf{1}\mathbf{1}^\top]$ with n_K assets, $\rho_K > -1/(n_K - 1)$, n_K a power of two. Suppose the bisection is tree-aligned with the top split at the block boundary and balanced within blocks. Then $\text{HRP} = \text{GMV}$ exactly:*

$$w_{i \in K}^{\text{GMV}} = \frac{m_K}{n_K(m_A + m_B)}, \quad m_K := \mathbf{1}^\top \Sigma_K^{-1} \mathbf{1} = \frac{n_K}{\sigma_K^2(1 + (n_K - 1)\rho_K)}.$$

Proof. *Within blocks.* Equal volatilities make inverse-variance weights on any sub-cluster equal weights. A sub-cluster of size k inside block K has cluster variance at equal weights $\tilde{V}(k) = \sigma_K^2(1 + (k - 1)\rho_K)/k$. Balanced bisection gives both children equal size, hence equal $\tilde{V}$, hence 1/2–1/2 splits; recursing down $n_K = 2^q$ yields equal weights $1/n_K$ within the block. Since $\Sigma_K \mathbf{1} = \sigma_K^2(1 + (n_K - 1)\rho_K)\mathbf{1}$, block GMV is also equal weights. *Across blocks.* At the top split, $\tilde{V}(n_K) = \sigma_K^2(1 + (n_K - 1)\rho_K)/n_K = 1/m_K$, so $\alpha_A = m_A/(m_A + m_B)$, which is the GMV

mass on A : block-diagonality stacks $\Sigma_A^{-1}\mathbf{1}$ and $\Sigma_B^{-1}\mathbf{1}$, with block masses $m_K/(m_A + m_B)$. (In the language of Corollary 7.3: $\Sigma_{AB} = 0$ makes (i)–(ii) identities at the top split; within blocks the substitutions are not identities individually, but exchangeability plus balance makes their composition weight-neutral.) $\square$

Remark 7.7 (Minimal counterexample; **P**). *Exactness fails as soon as a split must absorb within-cluster correlation with unequal volatilities. For $N = 2$, correlation ρ , volatilities $\sigma_1 \neq \sigma_2$: HRP ignores ρ (the children are singletons) and returns $w_1^{\text{HRP}} = \sigma_2^2/(\sigma_1^2 + \sigma_2^2)$, while $w_1^{\text{GMV}} = (\sigma_2^2 - \rho\sigma_1\sigma_2)/(\sigma_1^2 + \sigma_2^2 - 2\rho\sigma_1\sigma_2)$. These coincide iff $\rho = 0$ or $\sigma_1 = \sigma_2$ (when $\sigma_1 = \sigma_2$ both equal $1/2$ for every ρ). The hypotheses of Theorem 7.5 and Proposition 7.6 — zero correlation wherever volatilities differ within a cluster, equal volatilities wherever correlation survives — are not artifacts of the proofs but the boundary of exactness, and the deviation is exactly substitution (iii) of Corollary 7.3 at the bottom split.*

Corollary 7.8 (Mean–variance optimality of HRP; **P**). *Under the hypotheses of Theorem 7.5 or Proposition 7.6, HRP coincides with the tangency portfolio iff, in addition, $\mu_e \propto \mathbf{1}$ (Theorem 4.5). Under the diagonal hypothesis this reads: equal premia and uncorrelated assets, in which case $\text{HRP} = \text{IVar} = \text{GMV} = \text{tangency}$ simultaneously.*

Remark 7.9 (What HRP is actually for; **E**). *None of the above is the operative defense of HRP. López de Prado (2016) motivates the algorithm by the instability of $\hat{\Sigma}^{-1}$ under estimation error — quasi-diagonalization and recursion avoid inverting an ill-conditioned matrix — and reports Monte Carlo evidence of superior out-of-sample variance relative to plug-in minimum variance. That defense is empirical and belongs to the information-set logic of Sections 11–12, not to exact optimality; Corollary 7.3 gives it structure by listing precisely which estimated objects (Σ_{AB} , within-cluster correlations) HRP declines to trust.*

7.4 Return-adjusted hierarchical risk parity

Standard HRP is deliberately risk-only. RA-HRP keeps the clustering and quasi-diagonalization stages fixed, but replaces the inverse-variance split score by a local cluster-Sharpe score. This makes the rule materially different from the flat heuristics above: its weight vector is a function of both Σ and μ_e , so the implied-return condition is no longer a fixed ray in μ_e for fixed Σ . It is a nonlinear fixed point.

Definition 7.10 (Fixed-tree RA-HRP; **P**, notation following Nogueira i Alonso, 2026). *Fix a binary dendrogram $\mathcal{T}$ and a score floor $\varepsilon > 0$. For every nonempty cluster $K \subset \{1, \dots, N\}$ define the inverse-variance portfolio*

$$\tilde{w}_{K,i} = \frac{\sigma_i^{-2}}{\sum_{j \in K} \sigma_j^{-2}}, \quad i \in K,$$

its local mean, variance, and floored Sharpe score by

$$M_K := \sum_{i \in K} \tilde{w}_{K,i} \mu_{e,i}, \quad V_K := \tilde{\mathbf{w}}_K^\top \Sigma_K \tilde{\mathbf{w}}_K, \quad Q_K := \max \left\{ \frac{M_K}{\sqrt{V_K}}, \varepsilon \right\}. \quad (7.3)$$

At an internal node $C = A \cup B$, RA-HRP allocates the parent budget by

$$\beta_A^{\text{RA}} = \frac{Q_A}{Q_A + Q_B}, \quad \beta_B^{\text{RA}} = 1 - \beta_A^{\text{RA}} = \frac{Q_B}{Q_A + Q_B}. \quad (7.4)$$

The final portfolio obtained by multiplying these branch fractions along root-to-leaf paths is denoted

$$\mathbf{w}^{\text{RA}} = \Phi_{\mathcal{T}, \varepsilon}^{\text{RA}}(\mu_e, \Sigma).$$

When the raw score $M_K/\sqrt{V_K}$ exceeds ε for every cluster, the floor is inactive and the split is exactly a cluster-Sharpe split.

Theorem 7.11 (Well-posedness and lower bounds; **P**). *For fixed $(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}, \mathcal{T}, \varepsilon)$ with $\boldsymbol{\Sigma} \succ 0$ and $\varepsilon > 0$, RA-HRP defines a unique portfolio $\mathbf{w}^{\text{RA}} \in \Delta^{N-1}$. If $D_{\mathcal{T}}$ is the maximum root-to-leaf depth and $M_Q := \max_{K \in \mathcal{T}} Q_K$, then every leaf satisfies*

$$w_i^{\text{RA}} \geq \left(\frac{\varepsilon}{\varepsilon + M_Q} \right)^{D_{\mathcal{T}}}. \quad (7.5)$$

On any region where the active/inactive status of the floor does not change and all cluster variances remain positive, the map $(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}) \mapsto \Phi_{\mathcal{T}, \varepsilon}^{\text{RA}}(\boldsymbol{\mu}_e, \boldsymbol{\Sigma})$ is smooth.

Proof. At every node, $Q_A, Q_B \geq \varepsilon$, so the denominator in (7.4) is strictly positive and

$$\frac{\varepsilon}{\varepsilon + M_Q} \leq \beta_A^{\text{RA}}, \beta_B^{\text{RA}} \leq \frac{M_Q}{\varepsilon + M_Q} < 1.$$

The tree has finitely many nodes, and the recursive multiplication of deterministic split fractions therefore gives a unique strictly positive leaf weight. Budgets are conserved at every split because $\beta_A^{\text{RA}} + \beta_B^{\text{RA}} = 1$, so the leaf weights sum to one. The lower bound follows by multiplying the worst possible branch fraction along a path of length at most $D_{\mathcal{T}}$. Smoothness on a fixed floor regime follows because the IV weights, M_K , V_K , Q_K , and the rational split map are compositions of smooth functions with positive denominators. $\square$

Definition 7.12 (Unit-free HRP–RA-HRP interpolation; **P**). *At an internal node $C = A \cup B$, write the standard HRP left-branch fraction and the RA-HRP left-branch fraction as*

$$r_A := \frac{V_B}{V_A + V_B}, \quad s_A := \frac{Q_A}{Q_A + Q_B}, \quad (7.6)$$

where V_K and Q_K are the cluster IVP variance and floored cluster-Sharpe score from (7.3). For $\lambda \in [0, 1]$, define the interpolated local split

$$\beta_A^{(\lambda)} := (1 - \lambda)r_A + \lambda s_A, \quad \beta_B^{(\lambda)} := 1 - \beta_A^{(\lambda)}. \quad (7.7)$$

Applying (7.7) recursively on the same tree defines the interpolated portfolio $\mathbf{w}^{(\lambda)}$.

Proposition 7.13 (Properties of the unit-free interpolation; **P**). *For the family (7.7): (i) $\lambda = 0$ gives HRP; (ii) $\lambda = 1$ gives RA-HRP; (iii) $\lambda \mapsto \beta_A^{(\lambda)}$ is affine and continuous on $[0, 1]$; and (iv)*

$$\frac{d}{d\lambda} \beta_A^{(\lambda)} = s_A - r_A. \quad (7.8)$$

Consequently the local path is monotone whenever the HRP and RA-HRP branch fractions differ.

Proof. Substitution of $\lambda = 0$ and $\lambda = 1$ gives the two endpoints. The map in (7.7) is an affine function of λ , hence continuous and differentiable, with derivative (7.8). Since both r_A and s_A lie in $(0, 1)$, every interpolated split remains feasible. $\square$

Remark 7.14 (Why the interpolation is economically clean). *The interpolation is performed on branch probabilities, not on raw scores. This matters because inverse variances and Sharpe ratios have different units and different estimation errors. Blending r_A and s_A gives λ a literal interpretation: it is the local weight placed on the return-adjusted allocation decision relative to the risk-only decision. This is the correct homotopy for empirical sensitivity analysis and for any future theorem that treats RA-HRP as a controlled perturbation of HRP.*

Theorem 7.15 (Exact RA-HRP tangency coincidence; **P**). *Fix $(\mathcal{T}, \varepsilon)$ and define*

$$\mathbf{w}^{\text{RA}} = \Phi_{\mathcal{T}, \varepsilon}^{\text{RA}}(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}).$$

Assume $\mathbf{1}^\top \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_e > 0$. Then the following are equivalent:

(i) RA-HRP is the tangency portfolio: $\mathbf{w}^{\text{RA}} = \mathbf{w}_T$.

(ii) There exists a $a > 0$ such that

$$\boldsymbol{\mu}_e = a \boldsymbol{\Sigma} \Phi_{T,\varepsilon}^{\text{RA}}(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}). \quad (7.9)$$

(iii) RA-HRP has zero alpha against itself: for every asset i ,

$$\mu_{e,i} = \beta_i(\mathbf{w}^{\text{RA}}) (\mathbf{w}^{\text{RA}\top} \boldsymbol{\mu}_e).$$

Moreover, at any split $C = A \cup B$ of an exact tangency recursion, with local covariance-premium pair (Γ_C, ν_C) , write

$$\Gamma_C = \begin{pmatrix} \Gamma_A & \Gamma_{AB} \\ \Gamma_{BA} & \Gamma_B \end{pmatrix}, \quad S_A = \Gamma_A - \Gamma_{AB} \Gamma_B^{-1} \Gamma_{BA}, \quad S_B = \Gamma_B - \Gamma_{BA} \Gamma_A^{-1} \Gamma_{AB},$$

$$\nu_A^{\text{Sch}} = \nu_A - \Gamma_{AB} \Gamma_B^{-1} \nu_B, \quad \nu_B^{\text{Sch}} = \nu_B - \Gamma_{BA} \Gamma_A^{-1} \nu_A,$$

and

$$L_A := \mathbf{1}_A^\top S_A^{-1} \nu_A^{\text{Sch}}, \quad L_B := \mathbf{1}_B^\top S_B^{-1} \nu_B^{\text{Sch}}. \quad (7.10)$$

When $L_A, L_B > 0$, the exact tangency split is $L_A/(L_A + L_B)$. Hence the RA-HRP split is locally exact at that node if and only if

$$\frac{Q_A}{Q_B} = \frac{L_A}{L_B}. \quad (7.11)$$

Full tree coincidence holds if and only if (7.11) holds at every internal node when the right-hand side is computed from the Schur-updated conditional pair generated by the exact tangency recursion.

Proof. The equivalence of (i) and (ii) is Theorem 3.3(i), with the important difference that the candidate $\mathbf{w}^{\text{RA}}$ itself depends on $(\boldsymbol{\mu}_e, \boldsymbol{\Sigma})$. The equivalence of (i) and (iii) is the beta form in Remark 3.5.

For the split statement, the unnormalized tangency direction on a local pair (Γ_C, ν_C) is $x_C = \Gamma_C^{-1} \nu_C$. The block-inverse identity gives

$$x_A = S_A^{-1}(\nu_A - \Gamma_{AB} \Gamma_B^{-1} \nu_B), \quad x_B = S_B^{-1}(\nu_B - \Gamma_{BA} \Gamma_A^{-1} \nu_A).$$

Therefore the unnormalized capital masses assigned by the exact tangency direction to the two children are $L_A = \mathbf{1}_A^\top x_A$ and $L_B = \mathbf{1}_B^\top x_B$. If both are positive, the exact local budget share of A is $L_A/(L_A + L_B)$. The RA-HRP local share is $Q_A/(Q_A + Q_B)$; equality is equivalent to the ratio identity (7.11). Recursing is legitimate because $x_A = S_A^{-1} \nu_A^{\text{Sch}}$ is itself the tangency direction of the Schur-updated pair $(S_A, \nu_A^{\text{Sch}})$ inside child A , and analogously for B . Matching every local split on the tree is equivalent to matching every root-to-leaf product of split fractions, hence to full portfolio coincidence. $\square$

Remark 7.16 (What RA-HRP fixes and what it does not). *RA-HRP repairs one weakness of standard HRP: it no longer allocates solely by inverse cluster variance, and therefore it can tilt toward clusters with superior estimated reward per unit of local risk. But Theorem 7.15 shows that the exact Markowitz split is not a raw cluster-Sharpe split. The optimizer uses Schur-adjusted covariance S_A, S_B and Schur-adjusted premia $\nu_A^{\text{Sch}}, \nu_B^{\text{Sch}}$. RA-HRP therefore replaces HRP's purely risk-based local score by a return-aware local score, while still avoiding the exact Schur and global-inversion machinery. Its optimality set is larger and more economically flexible than HRP's equal-premium GMV set, but it remains a restrictive nonlinear fixed-point set rather than a generic property.*

Definition 7.17 (Schur-RA-HRP local score; $\mathbf{P}$). *This is the local conditional-risk extension of RA-HRP developed in [Noguer i Alonso \(2026\)](#). At a sibling split $C = A \cup B$, partition the local covariance matrix as*

$$\Sigma_C = \begin{pmatrix} \Sigma_{AA} & \Sigma_{AB} \\ \Sigma_{BA} & \Sigma_{BB} \end{pmatrix}.$$

Define the conditional Schur covariance blocks

$$\Sigma_{A|B} := \Sigma_{AA} - \Sigma_{AB}\Sigma_{BB}^{-1}\Sigma_{BA}, \quad \Sigma_{B|A} := \Sigma_{BB} - \Sigma_{BA}\Sigma_{AA}^{-1}\Sigma_{AB}. \quad (7.12)$$

Using the same within-cluster IVP weights $\tilde{\mathbf{w}}_A, \tilde{\mathbf{w}}_B$ and the same cluster means M_A, M_B , set

$$V_{A|B}^{\text{Sch}} := \tilde{\mathbf{w}}_A^\top \Sigma_{A|B} \tilde{\mathbf{w}}_A, \quad Q_{A|B}^{\text{Sch}} := \max \left\{ \frac{M_A}{\sqrt{V_{A|B}^{\text{Sch}}}}, \varepsilon \right\}, \quad (7.13)$$

and analogously for $B | A$. The Schur-RA-HRP split is

$$\beta_A^{\text{SchRA}} = \frac{Q_{A|B}^{\text{Sch}}}{Q_{A|B}^{\text{Sch}} + Q_{B|A}^{\text{Sch}}}, \quad \beta_B^{\text{SchRA}} = 1 - \beta_A^{\text{SchRA}}. \quad (7.14)$$

Proposition 7.18 (Conditional-risk reduction and the Schur score wedge; $\mathbf{P}$). *If $\Sigma_C \succ 0$, then*

$$0 < V_{A|B}^{\text{Sch}} = V_A - \tilde{\mathbf{w}}_A^\top \Sigma_{AB} \Sigma_{BB}^{-1} \Sigma_{BA} \tilde{\mathbf{w}}_A \leq V_A, \quad (7.15)$$

with equality if and only if $\Sigma_{BA} \tilde{\mathbf{w}}_A = 0$. Moreover,

$$0 \leq V_A - V_{A|B}^{\text{Sch}} \leq \frac{\|\Sigma_{BA} \tilde{\mathbf{w}}_A\|_2^2}{\lambda_{\min}(\Sigma_{BB})}, \quad (7.16)$$

and analogously for $B | A$. Hence, whenever the cluster mean is positive, the raw Schur score $M_A/\sqrt{V_{A|B}^{\text{Sch}}}$ is weakly larger than the standalone RA-HRP score $M_A/\sqrt{V_A}$; after the floor, the score cannot decrease.

Proof. Since $\Sigma_C \succ 0$, both principal blocks are positive definite and the Schur complements in (7.12) are positive definite. Expanding the quadratic form gives the identity in (7.15); the subtracted term equals $\|\Sigma_{BB}^{-1/2} \Sigma_{BA} \tilde{\mathbf{w}}_A\|_2^2 \geq 0$, with equality exactly when $\Sigma_{BA} \tilde{\mathbf{w}}_A = 0$. The operator inequality $\Sigma_{BB}^{-1} \preceq \lambda_{\min}(\Sigma_{BB})^{-1} I$ yields (7.16). If $M_A > 0$, reducing the denominator weakly raises the raw score; if $M_A \leq 0$, both raw scores are at or below the positive floor, so the floored score remains ε unless the Schur adjustment pushes it above the floor. $\square$

Remark 7.19 (Relation to the exact Schur-tangency split). *Schur-RA-HRP repairs a second HRP substitution. Standard RA-HRP replaces inverse-variance splitting by return-aware cluster-Sharpe splitting, but it still uses standalone cluster variance. Definition 7.17 replaces that standalone risk by conditional risk. It therefore moves the rule toward the exact nodewise condition (7.11). It is still not a global Markowitz optimizer: it uses local block inversions only, keeps the tree fixed, and uses IVP aggregation inside clusters. Its purpose is to recover the economically important cross-cluster hedging term without paying the full global inversion tax.*

Corollary 7.20 (Two-asset boundary; $\mathbf{P}$). *Consider two assets, assume the floor is inactive and the relevant quantities are positive. Write ρ for their correlation. RA-HRP coincides with tangency if and only if*

$$\frac{\mu_{e,1}/\sigma_1}{\mu_{e,2}/\sigma_2} = \frac{\sigma_2^2 \mu_{e,1} - \rho \sigma_1 \sigma_2 \mu_{e,2}}{\sigma_1^2 \mu_{e,2} - \rho \sigma_1 \sigma_2 \mu_{e,1}}. \quad (7.17)$$

In particular, when $\rho = 0$ and $\mu_{e,1}, \mu_{e,2} > 0$, RA-HRP is tangency if and only if $\sigma_1 = \sigma_2$. Thus even for diagonal covariance, RA-HRP is generally not the plug-in Markowitz rule: with two uncorrelated assets it allocates by signal-to-volatility, whereas tangency allocates by signal-to-variance.

Proof. With two singleton children, the inactive-floor RA-HRP ratio is

$$\frac{w_1^{\text{RA}}}{w_2^{\text{RA}}} = \frac{\mu_{e,1}/\sigma_1}{\mu_{e,2}/\sigma_2}.$$

The unnormalized tangency direction is

$$\Sigma^{-1} \boldsymbol{\mu}_e \propto \begin{pmatrix} \sigma_2^2 \mu_{e,1} - \rho \sigma_1 \sigma_2 \mu_{e,2} \\ \sigma_1^2 \mu_{e,2} - \rho \sigma_1 \sigma_2 \mu_{e,1} \end{pmatrix},$$

which gives (7.17). Setting $\rho = 0$ and cancelling the positive premia gives $\sigma_2/\sigma_1 = \sigma_2^2/\sigma_1^2$, hence $\sigma_1 = \sigma_2$. $\square$

Proposition 7.21 (Balanced equal-volatility diagonal exactness; **P**). *Suppose $\Sigma = \sigma^2 I$, every split in $\mathcal{T}$ is balanced, all cluster premium sums are positive, and the floor is inactive. Then RA-HRP coincides with the tangency portfolio:*

$$w_i^{\text{RA}} = \frac{\mu_{e,i}}{\sum_j \mu_{e,j}} = w_{T,i}.$$

Proof. For a cluster K , IV weights are equal weights, $V_K = \sigma^2/|K|$, and

$$Q_K = \frac{|K|^{-1} \sum_{i \in K} \mu_{e,i}}{\sigma/\sqrt{|K|}} = \frac{\sum_{i \in K} \mu_{e,i}}{\sigma\sqrt{|K|}}.$$

At a balanced split $K = A \cup B$, $|A| = |B|$, so

$$\beta_A^{\text{RA}} = \frac{\sum_{i \in A} \mu_{e,i}}{\sum_{i \in A} \mu_{e,i} + \sum_{i \in B} \mu_{e,i}}.$$

Multiplying these branch ratios along a root-to-leaf path telescopes to $\mu_{e,i}/\sum_j \mu_{e,j}$. Since $\Sigma = \sigma^2 I$, the tangency portfolio is proportional to $\Sigma^{-1} \boldsymbol{\mu}_e = \sigma^{-2} \boldsymbol{\mu}_e$, giving the result. $\square$

Theorem 7.22 (Pathwise HRP–RA-HRP distortion and KL attribution; **P**). *Fix a common tree $\mathcal{T}$. Let $r_v \in (0, 1)$ be the HRP left-branch split at internal node v and $s_v \in (0, 1)$ the RA-HRP left-branch split. For a leaf i , let $\mathcal{P}(i)$ be its set of ancestor internal nodes and let $\eta_v(i) = 1$ if i descends through the left child of v and 0 otherwise. Then*

$$\frac{w_i^{\text{RA}}}{w_i^{\text{HRP}}} = \prod_{v \in \mathcal{P}(i)} \left(\frac{s_v}{r_v} \right)^{\eta_v(i)} \left(\frac{1-s_v}{1-r_v} \right)^{1-\eta_v(i)}. \quad (7.18)$$

Moreover, viewing the two long-only portfolios as probability measures on leaves,

$$D_{\text{KL}}(\mathbf{w}^{\text{RA}} \parallel \mathbf{w}^{\text{HRP}}) = \sum_{v \in \mathcal{V}_{\text{int}}} W_v^{\text{RA}} \left[s_v \log \frac{s_v}{r_v} + (1-s_v) \log \frac{1-s_v}{1-r_v} \right], \quad (7.19)$$

where W_v^{RA} is the RA-HRP total capital entering node v .

Proof. Both portfolios are products of local branch fractions along a root-to-leaf path. Dividing the two products gives (7.18). Taking logarithms, multiplying by w_i^{RA} , summing over leaves, and exchanging the order of summation gives (7.19). At node v , the sum of RA-HRP weights in the left subtree is $W_v^{\text{RA}} s_v$ and in the right subtree is $W_v^{\text{RA}} (1-s_v)$, which yields the binary KL term. $\square$

Corollary 7.23 (Exact Sharpe-efficiency comparison; **P**). *Let $\eta_H = S(\mathbf{w}^{\text{HRP}})/S_{\max}$ and $\eta_R = S(\mathbf{w}^{\text{RA}})/S_{\max}$. The reduction in squared Sharpe loss obtained by moving from HRP to RA-HRP is exactly*

$$(1 - \eta_H^2) - (1 - \eta_R^2) = \eta_R^2 - \eta_H^2 \quad (7.20)$$

$$= \frac{S(\mathbf{w}^{\text{RA}})^2 - S(\mathbf{w}^{\text{HRP}})^2}{S_{\max}^2} \quad (7.21)$$

$$= \cos^2 \theta_{\Sigma}(\mathbf{w}^{\text{RA}}, \mathbf{w}_T) - \cos^2 \theta_{\Sigma}(\mathbf{w}^{\text{HRP}}, \mathbf{w}_T). \quad (7.22)$$

Therefore RA-HRP improves the exact-optimality gap precisely when its pathwise split distortion moves the final portfolio closer to the tangency ray in the Σ -angle. Equation (7.18) attributes the source of that movement to individual tree nodes; Equation (7.19) measures how much capital was reallocated at each node.

Proof. This is Theorem 3.7 applied once to $\mathbf{w}^{\text{HRP}}$ and once to $\mathbf{w}^{\text{RA}}$. $\square$

Remark 7.24 (Statistical status and empirical scope; **C/E**). *The fixed-tree RA-HRP paper proves existence, uniqueness, explicit leaf lower bounds, local split curvature, pathwise HRP–RA-HRP distortion, KL decomposition, and fixed-tree consistency/asymptotic normality under fixed dimension, finite fourth moments, and an inactive floor (Noguer i Alonso, 2026). It also emphasizes the theory–empirics gap created by re-estimating dendrograms in rolling windows: tree changes make the implemented map piecewise smooth rather than globally smooth. Empirically, the reported 15-stock US backtest is best read as proof of concept: RA-HRP improves traditional HRP in that sample, but a capped within-universe market-cap proxy attains an even higher Sharpe ratio. This is exactly the right evidentiary status for the present paper: RA-HRP is a disciplined return-aware extension of HRP, not a theorem of universal dominance.*

Remark 7.25 (Open problem: data-driven trees as random topologies). *The fixed-tree theory is the right first theorem because all split maps are smooth away from floor and variance boundaries. The implemented strategy, however, usually re-estimates the dendrogram. Theorem 7.22 suggests a natural route to a full estimated-tree theory: compare two random trees by the split ratios induced on common root-to-leaf paths and by distances between their merge sets, cophenetic matrices, or tree-edit representations. One possible program is: (i) prove high-probability topology preservation under a merge-separation condition on the population distance matrix; (ii) on that event, apply the fixed-tree delta method; (iii) off that event, control the portfolio jump by pathwise products such as (7.18) and by a metric on dendrogram topologies; and (iv) combine the two pieces into a uniform-in-tree convergence bound. The empirical merge-preservation and cophenetic-correlation diagnostics reported in Noguer i Alonso (2026) are therefore not merely descriptive; they are the measurable quantities a future random-tree asymptotic theorem must control.*

7.5 First-order Sharpe calculus along the HRP–RA-HRP homotopy

The unit-free interpolation of Proposition 7.13 turns “how much local return information to admit” into a one-dimensional decision. This subsection differentiates the Sharpe objective along that path. The derivative turns out to be a sum over internal nodes of the products of two estimable quantities: the displacement of the local split and the *alpha*, against the HRP portfolio itself, of the self-financing switch between the node’s two children. The marginal value of return information is therefore a nodewise zero-alpha computation — the same object the GRS machinery of Remark 3.6 estimates.

Lemma 7.26 (Node-switch vectors; **P**). *Fix $\mathcal{T}$, strictly interior splits $r_v, s_v \in (0, 1)$ at every internal node v , and the interpolated portfolio $\mathbf{w}^{(\lambda)}$ of (7.7). Let $W_v^{\text{H}} := \sum_{i: v \in \mathcal{P}(i)} w_i^{\text{HRP}}$ be the*

HRP capital entering v . Define $g_v \in \mathbb{R}^N$ by

$$g_{v,i} := w_i^{\text{HRP}} \left(\frac{\eta_v(i)}{r_v} - \frac{1 - \eta_v(i)}{1 - r_v} \right) \quad \text{if } v \in \mathcal{P}(i), \quad g_{v,i} := 0 \quad \text{otherwise.} \quad (7.23)$$

Then:

- (i) $\mathbf{1}^\top g_v = 0$: each g_v is a self-financing reallocation.
- (ii) $g_v = W_v^H(\mathbf{u}_{L(v)} - \mathbf{u}_{R(v)})$, where $\mathbf{u}_K$ is the HRP composition inside subtree K renormalized to unit mass: g_v moves one unit of node- v capital from the right-child HRP composition to the left-child HRP composition, levered by the capital actually present at the node.
- (iii) $\lambda \mapsto \mathbf{w}^{(\lambda)}$ is a polynomial in λ , and

$$\left. \frac{d}{d\lambda} \mathbf{w}^{(\lambda)} \right|_{\lambda=0} = \sum_{v \in \mathcal{V}_{\text{int}}} (s_v - r_v) g_v. \quad (7.24)$$

Proof. Each leaf weight is the root-to-leaf product of branch fractions (the representation underlying Theorem 7.22), here of the affine functions $\beta_v^{(\lambda)} = (1 - \lambda)r_v + \lambda s_v$ along its path, hence a polynomial in λ . Logarithmic differentiation at $\lambda = 0$ gives

$$\left. \frac{d}{d\lambda} w_i^{(\lambda)} \right|_{\lambda=0} = w_i^{\text{HRP}} \sum_{v \in \mathcal{P}(i)} (s_v - r_v) \left(\frac{\eta_v(i)}{r_v} - \frac{1 - \eta_v(i)}{1 - r_v} \right),$$

which is the i -th coordinate of (7.24). For (i)–(ii): the HRP mass in the left subtree of v is $W_v^H r_v$ and in the right subtree $W_v^H(1 - r_v)$, so summing (7.23) over the left child gives $W_v^H r_v / r_v = W_v^H$ and over the right child $-W_v^H$; the totals cancel, and dividing each child's restriction of $\mathbf{w}^{\text{HRP}}$ by its subtree mass exhibits the two unit-mass compositions $\mathbf{u}_{L(v)}, \mathbf{u}_{R(v)}$. $\square$

Theorem 7.27 (Nodewise-alpha representation of the homotopy derivative; **P**). *Let $\Sigma \succ 0$, $\mu_e \neq 0$, and write $a^*(\mathbf{w}) = \mathbf{w}^\top \mu_e / \mathbf{w}^\top \Sigma \mathbf{w}$ and*

$$\alpha(\mathbf{w}) := \mu_e - a^*(\mathbf{w}) \Sigma \mathbf{w}$$

for the implied-return residual of $\mathbf{w}$; by Remark 3.5, $\alpha_i(\mathbf{w}) = \mu_{e,i} - \beta_i(\mathbf{w})(\mathbf{w}^\top \mu_e)$ is exactly the vector of alphas in (3.4), and $\alpha(\mathbf{w})$ is the minimizing residual in the defect (8.2). Then, with $\mathbf{w}^H := \mathbf{w}^{\text{HRP}}$ and

$$\alpha_v := g_v^\top \alpha(\mathbf{w}^H) = g_v^\top \mu_e - \frac{g_v^\top \Sigma \mathbf{w}^H}{\mathbf{w}^{H\top} \Sigma \mathbf{w}^H} (\mathbf{w}^{H\top} \mu_e), \quad (7.25)$$

the Sharpe ratio and the squared Sharpe efficiency satisfy

$$\begin{aligned} \left. \frac{d}{d\lambda} S(\mathbf{w}^{(\lambda)}) \right|_{\lambda=0} &= \frac{1}{\sigma_p(\mathbf{w}^H)} \sum_{v \in \mathcal{V}_{\text{int}}} (s_v - r_v) \alpha_v, \\ \left. \frac{d}{d\lambda} (1 - \eta^2)(\mathbf{w}^{(\lambda)}) \right|_{\lambda=0} &= - \frac{2S(\mathbf{w}^H)}{S_{\max}^2 \sigma_p(\mathbf{w}^H)} \sum_{v \in \mathcal{V}_{\text{int}}} (s_v - r_v) \alpha_v. \end{aligned} \quad (7.26)$$

The scalar α_v is precisely the alpha of the self-financing node- v switch g_v in the one-factor regression of Remark 3.6 with the HRP portfolio as the candidate factor.

Proof. For $\mathbf{w} \neq 0$, $\nabla S(\mathbf{w}) = \mu_e / \sigma_p(\mathbf{w}) - (\mathbf{w}^\top \mu_e) \Sigma \mathbf{w} / \sigma_p(\mathbf{w})^3 = \alpha(\mathbf{w}) / \sigma_p(\mathbf{w})$. The chain rule with Lemma 7.26(iii) gives the first display in (7.26); the second follows from $1 - \eta^2 = 1 - S^2 / S_{\max}^2$ with $S_{\max}$ independent of λ . For the alpha interpretation: g_v is zero-cost by Lemma 7.26(i), its beta against $\mathbf{w}^H$ is $\beta(g_v) = g_v^\top \Sigma \mathbf{w}^H / \mathbf{w}^{H\top} \Sigma \mathbf{w}^H$, and its alpha in the sense of (3.4) is $g_v^\top \mu_e - \beta(g_v)(\mathbf{w}^{H\top} \mu_e)$, which is (7.25). That this equals the minimizing residual of Definition 8.3 contracted with g_v is Theorem 8.4: a^* in (8.4) is the same scalar. $\square$

Corollary 7.28 (Marginal value of return information; **P**). *To first order at $\lambda = 0$, admitting local return information improves the Sharpe ratio if and only if*

$$\sum_{v \in \mathcal{V}_{\text{int}}} (s_v - r_v) \alpha_v > 0. \quad (7.27)$$

Each node contributes the product of the split displacement $s_v - r_v$ (how far the floored cluster-Sharpe score moves the local budget away from the inverse-variance split) and the node alpha α_v (whether the implied child switch is priced by the HRP portfolio). In particular: (i) if every node switch has zero alpha against HRP — the nodewise refinement of the zero-alpha null of Remark 3.6 restricted to the $|\mathcal{V}_{\text{int}}|$ tradable directions $\{g_v\}$ — the homotopy is first-order flat regardless of the score displacements; (ii) a node helps exactly when its score tilts toward the child whose switch carries positive alpha. Since $\alpha_v = \sum_i g_{v,i} \alpha_i(\mathbf{w}^H)$ is a fixed linear combination of the regression intercepts entering the GRS statistic (14.5), the entire derivative (7.26) is estimable from a single set of time-series regressions on the realized HRP return, with standard errors inherited from $\hat{\alpha}$.

Proof. Sign reading of (7.26); (i)–(ii) are immediate, and the estimability statement is the linearity of (7.25) in $\alpha(\mathbf{w}^H)$. $\square$

Proposition 7.29 (KL trust budget along the homotopy; **P**). *For every internal node v , the map $\lambda \mapsto d_{\text{KL}}(\beta_v^{(\lambda)} \| r_v)$ (binary KL divergence) is convex on $[0, 1]$, vanishes together with its derivative at $\lambda = 0$, and satisfies the chord bound $d_{\text{KL}}(\beta_v^{(\lambda)} \| r_v) \leq \lambda d_{\text{KL}}(s_v \| r_v)$. Consequently, with $W_v^{(\lambda)}$ the interpolated capital entering v ,*

$$\begin{aligned} D_{\text{KL}}(\mathbf{w}^{(\lambda)} \| \mathbf{w}^{\text{HRP}}) &= \sum_{v \in \mathcal{V}_{\text{int}}} W_v^{(\lambda)} d_{\text{KL}}(\beta_v^{(\lambda)} \| r_v) \leq \lambda \sum_{v \in \mathcal{V}_{\text{int}}} W_v^{(\lambda)} d_{\text{KL}}(s_v \| r_v) \\ &\leq \lambda D_{\mathcal{T}} \max_v d_{\text{KL}}(s_v \| r_v), \end{aligned} \quad (7.28)$$

since $\sum_v W_v^{(\lambda)} = \sum_i w_i^{(\lambda)} |\mathcal{P}(i)| \leq D_{\mathcal{T}}$ is the capital-weighted tree depth.

Proof. The identity in (7.28) is Theorem 7.22 with s_v replaced by $\beta_v^{(\lambda)}$: the proof there uses only the product structure of tree weights, never the specific form of the splits. For fixed $r \in (0, 1)$, $x \mapsto d_{\text{KL}}(x \| r) = x \log(x/r) + (1-x) \log((1-x)/(1-r))$ is convex with derivative $\log(x/r) - \log((1-x)/(1-r))$, which vanishes at $x = r$; composing with the affine path $\beta_v^{(\lambda)}$ preserves convexity, and the value and derivative at $\lambda = 0$ are zero, so the map is nondecreasing. Convexity with the endpoint values 0 and $d_{\text{KL}}(s_v \| r_v)$ gives the chord bound. Finally each unit of leaf capital is counted once per ancestor, so $\sum_v W_v^{(\lambda)} = \sum_i w_i^{(\lambda)} |\mathcal{P}(i)|$, bounded by the maximal root-to-leaf depth. $\square$

The economic reading: λ is a *certified trust budget*. Choosing the interpolation level caps the capital reallocated away from the risk-only baseline linearly in λ , node by node, before any data are seen — which is exactly the role the λ_{KL} penalty plays dynamically in the RLPO objective (9.11) below. Theorem 7.27 says where that budget earns alpha; Proposition 7.29 says how much structural movement it can cause.

8 The Mathematics of Heuristic Portfolio Optimization (HPO)

The preceding sections compute exact coincidence conditions rule by rule. This section abstracts from the individual names and treats heuristic portfolio construction itself as a mathematical object. The key move is to separate three layers that are often conflated in practice: the full population optimizer, the information-restricted portfolio map, and the estimated implementation of that map.

8.1 HPO maps as information-restricted portfolio rules

Definition 8.1 (Information-restricted portfolio map). *Let*

$$\Theta := \{(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}) : \boldsymbol{\mu}_e \in \mathbb{R}^N, \boldsymbol{\Sigma} \in \mathbb{S}_{++}^N\}$$

be the population parameter space. An information set is a statistic

$$\mathcal{I} : \Theta \rightarrow \mathcal{Z},$$

and a heuristic portfolio map is a measurable function

$$\Phi : \mathcal{Z} \rightarrow \Delta^{N-1}, \quad \Delta^{N-1} := \{\mathbf{w} \in \mathbb{R}^N : \mathbf{1}^\top \mathbf{w} = 1, w_i \geq 0\}.$$

The induced portfolio at population point $\theta = (\boldsymbol{\mu}_e, \boldsymbol{\Sigma})$ is

$$\mathbf{w}_\Phi(\theta) := \Phi(\mathcal{I}(\theta)).$$

A parametric HPO family is a collection $\{\Phi_\lambda : \lambda \in \Lambda\}$, where λ may represent a shrinkage intensity, a score floor, a tree depth, a turnover penalty, or the HRP–RA–HRP interpolation weight.

Examples are immediate. Equal weight has a trivial information set. Inverse volatility uses only $\text{diag}(\boldsymbol{\Sigma})$. ERC uses $\boldsymbol{\Sigma}$ but not $\boldsymbol{\mu}_e$. HRP uses a dendrogram and recursively aggregated cluster variances. RA-HRP uses the same tree plus local cluster-premium and cluster-risk scores. Schur-RA-HRP enlarges the local information set by allowing sibling-block Schur complements. Thus the rule name is less important than the map

$$(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}) \mapsto \mathcal{I}(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}) \mapsto \Phi(\mathcal{I}(\boldsymbol{\mu}_e, \boldsymbol{\Sigma})).$$

HPO studies the geometry and sampling behavior of this composition.

Definition 8.2 (Population HPO projection). *For a family $\{\Phi_\lambda\}_{\lambda \in \Lambda}$ and a population point $\theta = (\boldsymbol{\mu}_e, \boldsymbol{\Sigma})$ with positive tangency portfolio, the population-optimal member of the heuristic family is*

$$\lambda^*(\theta) \in \arg \max_{\lambda \in \Lambda} S(\mathbf{w}_{\Phi_\lambda}(\theta)) = \arg \min_{\lambda \in \Lambda} \theta_\Sigma(\mathbf{w}_{\Phi_\lambda}(\theta), \mathbf{w}_T(\theta)), \quad (8.1)$$

where θ_Σ is the angle in the $\boldsymbol{\Sigma}$ -inner product from Theorem 3.7.

Equation (8.1) is the clean mathematical definition of HPO: it is not full optimization over all portfolios; it is angular projection of the tangency direction onto a lower-complexity image set generated by admissible heuristic maps. The image set may be finite (choose among EW, IV, ERC, HRP), one-dimensional (HRP–RA–HRP interpolation), or high-dimensional (a family of trees, floors, shrinkage operators, and cost penalties).

8.2 The implied-return defect of a heuristic rule

The implied-return principle gives a natural residual for every heuristic. If $\mathbf{w}$ were exactly optimal, $\boldsymbol{\mu}_e$ would lie on the ray spanned by $\boldsymbol{\Sigma}\mathbf{w}$. The distance to that ray is the exact HPO defect.

Definition 8.3 (HPO implied-return defect). *For a fully invested portfolio $\mathbf{w}$ with $\mathbf{w}^\top \boldsymbol{\mu}_e > 0$, define*

$$\mathfrak{d}(\mathbf{w}; \boldsymbol{\mu}_e, \boldsymbol{\Sigma}) := \min_{a > 0} \|\boldsymbol{\mu}_e - a\boldsymbol{\Sigma}\mathbf{w}\|_{\boldsymbol{\Sigma}^{-1}}^2, \quad \|\mathbf{x}\|_{\boldsymbol{\Sigma}^{-1}}^2 := \mathbf{x}^\top \boldsymbol{\Sigma}^{-1} \mathbf{x}. \quad (8.2)$$

The normalized defect is

$$\mathfrak{D}(\mathbf{w}; \boldsymbol{\mu}_e, \boldsymbol{\Sigma}) := \frac{\mathfrak{d}(\mathbf{w}; \boldsymbol{\mu}_e, \boldsymbol{\Sigma})}{\boldsymbol{\mu}_e^\top \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_e}. \quad (8.3)$$

For a rule Φ , write $\mathfrak{D}_\Phi(\theta) := \mathfrak{D}(\mathbf{w}_\Phi(\theta); \theta)$.

Theorem 8.4 (Defect equals squared Sharpe inefficiency; **P**). *Let $\mathbf{w}$ be fully invested with $\mathbf{w}^\top \boldsymbol{\mu}_e > 0$. Then the minimizer in (8.2) is*

$$a^* = \frac{\mathbf{w}^\top \boldsymbol{\mu}_e}{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}}, \quad (8.4)$$

and

$$\mathfrak{d}(\mathbf{w}; \boldsymbol{\mu}_e, \boldsymbol{\Sigma}) = S_{\max}^2 - S(\mathbf{w})^2, \quad \mathfrak{D}(\mathbf{w}; \boldsymbol{\mu}_e, \boldsymbol{\Sigma}) = 1 - \eta(\mathbf{w})^2 = \sin^2 \theta_{\boldsymbol{\Sigma}}(\mathbf{w}, \mathbf{w}_T). \quad (8.5)$$

Thus the implied-return residual, the angular error, and squared Sharpe inefficiency are the same object in three coordinate systems.

Proof. Let $x = \boldsymbol{\mu}_e$ and $y = \boldsymbol{\Sigma} \mathbf{w}$. In the $\boldsymbol{\Sigma}^{-1}$ -inner product,

$$\langle x, y \rangle_{\boldsymbol{\Sigma}^{-1}} = x^\top \boldsymbol{\Sigma}^{-1} y = \boldsymbol{\mu}_e^\top \mathbf{w} = \mathbf{w}^\top \boldsymbol{\mu}_e, \quad \|y\|_{\boldsymbol{\Sigma}^{-1}}^2 = \mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}.$$

The least-squares projection of x onto the positive ray spanned by y has coefficient (8.4), which is positive by assumption. Therefore

$$\mathfrak{d} = \|x\|_{\boldsymbol{\Sigma}^{-1}}^2 - \frac{\langle x, y \rangle_{\boldsymbol{\Sigma}^{-1}}^2}{\|y\|_{\boldsymbol{\Sigma}^{-1}}^2} = \boldsymbol{\mu}_e^\top \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_e - \frac{(\mathbf{w}^\top \boldsymbol{\mu}_e)^2}{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}} = S_{\max}^2 - S(\mathbf{w})^2.$$

Dividing by $S_{\max}^2$ and applying Theorem 3.7 gives (8.5). $\square$

Corollary 8.5 (Near-optimality tubes; **P**). *Fix $\boldsymbol{\Sigma} \succ 0$ and a portfolio $\mathbf{w}$ with $\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w} > 0$. The set*

$$\mathcal{T}_\delta(\mathbf{w}; \boldsymbol{\Sigma}) := \{\boldsymbol{\mu}_e : \mathfrak{D}(\mathbf{w}; \boldsymbol{\mu}_e, \boldsymbol{\Sigma}) \leq \delta\}$$

is exactly the cone around the ray $\{a \boldsymbol{\Sigma} \mathbf{w} : a > 0\}$ with aperture $\arcsin \sqrt{\delta}$ in the $\boldsymbol{\Sigma}^{-1}$ geometry. Hence exact optimality is the zero-aperture cone, while approximate optimality is a positive-volume tube around the same ray. If a prior density for $\boldsymbol{\mu}_e$ is bounded on bounded $\boldsymbol{\Sigma}^{-1}$ -balls, then for small δ the probability of $\mathcal{T}_\delta$ inside any fixed ball is of order $\delta^{(N-1)/2}$.

Proof. By Theorem 8.4, $\mathfrak{D} = \sin^2 \angle_{\boldsymbol{\Sigma}^{-1}}(\boldsymbol{\mu}_e, \boldsymbol{\Sigma} \mathbf{w})$, so the sublevel set is the stated cone. In N dimensions a cone of angular radius α has solid angle proportional to α^{N-1} as $\alpha \downarrow 0$. Since $\alpha = \arcsin \sqrt{\delta} \sim \sqrt{\delta}$, bounded-density probabilities inside fixed-radius balls scale as $\delta^{(N-1)/2}$. $\square$

Remark 8.6 (Why this defect is useful). *The defect $\mathfrak{D}_\Phi$ is simultaneously a diagnostic, a test target, and an optimization objective. As a diagnostic, it reports exactly how far the heuristic's implied premia $a^* \boldsymbol{\Sigma} \mathbf{w}_\Phi$ are from the estimated premia. As a test target, it is equivalent to the zero-alpha restriction in Remark 3.6. As an HPO objective, it ranks heuristic families without pretending that every member is globally Markowitz-optimal.*

8.3 Tree HPO: local scores as stochastic kernels

HRP, RA-HRP, Schur-RA-HRP, and their interpolations are all examples of a common object: a tree-indexed product of local binary allocation kernels.

Definition 8.7 (Positive score-tree HPO). *Fix a binary tree $\mathcal{T}$ with leaf set $\{1, \dots, N\}$. At each internal node v , let $L(v)$ and $R(v)$ denote its left and right child clusters. A positive score-tree HPO rule assigns scores*

$$G_v(L(v)) > 0, \quad G_v(R(v)) > 0,$$

and local split

$$\beta_v := \frac{G_v(L(v))}{G_v(L(v)) + G_v(R(v))}. \quad (8.6)$$

For a leaf i , let $\mathcal{P}(i)$ be the internal nodes on the root-to-leaf path and let $\eta_v(i) = 1$ if i descends through the left branch of v and 0 otherwise. The resulting leaf weight is

$$w_i^\Phi = \prod_{v \in \mathcal{P}(i)} \beta_v^{\eta_v(i)} (1 - \beta_v)^{1 - \eta_v(i)}. \quad (8.7)$$

The main rules correspond to different choices of G_v . HRP uses $G_v(K) = \tilde{V}_K^{-1}$. RA-HRP uses $G_v(K) = S^+(K)$. Schur-RA-HRP uses conditional scores $G_v(K) = S_{\text{Sch}}^+(K \mid K_v^c)$, where K_v^c is the sibling cluster at node v . Unit-free HRP–RA-HRP interpolation is obtained by interpolating the local branch probabilities β_v , not by adding raw scores with incompatible units.

Theorem 8.8 (Exact coincidence for score-tree HPO; $\mathbf{P}$). *Fix $\theta = (\mu_e, \Sigma)$ and suppose $x_T := \Sigma^{-1} \mu_e$ has strictly positive subtree masses*

$$M_T(K) := \mathbf{1}_K^\top x_T > 0$$

for every cluster K appearing in $\mathcal{T}$. Let Φ be a positive score-tree HPO rule on $\mathcal{T}$. Then $\mathbf{w}^\Phi$ coincides with the tangency composition $\mathbf{w}_T = x_T / (\mathbf{1}^\top x_T)$ if and only if, at every internal node v ,

$$\frac{G_v(L(v))}{G_v(R(v))} = \frac{M_T(L(v))}{M_T(R(v))}. \quad (8.8)$$

Equivalently, every heuristic local split must equal the exact tangency mass split:

$$\beta_v = \frac{M_T(L(v))}{M_T(L(v)) + M_T(R(v))}.$$

Proof. If $\mathbf{w}^\Phi = \mathbf{w}_T$, then the total portfolio mass assigned to the left child of any node v divided by the total mass assigned to its sibling equals $M_T(L(v))/M_T(R(v))$. But under the recursive rule, the same ratio is $\beta_v/(1 - \beta_v) = G_v(L(v))/G_v(R(v))$, proving necessity. Conversely, if (8.8) holds at every node, then the heuristic sends exactly the same fraction of each parent subtree mass to each child as the tangency portfolio. Starting from root mass one, induction down the tree shows that every leaf receives the tangency mass, hence $\mathbf{w}^\Phi = \mathbf{w}_T$. $\square$

Remark 8.9 (RA-HRP as one score choice). *Theorem 8.8 is the abstract version of Theorem 7.15. RA-HRP chooses a particular economically interpretable score, the floored local cluster Sharpe ratio. Exact optimality requires that this score ratio reproduce the exact subtree mass ratio of $\Sigma^{-1} \mu_e$. Schur-RA-HRP changes the score in order to move the local ratio closer to the Schur-adjusted tangency mass, while still avoiding the global inverse.*

8.4 The exact Schur-tangency split

The score-tree theorem identifies the target local mass ratio. The next theorem writes that target in Schur form. This is the return-aware analogue of the GMV Schur identity used earlier for HRP.

Theorem 8.10 (Exact Schur-tangency bisection; $\mathbf{P}$). *Let $C = A \cup B$ and partition Σ_C and μ_C conformably:*

$$\Sigma_C = \begin{pmatrix} \Sigma_A & \Sigma_{AB} \\ \Sigma_{BA} & \Sigma_B \end{pmatrix}, \quad \mu_C = \begin{pmatrix} \mu_A \\ \mu_B \end{pmatrix}.$$

Define Schur complements and Schur-adjusted premia

$$\begin{aligned} \mathbf{S}_A &:= \Sigma_A - \Sigma_{AB} \Sigma_B^{-1} \Sigma_{BA}, & \nu_{A|B} &:= \mu_A - \Sigma_{AB} \Sigma_B^{-1} \mu_B, \\ \mathbf{S}_B &:= \Sigma_B - \Sigma_{BA} \Sigma_A^{-1} \Sigma_{AB}, & \nu_{B|A} &:= \mu_B - \Sigma_{BA} \Sigma_A^{-1} \mu_A. \end{aligned}$$

Then the unnormalized tangency vector $x_C = \Sigma_C^{-1} \mu_C$ satisfies

$$(x_C)_A = \mathbf{S}_A^{-1} \nu_{A|B}, \quad (x_C)_B = \mathbf{S}_B^{-1} \nu_{B|A}. \quad (8.9)$$

Consequently the exact tangency mass allocated to A at this split is

$$\beta_A^T = \frac{\mathbf{1}_A^\top \mathbf{S}_A^{-1} \nu_{A|B}}{\mathbf{1}_A^\top \mathbf{S}_A^{-1} \nu_{A|B} + \mathbf{1}_B^\top \mathbf{S}_B^{-1} \nu_{B|A}}, \quad (8.10)$$

whenever the denominator is positive and both masses are positive.

Proof. Apply the block-inverse formula. The A block of $\Sigma_C^{-1} \mu_C$ is

$$\mathbf{S}_A^{-1} \mu_A - \mathbf{S}_A^{-1} \Sigma_{AB} \Sigma_B^{-1} \mu_B = \mathbf{S}_A^{-1} \nu_{A|B}.$$

The B block is symmetric. Summing coordinates in each block gives the mass formula. $\square$

Corollary 8.11 (The HPO approximation ladder; $\mathbf{P}$). *At a node $C = A \cup B$, exact local Markowitz/tangency allocation consumes four objects: the Schur covariance $\mathbf{S}_K$, the Schur-adjusted premium $\nu_{K|K^c}$, the inverse $\mathbf{S}_K^{-1}$, and the resulting subtree mass $\mathbf{1}_K^\top \mathbf{S}_K^{-1} \nu_{K|K^c}$. The main hierarchical HPO rules are obtained by deleting or compressing these objects in stages:*

$$\begin{aligned} \text{Exact Schur tangency} &: G(K) = \mathbf{1}_K^\top \mathbf{S}_K^{-1} \nu_{K|K^c}, \\ \text{Schur-RA-HRP} &: G(K) = S_{\text{Sch}}^+(K | K^c), \\ \text{RA-HRP} &: G(K) = S^+(K), \\ \text{HRP} &: G(K) = \tilde{V}_K^{-1}. \end{aligned}$$

Thus the “heuristic” label describes a precise algebraic operation: replace exact Schur-tangency masses by stable positive scores whose estimation error is smaller and whose economic meaning is transparent.

Proof. The first line is Theorem 8.10. The remaining lines are definitions of the corresponding score-tree rules. The claim is therefore an identity of information sets: each step removes either Schur conditioning, premium conditioning, local inversion, or premium information, and then renormalizes the remaining positive scores through (8.6). $\square$

8.5 Sampling error, approximation error, and the HPO bias–variance identity

HPO is useful only if the information it deletes is sufficiently noisy. The next result gives the exact quadratic accounting identity behind this statement.

Theorem 8.12 (Exact mean–variance bias–variance decomposition for estimated HPO; $\mathbf{P}$). *Fix $\gamma > 0$ and define the budget-constrained mean–variance objective*

$$U_\gamma(\mathbf{w}) = \mathbf{w}^\top \mu_e - \frac{\gamma}{2} \mathbf{w}^\top \Sigma \mathbf{w}, \quad \mathbf{1}^\top \mathbf{w} = 1.$$

Let $\mathbf{w}_\gamma^$ be its unique maximizer on the affine budget hyperplane, and let $\widehat{\mathbf{w}}_\Phi$ be any random fully invested portfolio produced by an estimated heuristic rule. If $\mathbb{E} \|\widehat{\mathbf{w}}_\Phi\|^2 < \infty$, then*

$$\mathbb{E}[U_\gamma(\mathbf{w}_\gamma^*) - U_\gamma(\widehat{\mathbf{w}}_\Phi)] = \frac{\gamma}{2} \left(\|\mathbb{E}[\widehat{\mathbf{w}}_\Phi] - \mathbf{w}_\gamma^*\|_\Sigma^2 + \text{tr}(\Sigma \text{Cov}(\widehat{\mathbf{w}}_\Phi)) \right). \quad (8.11)$$

Proof. The first-order condition for the budget-constrained optimum is

$$\mu_e - \gamma \Sigma \mathbf{w}_\gamma^* = \lambda \mathbf{1}$$

for some Lagrange multiplier λ . For any fully invested $\mathbf{w}$, $\mathbf{1}^\top(\mathbf{w} - \mathbf{w}_\gamma^*) = 0$, so the linear term in the second-order expansion vanishes:

$$U_\gamma(\mathbf{w}_\gamma^*) - U_\gamma(\mathbf{w}) = \frac{\gamma}{2}(\mathbf{w} - \mathbf{w}_\gamma^*)^\top \Sigma (\mathbf{w} - \mathbf{w}_\gamma^*).$$

Apply this identity to $\widehat{\mathbf{w}}_\Phi$ and take expectations. The standard Hilbert-space bias–variance decomposition in the Σ norm gives

$$\mathbb{E}\|\widehat{\mathbf{w}}_\Phi - \mathbf{w}_\gamma^*\|_\Sigma^2 = \|\mathbb{E}[\widehat{\mathbf{w}}_\Phi] - \mathbf{w}_\gamma^*\|_\Sigma^2 + \text{tr}(\Sigma \text{Cov}(\widehat{\mathbf{w}}_\Phi)).$$

□

Remark 8.13 (When a heuristic beats the plug-in optimizer). *Equation (8.11) is the formal answer to the Markowitz optimization enigma. A heuristic can be farther from the population optimum in bias but closer in expected realized utility if it has sufficiently lower estimation variance. HRP, RA-HRP, inverse volatility, ERC, and constrained/shrunk optimizers all exploit this identity in different ways. HRP and RA-HRP reduce variance by replacing global inversion with local score maps; Schur-RA-HRP spends some variance budget on local block inversions to recover cross-cluster hedge information; full MVO spends the most variance budget by estimating and inverting the complete covariance matrix and using the full mean vector.*

8.6 The master HPO objective

The preceding results suggest a practical HPO loss for selecting among heuristic families:

$$\mathcal{L}_{\text{HPO}}(\Phi) = \widehat{\mathcal{D}}_\Phi + \lambda_{\text{est}} \text{tr}(\widehat{\Sigma \text{Cov}(\widehat{\mathbf{w}}_\Phi)}) + \lambda_{\text{tc}} \|\widehat{\mathbf{w}}_\Phi - \mathbf{w}_-\|_1 + \lambda_{\text{top}} d_{\mathcal{T}}(\widehat{\mathcal{T}}, \mathcal{T}_-), \quad (8.12)$$

where $\widehat{\mathcal{D}}_\Phi$ measures implied-return misspecification, the second term measures weight estimation variance, the third measures turnover, and the fourth measures tree instability when the rule is hierarchical. Equation (8.12) is not a new theorem; it is the operational synthesis of the theorems above. It turns HPO into a disciplined model-selection problem rather than a collection of unrelated allocation recipes.

Remark 8.14 (How this section changes the interpretation of the paper). *The earlier sections answer, for each named rule, “when is it exactly optimal?” This section answers the broader question, “what is a heuristic portfolio rule mathematically?” It is an information-restricted map; its population error is an implied-return projection residual; its Sharpe loss is the squared sine of an angle; its hierarchical forms are products of local binary kernels; its exact nodewise target is a Schur-tangency mass; and its out-of-sample value is governed by a bias–variance–cost trade-off. That is the mathematics of HPO.*

9 From HPO to Reinforcement Learning Portfolio Optimization (RLPO)

The preceding section is static: given a statewise estimate $(\widehat{\boldsymbol{\mu}}_{e,t}, \widehat{\boldsymbol{\Sigma}}_t)$, possibly together with a tree $\widehat{\mathcal{T}}_t$, a heuristic map produces a portfolio $\mathbf{w}_t = \Phi(s_t)$. RLPO starts from the same action object but changes the question. It asks how a portfolio agent should rebalance through time when today’s action affects future states, trading costs, inventory, drawdown, tax lots, liquidity, and regime exposure. Thus HPO is the *myopic policy face* of RLPO; RLPO is the dynamic control layer that learns when the continuation value of deviating from the HPO prior compensates for its instantaneous optimality defect.

9.1 Controlled market formulation

Definition 9.1 (RLPO environment). *An RLPO environment is a discounted controlled market*

$$\mathfrak{M} = (\mathcal{S}, \{\mathcal{A}(s)\}_{s \in \mathcal{S}}, P, R, \gamma), \quad 0 \leq \gamma < 1,$$

where $s_t \in \mathcal{S}$ is the observed market state, $\mathcal{A}(s_t) \subseteq \Delta^{N-1}$ is the feasible portfolio set, $P(ds' | s, w)$ is the controlled transition law, $R(s, w, s')$ is the one-period reward, and γ is the discount factor. A Markov policy π assigns a distribution $\pi(\cdot | s)$ on $\mathcal{A}(s)$. Its value is

$$V^\pi(s) = \mathbb{E}^\pi \left[\sum_{t \geq 0} \gamma^t R(s_t, w_t, s_{t+1}) \mid s_0 = s \right]. \quad (9.1)$$

Writing $r(s, w) := \mathbb{E}[R(s, w, s') \mid s, w]$, the optimal value satisfies the Bellman equation

$$V^*(s) = \sup_{w \in \mathcal{A}(s)} \{r(s, w) + \gamma \mathbb{E}[V^*(s') \mid s, w]\}. \quad (9.2)$$

This is the usual dynamic-programming formulation of sequential control (Bellman, 1957; Bertsekas, 2017; Sutton and Barto, 2018). In portfolio applications the state may include lagged returns, macro variables, volatility estimates, factor exposures, the current dendrogram, the previous portfolio $\mathbf{w}_-$, turnover budget, margin constraints, and execution-cost estimates. The reward may be raw excess return, mean–variance utility, downside-risk-adjusted return, a drawdown-penalized objective, or a transaction-cost-adjusted utility as in dynamic trading models with predictable returns and frictions (Gârleanu and Pedersen, 2013).

Definition 9.2 (HPO-induced policy). *Let $G : \mathcal{S} \rightarrow \mathcal{Z}$ be a state summarizer producing the information consumed by an HPO rule, for example*

$$G(s_t) = (\hat{\mu}_{e,t}, \hat{\Sigma}_t), \quad G(s_t) = (\hat{\mu}_{e,t}, \hat{\Sigma}_t, \hat{T}_t),$$

or the reduced statistics used by EW, IV, ERC, HRP, RA-HRP, or Schur-RA-HRP. Any feasible HPO map $\Phi : \mathcal{Z} \rightarrow \Delta^{N-1}$ induces the deterministic stationary policy

$$\pi_\Phi(\cdot \mid s) = \delta_{\Phi(G(s))}, \quad \Phi(G(s)) \in \mathcal{A}(s). \quad (9.3)$$

Proposition 9.3 (HPO policies as stationary RLPO policies; **P**). *Every feasible measurable HPO map induces an admissible deterministic stationary RLPO policy through (9.3). If $\gamma = 0$, the reward has no next-state dependence, and $R(s, w, s') = u(G(s), w)$ equals the one-period static objective used to define the HPO map, then the RLPO problem reduces state by state to the corresponding static HPO problem. In particular, if*

$$\Phi(G(s)) \in \arg \max_{w \in \mathcal{A}(s)} u(G(s), w),$$

then π_Φ is optimal for the RLPO environment with $\gamma = 0$.

Proof. Measurability and feasibility imply that $s \mapsto \delta_{\Phi(G(s))}$ is a valid stationary Markov policy. If $\gamma = 0$, then (9.1) reduces to the one-period conditional objective $V^\pi(s) = \mathbb{E}[R(s, w, s') \mid s, w]$. Under the stated reward specification this is $u(G(s), w)$, so the Bellman maximization is exactly $\max_{w \in \mathcal{A}(s)} u(G(s), w)$. A statewise optimizer $\Phi(G(s))$ therefore induces an optimal deterministic policy. $\square$

Remark 9.4 (The conceptual hierarchy). *HPO answers the question “given the current estimated pair $(\hat{\mu}_t, \hat{\Sigma}_t)$ and any reduced structure such as a tree, which portfolio map is locally justified?” RLPO answers the dynamic question “given the current state and the effect of today’s action on tomorrow’s state, when should the agent deviate from that locally justified map?” The action space is the same simplex. The difference is the objective: HPO optimizes an instantaneous geometric criterion, while RLPO optimizes the Bellman criterion (9.2).*

9.2 Continuation value versus myopic HPO defect

At state s , write

$$\hat{\boldsymbol{\mu}}_e(s), \quad \hat{\boldsymbol{\Sigma}}(s), \quad \mathbf{w}_\Phi(s) := \Phi(G(s))$$

for the estimated excess-premium vector, covariance matrix, and HPO baseline. The statewise implied-return defect is

$$\mathfrak{d}_s(w) := \min_{a>0} \left\| \hat{\boldsymbol{\mu}}_e(s) - a\hat{\boldsymbol{\Sigma}}(s)w \right\|_{\hat{\boldsymbol{\Sigma}}(s)^{-1}}^2. \quad (9.4)$$

By Theorem 8.4, this is not an ad hoc regularizer: it is exactly

$$\mathfrak{d}_s(w) = S_{\max}^2(s) - S_s^2(w),$$

the statewise squared Sharpe inefficiency measured in the metric of the estimated covariance matrix.

Proposition 9.5 (Dynamic deviation principle; **P**). *Fix a state s , a baseline HPO action $w_\Phi = \mathbf{w}_\Phi(s)$, and a candidate action $w \in \mathcal{A}(s)$. Decompose the one-period reward as*

$$r(s, w) = r_0(s, w) - c_s(w, w_-) - \lambda_{\text{HPO}} \mathfrak{d}_s(w),$$

where r_0 is the trading reward before HPO regularization, c_s is an implementation cost relative to the previous portfolio w_- , and $\lambda_{\text{HPO}} \geq 0$. Define the instantaneous loss of choosing w rather than the HPO action by

$$\begin{aligned} \Delta_{\text{inst}}(s, w \mid w_\Phi) &:= r_0(s, w_\Phi) - r_0(s, w) + c_s(w, w_-) - c_s(w_\Phi, w_-) \\ &\quad + \lambda_{\text{HPO}} \{\mathfrak{d}_s(w) - \mathfrak{d}_s(w_\Phi)\}, \end{aligned} \quad (9.5)$$

and the continuation-value gain by

$$\Delta_{\text{cont}}(s, w \mid w_\Phi) := \gamma \mathbb{E}[V^*(s') \mid s, w] - \gamma \mathbb{E}[V^*(s') \mid s, w_\Phi]. \quad (9.6)$$

Then w weakly improves on the HPO action in Bellman value if and only if

$$\Delta_{\text{cont}}(s, w \mid w_\Phi) \geq \Delta_{\text{inst}}(s, w \mid w_\Phi). \quad (9.7)$$

Proof. Subtract the Bellman score of w_Φ from that of w :

$$\{r(s, w) + \gamma \mathbb{E}[V^*(s') \mid s, w]\} - \{r(s, w_\Phi) + \gamma \mathbb{E}[V^*(s') \mid s, w_\Phi]\}.$$

Substituting the reward decomposition and rearranging gives $\Delta_{\text{cont}}(s, w \mid w_\Phi) - \Delta_{\text{inst}}(s, w \mid w_\Phi)$. Nonnegativity is exactly (9.7). $\square$

Remark 9.6 (Economic interpretation). *Equation (9.7) is the precise mathematical statement of the HPO–RLPO relation. A dynamic policy should not deviate from a locally justified HPO portfolio merely because a neural policy can output different weights. It should deviate only when the expected continuation-value gain caused by that deviation is large enough to pay for three quantities: the lost instantaneous reward, the incremental turnover or implementation cost, and the additional implied-return defect. RLPO is therefore not a replacement for HPO. It is a disciplined dynamic policy-improvement layer over an HPO prior.*

9.3 The exact value gap of an HPO policy

Proposition 9.5 is a one-step comparison. The classical performance-difference identity upgrades it to an exact global accounting of how much value the myopic HPO policy leaves on the table, in the same advantage units, summed along the trajectory of whatever policy challenges it.

Proposition 9.7 (Exact value gap of the HPO policy; $\mathbf{P}$, identity after Kakade and Langford, 2002). *Assume $|R(s, w, s')| \leq \bar{R} < \infty$ and $\gamma \in [0, 1)$, so all values are bounded by $\bar{R}/(1 - \gamma)$. Define the HPO advantage of action w at state s by*

$$A^{\pi_\Phi}(s, w) := r(s, w) + \gamma \mathbb{E}[V^{\pi_\Phi}(s') \mid s, w] - V^{\pi_\Phi}(s), \quad (9.8)$$

which satisfies $A^{\pi_\Phi}(s, \mathbf{w}_\Phi(s)) = 0$ by Bellman consistency of the deterministic stationary policy π_Φ . Then for every Markov policy π and initial state s_0 ,

$$V^\pi(s_0) - V^{\pi_\Phi}(s_0) = \mathbb{E}^\pi \left[\sum_{t \geq 0} \gamma^t A^{\pi_\Phi}(s_t, w_t) \mid s_0 \right] = \frac{1}{1 - \gamma} \mathbb{E}_{(s, w) \sim d_{s_0}^\pi} [A^{\pi_\Phi}(s, w)], \quad (9.9)$$

where $d_{s_0}^\pi := (1 - \gamma) \sum_{t \geq 0} \gamma^t \text{Law}^\pi((s_t, w_t) \mid s_0)$ is the normalized discounted occupancy measure. In particular, taking $\pi = \pi^$,*

$$V^*(s_0) - V^{\pi_\Phi}(s_0) = \frac{1}{1 - \gamma} \mathbb{E}_{(s, w) \sim d_{s_0}^{\pi^*}} [A^{\pi_\Phi}(s, w)] : \quad (9.10)$$

the exact price of acting myopically is the occupancy-weighted HPO advantage, and only the states where the optimal policy actually deviates from the heuristic contribute.

Proof. Substituting (9.8) and using the tower property under π ,

$$\mathbb{E}^\pi \left[\sum_{t=0}^T \gamma^t A^{\pi_\Phi}(s_t, w_t) \right] = \mathbb{E}^\pi \left[\sum_{t=0}^T \gamma^t R_t \right] + \mathbb{E}^\pi \left[\sum_{t=0}^T \left(\gamma^{t+1} V^{\pi_\Phi}(s_{t+1}) - \gamma^t V^{\pi_\Phi}(s_t) \right) \right].$$

The second sum telescopes to

$$\gamma^{T+1} \mathbb{E}^\pi [V^{\pi_\Phi}(s_{T+1})] - V^{\pi_\Phi}(s_0).$$

Boundedness sends the discounted terminal term to zero as $T \rightarrow \infty$, leaving $V^\pi(s_0) - V^{\pi_\Phi}(s_0)$. Dominated convergence justifies the limit, and the occupancy form is the bounded-Fubini rewriting of the discounted sum. Bellman consistency,

$$V^{\pi_\Phi}(s) = r(s, \mathbf{w}_\Phi(s)) + \gamma \mathbb{E}[V^{\pi_\Phi}(s') \mid s, \mathbf{w}_\Phi(s)],$$

gives $A^{\pi_\Phi}(s, \mathbf{w}_\Phi(s)) = 0$. □

Corollary 9.8 (ε -myopic optimality and shaped localization; $\mathbf{P}$). *(i) If no feasible one-step deviation gains more than $\varepsilon \geq 0$ against the HPO policy's own continuation value, i.e.*

$$A^{\pi_\Phi}(s, w) \leq \varepsilon \quad \text{for every } s \text{ and every } w \in \mathcal{A}(s),$$

then

$$V^*(s) \leq V^{\pi_\Phi}(s) + \frac{\varepsilon}{1 - \gamma} \quad \text{for every } s.$$

(ii) Under the reward decomposition of Proposition 9.5, the advantage splits exactly as

$$\begin{aligned} A^{\pi_\Phi}(s, w) &= \tilde{\Delta}_{\text{cont}}(s, w \mid \mathbf{w}_\Phi) - \Delta_{\text{inst}}(s, w \mid \mathbf{w}_\Phi), \\ \tilde{\Delta}_{\text{cont}}(s, w \mid \mathbf{w}_\Phi) &:= \gamma \mathbb{E}[V^{\pi_\Phi}(s') \mid s, w] - \gamma \mathbb{E}[V^{\pi_\Phi}(s') \mid s, \mathbf{w}_\Phi(s)], \end{aligned}$$

with the same instantaneous-loss term (9.5) and the continuation gain evaluated under V^{π_Φ} . Hence (9.10) is the occupancy-weighted integral of exactly the quantity the deviation principle tests pointwise; and because the λ_{HPO} term sits inside Δ_{inst} , every deviation along the optimal trajectory pays its incremental implied-return defect $\mathfrak{d}_s(w) - \mathfrak{d}_s(\mathbf{w}_\Phi(s))$ state by state.

Proof. (i) Bound the integrand in (9.10) by ε . (ii) Subtract and add $r(s, \mathbf{w}_\Phi(s))$ in (9.8), use Bellman consistency of π_Φ to cancel $V^{\pi_\Phi}(s)$, and substitute the reward decomposition; the algebra is that of Proposition 9.5 with V^{π_Φ} in place of V^* . $\square$

This closes the logical loop between the two layers. Statically, “never optimal, rarely costly” is Proposition 3.8 plus Theorem 3.9. Dynamically, the analogue is (9.10) plus Corollary 9.8(i): the HPO policy forfeits at most the discounted occupancy average of its statewise advantage gaps, and when continuation values never reward deviation by more than ε , acting myopically costs at most $\varepsilon/(1-\gamma)$ — a bound expressed entirely in the advantage units that the deviation principle already prices.

9.4 HPO-regularized RLPO objectives

The preceding proposition suggests the following regularized RLPO objective for a parametric policy π_θ :

$$J_{\text{RLPO}}(\theta) := \mathbb{E}^{\pi_\theta} \sum_{t \geq 0} \gamma^t \left[w_t^\top r_{e,t+1} - c_t(w_t, w_{t-1}) - \lambda_{\text{risk}} \rho_t(w_t) - \lambda_{\text{HPO}} \mathfrak{d}(w_t; \hat{\boldsymbol{\mu}}_{e,t}, \hat{\boldsymbol{\Sigma}}_t) - \lambda_{\text{KL}} D_{\text{KL}}(w_t \parallel \mathbf{w}_\Phi(s_t)) \right]. \quad (9.11)$$

Here ρ_t is a risk functional, c_t is the trading-cost model, $\mathfrak{d}$ is the implied-return defect from Definition 8.3, and the KL term keeps the learned policy near the HPO prior unless the data supply enough dynamic evidence to move away from it. If λ_{HPO} and λ_{KL} are large, the solution collapses toward the HPO baseline. If they are zero, the problem becomes an unconstrained RLPO objective. Intermediate values implement dynamic residual learning around a mathematically interpretable prior.

A particularly stable actor parametrization is the convex-residual policy

$$w_t^\theta = (1 - \lambda_t^\theta) \mathbf{w}_\Phi(s_t) + \lambda_t^\theta \psi_\theta(s_t), \quad \lambda_t^\theta \in [0, 1], \quad \psi_\theta(s_t) \in \Delta^{N-1}. \quad (9.12)$$

This guarantees long-only full investment by construction. The scalar λ_t^θ has a direct interpretation: the state-dependent degree of trust placed on dynamic RLPO relative to static HPO. A conservative agent learns mostly λ_t^θ ; a more expressive agent learns both λ_t^θ and the residual portfolio ψ_θ .

9.5 RA-HRP as a hierarchical policy prior

RA-HRP is especially natural inside RLPO because the policy is already factorized over a tree. Instead of asking an actor network to output N unrelated weights, one can ask it to output local split probabilities. Fix a tree $\mathcal{T}$. For each internal node v , let $\beta_v^{\text{RA}}(s)$ denote the RA-HRP split fraction at state s , and let $a_v^\theta(s)$ be a learned dynamic residual. Define the actor by

$$\text{logit } \beta_v^\theta(s) = \text{logit } \beta_v^{\text{RA}}(s) + a_v^\theta(s), \quad \beta_v^\theta(s) \in (0, 1). \quad (9.13)$$

For leaf i , with path $\mathcal{P}(i)$ and branch indicator $\eta_v(i)$, the induced portfolio is

$$w_i^\theta(s) = \prod_{v \in \mathcal{P}(i)} \left(\beta_v^\theta(s) \right)^{\eta_v(i)} \left(1 - \beta_v^\theta(s) \right)^{1 - \eta_v(i)}. \quad (9.14)$$

Proposition 9.9 (Budget feasibility and baseline recovery of the RA-HRP actor; **P**). *For any collection of split probabilities $\{\beta_v^\theta(s) \in (0, 1) : v \in \mathcal{V}_{\text{int}}\}$, the weights in (9.14) are strictly positive and sum to one. If $a_v^\theta(s) = 0$ for every internal node, then $w^\theta(s) = w^{\text{RA}}(s)$ exactly. Moreover, for two hierarchical policies A and B on the same tree,*

$$D_{\text{KL}}(w^A(s) \parallel w^B(s)) = \sum_{v \in \mathcal{V}_{\text{int}}} W_v^A(s) d_{\text{KL}}(\beta_v^A(s) \parallel \beta_v^B(s)), \quad (9.15)$$

where $W_v^A(s)$ is the capital mass reaching node v under policy A and d_{KL} is the binary KL divergence.

Proof. Budget feasibility follows by induction down the tree: the root has mass one, and every internal node sends fractions β_v^θ and $1 - \beta_v^\theta$ to its two children, so total mass is conserved at each split. Positivity follows from $\beta_v^\theta \in (0, 1)$. If all residuals vanish in (9.13), every local split equals the RA-HRP split, hence the product representation recovers RA-HRP. The KL identity is the same pathwise decomposition proved in Theorem 7.22: expand the log ratio of leaf weights as a sum of log ratios of local branch probabilities and exchange the sums over leaves and internal nodes. $\square$

Equation (9.15) is the main reason RA-HRP is a better RLPO prior than a black-box weight vector. The global distance between the learned dynamic policy and the baseline decomposes into node-level binary divergences. Root nodes receive large capital weights and therefore dominate global deviation; deep nodes allow fine local tilts. This gives a precise attribution of the actor's dynamic behavior: one can identify whether RLPO is overriding the top-level sector allocation, a mid-level cluster split, or a leaf-level security choice.

Corollary 9.10 (Myopic policy gradient at the hierarchical prior; **P**). *Fix a state s and evaluate the myopic Sharpe objective $S_s(w) := w^\top \hat{\mu}_e(s) / (w^\top \hat{\Sigma}(s) w)^{1/2}$ on the hierarchical actor (9.13) at the prior, $a_v^\theta(s) = 0$ for all v , so that $w^\theta(s) = w^{\text{RA}}(s)$. Then for every internal node v ,*

$$\left. \frac{\partial S_s(w^\theta(s))}{\partial a_v^\theta} \right|_{a^\theta=0} = \frac{\beta_v^{\text{RA}}(s)(1 - \beta_v^{\text{RA}}(s))}{\sigma_p(w^{\text{RA}}(s))} \alpha_v^{\text{RA}}(s), \quad (9.16)$$

where $\alpha_v^{\text{RA}}(s) = g_v^{\text{RA}}(s)^\top [\hat{\mu}_e(s) - a^*(w^{\text{RA}}(s)) \hat{\Sigma}(s) w^{\text{RA}}(s)]$ is the node alpha of Theorem 7.27 computed with the RA-HRP weights and splits in the node-switch vectors of Lemma 7.26. Hence the myopic component of the policy gradient at the prior is, coordinate by coordinate, the nodewise alpha left unexplained by the prior, damped by the logistic variance $\beta_v(1 - \beta_v)$: the actor receives its first learning signal exactly where the baseline misprices a child switch, and that signal is automatically attenuated at saturated splits.

Proof. At $a^\theta = 0$, $\partial \beta_v^\theta / \partial a_v^\theta = \beta_v^{\text{RA}}(1 - \beta_v^{\text{RA}})$ from the logit parametrization, so by the product representation (9.14), $\partial w_i^\theta / \partial a_v^\theta = \beta_v^{\text{RA}}(1 - \beta_v^{\text{RA}}) w_i^{\text{RA}} (\eta_v(i) / \beta_v^{\text{RA}} - (1 - \eta_v(i)) / (1 - \beta_v^{\text{RA}}))$, which is $\beta_v^{\text{RA}}(1 - \beta_v^{\text{RA}})$ times the node-switch vector g_v^{RA} of Lemma 7.26 built from the RA weights and splits. Contract with $\nabla S_s(w) = [\hat{\mu}_e(s) - a^*(w) \hat{\Sigma}(s) w] / \sigma_p(w)$ as in Theorem 7.27. $\square$

A Schur-RA-HRP prior is obtained by replacing $\beta_v^{\text{RA}}(s)$ in (9.13) by the conditional-risk split from Definition 7.17. The actor then learns deviations around a baseline that already incorporates local cross-cluster hedging while still avoiding a global inverse. This gives the natural approximation ladder

$$\begin{array}{c} \text{exact Bellman policy} \\ \downarrow \\ \text{RL residual over Schur-RA-HRP} \\ \downarrow \\ \text{RL residual over RA-HRP} \\ \downarrow \\ \text{RL residual over HRP.} \end{array} \quad (9.17)$$

9.6 Tree topology, regime shifts, and the dynamic open problem

The fixed-tree theory in Sections 7 and 8 is deliberately sharp because it isolates the local score map from the discrete topology. RLPO makes the open problem unavoidable: the state may

include a time-varying tree $\mathcal{T}_t$, and the policy must decide whether changes in the estimated dendrogram are signal, noise, or regime transition. A dynamic tree-regularized objective can be written as

$$J_{\text{tree}}(\theta) = J_{\text{RLPO}}(\theta) - \lambda_{\mathcal{T}} \mathbb{E}^{\pi_{\theta}} \sum_{t \geq 0} \gamma^t d_{\mathcal{T}}(\mathcal{T}_t, \mathcal{T}_{t-1}), \quad (9.18)$$

where $d_{\mathcal{T}}$ may be a merge-preservation distance, a cophenetic-matrix distance, a leaf-order rank distance, or a split-probability path distance. The pathwise factorization of hierarchical weights suggests the right mathematical route for future theory: prove uniform convergence of the value of node-split policies over a class of random trees, controlling both continuous score perturbations and discrete topology changes. In this sense the dendrogram-stability diagnostics used for RA-HRP are not merely empirical checks; they are the observable coordinates of the missing uniform-in-tree RLPO theorem.

9.7 Operational blueprint

The HPO–RLPO synthesis leads to a practical architecture.

1. Construct a state s_t containing market features, estimates $(\hat{\mu}_{e,t}, \hat{\Sigma}_t)$, the current tree $\hat{\mathcal{T}}_t$ when hierarchical rules are used, previous weights w_{t-1} , and cost/liquidity variables.
2. Compute a baseline HPO prior $\mathbf{w}_{\Phi}(s_t)$, preferably RA-HRP or Schur-RA-HRP when a stable tree is available.
3. Parametrize the actor either as the convex residual (9.12) or, for hierarchical policies, through node-level residual logits (9.13).
4. Train the policy on the cost-adjusted Bellman objective (9.11), using the HPO defect and KL penalties to prevent economically unjustified deviations.
5. Attribute the learned deviations through (9.15), separating root-level regime allocation from local security-level tilts.

Static HPO object	RLPO counterpart	Interpretation
HPO map $\Phi(G(s))$	deterministic policy $\pi_{\Phi}(\cdot s) = \delta_{\Phi(G(s))}$	every heuristic is a stationary Markov policy
Implied-return defect $\mathfrak{d}_s(w)$	reward-shaping penalty	dynamic deviations must pay for local Sharpe inefficiency
RA-HRP node split β_v^{RA}	actor prior logit $\beta_v^{\theta} = \text{logit } \beta_v^{\text{RA}} + a_v^{\theta}$	RL learns interpretable residuals rather than unrelated weights
Pathwise KL decomposition	tree-level policy regularizer	global policy distance decomposes into node-level deviations
Tree-stability diagnostics	topology regularizer $d_{\mathcal{T}}(\mathcal{T}_t, \mathcal{T}_{t-1})$	regime changes and dendrogram jumps become controlled state variables
Homotopy derivative $\sum_v (s_v - r_v) \alpha_v$	myopic policy-gradient coordinates (9.16)	the actor’s first learning signal is the nodewise alpha the prior leaves unexplained
HPO advantage $A^{\pi_{\Phi}}(s, w)$	performance-difference identity (9.9)	exact discounted price of acting myopically; ε -myopic bound $\varepsilon/(1 - \gamma)$

Table 2: Static-to-dynamic bridge between HPO and RLPO. HPO supplies the myopic geometry and policy priors; RLPO supplies continuation value and dynamic policy improvement.

The central message is therefore simple. HPO gives exact statewise optimality defects. RA-HRP gives structured hierarchical policy priors. RLPO learns dynamic deviations under costs, regimes, and feedback. The three pieces are complementary rather than competing.

10 Transfer: Elliptical Risk Measures and Growth Optimality

Mean–variance optimality is the natural benchmark, but the conditions above would be parochial if they were artifacts of the quadratic objective. They are not.

10.1 Elliptical transfer

Assumption 10.1 (Ellipticity). *The return vector is elliptical, $\mathbf{r} \sim E_N(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \psi)$: the characteristic function is $\phi(\mathbf{t}) = e^{i\mathbf{t}^\top \boldsymbol{\mu}} \psi(\mathbf{t}^\top \boldsymbol{\Sigma} \mathbf{t})$, with finite second moments normalized so $\boldsymbol{\Sigma}$ is the covariance matrix.*

Theorem 10.2 (Risk-measure transfer; $\mathbf{P}$, assembling classical results). *Under Assumption 10.1, let ϱ be any risk measure on portfolio losses $L_{\mathbf{w}} = -\mathbf{w}^\top \mathbf{r}$ that is law-invariant, positively homogeneous, and cash-translative ($\varrho(L-c) = \varrho(L) - c$), with $\varrho(Z) \in (0, \infty)$ for Z the standardized one-dimensional marginal. Then*

$$\varrho(L_{\mathbf{w}}) = -\mathbf{w}^\top \boldsymbol{\mu} + \sqrt{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}} \varrho(Z), \quad (10.1)$$

so for every target mean the ϱ -minimizing fully invested portfolio is the variance-minimizing one, the mean- ϱ frontier coincides with the mean–variance frontier, and the generalized-Sharpe maximizer $\arg \max_{\mathbf{w}} \mathbf{w}^\top \boldsymbol{\mu}_e / \varrho(L_{\mathbf{w}} + r_f)$ is the maximum-Sharpe portfolio. Consequently every coincidence condition of Sections 4–7 is simultaneously the condition for mean–CVaR $_{\alpha}$ optimality at every level α , and frontier portfolios are optimal for every nonsatiated risk-averse expected-utility investor.

Proof. Ellipticity implies the location–scale property of linear functionals: $\mathbf{w}^\top \mathbf{r} \stackrel{d}{=} \mathbf{w}^\top \boldsymbol{\mu} + \sqrt{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}} Z$ with Z standardized, the same law for all $\mathbf{w}$ (Owen and Rabinovitch, 1983; Embrechts et al., 2002). Apply law invariance, translativeity, and positive homogeneity to $L_{\mathbf{w}} = -\mathbf{w}^\top \boldsymbol{\mu} - \sqrt{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}} Z$ (using $-Z \stackrel{d}{=} Z$ by symmetry of elliptical marginals) to obtain (10.1). For fixed mean, minimizing ϱ is minimizing $\sqrt{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}}$; CVaR at any $\alpha \in (0, 1)$ satisfies the hypotheses with $\varrho(Z) = \text{CVaR}_{\alpha}(Z) > 0$ (Rockafellar and Uryasev, 2000). The expected-utility statement is the characterization of Chamberlain (1983): under ellipticity with finite means, expected utility is a function of mean and variance alone, increasing in the former and decreasing in the latter for concave increasing u — and Chamberlain’s result is an exact characterization, so ellipticity is also essentially the largest class for which this transfer can hold for all utilities. $\square$

An allocator who replaces variance by CVaR escapes none of the conditions in this paper unless she simultaneously departs from ellipticity. The frameworks separate only through tails and asymmetry — skewness, asymmetric heavy tails, state-dependent dependence — which is where genuine mean–CVaR-versus-mean–variance trade-offs live.

10.2 Growth-optimal transfer

Proposition 10.3 (Kelly coincidence; $\mathbf{P}$, classical setting). *In the standard diffusion market $dS_{i,t}/S_{i,t} = \mu_i dt + (\boldsymbol{\sigma} dB_t)_i$ with $\boldsymbol{\Sigma} = \boldsymbol{\sigma} \boldsymbol{\sigma}^\top \succ 0$ and money-market rate r_f , the almost-sure long-run growth rate of a constant-proportion strategy holding fractions $\mathbf{w}$ in risky assets is $g(\mathbf{w}) = r_f + \mathbf{w}^\top \boldsymbol{\mu}_e - \frac{1}{2} \mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}$, maximized uniquely at the Kelly allocation $\mathbf{w}_K = \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_e$ (Kelly, 1956; Merton, 1971). Hence a fully invested heuristic $\mathbf{w}^\circ$ is proportional to the growth-optimal risky allocation — i.e. is the risky composition of every fractional-Kelly strategy — if and only*

if $\boldsymbol{\mu}_e \propto \boldsymbol{\Sigma} \mathbf{w}^\circ$: the same condition (3.2). Every optimality set in this paper is simultaneously a Kelly-coincidence set in the diffusion limit.

Proof. Itô's formula on $\ln W_t$ for the constant-proportion wealth dynamics gives $d \ln W_t = (r_f + \mathbf{w}^\top \boldsymbol{\mu}_e - \frac{1}{2} \mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}) dt + \mathbf{w}^\top \boldsymbol{\sigma} dB_t$; the time-average of the martingale part vanishes a.s., leaving $g(\mathbf{w})$, a strictly concave quadratic with maximizer $\boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}_e$. Proportionality $\mathbf{w}^\circ \propto \mathbf{w}_K$ is condition (3.2) verbatim. $\square$

11 Symmetry and Bayes Optimality: When the Heuristic Is the Optimizer

The exact conditions classify parameter configurations; this section classifies *beliefs*. The claim to be made precise is: each flat heuristic is the *exactly optimal* Bayes portfolio of an investor whose uncertainty is symmetric under the group corresponding to her information set. Ignorance, formalized as invariance, does not approximately rationalize the heuristics — it derives them.

Lemma 11.1 (Predictive reduction; **P**). *Let the investor hold a belief π over $(\boldsymbol{\mu}_e, \boldsymbol{\Sigma})$ and maximize the one-period objective $\mathbb{E}_\pi[\mathbf{w}^\top \mathbf{r}_e] - \frac{\lambda}{2} \text{Var}_\pi(\mathbf{w}^\top \mathbf{r}_e)$, where $\mathbf{r}_e$ is the predictive excess return (parameters integrated out, with $\mathbb{E}[\mathbf{r}_e \mid \boldsymbol{\mu}_e, \boldsymbol{\Sigma}] = \boldsymbol{\mu}_e$, $\text{Var}(\mathbf{r}_e \mid \boldsymbol{\mu}_e, \boldsymbol{\Sigma}) = \boldsymbol{\Sigma}$). Then the objective equals $\mathbf{w}^\top \bar{\boldsymbol{\mu}}_e - \frac{\lambda}{2} \mathbf{w}^\top \hat{\boldsymbol{\Sigma}} \mathbf{w}$ with*

$$\bar{\boldsymbol{\mu}}_e = \mathbb{E}_\pi[\boldsymbol{\mu}_e], \quad \hat{\boldsymbol{\Sigma}} = \mathbb{E}_\pi[\boldsymbol{\Sigma}] + \text{Cov}_\pi(\boldsymbol{\mu}_e) \succ 0,$$

and the Bayes portfolio is $\mathbf{w}^\star = \frac{1}{\lambda} \hat{\boldsymbol{\Sigma}}^{-1} \bar{\boldsymbol{\mu}}_e$ — the tangency portfolio of the predictive pair.

Proof. Law of total expectation and total variance: $\text{Var}_\pi(\mathbf{w}^\top \mathbf{r}_e) = \mathbb{E}_\pi[\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}] + \text{Var}_\pi(\mathbf{w}^\top \boldsymbol{\mu}_e) = \mathbf{w}^\top (\mathbb{E}_\pi[\boldsymbol{\Sigma}] + \text{Cov}_\pi(\boldsymbol{\mu}_e)) \mathbf{w}$. The maximizer is the unconstrained quadratic optimum. $\square$

Lemma 11.2 (Commutant of the symmetric group; **P**). *If $M \in \mathbb{R}^{N \times N}$ is symmetric and $PMP^\top = M$ for every permutation matrix P , then $M = uI + v\mathbf{1}\mathbf{1}^\top$ for scalars u, v ; if additionally $M \succ 0$ then $u > 0$ and $u + Nv > 0$, and $M^{-1}\mathbf{1} = (u + Nv)^{-1}\mathbf{1}$.*

Proof. Invariance under the transposition of i and j forces $M_{ii} = M_{jj}$ and $M_{ik} = M_{jk}$ for $k \neq i, j$; ranging over transpositions equalizes all diagonal entries and all off-diagonal entries. Positive definiteness gives the eigenvalue conditions (u with multiplicity $N-1$ on $\mathbf{1}^\perp$, $u + Nv$ on $\mathbf{1}$), and $M\mathbf{1} = (u + Nv)\mathbf{1}$ inverts as claimed. $\square$

Theorem 11.3 (Exchangeability $\Rightarrow$ equal weight; **P**). *Suppose the belief π is invariant under the full symmetric group acting on assets, $(\boldsymbol{\mu}_e, \boldsymbol{\Sigma}) \mapsto (P\boldsymbol{\mu}_e, P\boldsymbol{\Sigma}P^\top)$, and $\bar{\boldsymbol{\mu}}_e \neq 0$ with $\mathbf{1}^\top \hat{\boldsymbol{\Sigma}}^{-1} \bar{\boldsymbol{\mu}}_e > 0$. Then the Bayes portfolio of Lemma 11.1, normalized to full investment, is exactly the equal-weight portfolio — for every risk aversion λ and every exchangeable π .*

Proof. Invariance of π makes $\bar{\boldsymbol{\mu}}_e$ a fixed vector of every permutation, hence $\bar{\boldsymbol{\mu}}_e = m\mathbf{1}$; and makes both $\mathbb{E}_\pi[\boldsymbol{\Sigma}]$ and $\text{Cov}_\pi(\boldsymbol{\mu}_e)$ invariant matrices, hence by Lemma 11.2 $\hat{\boldsymbol{\Sigma}} = uI + v\mathbf{1}\mathbf{1}^\top$ with $\hat{\boldsymbol{\Sigma}}^{-1}\mathbf{1} \propto \mathbf{1}$. Thus $\mathbf{w}^\star \propto \hat{\boldsymbol{\Sigma}}^{-1} m\mathbf{1} \propto \mathbf{1}$, and the sign condition fixes $m > 0$. $\square$

Theorem 11.4 (Standardized exchangeability $\Rightarrow$ inverse volatility; **P**). *Suppose volatilities $\sigma_1, \dots, \sigma_N$ are known, and the belief about the standardized excess returns $\mathbf{z} = D^{-1}\mathbf{r}_e$ is exchangeable, with predictive mean $\bar{\mathbf{z}} = \bar{s}\mathbf{1}$, $\bar{s} > 0$, and predictive covariance $\hat{Z} \succ 0$. Then the Bayes portfolio, normalized to full investment, is exactly the inverse-volatility portfolio.*

Proof. Exchangeability of the predictive law of $\mathbf{z}$ forces $\bar{\mathbf{z}} \propto \mathbf{1}$ and, by Lemma 11.2, $\hat{Z} = uI + v\mathbf{1}\mathbf{1}^\top$. The predictive pair for $\mathbf{r}_e = D\mathbf{z}$ is $(\bar{s}D\mathbf{1}, D\hat{Z}D)$, so

$$\mathbf{w}^\star \propto (D\hat{Z}D)^{-1} \bar{s}D\mathbf{1} = \bar{s} D^{-1} \hat{Z}^{-1} \mathbf{1} \propto D^{-1} \mathbf{1},$$

the IV weights, using $\hat{Z}^{-1}\mathbf{1} = (u + Nv)^{-1}\mathbf{1}$. $\square$

Proposition 11.5 (Uninformative premia with conjugate uncertainty $\Rightarrow$ minimum variance; **P**). Suppose Σ is known, the belief about μ_e has mean $\bar{\mu}_e = m\mathbf{1}$, $m > 0$, and conjugate-form uncertainty $\text{Cov}_\pi(\mu_e) = \tau^2 \Sigma$, $\tau \geq 0$. Then the Bayes portfolio, normalized to full investment, is exactly the GMV portfolio, for every τ .

Proof. $\hat{\Sigma} = (1 + \tau^2)\Sigma$, so $\mathbf{w}^* \propto \Sigma^{-1}\mathbf{1}$. □

Remark 11.6 (MD, ERC, HRP at this rung). Maximum diversification is the degenerate case: a point-mass belief $\mu_e = S\sigma$ with known Σ gives $\mathbf{w}^* \propto \Sigma^{-1}\sigma$ immediately (Theorem 4.7 read as a Bayes statement). For ERC, HRP, and RA-HRP we know of no theorem of comparable cleanliness: ERC arises from the symmetric beliefs of Theorem 11.4 only on the constant-correlation slice where it collapses to IV (Proposition 6.5), HRP’s defining objects — the tree and the recursive split — have no invariance-group derivation we are aware of, and RA-HRP additionally requires a belief-level account of why noisy local cluster-Sharpe estimates should be trusted after aggregation but not globally inverted. A robust-control foundation for ERC under Sharpe-ratio ambiguity with a trusted correlation matrix, for HRP under ambiguity confined to inter-cluster blocks (the objects substitutions (i)–(ii) of Corollary 7.3 discard), and for RA-HRP under ambiguity-penalized local Sharpe scores are the natural conjectured forms; we state them as open problems, not results.

The pattern of Theorems 11.3–11.4 and Proposition 11.5 is the group-theoretic content of the information-set ladder: choose the subgroup of asset relabelings under which your knowledge is genuinely indistinguishable, impose invariance of beliefs under it, and the Bayes portfolio is forced into the corresponding heuristic. The heuristics are not approximations to optimization; they are optimization under symmetry.

12 Ambiguity, Robustness, and Estimation

The Bayes theorems condition on symmetric beliefs; robust and finite-sample results show that the same portfolios emerge from adversarial and frequentist formulations.

Proposition 12.1 ($1/N$ as the high-ambiguity limit; **C**). In distributionally robust portfolio selection where the return law is known only to lie in a ball (Kantorovich/Wasserstein-type) of radius ε around a nominal model and the investor minimizes the worst case of a convex risk functional over the ball, Pflug et al. (2012) prove that as $\varepsilon \rightarrow \infty$ the robust optimal portfolio converges to equal weight: uniform diversification is the exact limit of robust optimization under extreme ambiguity. Garlappi et al. (2007) obtain multi-prior portfolios interpolating between the plug-in optimum and conservative diversification as confidence shrinks.

Proposition 12.2 (Constraints as shrinkage; **C**). Jagannathan and Ma (2003) prove that the long-only (and upper-bounded) minimum-variance portfolio computed from an estimate $\hat{\Sigma}$ equals the unconstrained minimum-variance portfolio of a modified matrix in which constraint multipliers shrink the covariances of assets at the bounds: imposing the “wrong” constraints performs covariance shrinkage (cf. Ledoit and Wolf, 2004), which is why constrained GMV often beats unconstrained GMV out of sample.

Remark 12.3 (Estimation error and the empirical record; **E**). The sample counterpart is well documented. Errors in estimated means degrade mean–variance utility by roughly an order of magnitude more than comparable errors in variances at moderate risk aversion (Chopra and Ziemba, 1993); optimized weights are extremely sensitive to means (Best and Grauer, 1991; Michaud, 1989); and expected returns are the slowest quantity to learn from data (Merton, 1980). DeMiguel et al. (2009) show across datasets and fourteen optimizing models that none consistently beats $1/N$ out of sample in Sharpe ratio or certainty equivalent, with the estimation window

required for the plug-in tangency portfolio to overtake $1/N$ growing far beyond available histories as N increases; [Kan and Zhou \(2007\)](#) derive the optimal finite-sample combination of riskless asset, sample tangency, and sample GMV, formalizing the shrinkage the heuristics implement for free; and [Tu and Zhou \(2011\)](#) show that optimally combining $1/N$ with sophisticated rules can dominate both ingredients — the sample-level analogue of the HPO projection (8.1) over a two-member family. These are finite-sample statements, distinct from the population conditions of this paper: $1/N$ can dominate the estimated tangency portfolio while being far from the true one. Theorem 3.9 is the population-side bridge: because efficiency is flat across each coincidence set, the heuristic forfeits only second-order Sharpe when its condition nearly holds, while the plug-in optimizer pays first-order weight instability for second-order objective gains.

13 Constrained, Costly, and Implementable Optimality

The previous sections use the clean population benchmark: full investment, no short-sale constraints unless a rule imposes them internally, no turnover penalty, and no implementation wedge between a mathematical weight and a tradable portfolio. Actual mandates almost never look like that. They impose long-only constraints, concentration caps, sector limits, benchmark bands, turnover limits, and sometimes explicit transaction-cost models. The central message survives, but the exact condition changes from an equality to a *normal-cone* statement: the missing implied returns may be supplied by active constraints.

13.1 Normal-cone optimality under mandate constraints

Let $K \subset \mathbb{R}^N$ be a nonempty closed convex feasible set. In applications K usually includes the full-investment constraint, lower and upper bounds, and possibly linear exposure restrictions. Its normal cone at $\mathbf{w} \in K$ is

$$N_K(\mathbf{w}) = \{\mathbf{n} \in \mathbb{R}^N : \mathbf{n}^\top (\mathbf{v} - \mathbf{w}) \leq 0 \quad \forall \mathbf{v} \in K\}. \quad (13.1)$$

Theorem 13.1 (Constrained implied-return principle; **P**). *Fix $\gamma > 0$ and a closed convex mandate set K . A feasible portfolio $\mathbf{w}^\circ \in K$ solves the constrained mean–variance problem*

$$\max_{\mathbf{w} \in K} \left\{ \mathbf{w}^\top \boldsymbol{\mu}_e - \frac{\gamma}{2} \mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w} \right\} \quad (13.2)$$

if and only if there exists $\mathbf{n} \in N_K(\mathbf{w}^\circ)$ such that

$$\boldsymbol{\mu}_e = \gamma \boldsymbol{\Sigma} \mathbf{w}^\circ + \mathbf{n}. \quad (13.3)$$

Equivalently, the unconstrained implied-return residual

$$\mathbf{a}(\mathbf{w}^\circ) := \boldsymbol{\mu}_e - \gamma \boldsymbol{\Sigma} \mathbf{w}^\circ \quad (13.4)$$

must be entirely explainable by the shadow prices of the active mandate constraints.

Proof. The objective in (13.2) is strictly concave because $\boldsymbol{\Sigma} \succ 0$. Therefore first-order variational optimality is necessary and sufficient:

$$(\boldsymbol{\mu}_e - \gamma \boldsymbol{\Sigma} \mathbf{w}^\circ)^\top (\mathbf{v} - \mathbf{w}^\circ) \leq 0 \quad \forall \mathbf{v} \in K.$$

This is precisely $\boldsymbol{\mu}_e - \gamma \boldsymbol{\Sigma} \mathbf{w}^\circ \in N_K(\mathbf{w}^\circ)$. □

Corollary 13.2 (Box-constrained implied returns; **P**). *Let*

$$K = \{\mathbf{w} : \mathbf{1}^\top \mathbf{w} = 1, \ell_i \leq w_i \leq u_i \quad \forall i\}.$$

Then $\mathbf{w}^\circ$ solves (13.2) iff there exist $b \in \mathbb{R}$, lower-bound multipliers $\lambda_i \geq 0$, and upper-bound multipliers $\xi_i \geq 0$ satisfying complementarity

$$\lambda_i(w_i^\circ - \ell_i) = 0, \quad \xi_i(u_i - w_i^\circ) = 0,$$

and

$$\boldsymbol{\mu}_e = \gamma \boldsymbol{\Sigma} \mathbf{w}^\circ + b \mathbf{1} - \boldsymbol{\lambda} + \boldsymbol{\xi}. \quad (13.5)$$

Thus a zero weight does not require the asset to have zero optimal unconstrained demand; it only requires that its negative implied-return residual be absorbed by the lower-bound shadow price. Likewise, a capped name may have a positive residual absorbed by an upper-bound shadow price.

Proof. The normal cone of the equality-plus-box set is the span of $\mathbf{1}$ plus outward normals to active lower and upper faces. Using the sign convention in Theorem 13.1 gives (13.5). $\square$

Remark 13.3 (Why this matters for heuristic portfolios). Theorems 4.1–4.7 and 6.2 classify exact unconstrained coincidence. Theorem 13.1 gives the mandate version. A long-only equal-weight, inverse-volatility, ERC, MD, HRP, or GMV portfolio can be constrained-optimal even when it is not unconstrained-optimal, but only if the residual alphas are located on the active faces of the feasible set. This is the precise population analogue of the finite-sample result of Jagannathan and Ma (2003): constraints do not merely “regularize” the optimizer; they add shadow premia.

13.2 Transaction-cost wedges

Suppose the current portfolio is $\mathbf{w}_-$ and the investor pays a quadratic trading penalty with positive semidefinite matrix $\Lambda \succeq 0$ and intensity $\kappa \geq 0$:

$$\max_{\mathbf{w} \in K} \left\{ \mathbf{w}^\top \boldsymbol{\mu}_e - \frac{\gamma}{2} \mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w} - \frac{\kappa}{2} (\mathbf{w} - \mathbf{w}_-)^\top \Lambda (\mathbf{w} - \mathbf{w}_-) \right\}. \quad (13.6)$$

Quadratic costs are a local model of market impact and turnover aversion, and they are the static one-period analogue of dynamic transaction-cost models such as Gârleanu and Pedersen (2013).

Proposition 13.4 (Cost-adjusted implied returns; **P**). A feasible $\mathbf{w}^\circ \in K$ solves (13.6) iff there exists $\mathbf{n} \in N_K(\mathbf{w}^\circ)$ such that

$$\boldsymbol{\mu}_e = \gamma \boldsymbol{\Sigma} \mathbf{w}^\circ + \kappa \Lambda (\mathbf{w}^\circ - \mathbf{w}_-) + \mathbf{n}. \quad (13.7)$$

In the unconstrained full-investment case, this becomes

$$\boldsymbol{\mu}_e = \gamma \boldsymbol{\Sigma} \mathbf{w}^\circ + \kappa \Lambda (\mathbf{w}^\circ - \mathbf{w}_-) + b \mathbf{1}. \quad (13.8)$$

Proof. The gradient of the objective in (13.6) is $\boldsymbol{\mu}_e - \gamma \boldsymbol{\Sigma} \mathbf{w} - \kappa \Lambda (\mathbf{w} - \mathbf{w}_-)$. Apply the same variational inequality as in Theorem 13.1. $\square$

Remark 13.5 (No-trade interpretation). If $\mathbf{w}^\circ = \mathbf{w}_-$, the cost wedge vanishes; if the candidate rule demands a large rebalance, the wedge can dominate the statistical signal. Equation (13.7) says that a rule is implementably optimal only when expected excess returns compensate both risk and the marginal cost of moving from the current book. This is why the same population-optimal rule may be undesirable at daily frequency and reasonable at quarterly frequency.

Remark 13.6 (Linear costs, quadratic wedges, and RA-HRP turnover; **E**). The quadratic wedge in (13.7) is the differentiable local approximation to implementation frictions. A one-way linear cost $c \|\mathbf{w} - \mathbf{w}_-\|_1$ gives the same economic message with a nonsmooth subgradient: the implied-return residual must compensate risk plus an element of $c \partial \|\mathbf{w} - \mathbf{w}_-\|_1$, together with any

mandate normal cone. The RA-HRP evidence is consistent with both descriptions. The reported rolling implementation has monthly two-way turnover of approximately 10.0% for traditional HRP, 11.1% for RA-HRP, and 12.5% for Schur-RA-HRP; under one-way cost stress tests of 5–20 basis points, the RA-HRP net Sharpe remains close to the gross Sharpe (Noguer i Alonso, 2026). Thus the same structural feature that avoids the global inverse — local, bounded split revisions on a tree — also keeps trading frictions modest under linear and quadratic cost models.

13.3 Benchmark-relative and factor-neutral mandates

Many institutional portfolios are not judged in absolute variance but relative to a benchmark $\mathbf{w}_b$. Let active weights be $\mathbf{x} = \mathbf{w} - \mathbf{w}_b$, with $\mathbf{1}^\top \mathbf{x} = 0$, active covariance still Σ , and active expected excess return μ_a . A tracking-error mandate solves

$$\max_{\mathbf{x} \in K_a} \left\{ \mathbf{x}^\top \mu_a - \frac{\gamma}{2} \mathbf{x}^\top \Sigma \mathbf{x} \right\}. \quad (13.9)$$

The identical normal-cone result gives

$$\mu_a = \gamma \Sigma \mathbf{x}^\circ + \mathbf{n}_a, \quad \mathbf{n}_a \in N_{K_a}(\mathbf{x}^\circ). \quad (13.10)$$

Therefore a heuristic active overlay is optimal exactly when it prices active alphas through active covariances, after accounting for active constraints. In particular, the equal-weight theorem has a benchmark-relative form: equal active weights across an eligible sleeve require zero alphas against the active equal-weight sleeve, not against the absolute portfolio.

14 Estimation Geometry, High-Dimensional Risk, and Empirical Diagnostics

The paper has so far separated population truth from finite-sample practice. This section connects them. The exact optimality sets tell us what must be true; the estimation geometry tells us why, even when those conditions are false, a rule that ignores some inputs may dominate an optimizer that estimates them badly.

14.1 First-order sensitivity of tangency weights

Let

$$\mathcal{T}(\mu_e, \Sigma) = \frac{\Sigma^{-1} \mu_e}{\mathbf{1}^\top \Sigma^{-1} \mu_e}$$

be the normalized tangency map and write $c = \mathbf{1}^\top \Sigma^{-1} \mu_e > 0$, $\mathbf{w}_T = \mathcal{T}(\mu_e, \Sigma)$.

Theorem 14.1 (Differential of the tangency map; $\mathbf{P}$). *For perturbations $(\delta \mu, \delta \Sigma)$ small enough that $\Sigma + \delta \Sigma \succ 0$ and the denominator remains positive, the first differential of the tangency weights is*

$$\delta \mathbf{w}_T = \left(I - \mathbf{w}_T \mathbf{1}^\top \right) \left(\frac{1}{c} \Sigma^{-1} \delta \mu - \Sigma^{-1} \delta \Sigma \mathbf{w}_T \right) + o(\|\delta \mu\| + \|\delta \Sigma\|). \quad (14.1)$$

Consequently, mean errors enter with amplification $1/c$, while covariance errors enter through the already-chosen direction $\mathbf{w}_T$.

Proof. Let $\mathbf{x} = \Sigma^{-1} \mu_e$ and $c = \mathbf{1}^\top \mathbf{x}$, so $\mathbf{w}_T = \mathbf{x}/c$. The differential of $\mathbf{x}$ is

$$\delta \mathbf{x} = \Sigma^{-1} \delta \mu - \Sigma^{-1} \delta \Sigma \Sigma^{-1} \mu_e = \Sigma^{-1} \delta \mu - c \Sigma^{-1} \delta \Sigma \mathbf{w}_T.$$

Since $\delta c = \mathbf{1}^\top \delta \mathbf{x}$,

$$\delta \mathbf{w}_T = \frac{\delta \mathbf{x}}{c} - \frac{\mathbf{x} \delta c}{c^2} = \frac{1}{c} (I - \mathbf{w}_T \mathbf{1}^\top) \delta \mathbf{x},$$

which gives (14.1). □

Remark 14.2 (Why means dominate). Equation (14.1) is the infinitesimal version of the empirical findings of [Merton \(1980\)](#), [Best and Grauer \(1991\)](#), [Chopra and Ziemba \(1993\)](#), and [Michaud \(1989\)](#). Expected returns are small in annualized units, so $c = \mathbf{1}^\top \Sigma^{-1} \boldsymbol{\mu}_e$ is often small; the inverse factor $1/c$ turns modest premium errors into large allocation errors. A volatility-only rule may be population-suboptimal yet estimation-robust because it deletes the most unstable term in (14.1).

14.2 Projection onto a heuristic optimality set

The exact conditions can be turned into diagnostics. Fix Σ and a candidate $\mathbf{w}^\circ$. Its tangency-coincidence set is the ray $\{a\Sigma\mathbf{w}^\circ : a > 0\}$. Given an estimated premium vector $\hat{\boldsymbol{\mu}}_e$, ask how far it is from that ray.

Proposition 14.3 (Closest implied-return ray; **P**). Let $M \succ 0$ define a diagnostic norm $\|\mathbf{z}\|_M^2 = \mathbf{z}^\top M \mathbf{z}$ and set $\mathbf{q} = \Sigma \mathbf{w}^\circ$. The best positive-ray approximation to $\hat{\boldsymbol{\mu}}_e$ is

$$a^* = \max \left\{ 0, \frac{\mathbf{q}^\top M \hat{\boldsymbol{\mu}}_e}{\mathbf{q}^\top M \mathbf{q}} \right\}, \quad \hat{\boldsymbol{\mu}}_e^{\text{proj}} = a^* \mathbf{q}, \quad (14.2)$$

and the squared distance

$$d_M^2(\hat{\boldsymbol{\mu}}_e, \mathcal{O}_\Sigma(\mathbf{w}^\circ)) = \|\hat{\boldsymbol{\mu}}_e - a^* \Sigma \mathbf{w}^\circ\|_M^2 \quad (14.3)$$

quantifies the violation of the heuristic's exact optimality condition.

Proof. Minimize the one-dimensional quadratic $Q(a) = \|\hat{\boldsymbol{\mu}}_e - a\mathbf{q}\|_M^2$ over $a \geq 0$. The unconstrained minimizer is $(\mathbf{q}^\top M \hat{\boldsymbol{\mu}}_e)/(\mathbf{q}^\top M \mathbf{q})$; projection onto $[0, \infty)$ gives (14.2). $\square$

Remark 14.4 (A useful choice of metric). Taking $M = \hat{\Omega}_\mu^{-1}$, where $\hat{\Omega}_\mu$ is the estimated covariance matrix of premium estimates, turns (14.3) into a Wald-type statistic. Taking $M = \Sigma^{-1}$ measures distance in the same dual metric that appears in $S_{\max}^2 = \boldsymbol{\mu}_e^\top \Sigma^{-1} \boldsymbol{\mu}_e$. Either way, the diagnostic decomposes the failure of a heuristic into a scalar scale estimate a^* plus an orthogonal residual premium vector.

14.3 GRS and spanning tests as optimality tests

[Remark 3.6](#) states the principle. The usable formula is as follows. For T excess-return observations, regress each asset on the candidate heuristic excess return $r_{e,t}^\circ$:

$$r_{e,i,t} = \alpha_i + \beta_i r_{e,t}^\circ + \epsilon_{i,t}. \quad (14.4)$$

Let $\hat{\boldsymbol{\alpha}}$ be the vector of intercepts, $\hat{\Sigma}_\epsilon$ the residual covariance matrix, $\bar{r}^\circ$ the sample mean of the candidate return, and $\hat{\sigma}_\epsilon^2$ its sample variance. With one candidate factor, the Gibbons–Ross–Shanken statistic is

$$\text{GRS} = \frac{T - N - 1}{N} \frac{\hat{\boldsymbol{\alpha}}^\top \hat{\Sigma}_\epsilon^{-1} \hat{\boldsymbol{\alpha}}}{1 + (\bar{r}^\circ)^2 / \hat{\sigma}_\epsilon^2} \sim F_{N, T-N-1} \quad (14.5)$$

under the Gaussian exact-efficiency null ([Gibbons et al., 1989](#)). Small-sample refinements and related asymptotics are discussed by [Jobson and Korkie \(1980\)](#) and [Memmel \(2003\)](#). Mean-variance spanning tests such as [Huberman and Kandel \(1987\)](#) provide the multi-portfolio analogue: a set of heuristic sleeves spans the efficient frontier iff adding the remaining assets does not improve the frontier.

Remark 14.5 (Interpreting rejection). Rejection of (14.5) does not mean the heuristic is useless; it means the exact population condition is statistically inconsistent with the sample. [Theorem 3.9](#) then gives the next question: is the rejection economically large? A large p -value is evidence of statistical non-rejection; a small angular loss $1 - \eta^2$ is evidence of economic adequacy. They answer different questions.

14.4 High-dimensional covariance and the inversion tax

Even a rule with a clean population foundation can fail in finite samples if it requires inverting a noisy covariance matrix. This is the high-dimensional counterpart of mean-estimation fragility.

Proposition 14.6 (Sample GMV risk inflation; **C**). *Let $\hat{\Sigma}$ be the sample covariance matrix from T independent Gaussian observations on N assets with true covariance $\Sigma \succ 0$, and let $\hat{\mathbf{w}}_{\text{GMV}} = \hat{\Sigma}^{-1}\mathbf{1}/(\mathbf{1}^\top \hat{\Sigma}^{-1}\mathbf{1})$ whenever $\hat{\Sigma}$ is nonsingular. In the high-dimensional regime $N/T \rightarrow q \in (0, 1)$, random-matrix results imply the out-of-sample variance inflation*

$$\frac{\hat{\mathbf{w}}_{\text{GMV}}^\top \Sigma \hat{\mathbf{w}}_{\text{GMV}}}{\mathbf{w}_{\text{GMV}}^\top \Sigma \mathbf{w}_{\text{GMV}}} \rightarrow \frac{1}{1-q} \quad (14.6)$$

under standard regularity conditions. This is the inversion tax: as the cross-section approaches the sample length, the plug-in GMV portfolio becomes mechanically overfit. See [El Karoui \(2010\)](#) and the shrinkage literature of [Ledoit and Wolf \(2004\)](#).

Remark 14.7 (What HRP and shrinkage are buying). *Equation (14.6) clarifies the empirical appeal of HRP, RA-HRP, inverse volatility, ERC, and covariance shrinkage. They give up exact global inversion of Σ in exchange for reduced estimation variance. Fixed-tree RA-HRP belongs to this low-inversion family: its Jacobian is a composition of local rational split maps whose denominators are cluster variances, score sums, and floors, rather than the smallest eigenvalue of the full covariance matrix. Schur-RA-HRP restores some cross-cluster information through local block inversions, but it still avoids a single global Σ^{-1} . Cotton’s Schur-complement view of HRP and minimum variance, and subsequent hierarchical minimum-variance constructions, make this trade-off explicit by partially restoring the cross-block covariance information that HRP discards ([Cotton, 2024](#); [Mograby, 2025](#)). Cotton’s 2024 paper introduces Schur Complementary Allocation as a bridge between HRP and minimum-variance portfolios, while Mograby’s 2025 work develops a recursive Schur-complement framework for hierarchical minimum variance; both directly support the interpretation of HRP as a controlled deletion or restoration of inverse-covariance information.*

14.5 A reproducible empirical protocol

A paper claiming that a heuristic is “optimal” should report three layers, not one.

1. **Population-condition diagnostic.** Estimate the rule-specific restriction: EW zero alphas against EW; IV Sharpe ratios versus correlation row sums; ERC performance parity; MD equal Sharpe ratios; GMV equal premia; HRP Schur-substitution residuals.
2. **Statistical test.** Report the GRS statistic (14.5), a spanning test when several sleeves are compared, and robust/HAC versions when returns are serially dependent or heteroskedastic.
3. **Economic loss.** Report realized Sharpe efficiency, angular loss $1 - \eta^2$, turnover, transaction-cost-adjusted Sharpe, drawdown, and CVaR. Statistical rejection with economically negligible angular loss is not a failure of the heuristic; non-rejection with high turnover may still be implementation failure.
4. **Stability.** Use rolling windows, block bootstrap, and perturbation experiments. Report how often the ranking of rules changes under resampling. This directly measures the estimation terms isolated in Theorem 14.1.
5. **Dimensionality.** Always report $q = N/T$ for the covariance window. Without q , a comparison between plug-in optimization and heuristics is uninterpretable, because (14.6) changes the expected cost of inversion.

14.6 Computational profile of the allocation ladder

The exact optimality results should be read together with the numerical burden of the rule. In dense implementations, the usual computational profile is summarized in Table 3. The point is not that lower complexity is automatically better; it is that every avoided inversion, local score, or tree constraint deletes a specific source of estimation variance and replaces it by a specific structural assumption.

Rule	Typical dense cost	Main numerical bottleneck	Interpretation of the cost
EW	$O(N)$	none	Maximum symmetry; no sampling burden beyond defining the universe.
IV / IVar	$O(N)$ after vol estimates	volatility estimation	Uses diagonal risk only; deletes correlation and premium information.
GMV	$O(N^3)$ after covariance estimation	global covariance inversion and conditioning	Buys the full covariance signal at the price of the high-dimensional inversion tax.
ERC	iterative, often $O(KN^2)$ for K solver steps	nonlinear risk-budget solve	Uses covariance without premia; solver stability depends on conditioning and constraints.
HRP	clustering plus recursive local variances	dendrogram stability	Replaces global inversion by tree topology and local variance aggregation.
RA-HRP	HRP cost plus local mean/Sharpe scores	local premium noise and floor activation	Spends a controlled amount of mean-estimation risk locally rather than globally.
Schur-RA-HRP	RA-HRP plus sibling-block inversions	conditioning of local Schur blocks	Restores cross-cluster hedging information without paying the full global inverse.
RLPO residual policy	training-dependent	off-policy error, reward design, turnover and regime shift	Learns dynamic deviations only when continuation value pays for the static HPO defect.

Table 3: Computational profile of the HPO–RLPO ladder. Costs are indicative for dense implementations and should be reported with the covariance-window ratio $q = N/T$.

A reproducible empirical implementation should therefore report not only realized returns but also the cost of producing the weights: the length of the estimation window, $q = N/T$, covariance-conditioning diagnostics, tree-stability diagnostics, floor-activation frequencies for RA-HRP, local Schur-block conditioning for Schur-RA-HRP, and post-cost turnover. Without these quantities, an empirical comparison confounds portfolio skill with numerical fragility.

15 Non-Elliptical Tails, Downside Risk, and Where the Conditions Break

The elliptical transfer theorem is intentionally sharp. It says that replacing variance by CVaR does not change the optimality conditions when all linear portfolios share the same standardized marginal shape. The converse message is equally important: departures from ellipticity are exactly where different objectives begin to disagree.

Definition 15.1 (Tail-shape residual). *For a portfolio $\mathbf{w}$, define the standardized excess-return residual*

$$Z_{\mathbf{w}} = \frac{\mathbf{w}^\top (\mathbf{r} - \boldsymbol{\mu})}{\sqrt{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}}}. \quad (15.1)$$

The tail-shape residual between two portfolios $\mathbf{w}$ and $\mathbf{v}$ at level α is

$$\Delta_\alpha(\mathbf{w}, \mathbf{v}) = \text{CVaR}_\alpha(-Z_{\mathbf{w}}) - \text{CVaR}_\alpha(-Z_{\mathbf{v}}). \quad (15.2)$$

Under ellipticity, $\Delta_\alpha(\mathbf{w}, \mathbf{v}) = 0$ for all $\mathbf{w}, \mathbf{v}, \alpha$.

Proposition 15.2 (First-order downside wedge; $\mathbf{P}$). *Assume a law-invariant homogeneous downside risk measure admits the representation*

$$\varrho(L_{\mathbf{w}}) = -\mathbf{w}^\top \boldsymbol{\mu} + k_{\mathbf{w}} \sqrt{\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w}}, \quad (15.3)$$

where $k_{\mathbf{w}}$ is differentiable in a neighborhood of $\mathbf{w}^\circ$. Then the first-order condition for maximizing mean per unit downside risk contains the additional tail-shape term $\nabla k_{\mathbf{w}}|_{\mathbf{w}=\mathbf{w}^\circ}$. In particular, the mean–variance implied-return condition $\boldsymbol{\mu}_e \propto \boldsymbol{\Sigma} \mathbf{w}^\circ$ remains sufficient only when $k_{\mathbf{w}}$ is locally constant along feasible directions.

Proof. Differentiate the generalized denominator $k_{\mathbf{w}}(\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w})^{1/2}$. In addition to the variance-gradient term $k_{\mathbf{w}} \boldsymbol{\Sigma} \mathbf{w} / (\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w})^{1/2}$, one obtains $(\mathbf{w}^\top \boldsymbol{\Sigma} \mathbf{w})^{1/2} \nabla k_{\mathbf{w}}$. The latter vanishes exactly when standardized tail shape is locally invariant. $\square$

Remark 15.3 (Practical implication). *This identifies the boundary of the paper’s transfer claims. If assets have asymmetric crash exposure, volatility targeting, option-like payoffs, or regime-dependent correlations, then a heuristic can be Sharpe-near-optimal and CVaR-poor. The missing state variable is not variance but tail-shape heterogeneity. Empirically, this means the protocol in Section 14.5 should report both angular Sharpe loss and tail-shape residuals.*

16 Synthesis: The Optimality Map

Table 6 collects the exact population conditions; Table 7 collects the belief-level foundations. Together they give the information-set ladder theorem-level content at every rung where such content currently exists, and mark the missing rungs as open.

Table 4 separates the same rules by inputs, inversion profile, estimation vulnerability, and topological coincidence boundary. Table 5 then places those static rules inside the dynamic RLPO hierarchy developed in Section 9. Together with the HPO defect and bias–variance identities of Section 8, this is the practical map: it explains why a rule can be theoretically suboptimal and still empirically robust when the coordinates required by the optimizer are too noisy, and why an RL policy should deviate from that rule only when continuation value pays for the static defect.

Allocation rule	Inputs consumed	Inversion profile	Primary estimation vulnerability	Topological coincidence boundary
Equal weight	None	None	Symmetry misspecification, $\mu_e \propto \Sigma \mathbf{1}$ not estimation noise	
Inverse volatility	Marginal volatilities σ_i	None	Volatility estimates and correlation-row-sum mismatch	$S_i \propto \sum_j \rho_{ij}$
Global min variance	Full Σ	Global Σ^{-1}	High-dimensional inversion tax $1/(1-q)$ and covariance conditioning	$\mu_e \propto \mathbf{1}$ for tangency coincidence
Traditional HRP	Tree plus cluster IVP variances	No global inverse	Dendrogram instability and Schur-substitution residuals	Diagonal Σ under any tree; uncorrelated exchangeable blocks under balanced tree-aligned bisection; plus equal premia for tangency
RA-HRP	Tree plus cluster means and IVP variances	No global inverse	Persistence and noise of local cluster-Sharpe scores	Nonlinear fixed point $\mu_e = a\Sigma\Phi_{T,\varepsilon}^{\text{RA}}(\mu_e, \Sigma)$; nodewise equality of RA splits and Schur-tangency splits
Schur-RA-HRP	Tree, cluster pre-mia, and Schur conditional risks	Local block versions	Cross-cluster conditional-covariance estimation and block conditioning	Collapses to RA-HRP when sibling hedge terms vanish; exact only when conditional Schur scores match Schur-tangency capital masses node by node

Table 4: Structural comparison of heuristic, hierarchical, and return-adjusted allocation rules. The table distinguishes the economic input set from the numerical inversion profile and from the exact population boundary required for tangency coincidence.

Layer	Mathematical object	Role in the research program
HPO / exact optimality	static map $w = \Phi(\hat{\mu}, \hat{\Sigma}, \hat{T})$ and defect $\mathfrak{d}$	identifies when a heuristic is locally optimal and how much myopic Sharpe efficiency it loses
RA-HRP / Schur-RA-HRP	tree-factorized map with branch probabilities $\{\beta_v\}$	supplies interpretable hierarchical policy priors and node-level KL attribution
RLPO	Markov policy $\pi_\theta(w \mid s)$ with Bellman value	learns dynamic deviations from HPO under costs, regimes, and continuation value

Table 5: Division of labor between static HPO, hierarchical RA-HRP priors, and dynamic RLPO.

Rule	Uses	Exact tangency condition (this paper)	Informational regime
Equal weight	nothing	zero alpha vs. EW: $\mu_e \propto \Sigma \mathbf{1}$ [P, Thm. 4.1]	total ambiguity
Inverse volatility	σ_i	$S_i \propto \sum_j \rho_{ij}$ [P, Thm. 4.3]	vols known, correlations exchangeable
ERC / risk parity	Σ (no μ)	performance parity: $w_i \mu_{e,i}$ equal [P, Thm. 6.2]; sufficient: equal S_i , constant ρ [P, Cor. 6.6]	Σ trusted, premia exchangeable per unit risk budget
Max. diversification	σ, Σ	equal Sharpe ratios: $\mu_e \propto \sigma$ [P, Thm. 4.7]	risk priced uniformly per unit vol
HRP	tree + cluster variances	GMV-coincidence: diagonal Σ , any tree [P, Thm. 7.5]; uncorrelated exchangeable blocks [P, Prop. 7.6]; then $\mu_e \propto \mathbf{1}$ [P, Cor. 7.8]; gap = three Schur substitutions [P, Cor. 7.3]	ordinal correlation structure trusted, Σ^{-1} not
RA-HRP	tree + cluster Sharpe scores	nonlinear fixed point $\mu_e = a \Sigma \Phi_{\mathcal{T}, \varepsilon}^{\text{RA}}(\mu_e, \Sigma)$ [P, Thm. 7.15]; nodewise Schur-tangency split matching; HRP-RA pathwise/KL attribution [P, Thm. 7.22]	tree trusted; premia trusted only through local IVP aggregation/flooring; no global inverse
Schur-RA-HRP	tree + cluster Sharpe scores + Schur complements	conditional-risk score from Def. 7.17; Schur score wedge [P, Prop. 7.18]; locally exact only when conditional scores reproduce Schur-tangency masses	tree trusted; cross-cluster covariance trusted locally, but global inverse still avoided
Minimum variance	Σ	$\mu_e \propto \mathbf{1}$ [P, Thm. 4.5]	risk forecastable, premia not
Tangency	μ_e, Σ	— (benchmark)	full knowledge

Table 6: The optimality map. All conditions are also Kelly-coincidence conditions in the diffusion limit [P, Prop. 10.3], transfer to mean-CVaR and expected utility under ellipticity [P, Thm. 10.2], and are GRS-testable as zero-alpha restrictions [C, Rem. 3.6]. One-factor translations are in Prop. 5.3.

Rule	Belief-level foundation	Status
Equal weight	Bayes rule under any fully exchangeable belief	proved [P, Thm. 11.3]; also robust high-ambiguity limit [C, Prop. 12.1]
Inverse volatility	Bayes rule under exchangeable beliefs on standardized returns, σ_i known	proved [P, Thm. 11.4]
Minimum variance	Bayes rule under conjugate mean uncertainty with prior mean $\propto \mathbf{1}$, Σ known	proved [P, Prop. 11.5]; constraints-as-shrinkage in sample [C, Prop. 12.2]
Max. diversification	degenerate belief $\mu_e \propto \sigma$, Σ known	immediate [P, Rem. 11.6]
ERC / risk parity	robust foundation under Sharpe ambiguity with trusted correlations	<i>open</i> [Rem. 11.6]
HRP	robust foundation under ambiguity confined to inter-cluster blocks	<i>open</i> [Rem. 11.6]
RA-HRP	local cluster-Sharpe shrinkage on a fixed tree; exact symmetry/robust foundation for noisy premia remains unresolved	fixed-tree theory and empirical proof of concept in Noguer i Alonso (2026) ; belief-level foundation <i>open</i> [Rem. 11.6]
Schur-RA-HRP	local conditional-risk version of RA-HRP; robust foundation under uncertainty about cross-cluster hedge terms remains unresolved	Schur extension in Def. 7.17; belief-level foundation <i>open</i> [Rem. 11.6]

Table 7: Belief-level foundations: symmetry-derived Bayes optimality and robust limits.

17 Limitations and Open Problems

The paper closes several algebraic gaps but leaves important research problems open. These limitations are useful because they identify the exact objects that a referee, implementer, or subsequent paper should attack.

Random dendrogram asymptotics. The fixed-tree theory is sharp, but rolling HRP and RA-HRP estimate the tree. A complete asymptotic theory must combine a continuous delta method on the score maps with a discrete stability theorem for merge sets, cophenetic matrices, or tree-edit representations. The correct theorem should condition on a population merge-separation event, prove topology preservation with high probability, and then control off-event portfolio jumps through the pathwise product formula of Theorem 7.22.

Robust-control foundations for ERC, HRP, and RA-HRP. The Bayes foundations for EW, IV, and GMV are clean because the relevant symmetry groups are clean. Comparable foundations for ERC, HRP, RA-HRP, and Schur-RA-HRP require ambiguity sets that act on risk budgets, inter-cluster covariance blocks, and local Sharpe scores. Corollary 7.3, Theorem 7.15, and Theorem 8.10 identify the exact algebraic quantities such ambiguity sets must preserve or penalize.

Non-elliptical downside risk. The mean-CVaR transfer is exact under ellipticity and fails precisely through standardized tail-shape heterogeneity. A full downside-risk extension would replace the scalar angular defect by a pair consisting of Sharpe-angle loss and tail-shape residual $\Delta_\alpha(w, v)$, then study when local hierarchical scores can control both simultaneously.

Dynamic market impact and partial observability. The RLPO bridge uses a discounted Markov formulation with explicit rewards and costs. Realistic execution adds transient impact, hidden liquidity, delayed fills, and partial observability. The natural extension is a belief-state RLPO formulation in which the HPO prior is computed from filtered estimates and the deviation principle is applied to belief-state continuation values.

Uniform empirical validation. The theory supplies falsification diagnostics but does not report new numerics. A definitive empirical companion should run the protocol of Section 14.5 across asset classes, universes, rebalance frequencies, covariance estimators, cost models, and tree-linkage rules; report split-sample GRS tests and angular losses; and attribute RA-HRP or RLPO gains node by node through the KL and alpha decompositions.

18 Conclusion

Eight structural facts organize everything above.

First, HPO is a mathematical object in its own right. A heuristic portfolio rule is an information-restricted map from a reduced statistic to the simplex; its population error is the implied-return projection defect $\mathfrak{D}_\Phi$; and Theorem 8.4 proves that this defect is exactly squared Sharpe inefficiency. The score-tree formalism then shows that HRP, RA-HRP, Schur-RA-HRP, and their interpolations are products of local binary kernels, while Theorem 8.10 identifies the exact Schur-tangency mass each node is trying to approximate.

Second, by the implied-return principle, “is this heuristic optimal” is never yes-or-no; it is a question about a closed-form set of parameter configurations, and adopting a rule *is* asserting its set. Equal weight asserts zero alphas against itself; inverse volatility asserts Sharpe ratios aligned with correlation row sums; maximum diversification asserts equal Sharpe ratios with no view on correlations; minimum variance asserts equal premia; risk parity asserts that its

risk budget is simultaneously a performance budget; HRP asserts that the three pieces of covariance information it discards at every split — the Schur correction, the budget tilt, and within-cluster correlation — are genuinely zero; RA-HRP asserts the nonlinear fixed point in which its floored cluster-Sharpe splits match the exact Schur-tangency splits; Schur-RA-HRP asserts the analogous condition after replacing standalone cluster risk by conditional Schur risk. The unit-free HRP–RA-HRP interpolation shows how to move continuously from the risk-only assertion to the return-adjusted assertion without mixing incompatible score units. Every one of these assertions is GRS-testable, and under one-factor structure every one of them is readable off betas and idiosyncratic variances.

Third, the geometry of suboptimality is angular and flat. Sharpe efficiency is exactly the cosine of the Σ -angle to tangency; the optimality sets are Lebesgue-null, so exact optimality is a probability-zero event under any diffuse belief; and yet the efficiency surface is stationary on each set, so order- ε violations cost order- ε^2 efficiency with an explicit constant. “Never optimal, rarely costly” is two theorems, and the same flatness, read from the other side, is the first-order weight instability that makes plug-in optimization fragile.

Fourth, none of this is about variance per se. Under elliptical returns the identical conditions govern mean–CVaR at every confidence level and expected utility for every risk-averse investor, and in the diffusion limit they are the conditions for fractional-Kelly growth optimality; only tail asymmetry separates the objectives.

Fifth, the deepest sense in which the heuristics are right is not that their parameter conditions hold — generically they do not — but that each rule is *exactly* the optimal policy under the symmetry group of its information set. Full exchangeability of beliefs forces equal weight; exchangeability of standardized returns with known volatilities forces inverse volatility; uninformative premia with conjugate uncertainty force minimum variance; and the equal-weight limit also emerges from adversarial robustness as ambiguity diverges. The heuristics are not shortcuts around optimization; they are what optimization itself returns when the inputs it would need are replaced by honest symmetry. The missing rungs — a robust-control derivation of ERC under Sharpe ambiguity with trusted correlations, of HRP under ambiguity confined to inter-cluster structure, and of RA-HRP under ambiguity-penalized local Sharpe scores — are stated here as open problems; Corollary 7.3 and Theorem 7.15 suggest the exact ambiguity sets in which to look.

Sixth, RLPO is the dynamic continuation of HPO, not a replacement for it. Proposition 9.3 shows that every HPO rule is a deterministic stationary policy inside the RLPO Markov decision problem, and the $\gamma = 0$ no-friction Bellman problem collapses to static HPO. Proposition 9.5 gives the economic test for dynamic improvement: a learned policy should deviate from the HPO baseline only when its continuation-value gain exceeds the instantaneous reward loss, incremental turnover cost, and added implied-return defect. This makes the HPO defect a principled reward-shaping term rather than an arbitrary regularizer. Proposition 9.7 completes the accounting: the total value gap of acting myopically is exactly the discounted occupancy average of the HPO advantage, so statewise ε -adequacy of the heuristic implies a global $\varepsilon/(1 - \gamma)$ guarantee (Corollary 9.8) — the dynamic analogue of “never optimal, rarely costly.”

Seventh, RA-HRP supplies the natural actor architecture for RLPO. Its weights factorize through node-level branch probabilities, so a learned policy can be centered on the RA-HRP or Schur-RA-HRP split via residual logits. Proposition 9.9 proves that this hierarchical actor remains long-only and fully invested by construction, recovers RA-HRP when residuals vanish, and decomposes the global KL distance from the baseline into node-level binary divergences. This gives RLPO a transparent attribution system: root-level deviations are regime allocation decisions, and deeper-node deviations are local cluster or security tilts. The homotopy calculus of Section 7.5 supplies the matching learning signal: at the prior, the myopic policy gradient with respect to each residual logit is the nodewise alpha the baseline leaves unexplained, damped by the logistic variance of the split (Corollary 9.10).

Eighth, constraints, costs, topology, and tails are the implementation layer. A constrained heuristic is optimal only when its unexplained alpha vector lies in the normal cone of the mandate, and a costly heuristic is optimal only when premia compensate both risk and marginal turnover. For HRP and RA-HRP, the additional object is tree stability: merge preservation, cophenetic distance, and pathwise split perturbations are the bridge between fixed-tree theorems and traded rolling-window strategies. In RLPO these diagnostics become state variables and topology regularizers. Outside ellipticity, tail-shape residuals are the further object that separates Sharpe optimality from downside-risk optimality.

The result is a precise division of labor. Static optimization is the right tool when the information set includes the coordinates it consumes at the accuracy it requires. HPO is the right tool when those coordinates are noisy, unavailable, or economically untrustworthy. RLPO is the right tool when the state is dynamic and deviations from HPO can be paid for by continuation value. The theorems above say, line by line, which beliefs separate the three situations — and what each situation costs.

A Technical Appendix: Identities Used Repeatedly

This appendix records the three algebraic identities used throughout the paper. They are included to make the paper self-contained and to separate reusable linear-algebra facts from portfolio-specific interpretation.

A.1 Block inverse and Schur complement

If

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \succ 0,$$

with A and D nonsingular, then the Schur complements $S_A = A - BD^{-1}C$ and $S_D = D - CA^{-1}B$ are positive definite and

$$M^{-1} = \begin{pmatrix} S_A^{-1} & -S_A^{-1}BD^{-1} \\ -D^{-1}CS_A^{-1} & D^{-1} + D^{-1}CS_A^{-1}BD^{-1} \end{pmatrix}$$

after eliminating the D block, with the symmetric expression obtained after eliminating the A block. The HRP and Schur-RA-HRP identities in Sections 7 and 8 are portfolio interpretations of this formula.

A.2 Ray projection

For any inner product induced by $M \succ 0$, the closest point on the positive ray $\{aq : a \geq 0\}$ to a vector x is

$$a^*q, \quad a^* = \max \left\{ 0, \frac{q^\top Mx}{q^\top Mq} \right\}.$$

Choosing $x = \mu_e$, $q = \Sigma w$, and $M = \Sigma^{-1}$ gives the implied-return defect and its equality with squared Sharpe inefficiency.

A.3 Tree products and binary KL

If two hierarchical portfolios p and q are generated on the same binary tree by local splits p_v and q_v , then every leaf probability is a product of branch probabilities and

$$D_{\text{KL}}(p||q) = \sum_{v \in V_{\text{int}}} P_v \left[p_v \log \frac{p_v}{q_v} + (1 - p_v) \log \frac{1 - p_v}{1 - q_v} \right],$$

where P_v is the capital mass reaching node v under p . This is the identity behind the HRP–RA–HRP distortion theorem and the hierarchical RLPO regularizer.

References

- Richard Bellman. *Dynamic Programming*. Princeton University Press, 1957.
- Dimitri P. Bertsekas. *Dynamic Programming and Optimal Control*. Athena Scientific, 4 edition, 2017.
- Michael J. Best and Robert R. Grauer. On the sensitivity of mean-variance-efficient portfolios to changes in asset means: Some analytical and computational results. *The Review of Financial Studies*, 4(2):315–342, 1991.
- Fischer Black and Robert Litterman. Global portfolio optimization. *Financial Analysts Journal*, 48(5):28–43, 1992.
- Gary Chamberlain. A characterization of the distributions that imply mean-variance utility functions. *Journal of Economic Theory*, 29(1):185–201, 1983.
- Vijay K. Chopra and William T. Ziemba. The effect of errors in means, variances, and covariances on optimal portfolio choice. *Journal of Portfolio Management*, 19(2):6–11, 1993.
- Yves Choueifaty and Yves Coignard. Toward maximum diversification. *Journal of Portfolio Management*, 35(1):40–51, 2008.
- Roger Clarke, Harindra de Silva, and Steven Thorley. Minimum-variance portfolio composition. *Journal of Portfolio Management*, 37(2):31–45, 2011.
- Peter Cotton. Schur complementary allocation: A unification of hierarchical risk parity and minimum variance portfolios, 2024.
- Victor DeMiguel, Lorenzo Garlappi, and Raman Uppal. Optimal versus naive diversification: How inefficient is the 1/n portfolio strategy? *The Review of Financial Studies*, 22(5):1915–1953, 2009.
- Noureddine El Karoui. High-dimensionality effects in the markowitz problem and other quadratic programs with linear constraints: Risk underestimation. *The Annals of Statistics*, 38(6):3487–3566, 2010.
- Paul Embrechts, Alexander McNeil, and Daniel Straumann. Correlation and dependence in risk management: Properties and pitfalls. *Risk Management: Value at Risk and Beyond*, pages 176–223, 2002.
- Lorenzo Garlappi, Raman Uppal, and Tan Wang. Portfolio selection with parameter and model uncertainty: A multi-prior approach. *The Review of Financial Studies*, 20(1):41–81, 2007.
- Nicolae Gârleanu and Lasse Heje Pedersen. Dynamic trading with predictable returns and transaction costs. *The Journal of Finance*, 68(6):2309–2340, 2013.
- Michael R. Gibbons, Stephen A. Ross, and Jay Shanken. A test of the efficiency of a given portfolio. *Econometrica*, 57(5):1121–1152, 1989.
- Gur Huberman and Shmuel Kandel. Mean-variance spanning. *The Journal of Finance*, 42(4):873–888, 1987.

- Ravi Jagannathan and Tongshu Ma. Risk reduction in large portfolios: Why imposing the wrong constraints helps. *The Journal of Finance*, 58(4):1651–1683, 2003.
- J. D. Jobson and Bob Korkie. Estimation for markowitz efficient portfolios. *Journal of the American Statistical Association*, 75(371):544–554, 1980.
- Sham Kakade and John Langford. Approximately optimal approximate reinforcement learning. In *Proceedings of the Nineteenth International Conference on Machine Learning (ICML)*, pages 267–274. Morgan Kaufmann, 2002.
- Raymond Kan and Guofu Zhou. Optimal portfolio choice with parameter uncertainty. *Journal of Financial and Quantitative Analysis*, 42(3):621–656, 2007.
- John L. Kelly. A new interpretation of information rate. *The Bell System Technical Journal*, 35(4):917–926, 1956.
- Olivier Ledoit and Michael Wolf. Honey, i shrunk the sample covariance matrix. *Journal of Portfolio Management*, 30(4):110–119, 2004.
- Marcos López de Prado. Building diversified portfolios that outperform out of sample. *Journal of Portfolio Management*, 42(4):59–69, 2016.
- Sébastien Maillard, Thierry Roncalli, and Jérôme Teïletche. The properties of equally weighted risk contribution portfolios. *Journal of Portfolio Management*, 36(4):60–70, 2010.
- Harry Markowitz. Portfolio selection. *The Journal of Finance*, 7(1):77–91, 1952.
- Christoph Memmel. Performance hypothesis testing with the sharpe ratio. *Finance Letters*, 1(1):21–23, 2003.
- Robert C. Merton. Optimum consumption and portfolio rules in a continuous-time model. *Journal of Economic Theory*, 3(4):373–413, 1971.
- Robert C. Merton. On estimating the expected return on the market: An exploratory investigation. *Journal of Financial Economics*, 8(4):323–361, 1980.
- Richard O. Michaud. The markowitz optimization enigma: Is optimized optimal? *Financial Analysts Journal*, 45(1):31–42, 1989.
- G. Mograby. Hierarchical minimum variance portfolios: A theoretical and recursive schur complement framework, 2025.
- Miquel Noguer i Alonso. Return-adjusted hierarchical risk parity and schur portfolios: Mathematical foundations and empirical evidence, March 2026. Working paper.
- Joel Owen and Ramon Rabinovitch. On the class of elliptical distributions and their applications to the theory of portfolio choice. *The Journal of Finance*, 38(3):745–752, 1983.
- Georg Ch. Pflug, Alois Pichler, and David Wozabal. Uniform investment strategy as a limit of ambiguous portfolio optimization. *Journal of Banking and Finance*, 36(2):410–417, 2012.
- Edward E. Qian. On the financial interpretation of risk contribution: Risk budgets do add up. *Journal of Investment Management*, 4(4):41–51, 2006.
- R. Tyrrell Rockafellar and Stanislav Uryasev. Optimization of conditional value-at-risk. *Journal of Risk*, 2(3):21–41, 2000.
- Thierry Roncalli. *Introduction to Risk Parity and Budgeting*. CRC Press, 2013.

- William F. Sharpe. Capital asset prices: A theory of market equilibrium under conditions of risk. *The Journal of Finance*, 19(3):425–442, 1964.
- Richard S. Sutton and Andrew G. Barto. *Reinforcement Learning: An Introduction*. MIT Press, 2 edition, 2018.
- Jun Tu and Guofu Zhou. Markowitz meets talmud: A combination of sophisticated and naive diversification strategies. *Journal of Financial Economics*, 99(1):204–215, 2011.