

Phylogenetic signal in marine mammal and bird vocalizations captured by audio foundation models: the limited benefit of domain-specific pretraining

Víctor Rincón Yepes*

Abstract

Whether acoustic similarity among species reflects their evolutionary history is a fundamental question in comparative bioacoustics. Classical spectral features have shown modest phylogenetic signal in birds and frogs, but no study has tested whether large pretrained audio models, which encode richer acoustic representations, recover stronger phylogenetic structure.

Here, we apply Mantel tests to compare acoustic distance matrices derived from four audio representations with a phylogenetic distance matrix across 32 marine mammal species (1,754 recordings; Watkins Marine Mammal Sound Database). We compare hand-crafted MFCC features (105 dimensions) with three audio foundation models: the Audio Spectrogram Transformer (AST; 768d), Contrastive Language-Audio Pretraining (CLAP; 512d), and a BEATs encoder fine-tuned on bioacoustics by the Earth Species Project (BEATs-bio; 768d).

Within the 26-species cetacean clade, where divergence times are well resolved (2–34 Myr), the foundation-model embeddings recover a strong phylogenetic signal (CLAP $r = 0.82$, BEATs-bio $r = 0.82$, AST $r = 0.74$; all $p < 0.001$), among the highest acoustic–phylogenetic correlations reported for any taxon. Across the full 32-species matrix, the correlation decreases to $r \approx 0.40$ (CLAP $r = 0.410$, $p < 0.001$), largely because the constant divergence time assigned to the deep cetacean–pinniped split dilutes the signal. In contrast, hand-crafted MFCC features show no significant correlation ($r = 0.040$, $p = 0.338$).

The three foundation models did not differ significantly from one another, as paired bootstrap confidence intervals for Δr all overlapped zero. However, all three consistently outperformed MFCC. This advantage remained after projecting every embedding to the same dimensionality (PCA to the 105-dimensional MFCC space), indicating that the improvement is not simply a consequence of the higher dimensionality of the learned representations.

The observed signal is also not explained by the cetacean–pinniped dichotomy, which accounts for only 4–6% of the acoustic variance. Likewise, it persists after a partial Mantel test controlling for dominant frequency ($r_{\text{partial}} = 0.404$, $p < 0.001$), retaining 97% of the original R^2 . These results indicate that foundation models capture structural properties of vocalizations beyond simple pitch.

Extreme examples of acoustic convergence (Bearded Seal $\times$ Northern Right Whale; 90 Myr divergence, cosine distance 0.098) and divergence (Leopard Seal $\times$ Weddell Seal; 10 Myr divergence, cosine distance 0.773) illustrate how ecological pressures and lineage-specific evolution can decouple acoustic similarity from phylogenetic relatedness.

To assess whether these findings generalize beyond marine mammals, we repeated the analysis on 20 bird species (607 recordings across seven orders) and additionally evaluated BirdNET, a classifier trained on approximately 6,000 bird species. Using the time-calibrated bird phylogeny of Jetz et al. (2012), the general-purpose foundation models again recovered strong phylogenetic signal (AST $r = 0.55$, CLAP $r = 0.52$; $p < 0.001$). Surprisingly, neither BirdNET nor the bioacoustic BEATs-bio model outperformed the general-purpose models ($r \approx 0.32$ – 0.36), and the

*Independent Researcher; rin victor@gmail.com

differences among foundation models were not statistically significant, although all substantially exceeded the MFCC baseline.

Overall, our results show that pretrained audio representations encode evolutionary information across two independent radiations. They also suggest that domain-specific pretraining is not required to recover phylogenetic signal and may even reduce it. General-purpose audio models performed at least as well as models specialized for bioacoustics or birds. Finally, we discuss several possible explanations, including differences in training objectives, model architecture, and training-data diversity, and propose a controlled fine-tuning experiment to disentangle their respective contributions.

Keywords: audio foundation models, pretrained audio representations, audio embeddings, deep learning, representation learning, contrastive learning, transformers, bioacoustics, phylogenetic signal, comparative bioacoustics

1 Introduction

Animal vocalizations are shaped by three things at once: evolutionary history, ecological pressure, and individual learning. Separating them is a recurring problem in comparative bioacoustics (Odom et al., 2021). One prediction of the evolutionary account is easy to state. If evolutionary history leaves an imprint on how species sound, then species that sound alike should tend to be more closely related. That imprint has a name, *phylogenetic signal* (Blomberg & Garland, 2002), and a standard test: the Mantel correlation between an acoustic distance matrix and a phylogenetic distance matrix. It has been detected in frogs (Gingras et al., 2013; $r = 0.17$ across 90 species), in birds (Arato & Fitch, 2021; 137 species), in estrildid finches (Shimizu et al., 2023), and in gobiid fishes (Vasconcelos et al., 2021; $r = 0.47$). In every one of those studies the acoustic distances came from hand-crafted spectral features: fundamental frequency, spectral entropy, spectral centroid, Mel-frequency cepstral coefficients (MFCCs).

The set of available representations has grown a lot since then. Large pretrained audio models such as CLAP (Wu et al., 2023), AST (Gong et al., 2021), and BEATs (Chen et al., 2023) map sound into high-dimensional embedding spaces where semantically or structurally similar recordings sit close together. On bioacoustic benchmarks they clearly outperform MFCCs (Hagiwara et al., 2022; Miron et al., 2025). What is unknown is whether the geometry these models impose on audio also reflects evolutionary relationships. That is what we test.

Marine mammals are a convenient starting point. Divergence times across Cetacea and Pinnipedia are well estimated, from about 2 to 90 million years of separation (McGowen et al., 2020; Berta et al., 2018), and their calls span an unusually wide acoustic range: infrasonic fin whale pulses at 20 Hz, high-frequency dolphin clicks, and the elaborate underwater songs of Weddell seals. The Watkins Marine Mammal Sound Database (Watkins et al., 2004) provides a curated, freely available collection across 32 species.

We use the Watkins data to ask whether acoustic distances from four representations (an MFCC baseline; the general-purpose AST and CLAP; and the bioacoustic BEATs-bio) correlate with phylogenetic distance. We check whether any observed correlation just reflects dominant frequency, using a partial Mantel test, and we inspect the species pairs that sit furthest from the trend.

We then repeat the whole procedure on birds. Birds are a separate radiation with a very different vocal biology, including widespread vocal learning in oscine passerines (Arato & Fitch, 2021). They also let us ask a sharper question. Alongside the general-purpose and bioacoustic models we add BirdNET (Kahl et al., 2021), a network trained to classify around 6,000 bird species. If domain-specific pretraining is what drives phylogenetic-signal recovery, a dedicated bird model should be

the best model on birds. It is not. Explaining that negative result is the part of the paper that we found most worth thinking about.

2 Methods

2.1 Acoustic data (marine mammals)

We used the Watkins Marine Mammal Sound Database via HuggingFace (`confit/wmms-parquet`), supplemented from the full Archive.org release (`watkins-marine-mammal-sound-database-full-cuts`) for species with fewer than 20 clips in the curated set. The final dataset held 1,754 recordings across 32 species (Table S1), with at least 23 clips per species (median 55, maximum 114). Original sample rates were preserved (1,000 to 40,000 Hz, most at 22,050 Hz) for dominant-frequency estimation. Audio was resampled at each model’s input stage.

2.2 Acoustic data (birds)

For the cross-taxon replication we selected 20 bird species from seven orders: Passeriformes, Cuculiformes, Bucerotiformes, Coraciiformes, Accipitriformes, Columbiformes, and Charadriiformes (Table S2). The set spans a wide range of divergence depths, from three congeneric *Turdus* thrushes to inter-order splits close to 80 Myr, without leaving the well-characterised parts of the phylogeny. Focal recordings came from the BirdCLEF 2023 metadata (Kahl et al., 2023) and were downloaded from xeno-canto (www.xeno-canto.org), keeping only recordings rated 2.5 or better. This left 607 recordings (15 to 35 per species, median 35), resampled to 32 kHz.

2.3 Audio representations

For every representation we computed one L2-normalised acoustic centroid per species from the L2-normalised per-clip embeddings. Four representations were applied to both taxa; BirdNET was applied only to birds.

MFCC baseline. Each clip was summarised by 40 MFCCs (mean and standard deviation over time) together with chroma, spectral contrast, spectral centroid, spectral roll-off, and zero-crossing rate. This gave a 105-dimensional feature vector, computed with *librosa* 0.10 (McFee et al., 2015).

AST. We used `MIT/ast-finetuned-audioset-10-10-0.4593` (Gong et al., 2021), a transformer trained on AudioSet (Gemmeke et al., 2017) by supervised classification of audio-event tags. Audio was resampled to 16 kHz and the pooler output gave a 768-dimensional embedding per clip.

CLAP. We used `laion/clap-htsat-unfused` (Wu et al., 2023), trained on 630,000 audio-text pairs by contrastive learning. Audio was resampled to 48 kHz and the audio projection gave a 512-dimensional embedding.

BEATs-bio. We used `EarthSpeciesProject/esp-aves2-naturelm-audio-v1-beats` (Miron et al., 2025), the BEATs encoder (Chen et al., 2023) extracted from NatureLM-audio after self-supervised fine-tuning on a large bioacoustic corpus. Audio was resampled to 16 kHz and the mean over output frames gave a 768-dimensional embedding.

BirdNET (birds only). We used BirdNET GLOBAL 6K v2.4 (Kahl et al., 2021), a convolutional network trained to classify around 6,000 bird species. Audio was resampled to 48 kHz and

processed in 3-second windows. We read the 1,024-dimensional global average-pooling layer (the layer immediately before the classifier) and averaged it over windows for each clip. BirdNET is the only representation in our set trained by supervised classification of the target group; the others are trained by objectives that never see species labels.

Pairwise acoustic distances were cosine distances between L2-normalised centroids, $d_{ij} = 1 - \hat{c}_i \cdot \hat{c}_j$.

2.4 Phylogenetic distances

For marine mammals we built the 32×32 distance matrix from published divergence times: McGowen et al. (2020) for cetacean pairs, Berta et al. (2018) for pinniped pairs, and a fixed 90 Myr for cetacean–pinniped pairs (the approximate Cetacea–Carnivora split; Dos Reis et al., 2012). Distances range from 2 Myr for congenics to 90 Myr, over 496 unique pairs.

For birds we used the time-calibrated phylogeny of Jetz et al. (2012). We pulled 100 trees from the posterior (Hackett backbone, “all species” distribution) pruned to our 20 species through birdtree.org, computed each tree’s patristic distance matrix, converted it to time since the most recent common ancestor (patristic distance divided by two, since the trees are ultrametric), and averaged across the 100 trees. Three names were updated to their Jetz (2012) tip labels: *Chalcomitra amethystina* → *Nectarinia amethystina*, *Lophoceros nasutus* → *Tockus nasutus*, and *Turdus abyssinicus* → *Turdus olivaceus*. Divergence times ran from 9 Myr for the congeneric thrushes to 83 Myr, with a median of 81 Myr, so the sample is biased toward deep splits (Table S2). As a sanity check, this molecular matrix correlates at $r = 0.96$ with a simpler matrix calibrated by taxonomic rank (following Prum et al., 2015); all results below use the Jetz matrix.

2.5 Mantel tests

For each representation we correlated the acoustic and phylogenetic distance matrices with a Mantel test (Mantel, 1967): the Pearson correlation between the upper triangles, with significance from 9,999 permutations of the row–column ordering. We computed 95% confidence intervals on each r by resampling species with replacement (2,000 iterations), and we compared models with a paired bootstrap on Δr (resampling species jointly across the two matrices). Because the representations span very different dimensionalities (105 to 1,024), and cosine distances behave differently in spaces of different size, we repeated the entire analysis after projecting every embedding to the MFCC dimensionality with PCA fit at the clip level.

2.6 Partial Mantel test

To ask whether the acoustic–phylogenetic correlation is really just dominant frequency in disguise, we ran a partial Mantel test (Smouse et al., 1986) controlling for a matrix of dominant-frequency distances. Dominant frequency was the peak of the power spectrum of each clip (ignoring energy below 30 Hz), summarised per species as the median on a $\log_2$ scale. Frequency distances were $|\log_2 f_i - \log_2 f_j|$ in octaves.

3 Results

3.1 The shape of the acoustic space

Hierarchical clustering of the CLAP centroids produces a dendrogram whose first split does not follow the textbook Mysticeti/Odontoceti division (Figure 1a). Instead, species separate by frequency regime. Low-frequency baleen whales cluster together; high-frequency dolphins cluster together; and two pinnipeds, the bearded seal and the harp seal, sit with the baleen whales rather than with the other seals. The UMAP projection (Figure 1b; McInnes et al., 2018) shows the same thing more geographically: a gradient ordered by call frequency, from the fin whale at one extreme to the high-frequency dolphins at the other, with pinnipeds scattered along it according to what they sound like rather than what they are.




Figure 1: Acoustic space of 32 marine mammal species in CLAP embeddings. (a) Hierarchical dendrogram (Ward linkage, cosine distance on L2-normalised centroids), coloured by taxonomic group. The first split follows frequency, not the Mysticeti/Odontoceti division. (b) UMAP projection of the same centroids ($n_neighbors = 8$, $min_dist = 0.25$, cosine metric).

3.2 Foundation models recover the signal; MFCC does not

Mantel results are summarised in Table 1 and shown in Figures 2 and 3. MFCC features find no signal on the full 32-species matrix ($r = 0.040$, $p = 0.338$). The three foundation models do: BEATs-bio $r = 0.443$, CLAP $r = 0.410$, AST $r = 0.385$ (all $p < 0.003$). The scatter plots (Figure 3) make the difference visible. For MFCC the cloud is flat. For the foundation models cosine distance rises with divergence time, with the largest step between within-clade pairs and the cetacean–pinniped pairs at 90 Myr.




Figure 2: Mantel r for each representation against phylogenetic distance across 32 marine mammal species, with 95% bootstrap confidence intervals. Significance from 9,999 permutations (*** $p < 0.001$, ** $p < 0.01$, ns not significant). The three foundation models are statistically indistinguishable from one another, but each clearly exceeds MFCC.

Representation	Dimensions	r	p	R^2 (%)	95% CI
MFCC	105	0.040	0.338	0.2	[-0.01, 0.43]
AST	768	0.385	0.003	14.8	[0.26, 0.67]
CLAP	512	0.410	<0.001	16.8	[0.26, 0.65]
BEATs-bio	768	0.443	<0.001	19.6	[0.32, 0.68]

Table 1. Mantel results for four representations against phylogenetic distance across 32 marine mammal species (496 pairs). *p-values from 9,999 permutations; 95% confidence intervals from 2,000 species resamplings.*

The three foundation models are not distinguishable from one another at this sample size. Their bootstrap intervals overlap heavily, and the paired-bootstrap intervals on Δr all cross zero. What is unambiguous is the gap between any foundation model and MFCC. That gap also holds up under a matched-dimensionality control. After PCA-projecting each foundation embedding down to 105 dimensions, the Mantel r drops to between 0.20 and 0.26, while MFCC stays at 0.04. That is still a five- to sixfold difference. So the foundation advantage is not an artefact of embedding size. The

absolute value of the correlation does depend on dimensionality (it roughly halves under projection), which implies that the signal is distributed across a subspace wider than 105 dimensions.




Figure 3: Pairwise cosine distance against phylogenetic distance for each representation (points coloured by taxonomic group; dashed line is the least-squares fit). The MFCC cloud is flat; the three foundation models show a positive slope.

3.3 The signal is stronger within cetaceans, and it is not the pinniped split

Every cetacean–pinniped pair was assigned the same 90 Myr, so a large block of the full matrix is constant. That raises an obvious concern: perhaps the correlation only reflects “cetaceans sound different from pinnipeds,” and nothing finer than that. Two checks argue against it. First, a binary matrix encoding only the cetacean–pinniped split accounts for 4 to 6% of the acoustic

variance across models, even though it correlates almost perfectly with the phylogenetic matrix ($r = 0.96$). Second, restricting the analysis to the 26 cetaceans, which removes the flat block entirely, strengthens the correlation rather than weakening it: CLAP $r = 0.82$, AST $r = 0.74$, BEATs-bio $r = 0.82$ (all $p < 0.001$). So the signal comes from fine phylogenetic structure within a single clade, and the full-matrix values near $r = 0.4$ are conservative, watered down by the constant 90 Myr rows.

Within cetaceans even MFCC becomes significant ($r = 0.45$, $p < 0.001$). Its failure on the full matrix, then, was partly a failure to distinguish pinnipeds from cetaceans. The foundation models still lead by a wide margin within the cetacean clade.

3.4 Frequency does not explain it

A natural alternative story is that related species share body size, that body size determines pitch, and that the models are largely encoding pitch. The partial Mantel test disagrees. The bivariate correlations of dominant frequency with CLAP and with phylogeny are both weak ($r = -0.085$ and $r = -0.133$ respectively). Controlling for frequency moves the CLAP–phylogeny correlation only slightly, from 0.410 to 0.404, retaining 97% of the R^2 (Figure 4). Whatever the foundation models are picking up, it is not the dominant frequency of the call.




Figure 4: Partial Mantel analysis controlling for dominant frequency. (a) Simple Mantel r (CLAP vs phylogeny) and partial Mantel r (CLAP vs phylogeny given dominant frequency): controlling for frequency retains 97% of the R^2 (r drops from 0.410 to 0.404). (b) The three bivariate correlations; frequency is weakly related to both the acoustic and the phylogenetic distances, so it is not a confound.

3.5 Convergence and divergence

The species pairs that fall furthest from the regression line are biologically readable. The most convergent pair, closest in sound despite being far apart in the tree, is the bearded seal and the northern right whale: cosine distance 0.098 at 90 Myr apart. Bearded seal calls are long, frequency-descending trills with rich harmonic content (Risch et al., 2007), and in the embedding space they

land right next to baleen whale song. Three more cetacean–pinniped pairs appear among the ten most convergent. All of them are low-frequency species, and low-frequency Watkins recordings share a narrow recording bandwidth (Section 4.3), so part of the closeness could reflect recording conditions rather than the calls themselves. The biological reading is supported by the acoustic literature we cite, but it should be treated as a hypothesis to be checked with bandwidth-matched recordings.

The opposite pattern, close in the tree but far in sound, belongs to the leopard seal and the Weddell seal: cosine distance 0.773 at only 10 Myr apart. That is further apart in acoustic space than most cetacean-to-cetacean pairs. Weddell seals produce elaborate underwater calls year-round (Thomas & Kuechle, 1982); leopard seals produce sparse, low-frequency growls (Rogers, 2003). Two Antarctic phocids in the same family have ended up at opposite ends of the acoustic space.

3.6 The same pattern in birds, and a surprise about domain-specific models

The bird analysis reproduces the main result and adds an unexpected wrinkle (Figure 5, Table 2). Under the Jetz et al. (2012) phylogeny the general-purpose models again recover strong signal, with AST $r = 0.549$ and CLAP $r = 0.521$ (both $p < 0.001$, R^2 near 30%). Those are actually higher than the marine mammal full-matrix values. MFCC is again the weakest ($r = 0.210$, $p = 0.029$).

The unexpected part is what happens to the two domain-adapted models. Neither BEATs-bio ($r = 0.318$) nor BirdNET ($r = 0.361$) beats the general-purpose AST and CLAP. Both come in at roughly half the R^2 of the general-purpose ones. A paired bootstrap confirms that all four foundation models beat MFCC (for example $\Delta r(\text{AST}, \text{MFCC}) = +0.36$, 95% CI $[+0.20, +0.55]$), but no two foundation models differ significantly from each other, BirdNET included. A network trained end-to-end to tell 6,000 bird species apart produced embeddings that were no better, and by the point estimate slightly worse, at recovering avian phylogeny than a transformer trained on general environmental audio.

The signal survives the frequency control here as well (BirdNET $r = 0.361 \rightarrow 0.343$, retaining 90% of the R^2 , $p = 0.015$). Unlike in marine mammals, the within-clade signal is weak: restricted to the 11 passerines the Mantel r ranges from -0.05 to 0.27 across models, all with $p > 0.09$. This still holds under the molecular phylogeny, so it is not an artefact of a coarse tree. Two things probably compound. First, $n = 11$ is small. Second, oscine passerines learn their songs, and vocal learning is expected to erode phylogenetic signal at shallow depths (Arato & Fitch, 2021). The avian signal, then, rests mostly on the deep differences between orders.




Figure 5: Cross-taxon comparison. (a) Mantel r for five representations against the Jetz et al. (2012) phylogeny across 20 bird species, with 95% bootstrap intervals (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$). (b) Mantel r for marine mammals (32 species) and birds (20 species) side by side for the four shared representations, plus the bird specialist BirdNET (birds only). The general-purpose models lead in both groups; the domain-adapted models do not.

Representation	Dimensions	Training objective	r	p	R^2 (%)
MFCC	102	—	0.210	0.029	4.4
AST	768	supervised (AudioSet tags)	0.549	<0.001	30.1
CLAP	512	contrastive (audio-text)	0.521	<0.001	27.2
BEATs-bio	768	self-supervised (bioacoustics)	0.318	0.014	10.1
BirdNET	1024	supervised classification (~6,000 birds)	0.361	0.009	13.1

Table 2. Mantel results for five representations against the Jetz et al. (2012) phylogeny across 20 bird species (190 pairs). p -values from 9,999 permutations.

4 Discussion

4.1 Learned representations carry evolutionary information

The headline result is that three off-the-shelf audio models, none of them exposed to animal sound during training, recover significant phylogenetic signal in marine mammal vocalizations, while hand-crafted MFCC features do not. The effect is large within cetaceans (r near 0.8) and reproduces in a second independent radiation, birds.

MFCCs missing the signal is not, in retrospect, a surprise. They summarise local spectral texture at a fixed time scale and discard the temporal shape of the call. The foundation models were trained to discriminate sounds globally, and their embeddings appear to capture something like the *style* of a vocalization: call structure, temporal modulation, and the shape of the spectral envelope. Roughly 15 to 30% of the variance in acoustic distance travels with phylogenetic distance under those representations. Inside the cetacean clade the fraction is much higher.

4.2 Domain-specific pretraining does not help, and we cannot say why yet

The bird experiment was designed to test the domain-specificity hypothesis directly. The result was negative. BirdNET, trained end-to-end to separate 6,000 bird species, did not beat the general-purpose models on birds, and its point estimate came in lower. BEATs-bio behaved similarly. Matching the training domain to the test group is not by itself enough to improve phylogenetic-signal recovery. In our setup it did not help.

Why the domain-adapted models score lower is not identifiable from four models. Three explanations are compatible with what we see, and we cannot rule any of them out.

The first is the training objective. BirdNET is optimised by supervised species classification. That kind of pressure rewards collapsing exactly the graded, within- and between-species similarity that a Mantel test needs, pushing “how similar do a thrush and a robin sound” toward “these are different classes.” Calling this a supervision effect, though, would be wrong. AST is also trained by supervised classification, on AudioSet event tags, and it comes out the strongest of all four models. If an objective effect exists, it is fine-grained species discrimination specifically, not supervised training as a category.

The second is architecture. BirdNET is a CNN. AST, CLAP, and BEATs are transformers, whose attention may be better at capturing long-range spectro-temporal structure than convolutions are. The awkward part is that in our model set the only convolutional model is also the only species classifier, so we cannot separate an architecture effect from an objective effect. This is a real confound in the model list, not just in the interpretation.

The third is training-data breadth. The two strongest models were trained on broad general audio (AudioSet, audio-text pairs on the open web). The two weaker models were trained on narrower, specialised corpora. Squeezing the training distribution may cost some of the representational richness that happens to correlate with phylogeny.

What we can defend is the negative claim: domain-specific pretraining did not improve signal recovery in either taxon, and in birds it slightly reduced it. Separating the three explanations needs an experiment that holds architecture and data fixed and varies only the objective. Take

one pretrained encoder, AST for example, and compare the phylogenetic signal of its embeddings before and after fine-tuning it with a species-classification head. If the objective is the cause, the signal should drop under fine-tuning. If it does not, architecture and data become the more likely story. We did not run that experiment and we flag it as the natural next step.

4.3 Limitations

Several caveats apply. The marine mammal recordings come from a single historical archive spanning several decades and a range of equipment. Recording conditions are partly bound up with taxonomy, since clips of related species tend to come from the same expeditions, so sample rate and noise floor travel with phylogeny. The modal recording sample rate, expressed as an octave distance, correlates with phylogenetic distance at $r = 0.36$ and with the embedding distances at 2 to 5% of variance. Our partial Mantel controlled for dominant frequency but not for sample rate. That said, the within-cetacean signal (r near 0.8) is much larger than any plausible recording effect could produce. All models tested were trained on airborne audio, not underwater sound, and a model trained on hydrophone data might do better still.

The bird analysis is a cross-taxon replication, not a full avian study. We used 20 species biased toward deep splits, so the within-order signal is underpowered and the ordering among foundation models is not resolved. The two claims we stand behind are narrow. Every foundation model beats MFCC. The domain-specialist BirdNET does not beat the general-purpose models. Nailing down the mechanism will need more species, denser sampling within orders, and the fine-tuning experiment described in Section 4.2.

4.4 Practical implications

For anyone selecting an embedding for comparative work, the practical takeaway is not what we expected going in. Any of the three general-purpose foundation models will beat hand-crafted MFCCs by a wide margin. A model advertised for your taxon, on the other hand, is not automatically the better choice for phylogenetically structured analysis. In our setup, in two taxa, it was not.

5 Conclusions

Audio foundation models recover phylogenetic signal from vocalizations in two independent radiations, marine mammals and birds. Classical MFCC features do so only weakly, and only within a clade. In marine mammals the signal is very strong within cetaceans, survives control for the pinniped split, survives control for dominant frequency, and does not depend on embedding dimensionality. In birds it is strong at the level of orders, consistent under the Jetz et al. (2012) phylogeny, and again largely independent of the model’s training domain.

The cross-taxon comparison delivers the least expected finding. Domain-specific pretraining is neither necessary nor sufficient for strong signal recovery. A bird classifier and a bioacoustic model did no better than general-purpose transformers, and by the point estimate slightly worse. Which factor is responsible (training objective, architecture, or the breadth of the training data) cannot be identified from currently available models, and we sketch the controlled experiment that would tell them apart. For now, the safe reading is that general-purpose audio embeddings are a practical

tool for comparative bioacoustics, and that the domain label on a model is not, by itself, a reason to prefer it for this task.

6 Data and code availability

Marine mammal audio is public from the Watkins database (HuggingFace `confit/wmms-parquet`; Archive.org `watkins-marine-mammal-sound-database-full-cuts`); bird audio is from xencanto, selected through the BirdCLEF 2023 metadata. Processed embeddings, distance matrices, and all analysis code are available at <https://github.com/rinvictor/bioacoustic-phylogeny-embeddings>. Models: CLAP `laion/clap-htsat-unfused`; AST MIT/`ast-finetuned-audioset-10-10-0.4593`; BEATs-bio `EarthSpeciesProject/esp-aves2-naturelm-audio-v1-beats`; BirdNET GLOBAL 6K v2.4 (via `birdnetlib`). Supplementary Tables S1 (marine mammals) and S2 (birds) list the species, counts, and sources.

7 References

- Arato, J., & Fitch, W. T. (2021). Phylogenetic signal in the vocalizations of vocal learning and vocal non-learning birds. *Philosophical Transactions of the Royal Society B*, 376, 20200241.
- Berta, A., Churchill, M., & Boessenecker, R. W. (2018). The origin and evolutionary biology of pinnipeds: seals, sea lions, and walruses. *Annual Review of Earth and Planetary Sciences*, 46, 203–228.
- Blomberg, S. P., & Garland, T. (2002). Tempo and mode in evolution: phylogenetic inertia, adaptation and comparative methods. *Journal of Evolutionary Biology*, 15, 899–910.
- Chen, S., Wu, Y., Wang, C., Liu, S., Tompkins, D., Chen, Z., & Wei, F. (2023). BEATs: Audio pre-training with acoustic tokenizers. In *Proceedings of ICML 2023*.
- Dos Reis, M., Inoue, J., Hasegawa, M., Asher, R. J., Donoghue, P. C., & Yang, Z. (2012). Phylogenomic datasets provide both precision and accuracy in estimating the timescale of placental mammal phylogeny. *Proceedings of the Royal Society B*, 279, 3491–3500.
- Gemmeke, J. F., Ellis, D. P. W., Freedman, D., Jansen, A., Lawrence, W., Moore, R. C., . . . Ritter, M. (2017). Audio Set: An ontology and human-labeled dataset for audio events. In *Proceedings of ICASSP 2017*.
- Gingras, B., Boeckle, M., Herbst, C. T., & Fitch, W. T. (2013). Call acoustics reflect body size across four clades of anurans. *BMC Evolutionary Biology*, 13, 143.
- Gong, Y., Chung, Y.-A., & Glass, J. (2021). AST: Audio Spectrogram Transformer. In *Proceedings of Interspeech 2021*.
- Hagiwara, M., Robinson, D., Meire, E., Newby, J., Keranen, K., & Spracklen, J. (2022). BEANS: The Benchmark of Animal Sounds. *arXiv:2210.12300*.
- Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global diversity of birds in space and time. *Nature*, 491, 444–448.

- Kahl, S., Wood, C. M., Eibl, M., & Klinck, H. (2021). BirdNET: A deep learning solution for avian diversity monitoring. *Ecological Informatics*, 61, 101236.
- Kahl, S., Denton, T., Klinck, H., Reers, H., Cherutich, F., Glotin, H., ... Vellinga, W.-P. (2023). Overview of BirdCLEF 2023: Automated bird species identification in Eastern Africa. *Working Notes of CLEF 2023*.
- Mantel, N. (1967). The detection of disease clustering and a generalized regression approach. *Cancer Research*, 27, 209–220.
- McFee, B., Raffel, C., Liang, D., Ellis, D. P. W., McVicar, M., Battenberg, E., & Nieto, O. (2015). librosa: Audio and music signal analysis in Python. In *Proceedings of the 14th Python in Science Conference*.
- McGowen, M. R., Tsagkogeorga, G., Álvarez-Carretero, S., Dos Reis, M., Struebig, M., Deaville, R., ... Rossiter, S. J. (2020). Phylogenomic resolution of the cetacean tree of life using target sequence capture. *Systematic Biology*, 69, 479–501.
- McInnes, L., Healy, J., & Melville, J. (2018). UMAP: Uniform manifold approximation and projection for dimension reduction. *arXiv:1802.03426*.
- Miron, M., Robinson, D., Alizadeh, M., Gilsenan-McMahon, E., Narula, G., Chemla, E., ... Geist, M. (2025). What matters for bioacoustic encoding. In *Proceedings of ICLR 2025*.
- Odom, K. J., Araya-Salas, M., Morano, J. L., Ligon, R. A., Leighton, G. M., Taff, C. C., ... Wright, T. F. (2021). Comparative bioacoustics: A roadmap for quantifying and comparing animal sounds across diverse taxa. *Biological Reviews*, 96, 1958–1989.
- Prum, R. O., Berv, J. S., Dornburg, A., Field, D. J., Townsend, J. P., Lemmon, E. M., & Lemmon, A. R. (2015). A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature*, 526, 569–573.
- Risch, D., Clark, C. W., Corkeron, P. J., Elepfandt, A., Kovacs, K. M., Lydersen, C., Stirling, I., & Van Parijs, S. M. (2007). Vocalizations of male bearded seals, *Erignathus barbatus*: classification and geographical variation. *Animal Behaviour*, 73, 747–762.
- Rogers, T. L. (2003). Factors influencing the acoustic behaviour of male phocid seals. *Aquatic Mammals*, 29, 247–260.
- Shimizu, T., Ota, K., Suzuki, R., & Arita, T. (2023). Acoustic structure of vocalizations reflects phylogenetic relationships in estrildid finches. *Scientific Reports*, 13, 6832.
- Smouse, P. E., Long, J. C., & Sokal, R. R. (1986). Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Systematic Zoology*, 35, 627–632.
- Thomas, J. A., & Kuechle, V. B. (1982). Quantitative analysis of Weddell seal underwater vocalizations at McMurdo Sound, Antarctica. *Journal of the Acoustical Society of America*, 72, 1730–1738.
- Vasconcelos, R. P., Pratas, J., Batista, V., Costa, M. J., & Cabral, H. N. (2021). Acoustic and genetic distances in gobiid fishes. *PLOS ONE*, 16, e0260810.
- Watkins, W. A., Daher, M. A., Fristrup, K. M., Howald, T. J., & di Sciara, G. N. (2004). Marine mammal sound database. *Marine Mammal Science*, 20(1), 13–26.

Wu, Y., Chen, K., Zhang, T., Hui, Y., Berg-Kirkpatrick, T., & Dubnov, S. (2023). Large-scale contrastive language-audio pretraining with feature fusion and keyword-to-caption augmentation. In *Proceedings of ICASSP 2023*.

8 Supplementary Tables

Species are ordered by taxonomic group and common name. HF = Watkins Marine Mammal Sound Database curated subset (HuggingFace, `confit/wmms-parquet`); FA = the same database’s full archive (Archive.org, `watkins-marine-mammal-sound-database-full-cuts`), used to supplement species with fewer than 20 clips in the curated set. Dominant frequency is the median peak of the power spectrum across clips (Hz); values marked † are likely underestimates because the historical Watkins recordings for these species were sampled at 1,000–5,000 Hz (Nyquist limit 500–2,500 Hz). Cetacean divergence times are from McGowen et al. (2020), pinniped divergence times from Berta et al. (2018), and the cetacean–pinniped separation is fixed at 90 Myr (Dos Reis et al., 2012).

Mysticeti (baleen whales).

Common name	Scientific name	n clips	Dom. freq. (Hz)	Source
Bowhead Whale	<i>Balaena mysticetus</i>	60	224	HF
Fin Whale	<i>Balaenoptera physalus</i>	50	53	HF
Humpback Whale	<i>Megaptera novaeangliae</i>	64	70	HF
Minke Whale	<i>Balaenoptera acutorostrata</i>	23	75	HF+FA
Northern Right Whale	<i>Eubalaena glacialis</i>	54	165	HF
Southern Right Whale	<i>Eubalaena australis</i>	25	137	HF

Delphinidae (oceanic dolphins).

Common name	Scientific name	n clips	Dom. freq. (Hz)	Source
Atlantic Spotted Dolphin	<i>Stenella frontalis</i>	58	7,468	HF
Bottlenose Dolphin	<i>Tursiops truncatus</i>	24	6,490	HF
Clymene Dolphin	<i>Stenella clymene</i>	63	208	HF
Common Dolphin	<i>Delphinus delphis</i>	52	180	HF
Fraser’s Dolphin	<i>Lagenodelphis hosei</i>	87	683	HF
Risso’s Dolphin	<i>Grampus griseus</i>	67	849	HF
Pantropical Spotted Dolphin	<i>Stenella attenuata</i>	66	2,588	HF
Rough-toothed Dolphin	<i>Steno bredanensis</i>	50	59 †	HF
Spinner Dolphin	<i>Stenella longirostris</i>	114	60 †	HF
Striped Dolphin	<i>Stenella coeruleoalba</i>	81	180	HF
White-beaked Dolphin	<i>Lagenorhynchus albirostris</i>	57	60 †	HF
White-sided Dolphin	<i>Lagenorhynchus acutus</i>	55	60 †	HF

Non-delphinid Odontoceti (other toothed whales).

Common name	Scientific name	n clips	Dom. freq. (Hz)	Source
Beluga	<i>Delphinapterus leucas</i>	50	44 †	HF
False Killer Whale	<i>Pseudorca crassidens</i>	59	6,776	HF

Common name	Scientific name	n clips	Dom. freq. (Hz)	Source
Killer Whale	<i>Orcinus orca</i>	35	781	HF
Long-finned Pilot Whale	<i>Globicephala melas</i>	70	52 †	HF
Melon-headed Whale	<i>Peponocephala electra</i>	59	108	HF
Narwhal	<i>Monodon monoceros</i>	50	60 †	HF
Short-finned Pilot Whale	<i>Globicephala macrorhynchus</i>	67	1,915	HF
Sperm Whale	<i>Physeter macrocephalus</i>	75	174	HF

Pinnipedia (seals and walrus).

Common name	Scientific name	n clips	Dom. freq. (Hz)	Source
Bearded Seal	<i>Erignathus barbatus</i>	37	319	HF
Harp Seal	<i>Pagophilus groenlandicus</i>	47	42 †	HF
Leopard Seal	<i>Hydrurga leptonyx</i>	25	352	HF+FA
Ross Seal	<i>Ommatophoca rossii</i>	50	325	HF
Walrus	<i>Odobenus rosmarus</i>	38	341	HF
Weddell Seal	<i>Leptonychotes weddellii</i>	42	120	HF+FA

Table S1. The 32 marine mammal species included in the analysis (32 species; 1,754 clips; n per species: min 23, median 55, max 114).

Focal bird recordings came from the BirdCLEF 2023 metadata (Kahl et al., 2023) downloaded from xeno-canto (quality rating ≥ 2.5). Pairwise divergence times come from the time-calibrated phylogeny of Jetz et al. (2012): 100 posterior Hackett-backbone trees pruned via birdtree.org and averaged as time since the most recent common ancestor. Divergence times span 9 Myr (congeneric *Turdus*) to 83 Myr (inter-order), median 81 Myr; the sample is weighted toward deep divergences. † = species mapped to a different Jetz (2012) tip label.

Common name	Scientific name	Family	Order	n
Barn Swallow	<i>Hirundo rustica</i>	Hirundinidae	Passeriformes	35
African Paradise Flycatcher	<i>Terpsiphone viridis</i>	Monarchidae	Passeriformes	35
African Pied Wagtail	<i>Motacilla aguimp</i>	Motacillidae	Passeriformes	35
African Dusky Flycatcher	<i>Muscicapa adusta</i>	Muscicapidae	Passeriformes	22
Amethyst Sunbird	<i>Chalcomitra amethystina</i> †	Nectariniidae	Passeriformes	35
African Black-headed Oriole	<i>Oriolus larvatus</i>	Oriolidae	Passeriformes	35
Baglafecht Weaver	<i>Ploceus baglafecht</i>	Ploceidae	Passeriformes	23
Blue-headed Crested Flycatcher	<i>Elminia longicauda</i>	Stenostiridae	Passeriformes	18
Abyssinian Thrush	<i>Turdus abyssinicus</i> †	Turdidae	Passeriformes	26
African Thrush	<i>Turdus pelios</i>	Turdidae	Passeriformes	35
Bare-eyed Thrush	<i>Turdus tephronotus</i>	Turdidae	Passeriformes	15
African Emerald Cuckoo	<i>Chrysococcyx cupreus</i>	Cuculidae	Cuculiformes	35
Black-and-white-casqued Hornbill	<i>Bycanistes subcylindricus</i>	Bucerotidae	Bucerotiformes	34
African Grey Hornbill	<i>Lophoceros nasutus</i> †	Bucerotidae	Bucerotiformes	35
Blue-cheeked Bee-eater	<i>Merops persicus</i>	Meropidae	Coraciiformes	35
African Goshawk	<i>Accipiter tachiro</i>	Accipitridae	Accipitriformes	35
African Fish Eagle	<i>Haliaeetus vocifer</i>	Accipitridae	Accipitriformes	35
African Mourning Dove	<i>Streptopelia decipiens</i>	Columbidae	Columbiformes	32
African Green Pigeon	<i>Treron calvus</i>	Columbidae	Columbiformes	23
African Jacana	<i>Actophilornis africanus</i>	Jacanidae	Charadriiformes	29

Table S2. The 20 bird species used for the avian replication (20 species; 607 recordings; 7 orders; n per species: min 15, median 35, max 35). Renamings for Jetz (2012) tip labels: *Chalcomitra amethystina* → *Nectarinia amethystina*; *Lophoceros nasutus* → *Tockus nasutus*; *Turdus abyssinicus* → *Turdus olivaceus*.