

Pointwise Majorization for sub-Weibull and Mixed Tail Processes with Applications in Quadratic Chaos and Ergodic Diffusions

Haichen Hu^{‡1} David Simchi-Levi^{†1}

¹*Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA*

Abstract

Classical chaining controls an indexed stochastic process through a single worst-case bound, which can obscure substantial variation across the index set. We establish the first simultaneous pointwise majorization theory for Banach-valued processes with sub-Weibull or two-metric mixed-tail increments. For an anchored sub-Weibull process on a separable index space, write $v(t) := d(t, t_0)$. Given a reference measure μ , the envelope at t is governed by the pointwise Fernique-Talagrand functional of order α , $\Phi_{\mu,d}^{(\alpha)}(t) := \int_0^{4v(t)} (\log \frac{1}{\mu(B_d(t,r))})^{1/\alpha} dr$. $\forall \delta \in (0, 1)$, we obtain that

$$\mathbb{P}(\|Z_t\| \lesssim \{\Phi_{\mu,d}^{(\alpha)}(t) + v(t)(\log(e/\delta))^{1/\alpha}\}, \forall t) \geq 1 - \delta.$$

Our bound is determined by the pointwise complexity $\Phi_{\mu,d}^{(\alpha)}$ rather than a global quantity. The result holds for every $\alpha > 0$ and does not involve dyadic logarithmic terms from peeling. For mixed tail processes, with fixed measures μ_1, μ_2 and $v_j(t) := d_j(t, t_0)$, $\Phi_j(t) := \int_0^{4v_j(t)} (\log \frac{1}{\mu_j(B_{d_j}(t,r))})^{1/\alpha_j} dr, j = 1, 2$, for any $\delta \in (0, 1)$, we show that

$$\mathbb{P}(\|Z_t\| \lesssim \sum_{j=1}^2 \{\Phi_j(t) + v_j(t)(\log \frac{e}{\delta})^{1/\alpha_j}\}, \forall t) \geq 1 - \delta.$$

Although the two regimes are coupled in the mixed tail condition, each retains its own pseudo-metric, reference measure, pointwise Fernique-Talagrand functional, and tail exponent. The proof tracks the index-wise costs of measure-generated admissible chains and synchronizes them through a nested common refinement. For applications, we derive matrix-specific bounds for centered quadratic chaos under pseudo-metrics induced by the operator and Frobenius norms, and observable-specific finite-time bounds for diffusion empirical processes.

1 Introduction

Uniform control of stochastic processes is a basic problem in statistics and applied probability. Empirical processes govern uniform deviations of estimators and the validity of data-dependent

[‡]Email: huhc@mit.edu

[†]Email: dslevi@mit.edu

selection. This paper studies a refinement of this problem: rather than assigning one worst-case threshold to the full index set, we seek a simultaneous threshold adapted to each individual index.

Generic chaining estimates the supremum of an indexed stochastic process from the tail behavior of its increments and the metric complexity of its index set. Its classical background includes the entropy method of [Dudley \(1967\)](#), the pathwise inequality of [Garsia et al. \(1971\)](#), and the majorizing-measure program of [Fernique \(1971\)](#); [Preston \(1972\)](#); [Fernique \(1975\)](#). Talagrand’s converse and subsequent generic-chaining formulations identify the optimized worst-center ball-mass functional with the expected supremum of a Gaussian process, up to universal constants ([Talagrand, 1987, 1992, 1994, 1996, 2001, 2005](#)). This theory has also led to algorithmic constructions of majorizing measures ([Borst et al., 2021](#)) and to global bounds for general Orlicz, sub-Weibull, and mixed-tail processes ([Talagrand, 1990](#); [Dirksen, 2015](#)). Its standard output, however, is a single bound for the supremum, and therefore a single worst-case complexity for the entire index set.

However, in many problems, a global threshold is not always the desired output. One may inspect the same random field and then select an index whose local metric complexity is substantially smaller than the largest part of the parameter space. A fixed-index concentration inequality retains the scale of that index, but it cannot be substituted for a random index selected from the field. A bound for the supremum is valid after such a selection, but it charges every index with the worst-case complexity of the full set. We seek both properties at once. For any $\delta \in (0, 1)$, we have

$$\mathbb{P}(\forall t, \|Z_t\| \leq \mathfrak{M}_\delta(t)) \geq 1 - \delta, \quad (1.1)$$

where the deterministic threshold $\mathfrak{M}_\delta(t)$ reflects the local metric complexity at that same index. We refer to inequality (1.1) as a *simultaneous pointwise majorization*. It remains valid at any measurable index $\hat{t}$ chosen using the observed samples.

The purpose of this paper is to construct pointwise envelopes directly from increment inequalities. The basic idea is to use a fixed ambient measure to build one admissible chain while retaining its deterministic cost at every individual index instead of taking a supremum over indices during the construction. For mixed tails, we build one chain for each metric and combine them through a nested common refinement, so that all metric-specific chains telescope along the same process increments without losing their separate pointwise costs. This produces an additive local-complexity term and an anchor-radius confidence term at every index, without allocating failure probability over complexity or radius shells.

To illustrate the result, let $(Z_t)_{t \in T}$ be a Banach-valued process anchored at t_0 , so that $Z_{t_0} = 0$, and suppose that for some $\alpha > 0$ and pseudo-metric d , $\{Z_t\}_{t \in T}$ satisfied a sub-weibull increment, i.e.,

$$\mathbb{P}\left\{\|Z_t - Z_s\| > u^{1/\alpha} d(s, t)\right\} \leq 2e^{-u}, \quad s, t \in T, \quad u \geq 0.$$

For a Borel probability measure μ on T , fixed before observing the process, define

$$v(t) := d(t, t_0), \quad \Phi_{\mu, d}^{(\alpha)}(t) := \int_0^{4v(t)} \left(\log \frac{1}{\mu(B_d(t, r))} \right)^{1/\alpha} dr.$$

Our first theorem shows that, with probability at least $1 - \delta$, simultaneously for every $t \in T$,

$$\|Z_t\| \leq C_\alpha \left\{ \Phi_{\mu, d}^{(\alpha)}(t) + v(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha} \right\}. \quad (1.2)$$

The integral in (1.2) measures the ambient mass available around t over scales comparable to its distance from the anchor. The confidence term is scaled by the same individual radius $v(t)$, rather than by the diameter of T . Thus, the conclusion is uniform in its probability event but nonuniform in its deterministic threshold.

The finite Gaussian result of Xu (2026) is the direct precursor to our bound. The bound there is a fixed-measure ball-mass integral truncated at the index-dependent anchor radius, and the deviation term is scaled by that radius. Its stated upper bound, however, contains deterministic baseline scales and logarithmic losses arising from a common dyadic peeling argument. Our proof avoids this peeling by constructing a measure-generated admissible chain whose cost is tracked at each index rather than first replaced by a worst-case value over a radius or complexity shell; a single chaining event then yields the envelopes simultaneously for all indices. Thus, even in the Gaussian case $\alpha = 2$, inequality (1.2) sharpens the bound of Xu (2026) by removing the baseline scales and peeling logarithms while also extending the theory to sub-Weibull processes with order $\alpha > 0$, including the nonconvex range $0 < \alpha < 1$, and to separable infinite index spaces.

Our second theorem treats stochastic processes with genuinely mixed increment condition. Suppose that for two pseudo-metrics d_1, d_2 and exponents $\alpha_1, \alpha_2 > 0$, $\{Z_t\}_{t \in T}$ has a mixed tail increment:

$$\mathbb{P} \left\{ \|Z_t - Z_s\| > u^{1/\alpha_1} d_1(s, t) + u^{1/\alpha_2} d_2(s, t) \right\} \leq 2e^{-u}.$$

The same parameter u controls both regimes; the assumption neither yields the two single-metric increment inequalities separately nor presupposes a decomposition of Z into two processes. For fixed Borel probability measures μ_1, μ_2 , put

$$v_j(t) := d_j(t, t_0), \quad \Phi_j(t) := \int_0^{4v_j(t)} \left(\log \frac{1}{\mu_j(B_{d_j}(t, r))} \right)^{1/\alpha_j} dr, \quad j \in \{1, 2\}.$$

We prove that, with probability at least $1 - \delta$, every $t \in T$ satisfies

$$\|Z_t\| \leq C_{\alpha_1, \alpha_2} \sum_{j=1}^2 \left\{ \Phi_j(t) + v_j(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha_j} \right\}. \quad (1.3)$$

Each tail regime, therefore, retains its own metric, ambient measure, ball-mass term, and confidence scale. The common refinement is essential: it makes the two deterministic chains control the same process edges and is not an application of the single-metric theorem twice.

Related approaches include minorizing-metric and chaining bounds for sample-path regularity, suprema, and concentration (Kwapień and Rosiński, 2004; Bednorz, 2007, 2006, 2014). Learning-theoretic work develops local or problem-dependent complexity methods (Bartlett et al., 2005; Koltchinskii, 2006; Xu and Zeevi, 2025), as well as PAC-Bayesian and multiscale coding formulations (Audibert and Bousquet, 2007; Maurer, 2010; Chu and Raginsky, 2026). However, none of these results gives the particular simultaneous pointwise envelope proved here: on one event, every index is bounded by the sum of its own fixed-measure ball-mass integral and an anchor-distance confidence term, with this decomposition preserved across distinct mixed-tail metrics.

We illustrate our pointwise majorization theorem with two examples. The first is quadratic chaos, which arises in covariance estimation, random embeddings, and restricted-isometry analysis. Hanson–Wright inequalities and their fixed-form extensions provide the basic concentration theory (Hanson and Wright, 1971; Rudelson and Vershynin, 2013; Hsu et al., 2012; Adamczak, 2015), while

uniform chaos bounds underpin random-matrix and structured-embedding results (Mendelson et al., 2008; Krahmer et al., 2014; Dirksen, 2015; Klochkov and Zhivotovskiy, 2020). Our application retains matrix-specific information across a heterogeneous family without sacrificing simultaneous validity.

The second is the occupation empirical process of a stationary scalar diffusion, a basic object in invariant-distribution and density estimation. Its long-run and empirical process theory is classical (Kipnis and Varadhan, 1986; van Zanten, 2003; van der Vaart and van Zanten, 2005), while finite-time concentration, class-wide bounds, and extensions to broader diffusion models have been developed in Cattiaux and Guillin (2008); Redig and Völlering (2011); Gao et al. (2014); Pepin (2021); Aeckerle-Willems and Strauch (2021); Trottnier et al. (2023). Our application complements fixed-observable and worst-case class bounds with simultaneous finite-time control that adapts to the observable ultimately selected.

2 Pointwise Majorization for Sub-Weibull Processes

In this section, we establish a pointwise majorization theorem for sub-Weibull processes. Let $(\mathbb{B}, \|\cdot\|)$ be a separable Banach space, let $(T, \mathcal{A})$ be a standard Borel space. $d : T \times T \rightarrow [0, \infty)$ is a finite-valued, jointly $\mathcal{A} \otimes \mathcal{A}$ -measurable pseudo-metric such that (T, d) is separable. All random objects below are defined on a fixed probability space $(\Omega, \mathcal{H}, \mathbb{P})$.

Let $Z : \Omega \times T \rightarrow \mathbb{B}$ be jointly measurable from $(\Omega \times T, \mathcal{H} \otimes \mathcal{A})$ to $(\mathbb{B}, \mathcal{B}(\mathbb{B}))$, and write the stochastic process as $Z_t(\omega) := Z(\omega, t)$, $\omega \in \Omega$, $t \in T$. Thus, $(Z_t)_{t \in T}$ is a $\mathbb{B}$ -valued stochastic process indexed by T .

Now, we introduce some standard notation in studying empirical processes. First, we assume that $\{Z_t\}_{t \in T}$ has a continuous modification such that the map $t \mapsto Z_t(\omega)$ is continuous from (T, d) to $(\mathbb{B}, \|\cdot\|)$. Second, we assume that there exists an anchor $t_0 \in T$ such that $Z_{t_0} = 0$ almost surely.

To summarize, there exists $t_0 \in T$ and an event $\Omega_0 \in \mathcal{H}$ with $\mathbb{P}(\Omega_0) = 1$ such that, for every $\omega \in \Omega_0$, $Z_{t_0}(\omega) = 0$, the map $t \mapsto Z_t(\omega)$ is continuous from (T, d) to $(\mathbb{B}, \|\cdot\|)$. Throughout this paper, we will work conditioned on the event Ω_0 and thus we omit its dependence.

We say that $\{Z_t\}_{t \in T}$ is an *sub-Weibull process* of order $\alpha > 0$ with respect to d if, for every $s, t \in T$ and every $u \geq 0$,

$$\mathbb{P} \left\{ \|Z_t - Z_s\| > u^{1/\alpha} d(t, s) \right\} \leq 2e^{-u}.$$

The sub-Weibull process incorporates a wide range of stochastic processes. Specifically, if $\alpha = 2$, then $\{Z_t\}_{t \in T}$ is a sub-Gaussian process, and $\alpha = 1$ corresponds to the sub-exponential process.

To study the suprema of our sub-Weibull process, we recall some background about generic chaining. For the pseudo-metric space (T, d) , a sequence $\mathcal{T} = (T_n)_{n \geq 0}$ of subsets of T is *admissible* if $|T_0| = 1$, $|T_n| \leq 2^{2^n}$. The α -Talagrand functional (Dirksen, 2015) is defined as

$$\gamma_\alpha(T, d) := \inf_{(T_n)} \sup_{t \in T} \sum_{n \geq 0} 2^{n/\alpha} d(t, T_n), \quad d(t, T_n) = \inf_{s \in T_n} d(t, s).$$

The diameter of any set $S \subset T$ with respect to d is defined as $\text{diam}_d(S) = \sup_{s, t \in S} d(t, s)$. Moreover, we use $B_d(t, r)$ to denote the local neighborhood centered at t with radius r , i.e., $B_d(t, r) :=$

$\{s : d(t, s) \leq r\}$. We can also define the α -Talagrand functional on the subset S as

$$\gamma_\alpha(S, d) := \inf_{(S_n)} \sup_{t \in S} \sum_{n \geq 0} 2^{n/\alpha} d(t, S_n), \quad d(t, S_n) = \inf_{s \in S_n} d(t, s),$$

where S_n is an admissible sequence of subsets of S .

Now, we return to the discussion of the sub-Weibull process $\{Z_t\}_{t \in T}$. With the anchor t_0 such that $Z_{t_0} = 0$, we define the function $v(t) := d(t, t_0)$.

Fixing a Borel probability measure μ on $(T, \mathcal{B})$ before observing the process, for any $\alpha > 0$, the pointwise Fernique–Talagrand functional (Xu, 2026) is defined as

$$\Phi_{\mu, d}^{(\alpha)}(t) := \int_0^{4v(t)} \left(\log \frac{1}{\mu(B_d(t, u))} \right)^{1/\alpha} du.$$

Furthermore, we define the following function $J_b(\cdot)$. For any $a, b > 0$, we denote

$$J_b(x) := 1 + \lceil \log_2^+(x/b) \rceil, \quad \log_a^+(y) := \max \{ \log_a y, 0 \}.$$

Then, we have the following pointwise majorization theorem regarding the process $\{Z_t\}_{t \in T}$.

Theorem 2.1. Under the preceding setup, let $\{Z_t\}_{t \in T}$ be an anchored sub-Weibull process of order $\alpha > 0$, and fix a Borel probability measure μ on $(T, \mathcal{A})$. There exists a finite constant C_α , depending only on α , such that, for every $\delta \in (0, 1)$, with probability at least $1 - \delta$, simultaneously for every $t \in T$, we have

$$\|Z_t\| \leq C_\alpha \left[\Phi_{\mu, d}^{(\alpha)}(t) + v(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha} \right].$$

The conclusion of Theorem 2.1 is pointwise in its deterministic envelope but simultaneous in its probability event. More precisely, there exists a single event of probability at least $1 - \delta$ on which the displayed inequality holds for every $t \in T$, while the right-hand side depends on the local geometric quantities associated with the individual index t .

The first term,

$$\Phi_{\mu, d}^{(\alpha)}(t) = \int_0^{4v(t)} \left(\log \frac{1}{\mu(B_d(t, r))} \right)^{1/\alpha} dr,$$

measures the local multiscale complexity around t . At each scale r , the quantity $\mu(B_d(t, r))$ records the mass assigned by the ambient prior measure μ to the neighborhood of t . Thus, larger local ball masses give a smaller integrand and hence a smaller pointwise complexity. Moreover, the integral is taken only up to $4v(t)$, so the complexity profile involves only scales comparable with the distance from t to the anchor.

The second term is the pointwise confidence penalty. Since $Z_{t_0} = 0$ almost surely and $v(t) = d(t, t_0)$, the sub-Weibull increment condition gives, for every $u \geq 0$, we have

$$\mathbb{P} \left\{ \|Z_t\| > u^{1/\alpha} v(t) \right\} = \mathbb{P} \left\{ \|Z_t - Z_{t_0}\| > u^{1/\alpha} d(t, t_0) \right\} \leq 2e^{-u}.$$

Therefore, $v(t)$ is the natural fluctuation scale at the index t . Theorem 2.1 preserves this pointwise

scale in the simultaneous bound: the confidence penalty is proportional to $v(t) (\log \frac{e}{\delta})^{1/\alpha}$, rather than to the global diameter of T .

An important feature of the theorem is that no dyadic-shell terms appear in the conclusion. The local complexity $\Phi_{\mu,d}^{(\alpha)}(t)$ already records the multiplicity of the relevant scales through the ball masses $\mu(B_d(t,r))$. In the proof, we retain the pointwise cost of the measure-generated admissible chain. Thus, we do not first replace this cost by a supremum over a subset and then distribute the failure probability among dyadic shells. This is why no shell-allocation logarithms appear.

More generally, the power $1/\alpha$ in both the local profile and the confidence term reflects the order of the assumed sub-Weibull increments.

For $\alpha = 2$, Theorem 2.1 gives the sub-Gaussian bound

$$\|Z_t\| \leq C_2 \left[\Phi_{\mu,d}^{(2)}(t) + v(t) \sqrt{\log \frac{e}{\delta}} \right]$$

simultaneously for every $t \in T$. It improves the previous results in Xu (2026) by removing the auxiliary peeling logarithms from the upper envelope. For $\alpha = 1$, the corresponding sub-exponential confidence penalty is $v(t) \log \frac{e}{\delta}$.

The proof of Theorem 2.1 relies on a series of lemmas, and the full proof is deferred to Appendix B.

3 Pointwise Majorization for Mixed Tail Processes

In this section, we extend the pointwise majorization theorem in Section 2 to stochastic processes whose increments satisfy a two-regime tail bound. The two regimes may have different tail exponents and may be governed by different pseudo-metrics.

The same setup in this section is almost the same as that in Section 2. Let $(\mathbb{B}, \|\cdot\|)$ be a separable Banach space, and let $(T, \mathcal{A})$ be a standard Borel space. Let $d_1, d_2 : T \times T \rightarrow [0, \infty)$ be finite-valued, jointly $\mathcal{A} \otimes \mathcal{A}$ -measurable pseudo-metrics. Define the pseudo-metric ρ as

$$\rho(s, t) := d_1(s, t) + d_2(s, t), \quad s, t \in T,$$

and assume that (T, ρ) is separable. As in Section 2, all random objects below are defined on a fixed probability space $(\Omega, \mathcal{H}, \mathbb{P})$.

Let $Z : \Omega \times T \rightarrow \mathbb{B}$ be jointly measurable from $(\Omega \times T, \mathcal{H} \otimes \mathcal{A})$ to $(\mathbb{B}, \mathcal{B}(\mathbb{B}))$, and write

$$Z_t(\omega) := Z(\omega, t), \quad \omega \in \Omega, \quad t \in T.$$

We still assume that the process has a modification, which is also denoted by $(Z_t)_{t \in T}$. Similar to Section 2, there exists an anchor $t_0 \in T$ and an event $\Omega_0 \in \mathcal{H}$ with $\mathbb{P}(\Omega_0) = 1$ such that, for every $\omega \in \Omega_0$, $Z_{t_0}(\omega) = 0$ and the map $t \mapsto Z_t(\omega)$ is continuous from (T, ρ) to $(\mathbb{B}, \|\cdot\|)$, and we work with this modification throughout this section.

$(Z_t)_{t \in T}$ is said to be a two-metric mixed-tail process of orders $\alpha_1, \alpha_2 > 0$ with respect to (d_1, d_2) if, for every $s, t \in T$ and every $u \geq 0$, we have that

$$\mathbb{P} \left\{ \|Z_t - Z_s\| > u^{1/\alpha_1} d_1(t, s) + u^{1/\alpha_2} d_2(t, s) \right\} \leq 2e^{-u}. \quad (3.1)$$

The same parameter u controls the two terms in (3.1), but each term retains its own tail exponent and pseudo-metric.

The distinctive feature of (3.1) is that one deviation parameter activates two fluctuation regimes on the same probability event. The two terms are therefore not separate tail bounds that are assumed independently. In particular, neither d_1 nor d_2 is required by itself to control the increments. When $(\alpha_1, \alpha_2) = (1, 2)$, the threshold takes the familiar Bernstein form

$$ud_1(s, t) + \sqrt{u}d_2(s, t).$$

In this case, d_2 describes the variance-driven sub-Gaussian regime, whereas d_1 describes the sub-exponential correction that becomes decisive for larger deviations. More general pairs (α_1, α_2) allow the two geometries to have different sub-Weibull orders.

This structure occurs whenever an increment inequality involves two genuinely different scale parameters. For a centered empirical process based on m independent observations, a Bernstein inequality typically produces a variance metric of the form $m^{-1/2}\|f_t - f_s\|_{L^2(P)}$ and an envelope metric of the form $m^{-1}\|f_t - f_s\|_\infty$; this is the standard mixed sub-Gaussian-sub-exponential setting (Dirksen, 2015). For matrix-indexed centered quadratic forms of a vector with independent centered sub-Gaussian coordinates, the Hanson–Wright inequality yields a Frobenius-norm metric in the sub-Gaussian term and an operator-norm metric in the sub-exponential term (Rudelson and Vershynin, 2013). Centered additive functionals of symmetric or reversible Markov processes provide another example when a Poincaré or related functional inequality yields a finite-time Bernstein form (Gao et al., 2014). The two applications developed later in this paper: quadratic chaos and stationary scalar diffusions, are concrete instances of the latter two mechanisms.

Fix Borel probability measures μ_1 and μ_2 on $(T, \mathcal{A})$. For $j \in \{1, 2\}$, we define the pointwise anchor radius $v_j(t) := d_j(t, t_0)$ and the pointwise Fernique–Talagrand functional as

$$\Phi_{\mu_j, d_j}^{(\alpha_j)}(t) := \int_0^{4v_j(t)} \left(\log \frac{1}{\mu_j(B_{d_j}(t, r))} \right)^{1/\alpha_j} dr,$$

where $B_{d_j}(t, r) := \{s \in T : d_j(s, t) \leq r\}$. The integrals are understood in the extended-real sense, with $\log(1/0) := +\infty$. For notational simplicity, throughout this section and its appendix, we write

$$\Phi_j(t) := \Phi_{\mu_j, d_j}^{(\alpha_j)}(t), \quad j \in \{1, 2\}.$$

We now state the main theorem of this section.

Theorem 3.1. Under the preceding setup, let $\{Z_t\}_{t \in T}$ be an anchored two-metric mixed-tail process of orders $\alpha_1, \alpha_2 > 0$. There exists a finite constant C_{α_1, α_2} , depending only on α_1 and α_2 , such that, for every $\delta \in (0, 1)$, with probability at least $1 - \delta$, simultaneously for every $t \in T$, we have

$$\|Z_t\| \leq C_{\alpha_1, \alpha_2} \left\{ \Phi_1(t) + \Phi_2(t) + v_1(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha_1} + v_2(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha_2} \right\}.$$

If either $\Phi_1(t)$ or $\Phi_2(t)$ is infinite, the corresponding inequality holds trivially.

Theorem 3.1 holds on one event simultaneously for every $t \in T$, but it retains two different local geometries at each index. The measure μ_j and the term $\Phi_j(t)$ describe the local complexity in the

d_j -geometry, whereas $v_j(t) = d_j(t, t_0)$ is the corresponding pointwise fluctuation scale. In particular, the two reference measures need not be the same.

For the frequently occurring sub-exponential–sub-Gaussian ordering $(\alpha_1, \alpha_2) = (1, 2)$, the theorem gives

$$\|Z_t\| \leq C_{1,2} \left\{ \Phi_1(t) + \Phi_2(t) + v_1(t) \log \frac{e}{\delta} + v_2(t) \sqrt{\log \frac{e}{\delta}} \right\}$$

simultaneously for every $t \in T$. If one of the pseudo-metrics is identically zero, its profile and anchor radius vanish, and the result reduces to the sharpened single-metric theorem in Section 2.

The proof of Theorem 3.1 relies on a series of lemmas, and the full proof is deferred to Appendix C.

In the next two sections, we will apply the pointwise majorization theorems to important objects in probability and statistics to obtain new concentration bounds for them

4 Pointwise Envelopes for Quadratic Chaos Processes

In this section, we present the first application of our pointwise majorization results. We apply Theorem 3.1 to a matrix-indexed family of centered quadratic forms and derive a simultaneous pointwise envelope for the associated quadratic chaos process.

Quadratic chaos processes describe the simultaneous fluctuations of quadratic functionals of a common random vector. Uniform quadratic-form bounds arise in covariance-operator estimation, covariance estimation with missing observations, and variance-components estimation (Adamczak, 2015; Klochov and Zhivotovskiy, 2020; Dicker and Erdogdu, 2017). Suprema of second-order chaos processes are also central to structured random-matrix, random-embedding, and restricted-isometry problems (Krahmer et al., 2014; Dirksen, 2015).

The Hanson–Wright inequality reveals two different increment geometries about quadratic chaos processes: the Frobenius norm governs the sub-Gaussian regime, whereas the operator norm governs the sub-exponential regime. The application below combines this two-scale inequality with Theorem 3.1, producing one simultaneous bound that retains the local complexity of each individual matrix in both geometries.

For a real-valued random variable Y , we use the exponential Orlicz convention

$$\|Y\|_{\psi_2} := \inf \left\{ r > 0 : \mathbb{E}[\exp(Y^2/r^2)] \leq 2 \right\}.$$

Let $\xi = (\xi_1, \dots, \xi_n)^\top \in \mathbb{R}^n$ be a random vector with independent, mean-zero, real-valued coordinates such that $\max_{1 \leq i \leq n} \|\xi_i\|_{\psi_2} \leq M_0$ for some $M_0 > 0$. Let $(T, \mathcal{B})$ be a standard Borel space, and let $t \mapsto B_t$ be a Borel measurable map from T to the space of $n \times n$ deterministic real symmetric matrices. We use $B_t = (B_{ij}(t))_{1 \leq i, j \leq n}$ as the corresponding matrix class.

Definition 4.1. The centered *quadratic chaos process* indexed by $\{B_t\}_{t \in T}$ is defined by

$$C_t(\xi) := \xi^\top B_t \xi - \mathbb{E}_\xi \left[\xi^\top B_t \xi \right] = \sum_{i,j=1}^n B_{ij}(t) (\xi_i \xi_j - \mathbb{E}_\xi [\xi_i \xi_j]), \quad t \in T.$$

Following the setting in Section 3, we assume that there exists an anchor $t_0 \in T$ such that $B_{t_0} = 0$.

Then $C_{t_0}(\xi) = 0$ almost surely. For a matrix $B \in \mathbb{R}^{n \times n}$, we define $\|B\|_F = (\sum_{i,j=1}^n B_{ij}^2)^{1/2}$ to be the Frobenius norm and $\|B\|_{\text{op}} = \sup_{\|x\|_2=1} \|Bx\|_2$ to be the operator norm.

By the Hanson-Wright inequality (Lemma A.1), we have the following proposition about the quadratic chaos process.

Proposition 4.2. If $\max_{1 \leq i \leq n} \|\xi_i\|_{\psi_2} \leq M_0$, set $C_{HW} = 2c_{HW}$, where c_{HW} is the universal constant in Lemma A.1. Define two metrics

$$d_1(B_s, B_t) := C_{HW} M_0^2 \|B_s - B_t\|_{\text{op}}, \quad d_2(B_s, B_t) := C_{HW} M_0^2 \|B_s - B_t\|_F,$$

and set $\rho = d_1 + d_2$. Then, the maps d_1 and d_2 are finite-valued, jointly Borel pseudo-metrics, and (T, ρ) is separable. For every $s, t \in T$ and every $u \geq 0$, we have that

$$\mathbb{P}_\xi \{ |C_s(\xi) - C_t(\xi)| > u d_1(s, t) + \sqrt{u} d_2(s, t) \} \leq 2e^{-u}.$$

The process is jointly measurable and has almost surely ρ -continuous sample paths. Consequently, $\{C_t(\xi)\}_{t \in T}$ is an anchored mixed-tail process with order $(\alpha_1, \alpha_2) = (1, 2)$.

Fix two probability measures μ_1 and μ_2 on T before observing ξ . With Proposition 4.2, we define the pointwise anchor radii

$$v_1(t) := d_1(t, t_0) = C_{HW} M_0^2 \|B_t\|_{\text{op}}, \quad v_2(t) := d_2(t, t_0) = C_{HW} M_0^2 \|B_t\|_F.$$

For either pseudo-metric $d \in \{d_1, d_2\}$, write $B_d(t, r) := \{s \in T : d(s, t) \leq r\}$. The Talagrand functionals are

$$\begin{aligned} \Phi_1(t) &= \int_0^{4v_1(t)} \log \frac{1}{\mu_1(B_{d_1}(t, r))} dr = C_{HW} M_0^2 \int_0^{4\|B_t\|_{\text{op}}} \log \frac{1}{\mu_1\{s \in T : \|B_s - B_t\|_{\text{op}} \leq r\}} dr, \\ \Phi_2(t) &= \int_0^{4v_2(t)} \left(\log \frac{1}{\mu_2(B_{d_2}(t, r))} \right)^{1/2} dr = C_{HW} M_0^2 \int_0^{4\|B_t\|_F} \left(\log \frac{1}{\mu_2\{s \in T : \|B_s - B_t\|_F \leq r\}} \right)^{1/2} dr. \end{aligned}$$

Corollary 4.3. There exists a universal constant $C_{1,2} < \infty$ such that, for every $\delta \in (0, 1)$, with probability at least $1 - \delta$, simultaneously for every $t \in T$, we have

$$|C_t(\xi)| \leq C_{1,2} \left\{ \Phi_1(t) + \Phi_2(t) + v_1(t) \log \frac{e}{\delta} + v_2(t) \sqrt{\log \frac{e}{\delta}} \right\}.$$

The proof of Corollary 4.3 is deferred to Appendix D.

The first two terms describe the local complexity arising from the simultaneous control of the entire matrix family. More precisely, $\Phi_1(t)$ measures the multiscale complexity around B_t in the operator-norm geometry, while $\Phi_2(t)$ measures the corresponding complexity in the Frobenius-norm geometry.

The last two terms preserve the deviation scales of the fixed-matrix Hanson-Wright inequality. Indeed, for every $t \in T$, since $v_1(t) = C_{HW} M_0^2 \|B_t\|_{\text{op}}$, $v_2(t) = C_{HW} M_0^2 \|B_t\|_F$, we have that

$$v_1(t) \log \frac{e}{\delta} + v_2(t) \sqrt{\log \frac{e}{\delta}} = C_{HW} M_0^2 \left\{ \|B_t\|_{\text{op}} \log \frac{e}{\delta} + \|B_t\|_F \sqrt{\log \frac{e}{\delta}} \right\}.$$

These are precisely the sub-exponential and sub-Gaussian confidence scalings in the fixed-matrix Hanson-Wright inequality.

5 Pointwise Envelopes for Diffusion Empirical Processes

In this section, we give our second application of the pointwise majorization theorem. We first introduce a stationary scalar diffusion and the occupation empirical process indexed by a class of observables. We then identify its two increment geometries and apply Theorem 3.1.

Let W be a standard Brownian motion and consider

$$dX_s = b(X_s)ds + \sigma(X_s)dW_s, \quad s \geq 0. \quad (5.1)$$

We also use $a(x)$ to denote the square of the function $\sigma(x)$, i.e., $a(x) := \sigma^2(x)$.

Assumption 5.1. The functions $b, \sigma : \mathbb{R} \rightarrow \mathbb{R}$ are globally Lipschitz. There exist $0 < \underline{\sigma} \leq \bar{\sigma} < \infty$, $A > 0$, and $\gamma > 0$ such that $\underline{\sigma} \leq |\sigma(x)| \leq \bar{\sigma}$, if $x \in \mathbb{R}$, and that $\frac{b(x)}{a(x)} \operatorname{sgn}(x) \leq -\gamma$, if $|x| \geq A$.

We define $G(x) := \int_0^x \frac{2b(y)}{a(y)} dy$, $Z := \int_{\mathbb{R}} \frac{e^{G(x)}}{a(x)} dx$, $\pi(x) := \frac{e^{G(x)}}{Za(x)}$, $\ell(x) := a(x)\pi(x) = \frac{e^{G(x)}}{Z}$.

Global Lipschitz continuity implies the local Lipschitz and linear-growth conditions in [Aeckerle-Willems and Strauch \(2021\)](#). Therefore, we know that there is a unique global strong solution with ergodic properties, $0 < Z < \infty$, and the density π defined above is invariant. We start the diffusion in equilibrium, $X_0 \sim \pi$, independently of W , and write $\mathbb{P}_\pi$ and $\mathbb{E}_\pi$ for the resulting stationary law.

Now, we specify the diffusion empirical process that we are going to study. For any Borel bounded function class $\mathcal{F}$ and fixed observation horizon $\tau > 0$, given the realized diffusion process $\{X_s\}_{0 \leq s \leq \tau}$, we denote the empirical occupation measure as $\hat{\pi}_\tau := \frac{1}{\tau} \int_0^\tau \delta_{X_s} ds$.

Then, for any $f \in \mathcal{F}$, we denote

$$\hat{\pi}_\tau(f) := \frac{1}{\tau} \int_0^\tau f(X_s) ds, \quad \pi(f) := \int_{\mathbb{R}} f(x) \pi(x) dx, \quad \bar{f} := f - \pi(f).$$

Intuitively, $\hat{\pi}_\tau(f)$ is the sample path average of f over the trajectory $\{X_s\}_{0 \leq s \leq \tau}$. $\pi(f)$ is the expectation of f under the invariant probability measure, and $\bar{f}$ is its centralized version.

Then, our *diffusion empirical process* $\{Z_\tau(f)\}_{f \in \mathcal{F}}$ indexed by $\mathcal{F}$ is defined as

$$Z_\tau(f) := \sqrt{\tau} (\hat{\pi}_\tau(f) - \pi(f)) = \frac{1}{\sqrt{\tau}} \int_0^\tau \bar{f}(X_s) ds, \quad f \in \mathcal{F}.$$

Throughout this section, τ is fixed, and we apply pointwise majorization to the process $\{Z_\tau(f)\}_{f \in \mathcal{F}}$ indexed by the observable f . In particular, we have that $Z_\tau(0) = 0$, $Z_\tau(f) - Z_\tau(h) = Z_\tau(f - h)$.

Diffusion empirical processes provide a common framework for statistical procedures based on continuous observations of an ergodic diffusion. Indexing the process by indicator functions yields the empirical distribution function, kernel translates yield estimators of the invariant density, and classes of contrast functions arise in M -estimation ([Kutoyants, 1997](#); [van Zanten, 2003](#)). [van der Vaart and van Zanten \(2005\)](#) developed the corresponding Donsker theory, with applications to the empirical distribution function and the local-time density estimator, while [Kutoyants \(2010\)](#) used these objects to construct goodness-of-fit tests.

For finite observation horizons, [Aeckerle-Willems and Strauch \(2021\)](#) established uniform concentration inequalities for diffusion empirical processes and applied them to invariant-density estimation. They identified two natural increment geometries arising from a Poincaré-based Bernstein inequality:

the uniform-norm geometry and the asymptotic-variance geometry.

Building on these results, we provides a simultaneous observable-dependent bound that retains the local complexity of both geometries, rather than replacing them with a single class-wide supremum.

We now define the two scales. Let $0 < C_P < \infty$ be the explicit admissible Poincaré constant constructed in Lemma A.2 such that

$$\text{Var}_\pi(h) \leq \frac{C_P}{2} \int_{\mathbb{R}} a(x) |h'(x)|^2 \pi(x) dx \quad (5.2)$$

for every locally absolutely continuous $h \in L^2(\pi)$ for which the right-hand side is finite.

For a bounded Borel observable f , we define

$$R_f(x) := \int_x^\infty \bar{f}(y) \pi(y) dy = - \int_{-\infty}^x \bar{f}(y) \pi(y) dy, \quad \mathcal{V}_\pi(f)^2 := 4 \int_{\mathbb{R}} \frac{R_f(x)^2}{\ell(x)} dx.$$

The covariance formula of [van der Vaart and van Zanten \(2005\)](#) identifies $\mathcal{V}_\pi(f)^2$ as the scalar central-limit variance. Appendix E checks the normalization and finiteness, proves the seminorm property, and combines that formula with a finite-time Bernstein bound to identify the same quantity as the long-run variance.

Thus, for $f, h \in \mathcal{F}$, we define the following two pseodu-metrics:

$$d_1(f, h) := \frac{C_P}{\sqrt{\tau}} \|(f - h) - \pi(f - h)\|_\infty, \quad d_2(f, h) := \sqrt{2} \mathcal{V}_\pi(f - h),$$

where $\|g\|_\infty := \sup_{x \in \mathbb{R}} |g(x)|$.

Assumption 5.2. The set $\mathcal{F}$ is a standard Borel space whose elements are bounded Borel functions on $\mathbb{R}$. It contains the zero function, and the evaluation map $(f, x) \mapsto f(x)$ is jointly Borel on $\mathcal{F} \times \mathbb{R}$. The maps d_1 and d_2 are jointly Borel, and $(\mathcal{F}, d_1)$ is separable.

Proposition 5.3 (Mixed-tail property of the occupation process). Under Assumptions 5.1 and 5.2, d_1 and d_2 are finite-valued pseudo-metrics and $(\mathcal{F}, d_1 + d_2)$ is separable. The map $(\omega, f) \mapsto Z_\tau(f)(\omega)$ is jointly measurable, its sample paths are almost surely $(d_1 + d_2)$ -continuous, and $Z_\tau(0) = 0$ almost surely. Moreover, for every $f, h \in \mathcal{F}$ and $u \geq 0$,

$$\mathbb{P}_\pi \{|Z_\tau(f) - Z_\tau(h)| > u d_1(f, h) + \sqrt{u} d_2(f, h)\} \leq 2e^{-u}. \quad (5.3)$$

Consequently, $\{Z_\tau(f)\}_{f \in \mathcal{F}}$ is an anchored two-metric mixed-tail process of orders $(\alpha_1, \alpha_2) = (1, 2)$.

Fix Borel probability measures μ_1 and μ_2 on $\mathcal{F}$ before the trajectory is observed. For $j \in \{1, 2\}$, let

$$v_j(f) := d_j(f, 0), \quad B_{d_j}(f, r) := \{h \in \mathcal{F} : d_j(f, h) \leq r\}.$$

Thus, we have $v_1(f) = \frac{C_P}{\sqrt{\tau}} \|f - \pi(f)\|_\infty$, $v_2(f) = \sqrt{2} \mathcal{V}_\pi(f)$. With the convention $\log(1/0) := +\infty$, we define

$$\Phi_1(f) := \int_0^{4v_1(f)} \log \frac{1}{\mu_1(B_{d_1}(f, r))} dr, \quad \Phi_2(f) := \int_0^{4v_2(f)} \left(\log \frac{1}{\mu_2(B_{d_2}(f, r))} \right)^{1/2} dr.$$

Applying Theorem 3.1 to the diffusion empirical process $\{Z_\tau(f)\}_{f \in \mathcal{F}}$, we have the following corollary.

Corollary 5.4 (Simultaneous pointwise occupation bound). Suppose that Assumptions 5.1 and 5.2 hold. Fix $\tau > 0$, start the diffusion from $X_0 \sim \pi$, and fix μ_1, μ_2 before observing the trajectory. There exists a universal constant $C_{1,2} < \infty$ such that, for every $\delta \in (0, 1)$, with probability at least $1 - \delta$, simultaneously for every $f \in \mathcal{F}$,

$$|Z_\tau(f)| \leq C_{1,2} \left\{ \Phi_1(f) + \Phi_2(f) + v_1(f) \log \frac{e}{\delta} + v_2(f) \sqrt{\log \frac{e}{\delta}} \right\}. \quad (5.4)$$

If either local complexity term is infinite at f , the bound at that index is understood to be vacuous.

The terms $\Phi_1(f)$ and $\Phi_2(f)$ measure the local complexity of the observable class in the uniform-amplitude and long-run variance geometries, respectively. The last two terms retain the fixed-observable Bernstein scales $\frac{C_P}{\sqrt{\tau}} \|f - \pi(f)\|_\infty \log \frac{e}{\delta} + \sqrt{2} \mathcal{V}_\pi(f) \sqrt{\log \frac{e}{\delta}}$. Thus, the theorem does not replace the two scales of an individual observable with the worst-case diameters of $\mathcal{F}$.

Finally, notice that $|\hat{\pi}_\tau(f) - \pi(f)| = \frac{|Z_\tau(f)|}{\sqrt{\tau}}$. Hence, Inequality (5.4) also provides a simultaneous bound for all ergodic averages. Because the event is common to every f , it may be evaluated at a measurable trajectory-dependent choice $\hat{f} \in \mathcal{F}$ without another union bound; only the reference measures must have been fixed in advance. All proofs are given in Appendix E.

6 Discussion

In this paper, we develop the first simultaneous pointwise majorization theory for Banach-valued stochastic processes with sub-Weibull increments and with two-metric mixed-tail increments. In the single-metric setting, one high-probability event controls the entire process, while the bound at each index is the sum of a fixed-measure ball-mass integral and a confidence term scaled by the distance from that index to an anchor. This result extends the finite Gaussian upper-envelope construction of Xu (2026) to arbitrary sub-Weibull orders and separable index spaces. In the Gaussian setting, it also sharpens the previous upper bound (Xu, 2026) by removing the auxiliary baseline scales and peeling logarithms. Our two-metric theorem preserves a separate reference measure, local term, anchor radius, and tail exponent for each increment pseudo-metric. The main technical step is to construct measure-generated admissible chains while retaining their cost at every individual index, rather than first replacing that cost by a worst-case supremum. A nested common refinement then makes the two chains telescope along the same process increments without allocating failure probability over pointwise shells. We apply our bound to centered quadratic chaos processes and to diffusion empirical processes. The resulting bounds hold simultaneously over the full index class while adapting, respectively, to the matrix or observable being evaluated. These applications illustrate the main role of our theory: it complements global supremum bounds by preserving heterogeneity across the index set.

Several directions merit further study. First, the simultaneous nature of the bounds makes them potentially useful for the analysis of empirical risk minimization. A common event may be evaluated at a measurable predictor selected from the data without an additional adversarial union bound, provided that the index class, anchor, increment pseudo-metrics, and reference measures are already handled by other means. Turning this observation into excess-risk guarantees will require combining pointwise majorization with appropriate curvature, margin, or localization properties of the risk. It would be particularly interesting to determine whether this approach can complement recent model-free risk evaluation methods based on refitting and perturbation (Wainwright, 2025; Hu and

Simchi-Levi, 2025, 2026; Ni and Huo, 2026) by providing candidate-dependent stochastic certificates.

A second direction concerns the interpretation and choice of the ambient reference measure. The ball-mass term is not a complexity measure of the function class alone: it also depends on the increment pseudo-metric and the reference measure under consideration. Developing principled ways to choose or optimize this measure may lead to useful statistical complexity apart from existing notions such as local Rademacher complexity (Bartlett et al., 2005; Koltchinskii, 2006), VC dimension (Vapnik, 2013), covering numbers (Zhang, 2002) and spectral conditions (Yang et al., 2023; Simon et al., 2023; Hu et al., 2025).

Finally, matching lower bounds remain open. The Gaussian lower bounds of Xu (2026) provide a starting point, but an upper sub-Weibull or mixed-tail increment condition alone cannot imply a nontrivial lower bound, since it is also satisfied by degenerate processes. A meaningful lower-bound theory will therefore require suitable nondegeneracy, small-ball, or two-sided increment assumptions. Under such conditions, one may ask whether the local ball-mass terms, the anchor-dependent confidence scales, and their separation across distinct increment pseudo-metrics are unavoidable.

A Mathematical Tools

We record only the external results used directly in the two applications. The diffusion results are stated directly in the notation of Section 5.

Lemma A.1 (Lamperski (2023)). Let $\zeta = (\zeta_1, \dots, \zeta_n)^\top$ have independent, centered, real-valued coordinates satisfying $\max_i \|\zeta_i\|_{\psi_2} \leq b$, where

$$\|Y\|_{\psi_2} := \inf \left\{ r > 0 : \mathbb{E} \exp \left(\frac{Y^2}{r^2} \right) \leq 2 \right\}.$$

For every nonzero deterministic real matrix A and every $\varepsilon \geq 0$,

$$\mathbb{P} \left\{ \zeta^\top A \zeta - \mathbb{E} [\zeta^\top A \zeta] > \varepsilon \right\} \leq 2 \exp \left\{ -\frac{1}{c_{HW}} \min \left\{ \frac{\varepsilon^2}{b^4 \|A\|_{\mathbb{F}}^2}, \frac{\varepsilon}{b^2 \|A\|_{\text{op}}} \right\} \right\},$$

where $c_{HW} = 2048$ is a universal constant.

Lemma A.2 (Loukianov et al. (2011)). Consider the scalar diffusion in (5.1) under Assumption 5.1, with invariant density π and $\ell = a\pi$. Define

$$F(x) := \int_{-\infty}^x \pi(y) dy, \quad \bar{F}(x) := \int_x^{\infty} \pi(y) dy,$$

and $B_+ := \sup_{x>0} \bar{F}(x) \int_0^x \frac{2}{\ell(y)} dy$, $B_- := \sup_{x<0} F(x) \int_x^0 \frac{2}{\ell(y)} dy$.

If $B_+ \vee B_- < \infty$, then the stationary diffusion is reversible with respect to π , and we have the following Poincare inequality:

$$\text{Var}_\pi(h) \leq \frac{C_P}{2} \int_{\mathbb{R}} a(x) |h'(x)|^2 \pi(x) dx, \quad C_P := 4 \max \{B_+, B_-\}, \quad (\text{A.1})$$

for every locally absolutely continuous $h \in L^2(\pi)$ for which the right-hand side is finite.

Lemma A.3 (Gao et al. (2014)). Let X be the stationary diffusion in Section 5. Suppose that it is reversible with respect to π and that, $\exists 0 < C_P < \infty$ such that

$$\text{Var}_\pi(h) \leq \frac{C_P}{2} \int_{\mathbb{R}} a(x) |h'(x)|^2 \pi(x) dx \quad (\text{A.2})$$

for every locally absolutely continuous $h \in L^2(\pi)$ for which the right-hand side is finite. Let g be bounded and Borel with $\pi(g) = 0$. Then, for every $t, u > 0$,

$$\mathbb{P}_\pi \left\{ \left| \frac{1}{\sqrt{t}} \int_0^t g(X_s) ds \right| > 2\sqrt{C_P} \|g\|_{L^2(\pi)} \sqrt{u} + \frac{C_P \|g\|_\infty}{\sqrt{t}} u \right\} \leq 2e^{-u}. \quad (\text{A.3})$$

If the long-run variance of g equals $\mathcal{V}_\pi(g)^2$, i.e., $\mathcal{V}_\pi(g)^2 = \lim_{t \rightarrow \infty} \frac{1}{t} \text{Var}_{\mathbb{P}_\pi} \left(\int_0^t g(X_s) ds \right)$, then we have that

$$\mathcal{V}_\pi(g)^2 \leq 2C_P \|g\|_{L^2(\pi)}^2, \quad (\text{A.4})$$

and

$$\mathbb{P}_\pi \left\{ \left| \frac{1}{\sqrt{t}} \int_0^t g(X_s) ds \right| > \mathcal{V}_\pi(g) \sqrt{2u} + \frac{C_P \|g\|_\infty}{\sqrt{t}} u \right\} \leq 2e^{-u}. \quad (\text{A.5})$$

The two-sided statements follow by applying the one-sided result of Gao et al. (2014) to g and $-g$.

Lemma A.4 (van der Vaart and van Zanten (2005)). Let X be the stationary scalar diffusion considered in Section 5. Let f be a bounded Borel function such that

$$\mathcal{V}_\pi(f)^2 = 4 \int_{\mathbb{R}} \frac{R_f(x)^2}{\ell(x)} dx < \infty.$$

Then we have that

$$\frac{1}{\sqrt{t}} \int_0^t \bar{f}(X_s) ds \implies \mathcal{N}(0, \mathcal{V}_\pi(f)^2), \quad t \rightarrow \infty.$$

B Proofs in Section 2

Throughout this appendix, we use the conventions

$$\log \frac{1}{0} := +\infty, \quad J_b(+\infty) := +\infty.$$

Changing the process on a null event does not affect any of the claims below. We therefore set $Z_t(\omega) := 0$ for every $t \in T$ whenever $\omega \notin \Omega_0$. Thus, every sample path used below is d -continuous and satisfies $Z_{t_0} = 0$.

Lemma B.1. The map $(t, r) \mapsto \mu(B_d(t, r))$ is Borel measurable on $T \times [0, \infty)$. Consequently, v and $\Phi_{\mu, d}^{(\alpha)}$ are $[0, \infty]$ -valued Borel functions. Moreover, for every $r_0, s_0 > 0$, the maps

$$t \mapsto J_{r_0} \left(\Phi_{\mu, d}^{(\alpha)}(t) \right), \quad t \mapsto J_{s_0}(v(t))$$

are Borel measurable as extended-valued maps.

Proof of Lemma B.1. By the joint Borel measurability of d , the map $(t, r, s) \mapsto \mathbf{1}_{\{d(t,s) \leq r\}}$ is Borel measurable. Therefore, by parameterized integration, we have that

$$(t, r) \mapsto \int_T \mathbf{1}_{\{d(t,s) \leq r\}} \mu(ds) = \mu(B_d(t, r))$$

is Borel measurable. The map $v(t) = d(t, t_0)$ is also Borel measurable. We define

$$G(t, r) := \begin{cases} \left(\log \frac{1}{\mu(B_d(t, r))} \right)^{1/\alpha}, & 0 \leq r \leq 4v(t), \\ 0, & r > 4v(t). \end{cases}$$

Then G is a nonnegative, extended-valued Borel function and $\Phi_{\mu, d}^{(\alpha)}(t) = \int_0^\infty G(t, r) dr$. Another application of parameterized integration proves the measurability of $\Phi_{\mu, d}^{(\alpha)}$. The remaining claims follow by composition with the Borel function J_b . $\square$

The next lemma is a pointwise version of the usual finite-set chaining bound. The essential difference is that we do not replace the chaining cost at t by its maximum over K .

Lemma B.2. Suppose that $K \subset T$ is finite and $t_0 \in K$. Let $\mathcal{T} = (T_n)_{n \geq 0}$ be an admissible sequence of nonempty subsets of K such that $T_0 = \{t_0\}$ and $T_n = K$ for all sufficiently large n . Define the pointwise chaining cost

$$G_{\mathcal{T}}(t) := \sum_{n \geq 0} 2^{n/\alpha} d(t, T_n), \quad t \in K.$$

Then, there exists a finite constant A_α , depending only on α , such that, for every $u \geq 1$, with probability at least $1 - e^{-u}$, simultaneously for every $t \in K$, we have

$$\|Z_t\| \leq A_\alpha \left[G_{\mathcal{T}}(t) + u^{1/\alpha} v(t) \right].$$

Proof of Lemma B.2. For every $t \in K$ and every $n \geq 0$, choose $\pi_n(t) \in T_n$ such that

$$d(t, \pi_n(t)) = d(t, T_n).$$

Whenever $T_n = K$, we take $\pi_n(t) = t$. In particular, for every $t \in K$, the sequence $\pi_n(t)$ is eventually equal to t .

Fix $u \geq 1$ and let $\ell := \lfloor \log_2 u \rfloor$. Then, $2^\ell \leq u < 2^{\ell+1}$. We first control the coarse increments between T_ℓ and the anchor. By the sub-Weibull increment assumption and a union bound, we have

$$\mathbb{P} \left\{ \exists s \in T_\ell : \|Z_s - Z_{t_0}\| > (16u)^{1/\alpha} d(s, t_0) \right\} \leq 2|T_\ell| e^{-16u} \leq 2 \exp \left\{ (\log 2) 2^\ell - 16u \right\} \leq \frac{1}{2} e^{-u}.$$

For every $n > \ell$, we define the collection of distinct parent-child edges

$$E_n := \{(\pi_n(t), \pi_{n-1}(t)) : t \in K\}.$$

By admissibility, we obtain that

$$|E_n| \leq |T_n| |T_{n-1}| \leq 2^{2^n + 2^{n-1}} \leq \exp \left\{ 2^{n+1} \right\}.$$

Applying the increment assumption and another union bound over E_n implies

$$\mathbb{P} \left\{ \exists (s, r) \in E_n : \|Z_s - Z_r\| > (16 \cdot 2^n)^{1/\alpha} d(s, r) \right\} \leq 2 \exp \left\{ 2^{n+1} - 16 \cdot 2^n \right\} = 2e^{-14 \cdot 2^n}.$$

Moreover, since $2^{\ell+1} > u$, by some algebra, we have that

$$2 \sum_{n>\ell} e^{-14 \cdot 2^n} \leq \frac{2e^{-14 \cdot 2^{\ell+1}}}{1 - e^{-14 \cdot 2^{\ell+1}}} \leq \frac{1}{2} e^{-u}.$$

Combining the coarse and fine estimates, we obtain an event $\mathcal{E}_u$ such that $\mathbb{P}(\mathcal{E}_u) \geq 1 - e^{-u}$ and all the preceding increment inequalities hold simultaneously on $\mathcal{E}_u$.

We now fix $t \in K$ and work conditioning on $\mathcal{E}_u$. Since $\pi_n(t) = t$ for all sufficiently large n and $Z_{t_0} = 0$, we may telescope and obtain

$$\begin{aligned} \|Z_t\| &\leq \|Z_{\pi_\ell(t)} - Z_{t_0}\| + \sum_{n>\ell} \|Z_{\pi_n(t)} - Z_{\pi_{n-1}(t)}\| \\ &\leq (16u)^{1/\alpha} d(\pi_\ell(t), t_0) + 16^{1/\alpha} \sum_{n>\ell} 2^{n/\alpha} d(\pi_n(t), \pi_{n-1}(t)). \end{aligned}$$

For the coarse increment, by the triangle inequality, we have

$$d(\pi_\ell(t), t_0) \leq d(\pi_\ell(t), t) + d(t, t_0) = d(t, T_\ell) + v(t).$$

Consequently, using $u < 2^{\ell+1}$, we have that

$$\begin{aligned} (16u)^{1/\alpha} d(\pi_\ell(t), t_0) &\leq 16^{1/\alpha} u^{1/\alpha} v(t) + 16^{1/\alpha} u^{1/\alpha} d(t, T_\ell) \\ &\leq 16^{1/\alpha} u^{1/\alpha} v(t) + 32^{1/\alpha} 2^{\ell/\alpha} d(t, T_\ell) \\ &\leq 16^{1/\alpha} u^{1/\alpha} v(t) + 32^{1/\alpha} G_{\mathcal{T}}(t). \end{aligned}$$

For the fine increments, another application of the triangle inequality gives

$$\begin{aligned} \sum_{n>\ell} 2^{n/\alpha} d(\pi_n(t), \pi_{n-1}(t)) &\leq \sum_{n>\ell} 2^{n/\alpha} [d(t, T_n) + d(t, T_{n-1})] \\ &\leq (1 + 2^{1/\alpha}) \sum_{m \geq 0} 2^{m/\alpha} d(t, T_m) = (1 + 2^{1/\alpha}) G_{\mathcal{T}}(t). \end{aligned}$$

Combining the coarse and fine estimates, we obtain that

$$\|Z_t\| \leq 16^{1/\alpha} u^{1/\alpha} v(t) + \left[32^{1/\alpha} + 16^{1/\alpha} (1 + 2^{1/\alpha}) \right] G_{\mathcal{T}}(t).$$

Therefore, the assertion holds with

$$A_\alpha := \max \left\{ 16^{1/\alpha}, 32^{1/\alpha} + 16^{1/\alpha} (1 + 2^{1/\alpha}) \right\}.$$

Since the event $\mathcal{E}_u$ is common to every $t \in K$, the inequality holds simultaneously for all $t \in K$. $\square$

We next construct an admissible sequence from a probability measure on a finite pseudo-metric space. Unlike the usual bound, the following lemma retains the resulting chaining cost at each

individual index.

Lemma B.3. Let $K \subset T$ be finite and contain t_0 , let $d_K := d|_{K \times K}$, and let ν be a probability measure on K such that $\nu(\{t_0\}) \geq \frac{1}{2}$. For $t \in K$ and $n \geq 0$, define

$$r_n(t) := \min \left\{ r \in \{d_K(t, s) : s \in K\} : \nu(B_{d_K}(t, r)) \geq 2^{-2^n} \right\}.$$

Then, there exists an admissible sequence $\mathcal{T} = (T_n)_{n \geq 0}$ of subsets of K such that $T_0 = \{t_0\}$, $T_n = K$ for all sufficiently large n , and, for every $t \in K$,

$$G_{\mathcal{T}}(t) := \sum_{n \geq 0} 2^{n/\alpha} d_K(t, T_n) \leq v(t) + \kappa_{\alpha} \int_0^{r_0(t)} \left(\log \frac{1}{\nu(B_{d_K}(t, r))} \right)^{1/\alpha} dr,$$

where

$$\kappa_{\alpha} := \frac{2^{1+1/\alpha}}{(\log 2)^{1/\alpha} (1 - 2^{-1/\alpha})}.$$

Moreover, $r_0(t) \leq v(t)$ for every $t \in K$.

Proof of Lemma B.3. The minimum defining $r_n(t)$ exists because K is finite and the largest d_K -ball centered at t has ν -mass one. Moreover, the sequence $r_n(t)$ is nonincreasing in n .

For a fixed $n \geq 0$, we order the points of K by nondecreasing order of $\{r_n(t)\}_{t \in K}$, breaking ties according to an arbitrary fixed enumeration of K . We traverse this ordered list and retain a point t precisely when

$$d_K(t, s) > r_n(t) + r_n(s)$$

for every previously retained point s , and denote the set of retained points by S_n .

We first show that the balls $\{B_{d_K}(s, r_n(s)) : s \in S_n\}$ are pairwise disjoint. Indeed, suppose that two such balls, centered at $s, t \in S_n$, intersect. The triangle inequality would then give

$$d_K(s, t) \leq r_n(s) + r_n(t).$$

On the other hand, because both points were retained, the greedy rule gives

$$d_K(s, t) > r_n(s) + r_n(t),$$

which is a contradiction.

By the definition of $r_n(s)$, every one of these disjoint balls has ν -mass at least 2^{-2^n} . We have

$$1 \geq \sum_{s \in S_n} \nu(B_{d_K}(s, r_n(s))) \geq |S_n| 2^{-2^n}.$$

Consequently, we obtain $|S_n| \leq 2^{2^n}$.

We next prove that S_n approximates every point at its own mass radius. If $t \in S_n$, then $d_K(t, S_n) = 0$. If $t \notin S_n$, the point t was rejected because there exists a previously retained point $s \in S_n$ such that

$$r_n(s) \leq r_n(t), \quad d_K(t, s) \leq r_n(t) + r_n(s).$$

Therefore, in both cases, we have

$$d_K(t, S_n) \leq 2r_n(t), \quad t \in K.$$

For $t \in K$, we define $F_t(r) := \left(\log \frac{1}{\nu(B_{d_K}(t, r))} \right)^{1/\alpha}$. If $r < r_n(t)$, by the definition of $r_n(t)$, we have that $\nu(B_{d_K}(t, r)) < 2^{-2^n}$. It follows that $F_t(r) \geq (\log 2)^{1/\alpha} 2^{n/\alpha}$.

Since the intervals $[r_{n+1}(t), r_n(t))$, $n \geq 0$, are disjoint and contained in $[0, r_0(t)]$, we have

$$\begin{aligned} \int_0^{r_0(t)} F_t(r) dr &= \sum_n \int_{r_{n+1}(t)}^{r_n(t)} \log\left(\frac{1}{\nu(B_d(t, r))}\right)^{1/\alpha} dr \\ &\geq (\log 2)^{1/\alpha} \sum_{n \geq 0} 2^{n/\alpha} [r_n(t) - r_{n+1}(t)] \\ &\geq (\log 2)^{1/\alpha} \left(1 - 2^{-1/\alpha}\right) \sum_{n \geq 0} 2^{n/\alpha} r_n(t). \end{aligned}$$

We justify the last inequality. If the integral on the left-hand side is infinite, there is nothing to prove. If it is finite, then the d_K -equivalence class of t must have positive ν -mass; otherwise, because K is finite, F_t would be infinite on an interval of positive length. Hence, $r_n(t) = 0$ for all sufficiently large n . Summation by parts then gives

$$\sum_{n \geq 0} 2^{n/\alpha} [r_n(t) - r_{n+1}(t)] = \left(1 - 2^{-1/\alpha}\right) \sum_{n \geq 0} 2^{n/\alpha} r_n(t) + 2^{-1/\alpha} r_0(t) \geq \left(1 - 2^{-1/\alpha}\right) \sum_{n \geq 0} 2^{n/\alpha} r_n(t).$$

We now define the admissible sequence. Let $T_0 := \{t_0\}$, $T_n := S_{n-1}$, $n \geq 1$. Since K is finite, choose $n_K \geq 1$ such that $2^{2^{n_K}} \geq |K|$. We replace T_n by K for every $n \geq n_K$. This replacement can only decrease the distances to T_n , it preserves admissibility, and it ensures that $T_n = K$ for all sufficiently large n . By the approximation property of S_n , for every $t \in K$, we have

$$\begin{aligned} G_{\mathcal{T}}(t) &= d_K(t, t_0) + \sum_{n \geq 1} 2^{n/\alpha} d_K(t, T_n) \\ &\leq v(t) + 2 \sum_{n \geq 1} 2^{n/\alpha} r_{n-1}(t) \\ &= v(t) + 2^{1+1/\alpha} \sum_{m \geq 0} 2^{m/\alpha} r_m(t) \\ &\leq v(t) + \frac{2^{1+1/\alpha}}{(\log 2)^{1/\alpha} (1 - 2^{-1/\alpha})} \int_0^{r_0(t)} F_t(r) dr. \end{aligned}$$

This proves the first assertion.

Finally, the closed ball $B_{d_K}(t, v(t))$ contains the anchor t_0 . Since $\nu(\{t_0\}) \geq 1/2$, we have $\nu(B_{d_K}(t, v(t))) \geq \frac{1}{2} = 2^{-2^0}$. By the definition of $r_0(t)$, this implies that $r_0(t) \leq v(t)$.

We finish the proof. $\square$

The following lemma transfers the fixed ambient measure μ to an arbitrary finite set while retaining the pointwise ball masses.

Lemma B.4. Let $K \subseteq T$ be finite and contain t_0 , and let $d_K := d|_{K \times K}$. There exists a probability

measure $\tilde{\mu}_K$ on K such that, for every $t \in K$ and $r \geq 0$,

$$\tilde{\mu}_K(\{t_0\}) \geq \frac{1}{2}, \quad \tilde{\mu}_K(B_{d_K}(t, 2r)) \geq \frac{1}{2}\mu(B_d(t, r)).$$

Moreover, with $c_\alpha^{(0)} := \max\{1, 2^{1/\alpha-1}\}$, we have that

$$\int_0^{4v(t)} \left(\log \frac{1}{\tilde{\mu}_K(B_{d_K}(t, r))} \right)^{1/\alpha} dr \leq 2c_\alpha^{(0)}\Phi_{\mu,d}^{(\alpha)}(t) + 4c_\alpha^{(0)}(\log 2)^{1/\alpha}v(t).$$

Proof of Lemma B.4. Enumerate $K = \{k_1, \dots, k_N\}$ and let $p_K : T \rightarrow K$ be the nearest-point map, breaking ties by the smallest index. More precisely, the cell on which $p_K = k_j$ is

$$\bigcap_{i < j} \{s : d(s, k_j) < d(s, k_i)\} \cap \bigcap_{i > j} \{s : d(s, k_j) \leq d(s, k_i)\}.$$

It is Borel by the joint measurability of d . Therefore, the map p_K is Borel measurable. We define

$$\tilde{\mu}_K := \frac{1}{2}\delta_{t_0} + \frac{1}{2}(p_K)_\# \mu,$$

where $(p_K)_\# \mu$ is the pushforward of μ under p_K ; that is,

$$(p_K)_\# \mu(A) := \mu(p_K^{-1}(A)), \quad A \subseteq K.$$

It follows immediately that $\tilde{\mu}_K(\{t_0\}) \geq \frac{1}{2}$.

We next prove the ball-mass comparison. Fix $t \in K$ and suppose that $s \in B_d(t, r)$. By the nearest-point property of p_K , we have

$$d(s, p_K(s)) \leq d(s, t) \leq r.$$

Applying the triangle inequality and we obtain that

$$d(t, p_K(s)) \leq d(t, s) + d(s, p_K(s)) \leq 2r.$$

It implies that $B_d(t, r) \subseteq p_K^{-1}(B_{d_K}(t, 2r))$. Using the definition of $\tilde{\mu}_K$, we obtain

$$\tilde{\mu}_K(B_{d_K}(t, 2r)) \geq \frac{1}{2}(p_K)_\# \mu(B_{d_K}(t, 2r)) = \frac{1}{2}\mu(p_K^{-1}(B_{d_K}(t, 2r))) \geq \frac{1}{2}\mu(B_d(t, r)).$$

Applying the preceding inequality with radius $r/2$, we have

$$\log \frac{1}{\tilde{\mu}_K(B_{d_K}(t, r))} \leq \log 2 + \log \frac{1}{\mu(B_d(t, r/2))}.$$

For $x, y \geq 0$, by direct algebra, we obtain

$$(x + y)^{1/\alpha} \leq c_\alpha^{(0)} (x^{1/\alpha} + y^{1/\alpha}).$$

Therefore, we have

$$\int_0^{4v(t)} \left(\log \frac{1}{\tilde{\mu}_K(B_{d_K}(t, r))} \right)^{1/\alpha} dr \leq 4c_\alpha^{(0)} (\log 2)^{1/\alpha} v(t) + c_\alpha^{(0)} \int_0^{4v(t)} \left(\log \frac{1}{\mu(B_d(t, r/2))} \right)^{1/\alpha} dr.$$

By the change of variables $r = 2u$, we obtain

$$\begin{aligned} \int_0^{4v(t)} \left(\log \frac{1}{\tilde{\mu}_K(B_{d_K}(t, r))} \right)^{1/\alpha} dr &\leq 4c_\alpha^{(0)} (\log 2)^{1/\alpha} v(t) + 2c_\alpha^{(0)} \int_0^{2v(t)} \left(\log \frac{1}{\mu(B_d(t, u))} \right)^{1/\alpha} du \\ &\leq 4c_\alpha^{(0)} (\log 2)^{1/\alpha} v(t) + 2c_\alpha^{(0)} \Phi_{\mu, d}^{(\alpha)}(t). \end{aligned}$$

This proves the claimed integral inequality. $\square$

We now combine the pointwise tail estimate, the measure-generated chain, and the ambient restriction. The resulting finite-set estimate already has the same pointwise right-hand side as the main theorem.

Lemma B.5. Suppose that $K \subset T$ is finite and $t_0 \in K$. There exists a finite constant B_α , depending only on α , such that, for every $u \geq 1$, with probability at least $1 - e^{-u}$, simultaneously for every $t \in K$, we have

$$\|Z_t\| \leq B_\alpha \left[\Phi_{\mu, d}^{(\alpha)}(t) + u^{1/\alpha} v(t) \right].$$

Proof of Lemma B.5. Let $d_K := d|_{K \times K}$ and let $\tilde{\mu}_K$ be the probability measure obtained from Lemma B.4. In particular, we have

$$\tilde{\mu}_K(\{t_0\}) \geq \frac{1}{2}.$$

Therefore, Lemma B.3 gives an admissible sequence $\mathcal{T} = (T_n)_{n \geq 0}$, anchored at t_0 and eventually equal to K , such that, for every $t \in K$,

$$G_{\mathcal{T}}(t) \leq v(t) + \kappa_\alpha \int_0^{r_0(t)} \left(\log \frac{1}{\tilde{\mu}_K(B_{d_K}(t, r))} \right)^{1/\alpha} dr,$$

where $r_0(t) \leq v(t)$. Since $v(t) \leq 4v(t)$ and the integrand is nonnegative, we obtain

$$G_{\mathcal{T}}(t) \leq v(t) + \kappa_\alpha \int_0^{4v(t)} \left(\log \frac{1}{\tilde{\mu}_K(B_{d_K}(t, r))} \right)^{1/\alpha} dr.$$

Using Lemma B.4, it follows that

$$G_{\mathcal{T}}(t) \leq v(t) + 2\kappa_\alpha c_\alpha^{(0)} \Phi_{\mu, d}^{(\alpha)}(t) + 4\kappa_\alpha c_\alpha^{(0)} (\log 2)^{1/\alpha} v(t) \leq b_\alpha \left[\Phi_{\mu, d}^{(\alpha)}(t) + v(t) \right],$$

where

$$b_\alpha := \max \left\{ 2\kappa_\alpha c_\alpha^{(0)}, 1 + 4\kappa_\alpha c_\alpha^{(0)} (\log 2)^{1/\alpha} \right\}.$$

We apply Lemma B.2 to this admissible sequence. Outside an event of probability at most e^{-u} ,

simultaneously for every $t \in K$, we have

$$\|Z_t\| \leq A_\alpha \left[G_{\mathcal{T}}(t) + u^{1/\alpha} v(t) \right] \leq A_\alpha b_\alpha \Phi_{\mu,d}^{(\alpha)}(t) + A_\alpha (b_\alpha + u^{1/\alpha}) v(t).$$

Since $u \geq 1$, we have $u^{1/\alpha} \geq 1$, and hence $b_\alpha + u^{1/\alpha} \leq (b_\alpha + 1)u^{1/\alpha}$. Consequently, we obtain that with probability at least $1 - e^{-u}$,

$$\|Z_t\| \leq B_\alpha \left[\Phi_{\mu,d}^{(\alpha)}(t) + u^{1/\alpha} v(t) \right],$$

where $B_\alpha := A_\alpha \max \{b_\alpha, b_\alpha + 1\}$.

The proof is complete. $\square$

We are now ready to pass from finite sets to the full separable index space and prove Theorem 2.1. The passage uses countable dense subsets of rational upper-level sets. It does not allocate separate failure probabilities to these sets.

Proof of Theorem 2.1. Fix $u \geq 1$. For positive rational numbers $q, s \in \mathbb{Q}_{>0}$, define

$$H_{q,s} := \left\{ t \in T : \Phi_{\mu,d}^{(\alpha)}(t) \leq q, \quad v(t) \leq s \right\}.$$

By Lemma B.1, each $H_{q,s}$ is a Borel subset of T . More importantly, every $H_{q,s}$ is separable with respect to d . Indeed, the separable space (T, d) has a separable metric quotient, which is second countable; every subspace of a second countable space is second countable and hence separable.

For every pair $(q, s) \in \mathbb{Q}_{>0}^2$ such that $H_{q,s}$ is nonempty, choose a countable d -dense subset $D_{q,s}$ of $H_{q,s}$. The set $D := \{t_0\} \cup \bigcup_{(q,s) \in \mathbb{Q}_{>0}^2} D_{q,s}$ is countable. We enumerate its elements as

$$D = \{t_0, t_1, t_2, \dots\}$$

where, if D is finite, we repeat t_0 so that the enumeration is an infinite sequence. We then define the increasing sequence of finite sets

$$K_N := \{t_0, t_1, \dots, t_N\}, \quad N \geq 1.$$

For every $N \geq 1$, let E_N be the event on which

$$\|Z_t\| \leq B_\alpha \left[\Phi_{\mu,d}^{(\alpha)}(t) + u^{1/\alpha} v(t) \right]$$

holds for every $t \in K_N$. By Lemma B.5, we have

$$\mathbb{P}(E_N) \geq 1 - e^{-u}, \quad N \geq 1.$$

Since $K_N \subseteq K_{N+1}$ and the deterministic right-hand side does not depend on N , we have that

$$E_{N+1} \subseteq E_N.$$

Therefore, by the continuity of probability, we have

$$\mathbb{P} \left(\bigcap_{N \geq 1} E_N \right) = \lim_{N \rightarrow \infty} \mathbb{P}(E_N) \geq 1 - e^{-u}.$$

We intersect this event $\bigcap_N E_N$ with the probability-one event Ω_0 on which $t \mapsto Z_t$ is d -continuous and $Z_{t_0} = 0$. Now we condition on the event $\Omega_0 \cap \bigcap_N E_N$.

Fix $t \in T$ such that $\Phi_{\mu,d}^{(\alpha)}(t) < \infty$. Since d is finite-valued, $v(t) = d(t, t_0) < \infty$. Choose arbitrary positive rational numbers q and s satisfying

$$q > \Phi_{\mu,d}^{(\alpha)}(t), \quad s > v(t).$$

Then $t \in H_{q,s}$. Since $D_{q,s}$ is d -dense in $H_{q,s}$, there exists a sequence $(t_m)_{m \geq 1} \subseteq D_{q,s}$ such that $d(t_m, t) \rightarrow 0$. Every t_m belongs to the countable set D , and hence the inequality on $\bigcap_{N \geq 1} E_N$ gives

$$\|Z_{t_m}\| \leq B_\alpha \left[\Phi_{\mu,d}^{(\alpha)}(t_m) + u^{1/\alpha} v(t_m) \right] \leq B_\alpha \left[q + u^{1/\alpha} s \right].$$

By the path continuity of the process conditioned on Ω_0 , we have that $\|Z_t\| = \lim_{m \rightarrow \infty} \|Z_{t_m}\|$. It follows that

$$\|Z_t\| \leq B_\alpha \left[q + u^{1/\alpha} s \right].$$

Since q and s are arbitrary positive rationals strictly larger than $\Phi_{\mu,d}^{(\alpha)}(t)$ and $v(t)$, respectively, we may let

$$q \downarrow \Phi_{\mu,d}^{(\alpha)}(t), \quad s \downarrow v(t)$$

through rational values. Therefore, we obtain

$$\|Z_t\| \leq B_\alpha \left[\Phi_{\mu,d}^{(\alpha)}(t) + u^{1/\alpha} v(t) \right].$$

The assertion is vacuous at every point at which $\Phi_{\mu,d}^{(\alpha)}(t) = +\infty$. Consequently, with probability at least $1 - e^{-u}$, the preceding inequality holds simultaneously for every $t \in T$.

Finally, let $\delta \in (0, 1)$ and take $u := \log \frac{e}{\delta}$. Then we know that $u \geq 1$ and $e^{-u} = \frac{\delta}{e} \leq \delta$. Therefore, with probability at least $1 - \delta$, simultaneously for every $t \in T$, we have

$$\|Z_t\| \leq B_\alpha \left[\Phi_{\mu,d}^{(\alpha)}(t) + v(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha} \right].$$

The theorem follows with $C_\alpha := B_\alpha$. □

C Proofs in Section 3

Throughout this appendix, we use the convention $\log \frac{1}{0} := +\infty$. Changing the process on a null event does not affect any of the claims below. We therefore redefine

$$Z_t(\omega) := 0, \quad \omega \notin \Omega_0, t \in T,$$

Thus, every sample path used below is ρ -continuous and satisfies $Z_{t_0} = 0$.

We first verify the measurability of the pointwise radii and profiles.

Lemma C.1. For every $j \in \{1, 2\}$, the map

$$(t, r) \mapsto \mu_j(B_{d_j}(t, r))$$

is Borel measurable on $T \times [0, \infty)$. Consequently, v_j and Φ_j are $[0, \infty]$ -valued Borel functions on T .

Proof of Lemma C.1. Fix $j \in \{1, 2\}$. Since d_j is jointly $\mathcal{A} \otimes \mathcal{A}$ -measurable, the map

$$(t, r, s) \mapsto \mathbf{1}_{\{d_j(t, s) \leq r\}}$$

is measurable from $(T \times [0, \infty) \times T, \mathcal{A} \otimes \mathcal{B}([0, \infty)) \otimes \mathcal{A})$ to $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$. Therefore, by measurability of parameterized integrals, the map $(t, r) \mapsto \int_T \mathbf{1}_{\{d_j(t, s) \leq r\}} \mu_j(ds) = \mu_j(B_{d_j}(t, r))$ is Borel measurable. Moreover, since t_0 is fixed and d_j is jointly Borel, the map $v_j(t) = d_j(t, t_0)$ is Borel measurable. Again, we define

$$Q_j(t, r) := \begin{cases} \left(\log \frac{1}{\mu_j(B_{d_j}(t, r))} \right)^{1/\alpha_j}, & 0 \leq r \leq 4v_j(t), \\ 0, & r > 4v_j(t). \end{cases}$$

The preceding measurability statements show that Q_j is a nonnegative, extended-real-valued Borel function. Notice that the piecewise definition avoids the ambiguous product $0 \cdot \infty$ when a ball has zero μ_j -mass. By the definition of Φ_j , we have that $\Phi_j(t) = \int_0^\infty Q_j(t, r) dr$. Another application of measurability of parameterized integrals proves that Φ_j is Borel measurable. Since the same argument applies to $j = 1$ and $j = 2$, we finish the proof. $\square$

We next transfer each fixed ambient measure to an arbitrary finite subset. This is the nearest-point construction that permits the measure-generated chains to be built inside the finite set without losing the ambient ball masses.

Lemma C.2. Let $K \subseteq T$ be finite and contain t_0 . For $j \in \{1, 2\}$, write $d_{j,K} := d_j|_{K \times K}$. There exists a probability measure $\tilde{\mu}_{j,K}$ on K such that, for every $t \in K$ and every $r \geq 0$,

$$\tilde{\mu}_{j,K}(\{t_0\}) \geq \frac{1}{2}, \quad \tilde{\mu}_{j,K}(B_{d_{j,K}}(t, 2r)) \geq \frac{1}{2} \mu_j(B_{d_j}(t, r)).$$

Moreover, if $c_{\alpha_j}^{(0)} := \max \{1, 2^{1/\alpha_j - 1}\}$, then

$$\int_0^{4v_j(t)} \left(\log \frac{1}{\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, r))} \right)^{1/\alpha_j} dr \leq 2c_{\alpha_j}^{(0)} \Phi_j(t) + 4c_{\alpha_j}^{(0)} (\log 2)^{1/\alpha_j} v_j(t).$$

The proof of Lemma C.2 is very similar to the proof of Lemma B.4.

Proof of Lemma C.2. Fix $j \in \{1, 2\}$ and enumerate

$$K = \{k_1, \dots, k_N\}.$$

For every $s \in T$, let $q_{j,K}(s)$ be a point in K minimizing $d_j(s, \cdot)$, and break ties by choosing the point with the smallest index. More explicitly, define

$$C_i := \bigcap_{m < i} \{s \in T : d_j(s, k_i) < d_j(s, k_m)\} \cap \bigcap_{m > i} \{s \in T : d_j(s, k_i) \leq d_j(s, k_m)\}.$$

Every set C_i is Borel measurable because d_j is jointly Borel. The sets $C_1, \dots, C_N$ form a partition of T , and $q_{j,K}(s) = k_i$ for $s \in C_i$. Hence, $q_{j,K} : T \rightarrow K$ is Borel measurable and satisfies

$$d_j(s, q_{j,K}(s)) = d_j(s, K) := \min_{k \in K} d_j(s, k).$$

Define

$$\tilde{\mu}_{j,K} := \frac{1}{2}\delta_{t_0} + \frac{1}{2}(q_{j,K})_{\#}\mu_j,$$

where the pushforward measure is defined by

$$(q_{j,K})_{\#}\mu_j(A) := \mu_j(q_{j,K}^{-1}(A)), \quad A \subseteq K.$$

By construction, we have

$$\tilde{\mu}_{j,K}(\{t_0\}) \geq \frac{1}{2}.$$

We now prove the ball-mass comparison. Fix $t \in K$ and $r \geq 0$. If $s \in B_{d_j}(t, r)$, then $t \in K$ and the nearest-point property give

$$d_j(s, q_{j,K}(s)) \leq d_j(s, t) \leq r.$$

By the triangle inequality, we have

$$d_j(t, q_{j,K}(s)) \leq d_j(t, s) + d_j(s, q_{j,K}(s)) \leq 2r.$$

Therefore, we get $q_{j,K}(B_{d_j}(t, r)) \subseteq B_{d_{j,K}}(t, 2r)$. It follows that

$$\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, 2r)) \geq \frac{1}{2}(q_{j,K})_{\#}\mu_j(B_{d_{j,K}}(t, 2r)) = \frac{1}{2}\mu_j(q_{j,K}^{-1}(B_{d_{j,K}}(t, 2r))) \geq \frac{1}{2}\mu_j(B_{d_j}(t, r)).$$

It remains to prove the integral comparison. Recall that $p_j := \frac{1}{\alpha_j}$.

By the change of variable $r = 2u$, we have

$$\int_0^{4v_j(t)} \left(\log \frac{1}{\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, r))} \right)^{p_j} dr = 2 \int_0^{2v_j(t)} \left(\log \frac{1}{\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, 2u))} \right)^{p_j} du.$$

The ball-mass comparison gives us

$$\log \frac{1}{\tilde{\mu}_{j,K}(B_{d_j,K}(t, 2u))} \leq \log \frac{2}{\mu_j(B_{d_j}(t, u))} = \log \frac{1}{\mu_j(B_{d_j}(t, u))} + \log 2.$$

For nonnegative x, y , we have

$$(x + y)^{p_j} \leq c_{\alpha_j}^{(0)} (x^{p_j} + y^{p_j}).$$

Finally, combining these parts together, we obtain

$$\begin{aligned} \int_0^{4v_j(t)} \left(\log \frac{1}{\tilde{\mu}_{j,K}(B_{d_j,K}(t, r))} \right)^{p_j} dr &\leq 2c_{\alpha_j}^{(0)} \int_0^{2v_j(t)} \left(\log \frac{1}{\mu_j(B_{d_j}(t, u))} \right)^{p_j} du + 2c_{\alpha_j}^{(0)} \int_0^{2v_j(t)} (\log 2)^{p_j} du \\ &\leq 2c_{\alpha_j}^{(0)} \Phi_j(t) + 4c_{\alpha_j}^{(0)} (\log 2)^{p_j} v_j(t), \end{aligned}$$

where we used $2v_j(t) \leq 4v_j(t)$ in the final inequality. This proves the assertion for $j = 1, 2$. Applying the same construction to both metrics finishes the proof. $\square$

We now construct, for each metric, an admissible sequence whose chaining cost remains pointwise rather than being replaced by a worst-case supremum over K .

Lemma C.3. Let $K \subseteq T$ be finite and contain t_0 . For every $j \in \{1, 2\}$, there exists an admissible sequence $\mathcal{T}_j = (T_{j,k})_{k \geq 0}$ of subsets of K such that $T_{j,0} = \{t_0\}$ and $T_{j,k} = K$ for all sufficiently large k . Moreover, for every $t \in K$,

$$G_{j,K}(t) := \sum_{k \geq 0} 2^{k/\alpha_j} d_j(t, T_{j,k}) \leq C_{\alpha_j}^{(1)} (\Phi_j(t) + v_j(t)),$$

where $C_{\alpha_j}^{(1)} < \infty$ depends only on α_j .

Proof of Lemma C.3. Fix $j \in \{1, 2\}$ and write $p_j := \frac{1}{\alpha_j}$. Let $\tilde{\mu}_{j,K}$ be the probability measure supplied by Lemma C.2. For $t \in K$ and $n \geq 0$, define

$$r_{j,n}(t) := \min \left\{ r \in \{d_j(t, s) : s \in K\} : \tilde{\mu}_{j,K}(B_{d_j,K}(t, r)) \geq 2^{-2^n} \right\}.$$

The minimum exists because K is finite and the largest $d_{j,K}$ -ball centered at t has $\tilde{\mu}_{j,K}$ -mass one. Since the mass threshold 2^{-2^n} is nonincreasing in n , the sequence $(r_{j,n}(t))_{n \geq 0}$ is nonincreasing.

For a fixed $n \geq 0$, similar to the proof of Lemma B.3, we order the points of K by nondecreasing $r_{j,n}(t)$ and break ties according to an arbitrary fixed enumeration of K . Traverse the ordered list and retain a point t precisely when

$$d_j(t, s) > r_{j,n}(t) + r_{j,n}(s)$$

for every previously retained point s . Denote the set of retained points by $S_{j,n}$.

We first prove that the balls $\{B_{d_j,K}(s, r_{j,n}(s)) : s \in S_{j,n}\}$ are pairwise disjoint. Suppose that two such balls, centered at distinct points $s, t \in S_{j,n}$, intersect. Then there exists $z \in K$ such that

$$d_j(s, z) \leq r_{j,n}(s), \quad d_j(t, z) \leq r_{j,n}(t).$$

Applying the triangle inequality would gives us

$$d_j(s, t) \leq d_j(s, z) + d_j(z, t) \leq r_{j,n}(s) + r_{j,n}(t).$$

On the other hand, because both s and t were retained, the greedy rule gives

$$d_j(s, t) > r_{j,n}(s) + r_{j,n}(t),$$

which is a contradiction.

By the definition of $r_{j,n}(s)$, every disjoint ball has $\tilde{\mu}_{j,K}$ -mass at least 2^{-2^n} . Hence, we have

$$1 \geq \sum_{s \in S_{j,n}} \tilde{\mu}_{j,K} \left(B_{d_{j,K}}(s, r_{j,n}(s)) \right) \geq |S_{j,n}| 2^{-2^n}.$$

Consequently, we get $|S_{j,n}| \leq 2^{2^n}$. We next prove that $S_{j,n}$ approximates every point at its own mass radius. If $t \in S_{j,n}$, then $d_j(t, S_{j,n}) = 0$. If $t \notin S_{j,n}$, the point t was rejected because there exists a previously retained point $s \in S_{j,n}$ such that $r_{j,n}(s) \leq r_{j,n}(t)$ and

$$d_j(t, s) \leq r_{j,n}(t) + r_{j,n}(s).$$

Therefore, in both cases, we have

$$d_j(t, S_{j,n}) \leq 2r_{j,n}(t), \quad t \in K. \quad (\text{C.1})$$

For $t \in K$, we define

$$F_{j,t}(r) := \left(\log \frac{1}{\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, r))} \right)^{p_j}.$$

If $r < r_{j,n}(t)$, by the definition of $r_{j,n}(t)$, we have that $\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, r)) < 2^{-2^n}$, which implies that

$$F_{j,t}(r) \geq (\log 2)^{p_j} 2^{np_j}. \quad (\text{C.2})$$

Since the intervals $[r_{j,n+1}(t), r_{j,n}(t)]$, $n \geq 0$ are pairwise disjoint and contained in $[0, r_{j,0}(t)]$, applying inequality (C.2) gives us

$$\int_0^{r_{j,0}(t)} F_{j,t}(r) dr \geq (\log 2)^{p_j} \sum_{n \geq 0} 2^{np_j} (r_{j,n}(t) - r_{j,n+1}(t)). \quad (\text{C.3})$$

We next compare the last sum with the weighted sum of the mass radii. If the integral on the left-hand side is infinite, the desired estimate is immediate. Suppose that it is finite. Then we have $\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, 0)) > 0$. Indeed, if this mass were zero, the finiteness of K would imply that $F_{j,t}$ is infinite on an interval of positive length and the conclusion is trivial.

Since $2^{-2^n} \downarrow 0$ as n grows to infinity, the positive mass at radius zero implies that $r_{j,n}(t) = 0$ for sufficiently large n .

We may therefore apply summation by parts and obtain

$$\sum_{n \geq 0} 2^{np_j} (r_{j,n}(t) - r_{j,n+1}(t)) = (1 - 2^{-p_j}) \sum_{n \geq 0} 2^{np_j} r_{j,n}(t) + 2^{-p_j} r_{j,0}(t) \geq (1 - 2^{-p_j}) \sum_{n \geq 0} 2^{np_j} r_{j,n}(t).$$

Combining this identity with (C.3), we obtain

$$\sum_{n \geq 0} 2^{np_j} r_{j,n}(t) \leq \frac{1}{(\log 2)^{p_j} (1 - 2^{-p_j})} \int_0^{r_{j,0}(t)} F_{j,t}(r) dr. \quad (\text{C.4})$$

We now define the admissible sequence. Let $T_{j,0} := \{t_0\}$, $T_{j,k} := S_{j,k-1}$, $k \geq 1$. Thus, we have that $|T_{j,0}| = 1$, $|T_{j,k}| \leq 2^{2^{k-1}} \leq 2^{2^k}$. Thus, the sequence is admissible. Since K is finite, choose $k_K \geq 1$ such that $2^{2^{k_K}} \geq |K|$ and replace $T_{j,k}$ by K for every $k \geq k_K$. This replacement only decreases $d_j(t, T_{j,k})$, thus it preserves admissibility and ensures that $T_{j,k} = K$ for sufficiently large k .

Since $T_{j,0} = \{t_0\}$, we have

$$d_j(t, T_{j,0}) = d_j(t, t_0) = v_j(t).$$

Moreover, by (C.1), for every $k \geq 1$, we get

$$d_j(t, T_{j,k}) \leq 2r_{j,k-1}(t).$$

This inequality remains valid at the levels at which $T_{j,k}$ has been replaced by K , because the left-hand side is then zero. Consequently, we have

$$\begin{aligned} G_{j,K}(t) &= \sum_{k \geq 0} 2^{kp_j} d_j(t, T_{j,k}) = v_j(t) + \sum_{k \geq 1} 2^{kp_j} d_j(t, T_{j,k}) \\ &\leq v_j(t) + 2 \sum_{k \geq 1} 2^{kp_j} r_{j,k-1}(t) = v_j(t) + 2^{1+p_j} \sum_{n \geq 0} 2^{np_j} r_{j,n}(t). \end{aligned}$$

Using (C.4), we conclude that

$$G_{j,K}(t) \leq v_j(t) + \frac{2^{1+p_j}}{(\log 2)^{p_j} (1 - 2^{-p_j})} \int_0^{r_{j,0}(t)} F_{j,t}(r) dr. \quad (\text{C.5})$$

The atom at the anchor now gives the required pointwise truncation. The closed ball $B_{d_{j,K}}(t, v_j(t))$ contains t_0 . Since

$$\tilde{\mu}_{j,K}(\{t_0\}) \geq \frac{1}{2} = 2^{-2^0},$$

the definition of $r_{j,0}(t)$ implies that $r_{j,0}(t) \leq v_j(t)$. Therefore, we have

$$\int_0^{r_{j,0}(t)} F_{j,t}(r) dr \leq \int_0^{4v_j(t)} \left(\log \frac{1}{\tilde{\mu}_{j,K}(B_{d_{j,K}}(t, r))} \right)^{p_j} dr \leq 2c_{\alpha_j}^{(0)} \Phi_j(t) + 4c_{\alpha_j}^{(0)} (\log 2)^{p_j} v_j(t),$$

where the final inequality follows from Lemma C.2. Inserting this estimate into (C.5) and absorbing the constants into $C_{\alpha_j}^{(1)}$ proves

$$G_{j,K}(t) \leq C_{\alpha_j}^{(1)} (\Phi_j(t) + v_j(t)).$$

The construction is deterministic, and the inequality holds simultaneously for every $t \in K$. This holds for $j = 1$ and $j = 2$. We finish the proof. $\square$

We next combine the two admissible sequences. The common refinement is essential because the increment assumption controls the sum of the two metric contributions for a single process.

Lemma C.4. Let $K \subseteq T$ be finite and contain t_0 . For every $j \in \{1, 2\}$, let $\mathcal{T}_j = (T_{j,k})_{k \geq 0}$ be an admissible sequence of subsets of K such that $T_{j,0} = \{t_0\}$ and $T_{j,k} = K$ for all sufficiently large k . We define

$$G_j(t) := \sum_{k \geq 0} 2^{k/\alpha_j} d_j(t, T_{j,k}), \quad t \in K.$$

There exists a finite constant A_{α_1, α_2} , depending only on α_1 and α_2 , such that, for every $u \geq 1$, with probability at least $1 - e^{-u}$, simultaneously for every $t \in K$,

$$\|Z_t\| \leq A_{\alpha_1, \alpha_2} \left\{ G_1(t) + G_2(t) + u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) \right\}.$$

Proof of Lemma C.4. For $j \in \{1, 2\}$, $k \geq 0$, and $t \in K$, define the nearest point map $p_{j,k}(t) \in T_{j,k}$ such that

$$d_j(t, p_{j,k}(t)) = d_j(t, T_{j,k}).$$

Such a point exists because $T_{j,k}$ is a nonempty finite set. When $T_{j,k} = K$, we choose $p_{j,k}(t) := t$. For every $n \geq 2$, define an equivalence relation on K by

$$s \sim_n t \text{ iff } p_{j,k}(s) = p_{j,k}(t)$$

for every $j \in \{1, 2\}$ and every $0 \leq k \leq n-2$. Let $\mathcal{A}_n$ be the corresponding partition of K , and we write $\mathcal{A}_n(t)$ for the cell containing t . Equivalently, we define

$$P_n(t) := (p_{1,0}(t), \dots, p_{1,n-2}(t), p_{2,0}(t), \dots, p_{2,n-2}(t)).$$

Then two points lie in the same cell of $\mathcal{A}_n$ precisely when they have the same image under P_n .

The partitions are nested. Indeed, if $s \sim_n t$, then the nearest point labels of s and t agree at all levels $0 \leq k \leq n-2$, and therefore they also agree at all levels $0 \leq k \leq n-3$. Hence,

$$s \sim_n t \Rightarrow s \sim_{n-1} t.$$

Thus, $\mathcal{A}_n$ refines $\mathcal{A}_{n-1}$ for every $n \geq 3$.

We next bound the number of cells. By admissibility of the two sequences, we have that

$$|\mathcal{A}_n| \leq \prod_{j=1}^2 \prod_{k=0}^{n-2} |T_{j,k}| \leq \prod_{j=1}^2 \prod_{k=0}^{n-2} 2^{2^k} = 2^{2 \sum_{k=0}^{n-2} 2^k} = 2^{2(2^{n-1}-1)} = 2^{2^n-2} \leq 2^{2^n}.$$

For every $n \geq 2$, define the pointwise combined approximation cost

$$R_n(t) := 2^{n/\alpha_1} d_1(t, T_{1,n-2}) + 2^{n/\alpha_2} d_2(t, T_{2,n-2}).$$

For every cell $A \in \mathcal{A}_n$, choose one point $r_n(A)$ such that

$$r_n(A) \in \operatorname{argmin}_{s \in A} R_n(s),$$

and define $\pi_n(t) := r_n(\mathcal{A}_n(t))$. Thus, π_n is constant on every cell of $\mathcal{A}_n$.

Since t and $\pi_n(t)$ belong to the same cell, they have the same nearest-point label at level $n-2$. Therefore, we have that

$$p_{j,n-2}(t) = p_{j,n-2}(\pi_n(t)), \quad j \in \{1, 2\}.$$

By the triangle inequality, we obtain

$$d_j(t, \pi_n(t)) \leq d_j(t, p_{j,n-2}(t)) + d_j(p_{j,n-2}(t), \pi_n(t)) = d_j(t, T_{j,n-2}) + d_j(\pi_n(t), T_{j,n-2}).$$

Multiplying by $2^{n/\alpha_j}$ and summing over $j \in \{1, 2\}$, we obtain

$$2^{n/\alpha_1} d_1(t, \pi_n(t)) + 2^{n/\alpha_2} d_2(t, \pi_n(t)) \leq R_n(t) + R_n(\pi_n(t)).$$

Since $\pi_n(t)$ is the minimal distance, we have

$$R_n(\pi_n(t)) \leq R_n(t).$$

Consequently,

$$2^{n/\alpha_1} d_1(t, \pi_n(t)) + 2^{n/\alpha_2} d_2(t, \pi_n(t)) \leq 2R_n(t). \quad (\text{C.6})$$

Define $a_{\alpha_1, \alpha_2} := \max\{2^{1/\alpha_1}, 2^{1/\alpha_2}\}$. For every $n \geq 3$, the triangle inequality gives

$$d_j(\pi_n(t), \pi_{n-1}(t)) \leq d_j(\pi_n(t), t) + d_j(t, \pi_{n-1}(t)).$$

Using (C.6) at level n , we obtain $\sum_{j=1}^2 2^{n/\alpha_j} d_j(\pi_n(t), t) \leq 2R_n(t)$.

At level $n-1$, we use $2^{n/\alpha_j} \leq a_{\alpha_1, \alpha_2} 2^{(n-1)/\alpha_j}$ and obtain that

$$\sum_{j=1}^2 2^{n/\alpha_j} d_j(t, \pi_{n-1}(t)) \leq a_{\alpha_1, \alpha_2} \sum_{j=1}^2 2^{(n-1)/\alpha_j} d_j(t, \pi_{n-1}(t)) \leq 2a_{\alpha_1, \alpha_2} R_{n-1}(t).$$

Therefore, combining the inequalities above, we have

$$2^{n/\alpha_1} d_1(\pi_n(t), \pi_{n-1}(t)) + 2^{n/\alpha_2} d_2(\pi_n(t), \pi_{n-1}(t)) \leq 2R_n(t) + 2a_{\alpha_1, \alpha_2} R_{n-1}(t). \quad (\text{C.7})$$

Because the nearest-point maps are eventually the identity, the partitions $\mathcal{A}_n$ are eventually the singleton partition of K . Hence, for every $t \in K$, we have $\pi_n(t) = t$ for all sufficiently large n .

For every $n \geq 3$, define the set of distinct parent-child edges by

$$E_n := \{(\pi_n(t), \pi_{n-1}(t)) : t \in K\}.$$

Since $\mathcal{A}_n$ refines $\mathcal{A}_{n-1}$, every cell of $\mathcal{A}_n$ is contained in a unique cell of $\mathcal{A}_{n-1}$. Both π_n and π_{n-1} are constant on a cell of $\mathcal{A}_n$. Thus, every cell of $\mathcal{A}_n$ determines at most one edge in E_n , and

$$|E_n| \leq |\mathcal{A}_n| \leq 2^{2^n}. \quad (\text{C.8})$$

We first suppose that $u \geq 4$ and define $\ell := \lfloor \log_2 u \rfloor$. Then, we have

$$\ell \geq 2, \quad 2^\ell \leq u < 2^{\ell+1}.$$

We now apply the mixed-tail increment bound with parameter $16u$ to all distinct coarse edges $\{(\pi_\ell(t), t_0), t \in K\}$. The number of distinct points $\pi_\ell(t)$ is at most $|\mathcal{A}_\ell|$. Hence, by a union bound, we have

$$\begin{aligned} & \mathbb{P} \left\{ \exists t \in K \text{ s.t. } \|Z_{\pi_\ell(t)} - Z_{t_0}\| > (16u)^{1/\alpha_1} d_1(\pi_\ell(t), t_0) + (16u)^{1/\alpha_2} d_2(\pi_\ell(t), t_0) \right\} \\ & \leq 2|\mathcal{A}_\ell| e^{-16u} \leq 2 \exp \left\{ (\log 2) 2^\ell - 16u \right\} \leq 2e^{-15u}. \end{aligned}$$

For every $n > \ell$, apply the mixed-tail increment bound with parameter $16 \cdot 2^n$ to every edge in E_n . By (C.8) and another union bound,

$$\begin{aligned} & \mathbb{P} \left\{ \exists (s, r) \in E_n \text{ s.t. } \|Z_s - Z_r\| > (16 \cdot 2^n)^{1/\alpha_1} d_1(s, r) + (16 \cdot 2^n)^{1/\alpha_2} d_2(s, r) \right\} \\ & \leq 2|E_n| e^{-16 \cdot 2^n} \leq 2 \exp \left\{ -(16 - \log 2) 2^n \right\} \leq 2e^{-15 \cdot 2^n}. \end{aligned}$$

Since $2^{\ell+1} > u$, by direct algebra, we have

$$\sum_{n > \ell} 2e^{-15 \cdot 2^n} \leq 2 \sum_{r \geq 1} e^{-15u 2^{r-1}} \leq 2 \sum_{r \geq 1} e^{-15ur} = \frac{2e^{-15u}}{1 - e^{-15u}}.$$

For $u \geq 4$, we have that $2e^{-15u} \leq \frac{1}{2}e^{-u}$, $\frac{2e^{-15u}}{1 - e^{-15u}} \leq \frac{1}{2}e^{-u}$.

Therefore, outside an event $\mathcal{E}_u$ of probability at most e^{-u} , all the preceding coarse and fine increment bounds hold simultaneously. We now work conditioning on this event $\mathcal{E}_u$.

Since $\pi_n(t) = t$ for all sufficiently large n and $Z_{t_0} = 0$, we may telescope and obtain, simultaneously for every $t \in K$,

$$\begin{aligned} \|Z_t\| & \leq \|Z_{\pi_\ell(t)} - Z_{t_0}\| + \sum_{n > \ell} \|Z_{\pi_n(t)} - Z_{\pi_{n-1}(t)}\| \\ & \leq \sum_{j=1}^2 (16u)^{1/\alpha_j} d_j(\pi_\ell(t), t_0) + \sum_{n > \ell} \sum_{j=1}^2 (16 \cdot 2^n)^{1/\alpha_j} d_j(\pi_n(t), \pi_{n-1}(t)). \end{aligned}$$

We bound these two terms separately. For the first term, by the triangle inequality, we have

$$d_j(\pi_\ell(t), t_0) \leq d_j(\pi_\ell(t), t) + d_j(t, t_0) = d_j(t, \pi_\ell(t)) + v_j(t).$$

Define $c_{\alpha_1, \alpha_2} := \max \{16^{1/\alpha_1}, 16^{1/\alpha_2}\}$. Since $u < 2^{\ell+1}$, we have

$$u^{1/\alpha_j} < 2^{1/\alpha_j} 2^{\ell/\alpha_j} \leq a_{\alpha_1, \alpha_2} 2^{\ell/\alpha_j}.$$

It follows from (C.6) that

$$\sum_{j=1}^2 (16u)^{1/\alpha_j} d_j(t, \pi_\ell(t)) \leq c_{\alpha_1, \alpha_2} a_{\alpha_1, \alpha_2} \sum_{j=1}^2 2^{\ell/\alpha_j} d_j(t, \pi_\ell(t)) \leq 2c_{\alpha_1, \alpha_2}^{(16)} a_{\alpha_1, \alpha_2} R_\ell(t).$$

The anchor part satisfies that

$$\sum_{j=1}^2 (16u)^{1/\alpha_j} v_j(t) \leq c_{\alpha_1, \alpha_2} \left\{ u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) \right\}.$$

For the term $\sum_{n>\ell} \sum_{j=1}^2 (16 \cdot 2^n)^{1/\alpha_j} d_j(\pi_n(t), \pi_{n-1}(t))$, by inequality (C.7), we have that

$$\begin{aligned} \sum_{n>\ell} \sum_{j=1}^2 (16 \cdot 2^n)^{1/\alpha_j} d_j(\pi_n(t), \pi_{n-1}(t)) &\leq c_{\alpha_1, \alpha_2} \sum_{n>\ell} \{2R_n(t) + 2a_{\alpha_1, \alpha_2} R_{n-1}(t)\} \\ &\leq 2c_{\alpha_1, \alpha_2} (1 + a_{\alpha_1, \alpha_2}) \sum_{n \geq 2} R_n(t). \end{aligned}$$

Combining these inequalities together, we obtain

$$\|Z_t\| \leq C_{\alpha_1, \alpha_2}^{(2)} \left\{ u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) + \sum_{n \geq 2} R_n(t) \right\}. \quad (\text{C.9})$$

Finally, by the definition of $R_n(t)$ and the change of variables $k = n - 2$, we have the exact identity

$$\sum_{n \geq 2} R_n(t) = \sum_{n \geq 2} \sum_{j=1}^2 2^{n/\alpha_j} d_j(t, T_{j, n-2}) = \sum_{j=1}^2 2^{2/\alpha_j} \sum_{k \geq 0} 2^{k/\alpha_j} d_j(t, T_{j, k}) = 2^{2/\alpha_1} G_1(t) + 2^{2/\alpha_2} G_2(t).$$

Inserting this identity into (C.9) and enlarging the constant proves the conclusion for $u \geq 4$.

We now suppose that $1 \leq u < 4$. Apply the result already proved with the parameter 4. Its failure probability is $e^{-4} \leq e^{-u}$. Moreover, since $u \geq 1$, by some algebra, we have

$$4^{1/\alpha_j} v_j(t) \leq 4^{1/\alpha_j} u^{1/\alpha_j} v_j(t), \quad j \in \{1, 2\}.$$

Absorbing the constants $4^{1/\alpha_1}$ and $4^{1/\alpha_2}$ into A_{α_1, α_2} proves the result for every $u \geq 1$.

Thus, we finish the proof. $\square$

Combining the deterministic measure chains with the preceding probabilistic lemma gives the finite ambient-measure estimate used in the proof of the theorem.

Lemma C.5. There exists a finite constant B_{α_1, α_2} , depending only on α_1 and α_2 , such that, for every finite set $K \subseteq T$ containing t_0 and every $u \geq 1$, with probability at least $1 - e^{-u}$, simultaneously for every $t \in K$, we have

$$\|Z_t\| \leq B_{\alpha_1, \alpha_2} \left\{ \Phi_1(t) + \Phi_2(t) + u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) \right\}.$$

Proof of Lemma C.5. By Lemma C.3, for every $j \in \{1, 2\}$, there exists an admissible sequence $\mathcal{T}_j = (T_{j, k})_{k \geq 0}$, anchored at t_0 and eventually equal to K , such that, simultaneously for every $t \in K$,

$$G_j(t) := \sum_{k \geq 0} 2^{k/\alpha_j} d_j(t, T_{j, k}) \leq C_{\alpha_j}^{(1)} (\Phi_j(t) + v_j(t)).$$

Apply Lemma C.4 to these two sequences. Outside an event of probability at most e^{-u} , for every $t \in K$, we have

$$\begin{aligned} \|Z_t\| &\leq A_{\alpha_1, \alpha_2} \left\{ G_1(t) + G_2(t) + u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) \right\} \\ &\leq A_{\alpha_1, \alpha_2} \left[C_{\alpha_1}^{(1)} \Phi_1(t) + C_{\alpha_2}^{(1)} \Phi_2(t) + C_{\alpha_1}^{(1)} v_1(t) + C_{\alpha_2}^{(1)} v_2(t) + u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) \right]. \end{aligned}$$

Since $u \geq 1$, we have

$$v_j(t) \leq u^{1/\alpha_j} v_j(t), \quad j \in \{1, 2\}.$$

Therefore, we have

$$C_{\alpha_j}^{(1)} v_j(t) + u^{1/\alpha_j} v_j(t) \leq (C_{\alpha_j}^{(1)} + 1) u^{1/\alpha_j} v_j(t).$$

Absorbing the finitely many constants into B_{α_1, α_2} , we finish the proof. $\square$

We are now ready to prove the main theorem.

Proof of Theorem 3.1. For positive rational vectors $\mathbf{q} = (q_1, q_2) \in \mathbb{Q}_{>0}^2$, $\mathbf{s} = (s_1, s_2) \in \mathbb{Q}_{>0}^2$, we define

$$H_{\mathbf{q}, \mathbf{s}} := \{t \in T : \Phi_j(t) \leq q_j \text{ and } v_j(t) \leq s_j \text{ for } j = 1, 2\}.$$

By Lemma C.1, every set $H_{\mathbf{q}, \mathbf{s}}$ is Borel measurable.

We next choose countable dense subsets of these sets. Since (T, ρ) is a separable pseudo-metric space, its metric quotient by the relation $\rho(s, t) = 0$ is a separable metric space and hence is second countable. $H_{\mathbf{q}, \mathbf{s}}$ is a subset of T . Consequently, every $H_{\mathbf{q}, \mathbf{s}}$ is separable with respect to ρ .

For every pair $(\mathbf{q}, \mathbf{s})$, we choose a countable ρ -dense subset such that $D_{\mathbf{q}, \mathbf{s}} \subseteq H_{\mathbf{q}, \mathbf{s}}$. If the corresponding set is empty, take $D_{\mathbf{q}, \mathbf{s}} = \emptyset$.

The collection of positive rational vectors is countable. Hence, the union

$$D := \{t_0\} \cup \bigcup_{\mathbf{q} \in \mathbb{Q}_{>0}^2, \mathbf{s} \in \mathbb{Q}_{>0}^2} D_{\mathbf{q}, \mathbf{s}}$$

is countable. Beginning with the anchor, we enumerate it as $D = \{t_0, t_1, t_2, \dots\}$.

If D is finite, repeat t_0 in the enumeration so that the same notation remains valid. Define the increasing sequence of finite sets

$$K_N := \{t_0, t_1, \dots, t_N\}, \quad N \geq 1.$$

For every $N \geq 1$, let O_N be the event

$$O_N := \bigcap_{t \in K_N} \left\{ \|Z_t\| \leq B_{\alpha_1, \alpha_2} \left[\Phi_1(t) + \Phi_2(t) + u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) \right] \right\}.$$

These are the actual conclusion events in Lemma C.5. Then, we have that

$$\mathbb{P}(O_N) \geq 1 - e^{-u}, \quad N \geq 1.$$

Moreover, since $K_N \subseteq K_{N+1}$, the events are decreasing, i.e., $E_{N+1} \subseteq E_N$.

By the continuity of probability, we have that,

$$\mathbb{P} \left(\bigcap_{N \geq 1} O_N \right) = \lim_{N \rightarrow \infty} \mathbb{P}(O_N) \geq 1 - e^{-u}.$$

Fix an outcome in this event. By the null-set modification made at the beginning of the appendix, the sample path $t \mapsto Z_t$ is ρ -continuous. Let $t \in T$ satisfy $\Phi_1(t) < \infty$, $\Phi_2(t) < \infty$ and choose arbitrary positive rational numbers such that

$$q_j > \Phi_j(t), \quad s_j > v_j(t), \quad j \in \{1, 2\}.$$

Then, we have $t \in H_{\mathbf{q}, \mathbf{s}}$. Since $D_{\mathbf{q}, \mathbf{s}}$ is ρ -dense in this set, there exists a sequence $\{x_m\}_{m \geq 1} \subseteq D_{\mathbf{q}, \mathbf{s}}$ such that $\rho(x_m, t) \rightarrow 0$.

Every point x_m belongs to D , and hence it belongs to K_N for all sufficiently large N . Conditioned on the event $\bigcap_{N \geq 1} O_N$, we have that

$$\begin{aligned} \|Z_{x_m}\| &\leq B_{\alpha_1, \alpha_2} \left\{ \Phi_1(x_m) + \Phi_2(x_m) + u^{1/\alpha_1} v_1(x_m) + u^{1/\alpha_2} v_2(x_m) \right\} \\ &\leq B_{\alpha_1, \alpha_2} \left\{ q_1 + q_2 + u^{1/\alpha_1} s_1 + u^{1/\alpha_2} s_2 \right\}. \end{aligned}$$

Recall the path continuity and we have $Z_{x_m} \rightarrow Z_t$ in $(\mathbb{B}, \|\cdot\|)$.

Hence, we get

$$\|Z_t\| = \lim_{m \rightarrow \infty} \|Z_{x_m}\| \leq B_{\alpha_1, \alpha_2} \left\{ q_1 + q_2 + u^{1/\alpha_1} s_1 + u^{1/\alpha_2} s_2 \right\}.$$

Letting $q_j \downarrow \Phi_j(t)$ and $s_j \downarrow v_j(t)$ through rational values and we obtain

$$\|Z_t\| \leq B_{\alpha_1, \alpha_2} \left\{ \Phi_1(t) + \Phi_2(t) + u^{1/\alpha_1} v_1(t) + u^{1/\alpha_2} v_2(t) \right\}.$$

The event used above does not depend on t . Hence, the inequality holds simultaneously for every $t \in T$ for which both profiles are finite. If either profile is infinite, the assertion is vacuous.

Finally, set $u := \log \frac{e}{\delta}$. Then $u \geq 1$ and $e^{-u} = \frac{\delta}{e} \leq \delta$. The theorem follows with $C_{\alpha_1, \alpha_2} := B_{\alpha_1, \alpha_2}$. Therefore, with probability at least $1 - \delta$, simultaneously for every $t \in T$,

$$\|Z_t\| \leq C_{\alpha_1, \alpha_2} \left\{ \Phi_1(t) + \Phi_2(t) + v_1(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha_1} + v_2(t) \left(\log \frac{e}{\delta} \right)^{1/\alpha_2} \right\}.$$

We finish the proof. □

D Proofs in Section 4

Proof of Proposition 4.2. We first prove the mixed-tail increment inequality. Fix $s, t \in T$ and define

$$D_{s,t} := B_s - B_t.$$

Since B_s and B_t are symmetric, $D_{s,t}$ is symmetric. By Definition 4.1, we have

$$C_s(\xi) - C_t(\xi) = \xi^\top (B_s - B_t) \xi - \mathbb{E}_\xi [\xi^\top (B_s - B_t) \xi] = \xi^\top D_{s,t} \xi - \mathbb{E}_\xi [\xi^\top D_{s,t} \xi].$$

We next derive the explicit two-sided inequality needed below. Suppose that $D \neq 0$, and, for $x \geq 0$, define $q_D(x) := \min \left\{ \frac{x^2}{M_0^4 \|D\|_F^2}, \frac{x}{M_0^2 \|D\|_{\text{op}}} \right\}$.

Applying Lemma A.1 first to D and then to $-D$, and using a union bound, we obtain

$$\mathbb{P}_\xi \left\{ \left| \xi^\top D \xi - \mathbb{E}_\xi [\xi^\top D \xi] \right| > x \right\} \leq \min \left\{ 1, 4 \exp \left(-\frac{q_D(x)}{c_{HW}} \right) \right\}. \quad (\text{D.1})$$

We now claim that

$$\min \left\{ 1, 4 \exp \left(-\frac{q_D(x)}{c_{HW}} \right) \right\} \leq 2 \exp \left(-\frac{q_D(x)}{2c_{HW}} \right).$$

Indeed, if $q_D(x) \leq 2c_{HW} \log 2$, then the right-hand side is at least one, so the claim follows from the first term in the minimum. If $q_D(x) > 2c_{HW} \log 2$, then

$$4 \exp \left(-\frac{q_D(x)}{c_{HW}} \right) = 2 \exp \left(-\frac{q_D(x)}{2c_{HW}} \right) \left[2 \exp \left(-\frac{q_D(x)}{2c_{HW}} \right) \right] \leq 2 \exp \left(-\frac{q_D(x)}{2c_{HW}} \right).$$

Thus, the claim is true. Consequently, for every nonzero deterministic real matrix D and every $x \geq 0$, we have the explicit two-sided bound

$$\mathbb{P}_\xi \left\{ \left| \xi^\top D \xi - \mathbb{E}_\xi [\xi^\top D \xi] \right| > x \right\} \leq 2 \exp \left\{ -\frac{1}{2c_{HW}} \min \left\{ \frac{x^2}{M_0^4 \|D\|_F^2}, \frac{x}{M_0^2 \|D\|_{\text{op}}} \right\} \right\}.$$

Recall that $C_{HW} = 2c_{HW}$. For $u \geq 0$, set $x := C_{HW} M_0^2 \{ \sqrt{u} \|D_{s,t}\|_F + u \|D_{s,t}\|_{\text{op}} \}$. If $D_{s,t} = 0$, the desired assertion is immediate.

Suppose that $D_{s,t} \neq 0$. Since we have $x \geq C_{HW} M_0^2 \sqrt{u} \|D_{s,t}\|_F$, we obtain that $\frac{x^2}{M_0^4 \|D_{s,t}\|_F^2} \geq C_{HW}^2 u$.

Similarly, we have $x \geq C_{HW} M_0^2 u \|D_{s,t}\|_{\text{op}}$, which implies that $\frac{x}{M_0^2 \|D_{s,t}\|_{\text{op}}} \geq C_{HW} u$.

Therefore, since $C_{HW} \geq 1$, we have

$$q_{D_{s,t}}(x) = \min \left\{ \frac{x^2}{M_0^4 \|D_{s,t}\|_F^2}, \frac{x}{M_0^2 \|D_{s,t}\|_{\text{op}}} \right\} \geq \min \{ C_{HW}^2 u, C_{HW} u \} = C_{HW} u.$$

Applying (D.1) with $D = D_{s,t}$ and this choice of x , we obtain

$$\mathbb{P}_\xi \left\{ |C_s(\xi) - C_t(\xi)| > C_{HW} M_0^2 \{ \sqrt{u} \|B_s - B_t\|_F + u \|B_s - B_t\|_{\text{op}} \} \right\} \leq 2e^{-u}.$$

By the definitions of d_1 and d_2 , this is equivalent to

$$\mathbb{P}_\xi \left\{ |C_s(\xi) - C_t(\xi)| > u d_{\text{op}}(s, t) + \sqrt{u} d_F(s, t) \right\} \leq 2e^{-u}.$$

Thus, the required mixed-tail increment bound holds.

We next verify the pseudo-metric and measurability properties. The triangle inequalities for the

operator norm and the Frobenius norm show that d_{op} and d_{F} are pseudo-metrics. Since $t \mapsto B_t$ is Borel measurable, the map $(s, t) \mapsto B_s - B_t$ is Borel measurable from $T \times T$ to $\mathbb{R}^{n \times n}$. Both matrix norms are continuous. Consequently, d_{op} and d_{F} are jointly Borel measurable. They are finite-valued because each B_t is a finite-dimensional matrix.

We now prove separability. The finite-dimensional matrix space $\mathbb{R}^{n \times n}$ is separable under the Frobenius norm. Its subset $\mathcal{M} := \{B_t : t \in T\}$ therefore contains a countable Frobenius-dense subset $\mathcal{M}_0$. For each matrix in $\mathcal{M}_0$, choose one preimage in T , and let T_0 be the resulting countable subset of T . For every matrix D , we have $\|D\|_{\text{op}} \leq \|D\|_{\text{F}}$.

Hence, Frobenius convergence implies operator-norm convergence. It follows that T_0 is dense in T with respect to $\rho = d_{\text{op}} + d_{\text{F}}$. Thus, (T, ρ) is separable.

It remains to verify the process-level measurability and continuity. By the coordinate representation

$$C_t(\xi) = \sum_{i,j=1}^n B_{ij}(t) (\xi_i \xi_j - \mathbb{E}_{\xi} [\xi_i \xi_j]),$$

the map $(\omega, t) \mapsto C_t(\xi(\omega))$ is jointly measurable. Moreover, for every $s, t \in T$, we have

$$\begin{aligned} |C_s(\xi) - C_t(\xi)| &\leq \left| \xi^{\top} (B_s - B_t) \xi \right| + \mathbb{E}_{\xi} \left[\left| \xi^{\top} (B_s - B_t) \xi \right| \right] \\ &\leq \left\{ \|\xi\|_2^2 + \mathbb{E}_{\xi} [\|\xi\|_2^2] \right\} \|B_s - B_t\|_{\text{op}} \\ &= \frac{\|\xi\|_2^2 + \mathbb{E}_{\xi} [\|\xi\|_2^2]}{C_{\text{HW}} M_0^2} d_{\text{op}}(s, t) \\ &\leq \frac{\|\xi\|_2^2 + \mathbb{E}_{\xi} [\|\xi\|_2^2]}{C_{\text{HW}} M_0^2} \rho(s, t). \end{aligned}$$

The coefficient on the right-hand side is finite almost surely because the coordinates of ξ are sub-Gaussian. Therefore, with probability one, the map $t \mapsto C_t(\xi)$ is ρ -continuous. Finally, $B_{t_0} = 0$ implies $C_{t_0}(\xi) = 0$ almost surely. We have now verified all the assertions of the proposition. $\square$

Proof of Corollary 4.3. By Proposition 4.2, the centered quadratic chaos process satisfies all the assumptions of Theorem 3.1. Consequently, We apply Theorem 3.1 and obtain that with probability at least $1 - \delta$, simultaneously for every $t \in T$,

$$|C_t(\xi)| \leq C_{1,2} \left\{ \Phi_1(t) + \Phi_2(t) + v_1(t) \log \frac{e}{\delta} + v_2(t) \sqrt{\log \frac{e}{\delta}} \right\}.$$

This is precisely the claimed inequality. $\square$

E Proofs in Section 5

The diffusion results in Appendix A are stated directly in the notation of Section 5. We verify the explicit Poincaré constant, identify the long-run variance, and check the measurability and continuity hypotheses of the mixed-tail theorem.

E.1 The Poincaré constant and the long-run variance

Since b and σ are globally Lipschitz, they are locally Lipschitz and satisfy a linear-growth bound. Thus Assumption 5.1 is a special case of Definition 5 and equation (4.26) of [Aeckerle-Willems and Strauch \(2021\)](#). The statement following that equation and their equation (4.27) give the unique global strong solution, its ergodicity, and its invariant density π .

We first verify the quantities in Lemma A.2. Let

$$F(x) := \int_{-\infty}^x \pi(y) dy, \quad \bar{F}(x) := \int_x^{\infty} \pi(y) dy,$$

and define

$$B_+ := \sup_{x>0} \bar{F}(x) \int_0^x \frac{2}{\ell(y)} dy, \quad B_- := \sup_{x<0} F(x) \int_x^0 \frac{2}{\ell(y)} dy.$$

For $y \geq x \geq A$, by the drift assumption, we have that

$$G(y) - G(x) \leq -2\gamma(y - x). \quad (\text{E.1})$$

Since $a \geq \underline{\sigma}^2$, it follows that, for $x \geq A$, we have

$$\bar{F}(x) = \frac{1}{Z} \int_x^{\infty} \frac{e^{G(y)}}{a(y)} dy \leq \frac{e^{G(x)}}{2\gamma Z \underline{\sigma}^2} = \frac{\ell(x)}{2\gamma \underline{\sigma}^2}. \quad (\text{E.2})$$

Moreover, we have

$$\ell(x) \int_0^x \frac{2}{\ell(y)} dy \leq \ell(x) \int_0^A \frac{2}{\ell(y)} dy + 2 \int_A^x e^{-2\gamma(x-y)} dy \leq \ell(x) \int_0^A \frac{2}{\ell(y)} dy + \frac{1}{\gamma}.$$

Hence $B_+ < \infty$. Similarly, we have that $B_- < \infty$. According to Lemma A.2, we know that the stationary diffusion is reversible and that the Poincaré constant is

$$C_P := 4 \max \{B_+, B_-\} < \infty. \quad (\text{E.3})$$

We next identify the long-run variance. First, we claim that $\mathcal{V}_\pi(f)$ is a finite seminorm for every bounded Borel function f . Indeed, on $[A, \infty)$, equation (E.2) gives

$$\frac{R_f(x)^2}{\ell(x)} = \frac{(\int_x^\infty \bar{f}(y) \pi(y) dy)^2}{\ell(x)} \leq \frac{4\|f\|_\infty^2 (\int_x^\infty \pi(y) dy)^2}{\ell(x)} \leq 4\|f\|_\infty^2 \frac{\bar{F}(x)^2}{\ell(x)} \leq \frac{2\|f\|_\infty^2}{\gamma \underline{\sigma}^2} \bar{F}(x).$$

Taking the two arguments in (E.1) to be A and x , we have that $G(x) \leq G(A) - 2\gamma(x - A)$, $x \geq A$.

Combining this inequality with (E.2), we obtain $\bar{F}(x) \leq \frac{e^{G(A)}}{2\gamma Z \underline{\sigma}^2} e^{-2\gamma(x-A)}$, $x \geq A$.

Therefore, $\int_A^\infty \frac{2\|f\|_\infty^2}{\gamma \underline{\sigma}^2} \bar{F}(x) dx$ is finite because of the exponential tail above. Similarly, the left tail integral $\int_{-\infty}^{-A} \frac{2\|f\|_\infty^2}{\gamma \underline{\sigma}^2} \bar{F}(x) dx$ is finite. Lastly, by nature, the integral $\int_{-A}^A \frac{2\|f\|_\infty^2}{\gamma \underline{\sigma}^2} \bar{F}(x) dx$ is finite. Therefore, we obtain that $\int_{\mathbb{R}} \frac{R_f(x)^2}{\ell(x)} dx < \infty$ is finite and hence $\mathcal{V}_\pi(f) = 2\|\frac{R_f(x)}{\sqrt{\ell(x)}}\|_{L_2} < \infty$ is finite.

The centering map is linear, and hence we have that $R_{\lambda f + \eta h} = \lambda R_f + \eta R_h$ for all bounded Borel

functions f, h and all $\lambda, \eta \in \mathbb{R}$. Consequently,

$$\mathcal{V}_\pi(\lambda f) = |\lambda| \mathcal{V}_\pi(f), \quad \mathcal{V}_\pi(f + h) \leq \mathcal{V}_\pi(f) + \mathcal{V}_\pi(h),$$

where the second inequality follows from the triangle inequality in the L^2 norm. Together with $\mathcal{V}_\pi(0) = 0$, this proves that $\mathcal{V}_\pi$ is a finite seminorm.

Now, to apply Lemma A.3, it remains to show that $\mathcal{V}_\pi(f)^2 = \lim_{t \rightarrow \infty} \frac{1}{t} \text{Var}_{\mathbb{P}_\pi} \left(\int_0^t \bar{f}(X_s) ds \right)$. For arbitrary $t > 0$, we write $Z_t(f) := \frac{1}{\sqrt{t}} \int_0^t \bar{f}(X_s) ds$. By Lemma A.4, we have that

$$Z_t(f) \implies \mathcal{N}(0, \mathcal{V}_\pi(f)^2), \quad t \rightarrow \infty. \quad (\text{E.4})$$

We next upgrade this weak convergence to convergence of second moments. Set $A_f = 2\sqrt{C_P} \|\bar{f}\|_{L^2(\pi)}$, $B_f = C_P \|\bar{f}\|_\infty$. By Inequality (A.3) in Lemma A.3, for $t \geq 1$ and $u > 0$, we have that

$$\mathbb{P}_\pi \{ |Z_t(f)| > A_f \sqrt{u} + B_f u \} \leq 2e^{-u}.$$

Consequently, by direct algebra, for any t , we have

$$\mathbb{E}_\pi |Z_t(f)|^4 = 4 \int_0^\infty x^3 \mathbb{P}_\pi(|Z_t(f)| > x) dx \leq 8 \int_0^\infty x^3 [e^{-x^2/(4A_f^2)} + e^{-x/2B_f}] dx \leq 64A_f^4 + 768B_f^4.$$

Taking supremum over t , we obtain that $\sup_{t \geq 1} \mathbb{E}_\pi |Z_t(f)|^4 \leq 64A_f^4 + 768B_f^4 < \infty$. It follows that $\{Z_t(f)^2 : t \geq 1\}$ is uniformly integrable. Indeed, for every $M > 0$, by Markov inequality, we have

$$\sup_{t \geq 1} \mathbb{E}_\pi \left[Z_t(f)^2 \mathbf{1}_{\{Z_t(f)^2 > M\}} \right] \leq \frac{1}{M} \sup_{t \geq 1} \mathbb{E}_\pi |Z_t(f)|^4 \rightarrow 0.$$

By (E.4) and continuous mapping theorem, $Z_t(f)^2$ converges in distribution to Y^2 , where $Y \sim \mathcal{N}(0, \mathcal{V}_\pi(f)^2)$. Uniform integrability therefore implies that $\mathbb{E}_\pi Z_t(f)^2 \rightarrow \mathbb{E} Y^2 = \mathcal{V}_\pi(f)^2$.

Finally, since $\pi(\bar{f}) = 0$, we know that $\mathbb{E}_\pi Z_t(f) = 0$, which implies $\mathbb{E}_\pi Z_t(f)^2 = \frac{1}{t} \text{Var}_{\mathbb{P}_\pi} \left(\int_0^t \bar{f}(X_s) ds \right)$. Therefore, we have that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \text{Var}_{\mathbb{P}_\pi} \left(\int_0^t \bar{f}(X_s) ds \right) = \mathcal{V}_\pi(f)^2. \quad (\text{E.5})$$

Thus, the assumptions in Lemma A.3 are satisfied. We apply Inequality (A.4) and (A.5) to obtain

$$\mathcal{V}_\pi(f)^2 \leq 2C_P \|\bar{f}\|_{L^2(\pi)}^2, \quad (\text{E.6})$$

Now, we are ready to prove Proposition 5.3

Proof of Proposition 5.3. The centered uniform norm and $\mathcal{V}_\pi$ are seminorms, so d_1 and d_2 are finite-valued pseudo-metrics and they are joint Borel measurable.

For $g := (f - h) - \pi(f - h)$, equation (E.6) gives

$$d_2(f, h)^2 = 2\mathcal{V}_\pi(f - h)^2 \leq 4C_P \|g\|_{L^2(\pi)}^2 \leq 4C_P \|g\|_\infty^2.$$

Thus, we have $d_2(f, h) \leq 2\sqrt{\frac{\tau}{C_P}} d_1(f, h)$ and every countable d_1 -dense set is also dense for $d_1 + d_2$.

The jointly Borel evaluation map implies that $f \mapsto \pi(f)$ is Borel. Since the continuous adapted diffusion process X is measurable, the map

$$(\omega, s, f) \mapsto f(X_s(\omega)) - \pi(f)$$

is also jointly measurable. Parametrized integration therefore shows that $(\omega, f) \mapsto Z_\tau(f)(\omega)$ is jointly measurable. For every sample path, notice that $|Z_\tau(f) - Z_\tau(h)| \leq \sqrt{\tau} \|g\|_\infty = \frac{\tau}{C_P} d_1(f, h)$, so the sample paths are $(d_1 + d_2)$ -continuous. The zero function is an anchor because $Z_\tau(0) = 0$.

Recall that from the deduction above, the conditions in Lemma A.3 are satisfied. Hence, we apply Lemma A.3 to get that for every $u > 0$,

$$\mathbb{P}_\pi \left\{ |Z_\tau(f) - Z_\tau(h)| > \sqrt{2u} \mathcal{V}_\pi(f - h) + \frac{C_P \|g\|_\infty}{\sqrt{\tau}} u \right\} \leq 2e^{-u}.$$

This is exactly our result and we finish the proof. $\square$

Proof of Corollary 5.4. Proposition 5.3 verifies the hypotheses of Theorem 3.1 with

$$(\alpha_1, d_1, \mu_1) = (1, d_1, \mu_1), \quad (\alpha_2, d_2, \mu_2) = (2, d_2, \mu_2).$$

The anchor radii and local ball-mass terms in that theorem are precisely v_1, v_2 and Φ_1, Φ_2 . Substitution gives (5.4) on one event of probability at least $1 - \delta$, simultaneously for every $f \in \mathcal{F}$. $\square$

References

- Radosław Adamczak. A note on the Hanson–Wright inequality for random vectors with dependencies. *Electronic Communications in Probability*, 20(72):1–13, 2015.
- Cathrine Aeckerle-Willems and Claudia Strauch. Concentration of scalar ergodic diffusions and some statistical implications. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 57(4):1857–1887, 2021.
- Jean-Yves Audibert and Olivier Bousquet. Combining PAC-Bayesian and generic chaining bounds. *Journal of Machine Learning Research*, 8:863–889, 2007.
- Peter L. Bartlett, Olivier Bousquet, and Shahar Mendelson. Local Rademacher complexities. *The Annals of Statistics*, 33(4):1497 – 1537, 2005.
- Witold Bednorz. A theorem on majorizing measures. *The Annals of Probability*, 34(5):1771–1781, 2006.
- Witold Bednorz. Hölder continuity of random processes. *Journal of Theoretical Probability*, 20(4): 917–934, 2007.
- Witold Bednorz. Concentration via chaining method and its applications. *arXiv preprint arXiv:1405.0676*, 2014.
- Sander Borst, Daniel Dadush, Neil Olver, and Makrand Sinha. Majorizing measures for the optimizer. In *12th Innovations in Theoretical Computer Science Conference (ITCS 2021)*, volume 185 of *Leibniz International Proceedings in Informatics*, pages 73:1–73:20. Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2021.

- Patrick Cattiaux and Arnaud Guillin. Deviation bounds for additive functionals of Markov processes. *ESAIM: Probability and Statistics*, 12:12–29, 2008.
- Yifeng Chu and Maxim Raginsky. Majorizing measures, codes, and information. *IEEE Transactions on Information Theory*, 72(5):3301–3311, 2026.
- Lee H. Dicker and Murat A. Erdogdu. Flexible results for quadratic forms with applications to variance components estimation. *The Annals of Statistics*, 45(1):386–414, 2017.
- Sjoerd Dirksen. Tail bounds via generic chaining. *Electronic Journal of Probability*, 20(53):1–29, 2015.
- Richard M. Dudley. The sizes of compact subsets of Hilbert space and continuity of Gaussian processes. *Journal of Functional Analysis*, 1(3):290–330, 1967.
- Xavier Fernique. Régularité de processus gaussiens. *Inventiones Mathematicae*, 12:304–320, 1971.
- Xavier Fernique. Régularité des trajectoires des fonctions aléatoires gaussiennes. In *École d’Été de Probabilités de Saint-Flour IV–1974*, volume 480 of *Lecture Notes in Mathematics*, pages 1–96. Springer, Berlin, 1975.
- Fuqing Gao, Arnaud Guillin, and Liming Wu. Bernstein-type concentration inequalities for symmetric Markov processes. *Theory of Probability and Its Applications*, 58(3):358–382, 2014.
- Adriano M. Garsia, Eugene Rodemich, and Jr. Rumsey, Hans. A real variable lemma and the continuity of paths of some Gaussian processes. *Indiana University Mathematics Journal*, 20: 565–578, 1971.
- David L. Hanson and Farroll T. Wright. A bound on tail probabilities for quadratic forms in independent random variables. *The Annals of Mathematical Statistics*, 42(3):1079–1083, 1971.
- Daniel Hsu, Sham M. Kakade, and Tong Zhang. A tail inequality for quadratic forms of subgaussian random vectors. *Electronic Communications in Probability*, 17(52):1–6, 2012.
- Haichen Hu and David Simchi-Levi. Perturbing the derivative: Doubly wild refitting for model-free evaluation of opaque machine learning predictors. *arXiv preprint arXiv:2511.18789*, 2025.
- Haichen Hu and David Simchi-Levi. Interleaved resampling and refitting: Data and compute-efficient evaluation of black-box predictors. *arXiv preprint arXiv:2603.14218*, 2026.
- Haichen Hu, Rui Ai, Stephen Bates, and David Simchi-Levi. Contextual online decision making with infinite-dimensional functional regression. In *Forty-second International Conference on Machine Learning*, 2025.
- Claude Kipnis and S. R. Srinivasa Varadhan. Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions. *Communications in Mathematical Physics*, 104(1):1–19, 1986.
- Yegor Klochov and Nikita Zhivotovskiy. Uniform Hanson–Wright type concentration inequalities for unbounded entries via the entropy method. *Electronic Journal of Probability*, 25(22):1–30, 2020.
- Vladimir Koltchinskii. Local Rademacher complexities and oracle inequalities in risk minimization. *The Annals of Statistics*, 34(6):2593–2656, 2006.

- Felix Krahmer, Shahar Mendelson, and Holger Rauhut. Suprema of chaos processes and the restricted isometry property. *Communications on Pure and Applied Mathematics*, 67(11):1877–1904, 2014.
- Yury A. Kutoyants. Some problems of nonparametric estimation by observations of ergodic diffusion process. *Statistics & Probability Letters*, 32(3):311–320, 1997.
- Yury A. Kutoyants. On the goodness-of-fit testing for ergodic diffusion processes. *Journal of Nonparametric Statistics*, 22(4):529–543, 2010.
- Stanisław Kwapień and Jan Rosiński. Sample hölder continuity of stochastic processes and majorizing measures. In *Seminar on Stochastic Analysis, Random Fields and Applications IV*, volume 58 of *Progress in Probability*, pages 155–163. Birkhäuser, Basel, 2004.
- Andrew Lamperski. Nonasymptotic pointwise and worst-case bounds for classical spectrum estimators. *IEEE Transactions on Signal Processing*, 71:4273–4287, 2023.
- Oleg Loukianov, Dasha Loukianova, and Shiqi Song. Spectral gaps and exponential integrability of hitting times for linear diffusions. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 47(3):679–698, 2011.
- Andreas Maurer. Majorizing codes and measures, 2010. Unpublished manuscript.
- Shahar Mendelson, Alain Pajor, and Nicole Tomczak-Jaegermann. Uniform uncertainty principle for Bernoulli and subgaussian ensembles. *Constructive Approximation*, 28(3):277–289, 2008.
- Yijin Ni and Xiaoming Huo. Upper confidence bounds for the prediction error of kernel ridge regression via gaussian refitting. *arXiv preprint arXiv:2607.28846*, 2026.
- Bob Pepin. Concentration inequalities for additive functionals: A martingale approach. *Stochastic Processes and their Applications*, 135:103–138, 2021.
- Christopher J. Preston. Continuity properties of some Gaussian processes. *The Annals of Mathematical Statistics*, 43(1):285–292, 1972.
- Frank Redig and Florian Völlering. Concentration of additive functionals for Markov processes and applications to interacting particle systems. *arXiv preprint arXiv:1003.0006v2*, 2011.
- Mark Rudelson and Roman Vershynin. Hanson-Wright inequality and sub-gaussian concentration. *Electronic Communications in Probability*, 18(none):1 – 9, 2013.
- James B Simon, Madeline Dickens, Dhruva Karkada, and Michael R DeWeese. The eigenlearning framework: A conservation law perspective on kernel ridge regression and wide neural networks. *Transactions on Machine Learning Research*, 2023. ISSN 2835-8856.
- Michel Talagrand. Regularity of Gaussian processes. *Acta Mathematica*, 159:99–149, 1987.
- Michel Talagrand. Sample boundedness of stochastic processes under increment conditions. *The Annals of Probability*, 18(1):1–49, 1990.
- Michel Talagrand. A simple proof of the majorizing measure theorem. *Geometric and Functional Analysis*, 2(1):118–125, 1992.
- Michel Talagrand. Constructions of majorizing measures, Bernoulli processes and cotype. *Geometric and Functional Analysis*, 4(6):660–717, 1994.

- Michel Talagrand. Majorizing measures: the generic chaining. *The Annals of Probability*, 24(3): 1049 – 1103, 1996.
- Michel Talagrand. Majorizing measures without measures. *The Annals of Probability*, 29(1):411–417, 2001.
- Michel Talagrand. *The Generic Chaining: Upper and Lower Bounds of Stochastic Processes*. Springer, Berlin, 2005.
- Lukas Trottner, Cathrine Aeckerle-Willems, and Claudia Strauch. Concentration analysis of multivariate elliptic diffusions. *Journal of Machine Learning Research*, 24(106):1–38, 2023.
- Aad W. van der Vaart and Harry van Zanten. Donsker theorems for diffusions: Necessary and sufficient conditions. *The Annals of Probability*, 33(4):1422–1451, 2005.
- Harry van Zanten. On empirical processes for ergodic diffusions and rates of convergence of M -estimators. *Scandinavian Journal of Statistics*, 30(3):443–458, 2003.
- Vladimir Vapnik. *The nature of statistical learning theory*. Springer science & business media, 2013.
- Martin J Wainwright. Wild refitting for black box prediction. *arXiv preprint arXiv:2506.21460*, 2025.
- Yunbei Xu. Pointwise complexity for Gaussian fields: Upper envelopes, algorithmic lower bounds, and separation. *arXiv preprint arXiv:2606.07931v2*, 2026.
- Yunbei Xu and Assaf Zeevi. Towards optimal problem dependent generalization error bounds in statistical learning theory. *Mathematics of Operations Research*, 50(1):40–67, 2025.
- Greg Yang, James B Simon, and Jeremy Bernstein. A spectral condition for feature learning. *arXiv preprint arXiv:2310.17813*, 2023.
- Tong Zhang. Covering number bounds of certain regularized linear function classes. *Journal of Machine Learning Research*, 2(Mar):527–550, 2002.