

What Can Go Wrong During Caplet Stripping ?

Fabien Le Floc'h 

* Correspondence: fabien@2ipi.com

Received: May 2026

Abstract: We study exact and near exact extraction of caplet volatilities from market cap quotes and identify why some common choices produce extreme oscillations or negative vols. Interpolation scheme and node placement are shown to be the primary drivers of instability, which can be amplified by isolated bad quotes. We propose practical, production ready remedies: continuous flat-linear and C1 flat-smooth kernels that preserve bootstrap equivalence, midpoint node placement with a global solver, positivity enforcement via an exponential reparametrization or Hyman non-negative C1 splines. We also introduce simple data quality checks. Numerical experiments demonstrate substantially reduced oscillations, robust positive caplet curves, and negligible repricing error, delivering a fast and stable caplet stripping workflow suitable for real-world use.

Keywords: cap; floor; caplet; stripping; interpolation

1. Introduction

Interest rate caps and floors are widely used financial instruments that provide protection against interest rate fluctuations. A cap is a series of European call options on interest rates, while a floor is a series of European put options. Caplet stripping is the process of extracting the implied volatilities of individual caplets from the market prices of caps and floors. This process is crucial for pricing and risk management in the interest rate derivatives market.

Caps are typically quoted in terms of their implied volatilities. There are three conventions: the Black-Scholes implied volatility, which was the historical choice before negative interest rates became more widespread (also known as lognormal volatility), the Bachelier implied volatility (also called normal volatility or b.p. vol because it is quoted in basis points) and the shifted lognormal volatility. For the sake of simplicity, we will focus here on the Bachelier quotes, and the stripping will consist of finding the Bachelier implied volatilities of the caplets that are consistent with the market prices of the caps. Our observations also apply to the other conventions.

Let $\hat{\sigma}$ be the quoted Bachelier volatility for the cap, the cap market price is the sum of the caplet prices using a constant volatility $\sigma_{i,K} = \hat{\sigma}$ for each caplet. In the context of the Bachelier model, it reads:

$$P(K, T) = \sum_{i=1}^M \alpha_i B_i^p \left[(F_i - K) \Phi \left(\frac{F_i - K}{\sigma_{i,K} \sqrt{t_i}} \right) + \sigma_{i,K} \sqrt{t_i} \phi \left(\frac{F_i - K}{\sigma_{i,K} \sqrt{t_i}} \right) \right], \quad (1)$$

where α_i is the year fraction of the period, F_i is the forward rate for the period, K is the strike rate, t_i is the time to maturity of the caplet, B_i^p is the discount factor to payment date for the caplet with maturity t_i (for a Libor rate, this is close to the discount factor to t_{i+1} , for an RFR such as SOFR, this is close to the discount factor to t_i). Φ is the cumulative distribution function of the standard normal distribution, and ϕ is the probability density function of the standard normal distribution.

A standard 1Y USD Cap in the Libor world (on 3M Libor) has 3 caplets, the first one maturing in 3 months from today with payment 6 months from today. In the RFR world, a 1Y USD Cap on 3M SOFR has 4 caplets, the first one maturing in 3 months from today with payment¹ 3 months from today. Although

¹ There is typically an additional 2 business days payment delay.

forward looking rates (as was Libor) are less popular these days, we will only consider forward looking rates in this article. While for backward looking rates, the inclusion of a linear decay in the caplet volatilities during stripping (but not for quoting the cap price) may be important [1,2], we believe our observations presented here in the forward looking rate context still hold.

The goal of caplet stripping is to find the caplet volatilities $\sigma_{i,K}$ that are consistent with the market prices for a set of caps with increasing maturities. There are many ways to perform caplet stripping [3]. We will focus here on exact or nearly exact stripping, meaning that the caplet volatilities are chosen such that the model price of the caps matches the market price exactly or with a very small tolerance. Smoothness of the caplet volatilities may be desirable, especially since a discontinuity in caplet volatilities may lead to arbitrage opportunities and issues in the Theta sensitivity of the cap price. We will show that some choices of interpolation and node placement can lead to strong oscillations in the caplet volatilities, and we will propose remedies to alleviate this issue.

2. Stripping Methodology

2.1. Bootstrapping

We have N cap vol quotes $(\hat{\sigma}_1, \dots, \hat{\sigma}_N)$ at maturities $T_1 < \dots < T_N$, each defining a target cap price

$$P_q^{\text{mkt}} = \sum_{i=1}^{M_q} \text{Caplet}_i(\hat{\sigma}_q), \quad M_q = T_q/\Delta,$$

where Δ is the caplet frequency (e.g. 1 month). We parameterise the caplet vol curve by N node values $(v_1, \dots, v_N)$ at node times $(\tau_1, \dots, \tau_N)$, with an interpolation $\sigma(t) = I(t; \tau, v)$. The stripping problem is: find v such that

$$P_q^{\text{model}}(v) = \sum_{i=1}^{M_q} \text{Caplet}_i(\sigma(t_i)) = P_q^{\text{mkt}}, \quad q = 1, \dots, N.$$

The most common approach is to use a bootstrapping method, where the implied volatility of each caplet is determined sequentially, starting from the shortest maturity, along with a piecewise constant interpolation. A one dimensional root solver is used to find the caplet volatility that matches the market price of the cap.

2.2. Numerical Example

We consider the cap volatilities on Libor-1M as of February 2022, as shown in Table A1 with USD OIS discount curve from Table A2. We perform caplet stripping using linear interpolation with at-maturity nodes and midpoint nodes.

A classic bootstrap with piecewise-constant interpolation leads to Figure 1. The standard piecewise-constant (step-forward) interpolation of the caplet vol term structure produces sharp discontinuities at every node boundary. While the bootstrap is exact and sequential, the resulting vol curve is visually unappealing and can cause numerical issues in applications that evaluate the curve at non-caplet times (e.g. intra-month pricing or Theta like risk).

2.3. Child's Play

A key observation is that we calibrate only towards discrete points, and the interpolation between these points is not constrained by the market. If we remove any interpolation and just plot the node values for a typical bootstrap (assuming the caplet volatilities for maturities $T_{k-1} \leq t_i < T_k$ to be the same), we end up with Figure 2.




Figure 1. Caplet volatilities obtained with bootstrapping and piecewise-constant interpolation. The cap vols appear shifted by 1M because a cap maturity of 3M actually has its last fixing at 2M for a forward looking rate with 1M tenor.




Figure 2. Caplet volatilities after bootstrapping, zoom on the 6M-30M caplets.

How would a child fill in the gaps between the dots?

A child would simply join the dots with a line. This leads to the *flat-linear* interpolation: at each node boundary, instead of an instantaneous jump the vol transitions linearly over a short ramp of width $\beta\Delta$. Crucially, if $0 < \beta < 1$, all caplet maturities see exactly the same vol as the piecewise-constant scheme, so the sequential bootstrap produces identical node values.

Let Δ be the caplet frequency (e.g. $\Delta = 1$ month) and $1 \geq \beta > 0$ a ramp factor, we define $\bar{\beta} = \beta\Delta$ as the ramp width. A cap on a *Forward-looking* rate (e.g. LIBOR) of maturity T_k contains caplets fixing at $\Delta, 2\Delta, \dots, T_k - \Delta$. The last caplet fixes one period before T_k and pays at T_k . A cap on a *Backward-looking* rate (e.g. SOFR) of maturity T_k contains caplets fixing at $\Delta, 2\Delta, \dots, T_k$.

Let τ_k be the node times, for forward-looking rates the node is placed at the last fixing time of each cap, $\tau_k = T_k - \Delta$ while for backward-looking rates, $\tau_k = T_k$. The piecewise-constant interpolation is left-continuous on τ_k and reads

$$\sigma(t) = \sum_{q=1}^{M_q} \sigma_{q,K} 1_{t \in (\tau_{q-1}, \tau_q]} + \sigma_{M_q,K} 1_{t \in [\tau_{M_q}, \infty)},$$

with $\tau_0 = 0$.

Let $v_q = \sigma_{q,K}$ be the node value (the caplet normal vol), the flat-linear interpolant $\sigma(t)$ is defined as follows.

Boundary cases:

$$\sigma(t) = v_1 \quad \text{for } t \leq \tau_1, \quad \sigma(t) = v_N \quad \text{for } t \geq \tau_N.$$

Interior:

For t in the open interval (τ_{k-1}, τ_k) with $k \geq 2$, define the ramp centre

$$c_k = \tau_{k-1} + \frac{\Delta}{2}$$

and the ramp boundaries (clamped to the interval)

$$a_k = \max(\tau_{k-1}, c_k - \frac{\tilde{\beta}}{2}), \quad b_k = \min(\tau_k, c_k + \frac{\tilde{\beta}}{2}).$$

The interpolant is

$$\sigma(t) = \begin{cases} v_{k-1}, & \tau_{k-1} < t \leq a_k, \\ v_{k-1} + \frac{t - a_k}{b_k - a_k} (v_k - v_{k-1}), & a_k < t \leq b_k, \\ v_k, & b_k < t < \tau_k. \end{cases} \quad (2)$$

At the nodes themselves the step-forward convention gives $\sigma(\tau_k) = v_k$.

Interpretation:

On each interval $(\tau_{k-1}, \tau_k]$ the vol curve has three regimes:

1. A *flat* region at v_{k-1} from τ_{k-1} to a_k .
2. A *ramp* from a_k to b_k , centred at $c_k = \tau_{k-1} + \Delta/2$, during which the vol transitions from v_{k-1} to v_k .
3. A *flat* region at v_k from b_k to τ_k .

When $\beta = 1$, the ramp runs from τ_{k-1} to $\tau_{k-1} + \Delta$, leaving any remaining interval flat at v_k .

Figure 3 shows the impact of the ramp width β on the caplet volatilities.




Figure 3. Caplet volatilities with flat-linear interpolation and various ramp width β .

2.4. What might a draughtsman do?

The *flat-smooth* interpolant replaces the linear ramp with the cubic Hermite smoothstep $\Psi(s) = 3s^2 - 2s^3$, using the same centred placement:

$$\sigma(t) = \begin{cases} v_{k-1}, & \tau_{k-1} < t \leq a_k, \\ v_{k-1} + \Psi\left(\frac{t - a_k}{b_k - a_k}\right) (v_k - v_{k-1}), & a_k < t \leq b_k, \\ v_k, & b_k < t < \tau_k, \end{cases} \quad (3)$$

where $\Psi(s) = 3s^2 - 2s^3$.

Because $\Psi'(0) = \Psi'(1) = 0$, the resulting curve is C^1 (no slope discontinuities), while preserving bootstrap equivalence for $\beta \leq 1$.

Figure 4 shows the flat-smooth interpolation with $\beta = 1$.




Figure 4. Caplet volatilities with flat-smooth interpolation with $\beta = 1$.

2.5. Bootstrap equivalence

Proposition 1. *If $\beta \leq 1$, the flat-linear (or flat-smooth) interpolant evaluates to the same value as the piecewise-constant interpolant at every caplet fixing date. Consequently, sequential bootstrap produces identical node values.*

Proof. Caplet fixing times are integer multiples of Δ , and nodes are placed at cap boundaries (also multiples of Δ). The ramp centre is $c_k = \tau_{k-1} + \Delta/2$. For any caplet fixing time $t = \tau_{k-1} + j\Delta$ with $j \geq 1$ falling inside (τ_{k-1}, τ_k) , we have $t - c_k = (j - \frac{1}{2})\Delta \geq \Delta/2 \geq \tilde{\beta}/2$, so $t \geq b_k$ and $\sigma(t) = v_k$. At node times $t = \tau_k$ the step-forward convention gives $\sigma(t) = v_k$ directly. $\square$

2.6. Limiting case

- $\beta \rightarrow 0$: the ramp vanishes and $\sigma \rightarrow \sigma_{\text{flat}}$ (piecewise constant).
- $\beta = 1$ ($\tilde{\beta} = \Delta$): the ramp runs from τ_{k-1} to $\tau_{k-1} + \Delta$. The first caplet fixing inside the interval is at $\tau_{k-1} + \Delta = b_k$, where $\sigma = v_k$. Bootstrap equivalence holds.
- $\beta > 1$: the ramp boundaries are clamped to the interval (τ_{k-1}, τ_k) , so the ramp cannot extend further than the interval allows. Bootstrap is not applicable, a global solver is required.

2.7. Continuity

The flat-linear interpolant is continuous everywhere. At each node τ_k :

- *Left of τ_k* : t is in the post-ramp flat region of interval (τ_{k-1}, τ_k) (since $b_k = c_k + \bar{\beta}/2 \leq \tau_{k-1} + \Delta \leq \tau_k$), so $\sigma(\tau_k^-) = v_k$.
- *Right of τ_k* : t is in the pre-ramp flat region of interval (τ_k, τ_{k+1}) (since $a_{k+1} = c_{k+1} - \bar{\beta}/2 = \tau_k + (\Delta - \bar{\beta})/2 \geq \tau_k$), so $\sigma(\tau_k^+) = v_k$.

Both limits equal $\sigma(\tau_k) = v_k$, so σ is **globally continuous** for any $\beta \in [0, 1]$. The flat-smooth variant is additionally C^1 since the smoothstep has zero derivative at both ramp endpoints.

2.8. Generalisation: Compact Kernel Transitions

The flat-linear ramp is equivalent to convolving each step-function jump with a rectangular kernel of width $\bar{\beta}$. This generalises to any compact-support kernel K with $\int_0^1 K(s) ds = 1$. Define the transition function

$$\Psi(s) = \int_0^s K(u) du, \quad s \in [0, 1],$$

so that $\Psi(0) = 0$ and $\Psi(1) = 1$. The interpolant becomes

$$\sigma(t) = v_1 + \sum_{k \geq 2} (v_k - v_{k-1}) \Psi\left(\frac{t - a_k}{b_k - a_k}\right),$$

with Ψ clamped to $[0, 1]$ outside $[0, 1]$.

The flat-linear scheme uses $K \equiv 1$ (rectangular) giving $\Psi(s) = s$. The flat-smooth scheme uses the Hermite smoothstep kernel.

Kernel	Transition $\Psi(s)$	Smoothness
Rectangular ($K = 1$)	s	C^0 (flat-linear)
Hermite smoothstep	$3s^2 - 2s^3$	C^1 , $\Psi'(0) = \Psi'(1) = 0$
Cosine (Hann)	$\frac{1}{2}(1 - \cos \pi s)$	C^1
Quintic smotherstep	$6s^5 - 15s^4 + 10s^3$	C^2

3. Smoother Interpolations

3.1. Node Placement and Interpolation

Being overly enthusiastic about joining the dots, one may consider to directly interpolate linearly between the nodes, placed at the cap maturities. There is no dependency towards the next caplet (the matrix is lower triangular) and thus bootstrapping is still applicable. However, this approach can lead to strong oscillations in the caplet volatilities, especially for longer maturities. We will show that using midpoint nodes instead of at-maturity nodes can help reduce these oscillations. We will thus consider two kinds of node set:

- *At-maturity* nodes set $\tau_k = T_k - \delta$.
- *Midpoint* nodes set $\tau_1 = T_1 - \delta$ and $\tau_k = (T_{k-1} + T_k)/2 - \delta$ for $k \geq 2$.

The shift δ is typically one month for a forward looking rate with 1 month tenor, and zero for a backward looking rate.

The introduction of midpoint nodes leads to a non lower triangular matrix, and thus requires a global solver to find the caplet volatilities. We use the Levenberg-Marquardt method with backtracking line search as global solver [4]. With the global solver, we are not limited to piecewise linear interpolation, and we can also use a cubic spline interpolation.

3.2. Enforcing Positivity

A priori, there is no guarantee that the linear interpolation at midpoints will lead to positive caplet volatilities. In our example, the linear interpolation at mid-points actually leads to negative caplet volatilities

around 18 months. While we can just floor the negative volatilities to zero as in Figure 6, it is a better idea to enforce positivity during the optimization by using a change of variable $v_k = \exp(u_k)$ where u_k is the optimization variable. This way, the caplet volatilities are always positive, and we can still use the global solver to find the optimal u_k values.

For cubic splines, Dougherty *et al.* [5] propose a method to enforce non-negativity of the spline by filtering the first derivatives around offending nodes (while still imposing C1 continuity conditions). In our implementation, we use a C1 cubic spline based on Bessel (parabolic) estimate of derivatives along with the non-negative filter instead of a C2 cubic spline. We pay attention to use the corrected version for the filter coefficients as described in [6].

Figure 5 shows that both approaches work. With the exponential change of variable, the linear interpolation at midpoints does not reach zero anymore. The C2 cubic spline presents a small oscillation, while the C1 cubic spline with non-negative filter does not present any oscillation and is smoother than the linear interpolation with exponential transform.




Figure 5. Caplet volatilities obtained with non-negative constraint and midpoints nodes.

Table 1. Cap repricing error for different interpolation methods.

Method	Min vol (bp)	Min node	Reprice err
flat at mat.	39.75	39.75	9.12e-15
linear at mat.	0.0	-158.57	9.12e-15
linear mid	0.0	-13.87	8.21e-16
cubic mid	15.11	15.11	7.88e-15
hyman mid	4.95	4.95	5.93e-16
hyman mid floor=10 (shift)	10.0	10.0	4.14e-02
linear exp mid	19.06	19.06	4.93e-16
cubic exp mid	27.69	28.64	7.42e-16

4. Oscillations and Blow-ups

The resulting caplet volatilities obtained with linear interpolation and at-maturity nodes show strong oscillations, with a node near 18 months settling at -159 bp. With midpoint nodes, the oscillations are much reduced, and the node near 18 months settles at -14 bp instead of -159 bp (Figure 6). The oscillations grow stronger for longer maturities. The cubic spline interpolation exhibits even stronger oscillations.




Figure 6. Caplet volatilities obtained with linear interpolation at maturity nodes exhibit strong oscillations.

4.1. The Reason for the Oscillations

With linear interpolation and at-maturity nodes $\tau_k = T_k$, the vol at caplet time $t_i \in (T_{k-1}, T_k]$ is

$$\sigma(t_i) = v_{k-1} \frac{T_k - t_i}{T_k - T_{k-1}} + v_k \frac{t_i - T_{k-1}}{T_k - T_{k-1}}.$$

The sequential bootstrap solves for v_k given $(v_1, \dots, v_{k-1})$ already fixed. The incremental constraint (cap k minus cap $k-1$) involves caplets $i = M_{k-1}+1, \dots, M_k$ whose vols depend on v_k with weight

$$w_i = \frac{t_i - T_{k-1}}{T_k - T_{k-1}}, \quad w_i \in \left[\frac{\Delta}{T_k - T_{k-1}}, 1 \right].$$

The sensitivity of the incremental price to v_k is

$$\frac{\partial \Delta P_k}{\partial v_k} = \sum_{i=M_{k-1}+1}^{M_k} \frac{\partial \text{Caplet}_i}{\partial \sigma} \cdot w_i.$$

Since w_i increases linearly from ≈ 0 to 1, the average weight is $\bar{w} \approx 1/2$ which is roughly **half** the leverage of flat interpolation (where $w_i = 1$ for all i). When the target incremental vol drops sharply (e.g. the 2Y dip), v_k must *overshoot* to compensate for the diluted sensitivity. In our numerical example, node values of -175 bp emerge even though the target incremental vol is merely low, not negative.

4.2. Why midpoints help

With midpoint nodes $\tau_k = (T_{k-1} + T_k)/2$, the cap maturity T_q falls near the right edge of interval $[\tau_q, \tau_{q+1}]$ rather than coinciding with a node. Crucially, the Jacobian $J_{qk} = \partial P_q / \partial v_k$ is no longer lower-triangular: each node v_k has substantial influence on at least two cap prices. The global Newton system

$$J \Delta v = -r, \quad r_q = P_q^{\text{model}} - P_q^{\text{mkt}},$$

distributes corrections bilaterally across nodes.

In our numerical example, Node 8 (near 18M) settles at -19 bp instead of -175 bp because the optimizer can simultaneously adjust neighbouring nodes to absorb the 2Y dip.

5. The Real Issue With Our Example: The 2Y Dip

The quality of our cap quotes is in fact not great. The 2Y cap is very cheap compared to the 1Y and 3Y caps, which creates a dip in the caplet volatilities around 18 months. It is unlikely to be real, but may stem from a stale quote.

5.1. Modified Z-Score

Given a series of observations $x_1, \dots, x_n$, the *median absolute deviation* (MAD) is

$$\text{MAD} = \text{median}(|x_i - \tilde{x}|), \quad \tilde{x} = \text{median}(x_i).$$

Unlike the standard deviation, the MAD has a breakdown point of 50%: up to half the observations can be arbitrarily corrupted without affecting it.

To make the MAD comparable to the standard deviation for normally distributed data, Hampel [7] introduced the consistency factor $c = 1/\Phi^{-1}(3/4) \approx 1.4826$, so that $\hat{\sigma} = c \cdot \text{MAD}$ is a consistent estimator of σ .

The *modified Z-score* of Iglewicz and Hoaglin [8] is defined as

$$M_i = \frac{0.6745(x_i - \tilde{x})}{\text{MAD}}, \quad (4)$$

where $0.6745 = 1/c = \Phi^{-1}(3/4)$. This normalises the score so that $|M_i| \approx |z_i|$ when the data are Gaussian.

5.2. Outlier Criterion

An observation x_i is flagged as an outlier when

$$|M_i| > k.$$

Iglewicz and Hoaglin [8] recommend $k = 3.5$ as a conservative threshold. The choice $k = 3$ is also widely used and corresponds (for Gaussian data) to roughly the same false-positive rate as a 3σ rule applied with the classical mean and standard deviation.

5.3. Application to Cap Vol Stripping

In the cap vol stripping context, we compute residuals of each cap vol quote relative to a local median (e.g. a rolling window of five neighbouring quotes):

$$r_q = \hat{\sigma}_q - \text{median}(\hat{\sigma}_{q-2}, \dots, \hat{\sigma}_{q+2}).$$

The modified Z-score (4) is then applied to the residual series r_q . Quotes with $|M_q| > 3$ are excluded before stripping, and the remaining “clean” quotes are used to bootstrap caplet vols.

Running this logic on our example, we actually find 2 outliers corresponding to $T_2 = 3\text{M}$ and $T_8 = 2\text{Y}$. Figure 7 shows the resulting interpolated caplet volatilities with outliers removed. We clearly see that the oscillations are much reduced, but are still there when using linear or cubic interpolation at maturities. Midpoint node placement is very effective. The non-negative Hyman interpolation is close to the linear interpolation but much smoother.

5.4. Total Variance Check

Another ad-hoc criteria may be to compute the total variance of the cap $\hat{\sigma}_q^2 T_q$ (see Listing 1). In particular, we notice that it is not increasing around those two outliers. This does not mean that there is a calendar spread arbitrage opportunity, since caps of different maturities are not directly comparable as each caplet operates on a different forward rate. However, it is a strong indication that there may be an issue




Figure 7. Caplet volatilities obtained with outliers removed.

with the quotes, and that we should investigate further by looking at the outliers with the modified Z-score method.

6. Conclusion

When using a simple bootstrap algorithm to strip the caplet volatilities, there is no good reason to use a pure piecewise-constant interpolation. A flat-linear interpolation is always preferable as it produces a continuous vol curve. For smoother interpolations, the use of midpoint nodes instead of at-maturity nodes help to reduce oscillations significantly. In particular, we found the Hyman non-negative C1 spline to work well. Finally, it is important to check the quality of the cap quotes before stripping, as outliers can lead to strong oscillations in the caplet volatilities and we have provided a practical way of doing so.

Acknowledgments: I am grateful to Gary Kennedy for his precious feedback.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

1. Piterbarg, V. Interest rates benchmark reform and options markets. *Available at SSRN 3537925* **2020**.
2. Piterbarg, V. Benchmark reform goes non-linear. *Risk. net* **2020**, pp. 1–5.
3. White, R.; Iwashita, Y. Eight ways to strip your caplets: An introduction to caplet stripping. *OpenGamma Quantitative Research* **2014**.
4. Nocedal, J.; Wright, S.J. *Numerical optimization*; Springer, 2006.
5. Dougherty, R.L.; Edelman, A.S.; Hyman, J.M. Nonnegativity-, monotonicity-, or convexity-preserving cubic and quintic Hermite interpolation. *mathematics of computation* **1989**, *52*, 471–494.
6. Le Floc'h, F. Typo in Hyman non-negative constraint - 28 years later. <https://chasethedevil.github.io/post/typo-in-hyman-non-negative-constraint/>, 2017. Accessed: 2026-05-05.
7. Hampel, F.R. The influence curve and its role in robust estimation. *Journal of the american statistical association* **1974**, *69*, 383–393.
8. Iglewicz, B.; Hoaglin, D. Volume 16: how to detect and handle outliers, The ASQC basic references in quality control: statistical techniques, Edward F. Mykytka. *Ph. D. dissertation, Ph. D., Editor* **1993**.

Table A1. cap volatilities (in basis points) on Libor-1M as of February 2022 for a strike $K = 0\%$.

Maturity (months)	2	3	4	5	6	9	12	24	36	60	84	120	180
Cap Vol (bp)	79.30	111.83	93.10	91.83	94.51	102.25	99.86	60.42	109.15	96.34	84.37	76.90	76.34

Table A2. USD OIS discount curve as of February 2022.

Maturity (months)	0	1	2	3	6	12	24	36	60	84	120	180	240	360
Zero Rate (%)	0.05	0.06	0.20	0.35	0.69	1.01	1.44	1.62	1.71	1.81	1.83	2.04	2.25	2.17

Appendix A Market Data

Appendix B Total Variance Check

```

cap_maturities_months = Float64.([2, 3, 4, 5, 6, 9, 12, 24, 36, 60, 84, 120, 180])
cap_vols_bp = [79.30, 111.83, 93.10, 91.83, 94.51, 102.25, 99.86, 60.42,
109.15, 96.34, 84.37, 76.90, 76.34]

T = cap_maturities_months ./ 360
tv = (cap_vols_bp .^ 2) .* T
arb_violations = findall(diff(tv) .< 0)
println("\nTotal-variance arbitrage violations at indices: ", arb_violations)

```

Listing 1. Total variance check for arbitrage detection in *Julia* language.

Appendix C Outlier Detection

```

ref = [median(cap_vols_bp[max(1,i-2):min(end,i+2)]) for i in eachindex(cap_vols_bp)]
res = cap_vols_bp .- ref
mad_val = median(abs.(res .- median(res)))
mod_z = 0.6745 .* (res .- median(res)) ./ mad_val
outlier_idx = findall(abs.(mod_z) .> 3.0)
println("\nOutlier indices: ", outlier_idx)

```

Listing 2. Detect outliers in the cap volatilities using the MAD method in *Julia* language.