

First-Order Efficiency for Probabilistic Value Estimation via A Statistical Viewpoint

Ziqi Liu¹, Kiljae Lee², Yuan Zhang², Weijing Tang¹ *

¹Department of Statistics & Data Science, Carnegie Mellon University.

²Department of Statistics, The Ohio State University.

Abstract

Probabilistic values, including Shapley values and semivalues, provide a model-agnostic framework to attribute the behavior of a black-box model to data points or features, with a wide range of applications including explainable artificial intelligence and data valuation. However, their exact computation requires utility evaluations over exponentially many coalitions, making Monte Carlo approximation essential in modern machine learning applications. Existing estimators are often developed through different identification strategies, including weighted averages, self-normalized weighting, regression adjustment, and weighted least squares. Our key observation is that these seemingly distinct constructions share a common first-order error structure, in which the leading term is an augmented inverse-probability weighted influence term determined by the sampling law and a working surrogate function. This first-order representation yields an explicit expression for the leading mean squared error (MSE), which characterizes how the sampling law and the surrogate jointly determine statistical efficiency. Guided by this criterion, we propose an *Efficiency-Aware Surrogate-adjusted Estimator (EASE)* that directly chooses the sampling law and surrogate to minimize the first-order MSE. We demonstrate that EASE consistently outperforms state-of-the-art estimators for various probabilistic values.

Keywords: Cooperative games; Shapley value; Monte Carlo sampling; Regression adjustment.

*Corresponding author email: weijingt@andrew.cmu.edu.

1 Introduction

Shapley values [30], and the broader class of probabilistic values [10, 32], provide an axiomatic framework for quantifying the contribution of individual players in a cooperative game. This framework has become increasingly prominent in modern machine learning, including model-agnostic data attribution, where the goal is to understand how individual training examples affect model behavior, and explainable artificial intelligence, where the goal is to assess how input features contribute to a black-box model output [15, 17, 23, 31]. In these applications, data points or features are treated as players in a cooperative game. For a game with n players, the utility of a coalition is represented by a function $u : 2^{[n]} \rightarrow \mathbb{R}$, where $[n] = \{1, \dots, n\}$, and $u(S)$ measures the performance of a coalition $S \subseteq [n]$, such as the prediction accuracy of a model trained on the data points in S . Given u , a probabilistic value assigns to each player $i \in [n]$ a weighted average of its marginal contributions over all coalitions that do not contain i :

$$\phi_i(u) = \sum_{S \subseteq [n] \setminus \{i\}} \alpha_i^{(n)}(S) \{u(S \cup \{i\}) - u(S)\}, \quad (1)$$

where the weights are nonnegative and satisfy $\sum_{S \subseteq [n] \setminus \{i\}} \alpha_i^{(n)}(S) = 1$. When these weights depend only on the size of the coalition $|S|$, rather than on the identities of the players in S , the probabilistic value reduces to a semivalue [10]. As a further special case, when $\alpha_i^{(n)}(S) = \{n \binom{n-1}{|S|}\}^{-1}$, it becomes the Shapley value. Because the framework treats the utility function as a black box, it can be applied across a wide range of utility specifications.

Despite this model-agnostic appeal, exact computation of probabilistic values is generally infeasible. The definition in (1) requires evaluating the utility function over all 2^n coalitions, and in modern machine learning applications, a single utility evaluation may require retraining a model on a selected subset of data or running a costly inference procedure.

Exact computation therefore becomes impractical even for moderately large n .

The sampling-based approximation has therefore become the standard approach, and a broad literature has developed a range of Monte Carlo estimators for probabilistic values; see, for example, the surveys [4, 9]. These methods can be organized by how they identify the target under a sampling distribution. One representative family uses weighted-average identification, which directly represents the target as a weighted average of coalition utilities. This family includes weighted sample means [6, 11], self-normalized weighted sample means [16, 20, 31], and, more recently, regression-adjusted estimators [33]. Another widely used family exploits weighted least-squares (WLS) identification, which obtains the target through a projection of the utility function onto a structured feature space. This family includes empirical WLS estimators for Shapley values [5, 12, 23, 26] and for general semivalues [19].

Our key observation is that these seemingly different estimator constructions share a common first-order error structure. For both weighted-average and least-squares estimators, the first-order error term can be written as an augmented inverse-probability-weighted influence term indexed by the sampling law and a working surrogate function. This surrogate may be specified explicitly, as in regression-adjusted estimators, or induced implicitly, as in self-normalized or WLS estimators. This representation therefore provides a unified first-order characterization of many existing estimators and, more importantly, enables an explicit expression for the leading mean squared error (MSE).

In particular, following the first-order expansion, if n is fixed and $m \rightarrow \infty$, the leading term is the variance of the corresponding residual influence term divided by the Monte Carlo sample size m . For probabilistic-value estimation, however, this is not the primary regime of interest. The utility is defined on 2^n coalitions, whereas feasible sampling budgets are typically much smaller than 2^n , often only polynomial in n . Thus, the relevant question is

whether the first-order term already dominates the remainder with the polynomial sample sizes used in practice. To address this gap, we establish non-asymptotic remainder bounds for representative weighted-average and WLS estimators. Our results show that the first-order MSE approximation has a small relative error once $m = \Omega(n \text{polylog}(n))$, aligning with the regime studied in the literature [19, 20, 26]. This validation turns the first-order MSE expression into a practical design criterion, which makes explicit how the sampling law and the surrogate choice jointly determine statistical efficiency.

Motivated by this criterion, we propose the *Efficiency-Aware Surrogate-adjusted Estimator (EASE)*, which directly targets the first-order MSE. The optimization problem involves two coupled design choices: fitting the surrogate and choosing the sampling law. Since the objective depends on the unknown utility function, both components must be learned from sampled utility evaluations. Moreover, without structural restrictions, choosing the sampling law would require optimizing over all distributions on $2^{[n]}$, which is computationally infeasible for even moderate n . EASE makes this design problem tractable through a two-stage procedure. The initialization stage uses pilot samples to fit an initial surrogate and construct a residual-aware sampling law within a structured class. The estimation stage then draws samples from this learned law and constructs a cross-fitted augmented inverse-probability-weighted estimator, with the surrogate chosen to minimize the empirical first-order MSE criterion. Numerical results demonstrate that EASE consistently outperforms state-of-the-art estimators for a range of probabilistic-value targets.

The rest of the paper is organized as follows. Section 2 formulates the probabilistic-value estimation problem and reviews the weighted-average and WLS identifications. Section 3 develops the common first-order representation through working surrogates. Section 4 derives the first-order MSE criterion and non-asymptotic remainder bounds. We then introduce EASE and its two-stage sampling and estimation procedure in Section 5, followed

by extensive numerical experiments in Section 6.

2 Problem Setup for Probabilistic Value Estimation

In this section, we frame Monte Carlo estimation problem for a broad class of probabilistic-value targets. This framework covers the estimation of an individual probabilistic value as well as more general aggregate targets. Let $\boldsymbol{\phi}(u) = (\phi_1(u), \dots, \phi_n(u))^\top$, where $\phi_i(u)$ is the probabilistic value for each player i defined in (1). For any coefficient vector $\mathbf{a} = (a_1, \dots, a_n)^\top \in \mathbb{R}^n$, define

$$\tau_{\mathbf{a}}(u) := \mathbf{a}^\top \boldsymbol{\phi}(u) = \sum_{i=1}^n a_i \phi_i(u). \quad (2)$$

Let $\mathbf{e}_i \in \mathbb{R}^n$ denote the i -th standard basis vector. Then the special case $\mathbf{a} = \mathbf{e}_i$ recovers the probabilistic value of player i . More generally, this formulation also covers downstream summaries such as group-level totals. For example, if $\mathbf{a}$ is the indicator vector of a subset of players, then $\tau_{\mathbf{a}}(u)$ is the total probabilistic value over that subset, which is useful in group data attribution applications [18]. For notational simplicity, we focus on scalar targets, but the framework can be extended to vector-valued targets.

We estimate $\tau_{\mathbf{a}}(u)$ from Monte Carlo sampling. We first focus on the single-coalition sampling scheme and later extend the same logic to structured sampling designs in Appendix E. Let q be a sampling distribution on $2^{[n]}$, and suppose that we observe i.i.d. sampled coalitions $S_1, \dots, S_m \sim q$, together with their exact utility values. The observed data are therefore $\mathcal{D}_m = \{(S_t, u(S_t))\}_{t=1}^m$. Single-coalition sampling is widely used in practice because of its flexibility. Once the coalitions have been sampled and their utility values evaluated, the same evaluations can be reused for different targets. Changing the target from $\tau_{\mathbf{a}}$ to $\tau_{\mathbf{b}}$ only changes the coefficient vector from $\mathbf{a}$ to $\mathbf{b}$.

Given a sampling design, the target $\tau_{\mathbf{a}}(u)$ can often be identified as a functional of the observed data distribution in more than one way. Different identifications lead to different Monte Carlo estimators. We next revisit two common identifications that underlie many consistent estimators for probabilistic-value targets.

1. *Weighted Average Identification.* Probabilistic values are linear functionals of the utility function u . For each player i , define $\rho_i(S) := \alpha_i^{(n)}(S \setminus \{i\})\mathbf{1}\{i \in S\} - \alpha_i^{(n)}(S)\mathbf{1}\{i \notin S\}$, for $S \subseteq [n]$. Then $\phi_i(u) = \sum_{S \subseteq [n]} \rho_i(S)u(S)$. Therefore, any linear combination $\tau_{\mathbf{a}}(u)$ defined in (2) also admits the representation

$$\tau_{\mathbf{a}}(u) = \sum_{S \subseteq [n]} \rho_{\mathbf{a}}(S)u(S), \quad \rho_{\mathbf{a}}(S) := \sum_{i=1}^n a_i \rho_i(S). \quad (3)$$

Under single-coalition sampling from a distribution q , this becomes

$$\tau_{\mathbf{a}}(u) = \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)u(S)], \quad \gamma_{\mathbf{a},q}(S) := \frac{\rho_{\mathbf{a}}(S)}{q(S)}, \quad (4)$$

provided that $q(S) > 0$ whenever $\rho_{\mathbf{a}}(S) \neq 0$. Thus, the target is identified as an expectation under the sampling distribution q . This identification naturally motivates estimators that replace the expectation in (4) by its empirical counterpart, such as weighted sample means [6, 11], self-normalized weighted sample means [16, 20, 31], and regression-adjusted methods [33].

2. *Weighted Least-Squares Identification.* For Shapley values [5, 12, 23, 26] and more general semivalues [19], the target can also be characterized through a WLS problem:

$$\tau_{\mathbf{a}}(u) = \mathbf{c}_{\mathbf{a}}^\top \boldsymbol{\beta}^*(u), \quad \text{for every } \boldsymbol{\beta}^*(u) \in \arg \min_{\boldsymbol{\beta} \in \mathbb{R}^{d_{\mathbf{z}}}} \sum_{S \subseteq [n]} w(S) \{u(S) - \mathbf{z}(S)^\top \boldsymbol{\beta}\}^2, \quad (5)$$

where $\mathbf{c}_{\mathbf{a}}$ is a known vector that extracts the target from the fitted coefficient vector,

$w : 2^{[n]} \rightarrow [0, \infty)$ is a known nonnegative weight function, and $\mathbf{z} : 2^{[n]} \rightarrow \mathbb{R}^{d_z}$ is a particular feature map. In this characterization, the tuple $(w, \mathbf{z}, \mathbf{c}_a)$ determines the target. This tuple is not necessarily unique: different choices of weights, feature maps, readout vectors, and constraints may lead to the same target. For example, KernelSHAP [23] and PolySHAP [12] both identify Shapley values, but they use different WLS characterizations. Under single-coalition sampling from q , the same population criterion can be written as:

$$\tau_a(u) = \mathbf{c}_a^\top \boldsymbol{\beta}^*(u), \quad \text{for every } \boldsymbol{\beta}^*(u) \in \arg \min_{\boldsymbol{\beta} \in \mathbb{R}^{d_z}} \mathbb{E}_q \left[\frac{w(S)}{q(S)} \{u(S) - \mathbf{z}(S)^\top \boldsymbol{\beta}\}^2 \right]. \quad (6)$$

Thus, the target is no longer represented directly as a weighted average of $u(S)$. Instead, u is first projected onto a structured linear class and the target is then obtained by applying the readout $\mathbf{c}_a$. This identification motivates estimators that replace the population WLS criterion by its finite-sample counterpart and then apply the same readout to the fitted coefficient vector.

Toward a unified estimator-design view. The two identifications above lead to estimators with seemingly different structures. The weighted-average identification motivates estimators that reweight sampled utilities, possibly with regression adjustments. The weighted least-squares identification, in contrast, motivates estimators that solve a weighted regression problem and then map the fitted coefficients to the target. Thus, even for the same target $\tau_a(u)$ and the same sampled data $\mathcal{D}_m$, the resulting estimators can appear to rely on different principles. As we shall show in the next section, these constructions share a common first-order structure. Up to first-order terms, each can be viewed as an augmented weighted-average estimator indexed by a surrogate. The surrogate may be either specified explicitly, as in regression adjustment, or induced implicitly by self-normalization

or least-squares fitting. Under this view, differences among existing estimator constructions can often be traced to the *working surrogate classes* they use, either explicitly or implicitly, for first-order residual adjustment.

3 First-Order Representation via Working Surrogates

In this section, we formalize a first-order representation for a broad class of probabilistic-value estimators. Specifically, we show that many common weighted-average, self-normalized, regression-adjusted, and least-squares estimators admit an expansion whose leading term is an augmented weighted average indexed by a surrogate function h . The corresponding working surrogate class describes the set of first-order adjustments attainable by the estimation procedure. Let $\mathcal{H}$ be a working surrogate class consisting of functions $h : 2^{[n]} \rightarrow \mathbb{R}$. For each m , let $S_1, \dots, S_m \stackrel{\text{iid}}{\sim} q$, and let $\hat{\tau}_{\mathbf{a},m}$ denote an estimator based on these samples. We define regularity relative to $\mathcal{H}$ under the sampling law q through a common first-order error expansion and then place common estimators within this class.

Definition 1 (Regular estimator sequence). A sequence of estimators $\{\hat{\tau}_{\mathbf{a},m}\}_{m \geq 1}$ for $\tau_{\mathbf{a}}(u)$ is *regular* relative to $\mathcal{H}$ if there exists a sequence of nonrandom functions $h_m \in \mathcal{H}$ such that

$$\hat{\tau}_{\mathbf{a},m}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{j=1}^m \psi_{h_m}(S_j; u) + r_m \quad \text{with } \mathbb{E}_q(r_m^2) = o(m^{-1}), \quad (7)$$

where $\psi_h(S; u) := \gamma_{\mathbf{a},q}(S)\{u(S) - h(S)\} + \tau_{\mathbf{a}}(h) - \tau_{\mathbf{a}}(u)$.

Example 1 (Horvitz–Thompson estimator). The simplest example is the Horvitz–Thompson estimator [14] motivated by the weighted-average identification (4):

$$\hat{\tau}_{\mathbf{a},m}^{\text{HT}}(u) = \frac{1}{m} \sum_{t=1}^m \gamma_{\mathbf{a},q}(S_t) u(S_t). \quad (8)$$

By the definition that $\psi_0(S; u) = \gamma_{\mathbf{a},q}(S)u(S) - \tau_{\mathbf{a}}(u)$, we have $\hat{\tau}_{\mathbf{a},m}^{\text{HT}}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_0(S_t; u)$. Thus, $\hat{\tau}_{\mathbf{a},m}^{\text{HT}}$ is regular relative to $\mathcal{H} = \{0\}$, with $r_m = 0$ for every m . In the literature, this estimator coincides with unbiased KernelSHAP [6] and the order-one special case of SHAP-IQ [11].

Example 2 (Augmented inverse-probability weighted estimator). Another example is the augmented inverse-probability weighted (AIPW) estimator [28]. Let $\mathcal{H}^{\text{sur}}$ be an explicit working surrogate class consisting of functions $h : 2^{[n]} \rightarrow \mathbb{R}$ that can be evaluated at any coalition S and for which $\tau_{\mathbf{a}}(h)$ is efficiently computable (see Remark 1 for further discussion). For a fixed surrogate $h_m \in \mathcal{H}^{\text{sur}}$, define

$$\hat{\tau}_{\mathbf{a},m}^{\text{AIPW}}(u) = \tau_{\mathbf{a}}(h_m) + \frac{1}{m} \sum_{t=1}^m \gamma_{\mathbf{a},q}(S_t) \{u(S_t) - h_m(S_t)\}. \quad (9)$$

Then, by the definition of ψ_{h_m} , we have $\hat{\tau}_{\mathbf{a},m}^{\text{AIPW}}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_{h_m}(S_t; u)$. Thus, $\hat{\tau}_{\mathbf{a},m}^{\text{AIPW}}$ is regular relative to $\mathcal{H}^{\text{sur}}$, with $r_m = 0$ for every m . The regression-adjusted estimator proposed in [33] is equivalent to an AIPW estimator. This is also the with-replacement analogue of the generalized-difference estimator in model-assisted survey sampling [29].

Remark 1 (Explicit Working Surrogate Classes). *This class serves as a set of working models for variance reduction. We do not assume that the true utility function u belongs to $\mathcal{H}^{\text{sur}}$. Instead, the choice of $\mathcal{H}^{\text{sur}}$ is primarily constrained by computational tractability and statistical feasibility. First, for every $h \in \mathcal{H}^{\text{sur}}$, the target $\tau_{\mathbf{a}}(h)$ should be computable without requiring excessive additional utility evaluations. Otherwise, even if h approximates u well enough to reduce variance, exact calculation of the term $\tau_{\mathbf{a}}(h)$ may itself be too costly. Examples include tree ensembles in TreeSHAP [22], RKHS models in RKHS-SHAP [3], and low-degree polynomial classes in PolySHAP [12], for which probabilistic values can be computed in closed form or efficiently approximated. Second, the complexity of $\mathcal{H}^{\text{sur}}$ should be suitably restricted so that the surrogate model can be reliably learned from limited*

sampled coalitions. These desiderata parallel the requirements for auxiliary information used in model-assisted survey sampling: auxiliary variables should have tractable population totals [29], and the working model is used for variance reduction rather than assumed to be correctly specified [7].

The two examples above illustrate estimators in which the surrogate class is specified explicitly. The next two subsections demonstrate that the proposed regular estimator class also covers designs where the working surrogate class is introduced implicitly. This includes estimators whose original forms appear different, such as self-normalized weighted averages and weighted least-squares-based estimators.

3.1 Self-Normalized Weighted-Average Estimators

Several existing stratified weighted-average estimators share a common normalization structure [16, 20, 31]. Although they arise naturally from stratified sampling considerations in their original formulations, we recognize them as instances of a Hájek-type estimator [13]. The main difference across these estimators lies in the partition of the coalition space used for normalization.

To formalize this connection, suppose the coalition space $2^{[n]}$ is partitioned into K cells $\mathcal{C} = \{C_1, \dots, C_K\}$. Denote the design probability of cell C_k by $\pi_k := \mathbb{P}_q(S \in C_k)$ and the empirical frequency of cell C_k in the sample by $\hat{\pi}_k = N_k/m$, where $N_k = \sum_{t=1}^m \mathbf{1}\{S_t \in C_k\}$. Using this partition, the Horvitz–Thompson (HT) estimator in (8) can be rewritten as

$$\hat{\tau}_{\mathbf{a},m}^{\text{HT}}(u) = \sum_{k:N_k>0} \hat{\pi}_k \left\{ \frac{1}{N_k} \sum_{t:S_t \in C_k} \gamma_{\mathbf{a},q}(S_t) u(S_t) \right\}.$$

This representation reveals that the HT estimator weights each within-cell average by the *random* empirical frequency $\hat{\pi}_k$. A Hájek-type, or post-stratified, estimator improves upon

this by replacing this random frequency with the known design probability π_k :

$$\hat{\tau}_{\mathbf{a},m}^{\text{Hajek}}(u) = \sum_{k:N_k>0} \pi_k \left\{ \frac{1}{N_k} \sum_{t:S_t \in C_k} \gamma_{\mathbf{a},q}(S_t) u(S_t) \right\}. \quad (10)$$

This substitution eliminates the variance introduced by the random number of sampled coalitions falling into each cell, while preserving the information gathered from the sampled utilities within those cells. When $K = 1$, the partition consists of the entire coalition space, and the Hájek-type estimator in (10) reduces to the HT estimator in (8).

Under this formulation, the estimator of the Banzhaf value for player i in Wang and Jia [31] can be viewed as a Hájek-type estimator with the partition $\mathcal{C} = \{\{S : i \in S\}, \{S : i \notin S\}\}$. Stratified SVARM [16] refines this approach by further splitting these cells based on coalition size, i.e., $\mathcal{C} = \{\{S : i \in S, |S| = s\}, \{S : i \notin S, |S| = s\} : s = 0, \dots, n\}$. In a related vein, Li and Yu [20] use a size-indexed partition and a similar normalization idea to estimate more general probabilistic values.

As shown in the following proposition, once these estimators are recognized as Hájek-type estimators, their partitions determine the implicit working surrogate classes underlying them. This observation, in turn, places existing stratified weighted-average estimators within our regular-estimator framework. In particular, the corresponding surrogate class consists of functions h whose weighted version $\gamma_{\mathbf{a},q}h$ is constant on each cell of the partition:

$$\mathcal{H}_{\mathcal{C},\gamma_{\mathbf{a},q}} := \left\{ h : \gamma_{\mathbf{a},q}(S)h(S) = \sum_{k=1}^K \omega_k \mathbb{I}(S \in C_k) \text{ for some } \omega_1, \dots, \omega_K \in \mathbb{R} \right\}.$$

Proposition 1 (Regularity of Hájek-type Weighted Average). *Suppose that, on each cell $C_k \in \mathcal{C}$, the weight $\gamma_{\mathbf{a},q}$ is either identically zero or everywhere nonzero. Let $\mu_k := \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)u(S) \mid S \in C_k]$ if $\pi_k > 0$ and $\mu_k := 0$ otherwise. Define $h_{\mathcal{C}}$ by setting $h_{\mathcal{C}}(S) = \mu_k/\gamma_{\mathbf{a},q}(S)$ for $S \in C_k$ on cells where $\gamma_{\mathbf{a},q} \not\equiv 0$, and $h_{\mathcal{C}}(S) = 0$ on cells where $\gamma_{\mathbf{a},q} \equiv 0$. Then*

$h_{\mathcal{C}} \in \mathcal{H}_{\mathcal{C}, \gamma_{\mathbf{a}, q}}$, and the Hájek-type estimator $\hat{\tau}_{\mathbf{a}, m}^{\text{Hajek}}(u)$ in (10) satisfies that

$$\hat{\tau}_{\mathbf{a}, m}^{\text{Hajek}}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_{h_{\mathcal{C}}}(S_t; u) + r_m, \quad \text{with } \mathbb{E}_q(r_m^2) = o(m^{-1}). \quad (11)$$

Therefore, $\hat{\tau}_{\mathbf{a}, m}^{\text{Hajek}}(u)$ is regular relative to $\mathcal{H}_{\mathcal{C}, \gamma_{\mathbf{a}, q}}$.

3.2 Weighted Least-Squares Estimators

We next consider estimators motivated by the least-squares identification in (6). This formulation covers the widely used KernelSHAP estimator and related variants [5, 12, 23, 26], which estimate the target by replacing the population WLS criterion with its empirical analogue and then applying the fixed linear functional $\mathbf{c}_{\mathbf{a}}^{\top}$ to the fitted coefficients. Under limited observations, the sampled design may be rank deficient. To handle this issue and improve numerical stability, we study the WLS estimator with ridge penalty, which also facilitates regularity analysis. For $\lambda > 0$, define

$$\hat{\tau}_{\mathbf{a}, m}^{\text{WLS}, \lambda}(u) := \mathbf{c}_{\mathbf{a}}^{\top} \hat{\boldsymbol{\beta}}_{m, \lambda}, \quad \text{where } \hat{\boldsymbol{\beta}}_{m, \lambda} := \arg \min_{\boldsymbol{\beta} \in \mathbb{R}^d} \left\{ \sum_{t=1}^m \frac{w(S_t)}{q(S_t)} \{u(S_t) - \mathbf{z}(S_t)^{\top} \boldsymbol{\beta}\}^2 + \lambda \|\boldsymbol{\beta}\|_2^2 \right\}. \quad (12)$$

The ridge term makes the empirical problem well posed even under limited observations and provides a stable object for the first-order analysis. When m is large enough, the ridge penalty becomes asymptotically negligible.

The next proposition connects this ridge WLS estimator to the regular estimator framework.

In particular, the associated working surrogate class is the feature span

$$\mathcal{H}_{\text{WLS}} = \left\{ h : h(S) = \mathbf{z}(S)^{\top} \boldsymbol{\beta} \text{ for some } \boldsymbol{\beta} \in \mathbb{R}^d \right\}.$$

Proposition 2 (Regularity of Weighted Least-Squares Estimators). *Let $h_w \in$*

$\arg \min_{h \in \mathcal{H}_{\text{WLS}}} \sum_{S \subseteq [n]} w(S) \{u(S) - h(S)\}^2$ be a population WLS projection of u onto the feature span. Then

$$\hat{\tau}_{\mathbf{a},m}^{\text{WLS},\lambda}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_{h_w}(S_t; u) + r_m, \quad \text{with } \mathbb{E}_q(r_m^2) = o(m^{-1}).$$

Therefore, $\hat{\tau}_{\mathbf{a},m}^{\text{WLS},\lambda}(u)$ is regular relative to $\mathcal{H}_{\text{WLS}}$.

3.3 Composing Implicit and Explicit Surrogates

The previous subsections demonstrate that several weighted-average and least-squares estimators are regular relative to working surrogate classes that are induced implicitly by their estimation procedures. A complementary strategy is to combine these implicit classes with explicit surrogates. Let $h^{\text{sur}} \in \mathcal{H}^{\text{sur}}$ be an explicit surrogate whose target value $\tau_{\mathbf{a}}(h^{\text{sur}})$ is efficiently computable as discussed in Remark 1. Because $\tau_{\mathbf{a}}(\cdot)$, as defined in (2), is a linear functional on the space of utility functions, for any surrogate h^{sur} , we have $\tau_{\mathbf{a}}(u) = \tau_{\mathbf{a}}(h^{\text{sur}}) + \tau_{\mathbf{a}}(u - h^{\text{sur}})$. Thus, any estimator $\hat{\tau}_{\mathbf{a},m}$ for the residual target $\tau_{\mathbf{a}}(u - h^{\text{sur}})$ induces the *surrogate-adjusted* estimator

$$\hat{\tau}_{\mathbf{a},m}^{\text{sur}} := \tau_{\mathbf{a}}(h^{\text{sur}}) + \hat{\tau}_{\mathbf{a},m}(u - h^{\text{sur}}). \quad (13)$$

Here $\hat{\tau}_{\mathbf{a},m}(u - h^{\text{sur}})$ may be any of the weighted-average or least-squares estimators discussed above, applied to the residual observations $\{u(S_t) - h^{\text{sur}}(S_t)\}_{t=1}^m$.

Proposition 3 (Regularity under explicit surrogate adjustment). *Suppose $\hat{\tau}_{\mathbf{a},m}$ is regular relative to an implicit working surrogate class $\mathcal{H}_m^{\text{imp}}$. Then $\hat{\tau}_{\mathbf{a},m}^{\text{sur}}$ is regular relative to*

$$\mathcal{H}_m^{\text{sur}} + \mathcal{H}_m^{\text{imp}} := \{h^{\text{sur}} + h^{\text{imp}} : h^{\text{sur}} \in \mathcal{H}_m^{\text{sur}}, h^{\text{imp}} \in \mathcal{H}_m^{\text{imp}}\}.$$

The proposition shows that explicit and implicit surrogate adjustments compose additively. In one direction, an estimator with an implicit working surrogate class can be enriched by first subtracting an explicit surrogate and then applying the original estimator to the residual. In the other direction, regression-adjusted estimators such as RegressionMSR [33] can be extended by replacing the Horvitz–Thompson residual correction with a residual estimator that induces its own implicit surrogate class.

Fumagalli et al. [12] uses more expressive feature span in regression models, which can be interpreted as improving estimation by enriching the working surrogate class. The choice of this class is important for the estimator design. Its effectiveness depends on how well it captures the structure of u , while its practical choice is constrained by computation and statistical feasibility. Our focus in the next section is complementary. Rather than asking how to design or enrich the working surrogate class itself, we take a class $\mathcal{H}$ as given and study how to choose both the sampling distribution q and the surrogate $h \in \mathcal{H}$ to minimize estimation error.

4 Error Expansion and Design Principles

The regularity framework in the previous section connects many estimator constructions to a common first-order expansion indexed by a sampling law q and a working surrogate h . In this section, we study the leading term in the mean squared error implied by this expansion and its dependence on q and h . This dependence provides a design criterion for choosing sampling laws and surrogates that reduce first-order risk.

If $m \rightarrow \infty$ with the number of players n fixed, the regular expansion in (7) suggests that the leading MSE is the variance of ψ_h divided by m under some regularity conditions. In probabilistic-value estimation, however, this is not the main regime of interest. The utility

function is defined on 2^n coalitions, while feasible sampling budgets are usually much smaller than 2^n , often only polynomial in n [2, 15, 19, 20, 26]. Although the constants hidden in the lower-order remainder are independent of m , they may depend on n . A leading-order criterion is therefore informative only when this remainder is small relative to the leading term at the sample sizes used in practice.

We proceed in two steps. First, we identify the leading MSE term associated with a regular estimator, which captures the joint dependence on q and h . This reveals that the surrogate and the sampling law used by existing estimators are not necessarily the one that minimizes leading error. Second, we give non-asymptotic conditions under which the leading term dominates the remainder for polynomial sampling budgets.

4.1 First-Order MSE as a Design Criterion

We state the results for the general setting in which the same sampled coalitions $\mathcal{D}_m = \{(S_t, u(S_t))\}_{t=1}^m$ are used to estimate multiple linear transformations of probabilistic values, such as the probabilistic values of all players. The single target is a special case.

Let $\tau_{\mathbf{a}_1}(u), \dots, \tau_{\mathbf{a}_d}(u)$ be the targets of interest, where d is independent of m , and let $\mathbf{A} = [\mathbf{a}_1, \dots, \mathbf{a}_d] \in \mathbb{R}^{n \times d}$. The target vector $(\tau_{\mathbf{a}_1}(u), \dots, \tau_{\mathbf{a}_d}(u))^\top$ can then be written as $\boldsymbol{\tau}_{\mathbf{A}}(u) := \mathbf{A}^\top \boldsymbol{\phi}(u)$, where $\boldsymbol{\phi}(u)$ is the vector of probabilistic values. Consider the estimator $\hat{\boldsymbol{\tau}}_{\mathbf{A},m}(u) = (\hat{\tau}_{\mathbf{a}_1,m}(u), \dots, \hat{\tau}_{\mathbf{a}_d,m}(u))^\top$ constructed from $\mathcal{D}_m$. Suppose each coordinate $\hat{\tau}_{\mathbf{a}_j,m}$ is regular relative to the working surrogate class $\mathcal{H}_j$, for $j = 1, \dots, d$. Then there exists $h_j \in \mathcal{H}_j$ such that

$$\hat{\tau}_{\mathbf{a}_j,m}(u) - \tau_{\mathbf{a}_j}(u) := \frac{1}{m} \sum_{t=1}^m \psi_{\mathbf{a}_j,q,h_j}(S_t; u) + r_{m,j}, \quad \mathbb{E}_q(r_{m,j}^2) = o(m^{-1}), \quad (14)$$

where $r_{m,j}$ is the remainder term. Let $\mathbf{r}_m = (r_{m,1}, \dots, r_{m,d})$ and $\mathbf{h} = (h_1, \dots, h_d)$. Define

$V(\mathbf{a}_j; q, h_j) := \text{Var}_q[\gamma_{\mathbf{a}_j, q}(S)\{u(S) - h_j(S)\}]$ as the variance of $\psi_{\mathbf{a}_j, q, h_j}$ and $V(\mathbf{A}; q, \mathbf{h}) := \sum_{j=1}^d V(\mathbf{a}_j; q, h_j)$. The following lemma gives the first-order expansion of MSE.

Lemma 1 (First-order Error Expansion). *Assume that the coordinatewise expansion (14) holds for all $1 \leq j \leq d$. Suppose that $V^*(\mathbf{A}; q) := \inf_{h_j \in \mathcal{H}_j} V(\mathbf{A}; q, \mathbf{h}) > 0$, then*

$$\left| \frac{\mathbb{E}_q[\|\hat{\boldsymbol{\tau}}_{\mathbf{A}, m}(u) - \boldsymbol{\tau}_{\mathbf{A}}(u)\|_2^2]}{V(\mathbf{A}; q, \mathbf{h})/m} - 1 \right| \leq 2 \sqrt{\frac{m\mathbb{E}_q(\|\mathbf{r}_m\|_2^2)}{V(\mathbf{A}; q, \mathbf{h})}} + \frac{m\mathbb{E}_q(\|\mathbf{r}_m\|_2^2)}{V(\mathbf{A}; q, \mathbf{h})}. \quad (15)$$

Consequently, for any $0 < \epsilon \leq 1$, the relative error is at most ϵ whenever $\mathbb{E}_q(\|\mathbf{r}_m\|_2^2) \leq C\epsilon^2 \frac{V(\mathbf{A}; q, \mathbf{h})}{m}$ for some universal constant $C > 0$.

The condition $V^*(\mathbf{A}; q) > 0$ excludes degenerate cases in which the surrogate class can eliminate the first-order variance under the given sampling law. This condition is mild in practice. As discussed in Remark 1, surrogate classes are typically constrained by computational and statistical feasibility and are not assumed to contain the true utility function u .

Since $\mathbb{E}_q(\|\mathbf{r}_m\|_2^2) = \sum_{j=1}^d \mathbb{E}_q(r_{m,j}^2) = o(m^{-1})$, Lemma 1 implies that $V(\mathbf{A}; q, \mathbf{h})/m$ is the first-order MSE term. This expansion motivates using first-order MSE as a design criterion. It suggests choosing q and $\mathbf{h}$ to reduce $V(\mathbf{A}; q, \mathbf{h})$. Existing estimators often induce surrogate adjustments through criteria internal to their construction, such as self-normalization or weighted least squares, or through generic distance metrics between u and h . These choices need not minimize the first-order MSE criterion.

4.2 Non-Asymptotic Bounds for the Remainder

The bound in (15) shows that the accuracy of the first-order MSE approximation is governed by the scaled remainder $m\mathbb{E}_q(\|\mathbf{r}_m\|_2^2)$. The scale of this term depends on both the specific estimator design and the target. This subsection establishes explicit non-asymptotic upper bounds for $\mathbb{E}_q(\|\mathbf{r}_m\|_2^2)$ for two estimator classes: Hájek-type weighted averages and weighted

least-squares (WLS) estimators. We then apply these bounds to representative Shapley-value estimators [16, 20, 23, 26] and show that, for $0 < \epsilon \leq 1$, the first-order MSE approximation has relative error at most ϵ once $m = \Omega(n \text{polylog}(n)/\epsilon^2)$, which aligns with the regime of primary interest in the literature [19, 20, 26].

Hájek-type weighted average estimators. Consider the Hájek-type estimator for a scalar target in (10) with partition $\mathcal{C} = \{C_k\}_{k=1}^K$, and let $h_{\mathcal{C}}$ be the corresponding cellwise surrogate defined in Proposition 1. Denote the number of cells with positive sampling probability by $K_+ := |\{k : \pi_k > 0\}|$, and the smallest nonzero cell probability by $\pi_{\min} := \min_{k:\pi_k>0} \pi_k$. Define the average cellwise squared weight and the largest magnitude by $\bar{\Gamma}_{2,\mathcal{C}}^2 := \frac{1}{K_+} \sum_{k:\pi_k>0} \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)^2 \mid S \in C_k]$ and $\Gamma_{\infty} := \sup_{S:q(S)>0} |\gamma_{\mathbf{a},q}(S)|$, respectively.

Theorem 1 (Non-asymptotic remainder bound for Hájek-type estimators). *Assume the conditions in Proposition 1 hold. There exist universal constants $C_1, C_2 > 0$ such that*

$$\mathbb{E}_q(r_m^2) \leq C_1 \|u\|_{\infty}^2 \left\{ \frac{K_+ \bar{\Gamma}_{2,\mathcal{C}}^2}{m^2} + \Gamma_{\infty}^2 e^{-C_2 m \pi_{\min}} \right\}. \quad (16)$$

Theorem 1 bounds the remainder by two terms. The first term is the main polynomial term in m . It reflects the cost of estimating separate normalizing factors across the active cells in $\mathcal{C}$, and scales with both K_+ and $\bar{\Gamma}_{2,\mathcal{C}}^2$. The second term controls the rare event that some active cell receives no samples. It decays exponentially in $m\pi_{\min}$ and is governed by the largest weight Γ_{∞} . We interpret the bound through two representative Hájek-type estimators, OFA [20] and Stratified SVARM [16], specialized to Shapley-value estimation through their corresponding partitions and sampling laws. Details are given in Appendix B.1.

Corollary 1 (Remainder bound for OFA and Stratified SVARM). *Consider estimating the*

Shapley value ϕ_i of a fixed player i . There exist universal constants $C_1, C_2 > 0$ such that

$$\mathbb{E}_q(r_m^2) \leq C_1 \|u\|_\infty^2 \begin{cases} \frac{n \log n}{m^2} + n e^{-C_2 m / n^{3/2}}, & \text{for OFA [20]}, \\ \frac{n \log^2 n}{m^2} + \log^2 n e^{-C_2 m / (n \log n)}, & \text{for Stratified SVARM [16]}. \end{cases}$$

The polynomial terms in Corollary 1 determine the leading remainder scales. If $\|u\|_\infty$ is bounded and the first-order MSE is nondegenerate, then, for any $0 < \epsilon \leq 1$, the first-order MSE approximation has relative error at most ϵ once $m = \Omega(n \log n / \epsilon^2)$ for OFA and $m = \Omega(n \log^2 n / \epsilon^2)$ for Stratified SVARM. For OFA, this $n \log n$ dependence implies the non-asymptotic query complexity required for its Shapley-value approximation guarantee (see Appendix B.1 for more details).

WLS estimators. Consider a vector-valued extension of the WLS estimators. Suppose $\mathbf{z}(S) \in \mathbb{R}^{d_z}$ and the target vector $\boldsymbol{\tau}_A(u)$ is identified through the WLS projection

$$\boldsymbol{\tau}_A(u) = \mathbf{C}_A^\top \boldsymbol{\beta}^*(u), \quad \text{for every } \boldsymbol{\beta}^*(u) \in \arg \min_{\boldsymbol{\beta} \in \mathbb{R}^{d_z}} \mathbb{E}_q \left[\frac{w(S)}{q(S)} \{u(S) - \mathbf{z}(S)^\top \boldsymbol{\beta}\}^2 \right].$$

We estimate this target by $\hat{\boldsymbol{\tau}}_{A,m}^{\text{WLS},\lambda} = \mathbf{C}_A^\top \hat{\boldsymbol{\beta}}_{m,\lambda}$, where $\hat{\boldsymbol{\beta}}_{m,\lambda}$ is the ridge estimator in (12).

Each coordinate admits the regular expansion in (14) with the same surrogate $h_j = h_w$ for all j , where h_w is defined in Proposition 2. The next result establishes a non-asymptotic bound for the remainder $\mathbb{E}_q(\|\mathbf{r}_m\|_2^2)$, which depends on the leverage scores of the weighted feature design. Define the Gram matrix and the leverage scores as

$$\mathbf{G} = \mathbb{E}_q[\tilde{w}(S) \mathbf{z}(S) \mathbf{z}(S)^\top], \quad \ell(S) = \tilde{w}(S) \mathbf{z}(S)^\top \mathbf{G}^+ \mathbf{z}(S),$$

where $\tilde{w}(S) := w(S)/q(S)$ with $\text{supp}(w) \subseteq \text{supp}(q)$ and $\mathbf{G}^+$ is the Moore–Penrose pseudoinverse. Large values of $\ell(S)$ indicate that the WLS fit is more sensitive to the observation at S .

Theorem 2 (Non-asymptotic remainder bound for WLS estimators). *Let $L = \sup_{S:q(S)>0} \ell(S)$ be the maximum leverage score. Assume that $\mathbf{G}$ has nonzero rank. Let $\bar{\sigma}_{\min}(\mathbf{G})$ and $\sigma_{\max}(\mathbf{G})$ denote the minimum nonzero and maximum eigenvalues of $\mathbf{G}$, respectively, and define $\kappa_{\mathbf{G}} := \sigma_{\max}(\mathbf{G})/\bar{\sigma}_{\min}(\mathbf{G})$. Define $R_{\mathbf{A}}^2 := \sigma_{\max}(\mathbf{C}_{\mathbf{A}}^{\top} \mathbf{G}^+ \mathbf{C}_{\mathbf{A}})$, $\|u\|_{2,\tilde{w}}^2 := \mathbb{E}_q[\tilde{w}(S)u(S)^2]$, and $\|u\|_{\infty,\tilde{w}}^2 := \sup_{S:q(S)>0} \tilde{w}(S)u(S)^2$. If the ridge penalty parameter satisfies $\lambda \asymp \sigma_{\min}^+(\mathbf{G})$, then there exist universal constants $C_1, C_2 > 0$ such that*

$$\mathbb{E}_q(\|\mathbf{r}_m\|_2^2) \leq C_1 R_{\mathbf{A}}^2 L^2 \left[\frac{1}{m^2} \log d_{\mathbf{z}} \left(1 + \frac{L \log d_{\mathbf{z}}}{m} \right) \|u\|_{2,\tilde{w}}^2 + e^{-\frac{C_2 m}{L}} \cdot d_{\mathbf{z}} m^2 \kappa_{\mathbf{G}}^2 \|u\|_{\infty,\tilde{w}}^2 \right]. \quad (17)$$

The bound in Theorem 2 also has two terms. The first term is polynomial in m and is scaled by the maximum leverage L and by $R_{\mathbf{A}}^2$. The factor $R_{\mathbf{A}}^2$ captures how strongly the target readout $\mathbf{C}_{\mathbf{A}}^{\top} \boldsymbol{\beta}$ can amplify errors in weakly identified WLS directions. The second term is exponentially small in m/L and accounts for the event that the empirical weighted design is poorly conditioned.

We next specialize this bound to two representative WLS-based Shapley estimators: KernelSHAP [23] and LeverageSHAP [26]. We focus on the centered Shapley vector, $\boldsymbol{\tau}_{\mathbf{A}}(u) = \mathbf{A}\boldsymbol{\phi}(u)$, where $\mathbf{A} = \mathbf{I} - n^{-1}\mathbf{1}\mathbf{1}^{\top}$ and $\boldsymbol{\phi}(u)$ is the Shapley vector. This centered target $\boldsymbol{\tau}_{\mathbf{A}}(u)$ can be represented as a linear readout $\mathbf{C}_{\mathbf{A}}^{\top} \boldsymbol{\beta}$ with certain coefficients $\mathbf{C}_{\mathbf{A}}$.

Corollary 2 (Remainder bound for KernelSHAP and LeverageSHAP). *Consider estimating the centered Shapley vector $\mathbf{A}\boldsymbol{\phi}(u)$ through the WLS characterization. There exist universal constants $C_1, C_2 > 0$ such that*

$$\mathbb{E}_q(\|\mathbf{r}_m\|_2^2) \leq C_1 \|u\|_{\infty}^2 \begin{cases} \frac{n^2 \log^4 n}{m^2} \left(1 + \frac{n \log^2 n}{m} \right) + n^3 m^2 \log^3 n e^{-\frac{C_2 m}{n \log n}}, & \text{KernelSHAP,} \\ \frac{n^2 \log^2 n}{m^2} \left(1 + \frac{n \log n}{m} \right) + n^4 m^2 e^{-C_2 \frac{m}{n}}, & \text{LeverageSHAP.} \end{cases}$$

The polynomial terms in Corollary 2 yield sufficient sample sizes for making the remainder negligible relative to the first-order MSE. In particular, if $\|u\|_\infty$ is bounded and the first-order MSE has the natural full-vector scale $V(\mathbf{A}; q, \mathbf{h}) = \sum_{j=1}^n V(\mathbf{a}_j; q, h_w) = O(n)$, then it is sufficient to take $m = \Omega(n \log^4 n)$ for KernelSHAP and $m = \Omega(n \log^2 n)$ for LeverageSHAP. For LeverageSHAP, this scale implies the non-asymptotic $n \log n$ query guarantee up to one additional logarithmic factor (see Appendix B.2 for more details).

5 Efficiency-Aware Surrogate-Adjusted Estimation

So far, we have shown that many existing estimator designs are regular relative to working surrogate classes induced by their estimation procedures. The regularity framework gives a common first-order expansion, of which the leading MSE is given by $V(\mathbf{A}; q, \mathbf{h})/m$. This quantity depends on both the sampling law q and surrogate choice $\mathbf{h}$. Our goal in this section is to construct an estimator that targets this first-order MSE criterion directly.

The explicit surrogate-adjusted estimator, or AIPW estimator, in (9) provides a natural construction. For any fixed $(q, \mathbf{h})$, its MSE is exactly $V(\mathbf{A}; q, \mathbf{h})/m$, matching the leading MSE of a regular estimator with the same $(q, \mathbf{h})$. Therefore, given a working surrogate class $\mathcal{H}$, we can target the first-order efficiency by choosing the sampling law q and the surrogate $\mathbf{h} \in \mathcal{H}$ to minimize $V(\mathbf{A}; q, \mathbf{h})$.

This optimization problem involves two coupled design choices: how to fit the surrogate $\mathbf{h}$, and how to choose the sampling law q . Because the criterion depends on the unknown utility function u , both choices must be learned from sampled utility evaluations. In addition, choosing q without further structure would require optimizing over all distributions on $2^{[n]}$, which is computationally infeasible for even moderate n . We first study the surrogate-learning problem for a fixed sampling law q , and then turn to the remaining question of

how to choose q within a structured class using pilot data.

5.1 Surrogate Learning for a Fixed Sampling Law

Suppose that we observe $S_1, \dots, S_m \stackrel{\text{iid}}{\sim} q$ for a fixed sampling law q , which is positive on the support of each $\rho_{\mathbf{a}_j}$, and that we have access to the corresponding utility evaluations $Y_t = u(S_t)$. Consider the AIPW estimator with surrogate $\mathbf{h} = (h_1, \dots, h_d)$ for the target vector $\boldsymbol{\tau}_{\mathbf{A}}(u)$:

$$\hat{\tau}_{\mathbf{a}_j}^{\text{AIPW}}(u) = \tau_{\mathbf{a}_j}(h_j) + \frac{1}{m} \sum_{t=1}^m \gamma_{\mathbf{a}_j, q}(S_t) \{Y_t - h_j(S_t)\}, \quad j = 1, \dots, d. \quad (18)$$

For fixed q , the design question is to find $\mathbf{h} \in \mathcal{H}$ to minimize the MSE: $V(\mathbf{A}; q, \mathbf{h}) = \sum_{j=1}^d \text{Var}_q[\gamma_{\mathbf{a}_j, q}(S) \{u(S) - h_j(S)\}]$. Equivalently, this variance criterion can be written as a profiled squared-loss objective. Introduce an auxiliary centering vector $\boldsymbol{\mu} = (\mu_1, \dots, \mu_d)$ and define:

$$\mathcal{V}(\mathbf{A}; q, \mathbf{h}, \boldsymbol{\mu}) = \sum_{j=1}^d \mathbb{E}_q \left[\left\{ \gamma_{\mathbf{a}_j, q}(S) \{u(S) - h_j(S)\} - \mu_j \right\}^2 \right].$$

Then, $V(\mathbf{A}; q, \mathbf{h}) = \min_{\boldsymbol{\mu} \in \mathbb{R}^d} \mathcal{V}(\mathbf{A}; q, \mathbf{h}, \boldsymbol{\mu})$, where the minimizer centers each term at its mean. Thus, the oracle surrogate for fixed q can be equivalently obtained by solving $(\mathbf{h}_q^*, \boldsymbol{\mu}_q^*) \in \arg \min_{\mathbf{h} \in \mathcal{H}, \boldsymbol{\mu} \in \mathbb{R}^d} \mathcal{V}(\mathbf{A}; q, \mathbf{h}, \boldsymbol{\mu})$. Given the observed samples indexed by $\mathcal{T} \subseteq [m]$, we estimate this oracle target by the empirical least-squares problem

$$(\hat{\mathbf{h}}_{\mathcal{T}}, \hat{\boldsymbol{\mu}}_{\mathcal{T}}) \in \arg \min_{\mathbf{h} \in \mathcal{H}, \boldsymbol{\mu} \in \mathbb{R}^d} \sum_{t \in \mathcal{T}} \sum_{j=1}^d \left[\gamma_{\mathbf{a}_j, q}(S_t) \{Y_t - h_j(S_t)\} - \mu_j \right]^2. \quad (19)$$

The fitted surrogate $\hat{\mathbf{h}}_{\mathcal{T}}$ is then used in the AIPW estimator, while $\hat{\boldsymbol{\mu}}_{\mathcal{T}}$ is only an auxiliary variable used for training. To avoid using the same samples for surrogate learning and estimator evaluation, we use a cross-fitted version of (18), as summarized in Algorithm 1.

Remark 2. If the surrogate class is coordinate-separable, i.e., $\mathcal{H} = \mathcal{H}_1 \times \cdots \times \mathcal{H}_d$, then (19) decomposes into d independent problems:

$$(\hat{h}_{j,\mathcal{T}}, \hat{\mu}_{j,\mathcal{T}}) \in \arg \min_{h_j \in \mathcal{H}_j, \mu_j \in \mathbb{R}} \sum_{t \in \mathcal{T}} [\gamma_{\mathbf{a}_j, q}(S_t) \{Y_t - h_j(S_t)\} - \mu_j]^2, \quad j = 1, \dots, d.$$

This gives the most flexible surrogate within the product class, but requires fitting d separate models. To reduce computation, one may use a shared surrogate $h \in \mathcal{H}_0$ and solve

$$(\hat{h}_{\mathcal{T}}, \hat{\boldsymbol{\mu}}_{\mathcal{T}}) \in \arg \min_{h \in \mathcal{H}_0, \boldsymbol{\mu} \in \mathbb{R}^d} \sum_{t \in \mathcal{T}} \sum_{j=1}^d [\gamma_{\mathbf{a}_j, q}(S_t) \{Y_t - h(S_t)\} - \mu_j]^2.$$

The shared surrogate trades some statistical flexibility for lower computational cost.

Remark 3. Witter et al. [33] also use a cross-fitted AIPW construction, but their surrogate is learned by minimizing an unweighted approximation loss, $\sum_{t \in \mathcal{T}} \sum_{j=1}^d [Y_t - h(S_t)]^2$. Our first-order MSE analysis instead leads to the efficiency-aware objective in (19), which depends explicitly on both the weights $\gamma_{\mathbf{a}_j, q}$ and the sampling law q . Thus, the surrogate is trained to reduce the first-order risk, rather than only to approximate u uniformly over the sampled coalitions. The next subsection further chooses q to target the same first-order risk criterion.

5.2 Choice of Sampling Law

The first-order MSE criterion also suggests how q should be chosen. Since optimizing over all distributions on $2^{[n]}$ is infeasible even for moderate n , we restrict q to a structured class. Let $\mathcal{C} = \{C_1, \dots, C_K\}$ be a partition of the coalition space. We consider laws that are constant within each cell: $\mathcal{Q}_{\mathcal{C}} = \{q_{\boldsymbol{\pi}} : q_{\boldsymbol{\pi}}(S) = \pi_k \text{ for } S \in C_k, \pi_k \geq 0, \sum_{k=1}^K |C_k| \pi_k = 1\}$. This class includes, for example, coalition-size stratified designs, where coalitions of the same cardinality have the same sampling probability.

For a fixed surrogate $\mathbf{h} = (h_1, \dots, h_d)$, define the cellwise average squared residual

$$M_k(\mathbf{h}) := \frac{1}{|C_k|} \sum_{S \in C_k} \sum_{j=1}^d \rho_{\mathbf{a}_j}(S)^2 \{u(S) - h_j(S)\}^2, \quad \text{for } k = 1, \dots, K. \quad (20)$$

Within $\mathcal{Q}_{\mathcal{C}}$, the oracle sampling law is given by $q^*(S; \mathbf{h}) = \sqrt{M_k(\mathbf{h})} / \sum_{r=1}^K |C_r| \sqrt{M_r(\mathbf{h})}$, for $S \in C_k$. Thus, the per-coalition sampling probability in each cell is proportional to the square root of its average squared residual in that cell.

Since $M_k(\mathbf{h})$ is unknown before sampling, we use a two-stage plug-in procedure. We first initialize q by replacing the residual scale with a common constant:

$$M_k^{\text{init}} := \frac{1}{|C_k|} \sum_{S \in C_k} \sum_{j=1}^d \rho_{\mathbf{a}_j}(S)^2, \quad q^{\text{init}}(S) \propto \sqrt{M_k^{\text{init}}}, \quad \text{for } S \in C_k. \quad (21)$$

After collecting pilot samples $S_1, \dots, S_{m_{\text{init}}} \stackrel{\text{iid}}{\sim} q^{\text{init}}$ and evaluating $Y_t^{\text{init}} = u(S_t^{\text{init}})$, we fit an initial surrogate $\hat{\mathbf{h}}^{\text{init}}$ by solving (19). We then estimate the cellwise residual moments by

$$\hat{M}_k = \frac{1}{N_k} \sum_{t: S_t^{\text{init}} \in C_k} \sum_{j=1}^d \rho_{\mathbf{a}_j}(S_t^{\text{init}})^2 \{Y_t^{\text{init}} - \hat{h}_j^{\text{init}}(S_t^{\text{init}})\}^2, \quad \text{where } N_k = \sum_{t=1}^{m_{\text{init}}} \mathbb{I}(S_t^{\text{init}} \in C_k). \quad (22)$$

The post-pilot law is updated as $\hat{q}(S) \propto \sqrt{\hat{M}_k}$ for $S \in C_k$. The second stage samples from $\hat{q}$, refits the surrogate under this design, and applies the cross-fitted AIPW estimator. Algorithm 1 presents the simple version in which pilot samples are used only to construct $\hat{q}$ and are discarded from final estimation. We defer the handling of zero or extremely small estimated cell probabilities, empty pilot cells, on-the-fly surrogate estimates, and variants with complement-pair sampling to Appendix C.

Remark 4 (Residual-Aware Sampling). *This residual-aware oracle law can be viewed as a Neyman-type allocation [27] over the partition $\mathcal{C}$. Classical Neyman allocation assigns more samples to strata with larger variability, while our first-order analysis shows that, for*

our surrogate-adjusted estimator, the relevant variability is captured by the cellwise average squared residual $M_k(\mathbf{h})$. Thus, the allocation is driven not by raw utility variability alone, but by the residual variation left after surrogate adjustment. Because the working surrogate class may not contain the true utility function, this residual is generally non-negligible and can play an important role in sampling design. To the best of our knowledge, EASE is the first probabilistic-value estimation framework in which the sampling law is explicitly chosen as a function of the residuals induced by the fitted surrogate. The pilot stage in Algorithm 1 operationalizes this principle by estimating $M_k(\hat{\mathbf{h}})$ and using these estimates to choose the second-stage sampling law.

Algorithm 1: Efficiency-Aware Surrogate-Adjusted Estimation (EASE)

Input : Partition $\mathcal{C} = \{C_1, \dots, C_K\}$; pilot budget m_{init} ; estimation budget m_{est} ; fold partition $\{I_b\}_{b=1}^B$ of $\{1, \dots, m_{\text{est}}\}$; target matrix $\mathbf{A}$; surrogate class $\mathcal{H}$

Output : Estimator $\hat{\tau}_{\mathbf{A}}$

▷ **Stage 1: Initialization**

1 Draw $\{(S_t^{\text{init}}, Y_t^{\text{init}})\}_{t=1}^{m_{\text{init}}}$, where $S_t^{\text{init}} \stackrel{\text{iid}}{\sim} q^{\text{init}}$ is defined in (21) and $Y_t^{\text{init}} = u(S_t^{\text{init}})$

2 Estimate $\hat{M}_k$ for each cell using (22)

3 Set the final sampling law $q^{\text{final}}(S) \propto \sqrt{\hat{M}_k}$ for $S \in C_k$

▷ **Stage 2: Estimation**

4 Draw the estimation sample $\{(S_t, Y_t)\}_{t=1}^{m_{\text{est}}}$, where $S_t \sim q^{\text{final}}$ and $Y_t = u(S_t)$

5 **for** $b = 1, \dots, B$ **do**

6 Fit $(\hat{\mathbf{h}}^{(-b)}, \hat{\boldsymbol{\mu}}^{(-b)})$ by (19) on the training set $\mathcal{T} = \{1, \dots, m_{\text{est}}\} \setminus I_b$

7 Compute $\hat{\tau}_{\mathbf{A}}^{(b)}$ on the held-out fold I_b . Specifically, $\hat{\tau}_{\mathbf{A}}^{(b)} = (\hat{\tau}_{\mathbf{a}_1}^{(b)}, \dots, \hat{\tau}_{\mathbf{a}_d}^{(b)})^\top$, where, for $j = 1, \dots, d$,

$$\hat{\tau}_{\mathbf{a}_j}^{(b)} = \tau_{\mathbf{a}_j}(\hat{h}_j^{(-b)}) + \frac{1}{|I_b|} \sum_{t \in I_b} \gamma_{\mathbf{a}_j, q^{\text{final}}}(S_t) \{Y_t - \hat{h}_j^{(-b)}(S_t)\}.$$

8 **end**

9 Return $\hat{\tau}_{\mathbf{A}} = \sum_{b=1}^B \frac{|I_b|}{m_{\text{est}}} \hat{\tau}_{\mathbf{A}}^{(b)}$

6 Numerical Results

This section evaluates the proposed EASE method for probabilistic value estimation with two main goals. First, we compare EASE with representative state-of-the-art methods from three estimator families: RegressionMSR [33] as a surrogate-adjusted method, OFA [20] as a weighted-average method, and PolySHAP [12] as a WLS method. In each comparison, we keep the working surrogate class fixed to the class used either explicitly or implicitly by the baseline. EASE differs in how it learns the surrogate $\mathbf{h}$ and chooses the sampling law q according to the first-order efficiency criterion. This matched comparison isolates the gain from the EASE design rather than from a richer surrogate class. Second, we benchmark EASE against additional baselines across several probabilistic values, showing that the same efficiency-aware design principle remains effective beyond the matched comparisons. Code for reproducing the numerical experiments is available in the [EASE GitHub repository](#).

Design of cooperative games. We evaluate methods using structured sum-of-unanimity (SOU) games, which have been widely used for evaluating Monte Carlo estimators of Shapley values and related probabilistic values [18, 20]. SOU games are useful for this goal because their exact probabilistic values are available in closed form, allowing accurate evaluation of the estimation error. The utility takes the form $u(S) = \sum_{\tilde{S} \in \mathcal{S}} \theta_{\tilde{S}} \mathbb{I}(\tilde{S} \subseteq S)$, where $\mathcal{S}$ is a finite multiset of coalitions and each occurrence $\tilde{S} \in \mathcal{S}$ has coefficient $\theta_{\tilde{S}}$.

We design utilities with explicit control over structural complexity by decomposing the multiset of coalitions as $\mathcal{S} = \mathcal{S}_{\text{low}} \cup \mathcal{S}_{\text{high}}$. The low-order component $\mathcal{S}_{\text{low}} = \{\tilde{S} \subseteq [n] : 1 \leq |\tilde{S}| \leq 2\}$ contains all singleton and pairwise coalitions, which captures the main effects and pairwise interactions. The high-order component $\mathcal{S}_{\text{high}}$ consists of n^2 random coalitions with at least three players, representing more complex higher-order interactions that are harder to model. The coefficients $\theta_{\tilde{S}}$ determine the extent to which utilities can be explained

by low-order terms. Given $\eta \in (0, 1)$ and $\sigma^2 > 0$, we draw the coefficients independently as:

$$\theta_{\tilde{S}} \sim N\left(0, \frac{\eta\sigma^2}{|\mathcal{S}_{\text{low}}|}\right), \quad \tilde{S} \in \mathcal{S}_{\text{low}}; \quad \theta_{\tilde{S}} \sim N\left(0, \frac{(1-\eta)\sigma^2}{|\mathcal{S}_{\text{high}}|}\right), \quad \tilde{S} \in \mathcal{S}_{\text{high}}.$$

Under this scaling, $\eta\sigma^2$ is the total coefficient variance assigned to singleton and pairwise coalitions, while $(1-\eta)\sigma^2$ is assigned to higher-order coalitions. Thus, with larger η , utilities can be better approximated by surrogates that model low-order interactions. Varying η lets us compare EASE and baselines in a controlled range of structural complexity.

6.1 Matched Comparisons: Gains from Efficiency-aware Design

In this subsection, we set the target as the n -dimensional Shapley value vector $\phi(u)$ with $n = 40$. The ground truth Shapley values for SOU games can be computed analytically.

We compare EASE with three representative methods using matched working surrogate classes. For RegressionMSR [33], we match its linear version using the linear class spanned by $\mathbb{I}(i \in S)$, which captures main effects. For OFA [20], we match the player-by-size averaging structure used by the estimator. For each player i , OFA estimates averages of $u(S)$ over coalitions grouped by their size and by whether they contain i . Thus, the corresponding implicit surrogate class is spanned by indicators $\mathbb{I}(|S| = s, i \in S)$. For PolySHAP [12], the degree-two polynomial class is spanned by indicators $\mathbb{I}(\tilde{S} \subseteq S)$ with $|\tilde{S}| \leq 2$, which captures main effects and pairwise interactions.

We measure estimation accuracy by the relative squared error $\|\hat{\phi} - \phi\|_2^2 / \|\phi\|_2^2$. Figure 1 summarizes the relative estimation errors against the average number of utility evaluations per player. Across all three comparisons and three values of η , EASE achieves a smaller estimation error for Shapley values than the corresponding baseline at every measured budget. These results demonstrate the benefits of our efficiency-aware design: using the

same working surrogate class, the efficiency-guided selection of the surrogate $\mathbf{h}$ and the sampling law q yields consistent improvements over existing estimators.

Moreover, for each fixed game (i.e., a fixed η per row), EASE equipped with richer working classes, such as those used in OFA and PolySHAP, achieves lower estimation errors than EASE using the linear class from RegressionMSR, which is as expected. The advantage of the degree-two polynomial working class is particularly pronounced when $\eta = 0.75$; in this scenario, the utility function is well-captured by main effects and pairwise interactions, making this surrogate class highly effective. Interestingly, while OFA implicitly uses a richer surrogate class than RegressionMSR, this added complexity does not inherently guarantee better performance (as seen by comparing the orange lines in the first and second columns). However, when paired with our efficiency-guided surrogate and sampling law, EASE unlocks the potential of this added capacity and restores the expected performance hierarchy.

6.2 Benchmark Across Various Probabilistic Values

We further evaluate EASE across a broader benchmark by varying the target probabilistic values and the competing estimators. Using the same structured SOU games, we estimate six probabilistic value vectors: the Shapley value, Beta Shapley with parameters $(4, 1)$ and $(1, 4)$ [17], and weighted Banzhaf values with $p \in \{0.25, 0.5, 0.75\}$ [31].

For each target, we compare two EASE variants against existing baselines from the literature. The two EASE variants differ in the choice of the working surrogate class. *EASE-FO (First-Order)* uses a surrogate class spanned by the player indicators $\mathbb{I}(i \in S)$. To capture nonlinearities in the utility function, we augment this basis with an intercept and two coalition-size terms: $\log(1 + |S|)$ and $(|S|/n)^2$. *EASE-SP (Size-Player)* uses a richer surrogate class spanned by $\mathbb{I}(|S| = s, i \in S)$. While more flexible, this richer basis is more computationally expensive to fit due to the increased dimensionality. Baselines include OFA [20], sampling lift




Figure 1. Matched comparisons on SOU games for Shapley-value estimation. Each row corresponds to a value of $\eta \in \{0.25, 0.5, 0.75\}$, which controls the strength of the low-order component in the utility. The three columns compare EASE with RegressionMSR, OFA, and PolySHAP, respectively, using the same explicit or implicit working surrogate class as the corresponding baseline. The x-axis reports the average number of utility evaluations per player, and the y-axis reports the relative squared error $\|\hat{\phi} - \phi\|_2^2 / \|\phi\|_2^2$ on a log scale. Solid lines represent the mean relative error for each estimator, and the shaded bands indicate one standard deviation over 10 independent runs.

and weighted variants [17, 25], SHAP-IQ [11], GELS and WGELS [19], KernelSHAP [6, 23], permutation and weighted variants [2], complementary-contribution sampling [35], group testing [15], AME [21], RegressionMSR [33], and PolySHAP [12]. Methods restricted to specific probabilistic values are plotted only where applicable.

To quantify estimator performance, we summarize each convergence curve by the *area under the convergence curve* (AUCC), following recent work on probabilistic-value estimation [18, 20]. For a set of tracked utility-evaluation budgets $\mathcal{B} = \{50, 100, \dots, 5000\}$, the AUCC is defined as the average relative squared error along the convergence curve: $\text{AUCC} = \frac{1}{|\mathcal{B}|} \sum_{m \in \mathcal{B}} \|\hat{\phi}_m - \phi\|_2^2 / \|\phi\|_2^2$. Lower AUCC values indicate better sample efficiency.

Figure 2 reports the mean AUCC over 10 runs, with shaded bands representing one standard deviation. We present results for $\eta = 0.25$ in the main text and provide results for $\eta \in \{0.5, 0.75\}$ in Appendix D. For Beta-Shapley values (including the Shapley values), both EASE variants outperform all baselines, often by a significant margin, with EASE-SP typically providing the best performance. For weighted Banzhaf values, while AME and RegressionMSR also perform well, EASE remains highly competitive.

Overall, these results demonstrate that, by learning the surrogate $\mathbf{h}$ and choosing the sampling law q to minimize first-order MSE, the EASE framework offers consistently strong performance across a diverse range of probabilistic values.

7 Discussion: Beyond Single-Coalition Sampling

While we have focused on single-coalition sampling so far, the target $\tau_{\mathbf{a}}(u)$ can also be identified under *bundled sampling*, where each Monte Carlo draw returns a tuple of multiple coalitions $\mathbf{T} = (S_1, \dots, S_r) \sim Q$. Several established sampling designs fall into this category, including edge pairs $(S, S \cup \{i\})$ [17, 25], complement pairs (S, S^c) [12, 26, 35], and entire




Figure 2. AUCC benchmark on the SOU game with $\eta = 0.25$ comparing EASE against baseline estimators for various target probabilistic values. The x-axis denotes the specific target value being estimated, including Shapley values, Beta Shapley (BS), and weighted Banzhaf values (WB). The y-axis reports the Area Under the Convergence Curve (AUCC) on a log scale, with lower AUCC indicating better performance. Solid lines represent the mean AUCC for each estimator, and the shaded bands indicate one standard deviation over 10 independent runs.

permutation paths [1, 2, 24, 34]. In Appendix E, we extend the regular estimator class to the bundled sampling, and the first-order efficiency analysis for the regular estimators can be established similarly. However, unlike single-coalition sampling, the first-order MSE criterion depends on not only the surrogate h and the bundle law Q , but also the choice of a feasible weighting γ . Specifically, for the target $\tau_{\mathbf{a}}(u) = \sum_{S \subseteq [n]} \rho_{\mathbf{a}}(S)u(S)$, the weighting under a single-coalition law q is uniquely determined as $\gamma_{\mathbf{a},q}(S) = \rho_{\mathbf{a}}(S)/q(S)$. In contrast, under a bundled sampling law Q , the weighting vector $\gamma(\mathbf{T}) = (\gamma_1(\mathbf{T}), \dots, \gamma_r(\mathbf{T}))$ only needs to satisfy the condition $\rho_{\mathbf{a}}(S) = \mathbb{E}_Q[\sum_{\ell=1}^r \gamma_{\ell}(\mathbf{T}) \mathbf{1}\{S_{\ell} = S\}]$, for any $S \subseteq [n]$. Because this system is generally underdetermined, the feasible weighting is no longer unique, and the first-order MSE design involves selecting an optimal γ . In Appendix E, we study complement-pair sampling $\mathbf{T} = (S, S^c)$ with $S \sim q$ as a tractable example. For semivalues whose weights are symmetric with respect to coalition size, we demonstrate

that the weighting $\gamma(\mathbf{T}) = \left(\frac{\rho_{\mathbf{a}}(S)}{q(S)+q(S^c)}, -\frac{\rho_{\mathbf{a}}(S)}{q(S)+q(S^c)} \right)$, which has been implicitly used in the literature [12, 33, 35], represents the Rao–Blackwellized weighting. This choice minimizes the first-order MSE for a fixed complement-pair law and surrogate. A promising direction for future work is to generalize this understanding to design efficiency-aware estimators under a wider variety of bundled sampling designs.

Acknowledgements

Liu and Tang were supported by NSF DMS-2412853 and Jane Street Group, LLC. Lee and Zhang were supported by NSF DMS-2311109.

References

- [1] Castro, J., Gómez, D., Molina, E., et al. Improving polynomial estimation of the shapley value by stratified random sampling with optimum allocation. *Computers & Operations Research*, 82:180–188, 2017.
- [2] Castro, J., Gómez, D., and Tejada, J. Polynomial calculation of the shapley value based on sampling. *Computers & Operations Research*, 36(5):1726–1730, 2009.
- [3] Chau, S. L., Hu, R., Gonzalez, J., et al. RKHS-SHAP: Shapley values for kernel methods. In A. H. Oh, A. Agarwal, D. Belgrave, and K. Cho, editors, *Advances in Neural Information Processing Systems*. 2022.
- [4] Chen, H., Covert, I. C., Lundberg, S. M., et al. Algorithms to estimate shapley value feature attributions. *Nature Machine Intelligence*, 5(6):590–601, 2023.
- [5] Chen, T., Seshadri, A., Villani, M. J., et al. A unified framework for provably efficient algorithms to estimate shapley values. In *The Thirty-ninth Annual Conference on Neural Information Processing Systems*. 2025.
- [6] Covert, I. and Lee, S.-I. Improving KernelSHAP: Practical shapley value estimation using linear regression. In A. Banerjee and K. Fukumizu, editors, *Proceedings of The 24th International Conference on Artificial Intelligence and Statistics*, volume 130 of *Proceedings of Machine Learning Research*, pages 3457–3465. PMLR, 2021.

- [7] Dagdoug, M., Goga, C., and Haziza, D. Model-assisted estimation in high-dimensional settings for survey data. *Journal of Applied Statistics*, 50(3):761–785, 2023.
- [8] de la Peña, V. and Giné, E. *Decoupling: From dependence to independence*. Springer New York, 1999.
- [9] Deng, J., Hu, Y., Hu, P., et al. A survey of data attribution: Methods, applications, and evaluation in the era of generative AI. 2025.
- [10] Dubey, P., Neyman, A., and Weber, R. J. Value theory without efficiency. *Math. Oper. Res.*, 6(1):122–128, 1981. doi:10.1287/moor.6.1.122.
- [11] Fumagalli, F., Muschalik, M., Kolpaczki, P., et al. SHAP-IQ: Unified approximation of any-order shapley interactions. In *Thirty-seventh Conference on Neural Information Processing Systems*. 2023.
- [12] Fumagalli, F., Witter, R. T., and Musco, C. PolySHAP: Extending KernelSHAP with interaction-informed polynomial regression. In *International Conference on Learning Representations*. 2026.
- [13] Hájek, J. Comment on “an essay on the logical foundations of survey sampling, part one”. In V. P. Godambe and D. A. Sprott, editors, *Foundations of Statistical Inference*, page 236. Holt, Rinehart and Winston, Toronto, 1971.
- [14] Horvitz, D. G. and Thompson, D. J. A generalization of sampling without replacement from a finite universe. *Journal of the American Statistical Association*, 47(260):663–685, 1952.
- [15] Jia, R., Dao, D., Wang, B., et al. Towards efficient data valuation based on the shapley value. *AISTATS*, abs/1902.10275:1167–1176, 2019.
- [16] Kolpaczki, P., Bengs, V., Muschalik, M., et al. Approximating the shapley value without marginal contributions. *Proceedings of the AAAI Conference on Artificial Intelligence*, 38(12):13246–13255, 2024. doi:10.1609/aaai.v38i12.29225.
- [17] Kwon, Y. and Zou, J. Y. Beta shapley: A unified and noise-reduced data valuation framework for machine learning. *International Conference on Artificial Intelligence and Statistics*, 151:8780–8802, 2021.
- [18] Lee, K., Liu, Z., Tang, W., et al. Faithful group shapley value. In *The Thirty-ninth Annual Conference on Neural Information Processing Systems*. 2025.
- [19] Li, W. and Yu, Y. Faster approximation of probabilistic and distributional values via least squares. In B. Kim, Y. Yue, S. Chaudhuri, K. Fragkiadaki, M. Khan, and

- Y. Sun, editors, *International Conference on Representation Learning*, volume 2024, pages 51182–51216. 2024.
- [20] Li, W. and Yu, Y. One sample fits all: Approximating all probabilistic values simultaneously and efficiently. *Neural Information Processing Systems*, abs/2410.23808, 2024.
 - [21] Lin, J., Zhang, A., Lécuyer, M., et al. Measuring the effect of training data on deep learning predictions via randomized experiments. In K. Chaudhuri, S. Jegelka, L. Song, C. Szepesvari, G. Niu, and S. Sabato, editors, *Proceedings of the 39th International Conference on Machine Learning*, volume 162 of *Proceedings of Machine Learning Research*, pages 13468–13504. PMLR, 2022.
 - [22] Lundberg, S. M., Erion, G. G., and Lee, S.-I. Consistent individualized feature attribution for tree ensembles. *arXiv [cs.LG]*, 2018.
 - [23] Lundberg, S. M. and Lee, S.-I. A unified approach to interpreting model predictions. *Neural Information Processing Systems*, pages 4765–4774, 2017. doi:10.5555/3295222.3295230.
 - [24] Maleki, S., Tran-Thanh, L., Hines, G., et al. Bounding the estimation error of sampling-based shapley value approximation. *arXiv [cs.GT]*, 2013.
 - [25] Moehle, N., Boyd, S., and Ang, A. Portfolio performance attribution via shapley value. 2021.
 - [26] Musco, C. and Witter, R. T. Provably accurate shapley value estimation via leverage score sampling. *ArXiv*, abs/2410.01917, 2024. doi:10.48550/arXiv.2410.01917.
 - [27] Neyman, J. On the two different aspects of the representative method : The method of stratified sampling and the method of purposive selection. *Journal of the Royal Statistical Society*, 97(4):558–606, 1934.
 - [28] Rubin, D. B. Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology*, 66(5):688–701, 1974.
 - [29] Särndal, Swensson, B., and Wretman, J. *Model assisted survey sampling*. Springer Series in Statistics. Springer, New York, NY, 1992.
 - [30] Shapley, L. S. A value for n-person games. 2(28):307–317, 1953.
 - [31] Wang, J. T. and Jia, R. Data banzhaf: A robust data valuation framework for machine learning. *AISTATS*, 206:6388–6421, 2022.

- [32] Weber, R. J. Probabilistic values for games. In A. E. Roth, editor, *The Shapley Value: Essays in Honor of Lloyd S. Shapley*, pages 101–120. Cambridge University Press, Cambridge, 1988.
- [33] Witter, R. T., Liu, Y., and Musco, C. Regression-adjusted monte carlo estimators for shapley values and probabilistic values. In *Neural Information Processing Systems*. 2025.
- [34] Wu, M., Jia, R., Lin, C., et al. Variance reduced shapley value estimation for trustworthy data valuation. *Computers & Operations Research*, 159:106305, 2023. doi:10.1016/j.cor.2023.106305.
- [35] Zhang, J., Sun, Q., Liu, J., et al. Efficient sampling approaches to shapley value approximation. *Proceedings of the ACM on Management of Data*, 1(1):1–24, 2023. doi:10.1145/3588728.

SUPPLEMENTARY MATERIAL

Contents

A	Proofs	1
A.1	Proof of Theorem 1	1
A.2	Proof of Theorem 2	5
A.2.1	Proof of Lemma A.1	10
A.3	Proofs of Other Theoretical Results	13
A.3.1	Proof of Proposition 1	13
A.3.2	Proof of Proposition 2	14
A.3.3	Proof of Proposition 3	15
A.3.4	Proof of Lemma 1	15
B	Details for Non-Asymptotic Results	16
B.1	Details for Non-Asymptotic Results for Hájek Estimation	16
B.2	Details for Non-Asymptotic Results for Ridge Regression Estimation	19
C	Details of Algorithms	21
C.1	Initialization for a Fixed Sampling Partition	22
C.2	Flooring the Learned Sampling Law	22
C.3	Shared Linear Surrogate Learned On the Fly	23
C.4	Complement-Pair Sampling	24
D	Additional Experimental Results	25
E	Monte Carlo Estimation with Bundled Samples	25
E.1	Bundled Sampling and Identification	25
E.2	Bundle-Regular Estimators	27
E.3	First-Order Risk with a Fixed Bundle Weighting	29
E.4	Rao–Blackwell Weighting for Complement Pairs	30

A Proofs

A.1 Proof of Theorem 1

Proof. We prove the result for the scalar Hájek estimator $\hat{\tau}_{\mathbf{a},m}^{\text{Hájek}}(u)$ in (10). The proof has four steps. We first recall the cellwise surrogate from Proposition 1, then derive the exact remainder decomposition, and finally bound the normalization and missing-cell pieces separately.

Step 1: Recall the cellwise centering. For each cell C_k with $\pi_k > 0$, define

$$\mu_k := \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)u(S) \mid S \in C_k] = \frac{1}{\pi_k} \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)u(S)\mathbf{1}\{S \in C_k\}].$$

For cells with $\pi_k = 0$, set $\mu_k := 0$. By the compatibility condition in Proposition 1, on each cell C_k the weight $\gamma_{\mathbf{a},q}$ is either identically zero or everywhere nonzero. Hence the cellwise

surrogate $h_{\mathcal{C}}$ satisfies

$$\gamma_{\mathbf{a},q}(S)h_{\mathcal{C}}(S) = \sum_{k=1}^K \mu_k \mathbf{1}\{S \in C_k\},$$

with the convention $h_{\mathcal{C}}(S) = 0$ on cells where $\gamma_{\mathbf{a},q} \equiv 0$.

Taking expectations gives

$$\mathbb{E}_q[\gamma_{\mathbf{a},q}(S)h_{\mathcal{C}}(S)] = \sum_{k=1}^K \pi_k \mu_k = \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)u(S)] = \tau_{\mathbf{a}}(u),$$

where the last identity is the weighted-average identification (4). Therefore $\tau_{\mathbf{a}}(h_{\mathcal{C}}) = \tau_{\mathbf{a}}(u)$, and the influence-function term in Definition 1 simplifies to

$$\psi_{h_{\mathcal{C}}}(S; u) = \gamma_{\mathbf{a},q}(S)u(S) - \mu_{k(S)}.$$

Step 2: Decompose the estimator error. Let

$$\xi_t := \psi_{h_{\mathcal{C}}}(S_t; u) = \gamma_{\mathbf{a},q}(S_t)u(S_t) - \mu_{k(S_t)},$$

where $k(S_t)$ denotes the cell index of S_t . By construction,

$$\mathbb{E}(\xi_t \mid S_t \in C_k) = 0 \quad \text{for every } k.$$

Set

$$A_{mk} := \frac{1}{m} \sum_{t=1}^m \mathbf{1}\{S_t \in C_k\} \xi_t,$$

and, only within this proof,

$$\hat{\lambda}_k := \begin{cases} \pi_k / \hat{\pi}_k, & N_k > 0, \\ 0, & N_k = 0. \end{cases}$$

Define

$$r_{m,\text{norm}} := \sum_{k=1}^K (\hat{\lambda}_k - 1) A_{mk}, \quad r_{m,\text{miss}} := \sum_{k:N_k=0} \pi_k \mu_k.$$

Also, by the definition of $\hat{\lambda}_k$,

$$\hat{\lambda}_k \hat{\pi}_k = \pi_k \mathbf{1}\{N_k > 0\}.$$

Using the definition of $\hat{\tau}_{\mathbf{a},m}^{\text{Hajek}}(u)$, the identity

$$\hat{\pi}_k = \frac{1}{m} \sum_{t=1}^m \mathbf{1}\{S_t \in C_k\},$$

and $\tau_{\mathbf{a}}(u) = \sum_{k=1}^K \pi_k \mu_k$, we obtain

$$\begin{aligned}
\hat{\tau}_{\mathbf{a},m}^{\text{Hajek}}(u) - \tau_{\mathbf{a}}(u) &= \sum_{k=1}^K \hat{\lambda}_k \frac{1}{m} \sum_{t=1}^m \mathbf{1}\{S_t \in C_k\} \gamma_{\mathbf{a},q}(S_t) u(S_t) - \sum_{k=1}^K \pi_k \mu_k \\
&= \sum_{k=1}^K \hat{\lambda}_k \frac{1}{m} \sum_{t=1}^m \mathbf{1}\{S_t \in C_k\} \left\{ \gamma_{\mathbf{a},q}(S_t) u(S_t) - \mu_k \right\} + \sum_{k=1}^K (\hat{\lambda}_k \hat{\pi}_k - \pi_k) \mu_k \\
&= \sum_{k=1}^K \hat{\lambda}_k A_{mk} - r_{m,\text{miss}} \\
&= \sum_{k=1}^K A_{mk} + r_{m,\text{norm}} - r_{m,\text{miss}} \\
&= \frac{1}{m} \sum_{t=1}^m \xi_t + r_{m,\text{norm}} - r_{m,\text{miss}}.
\end{aligned}$$

Since $\xi_t = \psi_{h_{\mathcal{C}}}(S_t; u)$, this yields the exact decomposition

$$\hat{\tau}_{\mathbf{a},m}^{\text{Hajek}}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_{h_{\mathcal{C}}}(S_t; u) + r_m, \quad r_m := r_{m,\text{norm}} - r_{m,\text{miss}}. \quad (23)$$

Step 3: Bound the normalization remainder. Write the average cellwise second moment in the theorem display as

$$\Gamma_{2,\mathcal{C}}^2 := \frac{1}{K_+} \sum_{k:\pi_k > 0} \mathbb{E}_q \left[\gamma_{\mathbf{a},q}(S)^2 \mid S \in C_k \right], \quad \Gamma_{\infty} := \sup_{S:q(S) > 0} |\gamma_{\mathbf{a},q}(S)|.$$

Cells with $\pi_k = 0$ satisfy $N_k = 0$ and $A_{mk} = 0$ almost surely, so only cells with $\pi_k > 0$ contribute to the bounds.

For $N_k > 0$, write

$$\bar{\xi}_{mk} := \frac{1}{N_k} \sum_{t:S_t \in C_k} \xi_t,$$

and set $\bar{\xi}_{mk} := 0$ when $N_k = 0$. Then $A_{mk} = \hat{\pi}_k \bar{\xi}_{mk}$, and the normalization factor cancels:

$$(\hat{\lambda}_k - 1) A_{mk} = (\pi_k - \hat{\pi}_k) \bar{\xi}_{mk}.$$

When $N_k = 0$, both sides are zero by definition. When $N_k > 0$,

$$(\hat{\lambda}_k - 1) A_{mk} = \left(\frac{\pi_k}{\hat{\pi}_k} - 1 \right) \hat{\pi}_k \bar{\xi}_{mk} = (\pi_k - \hat{\pi}_k) \bar{\xi}_{mk}.$$

Let $\mathcal{F}_m$ be the sigma-field generated by the cell labels $\{k(S_t) : 1 \leq t \leq m\}$. Conditional on

$\mathcal{F}_m$, the within-cell averages $\bar{\xi}_{mk}$ have mean zero, and the cross terms vanish. Hence

$$\mathbb{E}(r_{m,\text{norm}}^2) = \sum_{k:\pi_k>0} \mathbb{E}[(\pi_k - \hat{\pi}_k)^2 \bar{\xi}_{mk}^2].$$

Moreover, conditional on N_k , the N_k observations falling in C_k are i.i.d. from $q(\cdot | C_k)$, so

$$\begin{aligned} \mathbb{E}(\bar{\xi}_{mk}^2 | N_k) &= \frac{1}{N_k} \text{Var}_q\{\gamma_{\mathbf{a},q}(S)u(S) | S \in C_k\} \mathbf{1}\{N_k > 0\} \\ &\leq \frac{\|u\|_\infty^2}{N_k} \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)^2 | S \in C_k] \mathbf{1}\{N_k > 0\}. \end{aligned}$$

For $N_k \sim \text{Bin}(m, \pi_k)$, the following binomial moment bound is uniform in m and k :

$$\mathbb{E}\left[\frac{(N_k - m\pi_k)^2}{N_k} \mathbf{1}\{N_k > 0\}\right] \leq C.$$

Indeed, on $\{N_k \geq m\pi_k/2\}$, the expectation is bounded by $2(m\pi_k)^{-1}\text{Var}(N_k) \leq 2$, while on $\{0 < N_k < m\pi_k/2\}$ it is bounded by $(m\pi_k)^2 P(N_k < m\pi_k/2) \leq (m\pi_k)^2 e^{-m\pi_k/8} \leq C$. Therefore,

$$\mathbb{E}[(\pi_k - \hat{\pi}_k)^2 \bar{\xi}_{mk}^2] \leq \frac{C\|u\|_\infty^2}{m^2} \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)^2 | S \in C_k],$$

and hence

$$\mathbb{E}(r_{m,\text{norm}}^2) \leq \frac{C\|u\|_\infty^2 K_+ \Gamma_{2,\mathcal{C}}^2}{m^2}.$$

Step 4: Bound the missing-cell remainder. Since $P(N_k = 0) = (1 - \pi_k)^m \leq e^{-m\pi_k}$,

$$|r_{m,\text{miss}}| \leq \|u\|_\infty \Gamma_\infty \sum_{k:\pi_k>0} \pi_k \mathbf{1}\{N_k = 0\}.$$

Because $\sum_{k:\pi_k>0} \pi_k \mathbf{1}\{N_k = 0\} \leq 1$, we have

$$\mathbb{E}(r_{m,\text{miss}}^2) \leq \|u\|_\infty^2 \Gamma_\infty^2 \sum_{k:\pi_k>0} \pi_k P(N_k = 0) \leq \|u\|_\infty^2 \Gamma_\infty^2 e^{-m\pi_{\min}}.$$

Combining the two bounds with $(x + y)^2 \leq 2x^2 + 2y^2$, and absorbing numerical constants into $C_1, C_2 > 0$, yields

$$\begin{aligned} \mathbb{E}(r_m^2) &\leq 2\mathbb{E}(r_{m,\text{norm}}^2) + 2\mathbb{E}(r_{m,\text{miss}}^2) \\ &\leq C_1 \|u\|_\infty^2 \left\{ \frac{K_+ \Gamma_{2,\mathcal{C}}^2}{m^2} + \Gamma_\infty^2 e^{-C_2 m\pi_{\min}} \right\} \end{aligned}$$

for universal constants $C_1, C_2 > 0$. This is (16). □

A.2 Proof of Theorem 2

The proof has two parts. We first establish a same-sample product bound that controls the interaction between the empirical Gram fluctuation and the empirical residual score. We then apply this lemma to the whitened WLS design to bound the remainder vector $\mathbf{r}_m$.

The good-event analysis below involves the product of two empirical averages computed from the same sampled coalitions. The next lemma isolates the bound needed for that product. We state it first and defer its proof to the end of this subsection.

Lemma A.1 (Same-sample covariance-score product). *Let $(\mathbf{x}, \mathbf{b})$ be a random pair in an r -dimensional Euclidean space. Assume*

$$\mathbb{E}(\mathbf{x}\mathbf{x}^\top) = \mathbf{I}_r, \quad \|\mathbf{x}\|_2^2 \leq L, \quad \mathbb{E}(\mathbf{b}) = 0, \quad M_{\mathbf{b}} := \mathbb{E}\|\mathbf{b}\|_2^2 < \infty.$$

For iid copies $(\mathbf{x}_t, \mathbf{b}_t)_{t=1}^m$, define

$$\bar{\mathbf{B}}_m := \frac{1}{m} \sum_{t=1}^m (\mathbf{x}_t \mathbf{x}_t^\top - \mathbf{I}_r), \quad \bar{\mathbf{b}}_m := \frac{1}{m} \sum_{t=1}^m \mathbf{b}_t.$$

Then, for a numerical constant $C > 0$,

$$\mathbb{E} \left[\|\bar{\mathbf{B}}_m\|_{\text{op}}^2 \|\bar{\mathbf{b}}_m\|_2^2 \right] \leq C \left\{ \frac{L \log(er)}{m^2} + \frac{L^2 \log^2(er)}{m^3} \right\} M_{\mathbf{b}}. \quad (24)$$

Proof of Theorem 2. We organize the proof into four steps. The first step identifies the common first-order term and the vector remainder. The remaining steps bound that remainder on a good conditioning event and on its complement.

Step 1: Setup and first-order decomposition. Fix m , write $\rho := \lambda/m$, and let

$$\mathbf{M} := \text{range}(\mathbf{G}).$$

All inverses below are taken on $\mathbf{M}$. Let $\boldsymbol{\beta}^\star$ be the minimum-Euclidean-norm element of

$$\arg \min_{\boldsymbol{\beta} \in \mathbb{R}^{d_{\mathbf{z}}}} \mathbb{E}_q \left[\tilde{w}(S) \{u(S) - \mathbf{z}(S)^\top \boldsymbol{\beta}\}^2 \right],$$

and define

$$h^\star(S) := \mathbf{z}(S)^\top \boldsymbol{\beta}^\star, \quad \varepsilon(S) := u(S) - h^\star(S).$$

Also define

$$\begin{aligned}\mathbf{b}_u &:= \mathbb{E}_q[\tilde{w}(S)\mathbf{z}(S)u(S)], & \hat{\mathbf{G}}_m &:= \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)\mathbf{z}(S_t)\mathbf{z}(S_t)^\top, \\ \hat{\mathbf{b}}_m &:= \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)\mathbf{z}(S_t)u(S_t), & \hat{\delta} &:= \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)\mathbf{z}(S_t)\varepsilon(S_t).\end{aligned}$$

Since $\mathbf{b}_u \in \mathbf{M}$, the minimum-norm normal-equation solution is

$$\boldsymbol{\beta}^\star = \mathbf{G}^+ \mathbf{b}_u, \quad \mathbb{E}_q[\tilde{w}(S)\mathbf{z}(S)\varepsilon(S)] = 0.$$

Let $\mathbf{a}_j$ and $\mathbf{c}_j$ denote the j -th columns of $\mathbf{A}$ and $\mathbf{C}_\mathbf{A}$, respectively. For an arbitrary set function v , let $\boldsymbol{\beta}^\star(v)$ be the minimum-norm population WLS solution. Then

$$\boldsymbol{\beta}^\star(v) = \mathbf{G}^+ \mathbb{E}_q[\tilde{w}(S)\mathbf{z}(S)v(S)].$$

By the assumed WLS identification, this implies

$$\tau_{\mathbf{a}_j}(v) = \mathbf{c}_j^\top \mathbf{G}^+ \mathbb{E}_q[\tilde{w}(S)\mathbf{z}(S)v(S)] \quad \text{for every } j \text{ and every } v.$$

Comparing with the weighted-average identification (4) for the scalar target $\tau_{\mathbf{a}_j}$ yields

$$\gamma_{\mathbf{a}_j, q}(S) = \tilde{w}(S)\mathbf{z}(S)^\top \mathbf{G}^+ \mathbf{c}_j \quad \text{for every } S \text{ with } q(S) > 0. \quad (25)$$

Applying the same readout identity to $v = h^\star$ gives

$$\boldsymbol{\tau}_\mathbf{A}(h^\star) = \mathbf{C}_\mathbf{A}^\top \boldsymbol{\beta}^\star = \boldsymbol{\tau}_\mathbf{A}(u),$$

so $h^\star$ is the common surrogate for all coordinates.

Since $u(S) = \mathbf{z}(S)^\top \boldsymbol{\beta}^\star + \varepsilon(S)$, the empirical right-hand side in the ridge normal equation decomposes as

$$\begin{aligned}\hat{\mathbf{b}}_m &= \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)\mathbf{z}(S_t)u(S_t) \\ &= \left\{ \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)\mathbf{z}(S_t)\mathbf{z}(S_t)^\top \right\} \boldsymbol{\beta}^\star + \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)\mathbf{z}(S_t)\varepsilon(S_t) \\ &= \hat{\mathbf{G}}_m \boldsymbol{\beta}^\star + \hat{\delta}.\end{aligned}$$

The ridge estimator satisfies

$$(\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z}) \hat{\boldsymbol{\beta}}_{m, \lambda} = \hat{\mathbf{b}}_m.$$

Therefore, using $\hat{\mathbf{G}}_m = (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z}) - \rho \mathbf{I}_{d_z}$,

$$\begin{aligned}\hat{\beta}_{m,\lambda} - \beta^* &= (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} (\hat{\mathbf{G}}_m \beta^* + \hat{\delta}) - \beta^* \\ &= \left[(\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} \hat{\mathbf{G}}_m - \mathbf{I}_{d_z} \right] \beta^* + (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} \hat{\delta} \\ &= (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} \hat{\delta} - \rho (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} \beta^*.\end{aligned}$$

Adding and subtracting the population linear term $\mathbf{C}_A^\top \mathbf{G}^+ \hat{\delta}$ in the target error gives

$$\begin{aligned}\hat{\tau}_{\mathbf{A},m}^{\text{WLS},\lambda} - \tau_{\mathbf{A}}(u) &= \mathbf{C}_A^\top \{ \hat{\beta}_{m,\lambda} - \beta^* \} \\ &= \mathbf{C}_A^\top (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} \hat{\delta} - \rho \mathbf{C}_A^\top (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} \beta^* \\ &= \mathbf{C}_A^\top \mathbf{G}^+ \hat{\delta} + \underbrace{\mathbf{C}_A^\top \left[(\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} - \mathbf{G}^+ \right] \hat{\delta} - \rho \mathbf{C}_A^\top (\hat{\mathbf{G}}_m + \rho \mathbf{I}_{d_z})^{-1} \beta^*}_{=: \mathbf{r}_m}.\end{aligned}$$

For each coordinate j , (25) gives

$$\mathbf{c}_j^\top \mathbf{G}^+ \hat{\delta} = \frac{1}{m} \sum_{t=1}^m \gamma_{\mathbf{a}_j,q}(S_t) \varepsilon(S_t) = \frac{1}{m} \sum_{t=1}^m \psi_{\mathbf{a}_j,q,h^*}(S_t; u).$$

Thus each coordinate has the regular expansion with surrogate h^* , and $\mathbf{r}_m$ is the common remainder vector. It remains to bound $\mathbb{E}_q(\|\mathbf{r}_m\|_2^2)$.

Step 2: Reduce the problem to a whitened coefficient error. For every S with $\tilde{w}(S) > 0$, we have $\mathbf{z}(S) \in \mathbf{M}$. Hence $\text{range}(\hat{\mathbf{G}}_m) \subseteq \mathbf{M}$, while $\hat{\delta}$ and β^* lie in $\mathbf{M}$. Let $r = \text{rank}(\mathbf{G})$, and choose an orthonormal basis $\mathbf{U} \in \mathbb{R}^{d_z \times r}$ for $\mathbf{M}$. Define

$$\mathbf{G}_M := \mathbf{U}^\top \mathbf{G} \mathbf{U}, \quad \hat{\mathbf{G}}_{M,m} := \mathbf{U}^\top \hat{\mathbf{G}}_m \mathbf{U}.$$

The matrix $\mathbf{G}_M$ is positive definite, and its eigenvalues are the nonzero eigenvalues of $\mathbf{G}$. Work in this r -dimensional coordinate system and define

$$\mathbf{H}_m := \mathbf{G}_M^{-1/2} \hat{\mathbf{G}}_{M,m} \mathbf{G}_M^{-1/2}, \quad \mathbf{y}_m := \mathbf{G}_M^{-1/2} \mathbf{U}^\top \hat{\delta}, \quad \boldsymbol{\theta}_m := \mathbf{G}_M^{1/2} \mathbf{U}^\top \beta^*.$$

Let

$$\mathcal{E}_m := \left\{ \|\mathbf{H}_m - \mathbf{I}_r\|_{\text{op}} \leq \frac{1}{2} \right\}.$$

In these coordinates, define the r -vector

$$\boldsymbol{\Delta}_m := \left[(\mathbf{H}_m + \rho \mathbf{G}_M^{-1})^{-1} - \mathbf{I}_r \right] \mathbf{y}_m - \rho (\mathbf{H}_m + \rho \mathbf{G}_M^{-1})^{-1} \mathbf{G}_M^{-1} \boldsymbol{\theta}_m.$$

Using $\mathbf{G}^+ = \mathbf{U}\mathbf{G}_M^{-1}\mathbf{U}^\top$ and the fact that $\hat{\delta}, \beta^* \in \mathbf{M}$, converting back from coordinates gives

$$\mathbf{r}_m = \mathbf{C}_A^\top \mathbf{U} \mathbf{G}_M^{-1/2} \Delta_m.$$

Consequently,

$$\|\mathbf{r}_m\|_2^2 \leq \sigma_{\max}(\mathbf{C}_A^\top \mathbf{U} \mathbf{G}_M^{-1} \mathbf{U}^\top \mathbf{C}_A) \|\Delta_m\|_2^2 = R_A^2 \|\Delta_m\|_2^2.$$

Therefore it suffices to bound $\mathbb{E}\|\Delta_m\|_2^2$.

On $\mathcal{E}_m$, the matrix $\mathbf{H}_m + \rho \mathbf{G}_M^{-1}$ has minimum eigenvalue at least $1/2$, so

$$\left\| \left(\mathbf{H}_m + \rho \mathbf{G}_M^{-1} \right)^{-1} - \mathbf{I}_r \right\|_{\text{op}} \leq 2 \left\{ \|\mathbf{H}_m - \mathbf{I}_r\|_{\text{op}} + \rho \|\mathbf{G}_M^{-1}\|_{\text{op}} \right\},$$

and

$$\left\| \left(\mathbf{H}_m + \rho \mathbf{G}_M^{-1} \right)^{-1} \right\|_{\text{op}} \leq 2.$$

Thus, on $\mathcal{E}_m$,

$$\|\Delta_m\|_2^2 \mathbf{1}(\mathcal{E}_m) \leq C \left[\|\mathbf{H}_m - \mathbf{I}_r\|_{\text{op}}^2 \|\mathbf{y}_m\|_2^2 + \frac{\lambda^2}{m^2 \{\sigma_{\min}^+(\mathbf{G})\}^2} \|\mathbf{y}_m\|_2^2 + \frac{\lambda^2}{m^2 \{\sigma_{\min}^+(\mathbf{G})\}^2} \|\boldsymbol{\theta}_m\|_2^2 \right].$$

Step 3: Bound the good event. Write

$$\mathbf{x}(S) := \tilde{w}(S)^{1/2} \mathbf{G}_M^{-1/2} \mathbf{U}^\top \mathbf{z}(S) \in \mathbb{R}^r.$$

Then $\mathbb{E}_q[\mathbf{x}(S)\mathbf{x}(S)^\top] = \mathbf{I}_r$ and $\|\mathbf{x}(S)\|_2^2 = \ell(S) \leq L$. Standard matrix Bernstein for bounded isotropic rows gives, after adjusting constants and using $d_z \geq 2$,

$$P(\mathcal{E}_m^c) \leq C \exp\{-cm/L + \log d_z\}.$$

Also,

$$\mathbf{H}_m - \mathbf{I}_r = \frac{1}{m} \sum_{t=1}^m \{\mathbf{x}(S_t)\mathbf{x}(S_t)^\top - \mathbf{I}_r\}, \quad \mathbf{y}_m = \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)^{1/2} \mathbf{x}(S_t) \varepsilon(S_t),$$

and both summands have mean zero. Applying Lemma A.1 in dimension $r \leq d_z$, with

$$\mathbf{b}(S) := \tilde{w}(S)^{1/2} \mathbf{x}(S) \varepsilon(S), \quad M_{\mathbf{b}} = \mathbb{E}_q[\tilde{w}(S) \ell(S) \varepsilon(S)^2],$$

gives

$$\mathbb{E}[\|\mathbf{H}_m - \mathbf{I}_r\|_{\text{op}}^2 \|\mathbf{y}_m\|_2^2] \leq \frac{CL \log d_{\mathbf{z}}}{m^2} \left(1 + \frac{L \log d_{\mathbf{z}}}{m}\right) \mathbb{E}_q[\tilde{w}(S) \ell(S) \varepsilon(S)^2].$$

Moreover,

$$\mathbb{E}\|\mathbf{y}_m\|_2^2 = \frac{1}{m} \mathbb{E}_q[\tilde{w}(S) \ell(S) \varepsilon(S)^2], \quad \|\boldsymbol{\theta}_m\|_2^2 = (\boldsymbol{\beta}^*)^\top \mathbf{G} \boldsymbol{\beta}^* \leq \mathbb{E}_q[\tilde{w}(S) u(S)^2],$$

where the last inequality is the Pythagorean property of weighted least squares. The same projection property also gives

$$\mathbb{E}_q[\tilde{w}(S) \ell(S) \varepsilon(S)^2] \leq L \mathbb{E}_q[\tilde{w}(S) \varepsilon(S)^2] \leq L \mathbb{E}_q[\tilde{w}(S) u(S)^2].$$

Taking expectations in the bound from Step 2, substituting the last three displays, and using $\lambda \asymp \sigma_{\min}^+(\mathbf{G})$, we obtain

$$\mathbb{E}[\|\boldsymbol{\Delta}_m\|_2^2 \mathbf{1}(\mathcal{E}_m)] \leq C \left[\frac{L \log d_{\mathbf{z}}}{m^2} \left(1 + \frac{L \log d_{\mathbf{z}}}{m}\right) \mathbb{E}_q[\tilde{w}(S) \ell(S) \varepsilon(S)^2] + \frac{\|u\|_{2, \tilde{w}}^2}{m^2} \right].$$

Because

$$\mathbb{E}_q[\tilde{w}(S) \ell(S) \varepsilon(S)^2] \leq L \|u\|_{2, \tilde{w}}^2,$$

the second term is absorbed into the first after adjusting constants. Hence

$$\mathbb{E}[\|\boldsymbol{\Delta}_m\|_2^2 \mathbf{1}(\mathcal{E}_m)] \leq \frac{CL^2 \log d_{\mathbf{z}}}{m^2} \left(1 + \frac{L \log d_{\mathbf{z}}}{m}\right) \|u\|_{2, \tilde{w}}^2.$$

Step 4: Bound the bad event. On $\mathcal{E}_m^c$, ridge still gives deterministic stability. Since

$$\left\| (\mathbf{H}_m + \rho \mathbf{G}_M^{-1})^{-1} \right\|_{\text{op}} \leq \frac{\sigma_{\max}(\mathbf{G}_M)}{\rho} = \frac{m \sigma_{\max}(\mathbf{G})}{\lambda},$$

we have

$$\left\| (\mathbf{H}_m + \rho \mathbf{G}_M^{-1})^{-1} - \mathbf{I}_r \right\|_{\text{op}} \leq 1 + \frac{m \sigma_{\max}(\mathbf{G})}{\lambda}.$$

Also, for every sample,

$$\|\mathbf{y}_m\|_2 \leq \frac{1}{m} \sum_{t=1}^m \tilde{w}(S_t)^{1/2} \|\mathbf{x}(S_t)\|_2 |\varepsilon(S_t)| \leq \left\{ L \sup_{S: q(S) > 0} \tilde{w}(S) \varepsilon(S)^2 \right\}^{1/2}.$$

Since $\varepsilon(S) = u(S) - \mathbf{z}(S)^\top \boldsymbol{\beta}^*$, for every S with $q(S) > 0$,

$$\tilde{w}(S) \varepsilon(S)^2 \leq 2 \tilde{w}(S) u(S)^2 + 2 \tilde{w}(S) \{\mathbf{z}(S)^\top \boldsymbol{\beta}^*\}^2.$$

Moreover,

$$\tilde{w}(S)\{\mathbf{z}(S)^\top \boldsymbol{\beta}^\star\}^2 \leq \ell(S)(\boldsymbol{\beta}^\star)^\top \mathbf{G} \boldsymbol{\beta}^\star \leq L \mathbb{E}_q[\tilde{w}(T)u(T)^2],$$

again by the Pythagorean property of weighted least squares. Hence

$$\sup_{S:q(S)>0} \tilde{w}(S)\varepsilon(S)^2 \leq 2\|u\|_{\infty,\tilde{w}}^2 + 2L\|u\|_{2,\tilde{w}}^2.$$

Moreover, $L \geq 1$ in the nonzero-rank case because $\mathbb{E}_q\{\ell(S)\} = \text{rank}(\mathbf{G})$. Since $\|u\|_{2,\tilde{w}}^2 \leq \|u\|_{\infty,\tilde{w}}^2$, the preceding display is at most

$$4L\|u\|_{\infty,\tilde{w}}^2.$$

Therefore

$$\begin{aligned} \|\boldsymbol{\Delta}_m\|_2^2 \mathbf{1}(\mathcal{E}_m^c) &\leq C \left\{ \left(1 + \frac{m\sigma_{\max}(\mathbf{G})}{\lambda}\right)^2 L \sup_{S:q(S)>0} \tilde{w}(S)\varepsilon(S)^2 \right. \\ &\quad \left. + \left(\frac{\sigma_{\max}(\mathbf{G})}{\sigma_{\min}^+(\mathbf{G})}\right)^2 \|u\|_{2,\tilde{w}}^2 \right\} \mathbf{1}(\mathcal{E}_m^c). \end{aligned}$$

Taking expectations, using $\|u\|_{2,\tilde{w}}^2 \leq \|u\|_{\infty,\tilde{w}}^2$, $\lambda \asymp \sigma_{\min}^+(\mathbf{G})$, and the Bernstein bound for $P(\mathcal{E}_m^c)$, we obtain

$$\mathbb{E}[\|\boldsymbol{\Delta}_m\|_2^2 \mathbf{1}(\mathcal{E}_m^c)] \leq CL^2 e^{-cm/L} d_{\mathbf{z}} m^2 \kappa_{\mathbf{G}}^2 \|u\|_{\infty,\tilde{w}}^2,$$

where the factor L^2 comes from

$$L \sup_{S:q(S)>0} \tilde{w}(S)\varepsilon(S)^2 \leq 4L^2 \|u\|_{\infty,\tilde{w}}^2.$$

Combining the good- and bad-event bounds yields

$$\mathbb{E}\|\boldsymbol{\Delta}_m\|_2^2 \leq C \left[\frac{L^2 \log d_{\mathbf{z}}}{m^2} \left(1 + \frac{L \log d_{\mathbf{z}}}{m}\right) \|u\|_{2,\tilde{w}}^2 + L^2 e^{-cm/L} d_{\mathbf{z}} m^2 \kappa_{\mathbf{G}}^2 \|u\|_{\infty,\tilde{w}}^2 \right].$$

Finally,

$$\mathbb{E}_q[\|\mathbf{r}_m\|_2^2] \leq R_{\mathbf{A}}^2 \mathbb{E}\|\boldsymbol{\Delta}_m\|_2^2,$$

so (17) follows. $\square$

A.2.1 Proof of Lemma A.1

This lemma is a same-sample version of a standard decoupling estimate for second-order U -statistic-type sums; see, for example, the decoupling theory for U -statistics and U -processes

in de la Peña and Giné [8, Chapter 3]. We give a self-contained proof because only this elementary split-sample form is needed here.

Proof. Write

$$\mathbf{B}_t := \mathbf{x}_t \mathbf{x}_t^\top - \mathbf{I}_r.$$

We view $\bar{\mathbf{B}}_m \otimes \bar{\mathbf{b}}_m$ as an element of the injective tensor product with norm

$$\|\mathbf{T}\|_{\vee} := \sup_{\|a\|_2=\|b\|_2=\|c\|_2=1} |\mathbf{T}(a, b, c)|.$$

For a rank-one tensor $\mathbf{B} \otimes \mathbf{b}$, this norm is

$$\|\mathbf{B} \otimes \mathbf{b}\|_{\vee} = \|\mathbf{B}\|_{\text{op}} \|\mathbf{b}\|_2.$$

Hence the left-hand side of (24) equals

$$\mathbb{E} \|\bar{\mathbf{B}}_m \otimes \bar{\mathbf{b}}_m\|_{\vee}^2.$$

Decompose

$$\bar{\mathbf{B}}_m \otimes \bar{\mathbf{b}}_m = \frac{1}{m^2} \sum_{i \neq j} \mathbf{B}_i \otimes \mathbf{b}_j + \frac{1}{m^2} \sum_{i=1}^m \mathbf{B}_i \otimes \mathbf{b}_i =: \mathbf{U}_m + \mathbf{D}_m.$$

Step 1: Control the off-diagonal term. Let $\delta_1, \dots, \delta_m$ be iid Bernoulli(1/2), independent of the data, and define

$$\mathbf{U}_m(\delta) := \frac{1}{m^2} \sum_{i \neq j} \delta_i (1 - \delta_j) \mathbf{B}_i \otimes \mathbf{b}_j.$$

Since $\mathbb{E}_{\delta}[\delta_i(1 - \delta_j)] = 1/4$ for $i \neq j$, conditional on the data,

$$\mathbf{U}_m = 4\mathbb{E}_{\delta}\{\mathbf{U}_m(\delta)\}.$$

Jensen's inequality therefore gives

$$\mathbb{E} \|\mathbf{U}_m\|_{\vee}^2 \leq 16 \mathbb{E} \mathbb{E}_{\delta} \|\mathbf{U}_m(\delta)\|_{\vee}^2.$$

For a fixed split, let $I = \{i : \delta_i = 1\}$ and $J = \{j : \delta_j = 0\}$. Then $I \cap J = \emptyset$, so

$$\mathbf{U}_m(\delta) = \frac{1}{m^2} \left(\sum_{i \in I} \mathbf{B}_i \right) \otimes \left(\sum_{j \in J} \mathbf{b}_j \right).$$

The two sums involve disjoint observations and are therefore independent. Thus, for every fixed split,

$$\mathbb{E}\|\mathbf{U}_m(\delta)\|_{\vee}^2 = \frac{1}{m^4} \mathbb{E}\left\|\sum_{i \in I} \mathbf{B}_i\right\|_{\text{op}}^2 \mathbb{E}\left\|\sum_{j \in J} \mathbf{b}_j\right\|_2^2.$$

Because $\mathbb{E}\mathbf{b} = 0$,

$$\mathbb{E}\left\|\sum_{j \in J} \mathbf{b}_j\right\|_2^2 = |J|M_{\mathbf{b}} \leq mM_{\mathbf{b}}.$$

The integrated matrix Bernstein bound for bounded isotropic rows, applied to the deterministic subset I , gives

$$\mathbb{E}\left\|\sum_{i \in I} \mathbf{B}_i\right\|_{\text{op}}^2 \leq C \left\{|I|L \log(er) + L^2 \log^2(er)\right\} \leq C \left\{mL \log(er) + L^2 \log^2(er)\right\}.$$

Consequently, uniformly over the split,

$$\mathbb{E}\|\mathbf{U}_m(\delta)\|_{\vee}^2 \leq C \left\{\frac{L \log(er)}{m^2} + \frac{L^2 \log^2(er)}{m^3}\right\} M_{\mathbf{b}}.$$

Averaging over δ gives the same bound for $\mathbb{E}\|\mathbf{U}_m\|_{\vee}^2$.

Step 2: Control the diagonal term. We next bound $\mathbf{D}_m$. First,

$$\mathbf{D}_m = \frac{1}{m} \mathbb{E}(\mathbf{B} \otimes \mathbf{b}) + \frac{1}{m^2} \sum_{i=1}^m \{\mathbf{B}_i \otimes \mathbf{b}_i - \mathbb{E}(\mathbf{B} \otimes \mathbf{b})\}.$$

For the mean term, for any unit a, b, c ,

$$\begin{aligned} \left|\mathbb{E}\left[(a^\top \mathbf{B} \mathbf{b})(c^\top \mathbf{b})\right]\right| &= \left|\mathbb{E}\left[(a^\top \mathbf{x})(b^\top \mathbf{x})(c^\top \mathbf{b})\right]\right| \\ &\leq \left\{\mathbb{E}[(a^\top \mathbf{x})^2(b^\top \mathbf{x})^2]\right\}^{1/2} \left\{\mathbb{E}[(c^\top \mathbf{b})^2]\right\}^{1/2} \leq (LM_{\mathbf{b}})^{1/2}, \end{aligned}$$

where the term involving $\mathbf{I}_r$ vanishes because $\mathbb{E}\mathbf{b} = 0$. Therefore

$$\left\|\frac{1}{m} \mathbb{E}(\mathbf{B} \otimes \mathbf{b})\right\|_{\vee}^2 \leq \frac{LM_{\mathbf{b}}}{m^2}.$$

For the centered diagonal fluctuation, a direct second-moment bound is enough. Use $\|\cdot\|_{\vee} \leq \|\cdot\|_F$. Since $\mathbb{E}\|\mathbf{x}\|_2^2 = r$ and $\|\mathbf{x}\|_2^2 \leq L$, we have $r \leq L$. Hence

$$\|\mathbf{B}\|_F = \|\mathbf{x}\mathbf{x}^\top - \mathbf{I}_r\|_F \leq \|\mathbf{x}\mathbf{x}^\top\|_F + \|\mathbf{I}_r\|_F \leq L + \sqrt{r} \leq CL.$$

Therefore, using independence and centering in the Hilbert space equipped with the Frobenius norm,

$$\begin{aligned}
\mathbb{E} \left\| \frac{1}{m^2} \sum_{i=1}^m \{ \mathbf{B}_i \otimes \mathbf{b}_i - \mathbb{E}(\mathbf{B} \otimes \mathbf{b}) \} \right\|_{\vee}^2 &\leq \frac{1}{m^4} \mathbb{E} \left\| \sum_{i=1}^m \{ \mathbf{B}_i \otimes \mathbf{b}_i - \mathbb{E}(\mathbf{B} \otimes \mathbf{b}) \} \right\|_F^2 \\
&= \frac{1}{m^3} \mathbb{E} \left\| \mathbf{B} \otimes \mathbf{b} - \mathbb{E}(\mathbf{B} \otimes \mathbf{b}) \right\|_F^2 \\
&\leq \frac{1}{m^3} \mathbb{E} \left\| \mathbf{B} \otimes \mathbf{b} \right\|_F^2 \leq \frac{CL^2 M_{\mathbf{b}}}{m^3} \leq \frac{CL^2 \log^2(er) M_{\mathbf{b}}}{m^3}.
\end{aligned}$$

Thus

$$\mathbb{E} \left\| \mathbf{D}_m \right\|_{\vee}^2 \leq C \left\{ \frac{L \log(er)}{m^2} + \frac{L^2 \log^2(er)}{m^3} \right\} M_{\mathbf{b}},$$

where we used $\log(er) \geq 1$.

Step 3: Combine the two pieces. Combining the off-diagonal and diagonal bounds proves (24). $\square$

A.3 Proofs of Other Theoretical Results

A.3.1 Proof of Proposition 1

Proof. First verify that the proposed function belongs to the stated working class. For $S \in C_k$, if $\gamma_{\mathbf{a},q}$ is not identically zero on C_k , then the compatibility condition makes $\gamma_{\mathbf{a},q}(S) \neq 0$ throughout that cell, and the definition of $h_{\mathcal{C}}$ gives

$$\gamma_{\mathbf{a},q}(S) h_{\mathcal{C}}(S) = \mu_k.$$

If $\gamma_{\mathbf{a},q} \equiv 0$ on C_k , then $\mu_k = \mathbb{E}_q[\gamma_{\mathbf{a},q}(S)u(S) \mid S \in C_k] = 0$ when $\pi_k > 0$, and $\mu_k = 0$ by convention when $\pi_k = 0$. In this case $\gamma_{\mathbf{a},q}(S) h_{\mathcal{C}}(S) = 0 = \mu_k$. Hence

$$\gamma_{\mathbf{a},q}(S) h_{\mathcal{C}}(S) = \sum_{k=1}^K \mu_k \mathbf{1}\{S \in C_k\},$$

so $h_{\mathcal{C}} \in \mathcal{H}_{\mathcal{C},\gamma_{\mathbf{a},q}}$.

The proof of Theorem 1 gives the exact decomposition

$$\hat{\tau}_{\mathbf{a},m}^{\text{Hajek}}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_{h_{\mathcal{C}}}(S_t; u) + r_m$$

with the remainder bound in (16). In the fixed finite-player setting, K_+ , $\Gamma_{2,\mathcal{C}}$, Γ_{∞} , $\pi_{\min}$,

and $\|u\|_\infty$ do not vary with m , and $\pi_{\min} > 0$ by definition over the active cells. Therefore

$$\mathbb{E}_q(r_m^2) = O(m^{-2}) + O\{e^{-cm}\} = o(m^{-1})$$

for some $c > 0$. This is the regularity condition in Definition 1, relative to $\mathcal{H}_{\mathcal{C}, \gamma_{\mathbf{a}, q}}$. $\square$

A.3.2 Proof of Proposition 2

Proof. Apply Theorem 2 to the scalar target by taking $d = 1$, $\mathbf{A} = \mathbf{a}$, and $\mathbf{C}_{\mathbf{A}} = \mathbf{c}_{\mathbf{a}}$. The proof of that theorem constructs the population WLS projection

$$h^*(S) = \mathbf{z}(S)^\top \boldsymbol{\beta}^*,$$

where $\boldsymbol{\beta}^*$ is a minimum-Euclidean-norm population WLS solution, and establishes the expansion

$$\hat{\tau}_{\mathbf{a}, m}^{\text{WLS}, \lambda}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_{h^*}(S_t; u) + r_m.$$

The function h^* is an element of $\mathcal{H}_{\text{WLS}}$, and it is a population WLS projection of u onto this feature span. If the population projection is not unique, let $h_w(S) = \mathbf{z}(S)^\top \boldsymbol{\beta}_w$ be any population WLS projection. Then $\boldsymbol{\beta}_w - \boldsymbol{\beta}^*$ lies in the null space of the population Gram matrix, so

$$\sum_{S \subseteq [n]} w(S) \{\mathbf{z}(S)^\top (\boldsymbol{\beta}_w - \boldsymbol{\beta}^*)\}^2 = 0.$$

Thus $h_w(S) = h^*(S)$ for every S with $w(S) > 0$. The score identity derived in the proof of Theorem 2,

$$\gamma_{\mathbf{a}, q}(S) = \tilde{w}(S) \mathbf{z}(S)^\top \mathbf{G}^+ \mathbf{c}_{\mathbf{a}},$$

also implies that $\gamma_{\mathbf{a}, q}(S) = 0$ whenever $w(S) = 0$. Hence $\gamma_{\mathbf{a}, q}(S) \{h_w(S) - h^*(S)\} = 0$ for all S with $q(S) > 0$. The WLS readout identification gives $\tau_{\mathbf{a}}(h_w) = \tau_{\mathbf{a}}(h^*)$. Consequently,

$$\psi_{h_w}(S; u) = \psi_{h^*}(S; u) \quad q\text{-almost surely.}$$

The same expansion may therefore be written with h_w .

Under the fixed-design ridge regime used in Theorem 2, the displayed bound (17) gives

$$\mathbb{E}_q(r_m^2) = O(m^{-2}) + O\{m^2 e^{-cm}\} = o(m^{-1})$$

for some $c > 0$. Therefore $\hat{\tau}_{\mathbf{a}, m}^{\text{WLS}, \lambda}$ is regular relative to $\mathcal{H}_{\text{WLS}}$. $\square$

A.3.3 Proof of Proposition 3

Proof. Let $g := u - h^{\text{sur}}$. By linearity of $\tau_{\mathbf{a}}$,

$$\tau_{\mathbf{a}}(u) = \tau_{\mathbf{a}}(h^{\text{sur}}) + \tau_{\mathbf{a}}(g).$$

Since $\hat{\tau}_{\mathbf{a},m}$ is regular relative to $\mathcal{H}_m^{\text{imp}}$, there exists $h_m^{\text{imp}} \in \mathcal{H}_m^{\text{imp}}$ such that, when the estimator is applied to the residual utility g ,

$$\hat{\tau}_{\mathbf{a},m}(g) - \tau_{\mathbf{a}}(g) = \frac{1}{m} \sum_{t=1}^m \psi_{h_m^{\text{imp}}}(S_t; g) + r_m, \quad \mathbb{E}_q(r_m^2) = o(m^{-1}).$$

The surrogate-adjusted estimator in (13) therefore satisfies

$$\begin{aligned} \hat{\tau}_{\mathbf{a},m}^{\text{sur}} - \tau_{\mathbf{a}}(u) &= \hat{\tau}_{\mathbf{a},m}(g) - \tau_{\mathbf{a}}(g) \\ &= \frac{1}{m} \sum_{t=1}^m \psi_{h_m^{\text{imp}}}(S_t; g) + r_m. \end{aligned}$$

It remains to identify the score. For each sampled coalition S ,

$$\begin{aligned} \psi_{h_m^{\text{imp}}}(S; g) &= \gamma_{\mathbf{a},q}(S) \{u(S) - h^{\text{sur}}(S) - h_m^{\text{imp}}(S)\} + \tau_{\mathbf{a}}(h_m^{\text{imp}}) - \tau_{\mathbf{a}}(u - h^{\text{sur}}) \\ &= \gamma_{\mathbf{a},q}(S) \{u(S) - [h^{\text{sur}}(S) + h_m^{\text{imp}}(S)]\} + \tau_{\mathbf{a}}(h^{\text{sur}} + h_m^{\text{imp}}) - \tau_{\mathbf{a}}(u) \\ &= \psi_{h^{\text{sur}} + h_m^{\text{imp}}}(S; u). \end{aligned}$$

Because $h^{\text{sur}} + h_m^{\text{imp}} \in \mathcal{H}_m^{\text{sur}} + \mathcal{H}_m^{\text{imp}}$, the adjusted estimator is regular relative to the stated sum class. $\square$

A.3.4 Proof of Lemma 1

Proof. For $t = 1, \dots, m$, define the d -vector

$$\Psi_t := \left(\psi_{\mathbf{a}_1, q, h_1}(S_t; u), \dots, \psi_{\mathbf{a}_d, q, h_d}(S_t; u) \right)^\top,$$

and set

$$\bar{\Psi}_m := \frac{1}{m} \sum_{t=1}^m \Psi_t.$$

Each coordinate has mean zero because

$$\mathbb{E}_q \{ \psi_{\mathbf{a}_j, q, h_j}(S; u) \} = \mathbb{E}_q [\gamma_{\mathbf{a}_j, q}(S) \{u(S) - h_j(S)\}] + \tau_{\mathbf{a}_j}(h_j) - \tau_{\mathbf{a}_j}(u) = 0.$$

The coordinatewise expansion (14) gives

$$\hat{\tau}_{\mathbf{A},m}(u) - \tau_{\mathbf{A}}(u) = \bar{\Psi}_m + \mathbf{r}_m.$$

Since the samples are iid and each coordinate of Ψ_t is centered,

$$\begin{aligned} \mathbb{E}_q \|\bar{\Psi}_m\|_2^2 &= \sum_{j=1}^d \mathbb{E}_q \left[\left(\frac{1}{m} \sum_{t=1}^m \psi_{\mathbf{a}_j, q, h_j}(S_t; u) \right)^2 \right] \\ &= \frac{1}{m} \sum_{j=1}^d \text{Var}_q[\gamma_{\mathbf{a}_j, q}(S)\{u(S) - h_j(S)\}] = \frac{V(\mathbf{A}; q, \mathbf{h})}{m}. \end{aligned}$$

The assumption $V^*(\mathbf{A}; q) > 0$ ensures that $V(\mathbf{A}; q, \mathbf{h}) > 0$, so the ratio below is well defined.

Using the decomposition $\|\bar{\Psi}_m + \mathbf{r}_m\|_2^2 - \|\bar{\Psi}_m\|_2^2 = 2\bar{\Psi}_m^\top \mathbf{r}_m + \|\mathbf{r}_m\|_2^2$ and Cauchy's inequality,

$$|\mathbb{E}_q \|\bar{\Psi}_m + \mathbf{r}_m\|_2^2 - \mathbb{E}_q \|\bar{\Psi}_m\|_2^2| \leq 2 \left\{ \mathbb{E}_q \|\bar{\Psi}_m\|_2^2 \right\}^{1/2} \left\{ \mathbb{E}_q \|\mathbf{r}_m\|_2^2 \right\}^{1/2} + \mathbb{E}_q \|\mathbf{r}_m\|_2^2.$$

Dividing both sides by $V(\mathbf{A}; q, \mathbf{h})/m$ yields (15).

Finally, if

$$\mathbb{E}_q \|\mathbf{r}_m\|_2^2 \leq C\epsilon^2 \frac{V(\mathbf{A}; q, \mathbf{h})}{m}$$

with a sufficiently small universal constant C , then the right-hand side of (15) is at most ϵ for every $0 < \epsilon \leq 1$. This proves the stated consequence. $\square$

B Details for Non-Asymptotic Results

B.1 Details for Non-Asymptotic Results for Hájek Estimation

We spell out the specialization of Theorem 1 to the Shapley value of a fixed player i . For the Shapley value, the weighted-average coefficient in (3) is

$$\rho_i(S) = \begin{cases} \frac{1}{|S| \binom{n}{|S|}}, & i \in S, 1 \leq |S| \leq n, \\ -\frac{1}{(n - |S|) \binom{n}{|S|}}, & i \notin S, 0 \leq |S| \leq n - 1. \end{cases} \quad (26)$$

The endpoint coalitions $\emptyset$ and $[n]$ are deterministic in the usual Shapley-value implementations, and their contribution $\{u([n]) - u(\emptyset)\}/n$ can be added without sampling error. Hence the rate calculations below concern the random part supported on the interior strata $1 \leq |S| \leq n - 1$.

Both OFA and Stratified SVARM use self-normalization over cells indexed by membership of player i and coalition size. For $s = 1, \dots, n-1$, define

$$C_{i,s}^+ := \{S \subseteq [n] : i \in S, |S| = s\}, \quad C_{i,s}^- := \{S \subseteq [n] : i \notin S, |S| = s\}.$$

Cells that are empty, such as $C_{i,n}^-$ and $C_{i,0}^+$, are ignored. If the sampling law first draws a size s with probability q_s and then draws uniformly among coalitions of that size, then

$$q(S) = \frac{q_{|S|}}{\binom{n}{|S|}}, \quad \pi_{i,s}^+ := \mathbb{P}_q(S \in C_{i,s}^+) = \frac{s}{n}q_s, \quad \pi_{i,s}^- := \mathbb{P}_q(S \in C_{i,s}^-) = \frac{n-s}{n}q_s.$$

For estimating ϕ_i , the score $\gamma_{i,q}(S) := \rho_i(S)/q(S)$ is constant on each membership-size cell:

$$\gamma_{i,q}(S) = \begin{cases} \frac{1}{sq_s}, & S \in C_{i,s}^+, \\ -\frac{1}{(n-s)q_s}, & S \in C_{i,s}^-. \end{cases} \quad (27)$$

Thus the compatibility condition in Proposition 1 holds for the membership-size partition.

Proof of Corollary 1. Let

$$K_+ = |\{C \in \mathcal{C} : \mathbb{P}_q(S \in C) > 0\}|.$$

For the membership-size partition above, $K_+ = 2(n-1) = O(n)$. We bound the two design quantities in Theorem 1,

$$\Gamma_{2,\mathcal{C}}^2 = \frac{1}{K_+} \sum_{C \in \mathcal{C} : \mathbb{P}_q(S \in C) > 0} \mathbb{E}_q[\gamma_{i,q}(S)^2 \mid S \in C], \quad \Gamma_\infty = \sup_{S: q(S) > 0} |\gamma_{i,q}(S)|.$$

Because $\gamma_{i,q}$ is constant on each cell, the conditional expectations are just the squared cell values in (27).

OFA. For Shapley-value estimation, the OFA size distribution is

$$q_s^{\text{OFA}} = \frac{\{s(n-s)\}^{-1/2}}{\sum_{r=1}^{n-1} \{r(n-r)\}^{-1/2}}, \quad s = 1, \dots, n-1.$$

The normalizing denominator is $\Theta(1)$. Substituting this q_s into (27) gives

$$|\gamma_{i,q}(S)|^2 \asymp \begin{cases} (n-s)/s, & S \in C_{i,s}^+, \\ s/(n-s), & S \in C_{i,s}^-. \end{cases}$$

Therefore

$$\Gamma_{2,\emptyset}^2 \lesssim \frac{1}{n} \sum_{s=1}^{n-1} \left\{ \frac{n-s}{s} + \frac{s}{n-s} \right\} = O(\log n),$$

$$\Gamma_{\infty}^2 = O(n).$$

Moreover,

$$\pi_{\min} = \min_{1 \leq s \leq n-1} \left\{ \frac{s}{n} q_s^{\text{OFA}}, \frac{n-s}{n} q_s^{\text{OFA}} \right\} = \Theta(n^{-3/2}).$$

Applying Theorem 1 yields

$$\mathbb{E}_q(r_m^2) \leq C_1 \|u\|_{\infty}^2 \left\{ \frac{n \log n}{m^2} + n e^{-C_2 m/n^{3/2}} \right\},$$

after absorbing constants into C_1, C_2 .

Stratified SVARM. Stratified SVARM uses the harmonic size allocation

$$q_s^{\text{SVARM}} = \frac{\{\min(s, n-s)\}^{-1}}{\sum_{r=1}^{n-1} \{\min(r, n-r)\}^{-1}}, \quad s = 1, \dots, n-1.$$

The denominator is $\Theta(\log n)$, so

$$q_s^{\text{SVARM}} \asymp \frac{1}{\min(s, n-s) \log n}.$$

Using (27),

$$|\gamma_{i,q}(S)| \lesssim \log n$$

on both $C_{i,s}^+$ and $C_{i,s}^-$. Hence

$$\Gamma_{2,\emptyset}^2 = O(\log^2 n), \quad \Gamma_{\infty}^2 = O(\log^2 n).$$

The smallest active cell probability satisfies

$$\pi_{\min} = \min_{1 \leq s \leq n-1} \left\{ \frac{s}{n} q_s^{\text{SVARM}}, \frac{n-s}{n} q_s^{\text{SVARM}} \right\} = \Theta((n \log n)^{-1}).$$

Theorem 1 therefore gives

$$\mathbb{E}_q(r_m^2) \leq C_1 \|u\|_{\infty}^2 \left\{ \frac{n \log^2 n}{m^2} + \log^2 n e^{-C_2 m/(n \log n)} \right\}.$$

This proves the two displayed bounds in Corollary 1. □

B.2 Details for Non-Asymptotic Results for Ridge Regression Estimation

We next instantiate Theorem 2 for the WLS geometry used by KernelSHAP and LeverageSHAP. Let

$$\mathbf{C} := \mathbf{I} - \frac{1}{n} \mathbf{1} \mathbf{1}^\top$$

be the projection onto the sum-zero subspace. For $1 \leq |S| \leq n-1$, define the centered coalition feature

$$\mathbf{z}(S) := \left(\mathbf{1}\{1 \in S\}, \dots, \mathbf{1}\{n \in S\} \right)^\top - \frac{|S|}{n} \mathbf{1} \in \mathbb{R}^n$$

and the Shapley kernel weight

$$w(S) := \frac{n-1}{\binom{n}{|S|} |S| (n-|S|)}.$$

The population WLS problem with this feature map identifies the centered Shapley vector $\mathbf{C}\phi(u)$; the deterministic efficiency component $\{u([n]) - u(\emptyset)\} \mathbf{1}/n$ can then be added separately if the full Shapley vector is needed.

For this design,

$$\mathbf{G} = \sum_{1 \leq |S| \leq n-1} w(S) \mathbf{z}(S) \mathbf{z}(S)^\top = \frac{n-1}{n} \mathbf{C}, \quad \mathbf{G}^+ = \frac{n}{n-1} \mathbf{C}. \quad (28)$$

Indeed, conditional on $|S| = s$, the centered inclusion vector has covariance

$$\frac{s(n-s)}{n(n-1)} \mathbf{C},$$

and summing the weighted contribution over $s = 1, \dots, n-1$ gives (28). Consequently, $\kappa_{\mathbf{G}} = 1$. Also,

$$\|\mathbf{z}(S)\|_2^2 = \frac{|S|(n-|S|)}{n}, \quad \mathbf{z}(S)^\top \mathbf{G}^+ \mathbf{z}(S) = \frac{|S|(n-|S|)}{n-1}. \quad (29)$$

For the centered readout, $R_{\mathbf{A}}^2 = \sigma_{\max}(\mathbf{C}^\top \mathbf{G}^+ \mathbf{C}) = O(1)$, and the feature dimension is $d_{\mathbf{z}} = n$.

Proof of Corollary 2. We evaluate the quantities in Theorem 2 under the two sampling laws.

KernelSHAP. KernelSHAP samples coalitions proportionally to the Shapley kernel weight:

$$q_{\text{KS}}(S) = \frac{w(S)}{W_{\text{KS}}}, \quad W_{\text{KS}} := \sum_{1 \leq |T| \leq n-1} w(T) = \frac{2(n-1)}{n} H_{n-1} = O(\log n),$$

where $H_{n-1} = \sum_{s=1}^{n-1} s^{-1}$. Equivalently,

$$q_{\text{KS}}(S) = \frac{n}{2H_{n-1}} \frac{1}{\binom{n}{|S|} |S| (n - |S|)}.$$

Thus $\tilde{w}(S) = w(S)/q_{\text{KS}}(S) = W_{\text{KS}}$. Combining this with (29),

$$\ell(S) = \tilde{w}(S) \mathbf{z}(S)^\top \mathbf{G}^+ \mathbf{z}(S) = W_{\text{KS}} \frac{|S|(n - |S|)}{n - 1},$$

so

$$L_{\text{KS}} = \sup_S \ell(S) = O(n \log n).$$

The weighted norms are bounded by

$$\|u\|_{2, \tilde{w}}^2 \leq W_{\text{KS}} \|u\|_\infty^2 = O(\log n) \|u\|_\infty^2, \quad \|u\|_{\infty, \tilde{w}}^2 \leq O(\log n) \|u\|_\infty^2.$$

Substituting $R_{\mathbf{A}}^2 = O(1)$, $d_{\mathbf{z}} = n$, $\kappa_{\mathbf{G}} = 1$, $L_{\text{KS}} = O(n \log n)$, and the preceding norm bounds into Theorem 2 gives

$$\begin{aligned} \mathbb{E}_q(\|\mathbf{r}_m\|_2^2) \leq C_1 \|u\|_\infty^2 & \left[\frac{n^2 \log^4 n}{m^2} \left(1 + \frac{n \log^2 n}{m} \right) \right. \\ & \left. + n^3 m^2 \log^3 n e^{-C_2 m / (n \log n)} \right]. \end{aligned}$$

LeverageSHAP. LeverageSHAP samples proportionally to the WLS leverage contribution

$$w(S) \mathbf{z}(S)^\top \mathbf{G}^+ \mathbf{z}(S).$$

By (29),

$$w(S) \mathbf{z}(S)^\top \mathbf{G}^+ \mathbf{z}(S) = \frac{1}{\binom{n}{|S|}}.$$

The total leverage mass is $n - 1 = \text{rank}(\mathbf{G})$, so

$$q_{\text{Lev}}(S) = \frac{1}{(n-1) \binom{n}{|S|}}, \quad 1 \leq |S| \leq n-1.$$

Equivalently, LeverageSHAP draws the size uniformly from $\{1, \dots, n-1\}$ and then draws a coalition uniformly within that size. For this law,

$$\tilde{w}(S) = \frac{w(S)}{q_{\text{Lev}}(S)} = \frac{(n-1)^2}{|S|(n-|S|)}.$$

Using (29),

$$\ell(S) = \frac{(n-1)^2}{|S|(n-|S|)} \frac{|S|(n-|S|)}{n-1} = n-1,$$

and hence $L_{\text{Lev}} = n-1 = O(n)$. The weighted norms satisfy

$$\|u\|_{2,\tilde{w}}^2 = \sum_{1 \leq |S| \leq n-1} w(S) u(S)^2 \leq W_{\text{KS}} \|u\|_{\infty}^2 = O(\log n) \|u\|_{\infty}^2,$$

and

$$\|u\|_{\infty,\tilde{w}}^2 \leq O(n) \|u\|_{\infty}^2,$$

because $|S|(n-|S|) \geq n-1$ on the interior strata. Substituting

$$R_{\mathbf{A}}^2 = O(1), \quad d_{\mathbf{z}} = n, \quad \kappa_{\mathbf{G}} = 1, \quad L_{\text{Lev}} = O(n)$$

into Theorem 2 gives

$$\mathbb{E}_q(\|\mathbf{r}_m\|_2^2) \leq C_1 \|u\|_{\infty}^2 \left[\frac{n^2 \log^2 n}{m^2} \left(1 + \frac{n \log n}{m} \right) + n^4 m^2 e^{-C_2 m/n} \right].$$

This proves Corollary 2. □

C Details of Algorithms

This appendix gives implementation details for the EASE construction in Section 5. We keep the notation of the main paper: the target vector is

$$\boldsymbol{\tau}_{\mathbf{A}}(u) = \left(\tau_{\mathbf{a}_1}(u), \dots, \tau_{\mathbf{a}_d}(u) \right)^\top, \quad \mathbf{A} = [\mathbf{a}_1, \dots, \mathbf{a}_d].$$

Define the vector of weighted-average coefficients

$$\boldsymbol{\rho}_{\mathbf{A}}(S) := \left(\rho_{\mathbf{a}_1}(S), \dots, \rho_{\mathbf{a}_d}(S) \right)^\top, \quad S \subseteq [n]. \quad (30)$$

Then $\boldsymbol{\tau}_{\mathbf{A}}(u) = \sum_{S \subseteq [n]} \boldsymbol{\rho}_{\mathbf{A}}(S) u(S)$. The first two subsections below apply to the general vector-surrogate setting $\mathbf{h} = (h_1, \dots, h_d)$. The shared-linear construction in Section C.3 is a computationally convenient special case.

C.1 Initialization for a Fixed Sampling Partition

Fix the partition $\mathcal{C} = \{C_1, \dots, C_K\}$ used to restrict the sampling law in Section 5. For a law that is constant within each cell, write $q(S) = \pi_k$ for $S \in C_k$, with $\sum_{k=1}^K |C_k| \pi_k = 1$. For a general vector surrogate $\mathbf{h} = (h_1, \dots, h_d)$, the cellwise residual scale is

$$M_k(\mathbf{h}) := \frac{1}{|C_k|} \sum_{S \in C_k} \sum_{j=1}^d \rho_{\mathbf{a}_j}(S)^2 \{u(S) - h_j(S)\}^2. \quad (31)$$

For fixed $\mathbf{h}$, the leading uncentered part of the first-order Euclidean risk within this partition is

$$\sum_{k=1}^K \frac{|C_k| M_k(\mathbf{h})}{\pi_k}.$$

Minimizing this expression over $(\pi_1, \dots, \pi_K)$ gives

$$q^*(S; \mathbf{h}) = \pi_k^*(\mathbf{h}), \quad \pi_k^*(\mathbf{h}) = \frac{\sqrt{M_k(\mathbf{h})}}{\sum_{r=1}^K |C_r| \sqrt{M_r(\mathbf{h})}}, \quad S \in C_k. \quad (32)$$

Before any informative surrogate has been learned, the residual terms in (31) are unknown. A simple initialization is obtained by treating the residuals $|u(S) - h_j(S)|$ as uniformly bounded by a common constant, or more generally as having the same order across cells and coordinates. Under either interpretation, the unknown common residual scale drops out of (32). We therefore initialize from

$$M_{k,0} := \frac{1}{|C_k|} \sum_{S \in C_k} \sum_{j=1}^d \rho_{\mathbf{a}_j}(S)^2, \quad q^{\text{init}}(S) = \pi_k^{\text{init}} := \frac{\sqrt{M_{k,0}}}{\sum_{r=1}^K |C_r| \sqrt{M_{r,0}}}, \quad S \in C_k. \quad (33)$$

Thus the pilot allocation is determined by the target weights and the chosen partition alone.

C.2 Flooring the Learned Sampling Law

The plug-in law learned from pilot data should also be defined in the general vector-surrogate setting. After drawing $\{(S_t^{\text{init}}, Y_t^{\text{init}})\}_{t=1}^{m_{\text{init}}}$ from q^{init} , fit $\hat{\mathbf{h}}^{\text{init}} = (\hat{h}_1^{\text{init}}, \dots, \hat{h}_d^{\text{init}})$ by the profiled empirical criterion in (19). For

$$N_k := \sum_{t=1}^{m_{\text{init}}} \mathbb{I}(S_t^{\text{init}} \in C_k),$$

estimate the cellwise residual scale by

$$\hat{M}_k = \frac{1}{\max\{N_k, 1\}} \sum_{t=1}^{m_{\text{init}}} \mathbb{I}(S_t^{\text{init}} \in C_k) \sum_{j=1}^d \rho_{\mathbf{a}_j}(S_t^{\text{init}})^2 \{Y_t^{\text{init}} - \hat{h}_j^{\text{init}}(S_t^{\text{init}})\}^2. \quad (34)$$

This convention sets $\hat{M}_k = 0$ for empty pilot cells. The unfloored plug-in law is

$$\tilde{q}(S) = \tilde{\pi}_k, \quad \tilde{\pi}_k = \frac{\sqrt{\hat{M}_k}}{\sum_{r=1}^K |C_r| \sqrt{\hat{M}_r}}, \quad S \in C_k,$$

when the denominator is positive; otherwise we keep $\tilde{q} = q^{\text{init}}$.

In finite samples, $\tilde{q}$ can be too aggressive. A small estimated $\hat{M}_k$ may come from pilot noise rather than from a genuinely negligible residual contribution, and a very small $q(S)$ inflates the correction weights $\rho_{\mathbf{a}_j}(S)/q(S)$. We therefore use a floored law

$$q^{\text{floor}}(S) = (1 - \epsilon)\tilde{q}(S) + \epsilon q^{\text{base}}(S), \quad q^{\text{base}}(S) := \frac{1}{K|C_k|}, \quad S \in C_k, \quad \epsilon \in (0, 1). \quad (35)$$

Thus every coalition $S \in C_k$ receives probability at least $\epsilon/(K|C_k|)$. The floor is a stability device: it prevents empty or underestimated cells from producing nearly infinite inverse-probability weights, while preserving the residual-aware allocation when ϵ is small.

C.3 Shared Linear Surrogate Learned On the Fly

We now specialize to a practical regime in which a single surrogate is shared across all coordinates and is linear in a chosen feature vector:

$$h_{\boldsymbol{\beta}}(S) = \mathbf{x}(S)^\top \boldsymbol{\beta}, \quad \mathbf{x}(S) \in \mathbb{R}^p.$$

Assume throughout this subsection that the sampling law q is fixed and positive on the support of $\boldsymbol{\rho}_A$. For observations $S_t \sim q$, set

$$\boldsymbol{\omega}_t := \frac{\boldsymbol{\rho}_A(S_t)}{q(S_t)}, \quad a_t := \|\boldsymbol{\omega}_t\|_2^2, \quad \mathbf{x}_t := \mathbf{x}(S_t).$$

Given the first t observations, the profiled shared-linear objective is

$$L_t(\boldsymbol{\beta}, \boldsymbol{\mu}) := \sum_{s=1}^t \left\| \boldsymbol{\omega}_s \{Y_s - \mathbf{x}_s^\top \boldsymbol{\beta}\} - \boldsymbol{\mu} \right\|_2^2 + \lambda \|\boldsymbol{\beta}\|_2^2, \quad \lambda > 0. \quad (36)$$

This is the empirical profiled criterion from (19) specialized to a shared linear surrogate

under a fixed sampling law.

The minimizer can be updated from four additive sufficient statistics:

$$\mathbf{R}_t := \sum_{s=1}^t a_s \mathbf{x}_s \mathbf{x}_s^\top, \quad \mathbf{d}_t := \sum_{s=1}^t a_s \mathbf{x}_s Y_s, \quad (37)$$

$$\mathbf{U}_t := \sum_{s=1}^t \boldsymbol{\omega}_s \mathbf{x}_s^\top, \quad \mathbf{v}_t := \sum_{s=1}^t \boldsymbol{\omega}_s Y_s. \quad (38)$$

Profiling out $\boldsymbol{\mu}$ gives

$$\hat{\boldsymbol{\beta}}_t = \left(\mathbf{R}_t - \frac{1}{t} \mathbf{U}_t^\top \mathbf{U}_t + \lambda \mathbf{I}_p \right)^{-1} \left(\mathbf{d}_t - \frac{1}{t} \mathbf{U}_t^\top \mathbf{v}_t \right), \quad (39)$$

and

$$\hat{\boldsymbol{\mu}}_t = \frac{1}{t} (\mathbf{v}_t - \mathbf{U}_t \hat{\boldsymbol{\beta}}_t).$$

Each statistic in (37)–(38) updates by addition when a new utility value is observed. Hence the shared linear surrogate can be updated on the fly without resolving the full least squares problem from scratch.

C.4 Complement-Pair Sampling

Complement-pair sampling observes $u(S)$ and $u(S^c)$ together. In this subsection, we only consider the common sign-symmetric case in which, for every target coordinate and every coalition,

$$\rho_{\mathbf{a}_j}(S^c) = -\rho_{\mathbf{a}_j}(S), \quad j = 1, \dots, d. \quad (40)$$

With the sign convention in (3), this is the relation satisfied by the Shapley values, Beta Shapley values with symmetric weights, and Banzhaf values. In practice, complement pair sampling is a natural choice for sign-symmetric targets [33].

Under (40), the target can be written as a sum over complement contrasts:

$$\boldsymbol{\tau}_\mathbf{A}(u) = \frac{1}{2} \sum_{S \subseteq [n]} \boldsymbol{\rho}_\mathbf{A}(S) \{u(S) - u(S^c)\}.$$

Suppose an ordered coalition S is drawn from a law q , but the observation reveals the pair (S, S^c) . The unordered pair mass is

$$q_{\text{pair}}(S) := q(S) + q(S^c),$$

which is the same for S and S^c . For a general vector surrogate $\mathbf{h} = (h_1, \dots, h_d)$, define the

coordinatewise residual contrast

$$\Delta_{\mathbf{h},j}(S) := \{u(S) - h_j(S)\} - \{u(S^c) - h_j(S^c)\}, \quad j = 1, \dots, d,$$

and let $\Delta_{\mathbf{h}}(S) = (\Delta_{\mathbf{h},1}(S), \dots, \Delta_{\mathbf{h},d}(S))^\top$. Then the paired AIPW score

$$\hat{\tau}^{\text{pair}}(S; \mathbf{h}) := \tau_{\mathbf{A}}(\mathbf{h}) + \frac{\boldsymbol{\rho}_{\mathbf{A}}(S) \odot \Delta_{\mathbf{h}}(S)}{q_{\text{pair}}(S)} \quad (41)$$

is unbiased when $S \sim q$, provided $q_{\text{pair}}(S) > 0$ whenever $\boldsymbol{\rho}_{\mathbf{A}}(S) \neq \mathbf{0}$. Indeed, the two ordered representatives S and S^c of the same pair contribute the same contrast because $\boldsymbol{\rho}_{\mathbf{A}}(S^c) = -\boldsymbol{\rho}_{\mathbf{A}}(S)$ and $\Delta_{\mathbf{h}}(S^c) = -\Delta_{\mathbf{h}}(S)$.

The same residual-aware sampling idea can be applied at the pair level. For a partition $\mathcal{P} = \{P_1, \dots, P_L\}$ of complement pairs, define

$$M_{\ell}^{\text{pair}}(\mathbf{h}) := \frac{1}{|P_{\ell}|} \sum_{\{S, S^c\} \in P_{\ell}} \|\boldsymbol{\rho}_{\mathbf{A}}(S) \odot \Delta_{\mathbf{h}}(S)\|_2^2.$$

The corresponding plug-in allocation assigns more pair mass to cells with large estimated residual contrasts. As in the single-coalition case, a floor should be added to the learned pair law to keep $q_{\text{pair}}(S)$ bounded away from zero on active pairs.

D Additional Experimental Results

This appendix provides the AUCC benchmark results for the higher low-order signal settings omitted from the main text. The qualitative pattern is consistent with the $\eta = 0.25$ AUCC benchmark in Figure 2: EASE-FO and EASE-SP remain among the most sample-efficient estimators, with EASE-SP typically strongest for Shapley and Beta-Shapley targets. Increasing η from 0.5 to 0.75 changes the scale of the errors but does not alter the main comparison trends.

E Monte Carlo Estimation with Bundled Samples

E.1 Bundled Sampling and Identification

The main text studies single-coalition sampling, where Monte Carlo draw $t = 1, \dots, m$ returns a pair $(S_t, u(S_t))$ with $S_t \subseteq [n]$. This appendix considers the more general case in which one draw reveals several coupled coalition values. Specifically, draw t returns




Figure 3. AUCC benchmark on the SOU games for $\eta = 0.5$ and $\eta = 0.75$. Each x-axis label is a target probabilistic value. Lower AUCC indicates better performance.

$(\mathbf{T}_t, u(\mathbf{T}_t))$, where

$$\mathbf{T}_t = (S_{t,1}, \dots, S_{t,r})$$

is a random bundle of r coalitions and

$$u(\mathbf{T}_t) = (u(S_{t,1}), \dots, u(S_{t,r}))$$

is the corresponding vector of utility values. The bundle is sampled from a known law Q on $(2^{[n]})^r$. We refer to this observation model as *bundled sampling*.

Bundled sampling covers several common Monte Carlo designs. Complement-pair sampling returns a random coalition S together with its complement S^c [35]. Edge-pair sampling returns a random coalition S and its one-player extension $S \cup \{i\}$ for some player i [17, 25]. Permutation-path sampling returns the full sequence of coalitions generated along a random permutation of the players [1, 2, 24, 34].

These designs differ in the bundle they observe, but the target remains the same linear functional $\tau_{\mathbf{a}}(u) = \sum_{S \subseteq [n]} \rho_{\mathbf{a}}(S) u(S)$. Under a fixed bundle law Q , the basic question is how to assign weights to the observed bundle entries so that their expectation recovers this target. We formalize this requirement below.

Definition E.1 (Feasible bundle weighting). Fix the target $\tau_{\mathbf{a}}(u) = \sum_{S \subseteq [n]} \rho_{\mathbf{a}}(S) u(S)$ and a bundle law Q on $(2^{[n]})^r$. An r -dimensional function $\gamma_{\mathbf{a},Q} : (2^{[n]})^r \rightarrow \mathbb{R}^r$ is a *feasible bundle*

weighting under Q if, for every utility function $u : 2^{[n]} \rightarrow \mathbb{R}$,

$$\tau_{\mathbf{a}}(u) = \mathbb{E}_Q \left[\boldsymbol{\gamma}_{\mathbf{a},Q}(\mathbf{T})^\top u(\mathbf{T}) \right].$$

We denote the class of all feasible bundle weightings by $\Gamma_{\mathbf{a}}(Q)$.

Single-coalition sampling is the special case in which the feasible weighting is unique. When $r = 1$, the definition reduces to the inverse-probability weighting $\boldsymbol{\gamma}_{\mathbf{a},Q}(S) = \rho_{\mathbf{a}}(S)/Q(S)$ for every $S \subseteq [n]$ with $Q(S) > 0$. When $r \geq 2$, feasibility instead imposes a system of linear constraints on a vector-valued weighting, and that system can have multiple solutions.

The complement-pair design makes this nonuniqueness explicit. Suppose $\mathbf{T} = (S, S^c)$ with $S \sim q$, and write the weighting as $\boldsymbol{\gamma}_{\mathbf{a},Q}(S, S^c) = (\gamma_1(S), \gamma_2(S))$. Matching the coefficient of each $u(S)$ in Definition E.1 gives the linear constraints

$$q(S)\gamma_1(S) + q(S^c)\gamma_2(S^c) = \rho_{\mathbf{a}}(S), \quad S \subseteq [n].$$

Thus, whenever both $q(S)$ and $q(S^c)$ are positive, $\gamma_1(S)$ can be chosen freely and $\gamma_2(S^c)$ adjusted to equal $\{\rho_{\mathbf{a}}(S) - q(S)\gamma_1(S)\}/q(S^c)$. Equivalently, if q has full support, any allocation function $\lambda : 2^{[n]} \rightarrow \mathbb{R}$ induces a feasible weighting through

$$\gamma_1(S) = \frac{\lambda(S)\rho_{\mathbf{a}}(S)}{q(S)}, \quad \gamma_2(S) = \frac{\{1 - \lambda(S^c)\}\rho_{\mathbf{a}}(S^c)}{q(S)}.$$

Different choices of λ distribute the same target coefficient across the two possible appearances of a coalition in the ordered complement pair. For instance, $\lambda \equiv 1$ assigns each coefficient to the first component when the coalition is sampled directly, whereas $\lambda \equiv 0$ assigns it to the second component when its complement is sampled. Hence the same target functional $\tau_{\mathbf{a}}(u)$ can be represented by many feasible bundle weightings.

E.2 Bundle-Regular Estimators

The same regularity idea used in the main text extends once the single coalition weighting is replaced by a feasible bundle weighting. We first record the coefficient form of Definition E.1. For $\boldsymbol{\gamma}_{\mathbf{a},Q} \in \Gamma_{\mathbf{a}}(Q)$, feasibility is equivalent to

$$\rho_{\mathbf{a}}(S) = \mathbb{E}_Q \left[\sum_{\ell=1}^r \boldsymbol{\gamma}_{\mathbf{a},Q,\ell}(\mathbf{T}) \mathbf{1}\{S_\ell = S\} \right], \quad S \subseteq [n]. \quad (42)$$

This form is useful because it separates the sampling law Q from the choice of weighting used for the entries of a bundle.

For a feasible weighting $\gamma \in \Gamma_{\mathbf{a}}(Q)$ and a surrogate $h : 2^{[n]} \rightarrow \mathbb{R}$, define the bundle influence term

$$\psi_{\gamma,h}(\mathbf{T}; u) := \gamma(\mathbf{T})^\top \{u(\mathbf{T}) - h(\mathbf{T})\} + \tau_{\mathbf{a}}(h) - \tau_{\mathbf{a}}(u). \quad (43)$$

By feasibility, $\mathbb{E}_Q\{\psi_{\gamma,h}(\mathbf{T}; u)\} = 0$. This gives the following analogue of Definition 1.

Definition E.2 (Bundle-regular estimator sequence). Fix a bundle law Q , a feasible weighting $\gamma \in \Gamma_{\mathbf{a}}(Q)$, and a working surrogate class $\mathcal{H}$ on $2^{[n]}$. A sequence of estimators $\{\hat{\tau}_{\mathbf{a},m}\}_{m \geq 1}$ is *bundle-regular relative to $(\gamma, \mathcal{H})$ under Q* if there exist nonrandom functions $h_m \in \mathcal{H}$ such that, for i.i.d. bundles $\mathbf{T}_1, \dots, \mathbf{T}_m \sim Q$,

$$\hat{\tau}_{\mathbf{a},m}(u) - \tau_{\mathbf{a}}(u) = \frac{1}{m} \sum_{t=1}^m \psi_{\gamma,h_m}(\mathbf{T}_t; u) + r_m, \quad \mathbb{E}_Q(r_m^2) = o(m^{-1}). \quad (44)$$

Definition E.2 keeps the same separation as in the single-coalition case: the weighting describes the identifying representation, whereas h_m describes the first-order surrogate adjustment.

This class includes several bundled estimators in the literature. Edge-pair methods such as sampling lift [25] and weighted sampling lift [17] are HT-type weighted averages with a design-specific γ and the working class $\mathcal{H} = \{0\}$. Permutation-based estimators and their stratified refinements [1, 2, 24, 34] can be viewed as bundled Hájek-type weighted averages, where the normalization over permutation strata induces the corresponding working class. Paired or complement-paired variants of WLS estimators, including LeverageSHAP [26] and PolySHAP [12], fit the same definition with the feasible weighting determined by the bundled sampling law and $\mathcal{H}$ given by the WLS feature span.

Example E.1 (Complement-pair sampling). Complementary-contribution sampling [35] evaluates coalition values in complement pairs. One draw returns $\mathbf{T} = (S, S^c)$ with $S \sim q$, so the same utility query batch contains both $u(S)$ and $u(S^c)$. This coupling is useful for symmetric semivalues, where the coefficients of a coalition and its complement have opposite signs, and the estimator can be written in terms of complementary differences. In our notation, a canonical feasible weighting for this design is

$$\gamma_1^{\text{comp}}(S, S^c) := \frac{\rho_{\mathbf{a}}(S)}{q(S) + q(S^c)}, \quad \gamma_2^{\text{comp}}(S, S^c) := \frac{\rho_{\mathbf{a}}(S^c)}{q(S) + q(S^c)}, \quad (45)$$

on pairs for which $q(S) + q(S^c) > 0$. Indeed, for every $S \subseteq [n]$,

$$q(S)\gamma_1^{\text{comp}}(S, S^c) + q(S^c)\gamma_2^{\text{comp}}(S^c, S) = \rho_{\mathbf{a}}(S).$$

Thus the contribution of a coalition can be assigned across its two possible appearances in an ordered complement pair. The construction in Zhang et al. [35] uses this paired

observation structure, together with allocation across coalition sizes, to reduce the variance of direct Monte Carlo estimation.

Example E.2 (Edge-pair sampling). Sampling lift [25] and weighted sampling lift [17] evaluate marginal-contribution pairs. Fix a player i , let q_i be a law on $2^{[n] \setminus \{i\}}$, and sample $\mathbf{T} = (S, S \cup \{i\})$ with $S \sim q_i$. The draw directly reveals the difference $u(S \cup \{i\}) - u(S)$. For the coordinate target $\tau_{\mathbf{e}_i}(u) = \phi_i(u)$, the marginal-contribution representation gives

$$\phi_i(u) = \sum_{S \subseteq [n] \setminus \{i\}} \alpha_i^{(n)}(S) \{u(S \cup \{i\}) - u(S)\}.$$

Thus a feasible edge weighting is

$$\gamma_1^{\text{edge}, i}(S, S \cup \{i\}) := -\frac{\alpha_i^{(n)}(S)}{q_i(S)}, \quad \gamma_2^{\text{edge}, i}(S, S \cup \{i\}) := \frac{\alpha_i^{(n)}(S)}{q_i(S)}. \quad (46)$$

The original sampling-lift construction uses this edge-pair observation to estimate Shapley-type targets from sampled marginal contributions, while weighted sampling lift modifies the sampling and weighting scheme to cover Beta Shapley and related probabilistic values.

More general edge-pair designs can sample both the base coalition and the added player. If the base coalition–player pair is sampled from a law λ on $\mathcal{E}_n = \{(S, i) : S \subseteq [n], i \notin S\}$, then feasibility requires

$$\rho_{\mathbf{a}}(S) = \sum_{i \notin S} \lambda(S, i) \gamma_1(S, i) + \sum_{i \in S} \lambda(S \setminus \{i\}, i) \gamma_2(S \setminus \{i\}, i), \quad S \subseteq [n]. \quad (47)$$

Unlike the fixed-player representation in Example E.2, this system couples the weighting choices across many directed edges.

E.3 First-Order Risk with a Fixed Bundle Weighting

For a bundle-regular estimator, the leading MSE is governed by the variance of the bundle influence term. Define

$$V_Q(\boldsymbol{\gamma}, h) := \text{Var}_Q\{\psi_{\boldsymbol{\gamma}, h}(\mathbf{T}; u)\} = \text{Var}_Q[\boldsymbol{\gamma}(\mathbf{T})^\top \{u(\mathbf{T}) - h(\mathbf{T})\}]. \quad (48)$$

If (44) holds and $\mathbb{E}_Q(r_m^2) = o\{V_Q(\boldsymbol{\gamma}, h_m)/m\}$, then

$$\mathbb{E}_Q[\{\hat{\tau}_{\mathbf{a}, m}(u) - \tau_{\mathbf{a}}(u)\}^2] = \frac{V_Q(\boldsymbol{\gamma}, h_m)}{m} \{1 + o(1)\}. \quad (49)$$

Equivalently,

$$V_Q(\gamma, h) = \inf_{t \in \mathbb{R}} \mathbb{E}_Q \left[\left\{ \gamma(\mathbf{T})^\top \{u(\mathbf{T}) - h(\mathbf{T})\} - t \right\}^2 \right]. \quad (50)$$

Thus, unlike the single-coalition case, the first-order criterion depends on three objects: the bundle law Q , the feasible weighting γ , and the surrogate h . For a fixed weighting, the oracle surrogate problem is

$$\inf_{h \in \mathcal{H}} V_Q(\gamma, h).$$

At the fully joint level one may instead consider

$$\inf_{\gamma \in \Gamma_{\mathbf{a}}(Q), h \in \mathcal{H}} V_Q(\gamma, h),$$

but the weighting optimization is generally a constrained problem over functions on the whole bundle space. Equation (47) illustrates why this optimization can be difficult: even for edge pairs, the identifying constraints are globally coupled.

E.4 Rao–Blackwell Weighting for Complement Pairs

Complement-pair sampling is an important exception in which the weighting optimization has an explicit solution. Let $r_h(S) := u(S) - h(S)$. For the design $\mathbf{T} = (S, S^c)$, $S \sim q$, any feasible weighting (γ_1, γ_2) must satisfy

$$q(S)\gamma_1(S, S^c) + q(S^c)\gamma_2(S^c, S) = \rho_{\mathbf{a}}(S), \quad S \subseteq [n]. \quad (51)$$

The canonical weighting in (45) is the Rao–Blackwellized version of any such feasible ordered-pair weighting.

Proposition E.1 (Rao–Blackwell optimality for complement pairs). *Assume the complement-pair design $\mathbf{T} = (S, S^c)$, $S \sim q$, and suppose $q(S) + q(S^c) > 0$ whenever the pair $\{S, S^c\}$ can contribute to the target. Let (γ_1, γ_2) be any feasible weighting satisfying (51). Then, for every fixed surrogate h ,*

$$\begin{aligned} & \text{Var}_q[\gamma_1^{\text{comp}}(S, S^c)r_h(S) + \gamma_2^{\text{comp}}(S, S^c)r_h(S^c)] \\ & \leq \text{Var}_q[\gamma_1(S, S^c)r_h(S) + \gamma_2(S, S^c)r_h(S^c)]. \end{aligned}$$

Proof. Let $S_0 \sim q$ denote the sampled base coalition, let $\mathcal{U}(S_0) = \{S_0, S_0^c\}$ denote the unordered complement pair, and define

$$X_\gamma(S_0) := \gamma_1(S_0, S_0^c)r_h(S_0) + \gamma_2(S_0, S_0^c)r_h(S_0^c).$$

Conditioning on $\mathcal{U}(S_0) = \{S, S^c\}$, the two possible ordered draws are (S, S^c) and (S^c, S) ,

with probabilities proportional to $q(S)$ and $q(S^c)$. Therefore,

$$\begin{aligned}
\mathbb{E}_q\{X_\gamma(S_0) \mid \mathcal{U}(S_0) = \{S, S^c\}\} &= \frac{\{q(S)\gamma_1(S, S^c) + q(S^c)\gamma_2(S^c, S)\}r_h(S)}{q(S) + q(S^c)} \\
&\quad + \frac{\{q(S)\gamma_2(S, S^c) + q(S^c)\gamma_1(S^c, S)\}r_h(S^c)}{q(S) + q(S^c)} \\
&= \frac{\rho_{\mathbf{a}}(S)r_h(S) + \rho_{\mathbf{a}}(S^c)r_h(S^c)}{q(S) + q(S^c)} \\
&= \gamma_1^{\text{comp}}(S, S^c)r_h(S) + \gamma_2^{\text{comp}}(S, S^c)r_h(S^c),
\end{aligned}$$

where the second equality uses (51) for S and S^c . The last display is measurable with respect to the unordered pair $\mathcal{U}(S_0)$. The variance inequality follows from the law of total variance:

$$\text{Var}_q\{X_\gamma(S_0)\} = \text{Var}_q[\mathbb{E}_q\{X_\gamma(S_0) \mid \mathcal{U}(S_0)\}] + \mathbb{E}_q[\text{Var}_q\{X_\gamma(S_0) \mid \mathcal{U}(S_0)\}].$$

□

Proposition E.1 shows that complement pairs differ from general bundled designs: one can remove the artificial randomness coming from the orientation of the ordered pair without changing the target. For symmetric semivalues, including the Shapley value, $\rho_i(S^c) = -\rho_i(S)$ for a single coordinate target. Hence the Rao–Blackwellized weighting reduces to a weighted difference,

$$\frac{\rho_i(S)}{q(S) + q(S^c)} [\{u(S) - h(S)\} - \{u(S^c) - h(S^c)\}].$$

When $h \equiv 0$, this is precisely the common complementary-contribution form based on $u(S) - u(S^c)$. From the present viewpoint, that form is not only natural for symmetric semivalues; among feasible ordered complement-pair weightings, it is the first-order variance-minimizing choice.